High-frequency sorghum transformation toolkit enhances *Cas9* efficiency and expands promoter-editing capability with *SpRY*

Jianqiang Shen[1,2,5,6], Kiflom Aregawi[1,2,5], Sultana Anwar[1], Tamara Miller[1], Evan Groover[1,2],

Mantra Rajkumar[3], David Savage[2,3,4], Peggy G. Lemaux[1,2,6]

[1]Department of Plant and Microbial Biology, University of California, Berkeley CA 94720,

USA.

[2] Innovative Genomics Institute, University of California, Berkeley CA 94720, USA.

[3]Department of Molecular and Cell Biology, University of California, Berkeley CA 94720, USA.

[4]Howard Hughes Medical Institute, University of California, Berkeley CA 94720, USA

[5] These two authors contributed equally.

[6] Correspondence: shenjq@berkeley.edu (J. Shen) or lemauxpg@berkeley.edu (P. G. Lemaux)

Technology readiness

This study demonstrates the development of an integrated genome engineering platform that combines morphogenic gene-assisted transformation with CRISPR/Cas9- and SpRY-mediated mutagenesis for multiplex genome editing, establishing a Technology Readiness Level (TRL) of 4/5. This workflow enables the generation of allelic variants through editing of both functional and regulatory elements in the plant genome. Advancing toward TRL 5 will require additional validation in relevant sorghum genotypes, including elite varieties. Moreover, the PAM-flexible editor SpRY is still under optimization, particularly at non-canonical PAM sites, which may

limit its application for the modification of certain cis-regulatory elements. Further

improvements in precision genome editing, such as the integration of prime editing or targeted

knock-in technologies, will also be necessary to enable desired and predictable modifications.

With continued optimization and incorporation of these advanced tools, this robust genome

engineering platform has the potential to accelerate plant breeding and enable basic research

supporting plant-based solutions to address climate change.

27 Abstract

Sorghum bicolor L. (Moench), the fifth most important cereal crop internationally, is used as

food, feed, forage, and fuels. Importantly, sorghum's natural tolerance to environmental stresses

leads to tolerance to climate variation. To optimize sorghum germplasm for crop improvement

requires highly efficient genetic transformation and genome editing; however, sorghum has

historically been recalcitrant to these genetic approaches. In this study, we report a high-

efficiency engineering toolkit, optimized for genome editing via *Agrobacterium*-mediated

transformation. Using CRISPR/Cas9-based editing machinery, the genetic tools led to editing

efficiencies up to 95.7% when monocistronic guide RNAs targeted the phytoene desaturase

(*SbPDS*) gene. We also tested a PAM (protospacer adjacent motif)-broadened *Cas9* variant,

intronized *SpRY* (*ZmSpRYi*), achievling comparable editing efficiencies. Using this toolkit

permits exploitation of advanced editing tools, like prime editors, base editors, and targeted

knock-in methods. These genetic advances offer new methods for sustainable crop improvement

in sorghum and potentially other cereals.

Keywords

Sorghum bicolor, Wuschel2, Transformation, Engineering, Promoter editing, CRISPR/Cas9,

SpRY

Introduction

Meeting future crop production demands, given more variable climate conditions and an increasing world population, requires higher crop yields with the same land usage. *Sorghum bicolor* L. (Monech) is an ideal crop for such future applications due to the diversity of its germplasm, particularly with respect to its multiple abiotic stress tolerances [1–3]. Breeding improved sorghum varieties, using precise genome editing, will accelerate development of improved cultivars [4–6]. However, genome editing with CRISPR/*Cas9* in sorghum has lagged in efficiency relative to other crops due to inefficient transformation systems for diverse genotypes, low editing percentage, and lack of optimized tools [7,8]. The current work seeks to address these stated limitations by utilizing a novel set of resources to achieve a high efficiency of sorghum transformation and genome editing.

Sorghum genome engineering has historically been a challenging process due to recalcitrance of most sorghum genotypes to *in vitro* tissue culture and a lack of technical expertise and effective tools for transformation [7]. A significant challenge for genome engineering in crop species is the requirement for germline transmission of mutations, necessitating gene delivery in pluripotent cells. Morphogenic factors, such as Babyboom (BBM) and Wuschel2 (WUS2) or WUS2 alone, increase the population of cells in target tissues that are competent for germline transmission [9–11]. Previously we demonstrated that transient expression from *Bbm* and *Wus2*

during Agrobacterium-mediated gene delivery facilitated transformation of several recalcitrant

sorghum varieties [8]. Concurrently, Che et al. demonstrated that during sorghum transformation

incorporation of the *Wus2* gene alone increased the rate of CRISPR/*Cas9*-mediated knockout of

bmr, the brown midrib gene [11]. The breakthrough in transformation technology due to

morphogene-assisted transformation has greatly expanded the possibility of genotype-

independent gene editing; however, developing tools for high efficiency genome editing with

CRISPR remains a challenge.

The CRISPR/*Cas9* genome editing system, first discovered as a component of the adaptive

immune response in *Streptococcus pyogenes* bacteria [12], is a powerful tool for genome editing

in plants that has potential for functional genomics efforts and accelerated breeding [13].

However, editing in non-model plants lags in efficiency compared to animal and microbial

systems, due to several factors, including the plant 'regeneration bottleneck' [7] and possibly

sub-optimal activity of editors *in planta* [8,14]. Optimization of Cas endonuclease and guide

RNA (gRNA) expression have been proposed to increase the efficiency of genome editing in

sorghum [14]. In addition, use of Cas9 variants with expanded Protospacer Adjacent Motif

(PAM) recognition sequences increases editing scopes [15-20]. To address these challenges

further in Arabidopsis and tobacco, codon optimization of the Cas9 nuclease was used to

increase gene knockout editing efficiency [21,22]. Use of specific promoters and gRNA

transcriptional configurations was shown to increase sorghum editing efficiency, when the

endogenous sorghum U6 promoter was used to drive gRNAs [23]. An additional limitation for

introducing genetic variations at specific sites is the fact that the canonical *Cas9* endonuclease

requires the NGG (N = A, T, C, G) PAM site, which limits the genome sites available for editing

at sites with suitable PAM [24]. This limitation is especially relevant for engineering genomic

regions such as *cis*-regulatory elements [25], where intra-populational sequence variations in

promoter regions contribute significantly to phenotypic variations [26].

Several Cas9 variants have been engineered to overcome the PAM limitation of *SpCas9*

[16,27,28]. *SpRY* endonuclease, an engineered variant of *Streptococcus pyogenes Cas9,* has a

nearly PAM-less requirement and has been used for base editing and functional genetics in rice

[19,20]. To demonstrate the utility of the SpRY endonuclease for editing cis-regulatory elements

in sorghum, we tested its efficiency for editing the promoter of the *CYP79A1* gene, a step in

dhurrin biosynthesis. Presence of dhurrin in sorghum limits its use as a forage because some

varieties produce significant levels of this cyanogenic compound that can be fatal to livestock

and accumulates under stress conditions [29–32]. Because dhurrin is also implicated in nitrogen

storage and biotic stress tolerance [33,34], fine-tuning dhurrin biosynthesis, rather than fully

eliminating it, is essential in balancing livestock safety with plant resilience and performance

[30].

The present work addresses three main goals for advancing transformation and gene editing in

Sorghum bicolor: increasing efficiency of CRISPR/*Cas9* editing, expanding available PAM

recognition sites, and providing efficient genome editing capability. We developed a flexible

functional genomics toolbox for Agrobacterium-mediated sorghum transformation and genome

editing. The toolbox includes a binary T-DNA vector incorporating the *Wus2* gene, along with

genes for *hygromycin B phosphotransferase* (*hph*) and *ZsGreen* (*ZsG*) used for selection and as a

visual marker, respectively. For CRISPR/*Cas9* genome editing, we utilized an intronized Cas9

endonuclease in a shuttle vector, enabling easy insertion into the final binary vector of designed

gRNA sequences and recombination. Additionally, for the first time in sorghum, we demonstrate

editing with the nearly PAM-less Cas9 variant ZmSpRYi, employing both poly- and

monocistronic gRNA configurations to target edits in the promoter region of the *CYP79A1*

dhurrin biosynthesis gene. Overall, this high-efficiency toolbox increases the scalability of

genome editing in sorghum, enhancing future efforts for accelerated breeding and synthetic

biology.

Results

Design of high efficiency binary and shuttle vectors

Our two-step strategy for vector development involves gRNA validation in protoplasts, followed

by stable plant genetic transformation (**Figure 1A**). This advance in using protoplasts can

enhance efficiency and accessibility for researchers to perform successful sorghum

transformation and editing. The editing machinery is integrated in shuttle vectors, which enable

gRNA cloning with insertion of either a poly- or monocistronic configuration (**Figure 1B**). The

gRNA scaffold and Pol III promoters necessary for driving gRNA expression are supplied by the

Level 0 constructs (**Figure S1**). The Level 0 and shuttle constructs are sequentially assembled to

form complete gRNA configurations, which are transfected into sorghum protoplasts to evaluate

editing efficiency of the gRNAs [35,36].

To enhance efficiency of sorghum transformation, a series of binary constructs were developed,

all of which are compatible with Gateway LR recombination and tailored to allow incorporation

of genes of interest for multiple applications (**Figure 1C**). A hygromycin resistance gene (*hph*),

used as a selectable marker in the foundational construct, pGL206 (**Figure 1C**), was chosen

because of the stringent selection properties of hygromycin during *in vitro* tissue culture of

sorghum [37]. pGL206 was used as a construct for testing a candidate morphogenic gene, i.e.,

maize *Wuschel2* (*Wus2*). To facilitate tracking of somatic embryogenesis and plantlet

development, a visible marker gene *ZsGreen1,* driven by the sorghum ubiquitin promoter, was

incorporated near the T-DNA left border in pGL216. Building on previous findings of enhanced

editing efficiency achieved with integration of maize *Wus2* [11], pGL216 was assembled with

Wus2, driven by the maize phospholipid transfer protein promoter (pLTP). This resulted in

pGL222 (**Figure 1C**), a binary vector optimized for delivering genome-editing machinery.

Expression of ZsGreen validates presence of at least one complete T-DNA copy and permits

easy visualization of T-DNA segregation in T_1 generations. The constructs developed provide a

robust framework for improving transformation and genome-editing efficiencies in sorghum.

High transformation frequency achieved through *Wus2* integration

To evaluate transformation frequency in genome editing trials, we knocked out the *Sorghum*

bicolor phytoene desaturase (*SbPDS*) gene, resulting in easy visualization of an albino

phenotype [8]. Previous studies indicated editing efficiency may vary in wheat (*Triticum*

aestivum) and setaria (*Setaria viridis*) depending on the nature of the configurations of gRNAs in

C3 and C4 grasses, respectively [38,39]; however, this has not been thoroughly investigated in

sorghum using CRISPR/*Cas9*. To optimize editing efficiency, constructs were designed

containing two gRNAs under two transcriptional configurations, polycistronic and monocistronic

(**Figure 1B**). In the polycistronic construct, gRNAs were driven by the sorghum snoRNA U6.2.3

promoter [23]. In the monocistronic configuration, gRNAs were expressed using the snoRNA

U6.2.3 and snoRNA U6.3.1 promoters [23]. Both validated gRNA transcriptional configurations

were combined with a Cas9 expression cassette and integrated into the binary vector pGL222,

resulting in two editing constructs, pGL220 (polycistronic) and pGL221 (monocistronic) (**Figure**

S2). The constructs, containing the morphogenic gene, *Wuschel2* (*Wus2*), and the selectable

marker gene, *hph*, were used to assess the impact on editing efficiency of the two gRNA

transcriptional configurations.

Both pGL220 and 221 were transformed into sorghum immature embryos (IEs) using a modified

protocol [8]. Following Agrobacterium infection, IEs were maintained at 25°C for one week on

co-cultivation medium. In the current approach (**Figure 1D**), unlike *Bbm*/*Wus2* assisted

transformation, the *Wus2* expression cassette does not need to be excised because *Wus2*

integration does not lead to severely abnormal phenotypes. Previous studies in wheat indicated

that using higher temperatures at this stage enhanced mutagenesis efficiency when employing

CRISPR/*Cas9* [40]. To explore this in sorghum, tissues were incubated, starting at the resting

stage, at either 25°C or 28°C until plant regeneration began. Development of IEs was monitored

to evaluate responses to *Wus2* expression at 1-, 2-, and 4-week intervals post-infection (**Figures**

1E–H). Fluorescence was detected as early as three days after infection (**Figure 1E**) and peaked

at the end of the resting stage (**Figure 1F**). Somatic embryogenesis was noticeable at two weeks,

with regenerated plantlets forming by week four (**Figures 1G-H**). Prior to transferring to rooting

media, fluorescence microscopy was used to screen for ZsGreen expression, confirming

integration of full-length T-DNA, given that the ZsGreen expression cassette is located adjacent

to the left border.

Putative transgenic plantlets were analyzed via PCR to further confirm T-DNA integration and

determine transformation frequencies. Transformation frequencies and editing efficiencies were

calculated using two strategies. The first was to calculate frequency using only one transgenic

plant regenerated from each IE; the second used all regenerated transgenic plants from each IE.

Both approaches have advantages. Considering a single plant per IE is helpful when using an

overexpression construct in order to get independent transgenic events. Conversely, calculating

transformation frequency by considering all regenerated plantlets from each IE, as in the current

approach, is beneficial when performing genome editing because multiple independently edited

events can be regenerated from a single IE and may have unique edits.

For the monocistronic strategy at 28^{o}C, when considering only one plant per IE, we generated 50

transgenic plants from two replications, obtaining a transformation frequency of 24.4% (50/205)

(**Table 1**). For the polycistronic strategy an average transformation frequency was 26.9%

(81/301). It is important to note that issues incubated at 28^{o}C appeared stressed and showed more

necrosis than those at 25^{o}C (**Figure S3**). Because of these phenotypic differences, more plants

were regenerated from tissues kept at 25^{o}C, increasing transformation frequency for the

monocistronic strategy to 31.1% versus 24.4% for tissues kept at 28^{o}C. When all regenerated

transgenic plants were considered, transformation frequency substantially increased (**Table 2**).

At 25^{o}C, the monocistronic strategy yielded transformation frequency of 79.4% (143/180).

Additionally, at 28^{o}C, it led to notable differences in transformation frequency, 66.8% (137/205)

(**Table 2**). This decrease in frequency may be due to the fact that tissues kept at 28^{o}C showed

more necrosis than those kept at 25^{o}C (**Figure S3**), thereby reducing the number of regenerated

plants for the monocistronic strategy thus lowering transformation frequencies (**Table 1**).

Enhancing CRISPR/Cas9 editing efficiency using WUS2

Previous work demonstrated that various factors can be used to improve CRISPR/*Cas9* editing

efficiency. One factor is intronized and codon-optimized *Cas9* [21,22]. Using this *Cas9* to

improve genome editing might result from enhanced mRNA stability and translation [22].

Another possible factor is the nature of promoters driving gRNAs (**Figure 1B, 2A**), as they play

an important role in regulating expression of gRNAs [23,41]. Additionally incubating tissues at

higher temperatures might lead to improved editing efficiency [40,42]. Using a *Wus2*-integrated

transformation strategy resulted in improved CRISPR/*Cas9* gene knockout frequency, which

might be due to enhanced cell division and/or chromatin accessibility [11]. In our study, different

factors were used that could lead to improved genome editing and knockout efficiencies. The

approaches include using: (i) the *Wus2*-integrated transformation strategy, (ii) a codon-optimized

and intronized *Cas9*, (iii) monocistronic and/or polycistronic tRNA-gRNA expression cassettes,

(iv) the *SbU6* 2.3 and *SbU6* 3.1 promoters driving two gRNAs, and/or (v) two temperatures,

25^{o}C and 28^{o}C, during certain tissue culture steps.

To assess the impact of the various factors stated above, we focused on *SbPDS* due to the ability

to observe varied albino phenotypes **(Figure 2B).** This phenotype was used to visually identify

plants with *SbPDS* gene knockouts and to calculate knockout efficiencies. In addition, we

amplified the *SbPDS* gene fragments from transgenic plants and used Sanger sequencing to

investigate genome editing outcomes. Results were further analyzed using the Inference of

CRISPR Edits (ICE) Analysis Tool [43]. Editing outcomes for individual plants, as measured by

knockout scores from ICE, ranged from 0% to 100% (**Figure 2C**). Samples with knockout scores

greater than 5% were considered successfully edited. Genotypic editing efficiency (**Tables 1, 2**)

was calculated as the percentage of edited plants among all transformants for that condition,

hereafter this will be termed editing efficiency. Phenotypic knockout efficiency (**Tables 1, 2**)

was defined as the percentage of plants showing an albino phenotype among all transformants,

hereafter this will be termed knockout efficiency. The combined effects of the factors mentioned

in the preceding paragraph resulted in substantially improved average editing and knockout

efficiencies reaching 65.4% (17/26) for polycistronic and 96.4% (54/56) for monocistronic

configurations in tissues maintained at 25^{o}C (**Table1**), compared to efficiencies of 22.6% and

16.7%, respectively, in our previous report [8]. Incubation at 28^{o}C versus 25^{o}C did not yield

notable differences in editing and knockout efficiencies (**Table 1**). Interestingly, somatic

embryos cultured at 25^{o}C appeared less stressed under hygromycin selection, perhaps due to

reduced selection pressure relative to those at 28^{o}C, as indicated by less tissue browning (**Figure**

S3). Furthermore, analysis of knockout scores for individual gRNAs revealed that gRNA1.1 in

the polycistronic constructs exhibited a notably reduced knockout score (**Figure 2C**), possibly

due to suboptimal processing efficiency during tRNA-mediated autocleavage. Overall, our

results highlight the robustness of the monocistronic strategy, although the polycistronic design

can also deliver reasonably high editing efficiencies.

T_0 plants were phenotyped based on the protortion of albino sectors observed. Fully albino

plantlets were scored as 100%, whereas full green plants as 0% represented. These phenotypic

assessments were compared with the knockout scores gathered through the ICE analysis,

enabling effective identification and quantification of the degree of penetrance and frequencies

of mutations in plantlets. For most plants, phenotypic and genotypic scores were highly

consistent, with a few discrepancies, such as in pGL220-2, -20 & -34 (**Table S2**). Overall, the

two metrics showed a strong positive correlation, with a Pearson's correlation coefficient of 0.81.

This correlation underscores the value of the ICE analysis as a reliable and affordable tool for

estimating penetrance based on genotypic data. particularly in cases where gene knockouts do

not produce visually detectable phenotypes in the T_0 generation.

High-efficiency editing using PAM-less Cas9 variant, ZmSpRYi

Previous studies demonstrated that CRISPR/*Cas9*-mediated editing of promoters can generate

diverse *cis*-regulatory alleles, providing valuable quantitative variation for crop breeding

applications [44–46]. However, the standard CRISPR/*Cas9* system relies on canonical gRNAs

with NGG as PAM sequences, which are less likely in promoter or 5' untranslated regions. In

addition, CRISPR/*Cas9*-mediated edits in our study predominantly resulted in insertions or

deletions of less than 3 bp (**Figure 2A**). These small modifications also present a challenge when

targeting *cis*-regulatory elements, which can span long stretches of DNA. These regions can

include multiple coordinated DNA motifs, which may not be effectively disrupted by such small

edits [47]. Taken together, these observations indicate a limitation of CRISPR/Cas9 in editing

cis-regulatory modules.

To expand the genome editing scope and to identify options with a less constrained PAM

sequence, a range of nucleases were considered, including *Cas9* variants and *Cas9* orthologs

[16–20]. We focused on a PAM-flexible *Cas9* variant, *SpRY* [24], which enables more frequent

cis-regulatory element editing and expands genome engineering in AT-rich regions. To create a

maize codon-optimized *SpRY* variant, *ZmSpRYi,* targeted mutations were introduced in the

intronized *ZmCas9i* by site-directed mutagenesis (**Figure S4**). Previous studies suggested that

SpRY is a less efficient endonuclease due to modified DNA unwinding activity [48]. To evaluate

the editing efficiency of ZmSpRYi, we did a direct comparison between ZmSpRYi and

ZmCas9i, using the same gRNAs for both (**Figure 2A, D**). Two constructs, featuring

polycistronic and monocistronic configurations of gRNAs, were transformed into sorghum. To

assess editing efficiency of ZmSpRYi, all independent transformants were genotyped by PCR

amplification, followed by Sanger sequencing. Analysis of the sequencing data using ICE

revealed that, when using the polycistronic transcription system, 26.7% (16/60) of independent

transformants were edited whereas, strikingly when using the monocistronic system, 88.0% (81/

92) of transformants were edited (**Table 3, Figure 2D**). Interestingly, the gRNA1.1 failed to

induce any edits, suggesting that ZmSpRYi may exhibit distinct editing activity or property,

compared to ZmCas9i. In contrast, the knockout score distribution of gRNA2.2 in the

monocistronic configuration showed a robust editing efficiency (85.9%) versus the polycistronic

configuration (26.7%) (**Table 3**), indicating that gRNAs expressed from monocistronic

configurations are more efficient This is consistent with the findings when using CRISPR/Cas9-

mediated genome editing (**Table 1, Figure 2C**). To investigate the editing characteristics of

ZmSpRYi, we compared the editing profile of ZmCas9i and ZmSpRYi using amplicon

sequencing from stable edits. Analysis of editing outcomes for gRNA2.2 revealed that deletions

peaked at the third nucleotide upstream of the NGG site when both ZmCas9i and ZmSpRYi were

used (**Figure 2F**). ZmSpRYi, however, generated a larger deletion range, spanning from -8 to

+3, particularly with the gRNA monocistronic configuration (**Figure 2F**). Further, examination

of indel sizes showed that both nucleases predominantly produced 2-bp deletions (**Figure 2G**),

although deletions mediated by ZmSpRYi were slightly larger than those from ZmCas9i. These

findings highlight the enhanced editing efficiency of the monocistronic system in sorghum when

using either ZmCas9i or ZmSpRYi. Moreover, ZmSpRYi directed by NGG gRNAs may induce

a wider spectrum of deletions, making it a promising tool for disrupting cis-regulatory elements.

Editing PDS with ZmSpRYi using non-canonical gRNAs

Canonical gRNAs for Cas9-mediated genome editing require NGG (N = A, T, C, G) as a PAM

sequence, whereas non-canonical PAMs refer to all other sequences that can be recognized and

bound by SpRY. To assess activity of gRNAs with noncanonical PAMs in combination with

ZmSpRYi within AT-rich regions of the *PDS* promoter and exons, we designed six gRNAs with

NAA, NAT, NTA or NTT PAMs (**Figure 3A**). Three gRNAs were positioned 168 bp, 100 bp,

and 66 bp upstream of the transcription start site, aiming to disrupt translational regulation. The

remaining three targeted coding sequences, two in the first exon and one in the second exon,

were positioned to interfere with protein translation (**Figure 3A**). To optimize gRNA expression

in monocots, we employed a variety of U6 and U3 snoRNA promoters derived from various

grass species (**Figure S1**). All editing components, including ZmSpRYi and the six gRNAs,

were delivered via Agrobacterium-mediated transformation.

A previous study reported that the reduced PAM specificity of SpRY might compromise editing

efficiency, potentially due to decreased gRNA-target binding stability and inefficient R-loop

formation [49]. To evaluate the *in vivo* activity of ZmSpRYi, the T_0 regenerants described above

were genotyped. Of the 64 independent transformants analyzed, 31 exhibited mutations at one or

more gRNAs target sites, resulting in an overall editing efficiency of 48.4% (**Table 4**). Notably,

ZmSpRYi-induced edits frequently resulted in larger deletions within the promoter region

(**Figure S5**), reflecting a distinct mutational profile compared with ZmCas9i, which generally

favors shorter indels in sorghum. Interestingly, Spg2 and Spg3, separated by 44 bp, exhibited

markedly higher dropout frequencies (**Figure S5**). The tendency of ZmSpRYi to generate larger

deletions may be particularly advantageous for promoter editing, as complete disruption of *cis-*

regulatory elements often requires near-total removal to achieve functional inactivation [50].

Furthermore, gRNAs targeting the promoter region, specifically Spg1, Spg2, and Spg3, showed

higher editing efficiencies (**Table 4; Figure 3B, C**), potentially due to factors such as chromatin

accessibility, gRNA activity, or gRNA expression levels. Overall, ZmSpRYi in sorghum

achieved editing efficiencies comparable to those reported in other crops, such as rice [19,20].

Given its ability to introduce larger deletions and to target AT-rich regions with greater

flexibility, ZmSpRYi serves as a highly effective tool for promoter editing.

Cis-regulatory element editing via ZmSpRYi in *CYP79A1* promoter

To test the potential of ZmSpRYi for promoter editing, we focused on *CYP79A1*, a key gene in

the cyanogenic dhurrin biosynthesis pathway [51,52]. *CYP79A1* encodes a tyrosine N-

monooxygenase that catalyzes the dedicated step in dhurrin production [53]. While complete

knockout of this gene eliminates dhurrin toxicity, it also removes dhurrin's benefit as a nitrogen

reservoir and natural defense compound [54]. Modulation of *CYP79A1* expression via promoter

editing would offer an approach for optimizing the agronomic potential of sorghum for diverse

applications.

To explore the generation of diverse promoter edits, we examined sequence conservation of

CYP79A1 promoter orthologs across C4 grasses, including *Sorghum bicolor* subspecies

drummondii and *verticilliflorum*, as well as *Sorghum propinquum*, *Saccharum officinarum* ×

spontaneum, *Miscanthus sinensis*, and *Andropogon gerardii* (**Figure 3D**). Based on variation

within the core promoter region, 1 kb upstream of the translational start site, compared to

sorghum genotype RTx430, we designed four gRNAs targeting the *CYP79A1* promoter and 5′

untranslated region (UTR) at positions 600 bp, 383 bp, 212 bp, and 23 bp upstream of the

translational start site (**Figure 3D**). These sites were selected with the aim of disrupting putative

cis-regulatory elements (**Figure 3E**). ZmSpRYi and all four gRNAs, Cpg 1, 2, 3, and 4, were

transformed into sorghum. Out of 40 independent T☐ transformants, 19 were successfully edited

at least by one gRNA, resulting in an overall editing efficiency of 47.5% (**Table 4**). Interestingly,

the editing patterns varied among the four gRNAs. Of the four gRNAs, Cpg2 and 3 demonstrated

the highest editing efficiencies, at 30.0% and 47.5%, respectively. In contrast, Cpg1 had 22.5%

editing, while Cpg4 gave rise to 0% editing (**Table 4**). Our current findings demonstrate the

feasibility of using ZmSpRYi-mediated promoter editing to attempt developing varieties with

varying dhurrin levels.

Discussion

Development of efficient, targeted, and versatile genome-editing tools is needed to advance

genetic engineering and genome editing in bioenergy crops like sorghum. As a C4 plant,

sorghum is naturally efficient in photosynthesis but additionally is highly resilient to

environmental stresses, such as drought, flooding, and salinity, which could increase with

climate changes [3,55]. However, the molecular mechanisms underlying such key agronomic

traits and the ability to validate appropriate genetic resources to confirm functions remain

significant challenges. To address these challenges, while introducing desirable genetic

variations in sorghum, we developed a shuttle-binary vector toolbox to address current

limitations in genome editing. This system simplifies genetic engineering by incorporating into

the transformation construct the morphogenic gene *Wus2*, which accelerates more effectual

delivery of editing machinery [11]. Additionally, several optimizations were also introduced to

improve efficiency of CRISPR/Cas9-based genome editing in sorghum, including a codon-

optimized Cas9 containing an intron, high-efficiency endogenous U3/6 promoters, and

monocistronic gRNA expression cassettes. These modifications addressed previous limitations

and significantly expanded the potential applications of this genome editing technology in

sorghum.

Wuschel integration simplifies the binary vector and improves

transformation frequency

Previous studies demonstrated that the morphogenic regulators, BBM and WUS2, significantly

enhance transformation frequency in monocot species, including sorghum [8,9]. However,

continued expression of the morphogenic regulators often results in developmental aberrations

during the vegetative stage. To address this issue, the CRE/LoxP recombination system was

employed to excise the morphogenic genes, following somatic embryo initiation [9]. While

effective, this approach requires increasing the binary vector size, complicating molecular

cloning and construct development. Several approaches were used to overcome this challenge

[8], including reducing construct size and development of systems that combine transient and

stable *Wus2* expression [11,56]. In our approach, we redesigned the binary vector into a versatile

delivery platform by incorporating an LR recombination site, thereby ensuring compatibility

with the Gateway system, which is widely used for construct development among the plant

research community. When paired with a *Wus2* expression cassette, we used this system to

achieve transformation frequencies from 18.2% to 31.4% (**Table1 and 2**), rates comparable to

those attained using the BBM/WUS2-assisted transformation approach [8].

Despite this success, limitations persisted with integration-based delivery of *Wus2*.

Approximately 10% of the T_0 transformants exhibited developmental deficiencies, such as curly

leaves during the vegetative stage, potentially caused by ectopic expression of WUS2. To

address this issue, pGL216 can be used as a platform for testing alternative morphogenic

regulator genes, including *Wuschel* orthologs from other species or members of the *Wuschel*

homeobox gene family (**Figure 1C**). These alternative candidate genes may enhance somatic

embryo initiation while minimizing developmental disruptions, offering a more refined approach

to transformation and regeneration.

Optimizing *Cas9* and guide RNA expression cassettes increases editing

efficiency

In our previous work, we successfully achieved *PDS* knockouts in sorghum using CRISPR/*Cas9*,

with an editing efficiency of 16.7% [8]. However, enhancing genome-editing efficiency further

requires overcoming certain challenges, particularly ensuring robust expression of the editing

machinery, including the *Cas9* protein and gRNAs [14]. To address the editing challenges, we

used a constitutive promoter to drive expression of the intronized maize codon-optimized

nuclease genes, including *Cas9* and *SpRY*. Intronization was shown to enhance protein

abundance at both transcriptional and translational levels in Arabidopsis [22]. For gRNA

expression optimization, we utilized the sorghum endogenous snoRNA U6 promoters, which

lead to high gRNA expression levels [23].

Previous studies suggest that transcriptional configurations, including polycistronic and monocistronic approaches, may affect editing efficiency [38]. However, it remained unclear whether a polycistronic configuration compromises editing efficiency in sorghum. To investigate this, we conducted a side-by-side comparison of polycistronic and monocistronic gRNA transcriptional configurations using Cas9 and SpRY nucleases. Results indicate that the monocistronic gRNA configuration led to significantly higher editing efficiency, up to 97.6% with Cas9 and 88.0% with SpRY, compared to 61.7% and 26.6% respectively observed with the polycistronic configurations (**Tables 1 and 3**). Additionally, gRNA1.1 exhibited lower editing efficiency in the polycistronic configuration compared to the monocistronic approach, suggesting that the temporal processing of the first exogenous tRNA in the polycistronic construct may be suboptimal in sorghum (**Figure 2D**). We also observed, when using the polycistronic constructs, a delayed onset of albino phenotypes resulting from *PDS* knockouts. This delay further supports the notion that transcriptional processing inefficiencies might contribute to the reduced editing efficiency observed with polycistronic gRNA configuration in sorghum. To improve editing outcomes, we optimized the monocistronic configuration for gRNA expression using a collection of monocot-derived snoRNA promoters. This collection includes U6 promoters from sorghum, wheat, and maize, as well as U3 promoters from wheat and rice (**Figure S1**), all of which have been experimentally validated in sorghum (**Figure 3B, Figure S5**). Promoters were incorporated into Level 0 constructs, providing flexible tools for expressing multiple gRNAs independently. This modular approach facilitates multiplex genome editing and is particularly well-suited for applications such as promoter editing and other complex genetic modifications.

To evaluate the impact of temperature on editing efficiency, we conducted an editing experiment

using different incubation temperatures, 25°C and 28°C, following Agrobacterium infection.

Interestingly, no apparent difference in editing efficiency was observed between the two

temperatures (**Table 1**). This outcome may be attributed to the high baseline editing efficiency

achieved through the modifications introduce into both the Cas9 and gRNA components.

Elevated temperatures did not confer further improvements, probably because the optimized

CRISPR/Cas9 constructs were already near their maximal efficiency. In contrast, tissues

incubated at 28°C exhibited significant necrosis; whereas those maintained at 25°C remained

healthier and produced more regenerants on average (**Figure S3**). Therefore, temperatures above

28°C were excluded from further experimentation. Our findings suggest that 25°C is a more

suitable incubation temperature for achieving efficient genome editing in sorghum.

431 Expanding the scope of genome editing using ZmSpRYi

One potentially impactful application of genome editing in crops is creating cis-regulatory

variations to diversify genetic resources for improving desirable traits [44]. Editing of cis-

regulatory elements in promoters and UTRs, which are typically AT-rich, is often constrained

when using Cas9, due to its strict NGG PAM requirement and in sorghum its tendency to

produce small deletions. To address the need for AT sequences, we considered several potential

genome editing candidates to target AT-rich regions, including Cas9 variants (Cas9-NG, xCas9,

and SpRY) and other Cas orthologs (Cas12a and Cas12b), which exhibit reduced dependency on

CG content at the PAM site [24,57–60]. Among these editing candidates, we chose SpRY which

has the highest flexibility due to its PAM-less feature [24].

Using the same gRNAs that targeted the PDS gene in our previous CRISPR/*Cas9*-mediated

editing experiments [8], in the present study Zm*SpRY*i achieved an editing efficiency of 88.0%

(**Table 3**). However, previous studies suggested that use of *SpRY* results in compromised double-

strand break activity, particularly with non-NGG PAM sequences [19,20]. To further investigate

ZmSpRYi activity with non-NGG PAM sequences, we tested its performance in combination

with WUS2. Previous research indicated that integration of *Wus2* can enhance CRISPR/*Cas9*-

mediated gene dropout frequency [11]. However, it remained unclear as to whether *Wus2*

integration can similarly enhance SpRY activity. To address this question, we designed six

gRNAs with NAA, NAT, NTA, and NTT PAM sequences (**Figure 3A**). Across 64 independent

transformants, an overall editing efficiency of 48.4% was observed (**Table 4**), which is

comparable to the editing efficiency reported in rice [19,20]. Interestingly, regardless of the

PAM sequences, gRNAs targeting promoter regions (48.4%, 46.9%, and 37.5%) exhibited

higher editing efficiencies than those targeting coding regions (21.9%, 26.6%, 32.8%) (**Table 4**),

probably due to differences in chromosome accessibility among these genomic regions.

Another concern for SpRY is the potential for off-target effects, including self-editing, which

refers to mutagenesis occurring within the gRNA expression cassette [19,20], possibly leading to

unintended edits. To assess this risk in edits with high knockout scores, we amplified and

sequenced potential off-target sites, including regions within the T-DNA. We found that 6.7%

(1/15) of plantlets from polycistronic configurations targeting PDS gRNAs with NGG PAMs

showed self-editing, whereas monocistronic configurations exhibited self-editing frequencies of

up to 33.3% (6/18) within the T-DNA (**Figure S6**). Moreover, gRNAs with non-canonical PAMs

showed 66.6% self-editing frequency on at least one gRNA, probably due to the increased

number of gRNAs, up to six in the construct (**Figure S6**). The target regions were also

successfully edited in these cases, suggesting that self-editing occurred either concurrently or

sequentially with the targeted editing process. Taken together the present study highlights the

promise of *SpRY* for expanding the genome editing toolbox in AT-rich regions, while

acknowledging challenges like compromised activity and self-editing risks.

Potential of ZmSpRYi for efficient promoter editing

Results of the present study demonstrates the potential use of SpRY, a PAM-flexible Cas9

variant, as a tool for promoter editing in plants. For example, by editing the CYP79A1 promoter,

it should be possible to generate quantitative variations to limit dhurrin production to increase the

safety of sorghum as livestock feed. Toward this end, the SpRY construct achieved a moderate

overall editing efficiency of 47.5%, with over half of the edited plants showing a knockout score

exceeding 50% at the second and third gRNA sites (**Table 3**). These editing results suggest

SpRY is a viable tool for editing AT-rich regulatory regions, where conventional Cas9 is

restricted by stringent PAM requirements. Furthermore, combining adjacent gRNAs, such as

Spg2 and Spg, enables the targeting of potential cis-regulatory elements and may promote larger

dropouts, due to the PAM flexibility of SpRY.

Despite its promise, SpRY remains under-optimized. The moderate editing rate suggests that

SpRY's PAM flexibility may compromise target engagement, possibly due to reduced DNA

binding affinity or inefficient R-loop formation [49]. Artificial intelligence-assisted optimization

represents a strategy to enhance the performance of PAM-less Cas9 variants, like SpRY, and

further alleviate PAM-related constraints [61]. While SpRY expands the range of targetable

sequences, gRNA performance was inconsistent. Although *in silico* predictions using certain algorithms indicated that all four gRNAs should be highly efficient, only the second and third showed moderate activity, while the fourth was non-functional. This discrepancy underscores the limitations of current gRNA efficiency prediction tools, which fail to consider biological variables, such as chromatin accessibility especially near transcription start sites [62]. Refining gRNA design algorithms that incorporate epigenomic features and consider the activity of engineered editors, for example SpRY, might improve future targeting accuracy and efficiency [63].

Concluding remarks

In the current study, we described high-efficiency transformation and editing approaches incorporating *wus2* integration with a robust Cas9-based genome editing toolkit, featuring intronized ZmCas9 and ZmSpRY. This platform is a versatile means for delivering genome editing machinery that can include base editors and prime editors. By integrating *wus2* with the Cas9 variant, SpRY, we expanded the editing scope to increase other desirable applications, such as promoter editing. These advancements provide a streamlined approach to generating diverse genetic variation in sorghum, facilitating crop improvement efforts to address food security challenges and contribute to carbon capture initiatives that address climate change.

Materials and Methods

Plant material and growth conditions

507   Seeds of *Sorghum bicolor* genotype, RTx430, were sourced from the Germplasm Resources

508   Information Network (GRIN, USDA-ARS, PI 655998). Plants were cultivated in three-gallon

509   pots containing SuperSoil (Rod Mclellan Company) in UC Berkeley's Oxford Research Unit

510   South Greenhouse, under controlled conditions of 30°C/24°C (day/night) with a 16-hour light/8-

511   hour dark photoperiod. Immature embryos (IEs) were harvested 12–14d post-anthesis.

Construct creation

To create pGL206 (**Figure 1C**), the *hygromycin B phosphotransferase* gene (*hph*), driven by the

Panicum virgatum Ubiquitin2 promoter (PvUbi2p), was amplified from pANIC10A [64] using

the primer SorEd1 and 2 (**Table S1**). All primers used for construct building are described in

Table S1. The terminator for sorghum PEPC was amplified from pPHP85425 [8] using the

primers SorEd3 and 4. These two parts were assembled by overlapping PCR. The vector

backbone of pPHP85425 was amplified using the primers SorEd5 and 6. In addition, the Buffer3

and attR4-attR3 fragments were amplified from pPHP85425 using the primers SorEd7 and 8.

These three fragments were assembled using Gibson assembly (NEBuilder® HiFi DNA

Assembly Master Mix). To make the construct pGL216 (**Figure 1C**), the *ZsG* expression

cassette was amplified from pGL190 [8] using the primer SorEd9 and 10. The amplified cassette

was then inserted into pGL206 through *Sfi*I digestion and subsequent ligation. To create pGL222

(**Figure 1C**), the partial backbone of pPHP96564 [11] and the *Wus2* expression cassette were

amplified from pPHP96564 using the primers SorEd11 and 12. The attR4-attR3 region was

amplified from pPHP85425 using the primers SorEd13 and 14. The *hygromycin B*

phosphotransferase (*hph*) gene was amplified from pGL206 using the primers SorEd15 and 16.

The rest of the backbone of pPHP96564 was amplified with the primers SorEd17 and 18. These

fragments were reassembled into the final vector using Gibson assembly. Additionally, the *ZsG*

gene, driven by the *Sorghum bicolor ubiquitin1* promoter, was amplified from pGL190 with the

primers SorEd19 and 20. The ZsG expression cassette was digested with *Nhe*I and *Pme*I and

ligated into the construct to finalize the construct, pGL222, validated using Nanopore

sequencing.

To optimize shuttle constructs, the sorghum U6.2.3 (SbU6.2.3) promoter [23] from sorghum

genotype RTx430 was amplified using the primers SorEd21 and 22. pGL198 [8] was linearized

with primers, SorEd23 and 24. The OsU3 promoter was replaced by SbU6.2.3 using Gibson

assembly. Additionally, ZmCas9 was replaced by ZmCas9i using primers SorEd25 and 26 for

vector linearization and primers SorEd27 and 28 for ZmCas9i amplification from pPHP96564,

resulting in the shuttle vector, pGL220. Based on the polycistronic shuttle vector construction,

the monocistronic shuttle vector was created by inserting two gRNA expression cassettes by

Gibson assembly with primers SorEd29-36 to create the monocistronic expression cassettes.

Cas9 and gRNA polycistronic and monocistronic cassettes were transferred to pGL222 to create

binary constructs pGL220 and 221 by LR recombination. To create pGL224, we used the same

strategy as described above to replace the OsU3 promoter and ZmCas9 with the SbU6.2.3

promoter and ZmCas9i. To streamline construct creation, level 0 constructs were developed.

These constructs were created by fusing the gRNA scaffold with various snoRNA promoters

(including sorghum U6_2.3 [SbU6_2.3], sorghum U6_3.1 [SbU6_3.1] [23], rice U3 [OsU3] [65],

maize U6 [ZmU6] [66], wheat U3 [TaU3] [67], and wheat U6 [TaU6]) [68] using overlapping

PCR with primers SorEd49 through 62. Fused DNA fragments were cloned into a pJET1.2

vector using CloneJET PCR cloning kit (ThermoFisher Scientific) and validated by Sanger

sequencing. These level 0 constructs served as templates for creating both polycistronic and

monocistronic gRNA editing constructs (**Figure S2**).

To create the ZmSpRYi construct, pGL224 was used as the template, and a total of 14 nucleotide

mutations were introduced. The first set of mutations on the first exon of ZmCas9i targeted the

removal of a BsaI recognition site and the substitution of alanine 61 with arginine. These

nucleotide mutations were incorporated using primers SorEd37 and SorEd38, enabling the

amplification of two overlapping fragments with primers SorEd39 and SorEd40. The

overlapping fragments were assembled through overlapping PCR. The modified sequence

replaced the original sequence via BamHI and BspEI digestion, followed by ligation. The

remaining ten mutations were introduced using a synthetic oligonucleotide chemically

synthesized by Genewiz (http://genewiz.com). All these mutations were inserted into the 3′ end

of ZmCas9i through Gibson assembly. Primers SorEd41 and SorEd42 were used to amplify the

insert, while primers SorEd43 and SorEd44 were employed for vector linearization.

Additionally, two unwanted BsaI restriction sites were eliminated via primer-directed

mutagenesis using primers SorEd45 through SorEd48, resulting in the construct, pGL225

(**Figure S1**). Designed for Agrobacterium-mediated transformation, the ZmCas9i gene in

pGL220 and pGL221 was replaced with the ZmSpRYi gene to create pGL226 and pGL227. To

generate the pGL236 construct targeting the *PDS* promoter and exons, gRNAs were designed

using the CRISPOR tool [69]. For gRNAs ending with NAN and NTN motifs, we amplified both

the gRNA scaffolds and the snoRNA promoter from level 0 constructs using primers that

incorporated the guide sequences. These primers also served as adaptors for overlapping PCR.

The primers used for assembling the six gRNA expression cassettes are listed as SorEd63-72.

The assembled PCR product was digested with *Bsa*I and ligated into the *Bsa*I-digested vector

pGL225. The integrity of the construct was validated via Nanopore sequencing and subsequently

recombined into the binary vector pGL222 through LR recombination, resulting in the final

construct, pGL236, to be used for Agrobacterium-mediated transformation.

For the pGL237 construct targeting the *CYP79A1* promoter, gRNAs were designed based on

conserved promoter and untranslated regions identified across homologs from the sorghum

genus, *Miscanthus sinensis*, *Saccharum officinarum*, and *Andropogon gerardi* (**Figure 3D**). The

CRISPOR tool was used to design gRNAs with the goal of disrupting transcription by targeting

predicted cis-regulatory elements. Considering the distribution of cis-elements and optimal

gRNA spacing, we selected gRNAs spaced approximately 200 bp apart within the promoter

region (**Figure 3D**). The gRNA expression cassettes were amplified using primers SorEd73-78,

assembled via overlapping PCR and ligated into pGL225, as previously described. The construct

was verified with Nanopore sequencing and finalized by LR recombination to generate pGL237

for further Agrobacterium-mediated transformation.

Agrobacterium-mediated transformation

The binary constructs were introduced into an auxotrophic strain of *Agrobacterium tumefaciens*

LBA4404 Thy- (Anand et al. 2017) by electroporation. Transformed Agrobacterium strains were

plated on yeast extract peptone (YEP) medium supplemented with 100 mg/L thymidine, 50 mg/L

gentamicin, and 50 mg/L spectinomycin. Details on the storage and preparation of the

Agrobacterium strains followed a previously published protocol [70]. In summary,

Agrobacterium stocks were streaked onto YEP plates containing appropriate antibiotics to

generate master plates. After a 2-day incubation at 28°C, 5 to 6 colonies from the master plates

were selected to establish working cultures on YEP plates. Following overnight incubation at

28°C and just prior to inoculation of sorghum immature embryos, Agrobacterium suspensions

were prepared in PHI-I medium [71], with optical density (OD550 nm) adjusted to 0.7; silwet L-

77 (0.005% v/v) and acetosyringone (0.2 mM) were added to finalize the Agrobacterium

inoculation medium. IEs were inoculated with Agrobacterium suspension culture for 5 minutes,

then placed on co-cultivation medium at 25 °C for one week. After co-cultivation, the IEs were

transferred to resting medium for an additional week and then moved to either 25 °C or 28 °C.

Selection followed on embryo maturation medium containing 15 mg/L hygromycin until

regeneration commenced. Regenerated plantlets at 1–2 cm in height were then transferred to

rooting medium until fully rooted (**Figure 1C**). Transformation frequency was calculated as the

percentage of IEs that successfully produced transformants.

608 Visualization of T-DNA integration

The ZsGreen reporter gene was used as a visible marker to track T-DNA integration and

segregation. To monitor T-DNA integration following Agrobacterium-mediated inoculation,

transformed IEs were examined weekly using a Lumar v12 epifluorescence stereo microscope

and a QImaging Retiga-SRV camera (Zeiss, UC Berkeley Biological Imaging Facility).

Response of IEs to WUS expression was assessed by analyzing the number and size of

fluorescent tissues. Prior to transferring shoots to rooting medium, regenerants exhibiting

fluorescence were selected for further genotyping and phenotyping.

616 Genotyping of editing events

To test putative T_0 transgenic lines for successful editing of target genes, genomic DNA from

young sorghum leaves was isolated using a urea buffer as described [8]. For the target regions in

putative PDS edits due to gRNA1.1 and 2.2, primers SorEd79 and 80 were employed. To

genotype editing by non-canonical gRNAs on the PDS gene, primers SorEd81 and 82 were used.

For genotyping CYP79a1 cis-regulatory elements, primers SorEd83 and 84 were used for PCR

amplification. Amplicons were amplified and then purified with the GeneJet PCR Purification

Kit (Thermo Scientific). Purified PCR products were Sanger-sequenced at the UC Berkeley

DNA Sequencing Facility, using the primers for PCR amplification. Sequencing data were

analyzed with the Inference of CRISPR Edits (ICE) tool [43], and knockout scores, referring to

proportion of indels that indicate a frameshift or are more than 21 bp in length, were used to

assess editing efficiency and the likelihood of transmission to the next generation. Editing

efficiency represents the percentage of plant cells within harvested somatic samples that

successfully undergo knockout events, as determined by the knockout score.

Phenotypic characterization of T_0 PDS plants

T_0 plantlets were phenotyped while growing in rooting media. Each plantlet was scored for the

percentage of albino phenotype, with 100% indicating plants that were completely albino and

0% indicating plants that were completely green. The phenotype score was compared with the

knockout percentage obtained using the ICE tool to determine correlation between the two

observations.

Resource availability

Lead contact

Requests for further information and resources should be directed to the lead contact, Peggy G.

Lemaux (lemauxpg@berkeley.edu)

Material availability

Data availability

646   All data generated in this work are reported in this manuscript. Any additional information will

647   be made available from the lead contact upon reasonable request.

Acknowledgments

This project was supported by the Chan Zuckerberg Initiative through the Innovative Genomics

Institute and by the Terraforming Soils EERC through Lawrance Livermore National Laboratory

under Award #SCW1841. PGL acknowledges support from the US Cooperative Extension

Service through the Division of Agriculture and Natural Resources of the University of

California. DFS is an investigator in the Howard Hughes Medical Institute. JS acknowledges

support from USDA-ARS Crop Improvement and Genetics Unit Albany, CA and ORISE

Research Participation Program. We thank Dr. William Gordon-Kamm at Corteva Agriscience

for helpful conversations and for the constructs, pPHP96564 and pPHP85425. We thank Dr.

Roger Thilmony at USDA-ARS Crop Improvement and Genetics Unit Albany for informative

discussions. We thank talented undergraduate students, Malia Ariyoshi, Amada Niemela, Emma

To, Nessa Kmetec, Isabella Lombardo, Dasha Wohlfarth, and Sanay Parikh, for their help in this

work. We are grateful for the help of the staff of the UC Berkeley Biological Imaging Facility

and the UC Berkeley Oxford Tract Greenhouse Facility.

Author contributions

663   Conceptualization: J.S., K.A., E.G., D.S. and P.G.L.. Funding acquisition: D.S. and P.G.L..

664   Methodology: J.S., K.A., and S.A.. Investigation: J.S., K.A., S.A., and M.R.. Formal analysis:

665   J.S. and K.A.. Visualization: J.S., M.R., and K.A. Writing—original draft: J.S., K.A., and T.M..

666   Writing—review and editing: J.S., K.A., S.A., T.M., and P.G.L.. Supervision: J.S., D.S. and

667   P.G.L.

Declaration of interest

669   Manuscript authors have no relationship with Corteva Agriscience. The authors declare that they

670   have no competing interests.

References

1.   Dahlberg, J.A. *et al.* (2004) Development of a sorghum core collection: Refinement and

evaluation of a subset from Sudan. *Econ. Bot.* 58, 556–567

2.   Paterson, A.H. *et al.* (2009) The Sorghum bicolor genome and the diversification of grasses.

Nature 457, 551–556

3.   Varoquaux, N. *et al.* (2019) Transcriptomic analysis of field-droughted sorghum from

seedling to maturity reveals biotic and metabolic responses. *Proc. Natl. Acad. Sci. U. S. A.*

116, 27124–27132

4. Ramstein, G.P. *et al.* (2019) Breaking the curse of dimensionality to identify causal variants

in Breeding 4. *Züchter Genet. Breed. Res.* 132, 559–567

5. Schindele, A. *et al.* (2020) CRISPR/Cas brings plant biology and breeding into the fast lane.

Curr. Opin. Biotechnol. 61, 7–14

6. Zhu, H. *et al.* (2020) Applications of CRISPR-Cas in agriculture and plant biotechnology.

Nat. Rev. Mol. Cell Biol. 21, 661–677

7. Altpeter, F. *et al.* (2016) Advancing crop transformation in the era of genome editing. *Plant*

Cell 28, 1510-1520

8. Aregawi, K. *et al.* (2022) Morphogene-assisted transformation of Sorghum bicolor allows

more efficient genome editing. *Plant Biotechnol. J.* 20, 748–760

9. Lowe, K. *et al.* (2016) Morphogenic regulators Baby boom and wuschel improve monocot

transformation. *Plant Cell* 28, 1998–2015

10. Lowe, K. *et al.* (2018) Rapid genotype "independent" Zea mays L. (maize) transformation

via direct somatic embryogenesis. *In Vitro Cell. Dev. Biol. Plant* 54, 240–252

11. Che, P. *et al.* (2022) Wuschel2 enables highly efficient CRISPR/Cas-targeted genome

editing during rapid de novo shoot regeneration in sorghum. *Commun. Biol.* 5, 344

12. Jinek, M. *et al.* (2012) A programmable dual-RNA-guided DNA endonuclease in adaptive

bacterial immunity. *Science* 337, 816–821

13. Zhang, Y. *et al.* (2019) The emerging and uncultivated potential of CRISPR technology in

plant science. *Nat. Plants* 5, 778–794

14. Liu, G. *et al.* (2019) Genome editing by CRISPR/Cas9 in sorghum through biolistic

bombardment. *Methods Mol. Biol.* 1931, 169–183

15. Endo, M. *et al.* (2019) Genome editing in plants by engineered CRISPR-Cas9 recognizing

703   NG PAM. *Nat. Plants* 5, 14–17

704 16. Ren, B. *et al.* (2019) Cas9-NG greatly expands the targeting scope of the genome-editing

705   toolkit by recognizing NG and other atypical PAMs in rice. *Mol. Plant* 12, 1015–1026

706 17. Zhong, Z. *et al.* (2019) Improving plant genome editing with high-fidelity xCas9 and non-

707   canonical PAM-targeting Cas9-NG. *Mol. Plant* 12, 1027–1036

708 18. Ge, Z. *et al.* (2019) Engineered xCas9 and SpCas9-NG variants broaden PAM recognition

709   sites to generate mutations in Arabidopsis plants. *Plant Biotechnol. J.* 17, 1865–1867

710 19. Ren, Q. *et al.* (2021) PAM-less plant genome editing using a CRISPR-SpRY toolbox. *Nat.*

711   *Plants* 7, 25–33

712 20. Xu, Z. *et al.* (2021) SpRY greatly expands the genome editing scope in rice with highly

713   flexible PAM recognition. *Genome Biol.* 22, 6

714 21. Castel, B. *et al.* (2019) Optimization of T-DNA architecture for Cas9-mediated mutagenesis

715   in Arabidopsis. *PLoS One* 14, e0204778

716 22. Grützner, R. *et al.* (2021) High-efficiency genome editing in plants mediated by a Cas9 gene

717   containing multiple introns. *Plant Commun.* 2, 100135

718 23. Massel, K. *et al.* (2022) Endogenous U6 promoters improve CRISPR/Cas9 editing

719   efficiencies in Sorghum bicolor and show potential for applications in other cereals. *Plant*

720   *Cell Rep.* 41, 489–492

721 24. Walton, R.T. *et al.* (2020) Unconstrained genome targeting with near-PAMless engineered

722   CRISPR-Cas9 variants. *Science* 368, 290–296

723 25. Hendelman, A. *et al.* (2021) Conserved pleiotropy of an ancient plant homeobox gene

724   uncovered by cis-regulatory dissection. *Cell* 184, 1724-1739.e16

725 26. Tao, Y. *et al.* (2021) Extensive variation within the pan-genome of cultivated and wild

sorghum. *Nat. Plants* 7, 766–773

27. Tang, X. *et al.* (2017) A CRISPR–Cpf1 system for efficient genome editing and

transcriptional repression in plants. *Nat. Plants* 3, 17018

28. Ming, M. *et al.* (2020) CRISPR-Cas12b enables efficient plant genome engineering. *Nat.*

Plants 6, 202–208

29. Wattenbarger, D.W. *et al.* (1968) Effects of frost and freezing on hydrocyanic acid potential

of sorghum plants[1]. *Crop Sci.* 8, 526–528

30. Busk, P.K. and Møller, B.L. (2002) Dhurrin synthesis in sorghum is regulated at the

transcriptional level and induced by nitrogen fertilization in older plants. *Plant Physiol.* 129,

1222–1231

31. O'Donnell, N.H. *et al.* (2013) Effects of PEG-induced osmotic stress on growth and dhurrin

levels of forage sorghum. *Plant Physiol. Biochem.* 73, 83–92

32. Giantin, S. *et al.* (2024) Overview of Cyanide Poisoning in Cattle from Sorghum halepense

and S. bicolor Cultivars in Northwest Italy. *Animals (Basel)* 14, 743

33. Adewusi, S.R. (1990) Turnover of dhurrin in green sorghum seedlings. *Plant Physiol.* 94,

1219–1224

34. Jenrich, R. *et al.* (2007) Evolution of heteromeric nitrilase complexes in Poaceae with new

functions in nitrile metabolism. *Proc. Natl. Acad. Sci. U. S. A.* 104, 18848–18853

35. Jores, T. *et al.* (2021) Synthetic promoter designs enabled by a comprehensive analysis of

plant core promoters. *Nat. Plants* 7, 842–855

36. Lee, J.S. *et al.* (2023) A streamlined guide RNA screening system for genome editing in

Sorghum bicolor. *Plant Methods* 19, 90

37. van den Elzen, P.J. *et al.* (1985) A chimaeric hygromycin resistance gene as a selectable

749  marker in plant cells. *Plant Mol. Biol.* 5, 299–302

750 38. Liang, Z. *et al.* (2021) Efficient genome editing in Setaria italica using CRISPR/Cas9 and

751  base editors. *Front. Plant Sci.* 12, 815946

752 39. Milner, M.J. *et al.* (2024) Differential editing efficiencies in cereal crops: a comparative

753  analysis of tRNA and ribozyme multiplexed guide delivery. *Front. Plant Sci.* 15, 1426184

754 40. Milner, M.J. *et al.* (2020) Turning up the temperature on CRISPR: Increased temperature

755  can improve the editing efficiency of wheat using CRISPR/Cas9. *Front. Plant Sci.* 11,

756  583374

757 41. Zhan, X. *et al.* (2021) Genome editing for plant research and crop improvement. *J. Integr.*

758  *Plant Biol.* 63, 3–33

759 42. LeBlanc, C. *et al.* (2018) Increased efficiency of targeted mutagenesis by CRISPR/Cas9 in

760  plants using heat stress. *Plant J.* 93, 377–386

761 43. Conant, D. *et al.* (2022) Inference of CRISPR edits from Sanger trace data. *CRISPR J.* 5,

762  123–130

763 44. Rodríguez-Leal, D. *et al.* (2017) Engineering quantitative trait variation for crop

764  improvement by genome editing. *Cell* 171, 470-480.e8

765 45. Zhou, J. *et al.* (2023) An efficient CRISPR-Cas12a promoter editing system for crop

766  improvement. *Nat. Plants* 9, 588–604

767 46. Patel-Tupper, D. *et al.* (2024) Multiplexed CRISPR-Cas9 mutagenesis of rice PSBS1

768  noncoding sequences for transgene-free overexpression. *Sci. Adv.* 10, eadm7452

769 47. Yocca, A.E. and Edger, P.P. (2022) Current status and future perspectives on the evolution

770  of cis-regulatory elements in plants. *Curr. Opin. Plant Biol.* 65, 102139

771 48. Hibshman, G.N. *et al.* (2024) Unraveling the mechanisms of PAMless DNA interrogation

by SpRY-Cas9. *Nat. Commun.* 15, 3663

49. Shi, H. *et al.* (2025) Rapid two-step target capture ensures efficient CRISPR-Cas9-guided

genome editing. *Mol. Cell* 85, 1730-1742.e9

50. Schmitz, R.J. *et al.* (2022) Cis-regulatory sequences in plants: Their importance, discovery,

and future challenges. *Plant Cell* 34, 718–741

51. Sibbesen, O. *et al.* (1994) Isolation of the heme-thiolate enzyme cytochrome P-450TYR,

which catalyzes the committed step in the biosynthesis of the cyanogenic glucoside dhurrin

in Sorghum bicolor (L.) Moench. *Proc. Natl. Acad. Sci. U. S. A.* 91, 9740–9744

52. Laursen, T. *et al.* (2016) Characterization of a dynamic metabolon producing the defense

compound dhurrin in sorghum. *Science* 354, 890–893

53. Laursen, T. *et al.* (2015) Plasticity of specialized metabolism as mediated by dynamic

metabolons. *Trends Plant Sci.* 20, 20–32

54. Blomstedt, C.K. *et al.* (2012) A combined biochemical screen and TILLING approach

identifies mutations in Sorghum bicolor L. Moench resulting in acyanogenic forage

production: Acyanogenic forage sorghum plants. *Plant Biotechnol. J.* 10, 54–66

55. Zhang, H. *et al.* (2023) A Gγ protein regulates alkaline sensitivity in crops. *Science* 379,

eade8416

56. Hoerster, G. *et al.* (2020) Use of non-integrating Zm-Wus2 vectors to enhance maize

transformation: Non-integrating WUS2 enhances transformation. *In Vitro Cell. Dev. Biol.*

Plant 56, 265–279

57. Zetsche, B. *et al.* (2015) Cpf1 is a single RNA-guided endonuclease of a class 2 CRISPR-

Cas system. *Cell* 163, 759–771

58. Nishimasu, H. *et al.* (2018) Engineered CRISPR-Cas9 nuclease with expanded targeting

space. *Science* 361, 1259–1262

59. Hu, J.H. *et al.* (2018) Evolved Cas9 variants with broad PAM compatibility and high DNA

specificity. *Nature* 556, 57–63

60. Strecker, J. *et al.* (2019) Engineering of CRISPR-Cas12b for human genome editing. *Nat.*

Commun. 10, 212

61. Ruffolo, J.A. *et al.* (2025) Design of highly functional genome editors by modelling

CRISPR-Cas sequences. *Nature* 645,518-525

62. Zhou, C. *et al.* (2021) Accessible chromatin regions and their functional interrelations with

gene transcription and epigenetic modifications in sorghum genome. *Plant Commun.* 2,

100140

63. Yuan, H. *et al.* (2025) An overview and comparative analysis of CRISPR-SpCas9 gRNA

activity prediction tools. *CRISPR J.* 8, 89–104

64. Mann, D.G.J. *et al.* (2012) Gateway-compatible vectors for high-throughput gene functional

analysis in switchgrass (Panicum virgatum L.) and other monocot species. *Plant Biotechnol.*

J. 10, 226–236

65. Xie, K. *et al.* (2015) Boosting CRISPR/Cas9 multiplex editing capability with the

endogenous tRNA-processing system. *Proc. Natl. Acad. Sci. U. S. A.* 112, 3570–3575

66. Qi, W. *et al.* (2016) High-efficiency CRISPR/Cas9 multiplex gene editing using the glycine

tRNA-processing system-based strategy in maize. *BMC Biotechnol.* 16, 58

67. Hahn, F. *et al.* (2020) A modular cloning toolkit for genome editing in plants. *BMC Plant*

Biol. 20

68. Debernardi, J.M. *et al.* (2020) A GRF–GIF chimeric protein improves the regeneration

efficiency of transgenic plants. *Nat. Biotechnol.* 38, 1274–1279

69. Concordet, J.-P. and Haeussler, M. (2018) CRISPOR: intuitive guide selection for

CRISPR/Cas9 genome editing experiments and screens. *Nucleic Acids Res.* 46, W242–

W245

70. Masters, A. *et al.* (2020) Agrobacterium-mediated immature embryo transformation of

recalcitrant maize inbred lines using morphogenic genes. *J. Vis. Exp.* 156, e60782

71. Wu, E. *et al.* (2014) Optimized Agrobacterium-mediated sorghum transformation protocol

and molecular data of transgenic sorghum plants. *In Vitro Cell. Dev. Biol. Plant* 50, 9–18

(A)

gRNAs   Shuttle vector

gRNA validation
in protoplasts

Argobacterium
transformation

Binary vector

(B)

Polycistronic shuttle vector

attL4   ZmUbi1p   ZmCas9i   SbU6.2.3   attL3

1,000 bp

tRNAs   gRNAs

Monocistronic shuttle vector

attL4   ZmUbi1p   ZmCas9i   attL3

1,000 bp

PolIII promoters   gRNAs

(C)

pGL206

RB   attR4   attR3   LB
Buf   *ccdB*   PvUbi2p   *Hph*

500 bp   SbPEPC1t

pGL216

RB   attR4   attR3   LB
Buf   *ccdB*   PvUbi2p   *Hph*   SbUbip   *ZsG*   OsUbit

500 bp   SbPEPC1t

pGL222

RB   attR4   attR3   LB
Buf   PLTp   *Wus2*   Buf   *ccdB*   PvUbi2p   *Hph*   SbUbip   *ZsG*   OsUbit

500 bp   IN2-1t   SbPEPC1t

(D)

Immature embryo
isolation

Co-cultivation
1 week

Resting
1 week

Regeneration
4-6 weeks

Rooting
2 weeks

Agrobacterium
infection

(E)   **(F)**   **(G)**   **(H)**

Figure 1. Workflow for construct design used for sorghum immature embryo

transformation.

(A) Diagram illustrating the workflow for construct creation and usage. **(B)** Maps of shuttle

vectors. Polycistronic shuttle vector contains a polycistronic gRNA transcription unit with tRNA

spacers; monocistronic shuttle vector contains monocistronic gRNA transcription units driven by

Pol III promoters. Rectangles with arrows indicate promoters. Other rectangles indicate exons

(dark-colored) or terminators (light-colored). Black horizontal lines indicate introns. Grey

vertical lines indicate Gateway recombination sites. **(C)** Maps of binary vectors. pGL206

represents the basic binary vector backbone. pGL216 is designed for testing the function of

morphogenic genes in enhancing transformation frequencies. pGL222 is a binary vector

backbone used in genome editing applications. Rectangles with arrows indicate promoters. Solid

rectangles indicate exons (dark-colored) or terminators (light-colored). Light gray rectangles

denote buffer regions; dark gray rectangles represent *ccdB* genes. Black vertical lines indicate T-

DNA borders; gray vertical lines indicate Gateway recombination sites; black vertical lines

indicate left and right borders. *Hygromycin phosphotransferase* (*Hph*) gene is driven by

switchgrass *ubiquitin 2* promoter (*PvUbi2*p) and terminated by sorghum *phosphoenolpyruvate*

carboxylase terminator (*SbPEPC1*t); *ZsGreen1* (*ZsG*) gene is driven by sorghum ubiquitin

promoter (*SbUbi*p) and terminated by rice *ubiquitin* terminator (*OsUbi*t); Maize *Wuschel2*

(*Wus2*) gene is driven by maize *phospholipid protein* promoter (*PLT*p) and terminated by maize

IN2-1 terminator (*IN2-1*t). **(D)** Diagram illustrating the transformation workflow, from immature

embryo isolation through rooted regenerants. **(E-H)** Sorghum tissues at various stages post-

Agrobacterium infection. **(E)** Immature embryos one week after infection; **(F)** two weeks after

infection and **(G)** four weeks after infection, visualized with overlays of brightfield and

fluorescence images captured with a Zeiss AxioZoom V16 low-magnification epifluorescence

microscope. Scale bars indicate 2 mm. **(H)** Sorghum tissue four weeks after Agrobacterium

infection, with two weeks of exposure to hygromycin selection. Scale bar indicates 2 mm.

Figure 2. CRISPR/*Cas9* and CRISPR/*SpRY*-mediated editing of the *SbPDS* gene using *Wus2*-integrated transformation

(**A**) Top panel: schematic representation of the structure of *SbPDS*, depicting *SbPDS* exons and gRNA target sequences. Light green rectangle indicates promoter. Light yellow rectangles represent untranslated regions. Dark gold rectangles represent exons. Gold lines indicate introns. Guide RNAs are indicated by brown lines. Bottom panel: representative genotypes of PDS edits in T_0 generation. Underlined sequences indicate gRNA Protospacer Adjacent Motif (PAM) sequences, and triangles represent potential double-strand break sites. Knockout scores for each gRNA are shown after each editing outcome. Knockout scores in percentages represent the dominant genotypes identified from amplicon sequencing, followed by analysis using Synthego Inference of CRISPR Edits (ICE) tools. (**B**) Phenotypes of PDS gene-edited plants. The left panel shows a non-edited plantlet; middle panel shows a fully albino plantlet; right panel shows a chimeric albino plantlet. Images were taken four weeks after initiation of rooting. Scale bar = 2 cm. (**C**) Violin plot showing the distribution of ZmCas9i-mediated knockout scores obtained from Sanger sequencing followed by Synthego ICE analysis. For PDS editing with polycistronic gRNA configurations (PLC), n=76; for PDS editing with monocistronic configurations (MNC), n=41. (**D**) Representative genotypes on gRNA2.2, resulting from ZmSpRYi-mediated editing of the SbPDS gene in the T_0 generation. Underlined sequences indicate gRNA PAM sequences; triangles represent potential double-strand break sites. Knockout scores for each gRNA are shown after each editing outcome. Knockout scores in percentage represent the dominant genotypes identified from amplicon sequencing, followed by analysis using Synthego ICE tools. For PDS editing with the polycistronic gRNA configuration, results are presented in the left panel; for PDS editing with monocistronic gRNA configuration, results are shown in the right

panel. **(E)** Violin plot showing distribution of ZmSpRYi-mediated knockout scores for *PDS*

gRNA2.2 edits mediated by CRISPR/SpRY, comparing polycistronic and monocistronic gRNA

expression cassettes. For PLC, n = 68; for MNC, n = 92. **(F)** Comparison of deletion positions

between ZmCas9i and ZmSpRYi at gRNA2.2 (NGG PAM sites) in T_0 plantlets. Each dot

represents a biological replicate. Distribution of deletion percentage from gRNA PLC and MNC

configurations are shown by highlighting the median and quartiles of each. ZmCas9i-PLC, n=76;

ZmCas9i-MNC, n=41. ZmSpRYi-PLC, n = 68; ZmSpRYi-MNC, n = 92. **(G)** Comparison of

indel size between ZmCas9i and ZmSpRYi at gRNA2.2 (NGG PAM sites) in T_0 plantlets. Each

dot represents a biological replicate. Distribution of indel percentages from gRNA PLC and

MNC configurations are shown by highlighting the median and quartiles of them. ZmCas9i-PLC,

n=76; ZmCas9i-MNC, n=41. ZmSpRYi-PLC, n = 68; ZmSpRYi-MNC, n = 92.

Figure 3. CRISPR/*SpRY*-mediated editing of cis-regulatory elements in promoter regions

(A) Schematic representation of CRISPR/*SpRY* target sites on the *SbPDS* gene. A light green rectangle represents the promoter region; light gold rectangles indicate untranslated regions (UTRs); dark gold rectangles denote exons; connecting light gold lines represent introns. Guide RNA (gRNA) target sites are shown as green lines, with PAM sequences highlighted in bold.

SbPDS gRNA (Spg) indicates the gRNA used for editing. **(B)** Violin plot showing distribution of

individual knockout scores, represented as dots, for non-canonical gRNAs targeting *SbPDS*, as

mediated by CRISPR/*SpRY*. $n = 31$. Spg was defined in the figure 3A legend. **(C)** Genotypic

outcomes of zSpRYi-mediated editing targeting of the *SbPDS* promoter and coding sequences.

Top panel: Diagram of *SbPDS* gene structure. A light green rectangle represents the promoter

region; light gold rectangles indicate untranslated regions (UTRs); dark gold rectangles denote

exons; connecting light gold lines represent introns. gRNA target sites are indicated by green

dots. *Bottom panels:* Representative T□ genotypes and mutation patterns. *Left:* Genotypes from

editing cis-regulatory elements in the promoter region. *Right:* Genotypes from edits within the

exons. Vertical dashed lines indicate predicted double-strand break sites. Black rectangles

represent deletions introduced via DNA repair; the width corresponds to indel length, and the

height indicates the frequency of edited alleles, based on Synthego ICE analysis. Spg: *Sb<u>PDS</u>*

gRNA. **(D)** Conservation of *CYP79A1* promoter regions across C$_4$ grasses. Bar graph shows

sequence similarity between the *SbCYP79A1* promoter (from *Sorghum bicolor* genotype

RTx430) and orthologous regions from *Miscanthus sinensis*, *Saccharum officinarum*, and

Andropogon gerardii. Promoter regions analyzed span 2,000 bp upstream of the translation start

site, with each bar representing a 30-nt window. gRNA locations within the promoter and UTR

are indicated by black dots; PAM sequences are in bold. **(E)** Overview of the dhurrin

biosynthetic pathway and cis-regulatory element editing strategy. *Left panel:* Simplified pathway

highlighting *CYP79A1* as the initial step in dhurrin biosynthesis. *Right panel:* Workflow

illustrating the use of cis-regulatory element editing to attempt to reduce dhurrin content by

targeting the *SbCYP79A1* promoter and UTR. **(F)** Genotypic outcomes of zSpRYi-mediated

editing of the *SbCYP79A1* promoter and UTR. *Top panel:* Gene diagram showing promoter,

UTR, exons, and gRNA target sites (dots below the gene structure). A light green rectangle

represents the promoter region; light gold rectangles indicate untranslated regions (UTRs); dark

gold rectangles denote exons; connecting light gold lines represent introns. gRNA target sites are

indicated by green dots. *Bottom panels:* Representative T□ genotypes reflecting mutation

patterns from cis-regulatory region editing. Deletions are marked by dark brown rectangles. The

width of the rectangles corresponds to indel length, and the height indicates the frequency of

edited alleles, based on Synthego ICE analysis. A magnified view of editing outcomes from the

cpg3 gRNA is also shown. Cpg: <u>C</u>YP79A1 <u>p</u>romoter <u>g</u>RNA.

Table 1. Transformation, editing and knockout efficiencies of Wus2-assisted transformation,

based on one plantlet per IE

gRNA system (Construct)	Temp (°C)	Replication	# IEs used	# Regenerants	# Escapes (%) [†]	+ for hph [‡]	Transformation frequency (%) [§]	# Edited plants	Editing efficiency (%) [¶]	# Plants with knockout [††]	Knockou efficiency (%
Polycistronic (pGL220)	28	1st	132	47	6 (12.8)	41	**31.0**	24	**58.5**	24	**58.5**
		2nd	169	44	4 (9.1)	40	**23.7**	25	**62.5**	25	**62.5**
		Total	301	91	10(10.9)	81	**26.9 (%)**	49	**60.5 (%)**	49	**60.5 (%**
	25	1st	143	30	4 (13.3)	26	**18.2**	17	**65.4**	17	**65.4**
Monocistronic (pGL221)	28	1st	93	24	1 (4.2)	23	**24.7**	22	**95.7**	22	**95.7**
		2nd	112	28	1 (3.6)	27	**24.1**	26	**96.3**	26	**96.3**
		Total	205	52	2(3.8)	50	**24.4 (%)**	48	**96.0 (%)**	48	**96.0 (%**
	25	1st	105	34	1 (2.9)	33	**31.4**	33	**100.0**	33	**100.0**
		2nd	75	23	0(0)	23	**30.7**	21	**91.3**	21	**91.3**
		Total	180	57	1(1.8)	56	**31.1 (%)**	54	**96.4 (%)**	54	**96.4 (%**

[†] # Escapes (%) = # and % of regenerated plants without transgene (T-DNA) = (# Escapes/# Regenerants) 100

[‡] + = # plants PCR-positive for *hph*

[§] Transformation frequency (%) = (# + for *hph* divided by # IEs used) x 100

[¶] Editing efficiency (%) = (# edited plants divided by # plants + for hph) x 100

[††] # Plants with knockout = # plants with frameshift mutation and albino phenotype

[‡‡] Knockout efficiency(%) = (# plants with knockout divided by # + for hph) x 100

Table 2. Transformation, editing and knockout efficiencies of Wus2-assisted transformation, based on multiple regenerated plantlets from each IE

gRNA system (Construct)	Temp (°C).	Replications	# IEs used	# Regenerants	# Escapes (%)[†]	+ for hph[‡]	Transformation frequency (%)[§]	# Edited plants	Editing efficiency (%)[¶]	# Plants with Knockout[††]	Knockout efficiency (%)
Polycistronic (pGL220)	28	1st	132	60	6 (10)	54	**40.9**	30	**55.6**	30	**55.6**
		2nd	169	48	4 (8.3)	44	**26.0**	27	**61.4**	27	**61.4**
		Total	301	108	10(9.2)	98	**32.6**	57	**58.2**	57	**58.2**
	25	1st	143	31	4 (12.9)	27	**18.9 (%)**	17	**63.0 (%)**	17	**63.0**
Monocistronic (pGL221)	28	1st	93	64	1 (1.6)	63	**67.7**	61	**96.8**	60	**95.2**
		2nd	112	77	3 (3.9)	74	**66.1**	66	**89.2**	63	**85.1**
		Total	205	141	4(2.8)	137	**66.8 (%)**	127	**92.7 (%)**	123	**89.8**
	25	1st	105	93	6 (6.5)	87	**82.9**	87	**100.0**	84	**96.6**
		2nd	75	60	4(6.7)	56	**74.7**	51	**91.1**	49	**87.5**
		Total	180	153	10(6.5)	143	**79.4 (%)**	138	**96.5 (%)**	133	**93.0**

[†] #Escapes (%) = # and % of regenerated plants without transgene (T-DNA) = (# Escapes/# Regenerants) X 100
[‡] + = # Plants PCR-positive for *hph*
[§] Transformation frequency (%) = (# + for *hph* divided by # IEs used) x 100
[¶] Editing efficiency (%) = (# Edited plants divided by # plants + for hph) x 100
[††] # plants with knockout = # plants with frameshift mutation and albino phenotype
[‡‡] Knockout efficiency (%) = (# Plants with knockout divided by # + for hph) x 100

943 **Table 3.** Transformation, editing and knockout efficiencies of *SpRY* using *Wus2*-assisted

944 transformation

gRNA system (Construct)	Replications	# IEs used	# Regenerants	# Escapes (%)[†]	+ for hph[‡]	Transformation frequency (%)[§]	# Edited plants	Editing efficiency (%)[¶]	# Plants with Knockout[††]	Knockout efficiency (%)
Polycistronic (pGL226)	1st	107	46	3 (6.5)	43	40.2	11	25.6	11	25.6
	2nd	60	21	4 (19.0)	17	28.3	5	29.4	5	29.4
Monocistronic (pGL227)	1st	119	49	0 (0)	49	41.2	40	81.6	40	81.6
	2nd	82	46	3 (6.5)	43	52.4	41	95.3	39	90.7

[†] # Escapes (%) = # and % of regenerated plants without transgene (T-DNA) = (# Escapes/# Regenerants) X 100

[‡] + = # Plants PCR-positive for *hph*

[§] Transformation frequency (%) = (# + for *hph* divided by # IEs used) x 100

[¶] Editing efficiency (%) = (# Edited plants divided by # plants + for hph) x 100

[††] # Plants with knockout = # Plants with frameshift mutations and albino phenotypes

[‡‡] Knockout efficiency (%) = (# Plants with knockout divided by # plants + for hph) x 100

 Table 4. Cis-regulatory element and gene exon editing efficiency of SpRY on phytoene desaturase and

953 cytochrome P450 genes

Target	Rep.[†]	#Reg.[‡]	#Edits %Efficiency[§]		%gRNA1 efficiency[§]		%RNA2 efficiency[§]		%gRNA3 efficiency[§]		%gRNA4 efficiency[§]		%gRNA5 efficiency[§]		%gRNA6 efficiency[§]	
SbPDS	1st	14	8	57.1	8	57.1	8	57.1	6	42.9	3	21.4	3	21.5	5	35.7
	2nd	50	23	46.0	21	42.0	22	44.0	18	36.0	11	22.0	14	28	16	32.0
SbCYP79A1	1st	40	19	47.5	9	22.5	12	30.0	19	47.5	0	0	NA[¶]	NA[¶]	NA[¶]	NA[¶]

954 [†] Rep.= Replication

955 [‡] #Reg.= Regenerant number

956 [§] %gRNA efficiency = (# Edited plants divided by # plants + for hph) x 100

957 [¶] NA = not applicable