High-frequency *Wus2*-integrated transformation toolkit enhances *Cas9* editing efficiency and expands capability with *SpRY* in sorghum

Jianqiang Shen[1,3,4], Kiflom Aregawi[1,3], Sultana Anwar[1], Tamara Miller[1], Peggy G. Lemaux[1,2,4]

[1]Department of Plant and Microbial Biology, University of California, Berkeley CA 94720, USA.

[2] Innovative Genomics Institute, University of California, Berkeley CA 94720, USA.

[3] These two authors contributed equally.

[4] Correspondence Email: shenjq@berkeley.edu or lemauxpg@berkeley.edu

Abstract

Sorghum bicolor L. (Moench) ranks as the fifth most important crop globally, serving diverse purposes such as forage, food, feed, and fuel. Its natural tolerance to various environmental stresses makes it a valuable resource for addressing climate resilience. Harnessing such genetic resources for crop improvement requires highly efficient genetic transformation and genome editing tools. However, sorghum has historically been recalcitrant to genetic transformation, and publicly available engineering tools remain limited. In this study, we report a high-efficiency engineering toolkit, including shuttle and binary vectors, optimized for genome editing via *Agrobacterium*-mediated transformation. Using this toolkit, we successfully introduced CRISPR/Cas9-based editing machinery to target the *phytoene desaturase* (*PDS*) gene, achieving editing efficiencies up to 95.7% with monocistronic gRNA expression cassettes. To further expand the genome editing repertoire, we employed a protospacer adjacent motif (PAM)-flexible *Cas9* variant, *SpRY* (*ZmSpRYi*), and achieved comparable editing efficiencies. Using our toolkit increases the potential for advanced editing tools to be used in engineering genetic variation of agronomically important quantitative traits. It also paves the way for deploying advanced genome editors, such as prime editors and base editors, to manipulate genetic and genomic resources. These advances provide a robust approach for advancing research in sorghum and other cereals, offering new opportunities for sustainable crop improvement.

Keywords

28 *Sorghum bicolor, Wuschel2*, transformation, editing, CRISPR, SpRY

Introduction

Meeting future demands for crop production with more variable climate conditions and an increasing
world population will require growing more food and feed on less land. *Sorghum bicolor* L. (Monech) is an
ideal crop for future applications because of the diversity of modern germplasm, particularly with respect
to its multiple abiotic stress tolerances (Dahlberg, Burke, and Rosenow 2004; Paterson et al. 2009;
Varoquaux et al. 2019). Breeding improved sorghum varieties using targeted genome editing and marker-
assisted selection will accelerate development of improved cultivars (Ramstein, Jensen, and Buckler 2019;
Schindele, Dorn, and Puchta 2020; Zhu, Li, and Gao 2020). However, utilizing CRISPR/*Cas9* genome editing
in sorghum lags in efficiency relative to other crops due to low editing efficiencies, a lack of efficient
transformation systems for diverse genotypes, and a lack of optimized, publicly available tools. The
current work seeks to address all three limitations by introducing a novel set of resources for high
throughput sorghum transformation and genome editing.

Sorghum genome engineering has historically been a challenging process due to recalcitrance of most
sorghum genotypes to *in vitro* tissue culture and a lack of technical expertise and effective tools for
transformation (Altpeter et al. 2016). A significant challenge of genome engineering in crop species is the
requirement for germline transmission of mutations necessitating gene delivery in pluripotent cells.
Transcription factors, such as Babyboom (BBM) and/or Wuschel2 (WUS2), increase the population of cells
in target tissues that are competent for germline transmission (Lowe et al. 2016), (Lowe et al. 2018).
Previously we demonstrated that incorporation of transient *Bbm* and *Wus2* expression during sorghum
Agrobacterium-mediated gene delivery increases transformation frequency for several recalcitrant
varieties (Aregawi et al. 2022). Concurrently, Che et al. demonstrated that during sorghum transformation
incorporation of the *Wus2* morphogenic gene alone increased CRISPR/*Cas9*-mediated gene dropout rate
of the *Brown midrib* gene (Che et al. 2022). The breakthrough in transformation technology due to
morphogene-assisted transformation has greatly expanded the possibility of genotype-independent gene
editing; however, developing tools for high efficiency genome editing with CRISPR remains a challenge.

The CRISPR/*Cas9* genome editing system**,** first discovered as a component of the adaptive immune

response in *Streptococcus pyogenes* bacteria (Jinek et al. 2012), is a powerful tool for genome editing that

has potential for functional genomics and accelerated breeding in plants (Zhang et al. 2019). Editing in

non-model plants lags in efficiency compared to animal and bacterial systems, due to several factors,

including the plant 'regeneration bottleneck' and sub-optimal *in planta* expression of editors.

Optimization of the Cas endonuclease, guide RNA expression and use of Cas variants with expanded PAM

recognition sequences have been proposed to increase the efficiency of gene editing in plants (Liu, Li, and

Godwin 2019; Q. Ren et al. 2021; Xu et al. 2021). To overcome these challenges, codon optimization of

the Cas9 nuclease was used to increase gene knockout editing efficiency in Arabidopsis and tobacco

(Castel et al. 2019; Grützner et al. 2021). The use of specific promoters and guide RNA transcriptional

configurations has been shown to increase sorghum editing efficiency as demonstrated when using the

endogenous U6 promoter driving gRNAs (Massel et al. 2022). Another inherent limitation to the capacity

for introducing genetic variation at specific sites is the fact that the canonical *Cas9* endonuclease requires

the NGG (N = A, T, C, and G) Protospacer Adjacent Motifs (PAMs) site, which severely limits the genome

sites available for editing. Several Cas variants have been engineered to overcome the PAM limitation of

SpCas9 (Tang et al. 2017; B. Ren et al. 2019; Ming et al. 2020). *SpRY* endonuclease is an engineered variant

of *Streptococcus pyogenes Cas9* with a nearly PAM-less requirement and has been used for base editing

and functional genetics in rice (B. Ren et al. 2019; Q. Ren et al. 2021; Xu et al. 2021).

The present work addresses three main goals for advancing transformation and editing of complex traits

in *Sorghum bicolor*: increasing the efficiency of CRISPR/*Cas9* editing, expanding available PAM recognition

sites, using the Cas9 variant, ZmSpRYi and providing needed genome editing capability to public

researchers. We developed a flexible functional genomics toolbox for Agrobacterium-mediated sorghum

transformation and genome editing. The toolbox includes a binary T-DNA vector incorporating the *Wus2*

gene, along with *hygromycin B phosphotransferase* (*hph*) and *ZsGreen* (*ZsG*) as selection and visual

markers. For CRISPR/*Cas9* genome editing, we utilized an intronized Cas9 endonuclease in a shuttle

vector, enabling easy insertion of designed guide RNA sequences. Additionally, for the first time in

sorghum, we demonstrate editing with the nearly PAM-less Cas9 variant ZmSpRYi, employing mono- or

poly-cistronic guide RNA expression. Overall, this high-efficiency toolbox enhances the scalability of

genome editing in sorghum, enabling faster breeding and supporting advancements in synthetic biology.

Results

Design of high efficiency binary and shuttle vectors

A two-step strategy was developed that involves gRNA validation in protoplasts, followed by stable plant genetic transformation (**Figure 1A**), that enhances efficiency and accessibility for researchers to perform sorghum transformation and editing. The editing machinery is integrated in a Level 1 construct, the shuttle vector, which facilitates guide RNA (gRNA) customization through insertion of either polycistronic or monocistronic transcriptional units (**Figure 1B**). The gRNA scaffold and Pol III promoters necessary for driving gRNA expression are supplied by the Level 0 constructs (**Figure S1**). These two elements are sequentially assembled to form complete gRNA transcription units, which are transfected into sorghum protoplasts to evaluate editing efficiency of the gRNAs (Jores et al. 2021; Lee et al. 2023).

To enhance efficiency of sorghum transformation, a series of binary constructs were developed, all of which are compatible with Gateway LR recombination and tailored to incorporate genes of interest for multiple applications (**Figure 1C**). The foundational construct, pGL206, includes a *hygro*mycin resistance gene (*hph*) as a selectable marker (**Figure 1C**). Hygromycin was chosen for its stringent selection properties during in vitro tissue culture of sorghum (van den Elzen et al. 1985). This construct serves as a platform for testing candidate morphogenic genes, i.e., maize *Wuschel2* (*Wus2*). To facilitate tracking of somatic embryogenesis and plantlet development, a visible marker *ZsGreen1,* driven by the sorghum ubiquitin promoter, was incorporated near the T-DNA left border; this construct is pGL216. Building on previous findings of enhanced editing efficiency with integration of maize *Wus2* (Che et al. 2022), the construct, pGL216, was assembled with *Wus2*, driven by the maize phospholipid transfer protein promoter (PLTP) (**Figure 1C**). This modification led to development of pGL222 (**Figure 1C**), a binary vector optimized for delivering genome-editing machinery. Presence of the fluorescence marker validates the presence of at least one complete T-DNA copy and easy visualization of T-DNA segregation in the T_1 generation. These constructs provide a robust framework for improving transformation and genome-editing efficiencies in sorghum.

Transformation and editing frequency using *Wuschel2* integration

To evaluate transformation frequency in genome editing trials, we targeted the *Sorghum bicolor phytoene desaturase* (*SbPDS*) gene using two gRNAs, PDS gRNA1.1 and 2.2 (**Figure 3A**), designed to knock out its function (Aregawi et al. 2022). Previous studies in wheat and Setaria indicated editing efficiency may vary depending on the nature of the transcriptional units of gRNAs in C3 and C4 grasses (Liang et al. 2021; Milner et al. 2024); however, this has not been thoroughly investigated in sorghum using CRISPR/*Cas9*. To study this, constructs were designed containing two gRNAs under two transcriptional configurations, polycistronic and monocistronic (**Figure 1B**). In the first construct, gRNAs were expressed from a polycistronic unit, driven by the sorghum snoRNA U6.2.3 promoter (Massel et al. 2022). In the second, gRNAs were expressed in separate monocistronic units driven by the snoRNA U6.2.3 and snoRNA U6.3.1 promoters (Massel et al. 2022). Both validated gRNA transcriptional units were combined with a Cas9 expression cassette and integrated into the binary vector pGL222, resulting in two editing constructs, pGL221 (polycistronic) and pGL220 (monocistronic) (**Figure S2**). These constructs, containing the morphogenic gene *Wuschel2* (*wus2*) and the selectable marker gene, *hph*, were used to assess the impact of gRNA transcriptional configurations on editing efficiency.

These two constructs were transformed into sorghum immature embryos (IEs) using a modified protocol (Aregawi et al. 2022). In this approach, the heat treatment typically applied after the callus resting stage was omitted, because the *Wus2* expression cassette was not excised after integration (**Figure 2A**). Following Agrobacterium infection, the IEs were maintained at 25°C for one week on co-cultivation media. Previous studies indicated that higher temperatures may enhance mutagenesis efficiency when employing CRISPR/*Cas9* in plants (Milner et al. 2020; Blomme et al. 2022). To explore this, tissues were incubated at either 25°C or 28°C until plant regeneration. Development of IEs were monitored to evaluate their responses to *Wus2* expression at 1-, 2-, and 4-weeks post-infection (**Figures 2B–E**). Fluorescence signaling was detected as early as one week after infection (**Figure 2B**) and peaked at the end of the resting stage (**Figure 2C**). Somatic embryogenesis was most prominent during the two-week time period, with regenerated plantlets forming by the fourth week (**Figures 2D–E**). Prior to transferring the putative regenerants to rooting media, fluorescence microscopy was used to screen for ZsGreen expression, confirming integration of the full-length T-DNA, given that the ZsGreen expression cassette is located adjacent to the right border. Subsequently, putative transgenic plantlets were analyzed via PCR to further confirm T-DNA integration.

Transformation and editing efficiencies were calculated using two strategies. The first was calculated

when including only one transgenic plant from each IE, the second when including all regenerated

transgenic plants. Each way has its advantages. Considering a single plant per IE is helpful when using

overexpression to get independent transgenic plants. Unlike when using a classical sorghum

transformation strategy, here we regenerated multiple plantlets from each IE. Transformation frequency

indicates the proportion of IEs within the total IEs used that successfully produce transformants.

Regarding the editing events, considering all regenerants derived from a single IE is beneficial as multiple

independently edited events can originate from the same IE.

When considering only one plant per IE at 28°C for the monocistronic strategy, we generated 50

transgenic plants from two replicates, giving a transformation frequency of 24.4%. Given 79

transformants, an average transformation frequency of 36.6% was obtained from the polycistronic

strategy. Unlike the 28°C result, tissues incubated at 25°C showed normal phenotype. Because of this

difference, more plants were regenerated, increasing the transformation frequency to 31.4% and 43.3%

for the monocistronic versus polycistronic strategies, respectively.

When all regenerated transgenic plants were considered, transformation frequency substantially

increased (**Table 2**). At 25°C, the monocistronic and polycistronic strategies yielded transformation

efficiencies of 82.9% and 100%, respectively. At 28°C, the monocistronic and polycistronic strategies led

to 66.9% and 62.7% transformation efficiencies, respectively. In general, tissues kept at 28°C showed more

necrosis than those kept at 25°C (**Figure S3**). This reduced the number of regenerated plants and thus the

transformation efficiencies were lower (**Table 1**).

Wuschel-mediated CRISPR/*Cas9* editing and editing efficiency

Previous work demonstrated that various factors are needed to improve CRISPR/*Cas9* editing efficiency.

One factor is intronized and maize codon-optimized *Cas9* (Castel et al. 2019; Grützner et al. 2021). This

version of *Cas9* could improve genome editing by enhancing the stability, translation, and expression of

Cas9 (Castel et al. 2019; Grützner et al. 2021). Another potential factor is the nature of the promoters

driving gRNAs, which play an important role in regulating expression of the gRNAs (Zhan et al. 2021;

Massel et al. 2022). Another might involve incubating tissues at higher temperatures, which might lead to

improved editing efficiency (LeBlanc et al. 2018; Milner et al. 2020). Finally, Che et al. (Che et al. 2022)

reported that the *Wus2*-integrated transformation strategy resulted in improved CRISPR/*Cas9* gene

dropout frequency, possibly due to enhanced cell division and/or chromatin accessibility.

In our work, different treatments were incorporated that could lead to improved genome editing and

knockout efficiencies, as described above. These include using the *Wus2*-integrated transformation

strategy, a maize codon-optimized and intronized *Cas9*, monocistronic and/or polycistronic tRNA-gRNA

expression cassettes, and/or the *SbU6* 2.3 and *SbU6* 3.1 promoters driving gRNAs. Effects of two

temperature conditions, 25°C and 28°C, during culturing of tissues were also tested.

To assess the impact of the totality of these various factors, *SbPDS* was amplified from transgenic plants

and characterized using Sanger sequencing to study the nature of the genome edits. Sequencing results

were further analyzed using the Synthego ICE tool. The editing efficiency of the edited plants, referring to

the knockout score, ranged from 0% to 100% (**Figure 3C**). Observed albino phenotypes **(Figure 3B)** were

used to visually identify plants with *SbPDS* gene knockouts and to calculate knockout efficiency. Based on

these analyses, it appears that, by combining the effects of all of these factors, editing and knockout

efficiencies greatly improved over those seen in our previous publication, which was 22.6% and 16.7%

((Aregawi et al. 2022)). When considering one plant per IE, the monocistronic strategy yielded a very high

editing and knockout efficiency of approximately 100% (**Table 1)**. Treating tissues at 28°C, compared to

25°C, showed no difference in editing and knockout scores. Although lower than the monocistronic

strategy, the polycistronic strategy still resulted in relatively high editing efficiencies of 76.6% and 88.5%

at 28°C and 25°C conditions, respectively (**Table 1)**. Most edited plants had a small, 1- or 2-bp deletion

upstream of the PAM region (**Figure 3D**). Only a small number (16.7 %) showed larger (~5-bp) deletions

or insertions.

Comparison of plant phenotype with knockout score in *SbPDS* edited plantlets

197   In our study, the *SbPDS* gene was selected as the target for genome editing, with the albino

198   phenotype serving as a visual phenotype for successful gene knockouts. T$_{0}$ plants were

199   phenotyped based on the percentage of the albino sectors observed. Fully albino plantlets were

scored as 100%, while 0% represented fully green plants. This approach enabled assessment of phenotype, which is not feasible for genes for which knockouts do not produce observable phenotypic changes. Mutations at the target sites were analyzed using Synthego ICE analysis, which uses Sanger sequencing data from amplicons encompassing the potential edited sites. This analysis enabled precise identification and quantification of the types and frequencies of mutations induced by genome editing. For the majority of plants, phenotypic and genotypic scores were closely aligned, with few discrepancies (**Table S2**). The two metrics showed a strong correlation, with a Pearson's correlation coefficient of 0.81. This strong correlation underscores the utility of Synthego ICE as a reliable tool for estimating the nature of genotypic changes, especially in cases where gene knockouts do not manifest visually detectable phenotypes in T_0 generation.

Editing efficiency using PAM-less intronized *Cas9* variant, ZmSpRYi

To enhance genome editing capabilities and to identify options with fewer PAM sequence constraints, a range of nucleases were considered, including *Cas9* variants and *Cas9* orthologs (Ge et al. 2019; B. Ren et al. 2019; Zhong et al. 2019; Q. Ren et al. 2021; Xu et al. 2021). Previous studies demonstrated that CRISPR/*Cas9*-mediated editing of promoters can generate diverse cis-regulatory alleles, providing valuable quantitative variation for crop breeding (Rodríguez-Leal et al. 2017; Zhou et al. 2023; Patel-Tupper et al. 2024). However, the standard CRISPR/*Cas9* system relies on canonical gRNAs with NGG as the PAM sequence, posing a significant limitation since promoter regions are typically AT-rich. This restricts the feasibility of targeted editing within cis-regulatory modules.

To address this restriction, PAM-flexible *Cas9* variants, such as *SpRY* (Walton et al. 2020), were used to enable targeted promoter editing and to expand genome engineering in AT-rich genomic regions. Building on an intronized *ZmCas9*, targeted mutations were introduced in eleven nucleotides to create a maize codon-optimized *SpRY* variant, *ZmSpRYi*, with an intron for enhanced transcription and translation (**Figure 4A**).

Previous studies suggested that *SpRY* activity may be less efficient due to its lack of an NGG PAM requirement (Hibshman et al. 2024). To evaluate the in vivo editing efficiency of ZmSpRYi, compared with

that of ZmCas9i, *ZmCas9i* was substituted with *ZmSpRYi* in constructs to enable a direct comparison using the same gRNAs. Two constructs were designed, featuring polycistronic and monocistronic transcriptional units of gRNAs, and these were transformed into sorghum RTx430. Regenerants from using the monocistronic editing construct (pGL227) exhibited an albino phenotype (**Figure 4B**) as early as four weeks post-Agrobacterium infection, whereas no albino phenotype was observed at that same time with the polycistronic editing construct (pGL226) (**Figure S2**). To assess editing efficiency of ZmSpRYi, all independent transformants were genotyped by PCR amplification, followed by Sanger sequencing. Analysis of the sequencing data using Synthego ICE revealed that, when using the polycistronic transcription system, 17 out of 60 (28.3%) independent transformants were edited, whereas 77 out of 89 (86.5%) transformants were edited using the monocistronic system (**Table 2, Figure 4C**). Further analysis showed that the monocistronic system yielded a higher knockout score (**Figure 4E**), suggesting that editing events occur earlier when individual promoters are used to drive gRNA expression. Previous studies demonstrated that *SpRY* exhibits self-editing activity, targeting its own expression cassettes within integrated T-DNA fragments (**Figure 4A**) (Q. Ren et al. 2021; Xu et al. 2021). Here, we analyzed gRNA expression cassettes within T-DNA and self-editing was observed in 1 out of 4 cases for polycistronic transformations and 3 out of 8 cases for monocistronic transformations (**Figure 4D**). These results underscore the enhanced efficiency in sorghum of the monocistronic system for genome editing using *ZmSpRYi*.

Discussion

Development of efficient, targeted, and versatile genome-editing tools is needed to advance genetic engineering and editing in bioenergy crops like sorghum. As a C4 plant, sorghum is naturally efficient in photosynthesis and additionally it is highly resilient to environmental stresses, such as drought, flooding, and salinity (Varoquaux et al. 2019). However, the molecular mechanisms underlying such key agronomic traits and the validation of appropriate genetic resources remain significant challenges. To address these issues and enable functional gene analysis while introducing desirable genetic variations in sorghum, we developed a shuttle-binary vector toolbox to address the current limitations of genome editing. This system simplifies genetic engineering by incorporating into the construct the morphogenic gene *Wus2*, which accelerates optimized delivery of editing machinery. Additionally, several optimizations were introduced to improve the efficiency of CRISPR/Cas9-based genome editing in sorghum. These included a codon-optimized Cas9 containing an intron, high-efficiency endogenous U6 promoters, and monocistronic

260 gRNA expression cassettes. These modifications addressed previous limitations and significantly expanded the

261 potential applications of this genome editing technology in sorghum.

262 Using Wuschel integration simplifies the binary vector and improves

263 transformation frequency

Previous studies demonstrated that the morphogenic regulators, BBM and WUS2, significantly enhance

transformation frequency in monocot plants, including sorghum (Lowe et al. 2016; Aregawi et al. 2022).

However, continued integration of these regulators often results in developmental aberrations during the

vegetative stage. To address this issue, the CRE/LoxP recombination system was employed to excise these

regulators, following somatic embryo initiation (Lowe et al. 2016). While effective, this approach increases

binary vector size, complicating molecular cloning and construct development. Efforts to overcome this

challenge include development of systems that combine transient and stable *Wus2* expression (Hoerster

et al. 2020), reducing construct size. In our approach, we redesigned the binary vector into a more

versatile delivery platform by incorporating the LR recombination site, making it compatible with the

Gateway system used frequently among plant biotechnologists for construct development. When paired

with a *Wus2* expression cassette, we used this system to achieve transformation frequencies of 24 to 42%,

comparable to those attained using BBM/WUS2-assisted transformation (Aregawi et al. 2022).

Despite this success, limitations persist with the integration-based delivery of Wus2. Approximately 10%

of the T_0 transformants exhibited developmental deficiencies, such as curly leaves during the vegetative

stage, potentially caused by ectopic expression of Wus2. To address this issue, pGL216 was considered as

a platform for testing alternative developmental regulators, including Wuschel orthologs from other

species or members of the Wuschel homeobox gene family (**Figure 1C**). These alternative candidates may

enhance somatic embryo initiation while minimizing developmental disruptions, offering a more refined

approach to transformation and regeneration.

Optimizing *Cas9* and guide RNA expression cassettes increases editing efficiency

In our previous work, we successfully achieved *PDS* knockouts in sorghum using CRISPR/*Cas9*, with an editing efficiency of 16.7% (Aregawi et al. 2022). However, enhancing genome-editing efficiency further requires overcoming certain challenges, particularly ensuring robust expression of the editing machinery, including *Cas9* protein and gRNAs (Liu, Li, and Godwin 2019). To address these challenges, we used a constitutive promoter to drive an intronized maize codon-optimized Cas9 (Che et al., 2022). Intronization has been shown to enhance protein abundance at both transcriptional and translational levels in Arabidopsis (Grützner et al. 2021). For gRNA expression optimization, we utilized the sorghum endogenous snoRNA U6 promoters, which lead to high gRNA expression levels (Massel et al. 2022).

Previous studies suggest that transcriptional differences between polycistronic and monocistronic gRNA transcription units can affect editing efficiency (Liang et al. 2021). However, it remains unclear in sorghum whether polycistronic transcription units compromise editing efficiency. To investigate this, we conducted a side-by-side comparison of polycistronic versus monocistronic gRNA transcription units. Analysis of results demonstrate that the monocistronic gRNA transcription unit achieved significantly higher editing efficiency, up to 91 to 100%, compared to 58.5 to 92% for the polycistronic unit. This indicates that temporal processing of exogenous tRNA may be compromised during the early embryonic development of cultured tissue. We also observed that the albino phenotypes resulting from PDS knockouts were delayed when using the polycistronic gRNA transcription unit. This delay further supports the notion that transcriptional processing inefficiencies might underlie the reduced editing efficiency associated with polycistronic gRNA systems in sorghum. To further investigate this, we assembled a collection of monocot snoRNA promoters, including U6 promoters from sorghum, wheat, and maize, as well as U3 promoters from wheat and rice (**Figure S1**). These promoters have been experimentally validated in their respective monocot species. Level 0 constructs, using these promoters, offer versatile options for combining multiple gRNAs, enabling simultaneous editing of multiple genomic targets. This approach holds promise for applications, such as promoter editing and other complex genetic modifications.

To evaluate the impact of temperature on editing efficiency, we conducted an editing experiment under in vitro tissue culture conditions. Incubation temperature following Agrobacterium infection was increased from 25°C to 28°C. Interestingly, no apparent difference in editing efficiency was observed

between the two temperatures (**Table 1**). This outcome may be attributed to the high baseline editing
efficiency achieved through the modifications made to the Cas9 and gRNA elements. As a result, the
elevated temperature did not provide additional improvements in editing, likely because the system was
already operating near its optimal efficiency. Also, tissues were healthier, and the number of regenerants
appears to be higher at 25 °C. (**Figure S3**). Based on these findings, we suggest 25°C as a more suitable
incubation temperature for efficient genome editing in sorghum.

Expanding the Scope of Genome Editing and Applications Using ZmSpRYi

One potentially impactful application of genome editing in crops is creating cis-regulatory variations to
diversify genetic resources for improving desirable traits (Rodríguez-Leal et al. 2017). Cis-regulatory
editing in promoters and untranslated regions (UTRs), which are typically AT-rich, is often limited, when
using Cas9, because of the strict requirement for NGG PAM sequences. To address this limitation, we
considered several potential genome editing candidates to target AT-rich regions, including Cas9 variants
(Cas9-NG, xCas9, and SpRY) and other Cas orthologs (Cas12a and Cas12b), which exhibit reduced
dependency on CG content at the PAM site (Zetsche et al. 2015; Nishimasu et al. 2018; Hu et al. 2018;
Strecker et al. 2019; Walton et al. 2020). Among these, SpRY demonstrated the highest flexibility due to
its PAM-less feature (Q. Ren et al. 2021; Xu et al. 2021).

Using the same gRNAs targeting the PDS gene, as in our previous *Cas9* experiments (Aregawi et al. 2022),
Zm*SpRY*i achieved a genotypic editing efficiency of 86.5%. However, previous studies suggested that use
of *SpRY* results in compromised double-strand break activity, particularly with non-NGG PAM sequences
(Q. Ren et al. 2021). To further investigate ZmSpRYi activity with non-NGG PAM sequences, we are
currently examining its performance in combination with WUS2. Previous research indicated that
integration of *Wus2* can enhance CRISPR/*Cas9*-mediated gene dropout frequency (Che et al. 2022).
However, it remains unclear as to whether *Wus2* integration can similarly enhance SpRY activity. To
address this question, we designed six gRNAs targeting the promoter, 5' UTR, and exons of the PDS gene
and are currently transforming sorghum with the appropriate editing constructs.

Another concern for SpRY is the potential for off-target effects, including self-editing. Self-editing refers
to mutagenesis occurring within the gRNA expression cassette (Q. Ren et al. 2021; Xu et al. 2021), which
could lead to unintended edits. To assess this risk in edits with high knockout scores we amplified and

sequenced potential off-target sites, including within the T-DNA region. We examined potential self-
editing regions and found that 25.0% to 37.5% of selected edits exhibited self-editing within the T-DNA
fragments (unpublished data). Notably, the target regions were also successfully edited in these cases,
suggesting that self-editing occurred either concurrently or sequentially with the targeted editing process.
The present study highlights the promise of *SpRY* for expanding the genome editing toolbox in AT-rich
regions while addressing challenges, such as compromised activity and off-target risks.
352
In this study, we developed a high-efficiency editing platform incorporating *wus2* integration with a robust
Cas9-based genome editing toolkit, featuring intronized ZmCas9 and ZmSpRY. This transformation
platform is a versatile tool for delivering genome editing machinery that can include base editors and
prime editors. By integrating *wus2* with the SpRY Cas9 variant, we expanded the editing scope to enable
other desirable applications, such as enabling promoter editing. These advancements provide a
streamlined approach to generating diverse genetic variation, facilitating crop improvement efforts that
address food security challenges and contribute to carbon capture initiatives that are addressing climate
change.
361

Materials and Methods

Plant material and growth conditions

364 Seeds of *Sorghum bicolor* genotype RTx430 were sourced from the Germplasm Resources Information
365 Network (GRIN, USDA-ARS, PI 655998). Plants were cultivated in three-gallon pots containing SuperSoil in
366 the Oxford Research Unit South Greenhouse, under controlled conditions of 30°C/24°C (day/night) with a
367 16-hour light/8-hour dark photoperiod. Immature embryos (IEs) were harvested 12–14 days post-
368 anthesis.

Construct creation

To create pGL206, the hygromycin phosphotransferase gene (*hph*), driven by the *Panicum virgatum*
Ubiquitin2 promoter (PvUbi2p), was amplified from pANIC10A (Mann et al. 2012) using the primer SorEd1
and 2 (**Table S1**). All primers used for construct building are described in **Table S1.** The terminator of

sorghum PEPC was amplified from pPHP85425 (Aregawi et al. 2022) using the primers SorEd3 and 4.

These two parts were assembled by overlapping PCR. The vector backbone of pPHP85425 was amplified

using the primers SorEd5 and 6. In addition, the Buffer3 and attR4-attR3 fragments were amplified from

pPHP85425 using the primers SorEd7 and 8. These three fragments were assembled using Gibson

assembly (NEBuilder® HiFi DNA Assembly Master Mix). To construct pGL216, the *ZsG* expression cassette

was amplified from pGL190 using the primer SorEd9 and 10. The amplified cassette was then inserted into

pGL206 through *Sfi*I digestion and subsequent ligation. To create pGL222, the partial backbone of

pPHP96564 (Che et al. 2022) and the *Wus2* expression cassette were amplified from pPHP96564 using the

primers SorEd11 and 12. The attR4-attR3 region was amplified from pPHP85425 (Aregawi et al. 2022)

using the primers SorEd13 and 14. The *hygromycin B phosphotransferase* gene was amplified from pGL206

using the primers SorEd15 and 16. The rest of the backbone of pPHP96564 was amplified with the primers

SorEd17 and 18. These fragments were reassembled into the final vector using Gibson assembly.

Additionally, the *ZsG* gene, driven by the *Sorghum bicolor ubiquitin1* promoter, was amplified from

pGL190 (Aregawi et al. 2022) with the primers SorEd19 and 20. The ZsG expression cassette was digested

by *Nhe*I and *Pme*I and ligated into the construct to finalize the construct, pGL222; it was validated using

Nanopore sequencing.

To optimize shuttle constructs, the sorghum U6.2.3 (SbU6.2.3) promoter (Massel et al. 2022) from

sorghum genotype RTx430 was amplified using the primers SorEd-21 and 22. pGL198 (Aregawi et al. 2022)

was linearized by primers, SorEd-23 and 24. The OsU3 promoter was replaced by SbU6.2.3 using Gibson

Assembly. Additionally, ZmCas9 was replaced by ZmCas9i using primers SorEd25 and 26 for vector

linearization and primers SorEd27 and 28 for ZmCas9i amplification from pPHP86564, resulting in the

shuttle vector, pGL220. Based on the polycistronic shuttle vector construction, the monocistronic shuttle

vector was created by inserting two gRNA expression cassettes by Gibson assembly with primers SorEd29-

36 to create the monocistronic expression cassettes. Cas9 and gRNA polycistronic and monocistronic

cassettes were transferred to pGL222 to create binary constructs pGL220 and 221 by LR recombination.

To create pGL224, we used the same strategy to replace the OsU3 promoter and ZmCas9 with the

SbU6.2.3 promoter and ZmCas9i as described above. To streamline construct creation, level 0 constructs

were developed. These constructs were created by fusing the gRNA scaffold with various snoRNA

promoters (including sorghum U6_2.3 [SbU6_2.3], sorghum U6_3.1 [SbU6_3.1], rice U3 [OsU3], maize U6

[ZmU6], wheat U3 [TaU3], and wheat U6 [TaU6]) using overlapping PCR with primers SorEd49 through 62.

Fused DNA fragments were cloned into a pJET1.2 vector using CloneJET PCR cloning kit (ThermoFisher

Scientific) and validated by Sanger sequencing. These level 0 constructs served as templates for creating

both polycistronic and monocistronic gRNA editing constructs (**Figure S2**).

To create the ZmSpRYi construct, pGL224 was used as the template, and a total of 14 nucleotide mutations

were introduced. The first set of mutations on the first exon of ZmCas9i targeted the removal of a BsaI

recognition site and the substitution of the alanine 61 with an arginine. These mutations were

incorporated using primers SorEd37 and SorEd38, enabling the amplification of two overlapping

fragments with primers SorEd39 and SorEd40. The overlapping fragments were assembled through

overlapping PCR. The modified sequence replaced the original sequence via BamHI and BspEI digestion,

followed by ligation. The remaining ten mutations were introduced using a synthetic oligonucleotide

chemically synthesized by Genewiz (http://genewiz.com). These mutations were inserted into the 3' end

of ZmCas9i through Gibson assembly. Primers SorEd41 and SorEd42 were used to amplify the insert, while

primers SorEd43 and SorEd44 were employed for vector linearization. Additionally, two unwanted BsaI

restriction sites were eliminated via primer-directed mutagenesis using primers SorEd45 through

SorEd48, resulting in the construct, pGL225 (**Figure S1**). Finally, the ZmCas9i gene in pGL220 and pGL221

was replaced with the ZmSpRYi gene to create pGL226 and pGL227, designed for Agrobacterium-mediated

transformation

Agrobacterium-mediated transformation

The binary constructs were introduced into an auxotrophic strain of *Agrobacterium tumefaciens* LBA4404

Thy- (Anand et al. 2017) by electroporation. Transformed Agrobacterium strains were plated on yeast

extract peptone (YEP) medium supplemented with 100 mg/L thymidine, 50 mg/L gentamicin, and 50 mg/L

spectinomycin. Details on the storage and preparation of the Agrobacterium strains follow a previously

published protocol (Masters et al. 2020). In summary, Agrobacterium stocks were streaked onto YEP

plates containing appropriate antibiotics to generate master plates. After a 2-day incubation at 28°C, 5–6

colonies from the master plates were selected to establish working cultures on YEP plates. Following

overnight incubation and just prior to inoculation of sorghum immature embryos, Agrobacterium

suspensions were prepared in PHI-I medium (Wu et al. 2014), with optical density (OD550 nm), adjusted

to 0.7; silwet L-77 (0.005% v/v) and acetosyringone (0.2 mM) were added to finalize the Agrobacterium

inoculation medium. Transformation frequency refers to the percentage of IEs used in the transformation
that successfully produce transformants.

Visualization of T-DNA integration

The ZsGreen reporter gene was used as a visible marker to track T-DNA integration and segregation. To
monitor T-DNA integration following Agrobacterium-mediated inoculation, transformed IEs were
examined weekly using a Lumar v12 epifluorescence stereo microscope and a QImaging Retiga SRV
camera (Zeiss, UC Berkeley Biological Imaging Facility). The response of IEs to WUS expression was
assessed by analyzing the number and size of fluorescent tissues. Prior to transferring shoots to rooting
medium, regenerants exhibiting fluorescence were selected for further genotyping and phenotyping.

Phenotypic characterization of T_0 transgenic plants

444 T_0 plantlets were phenotyped while growing in rooting medium. Each plantlet was scored for the
445 percentage of albino phenotype, with 100% indicating plants that were completely albino and 0%
446 indicating plants that were completely green. The phenotype score was compared with the knockout
447 percentage obtained using the Synthego Inference of CRISPR Edits (ICE) tool (Conant et al. 2022) to
448 determine correlation between the two observations.

Genotyping of editing events

To test putative T_0 transgenic lines for successful editing of target genes, genomic DNA from young
sorghum leaves was isolated using a urea buffer as described (Aregawi et al. 2022). Target regions of
potential edits were amplified using the primers SorEd63 and 64 and purified with the GeneJet PCR
Purification Kit (Thermo Scientific). Purified PCR products were sequenced with Sanger sequencing at the
UC Berkeley DNA Sequencing Facility. Sequencing data were analyzed using the Synthego ICE analysis tool
(Conant et al. 2022), and knockout scores, referring to proportion of indels that indicate a frameshift or
are more than 21 bp in length, were used to assess editing efficiency and the likelihood of transmission to
the next generation. The editing efficiency represents the percentage of plant cells within harvested
somatic samples that successfully undergo knockout events, as determined by the knockout score.

Acknowledgments

This project was supported by the Chan-Zuckerberg Initiative and the DOE EERC project through the Lawrence Livermore National Laboratory. PGL acknowledges support from the US Cooperative Extension Service through the Division of Agriculture and Natural Resources of the University of California. JS acknowledges support from USDA-ARS Crop Improvement and Genetics Unit Albany, CA and ORISE Research Participation Program. We thank Dr. William Gordon-Kamm at Corteva Agriscience for helpful conversations and for the constructs, pPHP96564 and pPHP85425. We thank Dr. Roger Thilmony at USDA-ARS Crop Improvement and Genetics Unit Albany for informative discussions. We thank talented undergraduate students, Malia Ariyoshi, Emma To, Eda Joy Ustuner and Mantra Rajkumar, for their help in this work. We are grateful for the help of the staff of the UC Berkeley Biological Imaging Facility and the UC Berkeley Oxford Tract Greenhouse Facility.

References

Altpeter, Fredy, Nathan M. Springer, Laura E. Bartley, Ann E. Blechl, Thomas P. Brutnell, Vitaly Citovsky, Liza J. Conrad, et al. 2016. "Advancing Crop Transformation in the Era of Genome Editing." *The Plant Cell* 28 (7): 1510–20.

Aregawi, Kiflom, Jianqiang Shen, Grady Pierroz, Manoj K. Sharma, Jeffery Dahlberg, Judith Owiti, and Peggy G. Lemaux. 2022. "Morphogene-Assisted Transformation of Sorghum Bicolor Allows More Efficient Genome Editing." *Plant Biotechnology Journal* 20 (4): 748–60.

Anand, A., Bass, S.H., Cho, H.-J., Klein, T.M., Lassner, M., McBridge, K.E. 2017. Methods and Compositions of Improved Plant Transformation. WIPO 2017/078836 Al. World Patent, filed August 26, 2016, and issued November 5, 2017.

Blomme, Jonas, Ward Develtere, Ayse Köse, Júlia Arraiza Ribera, Christophe Brugmans, Jessica Jaraba-Wallace, Ward Decaestecker, et al. 2022. "The Heat Is on: A Simple Method to Increase Genome Editing Efficiency in Plants." *BMC Plant Biology* 22 (1): 142.

Castel, Baptiste, Laurence Tomlinson, Federica Locci, Ying Yang, and Jonathan D. G. Jones. 2019. "Optimization of T-DNA Architecture for Cas9-Mediated Mutagenesis in Arabidopsis." *PloS One* 14 (1): e0204778.

Che, Ping, Emily Wu, Marissa K. Simon, Ajith Anand, Keith Lowe, Huirong Gao, Amy L. Sigmund, et al. 2022. "Wuschel2 Enables Highly Efficient CRISPR/Cas-Targeted Genome Editing during Rapid de Novo Shoot Regeneration in Sorghum." *Communications Biology* 5 (1): 344.

Conant, David, Tim Hsiau, Nicholas Rossi, Jennifer Oki, Travis Maures, Kelsey Waite, Joyce Yang, et al. 2022. "Inference of CRISPR Edits from Sanger Trace Data." *The CRISPR Journal* 5 (1): 123–30.

Dahlberg, J. A., J. J. Burke, and D. T. Rosenow. 2004. "Development of a Sorghum Core Collection: Refinement and Evaluation of a Subset from Sudan." *Economic Botany* 58 (4): 556–67.

Elzen, P. J. van den, J. Townsend, K. Y. Lee, and J. R. Bedbrook. 1985. "A Chimaeric Hygromycin
Resistance Gene as a Selectable Marker in Plant Cells." *Plant Molecular Biology* 5 (5): 299–302.
Ge, Zengxiang, Leiqi Zheng, Yuling Zhao, Jiahao Jiang, Emily J. Zhang, Tianxu Liu, Hongya Gu, and Li-Jia
Qu. 2019. "Engineered XCas9 and SpCas9-NG Variants Broaden PAM Recognition Sites to
Generate Mutations in Arabidopsis Plants." *Plant Biotechnology Journal* 17 (10): 1865–67.
Grützner, Ramona, Patrick Martin, Claudia Horn, Samuel Mortensen, Erin J. Cram, Carolyn W. T. Lee-
Parsons, Johannes Stuttmann, and Sylvestre Marillonnet. 2021. "High-Efficiency Genome Editing
in Plants Mediated by a Cas9 Gene Containing Multiple Introns." *Plant Communications* 2 (2):
100135.
Hibshman, Grace N., Jack P. K. Bravo, Matthew M. Hooper, Tyler L. Dangerfield, Hongshan Zhang, Ilya J.
Finkelstein, Kenneth A. Johnson, and David W. Taylor. 2024. "Unraveling the Mechanisms of
PAMless DNA Interrogation by SpRY-Cas9." *Nature Communications* 15 (1): 3663.
Hoerster, George, Ning Wang, Larisa Ryan, Emily Wu, Ajith Anand, Kevin McBride, Keith Lowe, Todd
Jones, and Bill Gordon-Kamm. 2020. "Use of Non-Integrating Zm-Wus2 Vectors to Enhance
Maize Transformation." *In Vitro Cellular & Developmental Biology. Plant: Journal of the Tissue
Culture Association* 56 (3): 265–79.
Hu, Johnny H., Shannon M. Miller, Maarten H. Geurts, Weixin Tang, Liwei Chen, Ning Sun, Christina M.
Zeina, et al. 2018. "Evolved Cas9 Variants with Broad PAM Compatibility and High DNA
Specificity." *Nature* 556 (7699): 57–63.
Jinek, Martin, Krzysztof Chylinski, Ines Fonfara, Michael Hauer, Jennifer A. Doudna, and Emmanuelle
Charpentier. 2012. "A Programmable Dual-RNA-Guided DNA Endonuclease in Adaptive Bacterial
Immunity." *Science (New York, N.Y.)* 337 (6096): 816–21.
Jores, Tobias, Jackson Tonnies, Travis Wrightsman, Edward S. Buckler, Josh T. Cuperus, Stanley Fields,
and Christine Queitsch. 2021. "Synthetic Promoter Designs Enabled by a Comprehensive
Analysis of Plant Core Promoters." *Nature Plants* 7 (6): 842–55.
LeBlanc, Chantal, Fei Zhang, Josefina Mendez, Yamile Lozano, Krishna Chatpar, Vivian F. Irish, and
Yannick Jacob. 2018. "Increased Efficiency of Targeted Mutagenesis by CRISPR/Cas9 in Plants
Using Heat Stress." *The Plant Journal: For Cell and Molecular Biology* 93 (2): 377–86.
Lee, Jeong Sun, Su-Ji Bae, Jin-Soo Kim, Changsoo Kim, and Beum-Chang Kang. 2023. "A Streamlined
Guide RNA Screening System for Genome Editing in Sorghum Bicolor." *Plant Methods* 19 (1): 90.
Liang, Zhen, Yuqing Wu, Lingling Ma, Yingjie Guo, and Yidong Ran. 2021. "Efficient Genome Editing in
Setaria Italica Using CRISPR/Cas9 and Base Editors." *Frontiers in Plant Science* 12: 815946.
Liu, Guoquan, Jieqing Li, and Ian D. Godwin. 2019. "Genome Editing by CRISPR/Cas9 in Sorghum through
Biolistic Bombardment." *Methods in Molecular Biology (Clifton, N.J.)* 1931: 169–83.
Lowe, Keith, Mauricio La Rota, George Hoerster, Craig Hastings, Ning Wang, Mark Chamberlin, Emily
Wu, Todd Jones, and William Gordon-Kamm. 2018. "Rapid Genotype 'Independent' Zea Mays L.
(Maize) Transformation via Direct Somatic Embryogenesis." *In Vitro Cellular & Developmental
Biology. Plant: Journal of the Tissue Culture Association* 54 (3): 240–52.
Lowe, Keith, Emily Wu, Ning Wang, George Hoerster, Craig Hastings, Myeong-Je Cho, Chris Scelonge, et
al. 2016. "Morphogenic Regulators Baby Boom and Wuschel Improve Monocot Transformation."
The Plant Cell 28 (9): 1998–2015.
Mann, David G. J., Peter R. Lafayette, Laura L. Abercrombie, Zachary R. King, Mitra Mazarei, Mathew C.
Halter, Charleson R. Poovaiah, et al. 2012. "Gateway-Compatible Vectors for High-Throughput
Gene Functional Analysis in Switchgrass (Panicum Virgatum L.) and Other Monocot Species."
Plant Biotechnology Journal 10 (2): 226–36.
Massel, Karen, Yasmine Lam, Jessica Hintzsche, Nicholas Lester, Jose R. Botella, and Ian D. Godwin. 2022.
"Endogenous U6 Promoters Improve CRISPR/Cas9 Editing Efficiencies in Sorghum Bicolor and
Show Potential for Applications in Other Cereals." *Plant Cell Reports* 41 (2): 489–92.

Masters, Alicia, Minjeong Kang, Morgan McCaw, Jacob D. Zobrist, William Gordon-Kamm, Todd Jones, and Kan Wang. 2020. "Agrobacterium-Mediated Immature Embryo Transformation of Recalcitrant Maize Inbred Lines Using Morphogenic Genes." *Journal of Visualized Experiments: JoVE*, no. 156 (February): e60782.

Milner, Matthew J., Melanie Craze, Matthew S. Hope, and Emma J. Wallington. 2020. "Turning up the Temperature on CRISPR: Increased Temperature Can Improve the Editing Efficiency of Wheat Using CRISPR/Cas9." *Frontiers in Plant Science* 11 (November): 583374.

Milner, Matthew J., Manisha Sharma, Ruth E. Bates, Michelle Whiting, Melanie S. Craze, Peter Miller, Jack Brooks, Allan Kouidri, and Emma J. Wallington. 2024. "Differential Editing Efficiencies in Cereal Crops: A Comparative Analysis of TRNA and Ribozyme Multiplexed Guide Delivery." *Frontiers in Plant Science* 15 (December): 1426184.

Ming, Meiling, Qiurong Ren, Changtian Pan, Yao He, Yingxiao Zhang, Shishi Liu, Zhaohui Zhong, et al. 2020. "CRISPR-Cas12b Enables Efficient Plant Genome Engineering." *Nature Plants* 6 (3): 202–8.

Nishimasu, Hiroshi, Xi Shi, Soh Ishiguro, Linyi Gao, Seiichi Hirano, Sae Okazaki, Taichi Noda, et al. 2018. "Engineered CRISPR-Cas9 Nuclease with Expanded Targeting Space." *Science (New York, N.Y.)* 361 (6408): 1259–62.

Patel-Tupper, Dhruv, Armen Kelikian, Anna Leipertz, Nina Maryn, Michelle Tjahjadi, Nicholas G. Karavolias, Myeong-Je Cho, and Krishna K. Niyogi. 2024. "Multiplexed CRISPR-Cas9 Mutagenesis of Rice PSBS1 Noncoding Sequences for Transgene-Free Overexpression." *Science Advances* 10 (23): eadm7452.

Paterson, Andrew H., John E. Bowers, Rémy Bruggmann, Inna Dubchak, Jane Grimwood, Heidrun Gundlach, Georg Haberer, et al. 2009. "The Sorghum Bicolor Genome and the Diversification of Grasses." *Nature* 457 (7229): 551–56.

Ramstein, Guillaume P., Sarah E. Jensen, and Edward S. Buckler. 2019. "Breaking the Curse of Dimensionality to Identify Causal Variants in Breeding 4." *Theoretical and Applied Genetics* 132 (3): 559–67.

Ren, Bin, Lang Liu, Shaofang Li, Yongjie Kuang, Jingwen Wang, Dawei Zhang, Xueping Zhou, Honghui Lin, and Huanbin Zhou. 2019. "Cas9-NG Greatly Expands the Targeting Scope of the Genome-Editing Toolkit by Recognizing NG and Other Atypical PAMs in Rice." *Molecular Plant* 12 (7): 1015–26.

Ren, Qiurong, Simon Sretenovic, Shishi Liu, Xu Tang, Lan Huang, Yao He, Li Liu, et al. 2021. "PAM-Less Plant Genome Editing Using a CRISPR-SpRY Toolbox." *Nature Plants* 7 (1): 25–33.

Rodríguez-Leal, Daniel, Zachary H. Lemmon, Jarrett Man, Madelaine E. Bartlett, and Zachary B. Lippman. 2017. "Engineering Quantitative Trait Variation for Crop Improvement by Genome Editing." *Cell* 171 (2): 470-480.e8.

Schindele, Angelina, Annika Dorn, and Holger Puchta. 2020. "CRISPR/Cas Brings Plant Biology and Breeding into the Fast Lane." *Current Opinion in Biotechnology* 61 (February): 7–14.

Strecker, Jonathan, Sara Jones, Balwina Koopal, Jonathan Schmid-Burgk, Bernd Zetsche, Linyi Gao, Kira S. Makarova, Eugene V. Koonin, and Feng Zhang. 2019. "Engineering of CRISPR-Cas12b for Human Genome Editing." *Nature Communications* 10 (1): 212.

Tang, Xu, Levi G. Lowder, Zhang Tao, Aimee A. Malzahn, Xuelian Zheng, D. Voytas, Zhaohui Zhong, et al. 2017. "A CRISPR–Cpf1 System for Efficient Genome Editing and Transcriptional Repression in Plants." *Nature Plants* 3 (February). https://doi.org/10.1038/nplants.2017.18.

Varoquaux, Nelle, Benjamin Cole, Cheng Gao, Grady Pierroz, Christopher R. Baker, Dhruv Patel, Mary Madera, et al. 2019. "Transcriptomic Analysis of Field-Droughted Sorghum from Seedling to Maturity Reveals Biotic and Metabolic Responses." *Proceedings of the National Academy of Sciences of the United States of America* 116 (52): 27124–32.

Walton, Russell T., Kathleen A. Christie, Madelynn N. Whittaker, and Benjamin P. Kleinstiver. 2020.
"Unconstrained Genome Targeting with Near-PAMless Engineered CRISPR-Cas9 Variants."
Science (New York, N.Y.) 368 (6488): 290–96.
Wu, Emily, Brian Lenderts, Kimberly Glassman, Maya Berezowska-Kaniewska, Heather Christensen,
Tracy Asmus, Shifu Zhen, Uyen Chu, Myeong-Je Cho, and Zuo-Yu Zhao. 2014. "Optimized
Agrobacterium-Mediated Sorghum Transformation Protocol and Molecular Data of Transgenic
Sorghum Plants." *In Vitro Cellular & Developmental Biology. Plant: Journal of the Tissue Culture
Association* 50 (1): 9–18.
Xu, Ziyan, Yongjie Kuang, Bin Ren, Daqi Yan, Fang Yan, Carl Spetz, Wenxian Sun, Guirong Wang, Xueping
Zhou, and Huanbin Zhou. 2021. "SpRY Greatly Expands the Genome Editing Scope in Rice with
Highly Flexible PAM Recognition." *Genome Biology* 22 (1): 6.
Zetsche, Bernd, Jonathan S. Gootenberg, Omar O. Abudayyeh, Ian M. Slaymaker, Kira S. Makarova,
Patrick Essletzbichler, Sara E. Volz, et al. 2015. "Cpf1 Is a Single RNA-Guided Endonuclease of a
Class 2 CRISPR-Cas System." *Cell* 163 (3): 759–71.
Zhan, Xiangqiang, Yuming Lu, Jian-Kang Zhu, and Jose Ramon Botella. 2021. "Genome Editing for Plant
Research and Crop Improvement." *Journal of Integrative Plant Biology* 63 (1): 3–33.
Zhang, Yingxiao, Aimee A. Malzahn, Simon Sretenovic, and Yiping Qi. 2019. "The Emerging and
Uncultivated Potential of CRISPR Technology in Plant Science." *Nature Plants* 5 (8): 778–94.
Zhong, Zhaohui, Simon Sretenovic, Qiurong Ren, Lijia Yang, Yu Bao, Caiyan Qi, Mingzhu Yuan, et al. 2019.
"Improving Plant Genome Editing with High-Fidelity XCas9 and Non-Canonical PAM-Targeting
Cas9-NG." *Molecular Plant* 12 (7): 1027–36.
Zhou, Jianping, Guanqing Liu, Yuxin Zhao, Rui Zhang, Xu Tang, Ling Li, Xinyu Jia, et al. 2023. "An Efficient
CRISPR-Cas12a Promoter Editing System for Crop Improvement." *Nature Plants* 9 (4): 588–604.
Zhu, Haocheng, Chao Li, and Caixia Gao. 2020. "Applications of CRISPR-Cas in Agriculture and Plant
Biotechnology." *Nature Reviews. Molecular Cell Biology* 21 (11): 661–77.
613

Author contributions

Jianqiang Shen designed the experiments, developed the constructs, conducted sorghum
transformations, and contributed to writing and editing. Kiflom Aregawi contributed to discussions on
experimental design, performed sorghum transformations, and contributed to writing and editing.
Sultana Anwar conducted sorghum transformations and contributed to editing. Tamara Miller contributed
to writing and editing. Peggy G. Lemaux supervised the project, secured funding, and contributed to
editing.

Competing interest statement

623 pGL and pPHP vectors are available for research purpose; using the pPHP vectors for commercial
624 applications requires a paid non-exclusive license from Corteva Agriscience. Manuscript authors have no
625 relationship with Corteva Agriscience.

Figure 1. Workflow for construct creation and construct maps.

(A) Diagram illustrating the workflow for construct creation and usage.

(B) Maps of shuttle vectors. Polycistronic shuttle vector contains a polycistronic gRNA transcription unit with tRNA spacers; monocistronic shuttle vector carries monocistronic gRNA transcription units driven by Pol III promoters. Rectangles with arrows indicate promoters. Other rectangles indicate exons (dark-colored) or terminators (light-colored). Black horizontal lines indicate introns. Grey vertical lines indicate Gateway recombination sites.

(C) Maps of binary vectors. pGL206 represents the basic binary vector backbone. pGL216 is designed for testing the function of morphogenic genes in enhancing transformation frequencies. pGL222 is a binary

vector backbone used in genome editing applications. Rectangles with arrows indicate promoters. Solid

rectangles indicate exons (dark-colored) or terminators (light-colored). Light gray rectangles denote

buffer regions; dark gray rectangles represent *ccdB* genes. Black vertical lines indicate T-DNA borders; gray

vertical lines indicate Gateway recombination sites.

Figure 2. Workflow for sorghum immature embryo transformation.

(A) Diagram illustrating the transformation workflow, from immature embryo isolation through rooted

regenerants.

(B-D) Sorghum tissues at various stages post-Agrobacterium infection. **(B)** Immature embryos one week

after infection; **(C)** two weeks after infection and **(D)** four weeks after infection, visualized with overlays

of brightfield and fluorescence images captured with a Zeiss AxioZoom V16 low-magnification

epifluorescence microscope. Scale bars indicate 2 mm.

 (E) Sorghum tissue four weeks after Agrobacterium infection, with two weeks of exposure to hygromycin

650   selection. Scale bars indicate 2 mm.

652
Figure 3: Wus2-assisted CRISPR/*Cas9* editing of *SbPDS* gene.

(A) Schematic representation of the structure of *SbPDS*, depicting *SbPDS* exons (green boxes) and gRNA target

sequences.

(B) T$_0$ edited plants showing chimeric (left) and complete (right) albino phenotypes.

(C) Violin plot showing the distribution of knockout scores obtained from Sanger sequencing and Synthego ICE

analysis.

(D) Sequencing results of targeted regions of *SbPDS* gene in five representative T$_0$ transgenic plants. gRNAs are

underlined, PAM sequences are shown in green, and green triangles indicate cut sites. Dashed lines indicate

deletions; insertion is shown in red.

C

			Editing
	gRNA1.1	gRNA2.2	efficiency
gRNAs	GAGGCAAGAGATGTTTTGGGCGG...GATGGGGATTGGTACGAGACTGG		
PDS-plc-12b	GAGGCAAGAGATGTTTTGGGCGG...GATGGGGATTGGTACGA_ _CTGG		0%; 27%
PDS-plc-23c	GAGGCAAGAGATGTTTTGGGCGG...GATGGGGATTGGTACGA_ _CTGG		0%; 100%
PDS-plc-41b	GAGGCAAGAGATGTTTTGGGCGG...GA_________________CTGG		0%; 36%
PDS-mnc-5b	GAGGCAAGAGATGTTTTGGGCGG...GATGGGGATTGGTAC_ _GACTGG		0%; 50%
PDS-mnc-28a	GAGGCAAGAGATGTTTTGGGCGG...GATGGGGATTGGTAC________		0%; 53%
PDS-mnc-31a	GAGGCAAGAGATGTTTTGGGCGG...GATGGGGATTGGTACG_GACTGG		0%; 30%

1.1 2.2
gRNA

D

			Self editing
	gRNA1.1	gRNA2.2	efficiency
gRNAs	GAGGCAAGAGATGTTTTGGGGTT...GATGGGGATTGGTACGAGACGTT		
PDS-plc-23c	GAGGCAAGAGATGTTTTGGGGTT...GATGGGGATTGGTAC_ _GACGTT		0%; 31%
PDS-mnc-5b	GAGGCAAGAGATGTTTTGGGGTT...GATGGGGATTGGTAC_ _GACGTT		0%; 16%
PDS-mnc-21a	GAGGCAAGAGATGTTT_GGGGTT...GATGGGGATTGGTAC_ _GACGTT		16%; 24%
PDS-mnc-22a	GAGGCAAGAGATGTTTTGGGGTT...GATGGGGATTGGTAC_ _GACGTT		0%; 30%

1.1 2.2
gRNA

Figure 4. Genome editing in sorghum mediated by CRISPR/SpRY

(A) Diagram illustrating self-editing and genome editing of target sites. Blue rectangles represent exons; gray rectangles denote untranslated regions (UTRs). Gray lines on *ZmSpRYi* indicate mutations. Dark lines above the exons indicate gRNAs. Arrows indicate gene expression or CRISPR/SpRY-mediated editing.

(B) Phenotypes of PDS gene-edited plants. The left panel shows a non-edited plantlet, the middle panel shows a fully albino plantlet, and the right panel shows a chimeric albino plantlet. Images were taken four weeks after initiation of rooting. Scale bar = 2 cm.

(C) Genotypes of edited *PDS* gene exons. Underlined sequences indicate gRNA core sequences, triangles

represent potential double-strand break sites, and blue-colored nucleotides indicate PAM sequences.

Editing efficiency refers to knockout scores in somatic tissues that were genotyped.

(D) Representative examples of self-editing in gRNA expression cassettes of integrated T-DNA. Underlined

sequences indicate gRNA core sequences, triangles represent potential double-strand break sites, and

blue-colored nucleotides indicate PAM sequences. Editing efficiency is based on knockout scores from

genotyping of somatic tissues.

(E) Violin plot showing distribution of knockout scores for *PDS* gRNA2.2 edits mediated by CRISPR/*SpRY*,

comparing polycistronic and monocistronic gRNA transcription units. For polycistronic units, n = 46; for

monocistronic units, n = 49.

Table 1. Transformation, editing and knockout efficiencies of Wus2-assisted transformation based on one
plant per IE

gRNA system (Construct)	Temp (°C).	Replication	# IEs used	# Regenerants	# Escapes (%) [†]	+ for hph [‡]	**Transformation frequency (%)** [§]	# edited plants	**Editing efficiency (%)** [¶]	# plants with Knockout [††]	**Knockout efficiency (%)** [‡‡]
Polycistronic (pGL220)	28	1st	90	40	2(0.05)	38	**42.2**	36	**94.7**	35	**92.0**
		2nd	132	47	6(0.12)	41	**31.0**	24	**58.5**	24	**58.5**
	25	1st	60	26	0(0.00)	26	**43.3**	23	**88.4**	22	**84.6**
Monocistronic (pGL221)	28	1st	93	24	1(0.04)	23	**24.7**	23	**100**	21	**91.3**
		2nd	112	28	1(0.03)	27	**24.1**	27	**100**	27	**100.0**
	25	1st	105	33	1(0.03)	33	**31.4**	33	**100**	33	**100.0**

[†] # Escapes (%) = number and percentage of regenerated plants without transgene (T-DNA) = (#Escapes/# Regenerant) X 100

[‡] + = plants PCR-positive for *hph*

[§] Transformation frequency (%) = (# + for *hph* divided by # IEs used) x 100

[¶] Editing efficiency (%) = (# edited plants divided by number of plants + for hph) x 100

[††] # plants with knockout = number of plants with frameshift mutation and albino phenotype

[‡‡] Knockout efficiency(%) = (# plants with knockout divided by + for hph) x 100

Table 2. Transformation, editing and knockout efficiencies of Wus2-assisted transformation based on
multiple regenerated plants from each IE

gRNA system (Construct)	Temp (°C).	Replications	# IEs used	# Regenerants	# Escapes (%)[†]	+ for hph[‡]	**Transformation frequency (%)**[§]	# edited plants	**Editing efficiency (%)**[¶]	# plants with Knockout[††]	**Knockout efficiency (%)**[‡‡]
Polycistronic (pGL220)	28	1st	90	82	6 (0.07)	76	**84.4**	68	**89.5**	58	**76.3**
		2nd	132	60	6 (0.1)	54	**40.9**	30	**55.5**	30	**55.5**
	25	1st	60	61	1 (0.02)	60	**100**	51	**85**	48	**80**
Monocistronic (pGL221)	28	1st	93	64	1 (0.02)	63	**67.7**	62	**98.4**	60	**95.2**
		2nd	112	77	3 (0.04)	74	**66.1**	66	**89.2**	63	**85.1**
	25	1st	105	93	6 (0.06)	87	**82.9**	87	**100.0**	84	**96.6**

[†] # Escapes (%) = Number and percentage of regenerated plants without transgene (T-DNA) = (#Escapes/# Regenerant) X 100

[‡] + indicates plants PCR-positive for hph

[§] Transformation frequency (%) = (+ for hph divided by # IEs used) x 100

[¶] Editing efficiency (%) = (# edited plants divided by number of plants + for hph) x 100

[††] # plants with knockout = Number of plants with frameshift mutation and albino phenotype

[‡‡] Knockout efficiency(%) = (# plants with knockout divided by # + for hph) x 100

Table 3. Transformation, editing and knockout efficiencies of *SpRY* using *Wus2*-assisted transformation
strategy

gRNA system (Construct)	Replications	# IEs used	# Regenerants	# Escapes (%)[†]	+ for hph[‡]	**Transformation frequency (%)**[§]	# edited plants	**Editing efficiency (%)**[¶]	# plants with Knockout[††]	**Knockout efficiency (%)**[‡‡]
Polycistronic (pGL226)	1st	107	46	3	43	**40.2**	11	**25.6**	11	**25.6**
	2nd	60	20	3	17	**28.3**	6	**35.3**	6	**35.3**
Monocistronic (pGL227)	1st	119	49	0	49	**41.2**	40	**81.6**	40	**81.6**
	2nd	82	43	3	40	**48.8**	37	**92.5**	37	**92.5**

[†] # Escapes (%) = number and percentage of regenerated plants without transgene (T-DNA) = (#Escapes/# Regenerant) X 100

[‡] + indicates plants PCR-positive for hph

[§] Transformation frequency (%) = (+ for hph divided by # IEs used) x 100

[¶] Editing efficiency (%) = (# edited plants divided by number of plants + for hph) x 100

[††] # plants with knockout = number of plants with frameshift mutations and albino phenotypes

[‡‡] Knockout efficiency(%) = (# plants with knockout divided by plants + for hph) x 100