Identification of Specialized tRNA Expression in Early Human Brain Development

Alex L. Bagi[1,2] (abagi@ucsc.edu), Todd M. Lowe[1] (tmjlowe@ucsc.edu), Sofie R. Salama[1,2,3] (ssalama@ucsc.edu)

[1] Department of Biomolecular Engineering, University of California Santa Cruz, Santa Cruz, California, USA.
[2] UC Santa Cruz Genomics Institute, University of California Santa Cruz, Santa Cruz, California, USA.
[3] Department of Molecular, Cell and Developmental Biology, University of California Santa Cruz, Santa Cruz, California, USA.

Lead Contact

Sofie R. Salama (ssalama@ucsc.edu)

Corresponding author

Todd M. Lowe (tmjlowe@ucsc.edu)
Sofie R. Salama (ssalama@ucsc.edu)

Abstract

Transfer RNAs (tRNAs) are traditionally known for their role in protein translation, yet recent discoveries highlight their broader functions in gene regulation, particularly through tRNA-derived small RNAs (tDRs). Studies have shown the singular importance of one unique ArgUCU tRNA isodecoder in mouse neural development, yet potential function(s) of tDRs derived from this and all other tRNAs remain largely unexplored in early human brain development. In this study, we employed cerebral cortical organoid models and AlkB-facilitated RNA methylation sequencing (ARM-seq) to profile tDRs across distinct stages of early human cerebral cortex development. Our analysis reveals dynamic expression patterns of diverse tDR groups derived from a wide range of isodecoders, with several distinct groups showing neural-specific expression. Computational analyses of these tDRs shows biased sequence motifs in over-represented tRNAs that are enriched with particular RNA modifications, giving initial clues to traits that change within the pool of tDRs during neural development. This expanded catalog of tDRs provides a framework for future studies on tRNA function in brain development and offers a deeper understanding of the complexity of tDR dynamics in neural differentiation.

Keywords

transfer RNA, brain organoid, neurodevelopment, tRNA fragments, RNA modification, RNA processing, pluripotent stem cells, comparative transcriptomics,

Introduction

A growing body of evidence suggests that tRNAs and their derived small RNAs (tDRs) play important regulatory roles that go beyond protein translation. Some of these extra-translational roles include global regulation of RNA silencing, immune system dysregulation, early developmental and cancer progression regulation.[1–5] tRNA-derived small RNAs (tDRNAs or tDRs) in particular have also been demonstrated to have many roles in regulation and disease.[6–9] Some specific angiogenin-induced tDRs also have the ability to form into G-quadruplexes, which have neuroprotective roles in motor neurons, affecting the pathogenesis of diseases such as amyotrophic lateral sclerosis.[10,11] Regulatory roles for tDRs are also poorly understood for a variety of novel mecianisms such as the regulation of noncoding RNA biogenesis and gene repression via protein interaction from one specific tDR-Gly-GCC.[12] Specific genes also highlight the complex interplay between tRNA modification and function, suggesting links between intellectual disability and translational efficiency of specific codons.[13] Further, the bioavailability of specific amino acids such as arginine, glycine, and selenocysteine have been known to play roles in neurological function, disorders, and development, which may be influenced by the expression of specific tRNAs and their associated tDRs.[14–16]

Spatial- and time-dependent expression of protein-coding genes is an essential function for cell patterning and differentiation of neural tissues in mammalian neurodevelopment, but the roles of small RNAs have only recently been explored. Recent studies have found several instances where tDRs may play roles in neurological diseases (reviewed in Qin et al. [17]). A landmark study found that a specific tRNA (tRNA Arg-TCT-4-1; *n-Tr20* in mouse) is abundant almost exclusively in the brain, and a single point mutation could lead to severe post-natal neurodegeneration.[18] The same tRNA has been shown that loss of the mature leads to changes

in seizure susceptibility and mTORC1 suppression leading to altered neurotransmission.[19] Except for this singular tRNA in mouse, no known cytosolic tRNAs have been shown to play a specialized role(s) in neuro-development or mature brain function. Notably, there are no studies of tDRs derived from this specialized tRNA that may also be part of its unique role in neurological tissues.

Because of the dearth of data on tDRs associated with neural development, we employed AlkB-facilitated RNA methylation sequencing (ARM-seq) [20] to enable comprehensive analysis of these small RNAs. This specialized RNA sequencing method uses the *Escherichia coli* dealkylating enzyme AlkB, to remove common tRNA modifications (primarily m^1A, m^1G, m^3C) that can cause reverse transcription to terminate before reaching the end of an RNA transcript. This and other similar methods, such as YAMAT-seq [21], hydro-seq [22], and QuantM-tRNA-seq [23], yield a much more complete picture of tDRs than standard small RNA sequencing methods. Furthermore, with a specialized tRNA analysis pipeline such as tRAX (tRNA Analysis of eXpression) [24], tDRs can be sorted into more complex categories than the common 5' and 3' halves and fragments distinction.

We applied ARM-seq to cerebral cortex brain organoids, an established model for early events in fetal brain development [25–28], to profile tDRs across various stages of pluripotent cells to formation of excitatory projection neurons of the developing cortical plate.[29–31] Characterization in this unique system enabled the identification of previously unknown brain-associated tDRs expressed during human cortical neurogenesis. We observed that, in addition to specific tDRs derived from the previously studied brain-specific Arg-UCU tRNA, a much wider array of tDRs exhibited changes, both in abundance and processing patterns. To further explore these dynamics, we employed dimensionality reduction, clustering, and classification techniques to group these tDRs based on fragment properties. Among the

identified classes including Arg-UCU tDRs, other highly expressed tDRs derived from selenocysteine, alanine, and glycine, collectively show sequence conservation at a number of specific positions.

Results

Validation of organoid model and RNA profiling

Human induced pluripotent stem cells (GM12878-c305 iPSCs) were grown into cerebral cortical organoids as a model to study tDR changes in early human embryogenesis. A standard time course of culturing conditions was used to achieve neural induction and organoid growth **(Figure 1A)**. Cells were aggregated into embryoid bodies and differentiated into cortical tissues over a period of 10 weeks (see Methods) to span the emergence of key cell types in the course of dorsal cortex development [29], sampled at four time points: day 0 (D0 - stem cells); day 14 (D14 - neural epithelium and radial glia neural progenitor cells), day 35 (D35 - emergence of deep layer neurons) and day 70 (D70 - continued generation of cortical projection neurons, emergence of outer radial glia). D70 organoids expressed expected neural cell-type markers using immunofluorescence staining (IF-staining) when compared against D0 stem cells **(Figure 1B-D)**. Circular structures known as neural rosettes (early neural tube-like formations) were observed throughout the perimeter of the organoid stained with antibodies for PAX6 (a radial glia neural stem cell marker), and vimentin (a radial glial fiber marker), confirming the presence of radial glia neural stem cells **(Figure 1B)**. Furthermore, these PAX6+ neural rosettes were surrounded by cells staining for TBR2 (an intermediate neural progenitor marker), and CTIP2 (a deep layer neuron marker) **(Figure 1D)**. The presence of different neural subtypes increased over time as the organoids developed and increased in size, validating expected neural organoid morphology.

To observe high-resolution tDR dynamics in this neural development model, we performed ARM-seq small RNA sequencing [20], followed by differential expression analysis using tRNA Analysis of eXpression (tRAX) [24] for samples representing the differentiation time course (D0, D14, D35, D70). We validated that our small RNA sequencing reflected neural differentiation with multiple classes of small RNA markers known to have specific expression in neurons or stem cells **(Figure 1E)**. Stem-associated mir302 family miRNAs were more highly expressed in D0 stem cells, whereas neural-associated miR-9 and miR-LET7 (miR-99a) family miRNAs were more highly expressed in the D70 organoids **(Table S1)**.[32–35] MT-RNR2, a mitochondrial ribosomal RNA found to play a neuroprotective role, is also upregulated in organoids relative to stem cells.[36–38] Many SNORD115 small nucleolar RNAs, known to be induced by neuronal differentiation [39], were also highly expressed in neural cell time points, confirming expected small RNA transcriptional changes during the organoid time course experiment. Principal Components Analysis (PCA) of small RNA read counts across all time points **(Figure 1F)** confirmed that biological replicates and adjacent time points cluster closest to one another.

Broad changes in tRNA Isotypes do not capture tDR-Specific Changes

The relative expression of the major classes of small RNAs was assessed across the developmental time course, with ARM-seq picking up many small RNAs in addition to tRNAs, and tRNA reads increasing in abundance with AlkB treatment **(Figure 2A; Figure S1)**. tDRs derived from Ala, Asn, iMet, Phe, SeC, and Ser showed greater than two-fold increases between D0 and D70 **(Figure 2B),** with the greatest expression change between D0 and D35. In contrast, more than 40% of Glu, His, Lys, Thr, and Tyr tDRs decreased over the full time

course, with His and Tyr showing the most immediate decreases between D0 and D35 conditions.

Relative fragment profiles and their expression as a function of time were dynamic. Gly, Thr, and Val tDRs showed changes in expression and diversity in tDRs between isodecoders as well as between developmental time points **(Figure 2C)**. In Gly a 5' fragment is evident in D0 samples, and this fragment diminishes over time, although at different rates in the three isodecoders. However, in Gly-CCC and to a lesser extent in Gly-GCC, a 3' fragment increases in expression as the cells differentiate. In the Thr isotype, a 3' fragment shape appears in all cases, however, both Thr-AGU and Thr-UGU isodecoders decrease in expression and share a decrease in coverage at position 58. In contrast, Thr-CGU isodecoders increase in expression over time and have coverage beyond position 58, suggesting that modifications at this position may play a role in the processing and abundance of this fragment. Val tRNA isotypes have the widest variety of tDR coverage profiles with a distinct pattern for each time point and isodecoder. In addition to Gly, Thr, and Val, other isotypes also had distinct relative fragment profiles among their isodecoders, but the changes in the relative expression of their codon pools were less dynamic **(Figure S2A).**

Expression patterns of neural tDRs are unique and change over time

Individual tDRs were chosen to be further assessed rather than isodecoder groups in order to find neural-specific fragments and their dynamic expression patterns **(Table S2)**. tDR reads were divided into five main subgroups consisting of whole tRNA reads (within 10 bases of the start and of the full tRNA sequence), 5' fragments (within 10 bases of the start position but do not reach end), 3' fragments (within 10 bases of end position, but do not reach start), other fragments (internal fragments), and pre-tRNAs (pre-partial tRNAs and pre-tRNAs). The

expression of D0 vs. D70 tDRs was compared for all tDRs and all pre-tRNA type reads **(Figure 3A)**. Between D0 and D70 samples, 27 tDRs were found to have a significant (p-value <= 0.05) and $\log_2$(fold-change) greater than 1.5 increase in expression in the cerebral cortex organoids relative to undifferentiated stem cells **(Table S2)**. The differences between D0 and D14 samples were far less significant, likely since neural induction has just started and the space between time points is much closer. As such, tDR changes after neural differentiation were mostly examined between D0/D35 and D0/D70 time points. Among the top rank-expressed neural tDRs, several amino families appeared multiple times: Ala (4), Arg (3), Gly (2), and Leu (3).

Nearly all tDRs that increase after neural differentiation are 3' end-counts (3' fragments that terminate at, or within 10 bp of the CCA tail), with several pre-tRNA counts also being detected. The top 15 neural- and top 15 stem-expressed tDRs between D0 and D70 were compared **(Figure 3B)** to determine overall significance and expression **(Figure S3A,B)**. While 5' end-counts (5' fragments that terminate within 10 bp of the tRNA start site) are less common when compared to 3' end-counts, they tend to appear in D0 samples and diminish entirely after neural differentiation in both D35 and D70 time points. With many 3' tDRs being expressed at D35 and D70, there is also great diversity in the shape of fragment coverage with drop-offs of coverage at different conserved tRNA positions. This suggests differential modifications of the mature tRNA the tDR is derived from since mature tRNA modifications often act as signals for cleavage creating tDRs.[40]

The diversity of the top three most neural-expressed tDR 3' fragments was explored in detail **(Figure 3C)**. While coverage drop-off commonly occurred near position 58, other coverage changes differed between tRNA genes. tRNA-SeC-TCA-1 and tRNA-Arg-TCT-4 derived tDRs, for example, have a unique coverage drop through the anticodon loop, making these tDRs match more traditional 3' half designations. Additionally, while both tRNA-Arg-TCT-4

and tRNA-Arg-TCT-1 increase in abundance over time, the former has a severe drop-off in coverage around position 40, the later persists into the D-arm before dropping off **(Figure 3C)**. tRNA-Ala-AGC-8-derived tDRs, however, had distinct drops in coverage between the anticodon and T-loop with residual coverage encompassing the remainder of the tRNA. tRNA-Gly-GCC-2- and tRNA-Gly-CCC-2-derived tDRs also had distinct drops in coverage and appeared in the top 10 most neural-expressed tDRs, in contrast to tRNA-Gly-CCC-1 (the most stem-expressed tDR) **(Figure 3D)**. Interestingly both neural-specific Gly tDRs (tRNA-GCC-2/CCC-2) were 3' specific whereas tRNA-Gly-CCC-1 appeared to be highly 5' biased with dropoff before position 58. tRNA-Sec-TCA-1, tRNA-Arg-TCT-4, and tRNA-Ala-AGC-8 were validated as increasing in full-length tRNA expression across neural differentiation via Northern blots **(Figure S3D)**. Most tDRs maintain the same fragment coverage over the time course with changes in relative abundance rather than the fragment that is expressed changing.

Additional tRNAs showed pronounced changes in the tDR fragments generated over the developmental time course. tRNA-Asp-GTC-1-derived tDRs look similar in shape and expression in the first two time points but gain a large increase in 3' fragment expression in the last two **(Figure 3E)**. tRNA-Gly-GCC-1-derived tDRs have a contrasting pattern with the diminishment of the 5' end tDR coverage as the organoids reach the last two time points. tRNA-Pro-TGG-2 tDRs exhibit a mix of both patterns, with a decrease in 5' fragments paired with an increase in 3' fragments over time. Since excitatory neuron formation increases between D35 and D70, 3' fragments may be associated with these neurons. tRNA-Arg-TCT-2 and tRNA-Ser-GCT-5 tDRs form a unique group that show higher expression in D0 and D70 relative to D14 and D35, which may suggest lower expression of these 3' tDRs in neural progenitor cells **(Figure S3E)**.

Multidimensional Analysis Reveals tDR Clusters Associated with Neural vs. Stem States

To assess the full diversity of unique tRNA expression, clustering and classification of fragment profiles split across different time points was performed. This was done by aggregating read coverage, read-starts/ends, deletions, and misincorporation information (suggesting modified positions) and clustering the combined dataset using UMAP (Uniform Manifold Approximation and Projection) followed by classification via HDBSCAN.[41,42] The resulting clusters were then manually annotated into neural and stem categories as described in the methods. This revealed a highly complex, diverse set of tRNAs, each comprising multiple isotypes, and neural-stem expression levels with no bias in time point observed **(Figure 4A)**. Three major groups of HDBScan clusters were identified (consisting of 21 specific subgroups) labeled A (stem), B (neural), and C (neutral) as well as the HDBSCAN unannotated category (U) for tRNAs that were too hard to class into a single cluster. Spatially clusters tended to correlate with isotype and neural/stem favored tRNAs with no observable bias in time point. We next explored the relationship between isotype and cluster to see if genomic traits could also be correlated to tDR expression.

The clusters were often dominated by specific tRNA isotypes with several clusters containing greater than 50% of specific isotypes: A1 (Glu), A2 (Gly), A3 (Gln), C1 (Lys), C2 (Asp), C3 (Leu), C5 (Val), C6 (His), C7 (Gly), C8 (Ser), C12 (Leu), C14 (Arg), B1 (Ala), B2 (Arg), and B3 (Ala) **(Figure 4B)**. The clusters often have distinct isotype **(Figure S4A)**, AlkB treatment **(Figure S4B)**, and neural expression patterns **(Figure S4D)** defined by their sequencing characteristics making the uniquely identifiable. Cluster C10 and C11 are exceptions to this both expressing very diverse isotype compositions and seeming to correlate more closely with the

unannotated (U) category. This can likely be explained by low mean read coverage in clusters C10 and C11 relative to other clusters making them hard to annotate **(Table S3)**. AlkB conditions were generally diverse with a slight preference for AlkB+ conditions in clusters with higher neural expression. We also checked subtle genomic features of tRNAs such as tRNAscanSE covariance, HMM, and secondary structure scores (Infernal - Inference of RNA Alignments) against the clusters and found that scores tended to match within clusters **(Figure S4C)**. These scores measure how much each individual tRNA sequence matches its canonical isotype which matches what we observed with individual isotypes defining certain clusters. This suggests that differential expression and tDR formation are influenced by these genomic features but most likely they are not the only driving factor.[43,44]

Looking further into the relationship between isotype and cluster, we looked at Arg specific tDRs. Disregarding U, C10, and C11 (due to aforementioned indiscernibility) Arg tDRs mainly are split across B1, B2, C4, and C14 clusters all residing in the same spatial area as one another **(Figure S4D)**. Interestingly, Arg-TCT-4 shows up in both B1 (neural) and C10 (neutral) clusters with D70 Arg-TCT-4 appearing in B1. This is most likely due to the extreme change in expression and coverage after neural differentiation with lower read coverage harder to discern. This indicates that other "neutral" tDRs appearing in C clusters may also be neurally associated but can be hard to discern with low read coverage. B1 clusters also contain Arg-TCG-1 and Arg-TCG-5 tDRs, with B2 clusters comprised almost entirely of Arg-CCT tDRs, all tRNAs that increase moderately over differentiation. Interestingly Arg-TCT-1 (which also increases across neural differentiation) appears in C10 (neutral) and U (unannotated) clusters with D70 appearing in C10, this mimics the change over the time course seen in Arg-TCT-4 albeit from U to C10 rather than C10 to B1. While both Arg-TCT-1 and Arg-TCT-4 increase in neural expression, the former has less of an expression increase than the later, and they express entirely different

fragments. This revealed that Arg tDR expression as a function of neural expression is correlated to specific tRNA transcripts rather than specific isotypes and subtle changes to genomic sequence correlate with broad effects in the type and expression levels of tDRs.

Sequence Motifs and Modifications Associate with Neural tDR Clusters

To further understand what causes specific neural tDR shape and expression we looked at tRNA modifications and their enzymes and how they correlated with underlying characteristics of the parent tRNA. Modifications were derived for all human tRNA transcripts from data in Zhang et al. [45] (see methods), and were projected onto the clusters showing that specific modifications such as acp^3U20, m^3C20, and m^1I37 correlate strongly with just neural clusters (B), and acp^3U20a, m^2_2G26, m^3C32, and I34 correlate with neural and neutral (B and C) clusters **(Figure 5A)**. Inosine (at position 37) has been associated with intellectual disabilities, and microcephaly and is added by ADAT3 in humans. m^2_2G (a modification found at position 26 in Arg-TCT-4) has been associated with intellectual disabilities and microcephaly via TRMT1.[46,47] m^1A58 is a ubiquitous modification found in nearly all tRNAs and is consistently expressed across all groups. While not present in our modification data, PUS3 and NSUN2 are known to play a roles in pseudouridylation and m^5C addition at positions 39 and 40, respectively, and are both tied to intellectual development.[48–50] We found that PUS3 and NSUN2 increased while ADAT3 decreased over a five week time course of human embryonic stem cell derived cortical organoids using data from Fiddes et al. [30] **(Figure S4E)**.

Sequence alignments of the tRNAs in the neural clusters (B) revealed conservation of tRNA sequence at positions 26, 30, 32, and 40 **(Figure S5A)**. This includes bases in close proximity to the cleavage at position 40. Finally, we wanted to see if sequence motifs (relative to background frequency) played a role in clusters as they have potential to affect tertiary structure

formation. Motifs such as GGGG (1-4, 68-71), and CCCC (2-5, 60-63) and GCC were observed to be more common in neural (B) clusters **(Figure S5B)**. The motif GCAU (26-29) also indicates a higher frequency of G at position 26 which most likely corresponds to the higher frequency of m^2_2G26 found in the neural (B) clusters. AUG (28-30), GUA (39-41) and GCC (39-41 in neural, 40-42 in stem) indicated that the G-C pair between 30 and 40 occurs more frequently in neural groups but a G is more likely at position 40 in stem. With neutral (C) and unannotated (U) clusters comprising the majority of tRNAs found in the clustering, relative motif frequency was much closer to expected background.

Finally we also looked at the internal arrangement of the tDRs for both Arg-TCT-1 and Arg-TCT-4 by looking at the ratio of 3' read ends (in relation to the tRNA structure) to 5' read ends **(Figure 5B)**. In Arg-TCT-4 read starts are grouped around position 39 and 40 indicating an association with having a C rather than U base changes at that position as found in other Arg-TCT isodecoders. Arg-TCT has five isodecoders, with Arg-TCT-4 having a unique sequence relative to the others containing a 20b position G as well as 2U, 20aC, 26G, 27C, 40C, 43G, 71A, and 72U **(Figure S5C)**. Further when we compared Arg-TCT-4 against the 15 top neural expressed tDRs we found conservation of C at position in 40 in all but Ala-TGC-3 and Ala-TGC-4 **(Figure 5D; Figure S5C)**. Interestingly Ala-TGC-3/4 both express more in D0 relative to D14 before increasing in D35 and D70 suggesting that they are active at different neural time points and might be part of a different neural expression pathway.

Discussion

tRNA gene expression has been previously examined in multiple neurological contexts [19,51,52], including neural cell lines and mature brain tissue, but none designed to capture key developmental transitions from human pluripotent stem cells to the early cerebral cortex and in

tDRs. This study unveils striking changes in tDR abundance and processing patterns during early human corticogenesis, the first requisite step to identifying neural-specific tRNA processing patterns which may have key roles in brain development. We have identified many unique tDRs that change their expression profile and have categorized them. Furthermore, we have categorized these complex tDR patterns by clustering transcripts based on processing patterns, sequence, and modifications to look for commonalities in processing and downstream function.

tRNA-Arg-TCT-4 has been previously identified in mouse as a neural specific tRNA with defects leading to truncal ataxia and severe neurodegeneration when position 50 has a C to T mutation [18], here we show that the Arg-TCT-4 derived tDRs abundance increases across early human neurogenesis as well. This profile of brain organoids has also allowed for the discovery of new tDRs that are induced during neural differentiation, such as those from SeC-TCA-1 and Ala-AGC-8. Further, while most studies focus on single tRNA transcripts, using ARM-seq combined with cerebral brain organoids has allowed for the discovery of many tDRs with dynamic expression in early human embryonic development. These fragments each have unique breakpoints in their read coverage as well, indicative of a unique modification profile on a tRNA by tRNA basis. The top hits for neural specific tDRs seemed to share a coverage profile similar to one another, suggesting they may be generated by a common mechanism.

The profiles of these tDRs tended to fall into similar patterns that could be visually grouped together. For example, some of the top neural favored tDRs (Arg-TCT-4 and Ala-AGC-8) each have a 3' fragmentation pattern that is visually unique, reflecting a combination of associated sequence factors that allows for the fragments to be classified into distinct clusters that are proximal to one another (B1 and C13). This suggests that advanced sequence characteristics beyond unique read coverage drive the grouping of neural expressed

tDRs. Pairing this information with known regulatory tDRs can be further used to understand the relationship between unique fragment profiles and tRNA regulation.

While this approach was used in the context of human cerebral cortex differentiation, identifying and classifying tDRs can be used in a broader context. Understanding the relationship of tDRs in a more expansive way than 3' and 5' derived can lead to a better understanding of tDR regulation overall. For example, the abundance of Arg-TCT-4 and Arg-TCT-1 both increase over neural development, however, Arg-TCT-4's associated 3' tDR spans from the anticodon to the end, whereas the one in Arg-TCT-1 starts in the D-loop instead. Arg-TCT-4 is also expressed at 1/10th the level of Arg-TCT-1, despite having a known regulatory role in the brain. Taken in conjunction, it is suggestive that the uniqueness of the Arg-TCT-4 fragment and not just the abundance of read coverage in mature neurons are what defines Arg-TCT-4 as a neural specific tDR regulator. As such tDRs roles in neurological development may go beyond just upregulation of expression across neural differentiation, and instead be a combination of genomic and RNA features (such as sequence motif and modifications) that play greater roles in regulation rather than translation.

Resource availability

Lead Contact

Requests for further information and resources should be directed to and will be fulfilled by the lead contact, Sofie R. Salama (ssalama@ucsc.edu).

Materials Availability

This study did not generate new unique reagents.

Data and Code Availability

All raw and processed sequencing data generated in this study have been submitted to the NCBI Gene Expression Omnibus (GEO; https://www.ncbi.nlm.nih.gov/geo/) under accession number GSE259250 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE259250). All computational methods are provided in a series of Jupyter notebooks available on GitHub (https://github.com/alba1735/organoidtRNAs/).

Acknowledgments

We received technical support with microscopy from Benjamin Abrams, UCSC Life Sciences Microscopy Center (RRID: SCR_02113). For Cell culture we received support from the CIRM Major Facility Award to UCSC (FA1-00617) and the IBSC Stem Cell Culture Facility (RRID:SCR_021353). Additionally, Dylan Baer assisted with tissue culture and cerebral cortical organoid growth, and Liam Tran helped with IF-staining methods and troubleshooting. We also received support from Jonathan Howard and Kristof Tyigi for advice on experimental methods, technical assistance, and support in RNA library preparation and general laboratory methods. Finally Aidan Manning collaborated in the creation of the modification data file for analysis.

Author contributions

A.B. performed all experiments, conducted the data analysis, generated the figures, tables, code, and wrote the manuscript. T.L. and S.S. provided critical review and editing of the manuscript.

Declaration of interests

The authors declare no competing interests.

Figure Legends

Figure 1: Validation of organoid model and RNA profiling

(A) Graphical abstract of the study depicting media and small molecule drug changes across the organoid developmental time course. Representations of organoids and relative size at crucial time points are shown with 500 μm scale bars. RNA collection, tRNA sequencing, analysis pipeline, and classification are also depicted. **(B)** D70 organoid stains using vimentin (VIM) and DAPI are used, showing cytoarchitecture of the organoid. Emphasis placed with white arrows showing locations or neural rosettes (early endotube-like formation) found throughout the perimeter of the organoid stains. **(C)** Antibody stains PAX6, a radial glial marker, CTIP2, a deep layer neuron marker, and TBR2, an intermediate neural progenitor marker, show neural marker gene expression with PAX6 staining the inside of neural rosettes, TBR2 surrounding the rosettes and CTIP2 encompassing the exterior of the organoid. **(D)** Neural marker stains were merged with a dotted white line for the exterior perimeter of the organoid. **(E)** Volcano plot showing D0 vs. D70 $\log_2$ fold-changes in the abundance of small RNAs against $-\log_{10}$(p-value) with tRNAs shown for scale. Targets of particular interest are labeled with horizontal dotted lines at p-value=0.05 and p-value=0.001 and vertical dotted lines at 1.5 fold difference. Small RNAs highly expressed in either stem or neural cells across multiple conditions (D0 vs. D14, D0 vs. D35, or D0 vs. D70) were highlighted in blue or yellow, respectively. Small RNAs known as stem or neural markers were highlighted in purple or magenta as well. **(F)** PCA plot of all ARM-seq samples for all measured small RNAs across organoid time points.

Figure 2: Broad changes in tRNA Isotypes do not capture tDR-Specific Changes

(A) Human small RNA read count distributions showing relative ratio of small RNAs across organoid time course, sequencing type, and AlkB treatment. **(B)** Human tRNA isotype reads-counts (total and fragment) distributions showing relative ratio of tRNA isotypes across organoid time course, sequencing type, and AlkB treatment. **(C)** Coverage profiles for the Gly, Thr, and Val isotype groups showing max expression across AlkB+ samples per position within each isotype across organoid time points. D-loop, anticodon-loop, and T-loop shown in gray, with lines added at specific coverage breakpoints in each plot.

Figure 3: Expression patterns of neural tDRs are unique and change over time

(A) Volcano plot showing D0 vs. D70 $\log_2$ fold-changes in the abundance of tDRs against $-\log_{10}$(p-value) with 3', 5' and internal fragment counts and full-length and pre-tRNA counts. Targets of particular interest are labeled with horizontal dotted lines at p-value=0.05 and p-value=0.001 and vertical dotted lines at 1.5 fold difference. tDRs highly expressed in either stem or neural cells across multiple conditions (D0 vs. D14, D0 vs. D35, or D0 vs. D70) were

highlighted in shades of blue or yellow, depending on how many conditions they appeared in. All samples were batch-corrected and normalized using DESeq2 [53] to account for differences in sample read counts. **(B)** Heatmap showing the top 15 neural and top 15 stem-favored tDRs across all conditions. Names in orange have less than 80 uniquely mapping read counts. **(C-E)** Coverage profiles for the top 3 highly expressed tDRs over organoid neurogenesis **(C)**, highly expressed stem and neural Gly tDRs **(D)**, and unique tDRs with coverage changes **(E)**. D-loop, anticodon-loop, and T-loop shown in gray, with lines added at specific coverage breakpoints in each plot.

Figure 4: Multidimensional Analysis Reveals tDR Clusters Associated with Neural vs. Stem States

(A) UMAP projections of tDR sequencing profiles, using unique coverage of tDR 3' and 5' read ends, read alignment deletions (see methods), and mismatched bases, by position in the tRNA, using HDBSCAN clustering and manual annotation of stem (A*n*), neural (B*n*), and neutral (C*n*) classes. tDR fragment profiles that could not be confidently annotated into clusters by HDBSCAN were masked (grey) and not included in further analysis. **(B)** Dot plot representation of the HDBSCAN clusters with dot size correlating to the proportion of isotype found in each cluster and the color based on the mean neural stem score (as defined in methods) with vertical lines at the top and bottom 25th percentiles defining the groups.

Figure 5: Sequence Motifs and Modifications Associate with Neural tDR Clusters

(A) Dot plot representation of the HDBSCAN clusters with dot size correlating to the proportion of modification at positions found in each cluster and the color based on the mean neural stem score (as defined in methods) with vertical lines at the top and bottom 25th percentiles defining the groups. **(B)** The percentage of 3' and 5' read ends of Arg-TCT-1 and Arg-TCT-4 tDRs. Positions with greater than 10% of relative normalized fragment read-starts/ends indicated with vertical dotted lines. **(C)** Secondary structure of Arg-TCT-4 showing divergence with Arg-TCT-1, key differences in bold and highlighted with arrows. **(D)** Sequence map of Arg-TCT tRNAs and top 15 most neural-expressed tDRs.

STAR Methods

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
See Table S4	Various	N/A

Chemicals, peptides, and kits		
AggreWell media	Stemcell Technologies	#05893
Agilent DNA High Sensitivity kit	Agilent	#5067-4626
Beta fibroblast growth factor (bFGF)	Miltenyi Biotec	#130-095-738
Brain-derived neurotrophic factor (BDNF)	Miltenyi Biotec	#130-093-811
Bovine Serum Albumin (BSA)	ThermoFisher Scientific	#B14
Chemiluminescent Nucleic Acid Detection Module	ThermoFisher Scientific	#89880
Direct-Zol RNA MiniPrep Kit	Zymo Research	#R2052
DMEM/F-12, GlutaMAX™ supplement	ThermoFisher Scientific	#10565018
DynaMarker Prestain Marker for Small RNA Plus	DiagnoCine	#DM253S
Epidermal growth factor (EGF)	R&D	#236-EG-01M
Essential-8 Flex media	ThermoFisher Scientific	#A2858501
Cytiva HyClone™ Characterized Fetal Bovine Serum (U.S.)	Fisher Scientific	#SH3007103
Gel Elution Buffer	New England Biolabs	#T3010
IWR-1	Millipore-Sigma	#681669
Linear Acrylamide	New England Biolabs	(supplied in NEBNext Kit)
Matrigel	Corning	#354277
mirVana™ miRNA Isolation Kit, without phenol	ThermoFisher Scientific	#AM1561
mTeSR™ Plus	Stemcell Technologies	#100-0276
N-(3-Dimethylaminopropyl)-N'-ethylcarbodiimide (EDC)	Sigma-Aldrich	#39391-10ML
N2 supplement	ThermoFisher Scientific	#17502048
NEBNext Multiplex Small RNA Library Prep Set	New England Biolabs	#E7300L
Neurotrophin-3 (NT-3)	Peprotech	#450-03-100ug
Paraformaldehyde (PFA)	ThermoFisher Scientific	#J19943.K2
PBS	Corning	#21031CM
ProLong™ Gold Antifade Mountant with DAPI	ThermoFisher Scientific	#P36935
RNA Clean and Concentrate-25	Zymo Research	#R1018
Sasai II media	N/A	(See composition in text)

Sasai III media	N/A	(See composition in text)
SB431542 (SB)	Millipore-Sigma	#S4317-5MG
Sucrose	Millipore-Sigma	#S0389-1KG
SuperSignal™ West Femto Maximum Sensitivity Substrate	ThermoFisher Scientific	#34094
SYBR™ Gold Nucleic Acid Gel Stain	ThermoFisher Scientific	#S11494
Tissue-Tek O.C.T. Compound	Sakura	#4583
TRI Reagent	Molecular Research Center, Inc.	#TR118
ULTRAhyb™ Ultrasensitive Hybridization Buffer	ThermoFisher Scientific	#AM8670
Vitronectin	ThermoFisher Scientific	#A14700
Y-27632 rock inhibitor	Stemcell Technologies	#72308
Materials, Devices		
AggreWell-800 plates	Stemcell Technologies	#34821
Low attachment 6-well dishes	Corning	#3471
Positively charged nylon membrane	Millipore-Sigma	#11209299001
Deposited data		
Primary sequencing data deposited	Gene Expression Omnibus (GEO)	GEO: GSE259250, https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE259250
Cell lines		
Human GM12878-c305 induced pluripotent stem cells (iPSCs)	Coriell	
Oligonucleotides		
See Table S4	IDT	N/A
Software and algorithms		
Bio-Rad Image Lab	Bio-Rad	https://www.bio-rad.com/en-us/product/image-lab-software?ID=KRE6P5E8Z

		https://doi.org/10.5281/zenodo.14675803
Custom Code	10.5281/zenodo.14675803	
ImageJ (Fiji)	Schindelin et al. [54]	https://imagej.net/software/fiji/
tRAX	Holmes et al. [24]	https://trna.ucsc.edu/tRAX/
tRNAgraph	10.5281/zenodo.14675737	https://doi.org/10.5281/zenodo.14675737
Zen Software suite	Zeiss	https://www.zeiss.com/microscopy/us/products/software/zeiss-zen.html

Organoid methods

Human GM12878-c305 induced pluripotent stem cells (iPSCs) were generated from the GM12878 lymphoblastoid cell line (Coriell) using a proprietary episomal vector-based reprogramming method by Cellular Dynamics International (fujifilmcdi.com) and were confirmed to have a normal karyotype using the KaryoStat Karyotyping service (ThermoFisher). Undifferentiated cells were grown in feeder-free conditions on matrigel (Corning) with mTeSR Plus (Stemcell Technologies) or on vitronectin with Essential-8 Flex media (ThermoFisher). Cerebral cortex organoids were generated using a protocol adapted from Kadoshima et al. [28]. 10,000 cells per embryoid body were aggregated using AggreWell-800 plates in AggreWell media (Stemcell Technologies) supplemented with 10 uM Y-27632 rock inhibitor (Stemcell Technologies), and transferred to low attachment 6-well dishes (Corning) on day 2. These methods supplemented the respective media with 10 uM SB431542 (SB, Millipore), and 1 uM IWR-1 (Millipore) for the first 14 days of differentiation. The media was then changed to Sasai II media (DMEM/F12 + glutamax supplemented with N2) on day 14. At this point, organoids were supplemented with 10ng/mL beta fibroblast growth factor (bFGF) and 10ng/mL epidermal

growth factor (EGF) to improve survival in Sasai II media. On day 18 the organoids were transferred to an in-incubator orbital shaker (100 rpm) where they remained for the duration of the experiment. After day 35, all cultures were grown in Sasai III media (Sasai II media supplemented with 50 mL of FBS (Hyclone)) supplemented with 10 ng/mL brain-derived neurotrophic factor (BDNF) and 10 ng/mL of neurotrophin-3 (NT-3).[55]

Organoid cultures exhibited heterogeneity in size and morphology. To mitigate this variability and ensure robust RNA-seq analysis, we initiated cultures with eight biological replicates in order to yield sufficient RNA for multiple independent ARM-seq libraries at each time point. Samples exhibiting low sequencing depth (<75k reads) were excluded from further analysis. This resulted in a larger number of biological replicates were available at earlier time points (n=8) compared to later time points (n=2).

Immunofluorescence Microscopy

Organoids were collected and fixed in 4% Paraformaldehyde (PFA) (ThermoFisher), washed 3x with PBS, then incubated in 500 μL of 30% Sucrose for 2-3 days at 4∘C until they began to float. They were then embedded in square cryomolds with Tissue-Tek O.C.T. Compound (Sakura) and placed at -80∘C for storage. They were then sectioned to 18 μm using a cryostat (Leica Biosystems) directly onto glass slides. After allowing samples to come to room temperature, 3 washes of 35 minutes in 1X PBS were performed. The sections were then incubated in a blocking solution of 10% BSA for 2 hours. The sections were then incubated in primary antibodies and blocking solution at 1:1000 dilution overnight at 4∘C **(Table S4)**. They were then washed 3 times for 30 minutes and incubated in secondary antibodies for 2 hours at room temperature. They were then washed 3 times for 30 minutes in PBS and sealed with ProLong™ Gold Antifade Mountant with DAPI (ThermoFisher Scientific # P36935). Microscopy

was performed using a Zeiss Axio Imager and Zen Software suite, with processing of the images performed using Fiji.[54]

RNA Isolation

Isolation of total RNA from cerebral cortical organoids and stem cells was performed using Direct-Zol RNA MiniPrep Kit (Zymo Research) with TRI Reagent (Molecular Research Center, Inc.). Since a single organoid yields far less total RNA than required for one sequencing library, approximately 3-8 organoids would be pooled in Trizol depending on organoid size for each sample replicate. The manufacturer's recommended volume of TRI Reagent was added to each sample (~1 mL). For RNA purification of stem cells, Trizol was directly added to cell culture plates on ice, scraped, and homogenized via pipetting. Organoids in Trizol were broken down via pipetting inside a 1mL Eppendorf tube on ice via a syringe. All total RNA was processed using a MirVana miRNA Isolation Kit (ThermoFisher Scientific), according to the manufacturer's instructions, to select for RNA <200 nt. This was followed by an RNA Clean and Concentrate-25 (Zymo Research).

Samples were divided into plus-AlkB experimental treatment and minus-AlkB control as previously described in ARM-seq methods.[20] AlkB sample treatment was used to effectively increase RT processivity of hyper-modified tRNA/tDRs while also removing sequencing bias favoring hypo-modified tRNA/tDRs **(Figure S1A)**. For example, the proportion of ARM-seq tRNA reads more than doubles (8.9% to 21.8%) when comparing untreated versus AlkB-treated samples. This was followed by a phenol-chloroform cleanup treatment. RNA samples were then used for library preparations.

ARM-seq library preparation

ARM-seq libraries were constructed as described previously [20] utilizing the NEBNext Multiplex Small RNA Library Prep Set (New England Biolabs). Treated RNA (Minus- or Plus-AlkB treatment; 100ng total small RNA) was used as input into the library preparation, and ¼ reaction volume of all reagents were used with the NEBNext kit. PCR-amplified libraries were purified using a phenol-chloroform extraction cleanup and then size-selected (140-250 nts) on a 6% non-denaturing TBE-acrylamide gel to remove unwanted primer dimer products. Libraries were eluted from sliced gel pieces using Gel Elution Buffer (New England Biolabs) and precipitated using 0.3 M NaOAc, 80% Ethanol, and 1 µL of Linear Acrylamide (supplied in NEBNext Kit) at final concentration. Samples were left in -80∘C freezer overnight to precipitate. Precipitated libraries were then pelleted, washed twice in 80% Ethanol, and resuspended in pure H2O. Libraries were then quantified using Qubit and Agilent DNA High Sensitivity kit.

RNA sequencing and differential expression analysis

Libraries prepared for ARM-seq were sequenced using either Illumina MiSeq or Illumina NextSeq 550. 75-nt pair-ended reads were produced as FASTQ files that were analyzed using tRNA Analysis of eXpression (tRAX).[24] Sequence adapters were trimmed, and pair-ended reads were merged using the tool trimadapters.py in tRAX. The reference database for tRAX was built with high-confidence tRNA predictions retrieved from the Genome tRNA Database[56] and the sequences of human genome assembly GRCh38. Other gene annotations were obtained from Ensembl release 102. Biological replicates of organoids at each time point were grouped as sample replicates for tRAX inputs and different time points were marked as pairs for differential expression comparison for each sequencing type. Default options of tRAX were used, and

samples with low sequence depth and outliers after DESeq2 correction were removed from analysis, ensuring duplicate at a minimum across all time points. To check for biases in organoid time points (due to unequal replicate distribution), total normalized reads **(Figure S1A,B)** were compared, showing no notable differences in abundance between time points. Additionally, we found the relative levels of tRNAs were consistent between sample groups, with no significant difference in total reads found between samples within AlkB treatment groups **(Figure S6A,B)**. AlkB treatment significantly increases tRNA read counts relative to overall RNA reads **(Figure S1B)**. ARM-seq also preferentially picks up more tRNA fragments and their read counts relative to full-length tRNAs **(Figure S1C)**. In order to standardize tRNA coverage profiles, each series of reads was aligned to conserved tRNA positions [57], dropping gap and extension positions. All visualizations were created using tRNAgraph (https://github.com/alba1735/tRNAgraph) and custom Python scripts (https://github.com/alba1735/organoidtRNAs).

Northern Blot Analysis

Northern blot analysis was conducted to assess the expression levels of specific tRNAs over a 70-day time course and was modified from Damm et al. [58]. Previously extracted day 0 and day 70 Total RNA was used. RNA samples (5-20 µg per lane) were resolved on a 15% TBE-urea denaturing polyacrylamide gel and transferred to a positively charged nylon membrane (Sigma) using a Trans-Blot® Turbo™ Transfer System (Bio-Rad). Membranes were then UV-crosslinked using an EDC (N-(3-Dimethylaminopropyl)-N'-ethylcarbodiimide)-mediated (Sigma) crosslinking protocol. Subsequently, membranes were pre-hybridized in ULTRAhyb Ultrasensitive Hybridization Buffer (Thermo Fisher) before overnight hybridization with custom 3' biotinylated LNA probes (IDT) designed against specific tRNA species: Arg-TCT-4, Ala-AGC-8, and SeC-TCA-1 at a final concentration of 50 pmol/mL **(Table S4)**. Following hybridization,

membranes were subjected to a series of stringent washes using low-stringency (2x SSC, 0.1% SDS) and high-stringency (0.1x SSC, 0.1% SDS) buffers. Detection was performed using the Chemiluminescent Nucleic Acid Detection Module (Thermo Fisher), and signal was visualized using a Chemi-Doc imager (Bio-Rad). Since specific tRNA/tDR concentration is low, a more sensitive substrate, SuperSignal™ West Femto Maximum Sensitivity Substrate (Thermo Fisher), was employed for signal detection. To ensure equal loading across lanes, the prestained DynaMarker Prestain Marker for Small RNA Plus (DiagnoCine) was used. Additionally, membranes were stained pre-transfer with SYBR™ Gold Nucleic Acid Gel Stain (Thermo Fisher) and imaged under UV light to verify the presence and integrity of the transferred RNA. Membranes were stripped using a boiling 0.1% SDS solution and reprobed as necessary following the overnight hybridization protocol described above. The Northern blots were normalized with a 5S probe for quantification.

tRNA Classification Analysis

Sequencing output data from tRAX was used for tDR classification. DESeq2 normalized read counts were combined into an AnnData object and were arranged by coverage-associated data across human GRCh38 tRNAs.[59] These were comprised of unique coverage (normalized read count with reads that are uniquely mapped to feature at the specific position), 3' and 5' read ends (with consideration to the tRNA structure), deletions (number of reads that have a gap at the specific position), and mismatched bases per position (normalized read count that does not match the reference base at specific position). Unique reads were chosen over total reads as this provided more specific coverage changes in the clusters and better tRNA uniformity therein. Thus, each cluster portrays the coverage makeup of component tDRs and gives a fragment profile in addition to other underlying sequence features. The data was further

preprocessed by removing read counts less than 20, regressing out the number of reads, and scaling and centering the data. Dimensionality reduction was performed using Uniform Manifold Approximation and Projection (UMAP) and clustering using HDBSCAN.[41,42] Classifying tRNAs into neural, neutral, and stem categories was based on the $\log_2$ fold-changes between day 0 and day 70 time points split into the top and bottom 20th percentiles of all tRNAs, after removing reads with low coverage and p-value >0.05. $\log_2$ fold-change was normalized between -1 and 1 with each tRNA transcript and assigned a "neural stem score" based on its relative expression between the differentiated and undifferentiated time points. To determine if a cluster was neural or stem specific, the mean of the "neural stem scores" was taken for each cluster, and the top and bottom 25 percentiles correlated to neural (B) and stem (A) clusters, respectively. Clusters from neither group were labeled as neutral (C) and unannotated (U) results from HDBSCAN, their own group. Sequence logos were generated using Logomaker [60] by using the mean normalized read counts per position and converting them into information scores.[61]

tRNA Modification Analysis

To derive tRNA modifications for analysis, we utilized data from Zhang et al. [45], specifically, we obtained the HEK293T cell line RNA sequencing data generated using OTTRseq, which provides full-length sequencing information necessary for modification analysis. We processed this data using the tMAP pipeline (https://github.com/Aimann/tMAP), as described in their paper. During this processing, we flagged any tRNA positions exhibiting a mismatch rate greater than 5% compared to the reference tRNA sequence. These mismatches are indicative of potential RNA modifications and, we cross-referenced the flagged tRNA positions with the Modomics database, a repository of known RNA modifications [62] as well commonly cited modifications in human.[63] Any flagged positions that matched a known

modification were then labeled and projected onto the clusters (described above) identified in

our primary experimental data.

References

1. Haussecker, D., Huang, Y., Lau, A., Parameswaran, P., Fire, A.Z., and Kay, M.A. (2010). Human tRNA-derived small RNAs in the global regulation of RNA silencing. RNA N. Y. N *16*, 673–695. https://doi.org/10.1261/rna.2000810.
2. Avcilar-Kucukgoze, I., and Kashina, A. (2020). Hijacking tRNAs From Translation: Regulatory Functions of tRNAs in Mammalian Cell Physiology. Front. Mol. Biosci. *7*, 610617. https://doi.org/10.3389/fmolb.2020.610617.
3. Nie, A., Sun, B., Fu, Z., and Yu, D. (2019). Roles of aminoacyl-tRNA synthetases in immune regulation and immune diseases. Cell Death Dis. *10*, 1–14. https://doi.org/10.1038/s41419-019-2145-5.
4. Su, Z., Frost, E.L., Lammert, C.R., Przanowska, R.K., Lukens, J.R., and Dutta, A. (2019). tRNA-derived fragments and microRNAs in the maternal-fetal interface of a mouse maternal-immune-activation autism model. bioRxiv, 2019.12.20.884650. https://doi.org/10.1101/2019.12.20.884650.
5. Goodarzi, H., Nguyen, H.C.B., Zhang, S., Dill, B.D., Molina, H., and Tavazoie, S.F. (2016). Modulated Expression of Specific tRNAs Drives Gene Expression and Cancer Progression. Cell *165*, 1416–1427. https://doi.org/10.1016/j.cell.2016.05.046.
6. Goodarzi, H., Liu, X., Nguyen, H.C.B., Zhang, S., Fish, L., and Tavazoie, S.F. (2015). Endogenous tRNA-Derived Fragments Suppress Breast Cancer Progression via YBX1 Displacement. Cell *161*, 790–802. https://doi.org/10.1016/j.cell.2015.02.053.
7. Dou, S., Wang, Y., and Lu, J. (2019). Metazoan tsRNAs: Biogenesis, Evolution and Regulatory Functions. Non-Coding RNA *5*. https://doi.org/10.3390/ncrna5010018.
8. Soares, A.R., and Santos, M. (2017). Discovery and function of transfer RNA-derived fragments and their role in disease. WIREs RNA *8*, e1423. https://doi.org/10.1002/wrna.1423.
9. Morisaki, I., Shiraishi, H., Fujinami, H., Shimizu, N., Hikida, T., Arai, Y., Kobayashi, T., Hanada, R., Penninger, J.M., Fujiki, M., et al. (2021). Modeling a human CLP1 mutation in mouse identifies an accumulation of tyrosine pre-tRNA fragments causing pontocerebellar hypoplasia type 10. Biochem. Biophys. Res. Commun. *570*, 60–66. https://doi.org/10.1016/j.bbrc.2021.07.036.
10. Yang, X.-L. (2014). tRNA-derived G-quadruplex protects motor neurons. Proc. Natl. Acad. Sci. *111*, 18108–18109. https://doi.org/10.1073/pnas.1420838111.
11. Ivanov, P., O'Day, E., Emara, M.M., Wagner, G., Lieberman, J., and Anderson, P. (2014). G-quadruplex structures contribute to the neuroprotective effects of angiogenin-induced tRNA fragments. Proc. Natl. Acad. Sci. U. S. A. *111*, 18201–18206. https://doi.org/10.1073/pnas.1407361111.
12. Boskovic, A., Bing, X.Y., Kaymak, E., and Rando, O.J. (2020). Control of noncoding RNA production and histone levels by a 5′ tRNA fragment. Genes Dev. *34*, 118–131. https://doi.org/10.1101/gad.332783.119.
13. Li, J., Wang, Y.-N., Xu, B.-S., Liu, Y.-P., Zhou, M., Long, T., Li, H., Dong, H., Nie, Y., Chen, P.R., et al. (2020). Intellectual disability-associated gene ftsj1 is responsible for

2′-O-methylation of specific tRNAs. EMBO Rep. *21*, e50095.
https://doi.org/10.15252/embr.202050095.

14. Pitts, M.W., Byrns, C.N., Ogawa, A.N., Kremer, P., and Berry, M.J. (2014). Selenoproteins in Nervous System Development and Function. Biol. Trace Elem. Res. *161*, 231–245. https://doi.org/10.1007/s12011-014-0060-2.

15. Ullah, R., Jo, M.H., Riaz, M., Alam, S.I., Saeed, K., Ali, W., Rehman, I.U., Ikram, M., and Kim, M.O. (2020). Glycine, the smallest amino acid, confers neuroprotection against d-galactose-induced neurodegeneration and memory impairment by regulating c-Jun N-terminal kinase in the mouse brain. J. Neuroinflammation *17*, 303. https://doi.org/10.1186/s12974-020-01989-w.

16. Mader, M.M.-D., and Czorlich, P. (2021). The role of L-arginine metabolism in neurocritical care patients. Neural Regen. Res. *17*, 1446–1453. https://doi.org/10.4103/1673-5374.327331.

17. Qin, C., Xu, P.-P., Zhang, X., Zhang, C., Liu, C.-B., Yang, D.-G., Gao, F., Yang, M.-L., Du, L.-J., and Li, J.-J. (2020). Pathological significance of tRNA-derived small RNAs in neurological disorders. Neural Regen. Res. *15*, 212. https://doi.org/10.4103/1673-5374.265560.

18. Ishimura, R., Nagy, G., Dotu, I., Zhou, H., Yang, X.-L., Schimmel, P., Senju, S., Nishimura, Y., Chuang, J.H., and Ackerman, S.L. (2014). Ribosome stalling induced by mutation of a CNS-specific tRNA causes neurodegeneration. Science *345*, 455–459. https://doi.org/10.1126/science.1249749.

19. Kapur, M., Ganguly, A., Nagy, G., Adamson, S.I., Chuang, J.H., Frankel, W.N., and Ackerman, S.L. (2020). Expression of the Neuronal tRNA n-Tr20 Regulates Synaptic Transmission and Seizure Susceptibility. Neuron *108*, 193-208.e9. https://doi.org/10.1016/j.neuron.2020.07.023.

20. Cozen, A.E., Quartley, E., Holmes, A.D., Hrabeta-Robinson, E., Phizicky, E.M., and Lowe, T.M. (2015). ARM-seq: AlkB-facilitated RNA methylation sequencing reveals a complex landscape of modified tRNA fragments. Nat. Methods *12*, 879–884. https://doi.org/10.1038/nmeth.3508.

21. Shigematsu, M., Honda, S., Loher, P., Telonis, A.G., Rigoutsos, I., and Kirino, Y. (2017). YAMAT-seq: an efficient method for high-throughput sequencing of mature transfer RNAs. Nucleic Acids Res. *45*, e70. https://doi.org/10.1093/nar/gkx005.

22. Cheng, Y.-H., Chen, Y.-C., Lin, E., Brien, R., Jung, S., Chen, Y.-T., Lee, W., Hao, Z., Sahoo, S., Min Kang, H., et al. (2019). Hydro-Seq enables contamination-free high-throughput single-cell RNA-sequencing for circulating tumor cells. Nat. Commun. *10*, 2163. https://doi.org/10.1038/s41467-019-10122-2.

23. Pinkard, O., McFarland, S., Sweet, T., and Coller, J. (2020). Quantitative tRNA-sequencing uncovers metazoan tissue-specific tRNA regulation. Nat. Commun. *11*, 4104. https://doi.org/10.1038/s41467-020-17879-x.

24. Holmes, A.D., Howard, J.M., Chan, P.P., and Lowe, T.M. (2022). tRNA Analysis of eXpression (tRAX): A tool for integrating analysis of tRNAs, tRNA-derived small RNAs, and tRNA modifications. Preprint at bioRxiv, https://doi.org/10.1101/2022.07.02.498565 https://doi.org/10.1101/2022.07.02.498565.

25. Eiraku, M., Watanabe, K., Matsuo-Takasaki, M., Kawada, M., Yonemura, S., Matsumura, M., Wataya, T., Nishiyama, A., Muguruma, K., and Sasai, Y. (2008). Self-organized formation of polarized cortical tissues from ESCs and its active manipulation by extrinsic signals. Cell Stem Cell *3*, 519–532. https://doi.org/10.1016/j.stem.2008.09.002.

26. Camp, J.G., Badsha, F., Florio, M., Kanton, S., Gerber, T., Wilsch-Bräuninger, M., Lewitus,

E., Sykes, A., Hevers, W., Lancaster, M., et al. (2015). Human cerebral organoids recapitulate gene expression programs of fetal neocortex development. Proc. Natl. Acad. Sci. *112*, 15672–15677. https://doi.org/10.1073/pnas.1520760112.

27. Kelava, I., and Lancaster, M.A. (2016). Dishing out mini-brains: Current progress and future prospects in brain organoid research. Dev. Biol. *420*, 199–209. https://doi.org/10.1016/j.ydbio.2016.06.037.

28. Kadoshima, T., Sakaguchi, H., Nakano, T., Soen, M., Ando, S., Eiraku, M., and Sasai, Y. (2013). Self-organization of axial polarity, inside-out layer pattern, and species-specific progenitor dynamics in human ES cell–derived neocortex. Proc. Natl. Acad. Sci. *110*, 20284–20289. https://doi.org/10.1073/pnas.1315710110.

29. Pollen, A.A., Bhaduri, A., Andrews, M.G., Nowakowski, T.J., Meyerson, O.S., Mostajo-Radji, M.A., Lullo, E.D., Alvarado, B., Bedolli, M., Dougherty, M.L., et al. (2019). Establishing Cerebral Organoids as Models of Human-Specific Brain Evolution. Cell *176*, 743-756.e17. https://doi.org/10.1016/j.cell.2019.01.017.

30. Fiddes, I.T., Lodewijk, G.A., Mooring, M., Bosworth, C.M., Ewing, A.D., Mantalas, G.L., Novak, A.M., Bout, A. van den, Bishara, A., Rosenkrantz, J.L., et al. (2018). Human-Specific NOTCH2NL Genes Affect Notch Signaling and Cortical Neurogenesis. Cell *173*, 1356-1369.e22. https://doi.org/10.1016/j.cell.2018.03.051.

31. Field, A.R., Jacobs, F.M.J., Fiddes, I.T., Phillips, A.P.R., Reyes-Ortiz, A.M., LaMontagne, E., Whitehead, L., Meng, V., Rosenkrantz, J.L., Olsen, M., et al. (2019). Structurally Conserved Primate LncRNAs Are Transiently Expressed during Human Cortical Differentiation and Influence Cell-Type-Specific Genes. Stem Cell Rep. *12*, 245–257. https://doi.org/10.1016/j.stemcr.2018.12.006.

32. Coolen, M., Katz, S., and Bally-Cuif, L. (2013). miR-9: a versatile regulator of neurogenesis. Front. Cell. Neurosci. *0*. https://doi.org/10.3389/fncel.2013.00220.

33. Cimadamore, F., Amador-Arjona, A., Chen, C., Huang, C.-T., and Terskikh, A.V. (2013). SOX2–LIN28/let-7 pathway regulates proliferation and neurogenesis in neural precursors. Proc. Natl. Acad. Sci. *110*, E3017–E3026. https://doi.org/10.1073/pnas.1220176110.

34. Zhao, C., Sun, G., Li, S., Lang, M.-F., Yang, S., Li, W., and Shi, Y. (2010). MicroRNA let-7b regulates neural stem cell proliferation and differentiation by targeting nuclear receptor TLX signaling. Proc. Natl. Acad. Sci. *107*, 1876–1881. https://doi.org/10.1073/pnas.0908750107.

35. Fairchild, C.L.A., Cheema, S.K., Wong, J., Hino, K., Simó, S., and La Torre, A. (2019). Let-7 regulates cell cycle dynamics in the developing cerebral cortex and retina. Sci. Rep. *9*, 15336. https://doi.org/10.1038/s41598-019-51703-x.

36. Hashimoto, Y., Niikura, T., Tajima, H., Yasukawa, T., Sudo, H., Ito, Y., Kita, Y., Kawasumi, M., Kouyama, K., Doyu, M., et al. (2001). A rescue factor abolishing neuronal cell death by a wide spectrum of familial Alzheimer's disease genes and Abeta. Proc. Natl. Acad. Sci. U. S. A. *98*, 6336–6341. https://doi.org/10.1073/pnas.101133498.

37. Hashimoto, Y., Niikura, T., Ito, Y., Sudo, H., Hata, M., Arakawa, E., Abe, Y., Kita, Y., and Nishimoto, I. (2001). Detailed characterization of neuroprotection by a rescue factor humanin against various Alzheimer's disease-relevant insults. J. Neurosci. Off. J. Soc. Neurosci. *21*, 9235–9245.

38. Ying, G., Iribarren, P., Zhou, Y., Gong, W., Zhang, N., Yu, Z.-X., Le, Y., Cui, Y., and Wang, J.M. (2004). Humanin, a newly identified neuroprotective factor, uses the G protein-coupled formylpeptide receptor-like-1 as a functional receptor. J. Immunol. Baltim. Md 1950 *172*, 7078–7085. https://doi.org/10.4049/jimmunol.172.11.7078.

39. Bratkovič, T., Modic, M., Camargo Ortega, G., Drukker, M., and Rogelj, B. (2018). Neuronal differentiation induces SNORD115 expression and is accompanied by post-transcriptional

changes of serotonin receptor 2c mRNA. Sci. Rep. *8*, 5101.
https://doi.org/10.1038/s41598-018-23293-7.

40. Lyons, S.M., Fay, M.M., and Ivanov, P. (2018). The role of RNA modifications in the regulation of tRNA cleavage. FEBS Lett. *592*, 2828–2844.
https://doi.org/10.1002/1873-3468.13205.

41. McInnes, L., Healy, J., and Melville, J. (2018). UMAP: Uniform Manifold Approximation and Projection for Dimension Reduction. ArXiv180203426 Cs Stat.

42. McInnes, L., Healy, J., and Astels, S. (2017). hdbscan: Hierarchical density based clustering. J. Open Source Softw. *2*, 205. https://doi.org/10.21105/joss.00205.

43. Chan, P.P., Lin, B.Y., Mak, A.J., and Lowe, T.M. (2019). tRNAscan-SE 2.0: Improved Detection and Functional Classification of Transfer RNA Genes. bioRxiv, 614032.
https://doi.org/10.1101/614032.

44. Nawrocki, E.P., and Eddy, S.R. (2013). Infernal 1.1: 100-fold faster RNA homology searches. Bioinforma. Oxf. Engl. *29*, 2933–2935.
https://doi.org/10.1093/bioinformatics/btt509.

45. Zhang, K., Manning, A.C., Lentini, J.M., Howard, J., Dalwigk, F., Maroofian, R., Efthymiou, S., Chan, P., Eliseev, S.I., Yang, Z., et al. (2025). Human TRMT1 and TRMT1L paralogs ensure the proper modification state, stability, and function of tRNAs. Cell Rep. *44*.
https://doi.org/10.1016/j.celrep.2024.115092.

46. de Crécy-Lagard, V., Boccaletto, P., Mangleburg, C.G., Sharma, P., Lowe, T.M., Leidel, S.A., and Bujnicki, J.M. (2019). Matching tRNA modifications in humans to their known and predicted enzymes. Nucleic Acids Res. *47*, 2143–2159. https://doi.org/10.1093/nar/gkz011.

47. Chujo, T., and Tomizawa, K. (2021). Human transfer RNA modopathies: diseases caused by aberrations in transfer RNA modifications. FEBS J. https://doi.org/10.1111/febs.15736.

48. Chen, J., and Patton, J.R. (2000). Pseudouridine synthase 3 from mouse modifies the anticodon loop of tRNA. Biochemistry *39*, 12723–12730. https://doi.org/10.1021/bi001109m.

49. Shaheen, R., Han, L., Faqeih, E., Ewida, N., Alobeid, E., Phizicky, E.M., and Alkuraya, F.S. (2016). A homozygous truncating mutation in PUS3 expands the role of tRNA modification in normal cognition. Hum. Genet. *135*, 707–713.
https://doi.org/10.1007/s00439-016-1665-7.

50. Tuorto, F., Liebers, R., Musch, T., Schaefer, M., Hofmann, S., Kellner, S., Frye, M., Helm, M., Stoecklin, G., and Lyko, F. (2012). RNA cytosine methylation by Dnmt2 and NSun2 promotes tRNA stability and protein synthesis. Nat. Struct. Mol. Biol. *19*, 900–905.
https://doi.org/10.1038/nsmb.2357.

51. Ishimura, R., Nagy, G., Dotu, I., Zhou, H., Yang, X.-L., Schimmel, P., Senju, S., Nishimura, Y., Chuang, J.H., and Ackerman, S.L. (2014). Ribosome stalling induced by mutation of a CNS-specific tRNA causes neurodegeneration. Science *345*, 455–459.
https://doi.org/10.1126/science.1249749.

52. Kapur, M., Molumby, M.J., Guzman, C., Heinz, S., and Ackerman, S.L. (2024). Cell-type-specific expression of tRNAs in the brain regulates cellular homeostasis. Neuron *112*, 1397-1415.e6. https://doi.org/10.1016/j.neuron.2024.01.028.

53. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550.
https://doi.org/10.1186/s13059-014-0550-8.

54. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-image analysis. Nat. Methods *9*, 676–682.
https://doi.org/10.1038/nmeth.2019.

55. Kindberg, A.A., Bendriem, R.M., Spivak, C.E., Chen, J., Handreck, A., Lupica, C.R., Liu, J., Freed, W.J., and Lee, C.-T. (2014). An in vitro model of human neocortical development using pluripotent stem cells: cocaine-induced cytoarchitectural alterations. Dis. Model. Mech. *7*, 1397–1405. https://doi.org/10.1242/dmm.017251.

56. Chan, P.P., and Lowe, T.M. (2016). GtRNAdb 2.0: an expanded database of transfer RNA genes identified in complete and draft genomes. Nucleic Acids Res. *44*, D184–D189. https://doi.org/10.1093/nar/gkv1309.

57. Sprinzl, M., Steegborn, C., Hübel, F., and Steinberg, S. (1996). Compilation of tRNA Sequences and Sequences of tRNA Genes. Nucleic Acids Res. *24*, 68–72. https://doi.org/10.1093/nar/24.1.68.

58. Damm, K., Bach, S., Müller, K.M.H., Klug, G., Burenina, O.Y., Kubareva, E.A., Grünweller, A., and Hartmann, R.K. (2015). Improved Northern blot detection of small RNAs using EDC crosslinking and DNA/LNA probes. Methods Mol. Biol. Clifton NJ *1296*, 41–51. https://doi.org/10.1007/978-1-4939-2547-6_5.

59. Virshup, I., Rybakov, S., Theis, F.J., Angerer, P., and Wolf, F.A. (2021). anndata: Annotated data. Preprint at bioRxiv, https://doi.org/10.1101/2021.12.16.473007 https://doi.org/10.1101/2021.12.16.473007.

60. Tareen, A., and Kinney, J.B. (2020). Logomaker: beautiful sequence logos in Python. Bioinforma. Oxf. Engl. *36*, 2272–2274. https://doi.org/10.1093/bioinformatics/btz921.

61. Schneider, T.D., and Stephens, R.M. (1990). Sequence logos: a new way to display consensus sequences. Nucleic Acids Res. *18*, 6097–6100.

62. Boccaletto, P., Machnicka, M.A., Purta, E., Piątkowski, P., Bagiński, B., Wirecki, T.K., de Crécy-Lagard, V., Ross, R., Limbach, P.A., Kotter, A., et al. (2018). MODOMICS: a database of RNA modification pathways. 2017 update. Nucleic Acids Res. *46*, D303–D307. https://doi.org/10.1093/nar/gkx1030.

63. Suzuki, T. (2021). The expanding world of tRNA modifications and their disease relevance. Nat. Rev. Mol. Cell Biol. *22*, 375–392. https://doi.org/10.1038/s41580-021-00342-0.

Figure 1

A GM12878-c305 Organoid Neural Development Overview

Aggregation
Day 2

Orbital Shaker
Day 18

Neural Induction | Neural Diff. | Neural Tissue

Aggrewell Media
(+SB/IWR)
Day 0

Sasai II (+FGF/EGF)
Day 14

Sasai III (+BDNF/NT3)
Day 35
Fragment Reads

Sasai IV
(+BDNF/NT3)
Day 70

RNA Extraction   ARM-seq   tRAX Analysis

GM12878
-c305

Day 0   Day 14   Day 35   Day 70

tDR Coverage
Profile Analysis

Visualization/
Interpretation

Classification   Preprocessing/Dimensionality Reduction

B Structure and Neural Rosettes

DAPI
VIM

DAPI
VIM

C Day 70 Organoid Neural Marker Gene Expression

CTIP2   TBR2

PAX6   DAPI

D D70 Organoid Neural Marker Merge

CTIP2 TBR2 PAX6

E ARM-seq (AlkB+) Small RNA Diff Expression D0 v D70

P-value <= 0.001
and log₂(FC) >= 1.5

- iPSC Favored
- Neural Favored
- Stem Markers
- Neural Markers
- tRNAs
- Small RNAs

miR-182-5p
miR-148a-3p
miR-200c-3p   SNORD93
miR-21-5p
miR-302a-5p   miR-302d-3p
miR-302c-3p   miR-378a-3p
miR-302a-3p   miR-151a-3p
miR-302b-3p   RN7SK

SNORD115
MT-RNR2
SNORA11
SNORD68   miR-9-5p
miR-9-3p
miR-99a-5p
RNA18SN

-log₁₀ (p-value)

log₂(fold change)

F C305 Small RNA PCA

Time point
- Day 70
- Day 35
- Day 14
- Day 0
- AlkB-
- AlkB+

Neural
Maturation

Early Neural
Diff.

Undiff.
Stem Cells

PCA 2 (15%)

PCA 1 (25%)

Figure 2

Figure 3

A Differential tRNA Fragment (tDR) Expression Day 0 vs Day 70

P-value <= 0.001
Log₂(FC) >= 1.5

- iPSC Favored (0-14 & 0-35 & 0-70)
- iPSC Favored (0-35 & 0-70 only)
- iPSC Favored (0-14 or 0-35 or 0-70)
- Neural Favored (0-14 & 0-35 & 0-70)
- Neural Favored (0-35 & 0-70 only)
- Neural Favored (0-14 or 0-35 or 0-70)

- ● 5' counts
- ■ 3' counts
- ▲ Other counts
- ◆ Whole counts
- ✕ Pre-tRNA counts

B Top Diff. Expressed tDRs Log₂ FC from D0

AlkB Treatment

C Top Neural-specific Fragment (tDR) Coverage

tRNA-SeC-TCA-1   tRNA-Arg-TCT-4   tRNA-Ala-AGC-8

D Unique Highly Expressed Gly Fragment (tDR) Coverage Change

tRNA-Gly-GCC-2   tRNA-Gly-CCC-2   tRNA-Gly-CCC-1

E Unique Fragment (tDR) Coverage Change

tRNA-Asp-GTC-1   tRNA-Gly-GCC-1   tRNA-Pro-TGG-2

Time point: Day 0, Day 14, Day 35, Day 70

Figure 4

A

Classification by Mean Sample Group Expressions

Figure 5