The two-process model: How simple, complex, unique is it from a mathematical perspective?

Anne C. Skeldon[1,2,3,*] and Derk-Jan Dijk[2,3,4]

[1]School of Mathematics & Physics, Faculty of Engineering and Physical Sciences, University of Surrey, Guildford, UK
[2]UK Dementia Research Institute, Care Research and Technology Centre at Imperial College, London and the University of Surrey, Guildford, UK
[3]NIHR Oxford Health Biomedical Research Centre, Oxford, UK
[4]Surrey Sleep Research Centre, School of Biosciences, Faculty of Health and Medical Sciences, University of Surrey, Guildford, UK
[*]Corresponding author

January 22, 2025

Abstract

    The two-process model (2pm) of sleep regulation is not only a conceptual framework but consists of mathematical equations that have meaning and structure. Despite the apparent simplicity of its graphical representation, the mathematical structure of the 2pm is complex and falls within the wider class of coupled oscillator models. The structure of the 2pm is very similar to models that have been used to capture other physiological phenomena such as cardiac arrhythmias, breathing and the integrate-and-fire behaviour of neurons. The 2pm is also intimately related to neuronal mutual inhibition models for the alternation between sleep and wake. Here we discuss how the relationship between neuronal models of sleep regulation and the 2pm model provides new insights into the physiological interpretation of the circadian thresholds of the 2pm and parameter values. We analyse the mathematical structure of the 2pm and how this structure relates to sleep patterns observed in the absence of 24-h time cues, in different species and in early childhood. We discuss how extending the 2pm to include a light-circadian pacemaker process allows for quantitative modelling of the effects of artificial light on sleep duration and sleep timing relative to the 24 hour day. Overall, we highlight how understanding the mathematical structure of the 2pm and its extensions provides alternative, not immediately intuitive, perspectives on the drivers of sleep phenotypes. We show that changes in sleep timing and the phase relationship between sleep and circadian rhythms may occur not only as a result of changes in the circadian pacemaker but also as a result of changes in sleep homeostasis, neuronal processes associated with excitability or the light environment. Analysis of the mathematical structure of the 2pm brings the opportunity to quantitatively predict the effect of interventions to support sleep and circadian alignment in individuals of all ages, including those with neurodegenerative disorders.

1 Introduction

One of the most widely reproduced figures in sleep and circadian science must be Fig. 4 from Alex Borbély's 1982 paper [1] that introduces the two-process model (2pm) of sleep regulation, re-plotted in Fig. 1a. Such is the strength of the idea of two processes regulating sleep and circadian rhythmicity that many data are interpreted within this conceptual framework. In the 1982 paper, this conceptual framework is presented as two processes which together capture sleep propensity and duration of recovery sleep. Subsequently, in 1984, Daan, Beersma & Borbély [2] introduced twin thresholds, one determining spontaneous sleep onset and one determining spontaneous sleep offset. The enduring appeal of the powerful two-process framework is perhaps in part due to the easily interpretable visual framing of a homeostatic sleep pressure that accumulates during wake and dissipates during sleep, see Fig. 1b. The separation into circadian and homeostatic components has repeatedly been shown to provide insight in sleep-wake regulation, for example [3, 4].

Although often represented graphically, the 2pm consists of mathematical equations with a precise meaning. For example, the model for the sleep process, process S, can be viewed as a chemical kinetic model in which the rate of change of homeostatic sleep pressure is a balance between clearance and production. Clearance is modelled as proportional to the instantaneous sleep pressure with rate constants $1/\chi_s$ h^{-1} and $1/\chi_w$ h^{-1} during sleep and wake respectively and production is at the fixed rate of μ_s/χ_s (units of homeostatic sleep pressure) h^{-1} during sleep and μ_w/χ_w (units of homeostatic sleep pressure) h^{-1} during wake. During wake, production is faster than clearance so homeostatic sleep pressure increases. During sleep, clearance is faster than production, so homeostatic sleep pressure decreases.[1]

The mathematical equations for the 2pm include a total of eight parameters: the time constants χ_s, χ_w; the 'accumulation' parameters μ_w and μ_s, also known as the upper and lower asymptotes; the mean levels of the upper (wake) threshold and lower (sleep) thresholds H_0^+ and H_0^- and the circadian amplitude a and period T_c. Daan, Beersma & Borbély considered non-dimensional versions of the equations which effectively sets μ_s to zero and $\mu_w \equiv \mu$ to 1. The full equations and a discussion of the scaling is given in the Supplementary Material Section 1.

---

[1]An alternative view is that there is no production during sleep, only clearance. This is equivalent to measuring sleep pressure relative to a baseline in which clearance and production are balanced.

The fact that the 2pm is not only a conceptual framework but also a mathematical model means it has quantitative predictive power. Indeed, Daan, Beersma & Borbély [2], showed that their model could explain sleep duration after sleep deprivation, internal desynchrony seen in time isolation experiments and enhancement and fragmentation of sleep in continuous bed rest. Subsequently, Beersma, Daan & Van den Hoofdakker [5] used the 2pm to argue that changes in sleep in depression could be explained by a reduced build-up rate of homeostatic sleep pressure during wake or increased noise in the wake threshold. Subsequently, extensive evaluation and modelling related to the build up and decay of homeostatic sleep pressure has been carried out in humans (e.g. [6]), rats (e.g. [7, 8]) as indexed by slow wake activity.

A key element of the 1982 paper is the use of slow wave activity (SWA, EEG power density in the 0.75-4.5 Hz range during NREM sleep) as a biomaker for sleep pressure. However, one potential criticism is that no physiological correlates of the levels of the thresholds or the circadian amplitude have yet been found. The precise shape of the threshold, mean values and amplitude of the circadian rhythm were selected to replicate patterns of human sleep. Daan, Beersma & Borbély postulate that the upper and lower thresholds may be due to the sensitivity of (unspecified) brain regions to sleep pressure. In general, they argue that in particular the upper threshold may be labile and respond to short term environmental and physiological changes. Subsequent work has shown that it is possible to relate the 2pm to mechanistic neuronal models, which gives further insight into possible physiological interpretations [9].

The 1984 paper [2] highlights the rich dynamics of the 2pm and its ability to simulate both monophasic and polyphasic sleep, see also Fig. 1c-e. From a mathematical perspective it is natural to ask what the underlying structure of the model is that leads to these different states, and then what does that mean for explaining data. In fact, around the time the 2pm was formulated, similar 'threshold' models were constructed for breathing (inspiration-expiration) [10], integrate and fire neuronal models [11], and cardiac arhymmias [12, 13]. All have the same underlying mathematical structures which can be formally analysed via circle maps [14]. This mathematical perspective highlights that the 2pm sits within the general class of systems which have a self-sustained oscillation with a so-called natural period (the sleep oscillator) which is driven by a periodic input (the circadian oscillator) [15]. All the structure that is standardly seen in such driven systems, such as tongue-like regions of entrainment, is then seen in the 2pm.

The 2pm describes the duration of sleep and the timing of sleep relative to the circadian minimum. When used to describe timing relative to the 24 hour day two important assumptions are made. First, that the circadian period T_c is entrained to 24 hours, so the periodicity of the circadian forcing T_f is 24 hours. Second, that the circadian minimum occurs at a specific clock-time. The 2pm does not include a mechanism to explain entrainment to 24 hours or the timing of the circadian pacemaker. Daan, Beersma & Borbély were aware of this limitation, but commented that there was insufficient knowledge of the phase response of humans to light to warrant inclusion of a light-circadian process. Consequently, that in its original form, the 2pm is limited in the extent to which it can explain sleep and circadian timing relative to the light-dark cycle. However, since 1984 more is known about the light-circadian entrainment process, which has led to extensions which include light [16].

In this paper we briefly discuss three aspects of the 2pm: first, the relation to neuronal mutual inhibition models of sleep-wake regulation, the consequent physiological interpretation of the thresholds and the wider implications; second, the mathematical structure that explains the internal desynchrony results, the appearance of bicircadian rhythms and polyphasic sleep, and third, extensions of the 2pm model to include interaction with the light-dark cycle. We end with a brief discussion of opportunities and future directions.

Figure 1: **The two process model**. S represents the homeostatic sleep pressure, $\overline{C}$ is the circadian drive for wakefulness. The grey shaded region highlights times when sleep occurs. **a** The two processes in the two-process model (2pm) as depicted in Fig. 4 from [1]. **b** The 2pm including upper and lower thresholds, as described in [2]. Homeostatic sleep pressure (black line) increases during wake and decreases during sleep. Switching between wake and sleep and sleep and wake occurs at upper and lower thresholds respectively (grey line). The thresholds are modulated according to a circadian rhythm. Here the circadian rhythm is assumed to be entrained to 24 hours. The timing of sleep is shown by the horizontal grey bars with the orange triangles indicating the times of the circadian minimum. Parameter values: $\chi_s = 4.2$ h, $\chi_w = 18.2$ h, $H_0^+ = 0.67$, $H_0^- = 0.17$, $a = 0.12$, $\mu = 1$. **c–e** Simulations for different values of the mean lower threshold and $\chi_s = 4.2$ h, $\chi_w = 18.2$ h, $H_0^+ = 0.75$, $a = 0.07$, $\mu = 1$. In each pair of panels the left hand panel shows the 2pm and the right hand panel shows a raster plot indicating the timing of sleep relative to the circadian minima (orange triangles). **c** Shows a repeating rhythm consisting of three sleep periods every four days, $H_0^- = 0.09$. **d** Shows a repeating rhythm consisting of three sleep periods every two days, $H_0^- = 0.50$. **e** Shows a polyphasic sleep pattern consisting of two sleep periods every day, $H_0^- = 0.58$.

2 Neuronal mechanisms underpining the two-process model

Since the 1980's we have learned much about the brain structures involved in sleep regulation. Saper, Scammell & Lu [17] conceptualised sleep-wake regulation as the result of the mutual inhibition of wake active and sleep active regions in the brain. This conceptual framework explains the stability of sleep and wake states: when awake, wake-active neurons fire, and by suppressing the firing of sleep-active neurons maintain wake. When asleep, sleep-active neurons fire, and by suppressing the firing of wake-active neurons, maintain sleep. Importantly, this mutual inhibition framework alone does not explain why switching between sleep and wake states occurs. Consquently, Saper, Scammell & Lu additionally proposed the existence of time dependent inputs to wake active and sleep active regions. These time dependent inputs reflect homeostatic sleep pressure and circadian rhythmicity. When these inputs are sufficiently strong, they overcome the inhibition and enable switching between states to occur.

The conceptual neuronal sleep-wake regulation framework was formulated in a mathematical model by Phillips and Robinson [18] with interactions as shown in Fig. 2a. Simulations of the Phillips-Robinson model capture the rise of sleep pressure during wake and decrease during sleep, the oscillations of the circadian rhythm and the so-called flip-flop between firing of sleep active and wake active neurons, see Fig. 2b-e.

The Phillips-Robinson model has two timescales. A fast timescale on which the neuronal dynamics act (seconds) and the slow timescales of sleep homeostasis and circadian rhythmicity (hours). This results in simulated sleep-wake cycles in which typically there are rapid transitions between sleep and wake, but that the states of sleep and wake last for hours, as in Fig. 2d,e. This multiple timescale structure means that an alternative representation of the Phillips-Robinson model is as a hysteresis loop, with transitions between wake and sleep and sleep and wake occurring at tipping points, see Fig.2f, [18, 19]. Furthermore, using a formal mathematical multiple timescale analysis, on the slow timescale the Phillips-Robinson model may be mapped to the 2pm [9]. In this mapping, the tipping points that occur in the neuronal model are equivalent to the switching points between wake and sleep in the 2pm.

Figure 2: **A neuronal mutual inhibition model and its relation to the two-process model. a** Neuronal structure underpinning sleep-wake regulation, based on the conceptual framework of Saper, Scammell & Lu [17] as framed by Phillips & Robinson [18]. **b–e** Simulation of the Phillips-Robinson model for standard parameter values. **b** Homeostatic sleep pressure increases during wake and decreases during sleep. **c** Sinusoidal representation of the circadian rhythm from the suprachiasmatic nucleus (SCN) entrained to 24 h. **d, e** Firing rate of wake active neurons. Firing rate is high during wake and low during sleep. **e** Firing rate of sleep active neurons. Firing rate is high during sleep and low during wake. **f** Hysteretic representation of the Phillips-Robinson model. In this representation, there is a folded surface which results from the mutual inhibition of the wake active and sleep active neuronal populations. The time dependent sleep drive results in a trajectory that evolves over the surface, switching between the wake (upper) surface to the sleep (lower) surface at the folds (tipping points) which are indicated by the black circles. **g** The Phillips-Robinson model in the form of the two-process model. The homeostatic sleep pressure (blue / red curve) increases during wake and decreases during sleep and is colour-coded as in panel **f**. The mean level of the upper (lower) threshold is given by the value of the drive at the upper (lower) fold in panel **f**. The oscillation of the thresholds is given by the circadian inhibitory input to the sleep drive.

Often the 2pm is described as the sleep homeostasis oscillating 'between' two thresholds, but the neuronal framework highlights that requiring sleep homeostasis to be between the thresholds is not a requirement for the model to successfully explain patterns of sleep behaviour. For example, Fig. 1b and Fig. 2g have the same sleep duration and relation between sleep and circadian timing but the gap between the thresholds in Fig. 2g is much smaller and the circadian amplitude much lower than in Fig.1b. In fact, the hysteretic structure suggests that at any one point, only one of the thresholds is of relevance i.e. the lower threshold when asleep (on the lower branch of the hysteresis loop) and the upper threshold when awake (on the upper branch of the hysteresis loop). This raises interesting interpretative differences. Based on a view that sleep and wake correspond to a hysteretic structure, the region between the thresholds is a region in which both sleep and wake states exist. Above the upper threshold, only the state of sleep can exist and below the lower threshold, only the state of wake can exist. The hysteretic structure may be alternatively viewed via a potential formulation, see Supplementary Fig. S1.

This hysteretic structure has at least three consequences. First, that between the two thresholds, switching between sleep and wake may occur through random perturbations. Second, when homeostatic sleep pressure reaches the upper threshold, either sleep ensues or some additional wake input (e.g. orexin) is required to shift the upper threshold. Third, if woken while still in the 'sleep only' region, then to remain awake requires additional wake drive to shift the upper threshold. This has been conceptualised as 'wake effort' [20, 21]. In modelling acute sleep deprivation, Daan, Beersma & Borbély viewed that the upper threshold was over-ridden. In the 'wake effort' view, the upper threshold is moved in order to allow wake to be maintained, as illustrated in Fig. 3. The fact that sleep pressure does not remain between the thresholds in the Phillips-Robinson parameter regime has implications for waking early. In the Daan, Beersma & Borbély parameter regime, waking early requires momentary stimulation to switch from one stable (sleep) to the other (wake). In the Phillips-Robinson parameter regime, wake can only be maintained by 'wake effort' shifting the position of the upper threshold. The requirement of additional effort to maintain wake remains until late enough in the day that the habitual rise in circadian drive for wakefulness means that wake is a stable state, see Fig. 3.

The correspondence between the 2pm and Saper's framework, via the Phillips-Robinson model, means that many of the results that have been found from simulating the Phillips-Robinson model can also be seen in simulations of the 2pm and vice versa.

Figure 3: **Wake effort.** In the neuronal interpretation of the two-process model, during wake, once the upper threshold has been reached requires effort i.e. additional input to wake active neurons which shifts the position of the upper threshold. If awoken while homeostatic sleep pressure is above the upper threshold, then again effort is required to shift the upper threshold. Regions in which wake effort is needed are highlighted in magenta.

Furthermore, there is a direct relationship between the parameters in the neuronal model and the levels of the thresholds in the 2pm, see Supplementary Fig. S2. For example, increasing the drive to wake active regions predominantly has the effect of raising the level of the upper threshold. Increasing the drive to sleep active regions lowers the mean value of the thresholds while leaving the distance between then unchanged, see Fig. 4. The positions of the thresholds are additionally affected by parameters that relate to the strength of the mutual inhibition and the neuronal firing rates.

Figure 4: **Increasing drive to sleep active neurons decreases the mean levels of the thresholds** Simulation of the Phillips-Robinson model [18] for different values of the drive to sleep active neurons. The gap between the thresholds in the 2pm model do not change, but lowering the thresholds increases sleep duration because of the exponential nature of the model for homeostatic sleep pressure.

3 The mathematical structure of the two-process model

The sleep-wake oscillator

In the absence of circadian rhythmicity (zero circadian amplitude), the 2pm reduces to a homeostatic sleep pressure oscillating between time-independent upper and lower thresholds, see Fig. 5a. This sleep-wake oscillator is also often described as a relaxation oscillator. The natural period T_nat of the oscillation can be formally derived in terms of parameters of the two-process model and is given by

$$T_\mathrm{nat} = T_\mathrm{sleep} + T_\mathrm{wake}, \tag{1}$$

where T_sleep is the duration of each sleep period and T_wake is the duration of each wake period. Here,

$$T_\mathrm{sleep} = \chi_s \log\left(\frac{H_0^+}{H_0^-}\right) \tag{2}$$

and

$$T_\mathrm{wake} = \chi_w \log\left(\frac{\mu - H_0^-}{\mu - H_0^+}\right), \tag{3}$$

where χ_s and χ_w are the homeostatic time constants for sleep and wake respectively, H_0^+ and H_0^- are the mean positions of the upper and lower thresholds and μ is the upper asymptote. For the standard parameters, $T_\mathrm{sleep} = 5.8$ h, $T_\mathrm{wake} = 16.8$ h, giving a natural period of $T_\mathrm{nat} = 22.6$ h in which approximately 26% is spent asleep.

The dependence of T_nat, T_sleep and T_wake on the parameters is shown in Fig. 5b-f. In line with the graphical and computational intuition described in Daan, Beersma & Borbély [2], χ_s changes sleep duration but has no impact on wake duration, χ_w and μ change wake duration but have no impact on sleep duration. The exponential nature of sleep homeostatic pressure, as modelled, means that the positions of the thresholds H_0+ and H_0^- alter both sleep and wake duration. The mean level of the thresholds also impact the average level of sleep pressure and the effects of sleep deprivation.

Figure 5: **Natural period of the sleep-wake oscillator. a** The two-process model (2pm) with the circadian amplitude set to zero. For the standard parameters ($\chi_s = 4.2$ h, $\chi_w = 18.2$ h, $H_0^+ = 0.67$, $H_0^- = 0.17$, $\mu = 1$) the period of the sleep-wake cycle is approximately 22.6 h, of which approximately 5.8. h is sleep ($\sim$26%). The right hand panel shows a raster plot with the timing of sleep on successive 24 h days. **b-f** Illustration of the dependence of the natural period on the parameters in the 2pm, as given by equations (1)-(3). The vertical red line indicates the position of the standard parameters.

Entrainment of the sleep-wake oscillator by the circadian oscillator

In the presence of circadian rhythmicity (circadian amplitude $a > 0$), the 2pm consists of a sleep-wake oscillator that is forced by the circadian oscillator. Characteristic of nonlinear forced oscillator systems, if the natural period T_{nat} of the sleep-wake oscillator is sufficiently close to the forcing period T_f, and the amplitude of the forcing is sufficiently high then entrained monophasic sleep-wake cycles of period T_f result. If the forcing is weak (i.e. low circadian amplitude) then only a small range of natural periods T_{nat} can be entrained to T_f. The stronger the forcing, the larger the range of natural periods can be entrained. This means that, in a (natural period)–(forcing amplitude) parameter plane, regions where entrained sleep-wake patterns consisting of one sleep-wake cycle occurs in every circadian period (monophasic sleep), form a tongue-like region, for example see Fig. 6a. We note that tongue-like entrainment regions are typical of forced systems of self-sustained oscillators: a similar tongue-like entrainment structure is seen in models of the circadian system forced by the light-dark cycle [22].

As observed by Daan, Beersma & Borbély [2], the ostensibly 'simple' 2pm exhibits many other solution types. Indeed, entrainment theory predicts that there exists an entrainment tongue in which n sleep-wake oscillations occur in m periods of the circadian drive for every integer pair (n, m). For example, the (2,1) tongue consists of sleep-wake cycles in which two sleeps occur in each circadian period. The (1,2) tongue consists of sleep-wake cycles in which one sleep occurs every two circadian periods. The largest tongue is the (1,1) tongue, corresponding to monophasic sleep patterns.

This mathematical structure has a very particular ordering, as shown in the example in Fig. 6a. For sufficiently small circadian amplitudes, this structure is precisely ordered, solutions are unique, complicated but not chaotic. For higher circadian amplitudes, it is to be expected that solutions become chaotic and that more than one solution may exist for the same parameter values. Within each tongue, the phase relationship between the sleep-wake cycle and the circadian rhythm changes. This is easiest to see in the large tongue for monophasic sleep. When the natural sleep-wake period is longer than the forcing period, the forcing has to speed up the sleep-wake oscillator, with the effect that the sleep-wake oscillator tends to be 'behind' the circadian oscillator in phase. Conversely, when the natural sleep-wake period is shorter than the forcing period, the forcing has to slow down the sleep-wake oscillator, with the effect that the sleep-wake oscillator tends to be 'ahead of' the circadian oscillator in phase. The net effect is that on

the side of the tongue with the longest natural period (right-hand edge for Fig. 6a) the circadian minimum occurs at the beginning of the night, see Fig. 6f. For the side of the tongue with the shortest natural period, the circadian minimum occurs at the end of the night, see Fig. 6e. The example shown comes from varying the mean level of the upper threshold. Varying the mean level of the lower threshold, the homeostatic time constants or the value of the upper asymptote similarly change the natural period. Varying any one of these parameters results in a similar set of tongues, but squashed or stretched because the relationship between the natural period T_{nat} and the parameters differs, see Supplementary Fig. S3.

This mathematical structure follows from entrainment theory because the 2pm is an example of a mathematical object called a circle map [23, 9]. Circle maps were first introduced by Arnold, [12], and the dynamics of circle maps have been extensively analysed since [24, 25, 26] and also in the particular context of the 2pm [15]. This 'tongue-like' structure is shared by threshold models for other systems [10, 11, 12, 13]. and can be formally analysed via circle maps [14]. Understanding the entrainment structure facilitates understanding and interpreting data, we discuss three examples below.

Figure 6: **Entrainment of the sleep-wake oscillator by the circadian oscillator: Arnold tongues.** **a** Number of sleeps per day as a function of the mean level of the upper threshold, H_0^+ and the circadian amplitude, generated by simulating the 2pm for approximately 110,000 different parameter values. The remaining parameters are: $\chi_s = 6$ h, $\chi_w = 18$ h, $H_0^- = 0.17$, $\mu = 1$. The largest region corresponds to monophasic sleep. In principle there are regions of n sleeps per day for m circadian periods for every integer pair (n,m) for every n/m. In practice, the upper and lower limits of the values for n/m are bounded by the choices of the other parameters. For the example shown $0.068 < n/m < 2.54$ days. Each region forms a tongue with the tip of the tongue occuring at zero circadian amplitude for the value of H_0^+ which gives a natural period of m/n. **b-g** Examples of simulations for different values of H_0^+ at a fixed circadian amplitude of $a = 0.07$, i.e. taking a horizontal line across panel **a**. In each case the two-process model along with a raster plot is shown. The orange triangles mark the time of the circadian minimum. **b** Polyphasic sleep pattern. **c** Two sleeps a day. **d** Monophasic sleep with the circadian minimum occuring near the end of the night. **e** Monophasic sleep with the circadian minimum occuring near the beginning of the night. **f** Four sleeps every five days. **g** One sleep every two days.

Internal desynchrony and bicircadian rhythms

Time isolation experiments in which knowledge of clock time and all access to environmental timing cues are removed have been critical in the understanding of circadian rhythmicity and sleep-wake regulation. Many experiments were carried out in purpose build facilities [27, 28, 29] and in caves [30]. Strogatz [31] summarised many of the available data and evaluated the ability of various oscillator models to explain the results. In time isolation experiments, an immediate effect is a loss of synchrony with the 24-hour day and are replaced by rhythms where sleep and circadian rhythms are in synchrony but typically have a 25 hour period. The phase of the minimum of the core body temperature was observed to shift from the end of the night to the beginning of the night. Subsequently, in many subjects internal desynchrony was observed, in which the circadian cycle typically became slightly shorter than 25 hours but the sleep-wake cycle desynchronised oscillating with a period that was either substantially longer or shorter than the circadian cycle. In some cases, a transition to bicircadian rhythms in which there is one sleep every two days, was observed.

Considering the underlying structure of the 2pm, as illustrated in Fig. 6 provides plausible explanations for how such sleep-wake patterns may occur. The initial loss of synchrony with the 24-h day results in an immediate shift in period of the circadian oscillator from its entrained value of 24 hours to 25 hours. The overall tongue-like entrainment structure remains, but with small quantitative differences, for example, the minimum of the monophasic sleep tongue occurs at 25/24 instead of one day. Indeed, a formal mathematical scaling gives that changing from a forcing period of $T_f = 24$ h to $T_f = 25$ h is equivalent to rescaling time to circadian days rather than calendar days and scaling the homeostatic time constants by a factor of 25/24 i.e. 4%. The immediate change from $T_f = 24$ h to $T_f = 25$ h therefore results in sleep-wake patterns which are entrained to the new forcing period of 25 h, see Supplementary Fig. S4a and b.

The observed change in phase of sleep relative to the circadian pacemaker can be explained by either a reduction in circadian amplitude or by a lengthening of the natural period T_{nat} see Supplementary Fig. S4c.

Note that changing the circadian amplitude has minimal change in the percentage of time spent asleep. In the model, changing the natural period can either increase of decrease the percentage of time spent asleep depending on which parameter is varied. The

occurrence of internal desynchrony occurs in regions close to the boundary of the monophasic sleep tongue. Formally, in the 2pm, the desynchronised patterns are generically periodic patterns of period much longer than the circadian period. For low circadian amplitude, the duration of sleep will be approximately the same every day. For higher circadian amplitudes, the duration of sleep will depend on the timing of sleep relative to the timing of the circadian rhythm resulting in occasional long sleeps, see Supplementary Fig. S4d and e. This latter phenomena is analogous to 'relative coordination' seen in desynchrony of circadian rhythms to the 24-hour day [32].

Finally, to explain the occurrence of bicircadian rhythms within the 2pm framework requires there to be a lengthening of the natural period of the sleep-wake oscillator T_{nat}. Since the period is given by equation (1), lengthening of the period occurs if the homeostatic time constants increase, the upper threshold increases, the lower threshold decreases (see Fig. 6a and Supplementary Fig. S4g) or the upper asymptote decreases. It is not immediately evident which parameter most plausibly explains the observed phenomena, although further using the percentage of time spent in sleep could establish which may be most plausible. Daan, Beersma & Borbély primarily focussed on varying the level of the upper threshold and the circadian amplitude [2]. Subsequently, Daan *et al* highlighted that low mean rectal temperature (low metabolic rate) was associated with long sleep-wake periods and high mean rectal temperature (high metabolic rate) was associated with short sleep-wake periods [33]. For the 2pm, this would suggest that higher metabolic rate are reflected in shorter homeostatic time constants, or a decreased gap between the thresholds (lower level of the upper threshold or higher level of the lower threshold) or a higher value of the upper asymptote.

In their neuronal model, Phillips-Robinson model [34] showed that internal desynchrony and bicircadian rhythms could be explained by changing the circadian amplitude and increasing the drive to wake active neurons during wake. The close correspondence between the Phillips-Robinson model and the 2pm, increasing the drive to wake promoting neurons is effectively the same as increasing the mean level of the upper threshold, see Supplementary Fig. S2.

Polyphasic sleep in different species

Daan, Beersma & Borbély highlighted that the 2pm can explain polyphasic sleep patterns [2], as also shown in Fig. 1 and Fig. 6. In particular they noted that when the thresholds

are close together polyphasic sleep occurs. Similarly, Phillips, Chen and Robinson [35] argued that their mutual inhibtion neuronal model can model observed patterns of sleep in different species. They showed that decreasing the homeostatic time constant for wake and sleep caused a shift from monophasic to polyphasic sleep and that increasing the mean drive to sleep active neurons increased the percentage of time spent asleep.

The mathematical structure of the 2pm adds further insight to both of these sets of observations. Reducing the gap between the thresholds reduces the natural period of the sleep-wake cycle T_{nat}, see equation (1) and Supplementary Fig. S6. When the natural period is short, periodic sleep-wake patterns which consist of multiple sleeps per day occur because of the underlying entrainment structure, as illustrated in Fig. 6.

The relation between neuronal model the 2pm, means that the same underlying mathematical structure is at work. Similarly, reducing the homeostatic time constants reduces the natural period, resulting in polyphasic rhythms for small values of the time constants. Increasing the mean level of the drive to sleep active neurons is equivalent to lowering the mean level of the thresholds, which increases the percentage of the natural period in which time is spent asleep.

The mathematical structure in addition highlights that when the circadian amplitude is low, polyphasic sleep will be distributed evenly across the 24-hour day with little variance in duration of one sleep-wake cycle and the next. In contrast, when the circadian amplitude is high, sleep episodes will predominantly occur only at some circadian phases and the duration of sleep-wake cycles will vary with time-of-day, see [36] and also Supplementary Fig. S5.

Transitions in early childhood

Developmental changes across early childhood result in changes from polyphasic sleep to monophasic sleep. The mathematical structure of the 2pm further suggests that this could be explained by physiological changes that lead to a gradual lengthening of the natural-sleep wake period rather than a change in the circadian process. In this progression, the theoretical structure predicts that periods of relatively regular sleep patterns followed by periods of transition, as the Arnold tongue structure is traversed [9]. For example in the shift from a pattern of three sleeps a day (a main nocturnal sleep, a morning nap, an afternoon nap) to two sleeps a day (a main nocturnal sleep and an

afternoon nap) it is to be expected that there will be a progression when first there are only occasional days on which only two sleeps occur through to a point where there are only occasional days in which three sleeps occur. The sequence follows a so-called Devil's staircase, see Fig. 7.

Figure 7: **Devil's staircase.** As the natural period is increased for fixed circadian amplitude the number of sleeps per day follows a pattern known as a 'Devil's staircase'. Here this is illustrated for the fixed parameter values $\chi_s = 4.2$ h, $\chi_w = 18.2$ h, $H_0^- = 0.17$, $\mu = 1$ and $a = 0.12$ by varying the value of the upper threshold H_0^+ i.e. taking a horizontal line across Fig. 6a. Note the curvature of the tongues means that the plateau at, for example, $T_{\mathrm{nat}} = 1/3$ corresponds to 2 sleeps a day. **b-f** Example raster plots for different values of H_0^+.

4 Timing of sleep relative to the 24 hour day: extending the two-process model to include the light-dark cycle

Extensive laboratory experiments have shown that light is the primary synchroniser ('zeitgeber' i.e. 'time giver') of the human circadian clock. Many experiments have quantified the effect of light, and mathematical equations that quantitatively describe the phase and dose response of the human circadian pacemaker to light have been derived, originally by Kronauer [37, 22] and subsequently modified by others, for example [38, 39]. The conceptual and mathematical understanding is that the light-circadian system is a forced oscillator system in which the intrinsic circadian oscillation is entrained by the light-dark cycle.

The circadian-light forced oscillator system, in common with other nonlinear forced oscillator systems including the sleep-circadian forced oscillator system, has an underlying mathematical structure in which there are entrainment regions that form tongue-like regions in the (natural period)–(forcing strength) parameter space [22, 40, 21]. If the light-dark cycle is only a weak zeitgeber i.e. there is insufficient contrast between light that is available during the day and light that is available during the night, then only a small range of intrinsic circadian periods close to 24 hours can be entrained. The stronger the forcing, the larger the range of circadian periods which can be entrained. Consequently, the entrainment regions form tongue-like structures. Typically of human relevance, the focus is usually on the (1,1) tongue in which one 24 hour light-dark cycle results in one 24 hour entrained circadian oscillation, but in principle other tongues exist. Analogous to internal desynchrony for the sleep-circadian oscillator system, desynchrony with respect the 24 hour day occurs when the light-dark cycle is too weak or the intrinsic period is too far from 24 hours. An additional feature, which is unique to humans and critical for understanding human chronotypes, is that access to electric light enables us to turn light on after dusk. This ability to self-select our lighting environment means that there is feedback between sleep-wake timing and the zeitgeber. This feedback results in a contraction of the entrainment region [21].

Within a neuronal framework, Phillips, Chen & Robinson extended the Phillips-Robinson model to include light [41]. This sleep-circadian-light neuronal framework has been used to model the effects of self-selection of light on sleep timing and further extended to include the effects of social constraints which restrict the opportunities to sleep [21]. This

sleep-circadian-light model has been shown to capture individual sleep phenotypes [42] and for scenario testing of, for example, the effect on social jetlag of moving school start times versus changing the light environment. Using the neuronal framework, plausible physiological reasons to explain changes in sleep duration and timing across the lifespan have been proposed [43]. Studying sleep in those living with schizophrenia, it has been suggested that a plausible reason for long sleep duration is a greater sleep propensity resulting in reduced levels of the mean thresholds in a two-process representation. Furthermore, fitting to individuals, highlighted that observed seasonal desynchrony could be plausibly explained by reduced light exposure in the winter months [42].

Inspired by the neuronal sleep-circadian-light neuronal framework and the close link between neuronal models and the 2pm, a homeostatic-light-circadian (HCL) model has been proposed, as represented schematically in Fig. 8. The HCL model combines the graphical simplicity of the original 2pm with a light-circadian entrainment process and social constraints [16]. At the same time, the shape of the thresholds was fitted to data for the circadian drive for wakefulness, as reported in [4]. The updated shape of the upper threshold better captures well-established features of sleep-wake regulation such as the wake maintenance zone [44]. It has been shown that combining the HCL model with data, personalised models which capture individual average sleep duration and timing can be constructed. In [16], the model was used to develop a method to separate out the relative contributions of physiology and environmental light in determining sleep timing. These personalised models could be used to design personalised light interventions to adjust sleep timing [42, 16].

Figure 8: **The homeostatic-circadian-light (HCL) model.** Reproduced from [16]. In the HCL model, light, gated by sleep, is fed into a model which captures the phase and dose response of light on the circadian pacemaker [38]. The circadian pacemaker then results in a work propensity rhythm which, when combined with sleep homeostasis, results in patterns of sleep and wake.

5 Discussion

One enduring appeal of the 2pm is its graphical simplicity which gives an intuitive understanding of how sleep duration depends on time-of-circadian-day. Aside from its use to frame experimental results, all current models of fatigue risk management are to a greater or lesser extent based on the principles of sleep homeostasis and circadian rhythmicity [45].

Interestingly, although the 2pm is often represented as a dichotomy of two separate processes, sleep homeostasis and circadian rhythmicity, and prompted the 'separation' of the processes in forced desynchrony protocols [46, 47], the 2pm itself is all about interactions. It is the interaction of the two processes that explains, for example, the duration of recovery sleep following sleep deprivation. It is the interaction that means that polyphasic sleep occurs at consolidated circadian phases when the circadian amplitude is high [36].

Indeed, it is also not so easy to separate out what are 'homeostatic' and 'circadian' parameters [48]. Specifically, of the eight parameters in the 2pm, the rate constants χ_w and χ_s which represent the clearance rate of homeostatic sleep pressure during wake and sleep respectively may be described as homeostatic parameters. Similarly, one may argue that μ_w and μ_s which represent the rate of build-up of homeostatic sleep pressure during wake and sleep respectively are homeostatic parameters. The circadian amplitude a is evidently a 'circadian' parameter. However, this leaves the two threshold parameters, the mean levels of the upper and lower thresholds, H_0^+ and H_0^- respectively, which have also commonly been considered circadian parameters [2]. Relating the 2pm to neuronal mutual inhibition models suggests that these thresholds may relate to properties of the neuronal mutual inhibition structure of sleep-wake regulation and inputs to wake promoting and sleep promoting regions — and may therefore be neither circadian or homeostatic parameters.

Linking the 2pm to neuronal mutual inhibition model also raises the question as to what is 'the two-process model', is it a specific set of parameters? Conceptualising the 2pm as instead a hysteretic loop suggests that there is no physiological reason for homeostatic sleep pressure to remain 'between' the thresholds. It also highlights the fact that the 2pm may alternatively be posed as a double-well potential problem in which the time dependent drives change the probability of states of wake and sleep occurring.

The 2pm has a deep mathematical structure that is inherent to nonlinear forced oscillator systems, as highlighted in Fig. 6 [15]. Understanding the mathematical entrainment structure establishes what determines different sleep-wake patterns in the 2pm. For sufficiently small circadian parameters (where 'sufficiently small' has a precise definition in terms of the parameters [15]) stable solutions are unique, so the only way that different patterns can be observed is by changing parameters. The sequence of transitions seen in early childhood has recently been mapped in two different neuronal models [49, 50]. Both inherit mathematical structure from the 2pm. For high circadian amplitudes, multiple solutions may be possible signifying that different patterns could occur as a change of state rather than a change of parameters. It is also likely that there are regions of parameter space in which monophasic sleep-wake patters are predicted but with alternating long and short sleep duration.

A further open question is the optimal parameters to describe sleep-wake regulation. For example, Fig. 1b and Fig. 2g slow wave activity (SWA) is a marker of sleep homeostasis [1, 51, 6], but it may or may not be that the time constants found by fitting to the timecourse of SWA across the night are the same as the time constants of a homeostatic sleep process which determines sleep timing. The time constants used in the Phillips-Robinson model are typically much longer (∼45 hours) yet can still replicate similar patterns of sleep. The use of longer time constants may be appropriate when, for example, simulating the effects of acute and partial sleep deprivation on performance and sleepiness [52, 53].

Understanding the mathematical structure can also challenge our pre-conceptions on what causes phenotypes of phenotypic change. For example, changes from polyphasic to monophasic sleep may not primarily be a consequences of maturation of the circadian system [9, 49]. Changes in sleep timing as we age may be a consequence of changes in circadian amplitude and / or sleep homeostasis and not changes in circadian period [43]. Differences in sleep duration may be a consequence of differences in drive to wake or drive to sleep [42], and not because of changes in homeostatic time constants. Desynchrony with the 24-hour day may not be due to clock-malfunction but due to insufficient zeitgeber strength [42].

Finally in Fig. 1 of their paper, Daan, Beersma & Borbély set out a scheme for their model [2]. In this scheme, the model included the effect of the light-dark cycle to synchronise the circadian oscillator and conscious decisions affecting the thresholds as well as the

interactions of sleep homeostasis and circadian rhythmicity. With extensions of the 2pm to include input from the light-dark cycle we are perhaps now somewhat closer to this vision [16]. Even though the 2pm does not capture basic features of sleep-wake regulation such as the REM—non-REM cycle, there remains much promise with a 'four-process' framework which includes sleep homeostasis, circadian rhythmicity, light and dynamic thresholds which encompass acute and chronic effects of stimulants [54]. Most of all, it is of value to recognise that the 2pm and its extensions are not only conceptual frameworks, but quantitative mathematical models which can be fitted and challenged by data. By challenging models in this quantiative way brings the opportunity to evaluate whether or not they can really explain the available data and what is missing. This, ultimately may inform the understanding the wide variety of sleep phenotypes, sleep-wake disorders and the development of countermeasures for situations in which the sleep-wake and circadian system is challenged by social constraints.

Acknowledgments

The authors are supported by the UK Dementia Research Institute [award number UKDRI-7206] through UK DRI Ltd, principally funded by the UK Medical Research Council, and additional funding partner Alzheimer's Society and also by the National Institute for Health Research (NIHR) Oxford Health Biomedical Research Centre (BRC), (NIHR203316). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health. We thank Stanisław Biber for feedback on the paper.

References

[1] Borbély, A. A. A two process model of sleep regulation. *Hum. Neurobiol.* **1**, 195–204 (1982).

[2] Daan, S., Beersma, D. G. M. & Borbély, A. A. Timing of human sleep: Recovery process gated by a circadian pacemaker. *Am. J. Physiol.* **246**, R161–R183 (1984).

[3] Dijk, D.-J. & Czeisler, C. A. Contribution of the circadian pacemaker and the sleep homeostat to sleep propensity, sleep structure, electroencephalographic slow waves, and sleep spindle activity in humans. *J. Neurosci.* **15**, 3526–3538 (1995).

[4] Dijk, D.-J., Duffy, J. F., Riel, E., Shanahan, T. L. & Czeisler, C. A. Ageing and the circadian and homeostatic regulation of human sleep during forced desynchrony of rest, melatonin and temperature rhythms. *J. Physiol.* **516**, 611–627 (1999).

[5] Beersma, D. G., Daan, S. & Van den Hoofdakker, R. H. The timing of sleep in depression: theoretical considerations. *Psychiatry Res.* **16**, 253–62 (1985).

[6] Achermann, P., Dijk, D.-J., Brunner, D. P. & Borbély, A. A. A model of human sleep homeostasis based on eeg slow-wave activity: Quantitative comparison of data and simulations. *Brain Res. Bull.* **31**, 97–113 (1993).

[7] Franken, P., Tobler, I. & Borbély, A. A. Sleep homeostasis in the rat: Simulation of the time course of eeg slow-wave activity. *Neurosci. Lett.* **130**, 141–144 (1991).

[8] Guillaumin, M. C. C. *et al.* Cortical region–specific sleep homeostasis in mice: effects of time of day and waking experience. *Sleep* **41**, zsy079 (2018).

[9] Skeldon, A. C., Dijk, D.-J. & Derks, G. Mathematical models for sleep-wake dynamics: Comparison of the two-process model and a mutual inhibition neuronal model. *PLOS ONE* **9**, 1–16 (2014).

[10] Glass, L. & Mackey, M. C. A simple model for phase locking of biological oscillators. *J. Math. Biology* **7**, 339–352 (1979).

[11] Bressloff, P. C. & Stark, J. Neuronal dynamics based on discontinuous circle maps. *Phys. Lett. A* **150**, 187–195 (1990).

[12] Arnold, V. I. Cardiac arrhythmias and circle mappings. *Chaos, reprinted from Arnold's thesis (Moscow 1959)* **1**, 20–21 (1991).

[13] Glass, L. Cardiac arrhythmias and circle maps - a classical problem. *Chaos* **1**, 13–19 (1991).

[14] Derks, G., Glendinning, P. A. & Skeldon, A. C. Creation of discontinuities in circle maps. *Proc. Roy. Soc.* **477**, 20200872 (2021).

[15] Bailey, M. P., Derks, G. & Skeldon, A. C. Circle maps with gaps: Understanding the dynamics of the two-process model for sleep–wake regulation. *Eur. J. App. Math.* **29**, 845–868 (2018).

[16] Skeldon, A. C. *et al.* Method to determine whether sleep phenotypes are driven by endogenous circadian rhythms or environmental light by combining longitudinal data and personalised mathematical models. *PLOS Comput. Biol.* **19**, 1–33 (2023).

[17] Saper, C. B., Scammell, T. E. & Lu, J. Hypothalamic regulation of sleep and circadian rhythms. *Nature* **437**, 1257–63 (2005).

[18] Phillips, A. J. K. & Robinson, P. A. A quantitative model of sleep-wake dynamics based on the physiology of the brainstem ascending arousal system. *J. Biol. Rhythms* **22**, 167–179 (2007).

[19] Fulcher, B. D., Phillips, A. J. K. & Robinson, P. A. Modeling the impact of impulsive stimuli on sleep-wake dynamics. *Phys. Rev. E* **78**, 051920 (2008).

[20] Fulcher, B. D., Phillips, A. J. K. & Robinson, P. A. Quantitative physiologically based modeling of subjective fatigue during sleep deprivation. *J. Theor. Biol.* **264**, 407–419 (2010).

[21] Skeldon, A. C., Phillips, A. J. K. & Dijk, D.-J. The effects of self-selected light-dark cycles and social constraints on human sleep and circadian timing: a modeling approach. *Sci. Rep.* **7**, 45158 (2017).

[22] Kronauer, R. E., Forger, D. B. & Jewett, M. E. Quantifying human circadian pacemaker response to brief, extended, and repeated light stimuli over the phototopic range. *J. Biol. Rhythms* **14**, 501–516 (1999).

[23] Nakao, M. & Yamamoto, M. Bifurcation properties of the two process model. *Psychiatry Clin. Neurosci.* **52**, 131–3 (1998).

[24] Katok, A. & de Hasselblatt, B. *Introduction to the Modern Theory of Dynamical Systems* (CUP, Cambridge, 1995).

[25] Boyland, P. Bifurcations of circle maps: Arnol'd tongues, bistability and rotation intervals. *Comm. Math. Phys.* **106**, 353–381 (1986).

[26] Mackay, R. S. & Tresser, C. Transition to topological chaos in circle maps. *Physica D* **19**, 206–237 (1986).

[27] Aschoff, J. Circadian rhythms in man. *Science* **148**, 1427–1432 (1965).

[28] Wever, R. *The circadian system of man: results of experiments under temporal isolation* (Springer Verlag, 1979).

[29] Czeisler, C. A., Weitzman, E. D., Moore-Ede, M. C., Zimmerman, J. C. & Knauer, R. S. Human sleep: Its duration and organization depend on its circadian phase. *Science* **210**, 1264–1267 (1980).

[30] Siffre, M. *Experiences Hors du Temps* (Fayard, Paris, 1972).

[31] Strogatz, S. *The Mathematical Structure of the Human Sleep-Wake Cycle* (Springer Verlag, 1986).

[32] Emens, J. S., Lewy, A. J., Lefler, B. J. & Sack, R. L. Relative coordination to unknown "weak zeitgebers" in free-running blind individuals. *J. Biol. Rhythms* **20**, 159–167 (2005).

[33] Daan, S., Honma, S. & Honma, K. Body temperature predicts the direction of internal desynchronization in humans isolated from time cues. *J. Biol. Rhythms* **28**, 403–411 (2013).

[34] Phillips, A. J. K., Czeisler, C. A. & Klerman, E. B. Revisiting spontaneous internal desynchrony using a quantitative model of sleep physiology. *J. Biol. Rhythms* **26**, 441–453 (2011).

[35] Phillips, A. J. K., Robinson, P. A., Kedziora, D. J. & Abeysuriya, R. G. Mammalian sleep dynamics: how diverse features arise from a common physiological framework. *PLoS. Comput. Biol.* **6**, e1000826 (2010b).

[36] Franken, P. & Dijk, D.-J. Sleep and circadian rhythmicity as entangled processes serving homeostasis. *Nat. Rev.Neurosci.* **25**, 43–59 (2024).

[37] Kronauer, R. E. A quantitative model for the effects of light on the amplitude and phase of the deep circadian pacemaker, based on human data. In Horne, J. (ed.) *Sleep '90, Proceedings of the Tenth European Congress on Sleep Research*, 306–309 (Bochum: Pontenagel, 1990).

[38] Forger, D. B., Jewett, M. E. & Kronauer, R. E. A simpler model of the human circadian pacemaker. *J. Biol. Rhythms* **14**, 533–537 (1999).

[39] St. Hilaire, M. A. *et al.* Addition of a non-photic component to a light-based mathematical model of the human circadian pacemaker. *J. Theor. Biol.* **247**, 583–599 (2007).

[40] Granada, A. E., Bordyugov, G., Kramer, A. & Herzel, H. Human chronotypes from a theoretical perspective. *PLoS One* e59464 (2013).

[41] Phillips, A. J. K., Chen, P. Y. & Robinson, P. A. Probing the mechanisms of chronotype using quantitative modeling. *J. Biol. Rhythms* **25**, 217–227 (2010).

[42] Skeldon, A. C., Dijk, D.-J., Meyer, N. & Wulff, K. Extracting Circadian and Sleep Parameters from Longitudinal Data in Schizophrenia for the Design of Pragmatic Light Interventions. *Schiz. Bull.* **48**, 447–456 (2022).

[43] Skeldon, A., Derks, G. & Dijk, D.-J. Modelling changes in sleep timing and duration across the lifespan: Changes in circadian rhythmicity or sleep homeostasis? *Sleep Med. Rev.* **28**, 96 – 107 (2016).

[44] Strogatz, S. H., Kronauer, R. E. & Czeisler, C. A. Circadian pacemaker interferes with sleep onset at specific times each day: role in insomnia. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* **253**, R172–R178 (1987).

[45] Riedy, S., Hursh, S. R., Dawson, D., Balkin, T. J. & Van Dongen, H. P. A. Sleep, sleepiness, and performance prediction modelling. In Kryger, M., Roth, T., Goldstein, C. A. & Dement, W. C. (eds.) *Principles and Practice of Sleep Medicine*, chap. 83, 780 –787 (Elsevier, 2022).

[46] Czeisler, C. A. *et al.* Stability, precision, and near-24-hour period of the human circadian pacemaker. *Science* **284**, 2177–2181 (1999).

[47] Wang, W. *et al.* Desynchronizing the sleep-wake cycle from circadian timing to assess their separate contributions to physiology and behaviour and to estimate intrinsic circadian period. *Nat Protoc* **18**, 579–603 (2023).

[48] Skeldon, A. C. & Dijk, D.-J. Modeling drosophila sleep: fly in the sky? *Sleep* **47**, zsad309 (2024).

[49] Webb, L., Phillips, A. J. K. & Roberts, J. A. Mapping the physiological changes in sleep regulation across infancy and young childhood. *PLoS Comput Biol.* **20**, e1012541 (2024).

[50] Athanasouli, C., Stowe, S. R., LeBourgeois, M. K., Booth, V. & Diniz Behn, C. G. Data-driven mathematical modeling of sleep consolidation in early childhood. *J. Theor. Biol.* **593**, 111892 (2024).

[51] Dijk, D.-J., Brunner, D. P. & Borbély, A. Time course of EEG power density during long sleep in humans. *Am. J. Physiol.* **258**, R650–61 (1990).

[52] McCauley, P. *et al.* A new mathematical model for the homeostatic effects of sleep loss on neurobehavioral performance. *J. Theor. Biol.* **256**, 227–239 (2009).

[53] Postnova, S., Lockley, S. W. & Robinson, P. A. Prediction of cognitive performance and subjective sleepiness using a model of arousal dynamics. *J. Biol. Rhythms* **33**, 203–218 (2018).

[54] Wyatt, J. K., Cajochen, C., Cecco, A. R.-D., Czeisler, C. A. & Dijk, D.-J. Low-dose repeated caffeine administration for circadian-phase-dependent performance degradation during extended wakefulness. *Sleep* **27**, 374–381 (2004).