1 **Early-life paternal relationships predict adult female survival in wild baboons**

2 David A.W.A.M. Jansen[1,2], J. Kinyua Warutere[3], Jenny Tung[4-9], Susan C. Alberts[4,5,6], Elizabeth A.
3 Archie[1*]
4
[1] Department of Biological Sciences, University of Notre Dame, Notre Dame, IN, USA
[2] Department of Pathobiological Science, School of Veterinary Medicine, University of Wisconsin-Madison,
United States
[3] Amboseli Baboon Research Project, Amboseli National Park, Kajiado, Kenya
[4] Department of Evolutionary Anthropology, Duke University, Durham, NC, USA
[5] Department of Biology, Duke University, Durham, NC, USA
[6] Duke University Population Research Institute, Duke University, Durham, NC, USA
[7] Department of Primate Behavior and Evolution, Max Planck Institute for Evolutionary Anthropology,
04103 Leipzig, Germany
[8] Canadian Institute for Advanced Research, Toronto, Canada M5G 1M1, Canada
[9] Faculty of Life Sciences, Institute of Biology, Leipzig University, Leipzig, Germany
16
17 * Corresponding author: earchie@nd.edu
18
19

ABSTRACT

Parent-offspring relationships can have profound effects on offspring behavior, health, and fitness in
adulthood. These effects are strong when parents make heavy investments in offspring care.
However, in some mammals, including several species of carnivores, rodents, and primates, fathers
live and socialize with offspring, but paternal care *per se* is subtle or indirect. Do these limited father-
offspring relationships also affect later-life outcomes for offspring? Working in a well-studied
baboon population where males contribute little direct offspring care, we found that juvenile female
baboons who had stronger paternal relationships, or who co-resided longer with their fathers, led
adult lives that were 2-4 years longer than females with weak or short paternal relationships. This
pattern did not differ between females who experienced high versus low levels of early-life adversity;
hence, paternal relationships were not especially protective against harsh early environments. Males'
relationships were strongest with juvenile females they were most likely to have sired and when
males had few current mating opportunities. Hence, father-daughter relationships may be
constrained by male mating effort. Because survival predicts female fitness, fathers and their
daughters may experience selection to engage socially and stay close in daughters' early lives.
35

36 **Keywords**: paternal care, parental care, mammals, early-life effects, fathers, adult survival

1. INTRODUCTION

In humans and other mammals, social environments are powerful determinants of individual health, survival, and fitness [1]. Social relationships in early life are especially important, both because of their immediate benefits to offspring—such as opportunities to learn social skills, gain resources, or receive protection—and also because these relationships have lasting consequences for adult health and survival [reviewed in 2]. Maternal relationships are especially well-studied in this regard [3-12], and across mammals, maternal loss and the quality of maternal care can have long-lasting consequences for offspring gene regulation, stress reactivity, social integration, and adult survival [3-9, 11-13].

But what about relationships with fathers? Early-life paternal social effects have received less attention, in part because it is rare for male mammals to make substantial investments in offspring care [14, 15]. However, in species with caring males, early-life father-offspring relationships can have profound effects on offspring in adulthood. In humans, paternal absence in childhood predicts negative outcomes for adult behavior, income, health, and mortality risk [e.g., 16, 17-20]. In rodents with biparental care, fathers affect the complexity of offspring social environments, with consequences for neurological development and adult behavior [reviewed in 21, 22]. But species in which fathers provide substantial care are unusual among mammals. In a wider (but still unusual) set of group-living mammals, fathers live and even socialize with offspring, but paternal care *per se* is subtle and often indirect [14, 23]. These species include several carnivores and equids, as well as gorillas, chimpanzees, baboons, and other primates [14, 18, 24-30]. Whether these more limited early-life paternal relationships have long-term consequences for offspring is largely unknown.

Here we test if early-life paternal relationships predict adult survival for female baboons—a species where fathers and their juvenile offspring may co-reside and interact, but where mothers provide all essential care. Baboons are useful for testing these relationships for three reasons. First, the baboons we studied live in polygynandrous mating systems where paternity certainty is incomplete, yet adult male baboons often maintain proximity to and interact with their offspring [31-38]. Further, lactating female baboons sometimes form close social bonds (i.e., "primary associations") with particular males, and these relationships are better explained by parenting than mating effort [33-35, 39-41]. In support, male primary associates typically do not sire their female associate's next infant [34, but see 35]. Furthermore, male primary associates are disproportionately the fathers of their female partner's current infant, intervene on behalf of females and their infants in conflicts, and may buffer infants from rough handling [31-38, 42].

Second, proximity to adult males and/or paternal presence in early life have developmental
and social consequences for young baboons. For instance, proximity to adult males, including
fathers and non-fathers, increases the complexity of the social environment for infant baboons [43].
Paternal presence is also correlated with earlier sexual maturity in daughters [31] and predicts
stronger social bonds between paternal half-siblings [44].
Third, adult lifespan, our outcome of interest, explains 80%-90% of the variance in lifetime
reproductive success for female baboons in our population [7, 45, 46]. Hence, if early-life paternal
relationships influence the lifespan of daughters, they may have important consequences for their
daughters' fitness and, by extension, the males themselves. If so, males and their daughters may
experience selection to form and maintain early-life social relationships with one another.
Working in the Amboseli baboon population in Kenya [47], we asked whether female
baboons who had stronger grooming relationships or longer co-residency with their fathers in early
life exhibit higher adult survival. We pursued three objectives. First, we measured patterns of
grooming and co-residency between juvenile females and their fathers. We measured grooming
because social bonds are often developed and maintained through grooming, a primary affiliative
behavior in many social species, including baboons [48-50]. We measured co-residency because
father-offspring pairs who live together for longer have more time to interact (co-residency varies
because males may disperse or die during their daughters' juvenile years).
Second, we tested whether females had stronger grooming relationships or longer co-
residency with their fathers in early life exhibit higher adult survival than females who had
weak/short paternal grooming or co-residency. In the Amboseli baboons, adult female longevity is
also predicted by an accumulation of harsh conditions in early life, such as drought, maternal loss, or
having a low-ranking or socially isolated mother [7, 45, 51]. Hence, we also used interaction effects
to test if early-life relationships with fathers might protect daughters from the negative effects of
cumulative early-life adversity.
Third, we tested why some fathers are more likely to groom or have longer co-residency
with their daughters than others. We predicted that males would have stronger relationships with
their daughters when they had high paternity certainty (e.g., spent more time mate guarding his
daughter's mother when she was conceived) and when reproductive tradeoffs are favorable (e.g.,
when the male had few current mating opportunities). Together, our results support the importance
of paternal presence and father-daughter social bonds to adult female baboons, lending context to
paternal effects on adult outcomes and the evolution of mammalian parental care.

2. METHODS

(a) Study population and subjects

Our subjects were wild baboons studied by the Amboseli Baboon Research Project (ABRP) in the Amboseli ecosystem, Kenya [47]. This population is admixed between yellow and anubis baboons (*Papio cynocephalus* and *P. anubis*), with majority yellow ancestry [52, 53]. ABRP observers collect behavioral and demographic data year-round on a near-daily basis, and all study animals are known based on visual recognition. Our analyses centered on 216 female baboons that: (i) survived the first 4 years of life, encompassing most or all of the juvenile period for females [median age at menarche in Amboseli = 4.5 years; 54]; (ii) had known mothers and fathers assigned using combined demographic and genetic data (see below); and (iii) had complete information on their experience of six sources of early-life adversity that together predict adult mortality [7, 51]: early maternal loss, maternal dominance rank, maternal social isolation, early-life drought, a close-in-age younger sibling, or large group size (see below). The females in our study were born into 13 different social groups, which are the fission or fusion products of two original study groups, first studied in 1971 and 1981.

(b) Assigning maternities and paternities

The 216 female subjects were born to 117 mothers and sired by 102 fathers. Maternities are known from near-daily demographic records. Paternity assignments were based on microsatellite genotypes from at least six microsatellite loci and demographic records used to identify an initial pool of candidate fathers. These methods are described in detail in previous studies [32, 55-57], but briefly, microsatellite genotypes for juvenile females, mothers, and potential fathers were analyzed in the likelihood-based paternity assignment program CERVUS [58, 59]. We first included all potential fathers residing in the mother's social group at the time of conception (potential fathers are any male in the adult male hierarchy), and then expanded the set of potential fathers to include all adult, ranked males in the population. These two sets of males typically identified the same father, but in the few cases where they disagreed, we assigned paternity based on the strength of additional evidence for each father (e.g., geographic distance between the candidates and the mother, male health status, or density of observations on the mother during the window of conception). In all cases, the assigned father resided in same social group as the mother at the time of conception [56]. Levels of confidence for all CERVUS analyses were set at 95%, and paternity assignments were robust across three rates of error, 1%, 5%, and 10% [32, 55-57].

(c) Measuring grooming and co-residency between juvenile females and adult males

We defined co-residency as the cumulative number of days each juvenile female lived in the same social group with her genetically confirmed father during the first 4 years of her life.

Following [6, 60], we measured annual pairwise grooming bonds for each year of the female's life, using the "dyadic sociality index" (DSI), which provides a numeric score for each juvenile female's dyadic relationship strength with individual adult males, scaled to be directly comparable to all other juvenile female-male pairs in the population in a given year of the female's life, birthday to birthday. We calculated three types of DSI scores for juvenile females, which differed in the males included in the calculations: DSI_{all} measured dyadic relationships between juvenile females and all adult males who lived in her social group for at least 30 days in the year in question, including her father (i.e., "co-resident" males); $DSI_{paternal}$ measured dyadic relationships between juvenile females and their fathers; and $DSI_{non-paternal}$ measured dyadic relationships between juvenile females and all co-resident males, excluding her father.

The grooming interactions used to calculate the DSI scores were collected via representative interaction sampling. During this sampling, observers moved through the group recording all grooming interactions in their line of sight, while conducting 10-min focal animal follows on a randomized list of females and juveniles [6, 60]. From these data, we calculated each dyad's log-transformed daily rate of grooming in a given year. All log-transformed daily rates were then regressed against a measure of observer effort in that year (the number of focal animal samples per female-day). The resulting residuals were z-scored within years to estimate the DSI. Negative DSI scores reflect dyads who groomed less than was typical in that year; positive DSI scores reflect dyads who groomed more than was typical in that year.

Our third objective required us to test why some fathers are more likely to groom their daughters than other males. For these analyses we compiled data on the presence or absence of male grooming directed to their daughters, in a given juvenile female year of life (contingent on ≥30 days of co-residency).

(d) Measuring early-life adversity

To test if fathers moderate early-life effects on female mortality in adulthood (objective 2), we measured early-life adversity using a cumulative adversity index developed in prior studies [6, 7, 45, 51, 61]. This index sums the presence of six sources of early-life adversity: (1) drought in the first

year of life (<200 mm of annual rainfall); (2) maternal death in the first four years of life; (3) being

born into a large group as an index of realized resource competition (group size in the top quartile;

≥36 adults); (4) the birth of a close-in-age younger sibling that may divert maternal resources

(interbirth interval in the shortest quartile, <1.5 years after the focal female's own birth); (5) being

born to a mother whose ordinal social dominance rank is in the bottom quartile of her group; and

(6) being born to a mother who is in the top quartile for social isolation over the first 2 years of the

juvenile's life, measured based on an overall index of her involvement in grooming [7]. For each of

the 216 juvenile females, we summed the number of these conditions that applied, resulting in a final

index that could range from zero to six. No subject experienced more than four sources of adversity

(20.3% of the 216 females experienced 0 sources of adversity; 40.2% experienced 1 source; 24.1%

experienced 2 sources; 11.6% experienced 3 sources, and 3.7% experienced 4 sources of adversity).

(e) Measuring predictors of father-daughter grooming and co-residency

For our third objective, we tested why some fathers are more likely to groom their juvenile

daughters than other fathers. For a sub-set of variables, we also tested whether they explained the

duration of father-daughter co-residency. Our sample sizes for these analyses were smaller than the

216 females in the first two objectives because we lacked information on some variables (see below).

Paternal age and the ages of daughters and their mothers were known from near daily demographic

records. All 216 daughters had ages accurate within a few days. Of the 102 fathers in our data, 43

(42.2%) were born into the study population and their ages were accurate within a few days. For the

remaining 59 fathers (57.8%), their ages were estimated to within a few years by comparing them to

known-age males from the population [62]. For the mothers, 101 (86.3%) had ages accurate within a

few days, 14 (12.0%) had ages accurate within 3 months, and 2 (1.7%) had ages accurate within 3

years.

Male ordinal dominance rank determines male priority of access to mates in our population, and

high-ranking males are expected to prioritize mating over parenting effort [63]. In Amboseli, male

ranks are calculated monthly based on decided dyadic agonistic encounters between adult males [64].

The *daily rate of fertile females* in the group could influence a male's mating opportunities and

time spent grooming daughters. This variable was calculated as the average daily number of peri-

ovulatory females in the group in a given juvenile female-year on the days the male was resident in

the social group [65]. Peri-ovulatory periods are inferred from continuous records of sexual skin

swellings that increase in size during the follicular phase and decrease during the luteal phase [65].

The *proportion of the mother's available consort time the male obtained* during the 5-day peri-ovulatory
period that led to the focal female's conception could affect male paternity certainty. This measure
was calculated by summing all observed consort time that a mother had with any adult male within 5
days before the likely conception date of the focal female. We then calculated the proportion of this
consort time that was monopolized by the male in question. Conception dates were calculated as
described previously based on obvious signs of female reproductive state [65, 66]. For 31 of the 216
females, no males were observed consorting the focal female's mother; hence analyses with this
variable only included 185 females.

The *number of potential fathers* present in the group at the juvenile female's conception could
influence male paternity certainty [56, 67]. This variable was calculated as the number of adult
ranked males present in the group during the 5-day peri-ovulatory period when the female was
conceived.

Following [34, 35]*, males who sired previous or future offspring of the juvenile female's mother* might be
more likely to groom their daughters if "primary associates" represent male mating effort. To test
this possibility, we identified all cases in which the focal female's father sired their mother's previous
or next offspring (n=185 females; see above). There were 25 cases (13.5%) where the father sired
the mother's previous offspring, 131 cases (70.8%) where he did not, 16 cases (8.6%) where the dad
was not present, and 13 cases (7%) where the paternity of the mother's previous offspring was
unknown. With respect to the mother's next offspring, there were 25 cases (13.5%) where the male
sired the mother's next offspring, 115 cases (62.2%) where he did not, 16 cases (8.6%) where the dad
was not present, and 29 cases (15.7%) where paternity was unknown. To avoid further reductions in
sample size, we imputed the 42 unknown paternities by randomly assigning paternity to one of the
adult ranked males present in the group when the female was conceived. We performed these
imputations 1000 times and ran our models on each imputed data set. We also repeated our analysis
on a smaller "complete cases" data set where paternities for the previous and subsequent offspring
were known.

The *number of offspring the male had in the social group (i.e., co-resident offspring-years)* could influence
his likelihood of remaining in his daughter's social group. This variable was calculated, for each
father, as the number of his juvenile offspring that were alive in the group in a given juvenile female-
year, scaled for days of co-residency. This variable may underestimate the male's true count of living
juvenile offspring because paternity is often missed for the youngest offspring (we generally obtain
the first fecal sample between 6 and 18 months of age).

The female's experience of *cumulative early-life adversity*, could also influence paternal

investment. Cumulative adversity was calculated as the sum of the 6 conditions a female might have

experienced prior to age 4 years (see above).

Finally, for our model of why some fathers groom their daughters more than others, *we also*

modeled observer effort, measured as the number of focal animal samples we collected per female-day.

We considered the effect of observer effort to account for the fact that observers are more likely to

observe grooming in groups where we conducted more intensive behavioral observations.

(f) Statistical Analyses

Most analyses were performed in R 4.4.0 using the packages lme4 [68], lmerTest [69], lmtest

[70], MuMIn [71], rptR [72], and survival [73]. For a full list of packages used in the analysis of this

paper see the GitHub repository in our data statement.

Objective 1: Characterizing patterns of grooming and co-residency between juvenile females, their fathers, and other

adult males

To measure grooming relationships between juvenile females and adult males, we calculated,

in each female-year (i) the average number of adult males each female groomed with, and (ii) the

percentage of grooming interactions she initiated with adult males (N=216 females; from 0 to 4

years of age). To test if female age predicted grooming initiation with males (both fathers and non-

fathers), we ran a binomial LMM where the response variable measured whether grooming with

adult males was initiated by females (1) or not (0), as a function of female age in each year of the

juvenile period. Female identity was modeled as a random effect. To test if grooming bonds were

stronger between father-daughter pairs than other male-female pairs, we also ran a LMM testing if

juvenile females' DSI_{all} scores with all adult, co-resident males, including their fathers, was predicted

by whether the male was the female's father or by female age (0 to 4 years of age), again modeling

female identity as a random effect.

Objective 2: Testing if paternal co-residency and social bonds predict adult female survival

To test if females who have longer juvenile co-residency or stronger grooming relationships

with their fathers exhibit higher adult survival than females who had short paternal co-residency or

weak paternal grooming relationships, we ran a series of Cox proportional hazards models where the

response variable was each female's age at death or censorship, contingent on survival to her 4[th] year

of life (N=216 females; 124 censored values). Models were fit using coxph in the R package survival

[73]. We tested which variables best explained variation in adult female mortality risk based on

Akaike information criteria (AICc), including each female's: (i) average annual dyadic grooming

bond strength with her father ($DSI_{paternal}$) during the first four years of her life (because grooming

patterns change with juvenile female age, these strengths were z-scored across females, within age

class [i.e., age 0 to 1 years, age 1 to 2 years etc.]); (ii and iii) average annual grooming bond strength

with all co-resident adult males, either excluding ($DSI_{non-paternal}$) or including her father (DSI_{all}; these

scores were also z-scored and averaged across the first four years of life); (iv) cumulative years of co-

residency with her father in the first four years of life; and (v) cumulative early-life adversity score in

the first four years of life.

To test if the effects of early-life adversity on adult female survival are moderated by paternal

grooming or co-residency, we added an interaction effect between females' early-life adversity scores

and (vi) their average annual $DSI_{paternal}$ scores, and (7) their cumulative years of paternal co-residency.

We also tested if juvenile females with strong paternal grooming bonds are socially well-

connected with females and males in adulthood. If so, early-life relationships between female

baboons and their fathers might be important for later life survival because they influence female

social connectivity in adulthood. Social connectivity to adult females and males predicts adult female

survival in this population [51, 60, 74]. Social connectivity was calculated as a social connectedness

index (SCI), which reflects the total amount of grooming the female, as an adult, gave and received

with other adult females (SCI_F) and adult males (SCI_M) in her social group [51, 60, 74]. For these

analyses, we first ran an LMM testing if females with stronger mean $DSI_{paternal}$ scores in the first 4

years of life had stronger SCI_F and SCI_M scores in adulthood, controlling for her age and dominance

rank and modeling female identity as a random effect. We then ran a series of survival models to test

if the observed association between $DSI_{paternal}$ and adult female survival is attenuated by adding SCI_F

and SCI_M to the model. The sample size for these models was 194 females because adult social

connectedness information was missing for 22 females.

Objective 3: Testing the predictors of father-daughter grooming and co-residency

To test why some fathers are more likely to groom their daughters than others, we ran a

binomial LMM where the response variable was whether a given father was observed to groom (1)

or did not groom (0) his daughter in each of the first four years of her life, contingent on 30 days of

co-residency in that year. The fixed effects were: (i-iii) the ages of the juvenile, father, and mother at

the start of the year; (iv) the father's average ordinal dominance rank in that year; (v) the average

number of fertile (i.e., peri-ovulatory) females in the group each day in that year; (vi) the proportion

of observed consort time the male had with the focal female's mother during the 5-day period when

she was conceived; (vii) the number of other adult males (i.e., potential fathers) in the group at the

daughter's conception; (viii and ix) whether the father sired a prior or future offspring with the

daughter's mother; (x) the number of juvenile paternal offspring the father had in the group in that

year ("co-resident offspring years"); (xi) the daughter's cumulative early-life adversity score; and (xii)

observer effort. Paternal identity was a random effect (N=495 father-years years involving 81 fathers

and 169 juvenile females). Because these models used imputed paternity assignments for some of

the prior/future offspring, we present the median of the model estimates, test statistics, and p-values

across 1000 imputed data sets. In the supplement, we also present a "complete cases" version of this

model without imputation (N=379 father-years years involving 70 fathers and 130 juvenile females).

We also performed a parallel analysis for all co-resident males (i.e., not just father-daughter pairs),

which included a binary fixed effect for if the adult male was the father (N=8,078 male-years years,

involving 297 males and 185 juvenile females). This analysis used imputed paternity assignments for

some of the prior/future offspring. We present the median of the model estimates, test statistics,

and p-values across 1000 imputed data sets.

To test why some fathers have longer co-residency with their daughters, we ran a LMM

where the response variable was the number of days the father resided in the same group as his

daughter in the first four years of her life. The fixed effects were: (i-ii) the father's and mother's ages

in the month that father-daughter co-residency ended (juvenile age was excluded because it was

colinear with the duration of co-residency); (iii) the father's dominance rank in the month co-

residency ended; (iv) the daily rate of fertile females in the group in the month co-residency ended;

(v) the proportion of consort time the male had with the focal female's mother during the 5-day

period when the daughter was conceived; (vi) the number of other adult males (i.e., potential fathers)

in the group at the daughter's conception; (vii) whether the father sired a prior offspring with the

daughter's mother (future offspring was excluded because only males with relatively long co-

residencies would be able to sire this future offspring); (viii) the number of juvenile paternal

offspring the father had in the group in the year the co-residency ended (co-resident offspring years);

and (ix) the daughter's cumulative early-life adversity score. Paternal identity was included as a

random effect (N=166 co-residencies involving 86 fathers and 166 juvenile females).

Before performing our analyses, we checked for multicollinearity using variance inflation

factor (VIF) analysis adapted for lmer models [75]. No variables had VIF > 2.5.

3. RESULTS

(a) Objective 1: Patterns of co-residency and grooming between juvenile females and their

fathers

The median co-residency between the 216 daughters and their fathers was 33 months (**Fig.**

1A; range=0-48 months). More than a third of these females (n=80; 37%) lived in the same group

with their father for ≥ 3 of their juvenile years, while for the remaining 63% of females (n=136),

their fathers either left the group or died sometime between their conception and 3 years of age

(**Fig. 1A**). Thirteen females (6%) never co-resided with their fathers because the male dispersed or

died between the focal female's conception and birth (**Fig. 1A**).

Grooming between juvenile females, their fathers, and other adult males changed in

frequency and directionality across females' juvenile years. From birth to 4 years of age, females

groomed with an increasing number of adult males (β=0.35; p=<0.001) and were more likely to

initiate grooming with adult males (**Fig. 1B**: binomial LMM: β=1.00; p<0.001). Among females in

the first year of life, 18.2% of the observed grooming interactions with adult males were initiated by

the females (**Fig. 1B**), and each female groomed on average with 1.15 adult males (range=1-3

males). However, by the 4th year of life, females had, on average, 1.54 male grooming partners

(range=1-6 males), and 83% of these interactions were initiated by the female (**Fig. 1B**).

Consistent with prior evidence that males and their offspring have differentiated

relationships [31-36], daughters' DSI$_{all}$ values were significantly stronger with their fathers than with

other co-resident adult males (**Table 1;** β=0.117; p=0.0047). However, the effect size was weak:

DSI$_{all}$ values between juvenile females and their fathers were just 4.29% stronger than DSI$_{all}$ between

juvenile females and non-paternal males, reflecting the fact that juvenile females also groom adult

males who are not their fathers, and these bonds are sometimes as strong (**Fig. S1**).

Table 1. LMM explaining dyadic bond strength between juvenile females and all co-resident adult males in a

given year (N=4,968 DSI$_{all}$ values between 216 females and 298 males, including 91 fathers).

Term	β (SE)	t	df	p-value	Interpretation
Age of juvenile female	0.134 (0.010)	13.103	10578.7	<0.001	↑juvenile age ↑bond strength
Male is the father	0.117 (0.042)	2.827	10527.1	0.0047	male is father ↑bond strength

Figure 1. Patterns of co-residency and grooming directionality between juvenile females, their fathers, and other adult males. **(A)** Cumulative paternal co-residency (x-axis) for the 216 juvenile females (y-axis). Each black bar represents the cumulative duration of time one female lived in the same group with her father. Red dashed lines demarcate the percentages of females who resided with their fathers for 1, 2, 3 or 4 years. **(B)** The average proportion of grooming interactions initiated by juvenile females (top: dark blue; bottom: red) or adult males (top: light blue; bottom: orange) as a function of female age. Top panel shows grooming initiation for fathers; bottom panel shows grooming initiation with all adult males.

(b) Objective 2: Early-life grooming and co-residency with fathers predicts adult female survival

We next tested whether daughters' early-life relationships with their fathers predicted their adult survival. We found that juvenile females who had relatively strong $DSI_{paternal}$ scores with their fathers, or who had relatively long co-residency with their fathers, or both, led longer adult lives than females with weaker paternal relationships (**Fig. 2**; **Table 2**). In support, the top three models

predicting adult female survival included either the female's average $DSI_{paternal}$ score in the first four

years of life, the duration of co-residency with her father, or both variables (**Fig. 2**; **Table 2 rows A-**

C). $DSI_{paternal}$ and co-residency were positively correlated with each other (**Fig. S2**; Pearson's

r=0.274, p=0.001), consistent with the idea that father-daughter pairs who co-reside for longer will

also have stronger grooming relationships. Models that included either one or both variables were

interchangeable in their ability to explain adult female mortality (**Fig. 2**; **Table 2 rows A-C;** range in

$\Delta AICc$=0.31-1.28). This effect was specific to $DSI_{paternal}$: strong social relationships during the

juvenile period with adult males in general (DSI_{all}), or with non-fathers ($DSI_{non-paternal}$), did not predict

adult survival (**Table 2 rows A-C** versus **rows D, F, and G**).

Early-life adversity also predicted adult female mortality (**Table 2 all models,** [7, 51]), but

the models that included $DSI_{paternal}$ and/or co-residency were a better fit to the data than a model

that only included early-life adversity (**Table 2 rows A-C vs row E**; range in $\Delta AICc$=3.228-4.56).

We therefore asked whether relationships with fathers predicted adult female survival more so for

females who experienced harsh early-life circumstances. However, adding an interaction effect

between female early-life adversity and either $DSI_{paternal}$ or paternal co-residency did not improve

model fit (range in $\Delta AICc$=1.66-3.57; **Table S1**). Hence, the effects of paternal relationships and

early-life adversity appear to be mostly independent. For the "median" female who experienced one

major source of early-life adversity, having a mean $DSI_{paternal}$ score in the top quartile for the

population predicted a median difference in survival of 1.8 years compared to females in the bottom

quartile of mean $DSI_{paternal}$ scores (**Fig. 2B**). Adult females who experienced one source of adversity

and lived with their fathers for all 4 years were predicted to live 2.6 years longer than females who

only lived with their father for 1 year (**Fig. 2C**). Females who experienced 3 or more sources of

early-life adversity were predicted to live 4.3 to 4.6 years longer if they co-resided with their father

for 4 years versus 1 year or had grooming relationships with their fathers in the top versus bottom

quartile (**Figs. 2B and 2C**).

Finally, we wondered whether some of the survival effects we observed could be explained

by adult females' social bonds with either sex in adulthood, both of which predict adult female

survival [51, 60, 74, 76]. This effect might arise if, for instance, juvenile females who have strong

$DSI_{paternal}$ scores tend to be more social overall, or if strong paternal bonds directly promote adult

social connectedness (e.g., a mediation effect that operates via social effects on health in adulthood).

We found that females who had stronger average $DSI_{paternal}$ scores across the first 4 years of life also

tended to have strong social bonds with both adult males (SCI_M; **Table S2**; β=0.16; p<0.001) and

adult females (SCI$_F$; **Table S2**; β=0.11; p=0.035) in adulthood. In support of the idea that DSI$_{paternal}$,

SCI$_F$, and SCI$_M$ all contribute to adult mortality risk, four of the seven best models in **Table S3**

(those with ΔAICc <2 from the best model) included a metric of adult social connectedness (SCI$_F$,

SCI$_M$, or both; **Table S3 rows 3, 4, 6 and 7**), while six of the best seven models included DSI$_{paternal}$,

paternal co-residency, or both variables (**Table S3 rows 1-6**).

Table 2. Results from seven alternative Cox proportional hazards models (n=216 females with 124 censored

values) showing predictors of adult female survival. Each cell shows the variable's hazard ratio (and 95%

confidence interval). Models are ordered by AICc. No model violated the proportional hazards assumption

(p-values ranged from 0.34 to 0.77).

Model	Cumulative early-life adversity	Mean DSI$_{paternal}$	Years of co-residency with father	Mean DSI$_{non-paternal}$	Mean DSI$_{all}$	AICc	ΔAICc
A	1.262 (1.004-1.586)	0.787 (0.603-1.026)	0.885 (0.759-1.032)	-	-	810.67	0.00
B	1.278 (1.016-1.608)	0.738 (0.573-0.951)	-	-	-	810.98	0.31
C	1.25 (0.995-1.571)	-	0.84 (0.725-0.973)	-	-	811.95	1.28
D	1.277 (1.014-1.608)	0.743 (0.576-0.959)	-	0.92 (0.675-1.255)	-	812.84	2.17
E	1.273 (1.012-1.601)	-	-	-	-	815.23	4.56
F	1.27 (1.008-1.6)	-	-	0.88 (0.648-1.195)	-	816.65	5.98
G	1.273 (1.01-1.604)	-	-	-	0.82 (0.616-1.092)	815.48	4.82

Figure 2. Juvenile females' paternal grooming relationships and co-residency predict their adult survival. (A) Estimates from the time-to-event effects on the hazard of death in adult female baboons. Effects and 95% CI are from **row A in Table 2**, the model that had the lowest AICc and that includes mean DSI$_{paternal}$ (green), paternal co-residency (blue) and early-life adversity (grey). **(B)** Predicted survival curves showing the effects of juvenile females' mean DSI$_{paternal}$ and early-life adversity on adult female survival (Wald test P= 0.006, N= 216; predictions from model B in Table 2). Dark green lines are females in the top quartile of DSI$_{paternal}$ scores; light green lines show females in the bottom quartile of DSI$_{paternal}$. Solid lines show females who experienced one source of early-life adversity; dashed lines show females who experienced 3 sources of adversity. **(C)** Predicted survival curves showing the effects of juvenile females' duration of co-residency with their fathers and early-life adversity on adult female survival (Wald test P= 0.007, N= 216; predictions from model C in **Table 2**). Dark blue lines show females who lived with their father for 1 year; light blue lines show females who lived with their fathers for all 4 juvenile years. Solid lines show females who experienced one source of adversity; dashed lines show females who experienced three sources of adversity.

(c) Objective 3: Fathers are more likely to groom and live with their daughters when paternity is more certain and reproductive opportunities are limited

Father-daughter relationships should be stronger when males have greater paternity certainty

and fewer reproductive opportunities. In support, males were most likely to groom their daughters

in a given year if the male was low-ranking, there were relatively few cycling females in the group

that year, the male had a higher proportion of consort time with the female's mother during the

cycle the daughter was conceived, and if the male had more offspring in the group (**Table 3; Table**

S4). Consistent with prior studies that suggest that male "primary associate" behavior is better

explained by parenting than mating effort [33, 35, 36], males who groomed their daughters were not

more likely to sire the mother's previous or next infant (**Table 3; Table S4**). A similar subset of

these variables also explained whether adult males had a grooming interaction with a juvenile female,

regardless of whether the male was the father (**Table S5**), suggesting that a male's rank and mating

behavior at the time of an infant's conception may influence his behavior towards that infant,

regardless of whether he is the father.

A slightly different set of variables predicted the duration of co-residency between fathers

and their daughters (**Table 4**). Adult males had longer co-residencies with their daughters if the

males were older, the mother was older, if there were more cycling females in the group that year of

the juvenile's life, and if they had more offspring in the group (**Table 4**). Longer co-residencies also

tended to occur if the male had a prior offspring with the female's mother, but this effect was not

significant (**Table 4**).

Table 3. Binomial GLMM of whether a male did (1) or did not groom (0) his juvenile daughter in a year of

her life (N=495 years; 81 fathers and 169 juvenile females with at least 30 days of co-residency >1 year).

Term	β (SE)	t-value	p-value	interpretation
Juvenile age	0.869 (0.209)	4.162	<0.001	↑juvenile age ↑grooming
Paternal age	0.201 (0.191)	1.057	0.291	
Maternal age	0.061 (0.085)	0.703	0.481	
Father's average ordinal rank	0.175 (0.058)	3.036	0.002	higher ordinal rank ↓grooming
Daily rate of fertile females	-9.342 (2.034)	-4.609	<0.001	↑fertile females ↓grooming
Proportion of consort time	2.334 (0.816)	2.881	0.004	↑consort time ↑grooming
Number of potential fathers at conception	0.030 (0.078)	0.381	0.704	
Father had future offspring with the mother	-0.470 (0.572)	-0.796	0.085	trend: will have offspring ↑grooming
Father had previous offspring with the mother	-0.835 (0.802)	-0.892	0.284	
Father's co-resident offspring-years	0.237 (0.045)	5.227	<0.001	↑co-resident offspring ↑grooming

	β (SE)	t-value	p-value	
Daughter's cumulative adversity	-0.059 (0.353)	-0.171	0.861	
Observer effort	1.711 (0.239)	7.168	<0.001	↑effort ↑grooming

Table 4. LMM of the duration of father-daughter co-residency during the daughter's 4-year juvenile period (N=166 co-residencies between 166 juvenile females and 86 fathers).

Term	β (SE)	t-value	p-value	interpretation
Maternal age	0.068 (0.021)	3.180	0.002	↑maternal age ↑co-residency
Paternal age	0.153 (0.051)	2.985	0.004	↑male age ↑co-residency
Father's ordinal rank	0.01 (0.040)	0.256	0.799	
Daily rate of fertile females	2.00 (0.994)	2.013	0.046	↑fertile females ↑co-residency
Proportion of consort time	0.009 (0.209)	0.045	0.9638	
Number of potential fathers at conception	-0.011 (0.022)	-0.523	0.602	
Father had previous offspring with the mother	-0.421 (0.238)	-1.769	0.080	trend: had offspring ↑co-residency
Father's co-resident offspring-years	0.102 (0.037)	2.795	0.006	↑offspring ↑ co-residency
Daughter's cumulative adversity score	-0.027 (0.097)	-0.284	0.777	

4. DISCUSSION

In many group-living mammals, males selectively interact with and provide low-cost forms of care to offspring [14, 23]. The selective pressures shaping these behaviors, and their importance to offspring health and survival, have received considerable attention in baboons [28, 31-42]. Here we report that the strength of early-life paternal social relationships predicts meaningful differences in adult survival for female baboons in Amboseli, Kenya. These differences are on the order of 2 to 4 years for adult females in the top versus bottom quartile for paternal grooming or co-residency time—effect sizes that are comparable to those for other major predictors of adult survival in Amboseli baboons, such as social isolation and early-life adversity [7, 60, 77]. This result joins prior evidence for early-life paternal effects in baboons, including the observations that paternal presence predicts earlier sexual maturity in daughters [31] and stronger social bonds between paternal half-siblings [44]. Other mammal species also have subtle or indirect forms of paternal care [14, 18, 24-30]; hence, early-life paternal social relationships may have long-term consequences for offspring in a greater number of species than is currently appreciated.

(a) What mechanisms explain early-life paternal social effects on female survival?

The two measures of father-offspring relationships we focused on—co-residency and
grooming—are not themselves parental care, raising the question: how do these measures lead to
early-life paternal effects on offspring? One possibility is that father-daughter grooming and co-
residency may be correlated with other male care behaviors, which in turn, directly benefit their
daughter's health and longevity. Prior studies show that male care in baboons includes intervening
on behalf of offspring in conflicts and providing a zone of safety that protects daughters and their
mothers from negative interactions with other group members, including threats from potentially
infanticidal males [78-80]. These forms of protection may reduce injury risk and improve offspring
and maternal health and survival [7, 61]. If strong early-life paternal relationships promote social
development [43], then fathers' presence may promote the development of adult female social
relationships. Grooming with fathers may also be one way that females learn to identify their
paternal sisters, who can themselves be important sources of social support in adulthood [81, 82].
However, early-life relationships with fathers do not necessarily have direct, causal effects on
offspring health and survival. Another possibility is that the strength of father-offspring
relationships reflects daughters' own phenotypic quality, which in turn, partly or completely explains
the relationship between these traits and female survival. Under this scenario, strong, healthy
juvenile females may build and maintain strong relationships with their fathers in early life, and also
lead long, healthy adult lives. In support, father-daughter grooming relationships are largely
maintained by daughters, not their fathers. Paternal effects may also be mediated by male health if
paternal health affects offspring health, even in the absence of a direct relationship between fathers
and offspring (through, for instance, epigenetic marks, semen quality, or other processes [19]). For
instance, in commercial pig farms, fathers kept in enriched conditions produced more live-born and
weaned piglets than those kept in pens—even though fathers had no contact with their offspring
[83]. Under this scenario, if fathers who are healthy and in good physical condition produce healthy
offspring, are less likely to disperse from groups in which they fathered offspring, and are more
socially engaged, including with their daughters, then these effects could create a correlation between
father-offspring social relationships and offspring health and survival that is not directly causal. In
support, the females in our data sometimes formed strong grooming relationships with males who
were not their fathers, but these relationships did not predict adult female survival.
508
(b) Do male baboons experience selection to provide parental care?

If male care behaviors have direct effects on their daughters' lifespans and fitness, then our

results help illuminate the selective pressures male baboons might experience with respect to

parenting effort. In Amboseli, adult female longevity is the largest contributor to female fitness for

baboons, explaining ~80-90% of individual variation in adult female lifetime reproductive success

[7, 45, 46]. Because female baboons typically produce one offspring every two years [84], the

increased years of survival predicted by paternal co-residency or grooming could translate to one to

two additional offspring over these females' lifespans. These results suggest that fathers and their

juvenile daughters experience selection to stay close and socially engage with each other.

Paternal care in baboons also lends context to the evolution of male parental behavior in

other primates, including humans [24]. Mating systems in early hominins are unknown, but the

presence of paternal care in modern humans has led many researchers to suggest that early hominins

lived in pair-bonded systems where males had limited alternative mating opportunities and high

paternity certainty [85-89]. However, our results, together with those from Assamese macaques and

mountain gorillas, which also provide evidence for subtle forms of paternal care in non-pair-bonded

systems [29, 90-92]—raise the possibility that male parental care could have emerged in early

hominins in the absence of strong pair-bonding [24].

While pair-bonds may not be a pre-requisite for male care in primates, male baboons are

more likely to invest in social bonds with offspring when the reproductive tradeoffs are favorable

[93-95]. In support, the males in our population were more likely to groom their daughters when

they had cues of paternity certainty (e.g., they consorted her mother near the time of her conception)

and few current mating opportunities (e.g., there were few fertile females in the group at the time of

her conception). Favorable reproductive tradeoffs may also explain why male "primary associates"

are more likely to be older than younger males, if the benefits males derive from mating effort may

decline as males age and their competitive abilities decline [34].

These observations, together with the survival patterns we observed, suggest that baboon

mothers may experience selection to increase males' paternity certainty. Doing so would be

advantageous to females if a prospective male sire meets other criteria for being a caring male (e.g.,

he sired a cohort of paternal siblings in the group). Selection for paternity certainty may also

contribute to the evolution of sexual swellings in female baboons, which provide obvious and

reliable signals of female ovulation and conception [35, 36, 65, 96, 97].

(c) How do early-life paternal relationships interact with other early-life conditions?

Paternal relationships did not seem to moderate the effects of early-life adversity on adult female survival. If early-life adversity and paternal relationships during the juvenile period have independent effects on adult female survival, then weak paternal relationships neither exacerbate the effects of other forms of early-life adversity, nor do paternal relationships have strong mitigating effects. However, the fact that they are independent means that, if a high adversity female has a strong early-life paternal relationship, she may be buffered against the effects of early-life adversity on adult survival. These results parallel those for the effects of adult social connectedness and fecal glucocorticoid hormones in our population, which also have effects on adult survival that are strong and appear to be mostly independent of early-life adversity [51, 77]. Hence, our data support the existence of multiple paths for mitigating early-life effects, including paternal relationships.

ACKNOWLEDGEMENTS

We are grateful to Jeanne Altmann for stewarding the ABRP and in designing many of its data collection protocols. We also acknowledge support from the National Science Foundation and the National Institutes of Health, currently through R01AG071684, R01AG075914, R01AG053330, R01AG053308, and R61AG078470. We thank the Max Planck Institute for Evolutionary Anthropology, Duke University, and the University of Notre Dame for financial and logistical support. In Kenya, our research was approved by the Wildlife Research Training Institute (WRTI), Kenya Wildlife Service (KWS), the National Commission for Science, Technology, and Innovation (NACOSTI), and the National Environment Management Authority (NEMA). We also thank the University of Nairobi, the Kenya Institute of Primate Research (KIPRE), the National Museums of Kenya, the members of the Amboseli-Longido pastoralist communities, the Enduimet Wildlife Management Area, and Ker & Downey Safaris for their cooperation and assistance in the field. We thank T. Wango and V. Oudu for their assistance in Nairobi. The ABRP database, is managed by J. Gordon and W. Wilbur, with contributions from N. Learn and K. Pinc. This research was approved by the IACUC at Duke University, University of Notre Dame, Princeton University, and the Ethics Council of the Max Planck Society. For further acknowledgments please visit http://amboselibaboons.nd.edu/acknowledgements/.

DATA STATEMENT

Our data are publicly available on Zenodo at https://doi.org/10.5281/zenodo.14590285. Our code

are available on GitHub at https://github.com/david-awam-

jansen/BaboonPaternalRelationshipsSurvival

REFERENCES

[1] Snyder-Mackler, N., Burger, J.R., et al. 2020 Social determinants of health and survival in humans
and other animals. *Science* **368**, 6493.
[2] Curley, J.P. & Champagne, F.A. 2023 Shaping the development of complex social behavior. *Ann*
N Y Acad Sci **1530**, 46-63.
[3] Goldenberg, S.Z. & Wittemyer, G. 2018 Orphaning and natal group dispersal are associated with
social costs in female elephants. *Animal Behaviour* **143**, 1-8.
[4] Goldenberg, S.Z. & Wittemyer, G. 2017 Orphaned female elephant social bonds reflect lack of
access to mature adults. *Scientific Reports* **7**.
[5] Kalcher-Sommersguter, E., Preuschoft, S., et al. 2015 Early maternal loss affects social
integration of chimpanzees throughout their lifetime. *Scientific Reports* **5**.
[6] Rosenbaum, S.R., Zeng, S., et al. 2020 Social bonds do not mediate the relationship between
early adversity and adult glucocorticoids in wild baboons. *Proceedings of the National Academy of Sciences*
117, 20052-20062.
[7] Tung, J., Archie, E.A., et al. 2016 Cumulative early adversity predicts longevity in wild baboons.
Nature Communications **7**, 11181.
[8] Stanton, M.A., Lonsdorf, E.V., et al. 2020 Consequences of maternal loss before and after
weaning in male and female wild chimpanzees. *Behavioral Ecology and Sociobiology*.
[9] Parker, J.M., Webb, C.T., et al. 2021 Poaching of African elephants indirectly decreases
population growth through lowered orphan survival. *Current Biology* **31**, 4156-+.
[10] Zipple, M.N., Altmann, J., et al. 2021 Maternal death and offspring fitness in multiple wild
primates. *Proc Natl Acad Sci U S A* **118**.
[11] Sear, R. & Mace, R. 2008 Who keeps children alive? A review of the effects of kin on child
survival. *Evolution and Human Behavior* **29**, 1-18.
[12] Lahdenpera, M., Mar, K.U., et al. 2016 Short-term and delayed effects of mother death on calf
mortality in Asian elephants. *Behav Ecol* **27**, 166-174.
[13] Patterson, S.K., Strum, S.C., et al. 2022 Early life adversity has long-term effects on sociality and
interaction style in female baboons. *Proc Biol Sci* **289**, 20212244.
[14] Kleiman, D. & Malcolm, J. 1981 The evolution of male parental investment in mammals. In
Parental Care in Mammals (eds. D. Gubernick & P. Klopfer), pp. 347–387. New York, Plenum Press.
[15] Clutton-Brock, T.H. 1991 *The evolution of parental care*. Princeton, Princeton University Press.
[16] Smith, K.R., Hanson, H.A., et al. 2014 Survival of offspring who experience early parental
death: early life conditions and later-life mortality. *Soc Sci Med* **119**, 180-190.
[17] Feldman, R. 2023 Father contribution to human resilience. *Development and Psychopathology* **35**,
2402-2419.
[18] Gettler, L.T., Boyette, A.H., et al. 2020 Broadening Perspectives on the Evolution of Human
Paternal Care and Fathers' Effects on Children. *Annu Rev Anthropol*.
[19] Watkins, A.J., Rubini, E., et al. 2020 Paternal programming of offspring health. *Early Hum Dev*
150, 105185.
[20] Todd, N., Valleron, A.J., et al. 2017 Prenatal loss of father during World War One is predictive
of a reduced lifespan in adulthood. *Proc Natl Acad Sci U S A* **114**, 4201-4206.

[21] Braun, K. & Champagne, F.A. 2014 Paternal influences on offspring development: behavioural
and epigenetic pathways. *J Neuroendocrinol* **26**, 697-706.
[22] Dimofski, P., Meyre, D., et al. 2021 Consequences of Paternal Nutrition on Offspring Health
and Disease. *Nutrients* **13**.
[23] Clutton-Brock, T. 2016 *Mammal Societies*, Wiley Blackwell.
[24] Rosenbaum, S. & Silk, J.B. 2022 Pathways to paternal care in primates. *Evol Anthropol* **31**, 245-
262.
[25] Sandel, A.A., Langergraber, K.E., et al. 2020 Adolescent male chimpanzees (*Pan troglodytes*) form
social bonds with their brothers and others during the transition to adulthood. *American Journal of*
Primatology **82**.
[26] Rosenbaum, S., Silk, J.B., et al. 2011 Male-immature relationships in multi-male groups of
mountain gorillas (Gorilla beringei beringei). *Am J Primatol* **73**, 356-365.
[27] de Bruin, R., Ganswindt, A., et al. 2016 From killer to carer: steroid hormones and paternal
behaviour. *Afr Zool* **51**, 173-182.
[28] Maestripieri, D. 1998 The evolution of male-infant interactions in the tribe Papionini (Primates:
Cercopithecidae). *Folia Primatol (Basel)* **69**, 247-251.
[29] Ostner, J., Vigilant, L., et al. 2013 Stable heterosexual associations in a promiscuous primate.
Animal Behaviour **86**, 623-631.
[30] Rosenbaum, S., Vigilant, L., et al. 2018 Caring for infants is associated with increased
reproductive success for male mountain gorillas. *Scientific Reports* **8**, 15223.
[31] Charpentier, M.J., Van Horn, R.C., et al. 2008 Paternal effects on offspring fitness in a
multimale primate society. *Proc Natl Acad Sci U S A* **105**, 1988-1992.
[32] Buchan, J.C., Alberts, S.C., et al. 2003 True paternal care in a multi-male primate society. *Nature*
425, 179-181.
[33] Nguyen, N., Van Horn, R.C., et al. 2009 "Friendships" between new mothers and adult males:
adaptive benefits and determinants in wild baboons (Papio cynocephalus). *Behavioral Ecology and*
Sociobiology **63**, 1331-1344.
[34] Silk, J.B., Stadele, V., et al. 2020 Shifts in Male Reproductive Tactics over the Life Course in a
Polygynandrous Mammal. *Curr Biol* **30**, 1716-1720 e1713.
[35] Stadele, V., Roberts, E.R., et al. 2019 Male-female relationships in olive baboons (Papio anubis):
Parenting or mating effort? *J Hum Evol* **127**, 81-92.
[36] Stadele, V., Vigilant, L., et al. 2021 Extended male–female bonds and potential for prolonged
paternal investment in a polygynandrous primate (Papio anubis). *Animal Behaviour*.
[37] Huchard, E., Alvergne, A., et al. 2010 More than friends? Behavioural and genetic aspects of
heterosexual associations in wild chacma baboons. *Behavioral Ecology and Sociobiology* **64**, 769-781.
[38] Moscovice, L.R., Di Fiore, A., et al. 2010 Hedging their bets? Male and female chacma baboons
form friendships based on likelihood of paternity. *Animal Behaviour* **79**, 1007-1015.
[39] Smuts, B.B. 1985 *Sex and Friendship in Baboons*. New York, Aldine Transaction.
[40] Palombit, R.A., Seyfarth, R.M., et al. 1997 The adaptive value of 'friendships' to female
baboons: experimental and observational evidence. *Animal Behavior* **54**, 599-614.
[41] Lemasson, A., Palombit, R.A., et al. 2008 Friendships between males and lactating females in a
free-ranging group of olive baboons (Papio hamadryas anubis): evidence from playback experiments.
Behavioral Ecology and Sociobiology **62**, 1027-1035.
[42] Moscovice, L.R., Heesen, M., et al. 2009 Paternity alone does not predict long-term investment
in juveniles by male baboons. *Behav Ecol Sociobiol* **63**, 1471-1482.
[43] Zipple, M.N., Southworth, C.A., et al. 2024 Infant spatial relationships with adult males in a
wild primate: males as mitigators or magnifiers of intergenerational effects of early adversity? *bioRxiv*.

[44] Lynch, E.C., Di Fiore, A., et al. 2017 Fathers enhance social bonds among paternal half-siblings
in immature olive baboons (Papio cynocephalus). *Behavioral Ecology and Sociobiology* **71**, 120.
[45] Weibel, C.J., Tung, J.T., et al. 2020 Accelerated reproduction is not an adaptive response to early
life adversity in wild baboons. *Proceedings of the National Academy of Sciences* **117**, 24909-24919.
[46] McLean, E.M., Archie, E.A., et al. 2019 Lifetime Fitness in Wild Female Baboons: Trade-Offs
and Individual Heterogeneity in Quality. *American Naturalist* **194**, 745-759.
[47] Alberts, S.C. & Altmann, J. 2012 The Amboseli Baboon Research Project: Themes of continuity
and change. In *Long-term field studies of primates* (eds. P. Kappeler & D.P. Watts), pp. 261-287, Springer
Verlag.
[48] Silk, J.B., Altmann, J., et al. 2006 Social relationships among adult female baboons (*Papio
cynocephalus*) I. Variation in the strength of social bonds. *Behavioral Ecology and Sociobiology* **61**, 183-195.
[49] Mitani, J.C. 2009 Male chimpanzees form enduring and equitable social bonds. *Animal Behaviour*
77, 633-640.
[50] Schino, G. & Aureli, F. 2008 Grooming reciprocation among female primates: a meta-analysis.
Biol Letters **4**, 9-11.
[51] Lange, E.C., Zeng, S., et al. 2023 Early life adversity and adult social relationships have
independent effects on survival in a wild primate. *Sci Adv* **9**, eade7172.
[52] Wall, J.D., Schlebusch, S.A., et al. 2016 Genome-wide ancestry and divergence patterns from
low-coverage sequencing data reveal a complex history of admixture in wild baboons. *Mol Ecol* **25**,
3469-3483.
[53] Vilgalys, T.P., Fogel, A.S., et al. 2022 Selection against admixture and gene regulatory divergence
in a long-term primate field study. *Science* **377**, 635-641.
[54] Onyango, P.O., Gesquiere, L.R., et al. 2013 Puberty and dispersal in a wild primate population.
Hormones and Behavior **64**, 240-249.
[55] McLean, E.M., Moorad, J.A., et al. 2023 Genetic variance and indirect genetic effects for
affiliative social behavior in a wild primate. *Evolution* **77**, 1607-1621.
[56] Alberts, S.C., Buchan, J.C., et al. 2006 Sexual selection in wild baboons: from mating
opportunities to paternity success. *Anim Behav* **72**, 1177-1196.
[57] Van Horn, R.C., Buchan, J.C., et al. 2007 Divided destinies: group choice of female savannah
baboons during social group fission. *Behavioral Ecology and Sociobiology* **61**, 1823-1837.
[58] Kalinowski, S.T., Taper, M.L., et al. 2007 Revising how the computer program CERVUS
accommodates genotyping error increases success in paternity assignment. *Molecular Ecology* **16**, 1099-
1006.
[59] Marshall, T.C., Slate, J., et al. 1998 Statistical confidence for likelihood-based paternity inference
in natural populations. *Molecular Ecology* **7**, 639-655.
[60] Campos, F.A., Villavicencio, F., et al. 2020 Social bonds, social status and survival in wild
baboons: a tale of two sexes. *Philos Trans R Soc Lond B Biol Sci* **375**, 20190621.
[61] Zipple, M.N., Archie, E.A., et al. 2019 Intergenerational effects of early adversity on survival in
wild baboons. *eLife* **8**, e47433.
[62] Alberts, S.C., Archie, E.A., et al. 2024 Monitoring guide for the Amboseli Baboon Research
Project: http://amboselibaboons.nd.edu/downloads/.
[63] Alberts, S.C., Watts, H.E., et al. 2003 Queuing and queue-jumping: long-term patterns of
reproductive skew in male savannah baboons, *Papio cynocephalus. Animal Behaviour* **65**, 821-840.
[64] Gordon, J.B., Jansen, D., et al. 2023 Ordinal dominance rank assignments: Protocol for the
Amboseli Baboon Research Project. (
[65] Gesquiere, L.R., Wango, E.O., et al. 2007 Mechanisms of sexual selection: sexual swellings and
estrogen concentrations as fertility indicators and cues for male consort decisions in wild baboons.
Hormones and Behavior **51**, 114-125.

[66] Beehner, J.C., Onderdonk, D.A., et al. 2006 The ecology of conception and pregnancy failure in
wild baboons. *Behavioral Ecology* **17**, 741-750.
[67] Altmann, J., Alberts, S.C., et al. 1996 Behavior predicts genetic structure in a wild primate
group. *Proceedings of the National Academy of Sciences* **93**, 5797-5801.
[68] Bates, D., Maechler, M., et al. 2014 lme4: Linear mixed-effects models using Eigen and S4. R
package version 1.1-5. In *http://CRAN.R-project.org/package=lme4* (
[69] Kuznetsova, A., Brockhoff, P.B., et al. 2017 lmerTest Package: Tests in Linear Mixed Effects
Models. *Journal of Statistical Software* **82**, 1-26.
[70] Zeileis, A. & Hothorn, T. 2002 Diagnostic Checking in Regression Relationships. *R News* **2**, 7–
10.
[71] Barton, K. 2023 R Package: MuMIn: Multi-model inference (1.47.5). (
[72] Stoffel, M.A., S., N., et al. 2017 rptR: Repeatability estimation and variance decomposition by
generalized linear mixed-effects models. *Methods in Ecology and Evolution* **8**, 1639-1644.
[73] Therneau. 2014 A Package for Survival Analysis in S. R package version 2.37-7,
http://CRAN.R-project.org/package=survival. (
[74] Archie, E.A., Tung, J., et al. 2014 Social affiliation matters: both same-sex and opposite-sex
relationships predict survival in wild female baboons. *Proceedings of the Royal Society* **281**, 20141261.
[75] Belsley, D.A., Kuh, E., et al. 1980 *Regression Diagnostics; Identifying Influence Data and Source of
Collinearity*. New York, Wiley.
[76] Silk, J.B., Beehner, J.C., et al. 2010 Strong and consistent social bonds enhance the longevity of
female baboons. *Current Biology* **20**, 1359-1361.
[77] Tung, J., Lange, E.C., et al. 2023 Social and early life determinants of survival from cradle to
grave: A case study in wild baboons. *Neurosci Biobehav Rev* **152**, 105282.
[78] Palombit, R.A. 2003 Male infanticide in wild savanna baboons: adap- tive significance and
intraspecific variation. In *Sexual selection and reproductive competition in primates: new perspectives and
directions* (ed. C.B. Jones), pp. 367-411, American Society of Primatologists.
[79] Collins, D.A., Busse, C.D., et al. 1984 Infanticide in two populations of savanna baboons. In
Infanticide: comparative and evolutionary perspectives (eds. G. Hausfater & S. Hrdy), pp. 193–216. New
York, Hawthorne.
[80] Zipple, M.N., Grady, J.H., et al. 2017 Conditional fetal and infant killing by male baboons. *Proc
Biol Sci* **284**.
[81] Smith, K., Alberts, S.C., et al. 2003 Wild female baboons bias their social behaviour towards
paternal half-sisters. *Proceedings of the Royal Society of London* **270**, 503-510.
[82] Alberts, S.C. 1999 Paternal kin discrimination in wild baboons. *Proceedings of the Royal Society of
London* **266**, 1501-1506.
[83] Sabei, L., Bernardino, T., et al. 2023 Life experiences of boars can shape the survival,
aggression, and nociception responses of their offspring. *Front Anim Sci* **4**.
[84] Gesquiere, L.R., Altmann, J., et al. 2018 Interbirth intervals in wild baboons: Environmental
predictors and hormonal correlates. *Am J Phys Anthropol* **166**, 107-126.
[85] Chapais, B. 2009 *Primeval kinship: how pair-bonding gave birth to human society*. Cambridge, MA,
Harvard University Press.
[86] Chapais, B. 2013 Monogamy, strongly bonded groups, and the evolution of human social
structure. *Evol Anthropol* **22**, 52-65.
[87] Rodseth, L., Wrangham, R.W., et al. 1991 The Human Community as a Primate Society [and
commentary]. *Curr Anthropol* **32**, 221-254.
[88] Swedell, L. & Plummer, T. 2019 Social evolution in Plio-Pleistocene hominins: Insights from
hamadryas baboons and paleoecology. *Journal of Human Evolution* **137**.
[89] Bales, K.L., Ardekani, C.S., et al. 2021 What is a pair bond? *Hormones and Behavior* **136**.

[90] Maestripieri, D. 1998 The evolution of male-infant interactions in the tribe Papionini (Primates: Cercopithecidae). *Folia Primatol* **69**, 247-251.

[91] Rosenbaum, S., Silk, J.B., et al. 2011 Male-Immature Relationships in Multi-Male Groups of Mountain Gorillas ( ). *American Journal of Primatology* **73**, 356-365.

[92] Rosenbaum, S., Vigilant, L., et al. 2018 Caring for infants is associated with increased reproductive success for male mountain gorillas. *Scientific Reports* **8**.

[93] Trivers, R.L. 1972 Parental investment and sexual selection. In *Sexual Selection and the Descent of Man 1871-1971* (Aldine Press.

[94] Stiver, K.A. & Alonzo, S.H. 2009 Parental and Mating Effort: Is There Necessarily a Trade-Off. *Ethology* **115**, 1101–1126.

[95] Kokko, H. & Jennions, M.D. 2008 Parental investment, sexual selection and sex ratios. *J Evol Biol* **21**, 919-948.

[96] Alberts, S.C. & Fitzpatrick, C.L. 2012 Paternal care and the evolution of exaggerated sexual swellings in primates. *Behavioral Ecology*.

[97] Nunn, C.L. 1999 The evolution of exaggerated sexual swellings in primates and the graded-signal hypothesis. *Anim Behav* **58**, 229-246.