Dissecting the molecular basis underlying mycobacterial cell-wall hydrolysis by the catalytic domains of *D29*LysA and *DS6A*LysA phage endolysins

Fernando Ceballos-Zúñiga[1], Laura Gálvez-Larrosa[1,#], Inés G. Muñoz[2], Lourdes[1] Infantes, Inmaculada Pérez-Dorado[1,*].

[1] Department of Crystallography and Structural Biology, Institute of Physical Chemistry Blas Cabrera, Spanish Research Council, Serrano 119, 28006 Madrid, Spain

[2] Protein Crystallography Unit. Structural Biology Programme. Spanish National Cancer Research Centre (CNIO). Melchor Fernández Almagro 3, 28009 Madrid, Spain.

[#] Center for Cooperative Research in Biosciences CIC bioGUNE, Basque Research and Technology Alliance (BRTA), Bizkaia Technology Park, Building 800, 48162, Derio, Bizkaia, Spain.

*Correspondence e-mail: jiperez@iqf.csic.es

These authors contributed equally: Fernando Ceballos-Zúñiga and Laura Gálvez-Larrosa

Abstract: Mycobacteria embrace a broad pool of microorganisms causing infections with renowned impact in human health causing altogether millions of deaths every year. From tuberculosis and lepra, caused by *Mycobacterium tuberculosis* and *Mycobacterium leprae* respectively, to infections caused by emergent/opportunistic pathogens such as *Mycobacterium abscessus*. The battle to confront this health burden is further challenged by limitations in the treatments currently available and the emergence of antimicrobial resistance, thus making necessary the search for new therapeutic strategies to fight these infections. Mycobacteriophage LysA endolysins are complex multi-domain peptidoglycan hydrolyses with reported antimicrobial relevance and potential to treat mycobacterial infections. However, and despite the therapeutic prospective of LysAs, a profound understanding of their mechanism of action still remains limited. The present work provides a thorough structural-functional study of the catalytic domains of two LysA endolysins encoded by the bacteriophages *D29* and *DS6A,* known to infect pathogenic mycobacteria including *M. tuberculosis*. As part of this study, we have characterised the four catalytic domains bared by both endolysins (*D29*N4/*D29*GH19 and *DS6A*GH19/*DS6A*Ami2B) alone and in complex with PG analogues. To do so, our approach was to combine protein engineering, X-ray crystallography, small-angle X-ray scattering and *in silico* tools that, to our knowledge, it has yielded the first experimental structures reported for mycobacteriophage endolysins. Our study unravels major aspects of peptidoglycan-binding and hydrolysis by *D29*LysA and *DS6A*LysA lysins and other homologue LysAs, including the hydrolase domains alike to the ones here examined. Altogether, this constitutes an important step forward for understanding how the mycobacterial cell-wall hydrolysis is done by this relevant class of endolysins and, opening the path for their future use in their therapeutic exploration as enzybiotics, allowing the rational design of *a la carte* enzymes with optimised lytic properties against mycobacterial pathogens.

Keywords: phage endolysins, mycobacterial infections, host-pathogen interactions, enzybiotics, antimicrobial resistance

1. Introduction

The *Mycobacterium* genus contains over 170 recognized species, many of which produce infectious diseases (IDs) in human such as Buruli ulcer, leprosy or tuberculosis (TB) (Tuberculosis; Buruli ulcer; Leprosy), caused by *Mycobacterium ulcerans* (*Mlc*), *Mycobacterium leprae*, and *Mycobacterium tuberculosis* (*Mtb*), respectively (Batt *et al*, 2020). Among them, human TB is the world's most devastating human ID, being the first cause of death in the world produced by bacteria according to the World Health Organization (Global Tuberculosis Report 2023, 2023). Buruli ulcer and leprosy are IDs associated with permanent deformation, disability and stigma, causing >200.000 new cases every year across >120 countries in the world (Global Leprosy (Hansen's disease) Strategy 2021-2030 | 2, 2021). In addition, various so-called non-tuberculous mycobacteria can cause IDs in immune-compromised individuals, such as *Mycobacterium abscessus* (*Mbs*), which constitutes an emergent pathogen (Johansen *et al*, 2020). Great part of the success of mycobacteria as pathogens relies on their mycobacterial cell envelop (MCE), which consists of a plasma membrane surrounded by complex structure formed by three distinct covalently linked macromolecules: the outer membrane (OM) rich in mycolic acids (MA), arabinogalactan (AG), and peptidoglycan (PG) (**Fig. 1**) (Catalão *et al*, 2019). Altogether this results in a very impermeable barrier essential for cell integrity that protects the bacilli from harsh conditions, being one of the main reasons for the antimicrobial resistance in mycobacteria (Catalão *et al*, 2019; Batt *et al*, 2020).

Among MCE components, PG provides the mechanical strength to resist the turgor pressure, at the time it plays a central role in regulating cellular shape, growth, and division (Batt *et al*, 2020; Vollmer *et al*, 2008). The PG molecule is built by linear saccharide strands of alternating units of *N*-acetyl glucosamine (NAG) and *N*-acetyl muramic acid (NAM) connected by ß-1,4 bonds (**Fig. 1**). These glycan strands are in turn covalently cross-linked via peptide stems formed by L-alanine (L-Ala), D-*iso*-glutamate (D-*iso*-Glu), *meso*-diaminopimelate (*m*DAP) acid, and D-Alanine (D-Ala), present at NAM residues. Mycobacterial species exhibit characteristic modifications in both glycan and peptide moieties, such as amidation of free carboxylic groups or glycolylation of NAM (*N*-glycolylmuramic acid or MurGlyc) (Catalão *et al*, 2019; Batt *et al*, 2020). While the former is important for the PG biosynthesis, glycolylation may confer with an additional strength of the PG molecule as well as resistance to lysozyme, thus protecting mycobacteria against the innate immune system (Raymond *et al*, 2005). Moreover, mycobacteria exhibit a higher degree of peptide cross-linking (up to 80%) compared to other bacteria, which results on a sturdier PG structure (Matsuhashi M., 1966; Vollmer & Höltje, 2004). PG constitutes an essential structural constituent required for cell integrity and its disruption

Figure 1. (A) Scheme representing the mycobacterial cell-wall architecture consisting of the inner membrane (IM) surrounded by the mycolyl-arabinogalactan-peptidoglycan (MA-AG-PG) complex (left). Mycobacteriophage LysA and LysB lysins targeting the PG and MA moieties are represented as blue and red packmen, respectively. Details of the peptidoglycan composition are shown (right), which consists of glycan strands formed by alternating N-acetylglucosamine (NAG) and N-acetylmuramic acid (NAM) and covalently cross-linked via peptide stems pending from NAM residues. The PG-hydrolase activities predicted in LysA endolysins are depicted as scissors. (B) Representation of the three-domain architecture predicted for *D29*LysA and *DS6A*LysA endolysins.

leads to bactericidal cell lysis (Vollmer *et al*, 2008), thus resulting in a major target for antimicrobials (Alderwick *et al*, 2015). However, the unique structure of the mycobacterial PG and the alarming emergence of resistance strategies make many of these antimicrobials ineffective in mycobacteria (Catalão *et al*, 2019; Batt *et al*, 2020). For *Mtb*, this pathogen is considered naturally resistant to most β-lactam antibiotics, which results of a highly active β-lactamase (BlaC) able to neutralise many β-lactams, and the presence of non-classical L,D-transpeptidases, inherently resistant to these antibiotics (Batt *et al*, 2020; Catalão *et al*, 2019). TB and other mycobacterial IDs still represent a major health burden worldwide, which is aggravated by an increasing emergence of multi-resistant strains, as well as high toxicity and long-term treatments associated with drugs currently in use (Catalão *et al*, 2019). This urgent the search for new bacteriostatics in a moment marked by an alarming shortage of new antibacterial classes. In this context, bacteriophage endolysins emerge as a novel and promising class of specific antimicrobials (enzybiotics) to treat bacterial IDs, including those caused by mycobacteria (Dams & Briers, 2019). Bacteriophage-encoded endolysins are PG-hydrolases targeting the bacterial cell-envelop, enabling host lysis and release of the viral progeny at the end of the lytic cycle (Dams & Briers, 2019). Phage endolysins are highly specific, and their exogenously application has been shown to yield a rapid and specific elimination of pathogenic bacteria *in vitro* and in animal models, of infection caused by different pathogens such as *Streptococcus sp.* (Nelson *et al*, 2001; Loeffler *et al*, 2003), *Bacillus anthracis* (Schuch *et al*, 2002) and *Staphylococcus aureus* (Haddad Kashani *et al*, 2018). Moreover, and given the essentiality of the PG molecule, endolysins exhibit poor prospects to develop resistance, further supporting enzybiotics as an attractive alternative to antibiotics (Dams & Briers, 2019).

Phages infecting mycobacteria (mycobacteriophages) produce an endolysin protein referred to as Lysin A (LysA), whose lytic action is assisted by a second enzyme with esterase activity, referred to as Lysin B (LysB), that cleaves the bonds connecting MAs and AG molecules (Catalão *et al*, 2019; Dams & Briers, 2019). The bactericidal activity of mycobacteriophage endolysins has been reported for LysA from *PDRPxv* mycobacteriophage, capable to lysate the non-pathogenic *Mycobacterium smegmatis*

(*Msm*), a lytic effect that was exacerbated by its combination with *PDRPxv*LysB (Eniyan *et al*, 2020). More recently, an enzyme cocktail combining LysA and LysB lysins with capsule degrading enzymes has exhibited bactericide effect on several pathogenic species of mycobacteria including *Mtb* (Bartlett *et al*, 2024). This finding reinforces the potential of mycobacteriophage LysA endolysins as antimicrobials, alone or in combination with other lysins and drugs (Eniyan *et al*, 2020; Bartlett *et al*, 2024). Thus, a profound understanding of the mechanisms underlying PG binding and hydrolysis by LysA endolysins at a molecular level is key for their rational development and application as effective enzybiotics. To this end, bioinformatic studies of LysA endolysins reveal them as complex as multi-domain enzymes. Predominantly, they exhibit a three-domain arrangement formed by N-terminal and central catalytic domains with putative peptidase, glycoside-hydrolase or/and amidase activities, followed by a C-terminal region reported to function as a PG-binding domain in *D29*LysA (Gangakhedkar & Jain, 2024; Payne & Hatfull, 2012). This complex architecture aligns with the intricate structure of the mycobacterial PG, complexity that has likely hampered their structural and functional analysis so far, proven by the scarce experimental structural information available. With, to our knowledge, only computational models reported for LysA from the bacteriophages *TM4* (Urdániz *et al*, 2022), *Che12* (Saadhali *et al*, 2016), *D29*LysA (Joshi *et al*, 2017, 2019; Gangakhedkar & Jain, 2024).

Among known mycobacteriophages, *D29* and *DS6A* are able to infect pathogenic species of mycobacteria including *Mtb* (W.B. Redmond & J. C. Carter, 1960; Yang *et al*, 2024; Froman *et al*, 1954). Based on this information, LysA endolysins encoded by *D29* and *DS6A* phages are expected to efficiently cleave the PG of *Mtb* and other pathogenic mycobacteria, hence emerging as relevant candidates for enzybiotic development targeting their infections. Both *DS6A*LysA (residues 1-486) and *D29*LysA (residues 1-493) exhibit a three-domain architecture (**Fig. 1**) that, in the case of *D29*LysA, consists of an N-terminal N4 domain with proposed cysteine protease activity, followed by a glycosyl hydrolase domain belonging to the family 19 (referred as GH19 domain), and a C1 domain reported to interact with the PG molecule (Payne & Hatfull, 2012; Gangakhedkar & Jain, 2024). *DS6A*LysA is comprised by an N-terminal GH19 domain followed by an

amidase domain predicted to be a member of Amidase_2 family (referred to as Ami2B domain), and a C2 domain suggested to function as a PG-binding region (Payne & Hatfull, 2012). Biochemical studies conducted in *D29*LysA endolysin show that *D29*N4 and *D29*GH19 are indeed domains exhibiting PG-hydrolase, where *D29*N4 is required for optimal bacterial lysis domain (Joshi *et al*, 2017; 2022; Gangakhedkar & Jain, 2024). Moreover analysis of *D29*N4, combining computational modelling with a thorough biochemical characterisation, has recently evidenced the proposed peptidase function predicted for this domain (Gangakhedkar & Jain, 2024). However, the lack of experimental structures to date have prevented the validation of *in silico* models available for *D29*N4, and other homologue GH19 and Ami2B domains (Joshi *et al*, 2017; Saadhali *et al*, 2016; Urdániz *et al*, 2022; Nair & Jain, 2024; Gangakhedkar & Jain, 2024); as well as acquiring details of their mechanism of action at a molecular level. Information that is most relevant for future investigations exploring the potential of *DS6A*LysA and *D29*LysA as enzybiotics.

In this work we present an extensive structural and functional study aiming at the characterisation of the catalytic domains of *D29*LysA and *DS6A*LysA endolysins by using X-ray crystallography, small-angle X-ray scattering and differential scanning fluorimetry in combination with site-directed mutagenesis and molecular docking approaches. To our knowledge, our results constitute the first experimental structures reported for mycobacteriophage endolysins, which unveil key aspects of their mode of interaction with the PG and hydrolysis. This work constitutes a solid groundwork for the future therapeutic application of *D29*LysA and *DS6A*LysA, including the rational design of optimised enzybiotics to fight mycobacterial infections.

2. Material and methods

2.1. Protein production
Protein construct bearing the catalytic domains of *D29*LysA (UniProt entry code O64203) and *DS6A*LysA (UniProt entry code G8I4E0) endolysins were designed for their recombinant production. These include each individual wild-type domain *D29*N4 (residues 1-174), *D29*GH19 (residues 179-370), *DS6A*GH19 (residues 1-196), and *DS6A*Ami2B (residues 206-395), the catalytically inactive mutant *D29*GH19[E228Q] (designed based on available activity studies of GH19 family (Hoell *et al*, 2006), and a construct bearing both GH19 and Ami2B domains of *DS6A*LysA, (*DS6A*GH19-Ami2B, residues 1-395). Constructs were designed as N-terminal His-tagged proteins followed by a 3C-protease site preceding *D29*LysA and *DS6A*LysA catalytic domains. The DNA coding for the constructs cloned into a pET-28a plasmid (between NcoI and XhoI restriction sites) was purchased to GenScript with codon optimization for *E. coli* expression. Proteins were produced as recombinant proteins in *E. coli* BL21 (DE3) Star (Invitrogen) using 2xTY media supplemented with kanamycin (50 μg/ml). Cells were grown at 37 °C until an $OD_{600\,nm}$ of 0.8 and then cooled down to 16 °C for an hour. Expression was then induced at 16 °C by the addition of 1 mM IPTG for 20 h. Cells were harvested by centrifugation at 5251 *g* for 20 min at 4 °C, and cell pellets were stored at -20 °C. Cell pellets were thawed and resuspended in 20 mM Tris-HCl, 500 mM NaCl, 10 mM imidazole (buffer A), supplemented with 1 mM $MgCl_2$, 1 mM $MnCl_2$ and 1 μg/ml DNAse (Sigma). This step was performed at RT while subsequent purification steps were

carried out at 4 °C. Cells were disrupted by sonication and the clarified extract was loaded into a 1 ml HisTrap HP crude column (Cytiva) previously equilibrated in buffer A. Protein samples were eluted using an imidazole gradient (from 10 mM to 500 mM) in buffer A, and then buffer exchanged to 20 mM Tris-HCl, 150 mM NaCl (Buffer B) using a PD-10 desalting column (Cytiva). When required, samples were diluted to 1 mg/ml in buffer B prior to overnight cleavage with the 3C protease. Cleaved and not cleaved proteins were concentrated using Amicon 10 and 30 KDa MWCO concentrators (Millipore), and further purified by size exclusion chromatography (SEC) using a Superdex 200 (16/60) or a Superdex 200 10/300 Increase GL column (Cytiva) preequilibrated in buffer B. Samples corresponding to *DS6A*GH19, *DS6A*Ami2B, *DS6A*GH19-Ami2B, *D29*N4 and *D29*GH19 (WT and mutant) were purified using buffers A and B at pH 8.0 supplemented with 4 mM βME (IMAC steps) and 1 mM DTT (SEC). Pure protein fractions obtained from SEC were pooled, concentrated, and flash-frozen in liquid nitrogen for their storage at -80 °C. Protein sample purity and quantification were assessed using PAGE-SDS and UV-Vis absorbance at 280 nm, respectively.

2.2. Crystallisation and structure determination
All crystallization experiments were carried out using the sitting-drop vapour diffusion method at 18ºC in 96-well MRC plates (Hampton Research) and employing an Oryx8 robot (Douglas Instruments). Crystals of *D29*N4, *D29*GH19[WT], *D29*GH19[E228Q], *DS6A*GH19, and *DS6A*Ami2B were initially grown in native conditions using the untagged proteins in all cases but for the *D29*N4 domain. *D29*N4 crystallised in 300 nl droplets formed by mixing 150 nl of protein at 11 mg/ml with 150 nl of precipitant solution (0.1 M HEPES pH 7.5, 1.8 M NaCl). *D29*GH19[WT] and *D29*GH19[E228Q] crystallised in up to four crystal forms (I-IV). Crystal form I grew in 200 nl droplets formed by mixing 100 nl of *D29*GH19[WT] at 8.25 mg/ml with 100 nl of precipitant solution (12% (v/v) ethylene glycol; 6% (w/v) PEG 8K, 0.1 M imidazole/MES buffer pH 6.5, and DL-amino-acids (glutamic acid, alanine, glycine; lysine, serine) at 20 mM); crystal form II grew in 200 nl droplets formed by mixing 100 nl of *D29*GH19[WT] at 8.25 mg/ml with 100 nl of precipitant solution (9% (v/v) MPD; 9% (v/v) PEG 1K; 9% (w/v) PEG 3350. 0.1 M imidazole/MES buffer pH 6.5, and DL-amino-acids (glutamic acid, alanine, glycine; lysine, serine) at 20 mM; crystal form III grew in 200 nl droplets formed by mixing 100 nl of *D29*GH19[WT] at 8.25 mg/ml with 100 nl of precipitant solution (9% (v/v) MPD, 9% (w/v) PEG 1K, 9% (w/v) PEG 3350, 0.1 M imidazole/MES buffer pH 6.1, and DL-amino-acids (glutamic acid, alanine, glycine; lysine, serine) at 20 mM; or 13% (v/v) MPD, 9% (v/v) PEG 1K, 9% (w/v) PEG 3350, 0.1 M imidazole/MES buffer pH 6.1, and DL-amino-acids (glutamic acid, alanine, glycine; lysine, serine) at 20 mM; crystal form IV grew in 200 nl droplets formed by mixing 100 nl of *D29*GH19[WT] at 8.25 mg/ml with 100 nl of precipitant solution (9% (v/v) MPD, 9% (w/v) PEG 1K, 9% (w/v) PEG 3350, 0.1 M Tris/BICINE pH 8.5, 30 mM $MgCl_2$, 30 mM $CaCl_2$); and crystal form V grew in 300 nl droplets formed by mixing 100 nl of *D29*GH19[E228Q] at 9 mg/ml with 200 nl of precipitant solution (20 % (v/v) PEG 500 MME, 10 % (w/v) PEG 20K, 0.1 M HEPES/MOPS buffer pH 7.5, and DL-amino-acids (glutamic acid, alanine, glycine; lysine, serine) at 20 mM. *DS6A*GH19[WT] crystallised in 300 nl droplets formed by mixing 150 nl of protein at 11 mg/ml with 100 nl of precipitant solution (2.2 M NaCl, Tris-HCl 50 mM pH 8.0), and 50 nl of a seed stock in stabilised in the precipitant solution with the butter at 100 mM. *DS6A*Ami2B[WT] crystallised in 300 nl droplets formed by mixing 150 nl of protein

at 16.5 mg/ml with 150 nl of precipitant solution (0.8 M NaH_2PO_4, 0.4 M KH_2PO_4 and 0.1 M HEPES at pH 7.5). Complexes of $D29$GH19^{WT}:NAG and $D29$GH19^{WT}:NAG$_2$:NAG were obtained by soaking native $D29$GH19^{WT} crystals into a precipitant solution supplemented with 31 mM NAG$_6$ for 30 min and 10 seconds, respectively. The $D29$GH19^{E228Q}:NAG$_4$ complex was obtained by *in situ* soaking, in which NAG$_4$ in powder was added to the crystallisation drop containing native $D29$GH19^{E228Q} crystals that were incubated for 45 min. A complex of $DS6A$Ami2B^{WT} bound to the PG fragment L-Ala-D-*iso*-Gln-L-Lys-D-Ala-D-Ala, here referred to as P5^{Lys} ($DS6A$Ami2B^{WT}:P5^{Lys} complex) was obtained by soaking native crystals of $DS6A$Ami2B in the precipitant solution supplemented with 100 mM of P5^{Lys} for 30 s. NAG$_4$, and NAG$_6$ were purchased to Megazyme and the P5^{Lys} peptide was purchased to GenScript. Crystals were flash-cooled in liquid nitrogen without cryoprotection for X-ray data collection, which enabled to measure complete X-ray data sets at 100K using synchrotron radiation at ALBA (Barcelona, Spain) and at ESRF (France). Data sets were integrated, scaled and reduced using XDS (Kabsch, 2010) and AIMLESS (Evans & Murshudov, 2013). Data collection statistics are summarized in Table 1SM. Structure resolution was carried out using the molecular replacement method with PHASER (McCoy *et al*, 2007). For native $D29$N4^{WT}, $D29$GH19^{WT}, $DS6A$GH19^{WT} and $DS6A$Ami2B structures, the coordinates corresponding to models of each individual domain predicted with AlphaFold2 (Evans *et al*, 2022; Jumper *et al*, 2021)were used as search models. For remaining structures, the experimental coordinates of $D29$GH19^{WT} and $DS6A$Ami2B were used as search models. A single and unambiguous solution was obtained in all cases. Initial models were subjected to alternate cycles of model building with COOT (Emsley *et al*, 2010), and refinement with PHENIX (Adams *et al*, 2010) and BUSTER (Bricogne G. *et al*, 2017). NCS restrictions were used during the refinement of structures with more than one protein monomer in the asymmetric unit. Regularised coordinates of the P5^{Lys} molecule were generated with MERCURY (MacRae *et al*, 2020). The geometry of the final models was validated using MOLPROBITY (Chen *et al*, 2009). Figures were generated using PYMOL (Version 3.1; Schrödinger). The atomic coordinates and structure factors have been deposited in the Protein Data Bank under the accession codes indicated in Table 1SM.

2.3. Sequence and tertiary structure alignment

Comparative analysis of $D29$N4, $D29$GH19, $DS6A$GH19, and $DS6A$Ami2B domains with homologue domains was performed using sequence alignment with Clustal Omega Multi Sequence Alignment (Sievers *et al*, 2011). Superimposition of three-dimensional structures was carried out using PYMOL (Version 3.1; Schrödinger). Search for structurally homologue domains of $D29$N4 and $DS6A$Ami2B were carried out using the DALI server (Holm *et al*, 2023).

2.4. Molecular docking with PG fragments

A complex of $D29$GH19^{WT} with a peptidoglycan fragment was modelled by molecular docking using the GOLD (Genetic Optimization for Ligand Docking) software (version 2024.3.0) (Verdonk *et al,* 2003). The PG fragment consisted on four saccharide units linked to a dipeptide moiety (NAG-NAM°[L-Ala-D-*iso*-Gln])$_2$ (4S2P), thus resulting in the complex referred to as $D29$GH19:4S2P. The coordinates of the 4S2P ligand were extracted from a Gram-negative-PG model provided by Prof. Shahrior Mobashery (Meroueh *et al*, 2006), and formatted with

MERCURY (MacRae *et al*, 2020). GOLD uses the Genetic algorithm (GA) method that allows a partial flexibility of the protein and full flexibility of ligand. Thirty independent GA runs, a maximum of 100.000 GA operations, were performed on a set of five groups with a population size of 100 individuals. An ensemble docking was performed using the four independent chains found in the $D29$GH19^{WT}:NAG$_2$:NAG crystal structure and a PG-fragment ligand, with a search efficiency of 200 % (very flexible docking). All water molecules and ligands were removed from the protein. Finally, the CHEMPLP scoring function was used with default parameters to evaluate the solutions.

A model of $DS6A$LysA catalytic domains ($DS6A$GH19-Ami2B) interacting with the bacterial PG was manually generated by superimposing the experimental complexes $D29$GH19^{E228Q}:NAG$_4$ and $DS6A$Ami2B:P5^{Lys} onto the SAXS model using PyMol (Version 3.1; Schrödinger). The resulting model was next correlated with a larger Gram-negative PG scaffold reported by NMR (Meroueh *et al*, 2006).

2.5. Small-angle X-ray scattering (SAXS)

SAXS experiments were performed at the beamline B21 of the Diamond Light Source (Harwell Campus, UK) (Cowieson *et al*, 2020). Samples of 45 µl of 6His-tagged and untagged $DS6A$GH19-Ami2B protein at 5.9 mg/ml were loaded into a SRT-C SEC-300 (Sepax) column equilibrated in buffer (20 mM Tris pH 8.0, 150 mM NaCl and 1 mM DTT) and connected to an Agilent 1200 HPLC system at 18 °C. Continuously eluting samples were exposed for 3 s in 10 s acquisition blocks using an X-ray wavelength of 1 Å, and a sample to detector (Eiger 4M) distance of 3.7 m. The data covered a momentum transfer range of $0.0032 < q < 0.34$ Å^{-1}. The frames recorded immediately before elution of the sample were subtracted from the protein scattering profiles. The Scåtter software package (www.bioisis.net) was used to analyse data, buffer-subtraction, scaling, merging and checking possible radiation damage of the samples. The R_g value was calculated with the Guinier approximation assuming that at very small angles $q < 1.3/R_g$. The particle distance distribution, D_{max}, was calculated from the scattering pattern with GNOM, and shape estimation was carried out with DAMMIF/DAMMIN, all these programs included in the ATSAS package (Manalastas-Cantos *et al*, 2021). The proteins molecular mass was estimated with GNOM. Interactively generated PDB-based homology models were made using the program COOT (Emsley *et al*, 2010) by manually adjusting the X-ray structures obtained in this work, into the envelope given by SAXS until a good correlation between the real-space scattering profile calculated for the homology model matched the experimental scattering data. This was computed with the program FoXS (Schneidman-Duhovny *et al*, 2016).

3. Results and discussion

3.1. *D29*N4 exhibits a cysteine peptidase-domain belonging to the C40 protease family.

3.1.1. Crystal structure of the *D29*N4 domain

The $D29$N4 structure consists of a papain-fold like core formed by a central six-stranded antiparallel β-sheet (β1'-β1-β5) flanked and one α-helix (α2) (**Fig. 2A**). This core is flanked, on one side, by three additional helices (α1, α3 and α3') neighbouring α2 and the loops interconnecting the four helices. A long-loop (β4-β5 loop) enfolds the β-sheet on the other side. The β-sheet is

capped by a small two-stranded β-sheet (βA-βB), and loops β1'-β1 and β2-β3 at one end. The other end of the β-sheet accommodates two additional loops connecting β1-β2 and β3-β4 strands. This structure is conserved in both molecules integrating the asymmetric unit (RMSD of 0.218 Å). A structural homology search across the Protein Data Bank using DALI server revealed *D29*N4 to exhibit homology with members of the NlpC/P60 large family of cell-wall related cysteine peptidases belonging to the peptidase family C40 (MEROPS data base). The latter include dipeptidyl-peptidases hydrolysing peptidoglycan peptide stem between L-Ala-γ-D-*iso*-Glu and L-Zaa-Yaa (where Zaa corresponds to L-lysine or *m*DAP; and Yaa is D-Ala or D-Ala-D-Ala) such as the autolysin Acd24020 from *Clostridium difficile* (*Cdf*Acd24020, PDB entry 7CFL) (Sekiya *et al,* 2021), YkfC endopeptidase from *Bacillus cereus* (*Bcr*YkfC, PDB entry 3H41) (Xu *et al*, 2010), and two RipA and RipB peptidoglycan hydrolases from *Mtb* (*Mtb*RipA and *Mtb*RipB; PDB entry codes 3PBC and 3PBI, respectively) (Böth *et al*, 2011). This aligns with the PG-hydrolase activity initially predicted for this domain (Payne & Hatfull, 2012), and with recent studies by Gangakhedkar and Jain, where the authors map key residues for *D29*N4 activity using site-directed mutagenesis based on sequence homology

studies and computational modelling of *D29*N4 (Gangakhedkar & Jain, 2024).

*The Bcr*YkfC peptidase has been shown to hydrolyse the L-Ala-γ-D-*iso*-Glu-L-Lys tetrapeptide between the glutamate and lysine residues, and its crystal structure has been determined in complex with the reaction product L-Ala-γ-D-Glu (*Bcr*YkfC:L-Ala-γ-D-Glu, PDB entry 3H41) that binds at positions 2 and 1 of the PG-binding channel (Xu *et al*, 2010). The structural comparison of *D29*N4 with *Bcr*YkfC:L-Ala-γ-D-Glu reveals a large and positively charged channel expected to accommodate the PG substrate in *D29*LysA (**Fig. 2AB**). This channel is formed by residues belonging to the α3 helix and the loops α1-α2, α2-α3, β1-β2, β3-β4 and β4-β5, and it accommodates the catalytic triad highly conserved across the C40 family of cysteine peptidases (Griffin *et al*, 2023; Ozhelvaci & Steczkiewicz, 2023). The latter consists, on the one hand of a nucleophilic cysteine residue (C41) located at the N-terminus of α2 and a histidine residue (H112) that acts as the base (**Fig. 2A**). The catalytic role of both residues has been validated by site-directed mutagenesis in *D29*N4 (Gangakhedkar & Jain, 2024) and in its homologue peptidases *Cdf*Acd24020 and *Mtb*RipA (Squeglia *et al*, 2014). Our crystal

structure of *D29*N4 shows that the position of the catalytic C41 at the active site is fastened by a network of interactions constituted by D40 and R62, forming a salt-bridge interaction, and their H-bond contacts with S42 and G43 residues, which are conserved in other homologue peptidases (**Fig. 2A**). The third constituent of the catalytic triad in C40 peptidases typically consists of a glutamate, histidine, glutamine or asparagine located at the C-terminus of the β3 strand and adjacent the catalytic histidine (Anantharaman & Aravind, 2003). This residue is conserved in *D29*N4 (E137) as well as in its homologue PG-hydrolases *Mtb*RipA (E444) and *Mtb*RipB (E213). The proposed role of this glutamate is to enhance the nucleophilic character of the catalytic cysteine by properly orienting the catalytic histidine towards its imidazole group (Ozhelvaci & Steczkiewicz, 2023), and it has been shown to be necessary for *D29*N4 activity (Gangakhedkar & Jain, 2024). In *D29*N4, the imidazole moiety of H112 orients favourably towards the C41 and E137 side-chains at H-bond distance, thus supporting the correct configuration of the catalytic triad, and the role proposed for E137. While these features align with the *in silico D29*N4 structure predicted by Gangakhedkar and Jain, their computational model shows deviations with respect to our crystallographic structure at the level of the N-terminus of α2 accommodating the catalytic cysteine. These deviations involve D40 and C41 main-chain carbonyl groups, which are stablishing H-bond contacts with the main-chain nitrogen of L44. This forces a distortion of the helical arrangement of α2 unseen in with our crystal structure, where the α-helix exhibits a canonical network of H-bonds.

Structural comparison of *D29*N4 with other N4 domains and homologue peptidases reveals variances in the loops forming the PG-binding channel. This includes longer connectors regions spanning α2-α3, α3-β1' and β1-β2 and β2-β3 in N4 domains. Also, a significantly longer β2-β3 loop is observed in only some LysA-N4 domains, including *D29*LysA. These prolonged loops contain glycine residues expected to confer them with flexibility, which is likely to assist on substrate interaction. This is particularly remarkable in the case of the β1-β2 loop, highly rich in glycine residues, which aligns with the high B-factor values observed in this region (**Fig. 2SM**). Interestingly, this loop fold towards the Y27 in *D29*N4, thus closing the substrate-binding channel. Of note, this closed conformation is preserved in both molecules of the crystal asymmetric unit. The superimposition of *D29*N4 and *Bcr*YkfC:L-Ala-γ-D-*iso*-Glu illustrates how the β1-β2 loop of *D29*N4 is adopting a close conformation that must undergo a conformational change in order to unlock the substrate-binding site and enable substrate interaction akin to the one observed in the *Bcr*YkfC:L-Ala-γ-D-*iso*-Glu complex (**Fig. 2A**). This points at the β1-β2 loop as a mobile element with a potential regulatory role in the interaction of *D29*N4 with the peptidoglycan.

Based on the superimposition of *D29*N4 and *Bcr*YkfC:L-Ala-γ-D-*iso*-Glu, we proceeded to inspect those interactions involved in the stabilisation of the L-Ala-γ-D-*iso*-Glu moiety, common in both mycobacterial and Gram-positive peptidoglycan molecules. This shows a conserved patter of contacts comprising hydrophobic (Y25 and Y27), charged (D40, R62), and neutral-polar (S42, G43, S65) residues decorating the subsites 1 and 2 of the PG-binding cavity (**Fig. 2AC**). Importantly, these residues are conserved across homologue peptidases including other LysA-N4 domains; as well as in *Cdf*Acd24020, where they are crucial in substrate interaction (Gangakhedkar & Jain, 2024), thus supporting a similar key role in

*D29*N4 (**Fig. 2C** and **Table 2SM**). This aligns with homology and mutagenesis studies reported by Gangakhedkar and Jain, where mutations of Y25A and S111A in *D29*N4 are shown to impair the protein activity (Gangakhedkar & Jain, 2024). Moreover, the superimposition of *D29*N4 and BcrYkfC:L-Ala-γ-D-*iso*-Glu also highlights the presence of two arginine residues (R139 and R154) in *D29*N4 that are likely contributing to the 1' site accommodating the carboxylate group of the mDAP (**Fig. 2**). This analysis differs from computational studies by Gangakhedkar and Jain, where the authors propose a *D29*N4:PG model predicting the mDAP moiety to orient towards R109 and Y110 residues of the β1-β2 loop, which seems to adopt an open conformation in the model (Xu *et al*, 2010; Gangakhedkar & Jain, 2024). Our sequence analysis shows that while R109 and Y110 are not conserved in the sequence of other LysA-N4 domains (**Fig. 2C**). However, R139 and R154 do exhibit conservation across mycobacteriophage endolysins thus supporting these two arginines as expected interactors of the mDAP moiety, contributing to stabilising the peptide moiety of the PG in a similar fashion to the one proposed for the *Cdf*Acd24020 autolysin (Sekiya *et al*, 2021).

3.2. *D29*GH19 and *DS6A*GH19 are flexible "short-loop" GH19 domains

3.2.1. General structure of *D29*GH19 and *DS6A*GH19 domains
We have determined the crystallographic structures of the wild-type *D29*GH19 domain (*D29*GH19^WT) solved in three different crystal forms (*D29*GH19^WT-I, *D29*GH19^WT-II and *D29*GH19^WT-III) at 1.6-2.0 Å resolution, and the crystal structure of the wild-type *DS6A*GH19 domain (*DS6A*GH19^WT) at 2.8 Å resolution. *D29*GH19^WT and *DS6A*GH19^WT exhibit a structure rich in alpha helices that arrange in two lobes (**Fig. 3A**). A first lobe is formed by residues provided by N-terminal (residues 179-233 in *D29*GH19^WT) and C-terminal (residues 307-370 in *D29*GH19^WT) regions, referred to as NC-lobe, that comprises helices α1-α4 and α7-α10. The second lobe is built by residues belonging to the middle region of the polypeptide chain (residues 234-308 in *D29*GH19^WT), here referred to as M-lobe, that comprises helices α6-α7. This fold is conserved in the four structures determined, which is reflected by the RMSD values calculated for: i) *D29*GH19^WT-I and *D29*GH19^WT-II (0.320 Å for 180 Cα atoms); ii) *D29*GH19^WT-I and *D29*GH19^WT-III (1.832 Å for 167 Cα atoms) and; *D29*GH19^WT-I and *DS6A*GH19^WT (1.071 Å for 154 Cα) (**Fig. 3A**). According to (Hoell *et al*, 2006), GH19 domains can be classified as "loopful" or "loopless" according to the presence or absence of insertions in loops I-V and C, located towards the substrate-binding groove, that results in larger or smaller number of substrate-interaction subsites, respectively (Hoell *et al*, 2006). The superimposition of *D29*GH19 with endochitinase C from *Secale cereale* (Rye) (*RSC-c*GH19, PDB entry 4DWX), as a representative of a "loop-full" GH19 domain, shows high degree of structural conservation in the two-lobe core structure but clear differences in the length of loops I-V and C, which are much shorter in *D29*GH19 (Fig. 4D and Fig. 4G). This is in line with the sequence alignment of homologue domains found across phages, bacteria and eukaryote, which highlights the presence of short I-V loops in both of *D29*GH19 and *DS6A*GH19 domains, and in GH19 domains present other LysA endolysins (**Fig. 3D**). Recently, GH19 hydrolases have been reclassified in two different subfamilies, referred to as GH19_1 and GH19_2, embracing enzymes with peptidoglycan endo-β-1,4-N-acetylmuramidase and chitin endo/exo-β-1,4-N-acetylglucosaminidase activities, respectively (see <u>CAZY – GH19</u>). Altogether this illustrates LysA-GH19s as "loopless" glycosyl-

Figure 3. General structure of *D29*GH19 and *D29*GH19 domains. (A) Superimposition of *D29*GH19[WT-I] (yellow), *D29*GH19[WT-II] (light grey) and *D29*GH19[WT-III] (dark grey) crystal structures depicted as cartoons. The catalytic residues and bound molecules coming from the crystallisation condition are represented as sticks, and calcium cations are shown as green spheres. MES and MPD molecules bound to *D29*GH19[WT-I] are coloured in black; and MES and EG molecules bound to *D29*GH19[WT-II] are coloured in light grey. (B) Superimposition of *D29*GH19[WT-I] (yellow) and *DS6A*GH19[WT] (blue) crystal structures depicted as cartoons, showing the conserved catalytic dyad as sticks. Calcium cations, MES and MPD are represented as in panel (A). (C) Overlay of the calcium-binding site in *D29*GH19[WT-I] (yellow) and the corresponding region in *DS6A*GH19[WT] (blue). (D) Details of the sequence alignment of *D29*GH19 and *DS6A*GH19 with other GH19 domains found across bacteria, eukaryota and phages, including mycobacteriophage LysA endolysins: *RSC-c*GH19 (UniProt entry Q9FRV0), endochitinase from *Carica papaya* (*Cpp*CHIT, UniProt entry P85084), chitinase 2 from *Oryza sativa subsp. japonica* (*Ost*CHI2, UniProt entry Q7DNA1), endochitinase A from *Zea mays* (*Zmy*CHIA, UniProt entry P29022), chitinase A from *Gemmabryum coronatum* (*Gcr*CHIA, UniProt entry A9ZSX9), chitinase A from *Streptomuces griseus* (*Sgr*CHIC, UniProt entry O50152), chitinase G from *Streptomyces coelicolor* (*Scl*CHIG, UniProt entry Q8CK55), endolysin from *Pseudomonas phage Ka2* (*Ka2*Lys, UniProt entry A0A9E7MJ36), endolysin from *Salmonella phage SPN1S* (*SPN1S*Lys, UniProt entry H2D0G4), placed Lysin A from *Mycobacteriophage Redi* (*Redi*LysA, UniProt entry G3M5B9), to be red accelerate displace Lysin A from *Mycobacteriophage BarrelRoll* (*BarrelRoll*LysA, UniProt entry G3MCI0), Lysin A from *Mycobacteriophage Ellie* (*Ellie*LysA, UniProt entry A0A7G8LLY0), Lysin A from *Mycobacteriophage MasterPo* (*MasterPo*LysA, UniProt entry A0A7G8LGT1), Lysin A from *Mycobacteriophage AnnaL29* (*AnnaL29*LysA, UniProt entry S5VWI7), Lysin A from *Mycobacteriophage Bombshell* (*Bombshell*LysA, UniProt entry A0A7G8LLL9), and Lysin A from *Mycobacteriophage Violet* (*Violet*LysA, UniProt entry G3MEJ3). (E) Structural superimposition of the *D29*GH19[WT-I] and the "loop-full" GH19 structure, corresponding to *RSC-c*GH19 (PDB entry 4DWX), represented as yellow and pink cartoons, respectively, with catalytic glutamates depicted as sticks. Loops I, II, III, IV, V and C of *RSC-c*GH19 are labelled and highlighted as thicker dark-pink loops. (F) and (G) show the electrostatic-potential surface representation of *D29*GH19[WT-I] and *DS6A*GH19[WT] domains, respectively. Positively and negatively charged residues are represented in colour blue and red, respectively. Catalytic residues and ligands bound at the substrate-binding groove are labelled.

hydrolases belonging to the GH19_1 subfamily, exhibiting smaller substrate-binding channels when compared to their "loopful" homologues belonging to the GH19_2 subfamily (Hoell *et al*, 2006).

Despite the high structural conservation observed between *D29*GH19 and *DS6A*GH19 models, significant differences are identified in the architecture of the M-lobe as well as subtle deviations in the inter-lobe arrangement. In *D29*GH19^{WT}, M-lobe exhibits an EF-hand-like calcium-binding site located at the apical region of the lobe, where the cation appears stabilised by interactions with five amino acids (T253, E255, D245, D257 and D259) resulting in a hepta-coordinated calcium complex (**Fig. 3C**). The presence of a calcium cation at this position is fully supported by a strong signal observed in the 2Fo-Fc electron density map (sigma = 15) that is well interpreted by the refinement of a calcium cation, as well as by the presence at this position of a peak of sigma > 9 in the anomalous electron density map calculated (**Fig. 3SM**). The calcium-interacting site is structurally conserved in both *D29*GH19^{WT-I} and *D29*GH19^{WT-II} structures, but it is not observed in the *D29*GH19^{WT-III} model, where this region is disordered. Interestingly, this calcium-binding site is not present in the *DS6A*GH19^{WT} domain, where three conserved residues (E70, T78 and D82) exhibit a different orientation, and two of the acidic residues interacting with calcium in *D29*GH19^{WT} (E255 and D257) are not conserved (W71 and A80 in *DS6A*GH19^{WT}) (**Fig. 3C-D**). To our knowledge, the calcium-binding site present in *D29*GH19 has not been observed in other GH19 structures reported. This is exemplified by the sequence alignment shown in Fig.3D that also indicates poor conservation of this calcium-binding site across LysA endolysins (**Fig. 3D**). Superimposition of *D29*GH19^{WT-I} and *D29*GH19^{WT-II} models with *D29*GH19^{WT-III} and *DS6A*GH19^{WT} structures highlights the presence of an inter-lobe shift, which aligns with higher RMSD values calculated (**Fig. 3AB**). In this shift, NC and M-lobes move as independent scaffolds, which agrees with the RMSD values calculated for the individual M-lobes (0.195 Å for 44 Cα atoms) and NC-lobes (0.240 Å for 103 atoms) of *D29*GH19^{WT-I} and *D29*GH19^{WT-III} coordinates; and with the RMSD values calculated for M-lobes (0.57 Å for 57 Cα atoms) and NC-lobes (1.115 Å for 84 Cα atoms) of *D29*GH19^{WT-I} and *DS6A*GH19^{WT} structures. In the case of *D29*GH19, the inter-lobe rotation observed supports an inherent flexibility of this domain relevant for the interaction with the PG substrate, as already proposed in other GH19_1 homologues (Ohnuma et al, 2013).

The substrate-binding site is embedded into a large grove formed by residues belonging to both M and NC-lobes that allocates the catalytic dyad and two glutamate residues (E228/E237 in *D29*GH19; E53/E62 in *DS6A*GH19), in the middle of the cavity (**Fig. 3**). This catalytic dyad is conserved across GH19 members, and their implication in catalysis has been verified through site-directed mutagenesis and biochemical studies in *D29*LysA, as well as the homologue Chitinase G from *Streptomyces coelicolor* and in the bacteriophage *SPN1S* endolysin, reported elsewhere and that we recover in **Table 3SM** (**Fig. 3D**) (Hoell *et al*, 2006; Park *et al*, 2014). Inspection of the electrostatic potential surface of *D29*GH19 and *DS6A*GH19 domains shows along PG-binding channels numerous negative charges, protruding from aspartate/glutamate side-chains as well as main-chain carboxyl groups, which is a typical feature of saccharide-interacting sites/grooves found in carbohydrate-binding domains/hydrolases (**Fig. 3FG**). In *D29*GH19^{WT-I} and *D29*GH19^{WT-II} structures, molecules coming from the crystallisation conditions bind adjacent the catalytic dyad, mimicking saccharide moieties of the substrate at positions -1 (a molecule of MES in *D29*GH19^{WT-I} and *D29*GH19^{WT-II}) and +2 (molecules of MPD or ED in *D29*GH19^{WT-I} and *D29*GH19^{WT-II}, respectively), thus defining a PG-binding groove with up to three positions (-1, +1 and +2) (**Fig. 3AF**).

3.2.2. *D29*GH19 conserves structural features required for PG hydrolysis and exhibits chitinase activity in *crystallo*

To gain more insights into substrate binding and hydrolysis, crystallographic complexes with NAG$_n$ oligosaccharides as glycan analogues were pursued by soaking using the catalytically-active *D29*GH19^{WT} domain and the catalytically inactive mutant E228Q (*D29*GH19^{E228Q}) domain. This led us to solve the structure of *D29*GH19^{E228Q} alone and bound to NAG$_4$ (*D29*GH19^{E228Q}:NAG$_4$) at 1.45 Å and 1.83 Å resolution, respectively. Comparison of *D29*GH19^{E228Q} and *D29*GH19^{WT} crystal models indicates high structural conservation (RMSD of 0.118 Å for 176 Cα atoms) (**Fig. 4SM**). The *D29*GH19^{E228Q}:NAG$_4$ structure consists of one molecule of the complex in the asymmetric unit, in which the saccharide binds from positions -3 to +1 of the PG-binding channel. In addition, the structure of two more complexes were obtained, by soaking *D29*GH19^{WT} crystals into a solution containing NAG$_6$, at 2.34 and 3.10 Å resolution (**Fig. 4** and **4SM**). The first one consists of four complexes of *D29*GH19^{WT} bound to NAG and NAG$_2$ (*D29*GH19^{WT}:NAG:NAG$_2$) in the asymmetric unit, where NAG$_2$ and NAG are products resulting of NAG$_6$ hydrolysis *in crystallo*. The four *D29*GH19^{WT}:NAG:NAG$_2$ complexes exhibit the same structure (RMSD values between 0.151 Å and 0.264 Å), with saccharide fragments binding with the same pose at positions -2 (NAG) and, +1 and +2 (NAG$_2$) of the PG-binding channel. The other complex consists of one *D29*GH19^{WT} domain bound to NAG (*D29*GH19^{WT}:NAG) in the asymmetric unit, where the monosaccharide is a product of NAG$_6$ hydrolysis *in crystallo* binding at position -2 of the PG-binding site (**Fig. 4** and **4SM**). Superimposition of complexes and native structures shows conservation of the fold across all the models, and the former evidences the capability of *D29*GH19^{WT} to interact with up to five saccharide rings at positions going from -3 to +2 (**Fig. 4A** and **4SM**). Interestingly, superimposition of *D29*GH19^{WT}:NAG:NAG$_2$ and *D29*GH19^{WT}:NAG complexes with native and *D29*GH19^{E228Q}:NAG$_4$ structures reveals an inter-lobe rearrangement, similar but larger to the one observed between wild-type native structures, resulting in a closure of the PG-binding channel by up to 5 Å in *D29*GH19^{WT}:NAG:NAG$_2$ and *D29*GH19^{WT}:NAG complexes (**Fig. 4AB**). This rearrangement can be explained as a rotation-in-block of the NC-lobe, which pivots towards a central tryptophan (W268) of the M-lobe that is partially conserved as a tryptophan or a proline in GH19 domains (**Fig. 4** and **4SM**). This movement involves the motion of several hydrophobic residues of helices α3, α4 and α7 that turn over the tryptophan to enable NC-lobe shift. This inter-lobe rearrangement is also observed in the structure of other GH19 chitinases such as *RSC-c*GH19 solved in native and NAG$_n$-bound conditions (**Fig. 4SM**), thus showing how this flexibility extends from "loopful" to "loopless" GH19 domains as a general feature of this family of hydrolases.

A more detailed inspection of the PG-binding channel in *D29*GH19^{WT}:NAG and *D29*GH19^{WT}:NAG:NAG$_2$ reveals that the binding mode of the NAG ring at position -2 is conserved in both complexes (**Fig. 4A**). Thus and for simplicity, subsequent structural analysis will focus on *D29*GH19^{E228Q}:NAG$_4$ and *D29*GH19^{WT}:NAG:NAG$_2$ complexes. Comparison of the PG-

binding channel in these complexes reveals a network of direct or water-mediated interactions participating in saccharide stabilisation that involves several hydrophobic residues (Y238, A239, Y244, L250, I269, M270, W315 and Y316) and polar amino-acids that include the catalytic residues (E/Q228, E237, T272, K274, N276 and N341), provided by both lobes (**Fig. 4C-D**). Interactions stablished via M-lobe are conserved in both *D29*GH19^{E228Q}:NAG$_4$ and *D29*GH19^{WT}:NAG:NAG$_2$ complexes, where it is worth highlighting an H-bond network bridging Y244, T272 and the catalytic E237 with a water molecule (W) and the acetyl group of the NAG ring at position -1. These contacts enable the correct orientation of the catalytic base E237 towards the water, thus allowing the CW molecule to locate at a distance (3.4 Å) and orientation to carry out the nucleophilic attach over the C1 atom of the substrate (**Fig. 4**). This H-bond network is observed in structures here reported for the WT and E228Q mutant of *D29*GH19, and other GH19s structures, constituting a structural hallmark of the catalytic mechanism of this family of glycosyl hydrolases (**Fig. 4SM**). The interaction pattern via NC-lobe residues does however vary between *D29*GH19^{E228Q}:NAG$_4$ and *D29*GH19^{WT}:NAG:NAG$_2$ structures (**Fig. 4**). This results of the close conformation of the PG-binding channel in *D29*GH19^{WT}:NAG:NAG$_2$, also observed in the *D29*GH19^{WT}:NAG complex. In this close conformation, the side chain of the catalytic E228 shifts towards the hydroxyl group at C4 of the NAG in position +1. This results in a strong H-bonding interaction (distance of 2.8 Å) between both hydroxyl and carboxyl groups compatible with proton exchange during substrate scission (**Fig. 4** and **4SM**). Moreover, the close arrangement of the PG-binding channel results in additional contacts and/or optimisation of pre-stablished

interactions of *D29*GH19 with the ligand. On the one hand, there is an approximation Y315 towards the NAG ring in position -2 that strengthens this interaction. On the other hand, the C-terminus of α4, the N-terminus of α10, and the liker connecting helices α9 and α10 shift to pack more tightly against the products, thus resulting in additional contacts via H227, R350, N341 and N346 side chains, and I340 and N341 main-chain carbonyl groups (**Fig. 4D**). These residues are conserved in *DS6A*GH19 but for Y238 and M270 (substituted by a distal tryptophan and a glutamine in *DS6A*GH19, respectively) and a tyrosine (Y98) edging the position -3 of the PG-binding channel in *DS6A*GH19 are missing in *D29*GH19 (D275) (**Fig. 4D** and **4SM**). Our sequence analysis shows high conservation of these amino-acids involved in ligand interaction, but for Y138, A139, K274 in *D29*GH19, that exhibit more divergent variations (**Fig. 4** and **4SM**). Importantly, mutation of several of these residues in *D29*GH19 and lysins *SPN1S* endolysin (reported elsewhere) evidence a clear impact in their interaction with the substrate or its hydrolysis (see **Table 2SM**), which altogether supports their relevance in *D29*GH19 and *DS6A*GH19 performance.

Next, we decided to further explore the mechanism of substrate recognition of a PG fragment by *D29*GH19 using molecular docking approaches. To do so, an unbiased docking search was performed using the close conformation of *D29*GH19^{WT} and the PG-fragment ligand (NAG-NAM[L-Ala-D-*iso*-Gln])$_2$ (4S2P). This yielded eleven top solutions (out of thirty) where the PG fragment consistently binds across positions -2 (NAG), -1 (NAM), +1 (NAG) and +2 (NAM), exhibiting a very conserved pose for the glycan strand and discreet variations in the orientation of the peptide moiety (**Fig. 4**

Figure 4. *D29*GH19 complexes with PG analogues. (A) Structural superimposition of *D29*GH19^{E228Q}:NAG$_4$ and *D29*GH19^{WT}:NAG:NAG$_2$ complexes with *D29*GH19^{WT-I}, represented as cartoons coloured in green, pink and yellow respectively. Bound saccharide molecules are represented as sticks coloured in black (NAG$_4$) and light white (NAG and NAG$_2$). (B) Detail of the electron density map 2Fo-Fc (blue mesh) corresponding to the NAG$_4$ molecule of the *D29*GH19^{E228Q}:NAG$_4$ complex contoured at sigma = 1.0. (C) and (D) show details of the interaction with NAG$_n$ fragments identified in *D29*GH19^{E228Q}:NAG$_4$ and *D29*GH19^{WT}:NAG:NAG$_2$ complexes, respectively. H-bonds are highlighted as grey dashed lines. PG analogues and residues involved in ligand binding are represented as sticks following the same colour code as in (A). Water molecules and calcium cations are presented as red and dark green spheres, respectively. (E) Detail of the complex *D29*GH19^{WT}:4S2P predicted by docking showing the substrate-binding cavity (protein in yellow). The catalytic residues and the PG fragment are represented as blue and yellow sticks, respectively.

and **4SM**). Inspection of the top docking solution, referred to as *D29*GH19^WT:4S2P complex, shows how the location of the four saccharide rings aligns well with the NAG$_n$ molecules interacting with *D29*GH19 *in crystallo*, which supports the reliability of the *in silico* model (see **Fig. 4SM**). The peptide steps protrude from the glycan strand exhibiting a conserved pose at the D-lactyl moieties, which interact with hydrophobic residues at position -1 (Y224 and I250) and position +2 (A239) via their methyl groups, and form H-bond contacts through the carbonyl group at position -1 with R247. L-Ala and D-*iso*-Gln residues of the ligand do show divergent orientations sustained by plausible interactions with residues NC- and/or M-lobes (see Fig. 4E). Importantly, the model shows the only glycan arrangement enabling the positioning of the peptide stems without forming clashes with the protein, as a shift of the NAG-NAM saccharides in the channel would generate collisions between the peptide steps pending from C3 carbon of the NAM units and the enzyme. While this further supports the likelihood of our computational supports, it also features the *D29*GH19 domain as an enzyme with muramidase, that not glycosaminidase, activity as the reducing saccharide after hydrolysis at position -1 would be a NAM residue and not NAG. To our knowledge, our studies provide the first evidence supporting the GH19_1 subfamily of hydrolases to function as muramidases cleaving the β-(1,4)-glycosidic bond connecting NAM and NAG saccharides of the PG.

3.3. *DS6A*Ami2B consists of a catalytically competent PG-amidase domain

3.3.1. *DS6A*Ami2B exhibits catalytic features characteristic of the Amidase_2 family

We have solved the crystal structure of *DS6A*Ami2B in native conditions, which consists of three polypeptide chains in the asymmetric unit. All of them exhibit the same fold (RMSD of 0.086-0.118 Å for >150 Cα) consisting of a central beta sheet built by two antiparallel (β1 and β4) and four parallel beta strands (β2, β3, β5 and β6) flanked by five alpha helices (α1-α5) (**Fig. 6A**). Structural homology analysis using DALI indicates highest homology with members of the Amidase 2 family, including the or the cell-wall remodelling amidases from *Staphylococcus aureus* AmiA (*Srs*AmiA) (PDB entry 4KNK; RMSD 3.194 for 36 Cα atoms) (Büttner et al, 2014) and AmpDh2 (*Srs*AmpDh2) (PDB entry 4BJ4; RMSD 0.996 for 60 Cα atoms) (Martínez-Caballero *et al*, 2013), or the amidase AmiD from *E. coli* (*Ecl*AmiD) (PDB entry 2WKX; 1.168A for 62 Cα atoms) (Kerff *et al*, 2010), which aligns with previous homology predictions reported (Urdániz *et al*, 2022). Members of the Amidase 2 family consist of zinc-dependent hydrolases exhibiting *N*-acetylmuramoyl-L-Alanine amidase activity (EC:3.5.1.28), and that cleave the PG molecule between the D-lactyl group of the NAM residue and the first L-Ala amino-acid of the peptide stem (Büttner *et al*, 2015). Activity that has been demonstrated for homologues including the Ami2B domain of the LysA encoded by the mycobacteriophage TM4 (*TM4*LysA) (Urdániz *et al*, 2022), the prophage Ba02 Ba02 PlyL endolysin encoded by *Bacillus anthracis* (*Bnt*PlyL) (Low *et al*, 2005), AmiE autolysin from

Figure 5. (A) Crystal structure of *DS6A*Ami2B domain represented as green cartoons and highlighting the loop β5-α4 loop in colour purple. Details of the active site are shown on the right box. Catalytic zinc and residues are depicted as a yellow sphere and sticks, respectively. A phosphate molecule found coordinating to the catalytic zinc is presented as sticks. (B) Electrostatic-potential surface representation of *DS6A*Ami2B with the loops contributing to form the PG-binding channel, and the zinc (yellow sphere) and residues involved in catalysis indicated with labels. (C) Details of the sequence alignment of *DS6A*Ami2B with other Amidase_2 domains: AmiA from *Staphylococcus aureus* (*Srs*AmiA; UniProt entry Q2FZK7), AmiE from *Staphylococcus epidermis* (*Spd*AmiE; UniProt entry O33635), LytA from *Streptococcus pneumoniae* (*Spn*LytA; UniProt entry P06653), PlyL from *Bacillus anthracis* (*Bnt*PlyL, UniProt entry A0A6H3AMF3), XlyA from *Bacillus subtilis* (*Bsb*XlyA; UniProt entry P39800), AmiD from *Escherichia coli* (*Ecl*AmiD; UniProt entry P75820), AmpDh2 from *Pseudomonas aeruginosa* (*Prg*AmpDh3; UniProt entry Q9I5D1), AmpDh2 from *Pseudomonas aeruginosa* (*Prg*AmpDh2; UniProt entry Q9HT86), AmpD from *Cytrobacter freundii* (*Cfr*AmpD; UniProt entry P82974), and LysA endolysins from the mycobacteriophages *Stinger* (*Stinger*LysA; UniProt entry G8I9G3), *Nigel* (*Nigel*LysA; UniProt entry B3VLX1), *Bxz1* (*Bxz1*LysA; UniProt entry Q852W9), *Pacc40* (*Pacc40*LysA; UniProt entry B5U576), *SG4* (*SG4*LysA; UniProt entry G8I9P6), *Che9d* (Che9dLysA; UniProt entry Q855S8), *Barnyard* (*Barnyard*LysA; UniProt entry Q856D3), *TM4* (*TM4*LysA; UniProt entry Q9ZX49), *Corndog* (*Corndog*LysA; UniProt entry Q856M5), *Marvin* (*Marvin*LysA; UniProt entry G1BNC1), *Wildcat* (*Wildcat*LysA; UniProt entry Q19Y11)

```
                 α1              β1              β2            α2              β3          β4
DS6ALysA     AWTGDPVWLEDVLRPVLG--------DRLKT-----LPSWQMYGHG-DFKDIR--GVMHIHHTGNA-----RETAESIRKGRPDLAGPL----------SNIHIAP--------DGTVTLVA  289
MarvinLysA   PVSGDPVWLEVLRRKRLG--------DRLVV-----HDGWKEYGTGGVMGEIW--GVMIHHTGNR-----NASWESIRNGRPDLAGPL----------SQALIAP--------DGKFHLVA  279
Pacc40LysA   GFSGDPVWLEDVLREALG--------DRLVVA----QADWKERGTGGVMGDIW--GVMIHHTGNR-----RETVAGIRDGRPDLAGPL----------SQCLITP--------DGKCHLIA  231
SG4LysA      AWTGDPVWLEEVLRPALG--------DRLKT-----LPGWQNSGHG-DFKDIR--GVMHIHHTGNA-----RETAESIRKGRPDLAGPL----------SNIHIAP--------DGTVTLVA  256
Che9dLysA    -VGVCWNHAGRGSYPWLPTDN--------DRLKT-----LPGWQNSGHG-DFKDIR--GVMHIHTGDS-----RESAQSIRNGRPDLAGPL----------ANIHIAP--------DGTVTIVA  280
TM4LysA      PWTGDPVWLADVLRA-BG--------LNVVE-----LPGWLDRGHG-DMGRLW--GVVCHHTGDS-----NTPSSEIAFH-PSL-GLC---------SQIHLAR--------NGTVTLCG  262
BarnyardLysA SRRGDVVWLPDILRS-AG--------LPVQV-----ADGANNRGHG-DMADVW--GIVDHHTGSD-----NASWQSIAFH-PSL-GLA---------SQLHLSY--------GGLFTVCG  80
Bxz1LysA     NWRGDPVWLKEVLEA-AG--------LVCHV-----TDGAYNRGHG-DFGEIW--GVVAHHTGSY-----GETPKGIAQH-PSL-GLA---------SQLYLGR--------NGEYTLCG  251
NigelLysA    AWRGDPLFLPQLLRA-FGV--------SVTTYT---DADGIPWDQRHG-DFGATI--WVLWHHTGSV-----NETDNGIAHH-PAL-GLA---------ANMLIHP--------DGHVVLTG  296
WildcatLysA  GFTGDDPVWMLADVLRA-BG--------LNVVE-----VDGWQNRGGQ-DQGTFW--GVMFHHTGSV-----GETPGIAQH-PTL-GLA---------AHILIKP--------DGHVVMGE  274
PrgAmpDh3    ------------------MLTIDY-----------NSYRTTTPYGKRVR---FLVLHYTALD-------FAASVKALTT-GAAS--------------AHYLIPAPHDPSYKAAGFKGQRIFNLVA  72
PrgAmpDh2    SFVLLAFTLSLLAGCSSG----------PRL--------FDYTSANQDSRVQ--FIVLHYTSTD---------FIVHVNHWRKDPEL---------------AHYLIGDDEP----------ATVYRLVD  66
EclAmiD      --MRRFFWLVAAALLLAGCAG-----EKGIVEKEGYQL--DTHRQAQAAYPRIK--VLVIHYTANDPSTW--FDSSLATLTD-KQVS---------------SHYLVPAVPPRYN------GKPRIWQLVP  92
SpnLytA      ----------MEINV---------SKLRTDLPQVGVQFPYRQVRAHSTDNPSTV-QNEADYHWRKDPEL---------------GNGCIMQVGP  66
SpdAmiE      KYTFKVNSGINNYIRKK-------NMKAPRIEEDYTSYFP--KYGYRNG-VGRPE--GIIVHDTANDRSTING--ELSYMKNNYQNAF----------VHAFV----------DGDRIIETAP  401
SrsAmiA      PKTSLPKYKFPQVNSSINDYICKNNLKAPKIEEDYTSYFP--KYAYRNG-VGRPE--GIIVHDTANDRSTING--EISYMKNNYQNAF----------VHAFV----------DGDRIIETAP  305
BsbXlyA      ------------------MVNILQD--------FIPVGANNRPGYAMTPL--YITVHNTANTAVGADAAAHARYLKNPDT----------TSWHFTV----------DDTEIYQHLP  70
BntPlyL      ------------------MEIRKKL,---VVPSKYGTKCPYTMKPK--YITVHNTYND-APAENEVNYMITNNNE-----------VSFHVAV----------DDKQAIQGIP  67
CfrAmpD      -MLLDEGWL---------AEARRV-----PSPHYDCRPDDENPS--LLVVHNISLPPGEFGGPWIDALFTGTIDPNNAHPYFAGIAHLRVSAHCLIRR---DGEIVQYVPN  89

                       α3  β5          Loop β5-α4          α4          β6              α5
DS6ALysA     -AGVCWNHAGAGSYPWLPTNNAWHMIGICEAMPTIRP--NGTYDEREPWPDAQIIAMRDTCAALTKRLGWDAG---------RVIGHKEYAGA------SQGKWDPGNDLMGWFRGEVAKAMRG---  395
MarvinLysA   -VGPCWNHAGVGSYPGLGSD-GNRRAIGFECAMPTIRP--DGSFDKGERWPDAQILTMRDATAVLEHLGHDHT---------HVCGHKEFNK------YQGKWDPGNGMDMNWFRGEVRKALAG---  383
Pacc40LysA   -VGPCWNHAGVGSYPGVGTNNGQRLIGFECAMPTIRP--DGSFDPAQRWPDAQILTMRDATAAVLKRLGHDSK---------HVIGHKEWAGA------TQGKWDPGNDIMPWFRGEVAKARG----  338
SG4LysA      -VGPCWNHAGRGSYPMLPTDNAWHMIGICEAMPTVT---NGKYDPAERWPDAQILSMRDTCAALATKLGHPAT---------RVIGHKEYAGA------AQGKWDPGNDIMPWFRGEVAKAMRG---  361
Che9dLysA    -VGVCWNHAGGGSYPWLPTDNAWHMIGIECAMPTIRP--PKFDPLPKGYDPAERWPDAQIISMRDTCAALALELPAT------RVIGHKEWAGT------AQGKWDPGGDIMNLFRDKEIDAMG----  388
TM4LysA      -VGIAWNHAGVGSYPWPGLGTNNANPRTIGIEAANDGGGR-PGRPH--RADWPAVQYDHYVEGNAAILRFLKHNEPVT-----HSIGHKEWAGA------AQGKWDPGGIDMNILFRKDIESAMRG---  366
BarnyardLysA -VGIAWNHAGNGSWPGLGTNNANPRTTGIEAANDGGGR-PGRPH--RADWPAVQYDHYVEGNAAILRFLKHNEPVT-----HSIGHKEWAGA------AQGKWDPGGIDMNILFRKDIESAMRG---  185
Bxz1LysA     -VGIAWNHAGGGSYPGLGTNNANRLTIGIEAANDGGGR-PGRPH--SMDSDVQYNAYVRGVAAILRKLGRDSS---------HVEGHKEWAGT------AQGKWDPGGIDMNITFRADVARVMGE---  355
NigelLysA    -SGIAWNHBGAGIYPGIPEDGIQVSIGIECSYG----PDRNGQY--TLPWFEAQMNAMIAVGGAISWFLGDTLPPS------RQIAHKEWAGRDNPLGVNKQGKPDPGNLDMTWFRAQIAARAAA---  404
WildcatLysA  -IGKANHAGRDYPGLPTNNANPVMIGVEVAIL---PRKDAPH--KENNAVDVVRQLMSMYNIPIE------VNNRTHQSWSG-----KYCPHRMLAEGRWGAFIQKVKSGNV  377
PrgAmpDh3    EEDRAWNHAGVSGWAR--RDMLNDTSIGIEIVNLA--RDDDGVF-TFPDYERSQINALKQLAKNILQRYPDMTPK------NVVGHSDIAV--------GNKSDPGPK-LPWKELYEAGIGAW---  174
PrgAmpDh2    ENRRAWNHAGVSEWQG--RTWLNATSIGIEIVNGQYRDTPQGRV--WPSFDQAQAIPLLKDIAKRHGITPD------RIIGHSDIAP-----------NGTHADIAP--------NGTHADIAP----  189
EclAmiD      --ATRLNDTSIGIELENRGWQKSAGVKY--FAPPEPAQIQALIPLAKDIYARYHIKPE----------QRKQDPGDLDMNIFRDVQQEIGAW---  194
SpnLytA      VDNGAWDVGGG--W-----NAETYAAVELIESHSTK---------EEFM-TDYRLYIELLRNLADEAGLPKTLDTGSLA-GIKTHEYCTNNQPN----NHSDHVDPYPYLQFKAKWGISREHDIENG  170
SpdAmiE      TDYLSWGAGP----Y-----GNQRFINVEIVHTHDYD--------SFA-RSMNNYADYAATQLQYYNLKPDQAGDSNGTVWTHAAISNFL------GGTDHADPHGYLRSHNYSYAELYDLIYEK  503
SrsAmiA      TDYLSWGVGA----V-----GNPRFINVEIVHTHDYA--------SFA-RSMNNYADYAATQLYYGLKPDSAEYDGNGTVWTHYAVSKYL------GGTDHADPHGYLRSHNYSYDQLYDLINEK  406
BsbXlyA      LNENGWACDGDGNGS--------GNRASIGIEICENAGDD-------------FA-KATANAQWLIKTLMAEHNISLA-----------VNVVPHKYWSG----------KECPRKLLDTWDSFKAGIGGGQ  159
BntPlyL      WERNAWACGDGNGP--------GNRESIEVEICYSKSGG---------DRYY-KAENNAVDVVRQLMSMYNIPIE------VNNRTHQSWSG-------KYCPHRMLAEGRWGAFIQKVKSGNV  160
CfrAmpD      FDKRAWNHAGVSSYQG--RERCNDFSIGIELEGTDT----------LAYTDAQYQQLAAVTNALITRYPAIAN------NMTGHCNIAP--------ERKTDPGPS-FDWARFRALVTPSS---  182
```

Staphylococcus epidermis (*Spd*AmiE) (Zoll *et al*, 2010) and LytA autolysin from *Streptococcus pneumoniae* (*Spn*LytA) (Mellroth *et al*, 2014; Sandalova *et al*, 2016).

Inspection of the structure of *DS6A*Ami2B reveals a large T-shape channel located at the C-terminus of the beta-sheet core formed by loops β2-α2, α2-β3, β4-α3, β5-α4, and β6-α5 (**Fig. 5B**). The latter constitutes the substrate-binding site that can be described as a transversal glycan-binding groove exhibiting a negatively charged core, and a perpendicular channel involved in the interaction with the peptide moiety and rich in positively charged residues. In line with our homology analysis, a zinc cation locates at the centre of this channel, where it coordinates to two histidines (H251 and H364) and one aspartate (D376) located at the loop β6-α5 and the C-terminus of β2, respectively. These residues are highly conserved across Amidase_2 members, and they have been shown to be essential for the activity of members of this family amidases including *TM4*LysA (Urdániz *et al*, 2022), *Spd*AmiE (Zoll *et al*, 2010), and *Spn*LytA (Mellroth *et al*, 2014; Sandalova *et al*, 2016) (**Fig. 5C** and **Table 4SM**). In *DS6A*Ami2B structure, the metal is also interacting with a phosphate molecule, present in the crystallization condition, resulting in a tetrahedral coordination of the zinc. The proposed role of the zinc is to polarize the water molecule meant to perform the nucleophilic attack onto the scissile bond of the substrate (Büttner et al, 2014; Sandalova et al, 2016). In addition, a catalytic dyad has been proposed to assist the zinc during PG hydrolysis (Zoll *et al*, 2010; Mellroth *et al*, 2014; Sandalova *et al*, 2016). This dyad consists of a glutamate residue located at the C-terminus of β5, which would act as an acid donating a proton to the newly formed N-terminus of the peptide (Zoll *et al*, 2010; Mellroth *et al*, 2014; Sandalova *et al*, 2016). And a histidine charged residue, located two residues upstream the zinc-coordinated aspartate, and proposed to participate in the stabilization of transition state (Zoll *et al*, 2010; Mellroth *et al*, 2014; Sandalova *et al*, 2016). Both residues are conserved in *DS6A*Ami2B (E317 and K374), which lays adjacent the catalytic zinc orienting its side chain towards the cofactor. Altogether these findings support that *DS6A*Ami2B is a catalytically competent amidase exhibiting conserved features of the Amidase 2 family (**Fig. 5**) (Büttner *et al*, 2015). This is well illustrated by the sequence and structural conservation of *DS6A*Ami2B with other homologue amidases depicted in **Fig. 5 and 5SM**. The latter also highlights the fourteen-residues long β5-α4 loop flanking the PG-binding channel. This long loop resembles the one present in other homologues such as *Srs*AmpDh2, where it has been shown to act as a PG-fastening element (Martínez-Caballero *et al*, 2013). Interestingly, and despite differences in sequence, the length of this loop is conserved across other LysA-Ami2B domains that frequently exhibit an aromatic residue located at the middle of the loop, which emerges as a plausible feature participating in PG interaction (**Fig. 5C**).

3.3.2. *DS6A*Ami2B substrate binding can accommodate a tetrasaccharide-pentapeptide substrate

To investigate the molecular bases of substrate binding and hydrolysis, we determined the crystal structure of *DS6A*Ami2B in complex with the peptide L-Ala-D-*iso*-Gln-L-Lys-D-Ala-D-Ala (P5^{Lys} peptide), bearing a L-Lys residue instead of the mDAP found in the mycobacterial PG. The structure of *DS6A*Ami2B:P5^{Lys} was solved at 2.74 Å resolution, and it consists of three molecules of complex in the asymmetric unit exhibiting the same fold (RMSD of 0.139-0.176 Å for 165 Cα atoms), also conserved with respect to the native structure (RMSD of 0.190-0.230 Å for 175 Cα atoms) (**Fig.**

6 and **6SM**). The three molecules of P5^{Lys} exhibit a conserved pose. The latter is unambiguously defined by the electron density map, but for the Cδ, Cε and Nξ side-chain atoms of L-Lys and the C-terminal carboxylate group exhibiting none or poor density (**Fig. 6AB**). The *DS6A*Ami2B: P5^{Lys} structure reveals a peptide-binding groove shaped by loops α2-β3, β4-α3 and β6-α5 and up to five pockets, from *p*1 to *p*5, accommodating each amino acid of the ligand (**Fig. 6AB**). The *p*1 pocket is formed by two leucine (L270 and L274) side-chains and R265, which stabilizing the methyl group and the main chain carboxyl group of the L-Ala. While L270 is conserved in all LysAs inspected, L274 and R265 are frequently substituted by Ala. In the latter case, a His residue located downstream (residue aa$_{i+2}$) might play the role of R265. The *p*2 pocked interacts with the D-*iso*-Gln through an H-bond network involving W294 and H295 main-chains, and N310 and Q372 side-chains, with these latter exhibit high conservation across the LysAs analysed (**Fig. 5C**). The mycobacterial PG may contain a D-*iso*-Glu or a D-*iso*-Gln in the second amino acid of the peptide moiety, which exhibit amide or carboxylate groups at the Cα atom of the residue, respectively (Vollmer *et al*, 2008). Interestingly, while Q372 forms a bidentate H-bond contact with the amide group of the D-*iso*-Gln, a carboxylate group would only enable a single H-bond interaction. However, and in such a case, this lost contact might be replaced by another H-bond interaction with K374, whose side chain locates near enough to the D-*iso*-Gln in our structure (**Fig. 6M**). Although further studies will be required to validate this proposal, our analysis points at *DS6A*Ami2B and LysA-Ami2B homologues as amidase domains able to recognize D-*iso*-Gln and D-*iso*-Glu at *p*2 of the peptide moiety. At *p*3, the side and main chains of L-Lys are stabilized by hydrophobic contacts formed with L270 and W294, and H-bond interactions with N310, respectively, all conserved across LysA-Ami2B domains. The side chain of L-Lys does not stablish specific contacts with Ami2B, which aligns with the flexibility observed for atoms Cδ, Cε and N (**Fig. 6**). This is not surprising, as the mycobacterial PG presents mDAP at the third amino-acid of the peptide moiety, which bears a carboxylate group absent in the L-Lys side-chain. Interestingly, inspection of residues nearby *p*3 does reveal the presence of an arginine (R271) adjacent to *p*3 as possible interactor of the mDAP side chain. At the end of the peptide-binding groove, the channel widens at *p*4 and *p*5 pockets that accommodate the D-Ala-D-Ala moiety via H-bond contacts of the ligand backbone with the N307 main-chain and the highly conserved N308 side-chain. Comparison of *DS6S*Ami2B:P5^{Lys} complex with the native structure does not show significant changes in the residues involved in P5^{Lys} interaction, thus suggesting the peptide-binding to be pre-formed for PG recognition. Nevertheless, we must bear in mind the carboxylate group missing in our peptide due to the presence of L-Lys, which might affect potential rearrangements at loop α2-β3 the proposed interactions of R271 with the mDAP side chain.

We also attempted the resolution of a *DS6A*Ami2B crystal structure bound to saccharide fragments. This did not yield any structure, which is likely due to steric hindrance of the glycan-binding site resulting of contacts with another molecule of the crystal. Thus, and to analyse the glycan-binding groove, we superimposed the *DS6A*Ami2B:P5^{Lys} complex with the crystal structure of the amidase domain of the *Spn*LytA autolysin in complex with a PG fragment (*Spn*LytA:4S5P, PDB entry 5CTV) (Sandalova *et al*, 2016) (**Fig. 6C**). This evidences a large glycan-binding channel in *DS6A*Ami2B able to accommodate up a tetra-saccharide NAG-NAM-NAG-NAM occupying positions -1, 0, +1 and +2. This comparison also reveals significant differences and similitudes

Figure 6. (A) Cartoon representation of the *DS6A*Ami2B:P5^Lys complex (in orange) superimposed to the native *DS6A*Ami2B structure (in green). Detail of the 2Fo-Fc electron density map (blue mesh) contoured at sigma=1.0 and defining the P5^Lys molecule (black sticks) is shown. The zinc cation bound in both structures is represented as a yellow sphere. The loops forming the both glycan and peptide-binding channels and *p*1-*p*5 amino-acid binding pockets are labelled. (B) View of the residues participating in the interaction with the P5^Lys molecule in *DS6A*Ami2B:P5^Lys and native *DS6A*Ami2B structures, coloured in orange and green, respectively. P5^Lys and Zn(II) are represented as in (A). (C) Superimposition of *DS6A*Ami2B:P5^Lys (in orange) and *Spn*LytA:4S5P (in cyan, PDB entry 5CTV) with P5^Lys and 4S5P molecules represented as black and white sticks, respectively. (D) Detail of the active sites in *DS6A*Ami2B bound to phosphate molecule, *DS6A*Ami2B:P5^Lys zinc-interacting water molecule, and *Spn*LytA:4S5P. (E) Proposed catalytic cycle in *DS6A*Ami2B.

between both glycan-binding channels. Related to the former, *DS6A*Ami2B presents a tryptophan residue (W329) edging the channel at -1 position in (V148 in *Spn*LytA), thus locking the substrate-binding channel beyond the -1 position. The W329 conformation is stabilised through contacts stabilised with residues of a hydrophobic pocket, and through cation-pi contacts with the arginine of an adjacent *DS6A*Ami2B:P5^Lys complex and that is

additionally forming a salt-bridge interaction with a glutamate (E331) of the β5-α4 loop (**Fig. 6SM**). The location and contacts stabilising W329 side-chain opens an interrogation regarding its conformation and potential flexibility in the isolated monomer. This and the conservation observed for W329 across LysAs suggest this residue as plausible element regulating the opening of the PG-binding channel, which would enable interactions with the glycan

moiety beyond position -1 in LysA-Ami2B domains. This is an interesting posit that will require more studies to be validated. Other differences observed between *DS6A*Ami2B:P5^{Lys} and *Spn*LytA:4S5P complexes regards the α2 helix and the α2-β3 loop framing positions 0, +1 and +2. On the one hand, we observed narrower pockets in *DS6A*Ami2B at positions 0 and +1, where R267 would orient towards the NAM ring at position 0 to likely form favourable contacts. The R267 orientation would be safeguarded by restrictions imposed by a preceding proline and contacts stablished with the peptide moiety. On the other hand, a tyrosine residue found in other Amidase_2 members at position +2 (Y41 in *Spn*LytA) is missing in *DS6A*Ami2B and other LysA-Ami2B domains, thus resulting in a shallower +2 site. This seems to compensate the presence of a bulkier α2-β3 loop, thus resulting in enough room to accommodate the NAM and nascent peptide stem expected to bind at this position. Despite the differences observed, we identify three residues of *Spn*LytA (N30, E38 and R44) forming key contacts with the PG, and whose mutation has been shown to significantly decrease the catalytic activity of the enzyme (Table 4SM) (Mellroth *et al*, 2014). These residues are conserved across Amidase_2 domains, including *DS6A*Ami2B (N255, E258 and K265) and LysA-Ami2B inspected, thus pointing at a relevant role in PG binding by LysA-Ami2B amidases (**Fig. 5C**). Finally, the β5-α4 loop of *DS6A*Ami2B presents an aromatic residue (Y329), occupying a central position of the loop, that orients toward position -1. This residue is also found in other LysA-Ami2B domains, as tyrosine, phenylalanine and histidine (**Fig. 5C**). And in Amidase_2 homologues exhibiting a long β5-α4 loop such as *Srs*AmpDh2, where it has been shown to participate in substrate interaction at positions -1 and -2 (Martínez-Caballero *et al*, 2013), thus supporting the proposed PG-binding role of β5-α4 loop in LysA-Ami2B amidases (**Fig. 5C** and **6SM**).

To gain insights into *DS6A*Ami2B catalytic mechanism, we inspected the active site, in both native and peptide-bound structures, and compared it with the high-resolution structure of *Spn*LytA:S4P5 (**Fig. 6B**). This highlights, on the one hand, the zinc-interacting phosphate and water molecules laying adjacent the scissile bond in *DS6A*Ami2B native and complex structures, respectively. On the other hand, the phosphate lays adjacent the scissile bond in *DS6A*Ami2B and *Spn*LytA:S4P5 complexes mimicking the tetrahedral intermediate (TI) of the reaction to be formed upon the attack of w^{cat} onto the lactyl group of the PG. In this line, the water molecule adjacent the scissile bond (< 3 Å) nicely overlays with one of the oxygens of the phosphate, thus emerging as the potential catalytic water (w^{cat}) involved in catalysis. The oxyanion formed upon the catalytic attack is nicely mimicked by a second oxygen of the phosphate that locates at H-bond distance from the catalytic E317 side-chain, which supports E317 to function as an acid stabilising the TI. The TI formation would drag the peptide moiety towards the catalytic zinc, thus approaching the scissile bond towards His295 main-chain to a distance favourable for H-bond interactions, thus contributing to stabilise the TI state. In this configuration, the side chain of K374 would lay close enough to supply the proton necessary to form the amine group of the peptide product, thus favouring the final rupture of the substrate. While this catalytic mechanism partially aligns with the proposed for members of the Amidase_2 family (Zoll *et al*, 2010; Mellroth *et al*, 2014; Sandalova *et al*, 2016) it also features new posits regarding the roles played by E317 and K374 catalytic residues and assignment of the catalytic water in LysA-Ami2B and other Amidase_2 domains. Hypothesis that should be further investigated in order to unambiguously validate this mechanism.

3.4. Model of interaction of *DS6A*LysA catalytic domains with the mycobacterial cell-wall

We have determined the molecular envelop corresponding to the two catalytic domains of *DS6A*LysA in solution by SAXS and using the 6His-tagged and untagged versions of the *DS6A*GH19-Ami2B construct. Initially, we docked both crystallographic *DS6A*GH19 and *DS6A*Ami2B domains onto the molecular envelop calculated for the 6His-*DS6A*GH19-Ami2B construct. Using the protrusion resulting of the presence of the 6His-tag, we could unambiguously assign and dock each domain, which were oriented based on the location of the His-tag and the linker connecting GH19 and Ami2B domains (**Fig. 7SM**). The scattering curve calculated for this initial model shows a good fitting with the experimental data (**Fig. 7SM**). Thereafter, this initial model was docked onto the molecular envelop determined for the untagged *DS6A*GH19-Ami2B construct. *DS6A*GH19 and *DS6A*Ami2B domains were slightly readjusted in order optimise their fitting to the SAXS envelop, resulting in a model showing a good fitting with the experimental data (**Fig. 7** and **7SM**). Interestingly, examination of this final model shows that the PG-binding channels of both GH19 and Ami2B are facing the same side envelop, thus suggesting that both catalytic domains may orient towards a narrow region of the substrate to, concomitantly, proceed to its digestion (**Fig. 7**). To explore this hypothesis, we first superimposed the *D29*GH19^{E228Q}:NAG$_4$ and *DS6A*Ami2B:P5^{Lys} complexes onto the SAXS model. Next, we measured the distance between both glycan strands bound at *D29*GH19 and *DS6A*Ami2B PG-binding channels, which consisted of 40 Å approximately. Interestingly, the distance spanning glycan strands measured in a model of the PG molecule reported by NMR (Meroueh *et al*, 2006) falls very close to the one observed in our model, which suggest that both catalytic domains of *DS6A*LysA are likely to act cooperatively in the hydrolysis of the PG molecule *in vivo* (**Fig. 7**).

4. Conclusions

Our work gathers fundamental details of peptidoglycan hydrolysis by *D29*LysA and *DS6A*LysA mycobacteriophage endolysins with therapeutic relevance for enzybiotics development to combat mycobacterial infections. Starting with the crystallographic characterisation of *D29*N4, our study has enabled us to experimentally validate the dipeptidyl-peptidase function of this domain and narrow its membership to the C41 family of peptidases, which aligns with previously reported *in silico* and biochemical studies. Importantly, we provide with new details about the mechanism of PG binding and hydrolysis by *D29*N4, including the precise arrangement of the active site and the identification of the β1-β2 loop as a potential regulatory element in peptidoglycan interaction. Moreover, and based on biochemical studies reported elsewhere, we have mapped key residues for substrate interaction. Also, we have identified new residues potentially involved in the stabilisation of the mDAP moiety of substrate, as particular feature of this domain with implications in substrate selectivity. We also show conservation of these structural-functional features in other homologue LysA, thus shedding light into the function of this class of N4 peptidases.

Our study of *D29*GH19 and *DS6A*GH19 shows that LysA-GH19s domains are "short-loop" glycosyl-hydrolases belonging to the GH19_1 subfamily, and characterised by smaller substrate-binding channels when compared to their "loopful" chitinase homologues belonging to the GH19_2 subfamily. Our crystallographic characterisation of several *D29*GH19 complexes with glycan

Figure 7. (A) *Ab initio* determined SAXS envelop for the untagged *DS6A*GH19-Ami2B construct in solution by SAXS (in grey) where the domains GH19 (in blue) and Ami2B (in green) have been fitted. The N-terminus of the GH19 domain (in yellow) is highlighted as thicker ribbons. The C-terminus of the GH19 domain and the N-terminus of the Ami2B domain (coloured in magenta) are highlighted as thicker ribbons. The dashed lines in pink do presume the location of the inter-domain linker. (B) Plot showing the normalized pair-distance distribution function P(r) for *DS6A*GH19-Ami2B (a.u., arbitrary units). (C) Experimental scattering curve (dots) and theoretical scattering curve computed for the GH19-Ami2B model (smooth). (D) View of the model corresponding of DS6ALysA catalytic domains, resulting of the superimposition of D29GH19:NAG4 (in blue) and *DS6A*Ami2B:P5^{Lys} (in green) complexes onto the SAXS model. Both PG fragments are depicted as black sticks, with the distance between their glycan moieties is indicated and compared with the dimensions of a model corresponding to the Gram-negative PG molecule determined by NMR (on the right).

analogues has enabled us to capture snapshots of the catalytic process. The latter transits from an initial open state of the domain (*D29*GH19^{E228Q}:NAG$_4$), necessary for substrate binding but not suitable for hydrolysis, to a close state of the domain (*D29*GH19^{WT}:NAG:NAG$_2$ and *D29*GH19^{WT}:NAG) indispensable for correct substrate binding and hydrolysis, that we have captured as post-hydrolysis stages *in crystallo*. Our structural analysis evidences the significant inter-lobe rearrangement required for this transition that only occurs when the enzyme is catalytically ready. Moreover, our data supports that the M-lobe functions as a pre-formed structural platform for substrate interaction in the open state that adequately place the substrate towards its catalytic base. This crucial step will be followed by the closure of the PG-binding groove around the substrate to enable a correct approximation of the catalytic acid towards the glycosidic bond, in line with the GH19 family hydrolase mechanism with inversion of the configuration (Iseli et al, 1996). Moreover, a detailed analysis of our crystallographic and docking structures has led us to identify essential residues involved in PG binding and hydrolysis, as well as evidences supporting *D29*GH19 to operate as muramidase enzyme cleaving the glycosidic bond connecting NAM and NAG saccharides of the PG. These features extend to other LysA-GH19 domains thus unravelling key characteristics for this class of endolysin domains and for the whole subfamily of GH19_1 hydrolases.

We also report the crystal structures of the *DS6A*Ami2B domain, alone and in complex with an analogue of the product of the reaction. Our structural analysis evidence *DS6A*Ami2B as member of the Amidase_2 family, conserving the characteristic catalytic features described for these group of amidases. Our structures reveal central details of the mode of interaction of this enzyme with the peptide moiety of the PG, which takes place through a large cavity able to accommodate a substrate of up to four saccharide rings and five amino acids of the peptide stem. We also uncover particular features of the PG-binding channel of *DS6A*Ami2B with respect to other Amidase_2 homologues, such as: i) a rigid loop α2-β3 loop preforming *p*1, *p*2, 0 and +1 pockets; ii) the presence of a tryptophan locking the potion -2 of the glycan-binding groove, which emerges as a potential gate-keeper regulating accessibility of the PG-biding site; and iii) the presence of a long β5-α4 loop, potentially involved in interaction with the glycan moiety. Moreover, a phosphate molecule mimicking the tetrahedral intermediate state of the reaction in the native structure has let us to explain potential interactions at the active site during catalysis, allowing us to propose a model of the catalytic mechanism for *DS6A*Ami2B and other LysA-Ami2B. In this study, we have further explored the arrangement of *DS6A*LysA catalytic domains in solution using SAXS. In order to full-proof the relative orientation of both muramidase and amidase domains, we characterised the enzyme in-solution by SAXS with and without the presence of the His-tag.

The results permitted to understand the manner and orientation adopted by both domains with respect to each other. Our SAXS model supports both PG-binding sites to be located at the same side of the molecular envelop, thus supporting both catalytic activities to functions together, explaining a sequential model of digestion of the cell-wall. Altogether, the present study constitutes a solid groundwork for future investigations on this relevant class of endolysins, including the rational design of LysA lysins with optimised enzybiotic activity to treat mycobacterial IDs and to combat emergent antimicrobial resistances associated to this class of pathogens.

5. References

Adams PD, Afonine P V., Bunkóczi G, Chen VB, Davis IW, Echols N, Headd JJ, Hung LW, Kapral GJ, Grosse-Kunstleve RW, *et al* (2010) PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta Cryst D* 66: 213–221

Alderwick LJ, Harrison J, Lloyd GS & Birch HL (2015) The mycobacterial cell wall—peptidoglycan and arabinogalactan. *Cold Spring Harb Perspect Med* 5: 1–16

Anantharaman V & Aravind L (2003) Evolutionary history, structural features and biochemical diversity of the NlpC/P60 superfamily of enzymes

Bartlett HP, Dawson CC, Glickman CM, Osborn DW, Evans CR, Garcia BJ, Frost LC, Cummings JE, Whittel N, Slayden RA, *et al* (2024) Targeting intracellular nontuberculous mycobacteria and M. tuberculosis with a bactericidal enzymatic cocktail . *Microbiol Spectr* 12

Batt SM, Burke CE, Moorey AR & Besra GS (2020) Antibiotics and resistance: the two-sided coin of the mycobacterial cell wall. *Cell Surface* 6 doi:10.1016/j.tcsw.2020.100044 [PREPRINT]

Böth D, Schneider G & Schnell R (2011) Peptidoglycan Remodeling in Mycobacterium tuberculosis: Comparison of Structures and Catalytic Activities of RipA and RipB. *J Mol Biol* 413: 247–260

Bricogne G., Blanc E., Brandl M., Flensburg C., Keller P., Paciorek W., Roversi P, Sharff A., Smart O.S., Vonrhein C., *et al* (2017) Welcome to Global Phasing Limited. *BUSTER v2112*

Buruli ulcer

Büttner FM, Renner-Schneck M & Stehle T (2015) X-ray crystallography and its impact on understanding bacterial cell wall remodeling processes. *International Journal of Medical Microbiology* 305: 209–216

Büttner FM, Zoll S, Nega M, Götz F & Stehle T (2014) Structure-function analysis of Staphylococcus aureus amidase reveals the determinants of peptidoglycan recognition and cleavage. *Journal of Biological Chemistry* 289: 11083–11094

Catalão MJ, Filipe SR & Pimentel M (2019) Revisiting anti-tuberculosis therapeutic strategies that target the peptidoglycan structure and synthesis. *Front Microbiol* 10

CAZy - GH19

Chen VB, Arendall WB, Headd JJ, Keedy DA, Immormino RM, Kapral GJ, Murray LW, Richardson JS & Richardson DC (2009) MolProbity: all-atom structure validation for macromolecular crystallography. *Acta Cryst D* 66: 12–21

Cowieson NP, Edwards-Gayle CJC, Inoue K, Khunti NS, Doutch J, Williams E, Daniels S, Preece G, Krumpa NA, Sutter JP, *et al* (2020) Beamline B21: High-throughput small-angle X-ray scattering at Diamond Light Source. *J Synchrotron Radiat* 27: 1438–1446

Dams D & Briers Y (2019) Enzybiotics: Enzyme-Based Antibacterials as Therapeutics. *Adv Exp Med Biol*: 1148:233–253

Emsley P, Lohkamp B, Scott WG & Cowtan K (2010) Features and development of Coot. *Acta Cryst D* 66: 486–501

Eniyan K, Sinha A, Ahmad S & Bajpai U (2020) Functional characterization of the endolysins derived from mycobacteriophage PDRPxv. *World J Microbiol Biotechnol* 36

Evans PR & Murshudov GN (2013) How good are my data and what is the resolution? *Acta Cryst D* 69: 1204–1214

Evans R, O'Neill M, Pritzel A, Antropova N, Senior A, Green T, Žídek A, Bates R, Blackwell S, Yim J, *et al* (2022) Protein complex prediction with AlphaFold-Multimer. *bioRxiv*: 2021.10.04.463034

Froman S;, Drake W, Will E & Bogen FAPHA (1954) Bacteriophage Active Against Virulent Mycobacterium tuberculosis 1. Isolation and Activity

Gangakhedkar R & Jain V (2024) Construing the function of N-terminal domain of D29 mycobacteriophage LysA endolysin in phage lytic efficiency and proliferation. *Mol Microbiol* 122: 243–254

Global Leprosy (Hansen's disease) Strategy 2021-2030 | 2 (2021)

Global Tuberculosis Report 2023 (2023) World Health Organization

Griffin ME, Klupt S, Espinosa J & Hang HC (2023) Peptidoglycan NlpC/P60 peptidases in bacterial physiology and host interactions. *Cell Chem Biol* 30: 436–456 doi:10.1016/j.chembiol.2022.11.001 [PREPRINT]

Haddad Kashani H, Schmelcher M, Sabzalipoor H, Seyed Hosseini E & Moniri R (2018) Recombinant endolysins as potential therapeutics against antibiotic-resistant staphylococcus aureus: Current status of research and novel delivery strategies. *Clin Microbiol Rev* 31

Hoell IA, Dalhus B, Heggset EB, Aspmo SI & Eijsink VGH (2006) Crystal structure and enzymatic properties of a bacterial family 19 chitinase reveal differences from plant enzymes. *FEBS Journal* 273: 4889–4900

Holm L, Laiho A, Törönen P & Salgado M (2023) DALI shines a light on remote homologs: One hundred discoveries. *Protein Science* 32

Iseli B, Armand S, Boller T, Neuhaus JM & Henrissat B (1996) Plant chitinases use two different hydrolytic mechanisms. *FEBS Lett* 382: 186–188

Johansen MD, Herrmann JL & Kremer L (2020) Non-tuberculous mycobacteria and the rise of Mycobacterium abscessus. *Nature Reviews Microbiology 2020 18:7* 18: 392–407

Joshi H, Nair G, Gangakhedkar R & Jain V (2019) Understanding the role of the lysozyme-like domain of D29 mycobacteriophage-encoded endolysin in host cell lysis and phage propagation. *Microbiology (United Kingdom)* 165: 1013–1023

Joshi H, Seniya SP, Suryanarayanan V, Patidar ND, Singh SK & Jain V (2017) Dissecting the structure–function relationship in lysozyme domain of mycobacteriophage D29-encoded peptidoglycan hydrolase. *FEBS Lett* 591: 3276–3287

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Žídek A, Potapenko A, *et al* (2021) Highly accurate protein structure prediction with AlphaFold. *Nature 2021 596:7873* 596: 583–589

Kabsch W (2010) *XDS. Acta Cryst D* 66: 125–132

Kerff F, Petrella S, Mercier F, Sauvage E, Herman R, Pennartz A, Zervosen A, Luxen A, Frère JM, Joris B, *et al* (2010) Specific Structural Features of the N-Acetylmuramoyl-l-Alanine Amidase AmiD from Escherichia coli and Mechanistic Implications for Enzymes of This Family. *J Mol Biol* 397: 249–259

Leprosy

Loeffler JM, Djurkovic S & Fischetti VA (2003) Phage Lytic Enzyme Cpl-1 as a Novel Antimicrobial for Pneumococcal Bacteremia. *Infect Immun* 71: 6199–6204

MacRae CF, Sovago I, Cottrell SJ, Galek PTA, McCabe P, Pidcock E, Platings M, Shields GP, Stevens JS, Towler M, *et al* (2020) Mercury 4.0: from visualization to analysis, design and prediction. *urn:issn:1600-5767* 53: 226–235

Martínez-Caballero S, Lee M, Artola-Recolons C, Carrasco-López C, Hesek D, Spink E, Lastochkin E, Zhang W, Hellman LM, Boggess B, *et al* (2013) Reaction products and the X-ray structure of AmpDh2, a

virulence determinant of Pseudomonas aeruginosa. *J Am Chem Soc* 135: 10318–10321

Matsuhashi M. (1966) [Biosynthesis in the bacterial cell wall] - PubMed. *Tanpakushitsu Kakusan Koso*: 875–886

McCoy AJ, Grosse-Kunstleve RW, Adams PD, Winn MD, Storoni LC & Read RJ (2007) Phaser crystallographic software. *J Appl Crystallogr* 40: 658–674

Meroueh SO, Bencze KZ, Hesek D, Lee M, Fisher JF, Stemmler TL & Mobashery S (2006) Three-dimensional structure of the bacterial cell wall peptidoglycan PNAS

Nelson D, Loomis L & Fischetti VA (2001) Prevention and elimination of upper respiratory colonization of mice by group A streptococci by using a bacteriophage lytic enzyme

Ozhelvaci F & Steczkiewicz K (2023) Identification and classification of papain-like cysteine proteinases. *Journal of Biological Chemistry* 299: 104801

Park Y, Lim JA, Kong M, Ryu S & Rhee S (2014) Structure of bacteriophage SPN1S endolysin reveals an unusual two-module fold for the peptidoglycan lytic and binding activity. *Mol Microbiol* 92: 316–325

Payne KM & Hatfull GF (2012) Mycobacteriophage Endolysins: Diverse and Modular Enzymes with Multiple Catalytic Activities. *PLoS One* 7: e34052

Raymond JB, Mahapatra S, Crick DC & Pavelka MS (2005) Identification of the namH gene, encoding the hydroxylase responsible for the N-glycolylation of the mycobacterial peptidoglycan. *Journal of Biological Chemistry* 280: 326–333

Saadhali SA, Hassan S, Elizabeth Hanna L, Devi Ranganathan U & Kumar V (2016) Homology modeling, substrate docking, and molecular simulation studies of mycobacteriophage Che12 lysin A. *J Mol Model*

Sandalova T, Lee M, Henriques-Normark B, Hesek D, Mobashery S, Mellroth P & Achour A (2016) The crystal structure of the major pneumococcal autolysin LytA in complex with a large peptidoglycan fragment reveals the pivotal role of glycans for lytic activity. *Mol Microbiol* 101: 954–967

Schneidman-Duhovny D, Hammel M, Tainer JA & Sali A (2016) FoXS, FoXSDock and MultiFoXS: Single-state and multi-state structural modeling of proteins and their complexes based on SAXS profiles. *Nucleic Acids Res* 44: W424–W429

Schuch R, Nelson D & Fischetti VA (2002) A bacteriolytic agent that detects and kills Bacillus anthracis. *Nature 2002 418:6900* 418: 884–889

Sekiya H, Tamai E, Kawasaki J, Murakami K & Kamitori S (2021) Structural and biochemical characterizations of the novel autolysin Acd24020 from Clostridioides difficile and its full-function catalytic domain as a lytic enzyme. *Mol Microbiol* 115: 684–698

Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M, Söding J, *et al* (2011) Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol Syst Biol* 7: 539

Squeglia F, Ruggiero A, Romano M, Vitagliano L & Berisio R (2014) Mutational and structural study of RipA, a key enzyme in Mycobacterium tuberculosis cell division: Evidence for the l-to-d inversion of configuration of the catalytic cysteine. *Acta Crystallogr D Biol Crystallogr* 70: 2295–2300

Tuberculosis

Urdániz E, Martín M, Payaslián F, Defelipe LA, Dodes M, Martinez M, Alzari PM, Cabrera G, Martí MA & Piuri M (2022) Gp29 LysA of mycobacteriophage TM4 can hydrolyze peptidoglycan through an N-acetyl-muramoyl-L-alanine amidase activity. *Biochim Biophys Acta Proteins Proteom* 1870

Verdonk ML, Cole JC, Hartshorn MJ, Murray CW & Taylor RD (2003) Improved protein–ligand docking using GOLD. *Proteins: Structure, Function, and Bioinformatics* 52: 609–623

Vollmer W, Blanot D & De Pedro MA (2008) Peptidoglycan structure and architecture. *FEMS Microbiol Rev* 32: 149–167 doi:10.1111/j.1574-6976.2007.00094.x [PREPRINT]

Vollmer W & Höltje JV (2004) The architecture of the murein (peptidoglycan) in gram-negative bacteria: Vertical scaffold or horizontal layer(s)? *J Bacteriol* 186: 5978–5987

W.B. Redmond & J. C. Carter (1960) A Bacteriophage Specific for Mycobacterium Tuberculosis, Varieties Hominis and Bovis | American Review of Respiratory Disease. *American Review of Respiratory Disease*: 82(6).781-786

Xu Q, Abdubek P, Astakhova T, Axelrod HL, Bakolitsa C, Cai X, Carlton D, Chen C, Chiu HJ, Chiu M, *et al* (2010) Structure of the γ-d-glutamyl-l-diamino acid endopeptidase YkfC from Bacillus cereus in complex with l-Ala-γ-d-Glu: insights into substrate recognition by NlpC/P60 cysteine peptidases. *urn:issn:1744-3091* 66: 1354–1364

Yang F, Labani-Motlagh A, Bohorquez JA, Moreira JD, Ansari D, Patel S, Spagnolo F, Florence J, Vankayalapati A, Sakai T, *et al* (2024) Bacteriophage therapy for the treatment of Mycobacterium tuberculosis infections in humanized mice. *Commun Biol* 7

Acknowledges

This work was supported by a Atracción de Talento program (Modalidad 1) grant from the Madrid Regional Government (Spain). The authors would like to also thank Prof. Martín Martínez-Ripoll and Prof. Juan A. Hermoso for their precious discussion about this work and manuscript revision; to Prof. Sharhiar Mobashery for kindly sharing with us the coordinate file corresponding to the PG structural-model; and to Dr. Beatriz González as MB-BAG coordinator, for facilitating us with access to MX beamtime at the European Synchrotron Radiation Facility (ESRF). We would like to thank the ESRF (beamline ID30A-3, proposals mx33176 and mx35926), Diamond Light Source (beamline B21, proposals mx33176 and mx35926) and the ALBA Synchrotron (XALOC-BL13 beamline proposals 2021085244 and 2022097082) for beamtime and their staff for assistance during data collection.

Author contribution

I.P.D. conceived the project and supervised the experiments. F.C.Z. performed the expression, purification, structural characterization of (PDB entries of structures 9HNV, 9HNU, 9HP7, 9HNA and 9HTY). L.G.L. performed the expression, purification, structure characterization of (PDB entries of structures 9HYR, 9HQW, 9HR1, 9HRM, 9HV2 and 9HV0). F.C.Z. carried out the final validation and structure deposition of all the crystallographic structures in the Protein Data Bank. I.P.D. carried out the analysis of the crystallographic structures with the contribution of F.C.Z. and L.G.L. I.G.M. performed the SAXS studies, including data analysis, figure preparation and data deposition. L.I.S. carried out the computational docking studies and model analysis. I.P.D. wrote the manuscript and prepared the figures with involvement of all the other authors.

Figures Supplemental Material

Figure 2SM. Normalised representation of the B-factor values observed in chains A (left) and B (right) of the crystal structure of *D29*N4, which range from colour blue (lowest B-factor values) to red (highest B-factor values). The catalytic triad (represented as sticks) and the β1-β2 loop are indicated with a labels.

Figure 3SM. Detail of the EF-hand-like calcium-binding site of *D29*GH19. Electron density peaks corresponding to the calcium cation identified at both non anomalous and anomalous 2Fo-Fc maps are depicted as a mesh.

Figure 4SM. (A) Superimposition of *D29*GH19^WT and *D29*GH19^E288Q represented as cartoons in yellow and cyan, respectively, depicting the catalytic residues as sticks. (B) Detail of the hydrophobic core involved in NC-lobe pivoting towards α7 between *D29*GH19^E228Q:NAG4 (in green) and *D29*GH19^WT:NAG:NAG2 (in pink). (C) Superimposition the of crystal structures of *Rsc*-C, in the absence (PDB entry 4DWX) and bound to a NAG4 (PDB entry 4J0L), illustrating the closure of the substrate-binding cavity upon ligand binding in a "loop-full" GH19 domain. (D) and (E) detail of the 2Fo-Fc electron density map (blue mesh) corresponding to the NAG:NAG2 and NAG molecules bound in complexes *D29*GH19^WT:NAG

(white) and $D29$GH19^{WT}:NAG:NAG$_2$ (grey), respectively, countered at sigma = 1.0. (F) Superimposition of $D29$GH19^{E228Q}:NAG$_4$ (protein in green, NAG$_4$ in black) and $DS6A$GH19 (white) structures showing residues involved in substrate interaction in both domains as sticks. (G) Detail of the active site in $D29$GH19^{WT-}^{I}, $D29$GH19^{E228Q}:NAG$_4$ and $D29$GH19^{WT}:NAG:NAG$_2$ superimposed structures. Key residues involved in catalysis and NAG$_n$ fragments are represented as sticks following the same colour code as in (A). The catalytic water (CW) molecule activated by E237 is represented as spheres coloured in yellow (in $D29$GH19^{WT-}^{I}), green (in $D29$GH19^{E228Q}:NAG$_4$) and pink (in $D29$GH19^{WT}:NAG:NAG$_2$). (H) Representation of the eleven top solutions obtained in the docking of 4S2P PG-fragment (lines coloured in grey) onto $D29$GH19^{WT}. (I) Superimposition of the top docking solution calculated for $D29$GH19^{WT}:4S2P complex (protein in yellow, ligand in blue) with ligands bound to the crystallographic complexes $D29$GH19^{E228Q}:NAG$_4$ (black sticks) and $D29$GH19^{WT}:NAG:NAG$_2$ (white sticks).

Figure 6SM. (A) Superimposition of the three complexes showing the peptide pose.

Figure 7SM. (A) *Ab initio* molecular envelope determined by SAXS for a *DS6A*Ami2B construct containing a N-terminal 6His-tag. (B) Normalized pair-distance distribution function P(r) for the protein without the his-tag (green graph). (C) Experimental scattering curve (lime green dots) and theoretical scattering curve, in emerald green, computed for the model (smooth), where the protein adopts a conformation that could be explained by the model resulting from joining the PDBs 9HQW and 9HNA, using the program Coot. The data shows an extra volume that clarifies where the N-terminal domain of *DS6A*GH19 is located with respect to the *DS6A*Ami2B domain. (D) *Ab initio* molecular envelope determined by SAXS for the untagged DS6AAmi2B construct.

Table 1SM. X-ray data collection and refinement statistics.

	$D29N4^{WT}$ native	$D29GH19^{WT}$ native (Form I)	$D29GH19^{WT}$ native (Form II)	$D29GH19^{WT}$ native (Form III)	$D29GH19^{WT}$ with NAG2/NAG (Form IV)	$D29GH19^{WT}$ with NAG (Form III)	$DS6AGH19^{WT}$ native	$D29GH19^{E228Q}$ native (Form V)	$D29GH19^{E228Q}$ with NAG4 (Form I)	$DS6AAmi2B^{WT}$ native	$DS6AAmi2B^{WT}$ with P5^{Lys}
Data collection statistics											
PDB entry code	9HYR	9HQW	9HR1	9HRM	9HV2	9HV0	9HNV	9HNU	9HP7	9HNA	9HTY
Space group	$P2_12_12_1$	$C2$	$P2_12_12_1$	$P2_12_12_1$	$P2_12_12$	$P2_12_12_1$	$I222$	$I2$	$C2$	$C2$	$C2$
No. monomers in ASU	2	1	2	2	4	2	3	1	1	3	3
a, b, c (Å)	65.2, 67.0, 83.0	91.3, 44.8, 69.7	44.6, 89.0, 113.4	37.0, 57.0, 151.3	78.7, 84.2, 135.3	37.2, 56.5, 152.6	89.2, 145.9, 147.7	70.1, 44.9, 76.6	91.1, 44.8, 69.5	72.0, 121.0, 80.7	70.7, 121.2, 80.0
α, β, γ (°)	90.0, 90.0, 90.0	90.0, 126.1, 90.0	90.0, 90.0, 90.0	90.0, 90.0, 90.0	90.0, 90.0, 90.0	90.0, 90.0, 90.0	90.0, 90.0, 90.0	90.0, 103.0, 90.0	90.0, 126.4, 90.0	90.0, 107.2, 90.0	90.0, 106.3, 90.0
Wavelength (Å)	0.9793	0.9793	0.9793	0.9677	0.9677	0.9677	0.9792	0.9792	0.9792	0.9793	0.9792
Beamline	XALOC	XALOC	XALOC	ID30A-3	ID30A-3	ID30A-3	XALOC	XALOC	XALOC	XALOC	XALOC
Resolution range (Å)	82.97-3.20 (3.42-3.20)	56.27-1.63 (1.66-1.63)	113.39-2.00 (2.05-2.00)	57.04-2.02 (2.10-2.04)	45.81-3.10 (3.31-3.10)	76.31-2.43 (2.52-2.43)	76.34-2.75 (2.88-2.75)	45.56-1.45 (1.47-1.45)	38.25-1.83 (1.86-1.83)	77.14-2.04 (2.07-2.04)	60.62-2.74 (2.79-2.74)
R_{merge}	0.321 (0.701)	0.059 (0.597)	0.117 (0.895)	0.220 (1.041)	0.271 (1.448)	0.174 (0.891)	0.344 (2.156)	0.047 (0.312)	0.073 (0.758)	0.134 (1.240)	0.112 (0.415)
R_{pim}	0.173 (0.376)	0.036 (0.388)	0.033 (0.246)	0.160 (0.754)	0.136 (0.710)	0.097 (0.486)	0.128 (0.799)	0.021 (0.158)	0.031 (0.346)	0.055 (0.497)	0.072 (0.258)
No. total reflections	27415 (4952)	101079 (4353)	434589 (33922)	81875 (6595)	73573 (13776)	47708 (5232)	210189 (27695)	236632 (9170)	126584 (5698)	283455 (14979)	56465 (2970)
No. unique reflections	6267 (1110)	27799 (1355)	30434 (2255)	20836 (1595)	16002 (2889)	12304 (1286)	25458 (3356)	41013 (1973)	19241 (991)	41716 (2098)	16754 (845)
Mean ([I]/SD[I])	4.7 (2.4)	14.2 (2.2)	17.2 (3.5)	5.5 (1.8)	5.2 (1.2)	6.1 (1.8)	6.7 (1.2)	16.8 (3.2)	15.1 (2.2)	10.2 (2.1)	5.6 (2.3)
$CC_{1/2}$	0.934 (0.728)	0.998 (0.816)	0.999 (0.880)	0.979 (0.514)	0.985 (0.523)	0.983 (0.516)	0.984 (0.512)	0.999 (0.970)	0.999 (0.872)	0.994 (0.666)	0.991 (0.864)
Completeness (%)	99.1 (99.7)	98.4 (97.6)	96.9 (100.0)	98.5 (99.6)	95.2 (96.4)	96.6 (97.9)	100.0 (100.0)	99.3 (97.6)	95.1 (99.9)	98.7 (97.9)	98.0 (99.8)
Multiplicity	4.4 (4.5)	3.6 (3.2)	14.3 (15.0)	3.9 (4.1)	4.6 (4.8)	3.9 (4.1)	8.3 (8.3)	5.8 (4.6)	6.6 (5.7)	6.8 (7.1)	3.4 (3.5)
Refinement statistics											
Resolution range (Å)	51.25-3.20	45.49-1.64	70.04-2.00	53.37-2.04	43.81-3.10	76.31-2.43	76.34-2.75	45.56-1.45	38.25-1.83	77.14-2.04	60.62-2.74
No. reflections		27776	30368	20789	15962	12264	25438	40948	19225	41701	16749
R_{work} (%)	0.25	0.18	0.17	0.18	0.24	0.21	0.24	0.15	0.18	0.18	0.16
R_{free} (%)	0.31	0.21	0.20	0.24	0.28	0.26	0.28	0.18	0.20	0.21	0.20
No. of atoms	2714	1763	3463	2922	6195	2740	4630	1770	1688	4663	4515
Protein	2714	1515	3064	2681	6007	2682	4535	1542	1519	4368	4326
Ion	–	1	2	–	4	–	–	1	1	28	18
PG analogue	–	–	–	–	180	15	–	–	57	–	102
Water	–	227	355	241	8	43	95	213	111	267	69
Other (MES, PEG, MPD, Ala, D-Ser)	–	12/_/8/_	24/12/_/6	–	–	–	–	14	–	–	–
B factors (Å^2)											
Protein	34.5	23.9	28.4	21.5	62.0	32.2	44.0	24.4	30.87	36.81	34.35
Other (MES, PEG, MPD, Ala, D-Ser)	–	21.8/_/34.5/_	34.4/38.1/_/35.1	–	–	–	–	46.6	–	–	–
PG analogue	–	–	–	–	64.1	29.0	–	–	43.86	–	43.43
Ion	–	16.1	24.4	–	71.9	–	–	17.33	21.73	28.02	27.48
Water	–	34.4	36.9	29.2	59.6	32.2	36.41	34.78	36.68	43.78	34.96
R.m.s. deviations											
Bond lengths (Å)	0.002	0.005	0.006	0.006	0.002	0.003	0.003	0.011	0.004	0.005	0.005
Angle lengths (°)	0.545	0.750	0.735	0.776	0.446	0.666	0.614	1.103	0.659	0.788	0.741

Table 2SM. Mutants reported in *D29*N4 and homologue domains

Peptidase domain	Mutant reported	Equivalent residue in *D29*LysA	Effect in PG-hydrolase activity	Impact of the mutation	Reference
Mycobacterium tuberculosis RipA peptidase	Cys383A	C41	Inactive	Active site	Both et al. 2011
Clostridium difficile Acd24020 autolysin	C229A	C41	Inactive	Active site	Sekiya et al. 2021
	H354A	H112	Inactive	Active site	
	Y287A	Y25A	Inactive	PG-binding	
	W289A	Y27A	Activity decreased	PG-binding	
	D298A	D40	Activity decreased	Active site architecture	
	R317A	R82	Inactive	PG-binding	
	S300A	S42	Activity decreased	Active site architecture	
	V318A	S65	Activity decreased	PG-binding	
	S351Y	R109	Activity decreased	PG-binding	
	S353A	S111	Activity decreased	PG-binding	
Bacteriophage *D29* endolysin	Y25A	Y25	Inactive	PG-binding	Gangakhedkar & Lain 2024
	C41A	C41	Inactive	Active site	
	S111A	S111	Inactive	Active site architecture	
	H112A	H112	Inactive	Active site	
	E137A	E137	Inactive	Active site	

Table 3SM. Mutants reported in *D29*GH19 and homologue domains

Glycosyl hydrolase 19	Mutant reported	Equivalent residue in *D29*LysA	Effect in PG-hydrolase activity	Impact of the mutation	Reference
Streptomyces coelicolor Chitinase G	E68Q	E228	Inactive	Active site	Hoel et al. 2006
	E77Q	E237	Inactive	Active site	
Bacteriophage *SPN1S* endolysin	E49Q	E228	Inactive	Active site	Park et al. 2014
	E58Q	E237	Inactive	Active site	
	N59A	Y238	Decreased	PG-binding	
	Q128A	M270	Decreased	PG-binding	
	T130A	T272	Decreased	Active site architecture	
	N134A	N276	Decreased	PG-binding	
	F166A	Y316	Decreased	PG-binding	
	K170A	R321	Decreased	PG-binding	
	N187A	N341	Decreased	PG-binding	
Bacteriophage *D29* endolysin	E228Q	E228	Inactive	Active site	This study
	T345A	T345	Decreased	PG-binding	Joshi et al. 2019
	N346A	N346	Decreased	PG-binding	
	L347A	L347	Decreased	PG-binding	
	R350A	R350	Inactive	Active site architecture	

Table 4SM. Mutants reported in *DS6A*Ami2B homologue domains

Amidase 2B	Mutant reported	Equivalent residue in *DS6A*LysA	Effect in PG-hydrolase activity	Impact of the mutation	Reference
Bacteriophage *TM4* LysA endolysin	H226S	H251	Inactive	Active site	Urdáiz et al. 2022
	H335S	H335	Inactive	Active site	
	D347N	D376	Inactive	Active site	
	E290Q	E317	Inactive	Active site	
Streptococcus pneumoniae LytA autolysin	His26A	H251	Inactive	Active site	Mellroth et al. 2014
	His133A	H335	Inactive	Active site	
	D149A	D376	Inactive	Active site	
	E87A	E317	Inactive	Active site	
	H147A	K374	Decreased	Active site	
	N30L/Q	N255	Decreased	PG-binding	
	R44E/Q	K265	Decreased	PG-binding	

Table 5SM. SAS sample details, data collection, analysis, and 3D modelling details for biomolecules in solution.

Scattering particle composition		
Protein	Untagged *DS6A*GH19-Ami2B	His-tagged *DS6A*GH19-Ami2B
Sample environment/configuration		
Solvent composition	20 mM Tris pH 8.0, 150 mM NaCl, 1 mM DTT	
Sample temperature (°C)	4	
In beam sample cell (°C)	18	
Size Exclusion Chromatography SEC-SAS		
Sample injection concentration, (mg/ml)	5.9	5.6
Sample injection volume, mL	45	
SEC column type	SRT-C SEC-300 (Sepax)	Shodex KW-403
SEC flowrate, mL/min	0,16	0.16
Data acquisition/reduction software	GDA/Scåtter	GDA/Scåtter
Source/instrument description or reference	B21, Diamond Light Source	B21, Diamond Light Source
Measured q-range ($q_{min} - q_{max}$; Å^{-1})	0.0031-0.37	0.0031-0.37
Exposure time (s) and final number of sample frames	3, 600	3, 600
Relevant details		
Methods/Software	PRIMUS, AUTORG, GNOM	
Guinier Analysis		
I (0) ± σ (cm^{-1})	0,03900	0,03400
R_g ± σ (Å)	28,45±0,04	28,43±0,06
Data point range	80-323	80-315
PDDF/P(r) analysis		
I (0) ± σ (cm^{-1})	0,03921	0,03397
R_g ± σ (Å)	27,89±0,04	29,15±0,03
d_{max} (Å)	95	103
Data point range	54-2164	28-1600
q-range (Å^{-1})	0,014-0,2866	0,0080-0,2132
Methods/Software		
Volume estimates		
Porod volume, V_p (Å^3)	500,900	621,000
Molecular weight (M) estimates (kDa)		
From chemical composition	43.5	45,4
From SAS, concentration independent method (MoW)	39,8	49,2
From SAS-independent measure (MALLS)	**XXXX**	**XXXX**
Shape modelling method		
Software	DAMMIN, DAMMIF, FoXS	
q-range for fit ($q_{min} - q_{max}$; Å^{-1})	0,0045-0,3398	
Symmetry/anisotropy assumptions	P1	P1
Number of individual model reconstructions	20	20
χ^2, CorMap P-values for fit	1,44, 1,03	0,64, 1.05
Atomistic modelling methods		
Software	FoXS	
q-range for fit ($q_{min} - q_{max}$; Å^{-1})	0,0045-0,3398	0,0045-0,3398
Symmetry/anisotropy assumptions	N/A	N/A
Number of individual model reconstructions	1 (Based on PDBs: 9HNA y 9HNV)	1 (Based on PDBs: 9HNA y 9HNV)
χ^2, CorMap P-values for fit	0,90, 1,04	1,24, 1,02
Data and model deposition		
SASBDB IDs	**SASDXXX ID 6560**	**SASDXXX ID 6561**

References

- Böth D, Schneider G & Schnell R (2011). J. Mol. Biol. 413: 247–260

- Sekiya H, Tamai E, Kawasaki J, Murakami K & Kamitori S (2021). Mol. Microbiol. 115: 684–698

- Gangakhedkar R & Jain V (2024). Mol. Microbiol. 122: 243–254

- Hoell IA, Dalhus B, Heggset EB, Aspmo SI & Eijsink VGH (2006). FEBS J. 273: 4889–4900

- Park Y, Lim JA, Kong M, Ryu S & Rhee S (2014). Mol. Microbiol. 92: 316–325

- Joshi H, Nair G, Gangakhedkar R & Jain V (2019). Microbiology (United Kingdom) 165: 1013–1023

- Urdániz E, Martín M, Payaslián F, Defelipe LA, Dodes M, Martinez M, Alzari PM, Cabrera G, Martí MA & Piuri M (2022). Biochim. Biophys. Acta Proteins Proteom. 1870(2): 140745

- Mellroth P, Sandalova T, Kikhney A, Vilaplana F, Hesel D, Lee M, Mobashery S, Normark S, Svergun D, Henriques-Normark B, et al (2014). mBio 5(1): e01120-13

- Hajizadeh NR, Franke D, Jeffries CM & Svergun DI (2018). Scientific Reports 8, 7204.

- Piiadov V, Ares de Araujo E, Oliveira Neto M, Craievich AF & Polikarpov I. (2019). Protein Science 28, 454-463.

- Rambo RP & Tainer JA (2013). Nature 496, 477-481.

- Trewhella J, Duff AP, Durand D, Gabel F, Guss JM, Hendrickson WA, Hura GL, Jacques DA, Kirby NM, Kwan AH, Perez J, Pollack L, Ryan TM, Sali A, Schneidman-Duhovny D, Schwede T, Svergun DI, Sugiyama M, Tainer JA, Vachette P, Westbrook J. & Whitten AE (2017). Acta Crystallogr. Sect. D, Structural Biology 73, 710-728.