Spatially explicit capture–recapture models for relative density

Murray G. Efford[1]*

[1]Department of Mathematics and Statistics, University of Otago, P.O. Box 56, Dunedin 9054,

New Zealand.

* E-mail: murray.efford@otago.ac.nz

Open research statement: Data, R code and simulation results (Efford, 2025c,d) have been

archived on Zenodo at https://doi.org/10.5281/zenodo.17127944 and

https://doi.org/10.5281/zenodo.17402993.

Keywords: conditional likelihood; density surfaces; individual covariates; *Oligosoma*

infrapunctatum; population density; relative density; spatial capture–recapture; trend analysis;

Trichosurus vulpecula

Abstract

Spatially explicit capture–recapture (SECR) methods are used widely to estimate animal population density and related parameters. Maximum likelihood has been applied to two flavors of SECR model - a full model that includes absolute density as a spatially varying parameter, and a model conditional on the number of detected individuals that assumes uniform density. We show how the conditional model may be extended to estimate relative density as a function of habitat and other spatial variables. Unlike absolute density, coefficients of relative density are robust to bias from heterogeneous individual detection. The conditional model has the further advantage of seamlessly including individual covariates of detection, and it is faster to fit than the full model. The spatial absolute density surface may be derived from the conditional model fit. Population growth rate (relative density with respect to time) may also be estimated from a conditional model, given temporal data. These properties are demonstrated by examples and simulation. It is suggested that conditional-likelihood models, including those for relative density, play a central role in SECR.

Introduction

Capture–recapture is used to estimate parameters of animal populations from detections of marked individuals. In spatially explicit capture–recapture (SECR) the population is conceived as a spatial distribution of individual activity centers (AC) (Borchers & Efford, 2008; Efford & Fewster, 2013; Royle et al., 2014; Borchers & Fewster, 2016; Efford, 2025a). The method is well-suited to data from automatic cameras, given that individuals can be distinguished by their

natural marks, or from passive DNA samplers such as hair snares. By modeling the

distance-dependent spatial detection process, SECR avoids the assumption of constant detection

probability or effective sampling area that limits inference from raw counts used as population

indices (Kéry & Royle, 2021 Chapter 1).

We are concerned with variation in population density relative to factors of biological interest. In

the SECR model these factors are spatial or temporal covariates in a model for the intensity of a

point process for AC. A spatial covariate may be any known attribute of points in the potential

habitat; candidates include vegetation type, disturbance history, distance to human habitation, and

the spatial coordinates themselves. Spatial variation in population density is of general interest, as

shown by the frequency with which it appears in published SECR models (Tourani, 2022), and

recent papers have identified specific reasons for modeling density. For example, correlated

spatial variation in density and detection parameters causes bias in estimates of average density if

not modeled (McLellan et al., 2023; Efford, 2025a). Variation in animal density provides

important data for population management (e.g., Boulanger, Nielsen & Stenhouse, 2018; Lamb et

al., 2018; Ferrão da Costa et al., 2025). As an aside, many papers include illustrations of so-called

'density surfaces' that are not based on modeled density, but rather aggregate the estimated

sampling distributions of the AC of detected individuals; these have inferior statistical properties

and are no substitute for a density model (Durbach et al., 2024).

The rich possibilities in SECR for modeling density across space and time have appeared to run up

against constraints intrinsic to capture–recapture. For example, both spatial estimates of absolute

density and non-spatial estimates of population size are negatively biased by unmodelled variation

of detection parameters among individuals (e.g., Otis et al. 1978; Royle et al. 2013; Efford 2014;

Efford and Mowat 2014; Efford 2025c). A workaround is to include in the model individual characteristics (size, age, sex, etc.) that may account for the variation. Individual covariates are readily included in models that use a likelihood conditional on the number of detected individuals. Such models in turn provide individual-specific estimates of capture probability or effective sampling area, leading by Horvitz–Thompson methods to robust estimates of non-spatial population size (Huggins 1989; Alho 1990) or population density (Borchers & Efford, 2008). However, individual covariates have been considered incompatible with modeling spatial variation in SECR because that requires maximization of the full (unconditional) likelihood (Borchers & Efford, 2008; Borchers & Fewster, 2016; Sutherland, Royle & Linden, 2019), or an equivalent Bayesian method (Royle et al., 2014). This report shows that relative density (i.e. variation with respect to covariates) can be described by fitting the conditional model alone, leaving absolute density as an optional derived parameter, and that this approach has advantages for ecologists.

Spatial pattern is usually represented in SECR models by a log-linear model for density as a function of location-specific covariates. For the absolute density $D(\mathbf{x})$ at a point $\mathbf{x}$ (coordinates x, y) this takes the form

$$\log D(\mathbf{x}|\phi) = \beta_0 + \sum_{v=1}^{V} \beta_v c_v(\mathbf{x}), \tag{1}$$

where the $c_v(\mathbf{x})$ are V spatial covariates measured at $\mathbf{x}$ and the βs are coefficients to be estimated that together comprise the parameter vector ϕ. Categorical attributes are coded using indicator variables. The bases of regression splines may be used for fitting smooth trend surfaces (Borchers & Kidney, 2014). The logarithmic transformation ensures that the modeled density is always positive. The transformation is analogous to the 'link' function in generalized linear models (McCullagh & Nelder, 1989) and we use that term for convenience.

Relative density differs from absolute density by a constant factor that is given by the intercept β_0.

The benefits of focusing on relative density are demonstrated in this report by simulation and the

analysis of example datasets. They may be summarized as follows. Fitting the conditional model

is significantly faster than fitting the full model. While estimates of absolute density are biased by

heterogeneity of detection parameters among individuals, heterogeneity has little effect on the

estimated coefficients of relative density. Location-specific absolute density may be estimated as a

derived parameter. Alongside any density covariates, the conditional model may include

individual covariates of detection that improve model fit and potentially reduce bias in derived

estimates of absolute density. Conditional models with spatial variation in density and models

with individual covariates may be compared directly by information-theoretic criteria such as AIC

as they are no longer in separate domains.

Change in population size or density over time (population growth rate) is central to demographic

studies in ecology, and this is another application for conditional capture–recapture models. In

non-spatial open population capture–recapture models, the emphasis has been on estimating per

capita survival by the Cormack–Jolly–Seber (CJS) method that conditions on the first release of

each marked animal (Lebreton et al. 1992). Conditioning on the total number of detected

individuals in an open population model leads to estimates of both survival and recruitment

(Pradel, 1996; Link & Barker, 2005; Schofield & Barker, 2016), where recruitment can be

parameterized as the finite population growth rate λ. Efford & Schofield (2020) described a

spatial version of the Pradel–Link–Barker (PLB) method in which λ measures change in

population density i.e. λ is the *relative* density at consecutive discrete time steps. Temporal

variation in density may also be inferred from a chain of samples ('sessions') analysed without

regard to individual survival, as an extension of closed-population SECR (Efford, Borchers &

Byrom, 2009; Borchers & Fewster, 2016; Sutherland, Royle & Linden, 2019; Efford, 2025a). A

conditional multi-session closed model is proposed here that leads directly to estimates of the

session-specific λ.

Concluding remarks emphasize the central place of relative-density methods in SECR. Many

questions can be answered directly by fitting the relative density model, and absolute density is

available as a derived parameter if needed. Certain limitations of the method are acknowledged.

Conditional likelihood for relative density

The full SECR likelihood has the form

$$L(\phi, \theta | \Omega) = \Pr(\mathbf{\Omega} | n, \phi, \theta) \, \Pr(n | \phi, \theta), \tag{2}$$

where Ω is the set of n observed spatial detection histories, ϕ is a vector of density parameters and

θ is a vector of detection parameters (Borchers & Efford, 2008). The first factor on the right of

Equation (2) concerns the joint probability of the n observed detection histories. The second

factor concerns the number of detected animals n. When individuals are detected independently

of each other, the first factor may be expanded to

$$\Pr(\mathbf{\Omega} | n, \phi, \theta) \propto \prod_{i=1}^{n} \int \Pr(\boldsymbol{\omega}_i | \theta, \mathbf{x}) \, f(\mathbf{x} | \boldsymbol{\omega}_i > 0, \phi, \theta) \, d\mathbf{x}. \tag{3}$$

The actual location of each AC is unknown, so the location $\mathbf{x}$ is 'marginalized' out of the

likelihood by integrating over all possible locations in two dimensions. A critical factor is the

probability density $f(\mathbf{x} | \boldsymbol{\omega}_i > 0, \phi, \theta)$ for the AC location $\mathbf{x}$ of individual i, given that the

individual was detected ($\omega_i > 0$). That probability density has components for (i) the spatial distribution of the population, modeled with the density function $D(\mathbf{x}|\phi)$ from Equation (1), and (ii) the probability $p.(\mathbf{x}|\theta)$ that an individual with AC at $\mathbf{x}$ appears in the sample. Formally:

$$f(\mathbf{x}|\omega_i > 0, \phi, \theta) = \frac{D(\mathbf{x}|\phi)p.(\mathbf{x}|\theta)}{\int D(\mathbf{x}|\phi)p.(\mathbf{x}|\theta)d\mathbf{x}}. \tag{4}$$

The probability $p.(\mathbf{x}|\theta)$ combines the probabilities of detection at each detector, each possibly compounding detection over multiple occasions (Borchers & Efford, 2008; Efford, Borchers, & Byrom, 2009; Appendix S1). If density is uniform then $D(\mathbf{x}|\phi)$ cancels out of Equation (4) and maximizing the likelihood conditional on n in Equation (3) provides an estimate of θ (Borchers & Efford, 2008). However, if density is not uniform then $D(\mathbf{x}|\phi)$ does not cancel out of Equation (4) and the spatial variation in density must be included to obtain an unbiased estimate of θ. This is not as onerous as it sounds.

We define relative density by $D'(\mathbf{x}|\phi^-) \equiv k^{-1}D(\mathbf{x}|\phi)$, where ϕ^- is a vector with one fewer coefficients than ϕ and k^{-1} is a constant of proportionality. For the log-linear model, $k = \exp(\beta_0)$ and ϕ^- is simply ϕ without the intercept. We simplify and maximize the conditional likelihood in Equation (3) to estimate both the detection parameters θ and the coefficients of relative density ϕ^-. Cancelling the constant k yields the conditional likelihood

$$\Pr(\mathbf{\Omega}|n, \phi^-, \theta) \propto \prod_{i=1}^{n} \frac{\int \Pr(\omega_i|\theta, \mathbf{x}) \, D'(\mathbf{x}|\phi^-) \, p.(\mathbf{x}|\theta) \, d\mathbf{x}}{\int D'(\mathbf{x}|\phi^-) \, p.(\mathbf{x}|\theta) \, d\mathbf{x}}. \tag{5}$$

Individual covariates

Spatial detection is governed by the parameter vector θ that typically comprises the parameters λ_0 and σ for the intercept and spatial scale of a function relating detection hazard to distance

(Appendix S1). This use of λ_0 is unrelated to the conventional use of λ for population growth rate.

Known predictors of variation, such as weather or trap type, are included by specifying a linear submodel for either parameter on its link scale, as for density in Equation (1). Each submodel may also include predictors that vary among individuals ('individual covariates'), but these raise issues as we describe next.

To include individual covariates for detection parameters in the full likelihood it is necessary to model their distribution in the population. That is likely to differ from the distribution in the sample due to sampling bias. The shape of the covariate distribution must be specified and each evaluation of the likelihood requires integration over potential values of the covariate (e.g., Pollock, 2002). This is feasible for a single binary variable (e.g., sex) as the only parameter is the mixing proportion (e.g., sex ratio). For multi-level or continuous individual covariates the added model complexity and computational burden have discouraged implementations using the full likelihood. The problem is bypassed in Bayesian SECR using the complete data likelihood, as the population distribution of each individual covariate is included in the model (Link & Barker, 2010); an arbitrary shape is specified for the distribution (e.g., normal) and integration happens implicitly.

The problem is also bypassed by maximizing the conditional likelihood, without the need to specify the population distributions of covariates (Huggins, 1989; Alho, 1990). The conditional model concerns only detected individuals whose covariates are observed directly. Thus maximizing the conditional likelihood allows us simultaneously to fit models for both relative density and covariate effects on individual detection. Further, Horvitz-Thompson-like methods for inferring absolute population size or density from a sample adjust for sampling bias with respect

^{143} to covariates.

Absolute density as a derived parameter

^{145} Absolute density may be computed directly from a fitted model for relative density. Derived

^{146} estimates of density are the same as those from maximization of the full likelihood if the AC

^{147} follow a Poisson distribution in two dimensions, whether that is homogeneous or inhomogeneous

^{148} (see Discussion).

^{149} Given uniform density, the effective sampling area a is a function of θ that may include a vector of

^{150} covariates $\mathbf{z}_i$ for each individual i, hence $a(\theta, \mathbf{z}_i) = \int p.(\mathbf{x}|\theta, \mathbf{z}_i)\,d\mathbf{x}$. This leads to the

^{151} Horvitz-Thompson-like estimate of the uniform density $\hat{D} = \sum_{i=1}^{n} 1/a(\hat{\theta}, \mathbf{z}_i)$ (Borchers & Efford,

^{152} 2008). For non-uniform density, observe that when $p.(\mathbf{x}|\theta)$ does not depend on individual

^{153} covariates, $\mathrm{E}(n) = k \int D'(\mathbf{x}|\phi^-)p.(\mathbf{x}|\theta)\,d\mathbf{x}$, and $\hat{k} = n/\int D'(\mathbf{x}|\phi^-)p.(\mathbf{x}|\theta)\,d\mathbf{x}$, where $\hat{k}$ provides

^{154} the missing intercept of the absolute density model ($\hat{\beta}_0 = \log \hat{k}$). When detection parameters

^{155} depend on individual covariates the factor n is replaced by summation over n:

$$\hat{k} = \sum_{i=1}^{n} 1/\int D'(\mathbf{x}|\hat{\phi}^-)p.(\mathbf{x}|\hat{\theta}, \mathbf{z}_i)\,d\mathbf{x}, \tag{6}$$

^{156} The sampling variance of β_0 and its covariance with other βs may be estimated by the delta

^{157} method using numerical estimates of the gradient with respect to the estimated βs ($\hat{\phi}^-$). For

^{158} uniform density $D'(\mathbf{x}|\phi^-)$ takes the value 1, and $\int D'(\mathbf{x}|\phi^-)p.(\mathbf{x}|\theta, \mathbf{z}_i)\,d\mathbf{x}$ collapses to the

^{159} effective sampling area of individual i. Equation (6) is then the previous Horvitz-Thompson-like

^{160} estimate. Further, a derived estimate of the absolute density at a point $\mathbf{x}$ is

$$\hat{D}(\mathbf{x}|n, \hat{\phi}^-, \hat{\theta}) = \hat{k}\, D'(\mathbf{x}|\hat{\phi}^-). \tag{7}$$

Multi-session conditioning

When the data span multiple independent sub-populations ('sessions'), and the density model describes variation among them such as a temporal trend, conditioning is on the total number observed, as in PLB models (Efford & Schofield, 2020). The conditional likelihood is then the product of the T session-specific likelihoods and a factor for the session-specific counts n_t, modelled as multinomial with size $n = \sum n_t$:

$$\Pr(\Omega|n, \phi^-, \theta) \propto \frac{n!}{n_1!, n_2!, ..., n_T!} \prod_{t=1}^{T} p_t^{n_t} \prod_{i=1}^{n_t} \frac{\int \Pr(\omega_{it}|\theta, \mathbf{x})\, D'_t(\mathbf{x}|\phi^-)\, p_{\cdot t}(\mathbf{x}|\theta)\, d\mathbf{x}}{\int D'_t(\mathbf{x}|\phi^-)\, p_{\cdot t}(\mathbf{x}|\theta)\, d\mathbf{x}}, \tag{8}$$

where subscript t indicates the session. The multinomial probabilities p_t are estimated from the normalized expected number of detections in each session, using session-specific derived estimates of absolute density.

In order to parameterize the temporal model as population growth rate we model the cumulative change in density:

$$\log D(t|\phi) = \beta_0 + \sum_{j=1}^{t-1} l_j, \tag{9}$$

where β_0 is the initial log density and l_t is a linear predictor such that $\lambda_t = \exp l_t$. In the simplest model, with equal time intervals and session-specific λ_t, each l_t corresponds to one β and the parameter vector for the conditional model is $\phi^- = (\beta_1, \beta_2, ..., \beta_{T-1})$. Otherwise, there may be a linear submodel for the l_t in terms of a reduced set of βs, and the l_t may be scaled by time interval.

Simulations

Simulations were used to evaluate the performance of conditional likelihood models when the detection parameter λ_0 varied among individuals. Spatial pattern was generated as a Gaussian

random field whose intensity $c_1(\mathbf{x})$ was scaled to (0,1) and treated as a known covariate. Expected density was modeled by $\log D(\mathbf{x}) = \beta_0 + \beta_1 c_1(\mathbf{x})$, where $\beta_0 = \log(0.2)$ and $\beta_1 = \log(10)$; density therefore ranged between $0.2\sigma^{-2}$ and $2\sigma^{-2}$ depending on the covariate value at $\mathbf{x}$. AC were generated from an inhomogeneous Poisson distribution with the modeled density, and detections with a hazard-halfnormal function were simulated on a 10×10 square grid of binary proximity detectors at 2-σ spacing (Fig. 1). A conditional SECR model was fitted with the same structure as the generating model. The fitted model estimated the coefficient $\hat{\beta}_1$ directly, and $\hat{\beta}_0$ was derived using Equation (6). The intercept λ_0 of the exponential hazard function was allowed to vary among individuals according to a log-normal distribution with mean 0.2 and CV 0, 0.2, 0.4, 0.6 or 0.8. Simulations were repeated 2000 times for each level of $\mathrm{CV}(\lambda_0)$. The average number of individuals detected declined with increasing CV from $n = 265$ individuals (SD = 71.6) ($\mathrm{CV}(\lambda_0) = 0$) to 232 (SD 60.7) ($\mathrm{CV}(\lambda_0) = 0.8$), while the average number of detections per individual increased from 1.77 to 1.97. Results were expressed as the average absolute bias of each coefficient (absolute bias on the log scale approximates relative bias on the natural scale). R code and detailed results are given in section 'MhS' of Efford (2025c).

Estimates of $\hat{\beta}_0$, the intercept that determines absolute density, were strongly biased by unmodelled variation in detection, but, estimates of relative density ($\hat{\beta}_1$) were unaffected (Fig. 2). Execution time for the conditional likelihood fit (3 parameters) averaged 59% to 70% of that for the full likelihood (4 parameters). The absolute difference between estimates of β_1 from the full and conditional models was less than 0.0003 in all simulations.

Simulations were also conducted to assess the effect of variation in λ_0 on estimates of density and population growth rate from temporal data. The pattern of bias followed that for spatial pattern

(density biased, trend unbiased) so detailed results are not presented here; they are available in section 'MhT' of Efford (2025c).

Example. Spatial distribution of skinks

Data were analysed from a study of diurnal lizards in the Upper Buller Valley, South Island, New Zealand. Pitfall traps (covered sunken cans baited with a morsel of fruit in sugar syrup) were set in two large grids, each with 231 traps about 5 m apart and operated intermittently through 1995–1996. Traps were checked daily and individuals were marked uniquely by toe clipping. We selected data on one species (*Oligosoma infrapunctatum*) trapped for 3 days in November 1995. Ground cover and vegetation were recorded in a 1-m radius plot at each trap site. Each site was assigned to one of two habitat classes: 'low cover' (open with bare ground or low-canopy vegetation and grasses; 52.6% of sites) and 'high cover' (more-closed vegetation with a higher canopy; 47.5% of sites). A subset of the data was used by Efford & Fewster (2013) and the full dataset is included in Efford (2025b), where it is described further.

SECR models were fitted conditional on the number of trapped individuals, $n = 253$. Individuals varied greatly in body size measured as snout-vent length (SVL). This was incorporated as a predictor of the detection parameters λ_0 and σ either as a log-linear effect or as non-linear spline smooths with 3 or 5 degrees of freedom. Each candidate detection model was fitted with and without an effect of habitat class on local density. The model with lowest AIC included both the habitat effect on density and a non-linear effect of body size on λ_0 (Fig. 3). A conditional-likelihood habitat model without individual covariates model fit in 58% of the time

taken by a matching full-likelihood model. Details are in Appendix S2.

Oligosoma infrapunctatum was concentrated at sites with higher vegetation: density in the 'high' class exceeded density in the 'low' class by a factor of $\exp \beta_1 = 43.2$, regardless of whether the model allowed for the effect of body size on detection. Allowing for body size considerably increased the estimate of density, but the precision of density estimates was much reduced (Table 1).

Example. Brushtail possum population dynamics

The conditional-likelihood method for relative density through time (i.e. population growth rate λ) is illustrated with data from a multi-year trapping study of the brushtail possum *Trichosurus vulpecula* in New Zealand. A population in mixed forest was monitored by live trapping (Efford & Cowan, 2004). Analysis focused on June data from 1996 to 2005, a period of constant trapping effort (167 traps were set at the same sites at 30-m spacing for 5 nights). In total, 698 individuals were captured 4408 times. Annual population growth rate was estimated by multi-session closed-population SECR, parameterized as in Equation 9 of the main text. Direct estimates of λ from the conditional model parameterized as relative density were indistinguishable from those computed as the ratio of successive density estimates from maximization of the full-likelihood (Fig. 4). The year-specific conditional-likelihood model fit in 78% of the time taken by a matching full-likelihood model. The direct approach enables further modeling of λ. Models were compared with fixed $\lambda_t = 0$ (constant density), constant λ_t, session-specific λ_t, and smooths with varying degrees of freedom. Appendix S3 gives further details and results, including comparisons

 with open-population (PLB) models fitted to the same data.

Implementation

The number of identifiable coefficients for relative density in the conditional model (ϕ^-) is one

less than the number for absolute density in the full model (ϕ). Given software for the full

likelihood, the conditional likelihood is maximized by omitting the likelihood component for n

and fixing the intercept β_0 to zero, assuming a log link for density. One may also use an identity

link truncated at zero (Efford & Fewster, 2013); then $k = \beta_0$ and ϕ^- is ϕ with the intercept

removed and remaining coefficients divided by β_0. For the log link, estimated coefficients of the

Poisson relative density model are identical within numerical error to the corresponding

coefficients of the full model. For the identity link, the estimated coefficients equal those from the

full model divided by its intercept. The density intercept on the natural scale may be derived from

the density-weighted effective sampling areas of Equation (6).

Estimation of relative density by maximizing the conditional likelihood is included in recent

versions of the R package 'secr' (R Core Team, 2025; Efford, 2025b). Integration is performed in

'secr' by summing the integrand over a set of pixels for which its value is non-negligible, known

as the habitat mask. The conditional likelihood in Equation (5) is maximized numerically

whenever the CL argument of function 'secr.fit' is TRUE and a model formula is specified for

density. The usual asymptotic variance-covariance matrix for coefficients in the model (ϕ^-, θ) is

obtained from the negative inverse Hessian evaluated at the MLE. Functions 'derivedDcoef' and

'derivedDsurface' implement Equations (6) and (7) respectively. The parameterization in

²⁶¹ Equation (9) is selected with the 'Dlambda' details option.

²⁶² Discussion

²⁶³ A model for the distribution of animal activity centers may be split into a model for the density

²⁶⁴ relative to spatial or temporal covariates and a scalar that converts relative density to absolute

²⁶⁵ density. Relative density captures effects of ecological interest and is estimated efficiently by

²⁶⁶ maximizing the likelihood conditional on n, the number of observed individuals. This report

²⁶⁷ emphasises the central role and advantages of the conditional model, while acknowledging some

²⁶⁸ limitations.

²⁶⁹ Spatial variation in relative density is fully captured by a conditional model. Coefficients for both

²⁷⁰ detection (θ) and covariate effects on density (ϕ^-) do not differ between equivalent full- and

²⁷¹ conditional-likelihood models when the AC come from an inhomogeneous Poisson process, as in

²⁷² the simulations. The key here is that the statistic n is then Poisson (e.g., Efford & Fletcher, 2025),

²⁷³ and hence n is S-ancillary for the model coefficients in θ and ϕ (Schofield & Barker, 2016).

²⁷⁴ S-ancillarity means that n need not be modeled, and that derived estimates of absolute density

²⁷⁵ from the conditional model are identical to full likelihood estimates, within the limits of the

²⁷⁶ maximization algorithm. This property does not hold exactly when the number of AC in a

²⁷⁷ specified region is assumed to be fixed rather than Poisson; then the fitted model for n is binomial,

²⁷⁸ and n is no longer S-ancillary for the coefficients (Schofield & Barker, 2016).

²⁷⁹ Simulations also demonstrated that the coefficients of *relative* density (i.e. ϕ^-, excluding β_0) are

²⁸⁰ robust to unmodelled individual heterogeneity. This is true whether they are estimated by

maximizing the full or conditional likelihoods. However, estimates of β_0 and hence *absolute*

density from either model are biased by such heterogeneity. If absolute density is required, a

partial solution unique to the conditional model is to include individual covariates in submodels

for the detection parameters. This was demonstrated in the skink example, although confidence

intervals of the resulting estimates were wide.

Ecological investigations using SECR typically compare multiple models by information-theoretic

criteria such as AIC to find a 'best' model or to provide weights for model averaging. Comparison

is possible only within a class of model (full or conditional). The greater scope of conditional

models, now including both relative density and individual covariates, suggests that they should

be used for all such comparisons. Estimating one fewer parameters than in the full model leads to

faster model fitting, with time savings of 42% and 22% in the examples.

The effectiveness of the conditional model for estimating spatial variation in relative density has a

direct equivalent in the Pradel–Link–Barker (PLB) open population model that also conditions on

n (Schofield & Barker, 2016; Efford & Schofield, 2020). Temporal variation of the finite

population growth rate in PLB models is analogous to spatial variation in relative density. A

temporal series of closed-population samples may be analysed jointly by SECR, and it is

convenient to parameterize the joint model in terms of population growth rate, as in the brushtail

possum example (Fig. 4). Marescot et al. (2011) drew attention to the robustness and utility of

non-spatial estimates of population growth rate. Simulations reported separately (Efford, 2025c)

confirmed the robustness to individual heterogeneity of the population growth rate estimated from

a temporal series of closed-population samples. This mirrors the robustness of relative density

with respect to spatial covariates (Fig. 2).

Limitations

A case has been made for using SECR to model spatial variation in density, but there are limits to the ability of SECR of whatever flavor to resolve patterns and attribute variation to habitat covariates. The underlying Poisson point process model is stochastic and large samples are required to resolve patterns. The spatial detection process (scale σ) blurs the picture, especially when the study area is small relative to σ. The precision of habitat-specific estimates of absolute density can be poor, and is exacerbated by uncertainty in the detection model, as in the skink example.

Derived estimates of absolute density are readily computed by the Horvitz-Thompson method from any CL model. However, their variances must be obtained by the delta method which relies on a numerical estimate of the gradient vector for density with respect to the fitted coefficients (Huggins, 1989). Profile likelihood confidence intervals are not an option.

It is not possible to treat absolute density as a derived parameter when another component of the likelihood depends upon it. Specifically, the conditional model does not allow the spatial scale of detection σ to be a function of absolute density (Efford et al., 2016).

Variance estimates for population growth rate from a sequence of closed populations assume that each comprises an independent Poisson sample of AC. Survival of some individuals, each keeping its AC, violates this assumption and seems likely to cause a spurious improvement in precision, relative to an open population model that includes survival. However, the reverse appeared to be the case in the brushtail possum example (Appendix S3), possibly because survival reduced the component to be estimated. Further investigation is warranted.

Conclusion

Pollock (2002) and Royle (2009) expressed unease about the use of the conditional likelihood in closed-population capture–recapture on the abstract ground that the parameter of interest (population size in their case) was not in the model. However, there are few negative consequences in practice (preceding section). The conditional SECR model allows relative density to be modeled with respect to covariates despite the omission of absolute density. Fitting is fast and models with any combination of spatial and individual covariates may be compared by AIC. Attention is focused on the robustly estimated coefficients of the density model rather than the non-robust intercept that determines absolute density. Models fit by maximizing the conditional likelihood are sufficient to address questions regarding spatial or temporal variation in density, and they are a logical first step when the goal is to estimate absolute density.

Acknowledgments

I thank the many people who helped collect field data. The Lake Station skink study was a collaboration with Bruce Thomas, Nick Spencer, Rob Mason and Peter Williams. Contributors to the possum study are listed in Appendix S3. I acknowledge the use of New Zealand eScience Infrastructure (NeSI) high performance computing facilities URL https://www.nesi.org.nz. I thank Matt Schofield, Pierre Dupont and two anonymous referees for their comments on drafts.

Conflict of Interest Statement

The author declares no conflict of interest.

References

Alho, J. M. 1990. "Logistic regression in capture-recapture models." *Biometrics* 46: 623–635.

Borchers, D. L., and M. G. Efford. 2008. "Spatially Explicit Maximum Likelihood Methods for Capture–Recapture Studies." *Biometrics* 64: 377–385. https://doi.org/10.1111/j.1541-0420.2007.00927.x

Borchers, D. L., and R. M. Fewster. 2016. "Spatial Capture–Recapture Models." *Statistical Science* 31: 219–232. https://doi.org/10.1214/16-sts557

Borchers, D. L., and D. Kidney. 2014. "Flexible Density Surface Estimation for Spatially Explicit Capture–Recapture Surveys." CREEM Technical Report No. 2014-1, University of St Andrews. https://research-repository.st-andrews.ac.uk/bitstream/handle/10023/9147/secrgam_techrep.pdf?sequence=1

Boulanger, J., S. E. Nielsen, and G. B. Stenhouse. 2018. "Using Spatial Mark–Recapture for Conservation Monitoring of Grizzly Bear Populations in Alberta." *Scientific Reports* 8: 5204. https://doi.org/10.1038/s41598-018-23502-3

Durbach, I., R. Chopara, D. L. Borchers, R. Phillip, K. Sharma, and B. C. Stevenson. 2024. "That's Not the Mona Lisa! How to Interpret Spatial Capture–Recapture Density Surface Estimates." *Biometrics* 80(1). https://doi.org/10.1093/biomtc/ujad020

Efford, M. G. 2014. "Bias from Heterogeneous Usage of Space in Spatially Explicit Capture–recapture Analyses." *Methods in Ecology and Evolution* 5: 599–602.

Efford, M. G. 2025a. *The SECR Book. A Handbook of Spatially Explicit Capture–Recapture Methods*. Version 1.0.0. Zenodo. https://doi.org/10.5281/zenodo.15109938.

Efford, M. G. 2025b. "secr: Spatially Explicit Capture–Recapture Models." R package version 5.3.0. https://doi.org/10.32614/CRAN.package.secr.

Efford, M. G. 2025c. "Spatially explicit capture–recapture simulations." Zenodo. https://doi.org/10.5281/zenodo.17127944.

Efford, M. G. 2025d. "Example data for paper 'Spatially explicit capture-recapture models for relative density'." Zenodo. https://doi.org/10.5281/zenodo.17402993.

Efford, M. G., and P. E. Cowan. 2004. "Long-term population trend of *Trichosurus vulpecula* in the Orongorongo Valley, New Zealand." In *The Biology of Australian Possums and Gliders* edited by R. L. Goldingay and S. M. Jackson, pp. 471–483. Chipping Norton, NSW: Surrey Beatty & Sons.

Efford, M. G., and R. M. Fewster. 2013. "Estimating Population Size by Spatially Explicit Capture–Recapture." *Oikos* 122: 918–928. https://doi.org/10.1111/j.1600-0706.2012.20440.x

Efford, M. G., and D. Fletcher. 2025. "Effect of Spatial Overdispersion on Confidence Intervals for Population Density Estimated by Spatial Capture–Recapture." *Peer Community Journal* 5: e70. https://doi.org/10.24072/pcjournal.578

Efford, M. G., and G. Mowat. 2014. "Compensatory Heterogeneity in Spatially Explicit
    Capture–Recapture Data." *Ecology* 95:, 1341–1348.

Efford, M. G., and M. R. Schofield. 2020. "A Spatial Open-population Capture–Recapture
    Model." *Biometrics* 76: 392–402. https://doi.org/10.1111/biom.13150

Efford, M. G., D. L. Borchers, and A. E. Byrom. 2009. "Density Estimation by Spatially
    Explicit Capture–Recapture: Likelihood-based Methods." In *Modeling Demographic
    Processes in Marked Populations* edited by D. L. Thomson, E. G. Cooch, and M. J. Conroy,
    pp. 255–269. New York: Springer.

Efford, M. G., D. K. Dawson, Y. V. Jhala, and Q. Qureshi. 2016. "Density-dependent
    Home-range Size Revealed by Spatially Explicit Capture–Recapture." *Ecography* 39:
    676–688. https://doi.org/10.1111/ecog.01511

Ferrão da Costa, G., M. Mascarenhas, C. Fonseca, and C. Sutherland. 2025. "Spatial
    capture–recapture can improve environmental impact assessments for large carnivores."
    Ecological Solutions and Evidence 6: e70069.
    https://doi.org/10.1002/2688-8319.70069

Huggins, R. M. 1989. "On the Statistical Analysis of Capture Experiments." *Biometrika* 76:
    133–140.

Kéry, M., and J. A. Royle. 2021. *Applied Hierarchical Modeling in Ecology*. Vol. 2. London,
    UK: Elsevier/Academic Press.

Lamb, C. T., G. Mowat, A. Reid, L. Smit, M. Proctor, B. N. McLellan, S. E. Nielsen, and S.
    Boutin. 2018. "Effects of Habitat Quality and Access Management on the Density of a

Recovering Grizzly Bear Population." *Journal of Applied Ecology* 55: 1406–1417. https://doi.org/10.1111/1365-2664.13056

Link, W. A., and R. J. Barker. 2005. "Modeling Association Among Demographic Parameters in Analysis of Open Population Capture– Recapture Data." *Biometrics* 61: 46–54.

Link, W. A., and R. J. Barker. 2010. *Introduction to Bayesian Inference*. London, UK: Elsevier/Academic Press.

Marescot, L., R. Pradel, C. Duchamp, S. Cubaynes, E. Marboutin, R. Choquet, C. Miquel, and O. Gimenez. 2011. "Capture–Recapture Population Growth Rate as a Robust Tool Against Detection Heterogeneity for Population Management." *Ecological Applications* 21: 2898–2907.

McLellan, B. A., E. Howe, R. R. Marrotte, and J. M. Northrup. 2023. "Accounting for Heterogeneous Density and Detectability in Spatially Explicit Capture–Recapture Studies of Carnivores." *Ecosphere* 14: e4669. https://doi.org/10.1002/

McCullagh P., and J. A. Nelder. 1989. *Generalized Linear Models*. London, UK: Chapman and Hall.

Otis, D. L., K. P. Burnham, G. C. White, and D. R. Anderson. 1978. "Statistical Inference from Capture Data on Closed Animal Populations." *Wildlife Monographs* 62: 1–135.

Pollock, K. H. 2002. "The use of auxiliary variables in capture–recapture modelling: an overview." *Journal of Applied Statistics* 29: 85–102.

Pradel, R. 1996. "Utilization of Capture–Mark–Recapture for the Study of Recruitment and Population Growth Rate." *Biometrics* 52: 703–709.

R Core Team. 2025. *R: A Language and Environment for Statistical Computing.* Vienna: R Foundation for Statistical Computing. https://www.R-project.org/

Royle, J. A. 2009. "Analysis of Capture–Recapture Models with Individual Covariates Using Data Augmentation." *Biometrics* 65: 267–274.

Royle, J. A., R. B. Chandler, C. C. Sun, and A. K. Fuller. 2013. "Integrating Resource Selection Information With Spatial Capture–Recapture." *Methods in Ecology and Evolution* 4: 520–530.

Royle, J. A., R. B. Chandler, R. Sollmann, and B. Gardner. 2014. *Spatial Capture–Recapture.* Waltham, MA: Academic Press.

Schofield, M. R., and R. J. Barker. 2016. "50-year-old Curiosities: Ancillarity and Inference in Capture–Recapture Models." *Statistical Science* 31: 161–174. https://doi.org/10.1214/16-STS550

Sutherland, C., J. A. Royle, and D. W. Linden. 2019. "oSCR: a Spatial Capture–Recapture R Package for Inference About Spatial Ecological Processes." *Ecography* 42: 1459–1469.

Tourani, M. 2022. "A Review of Spatial Capture–Recapture: Ecological Insights, Limitations, and Prospects." *Ecology and Evolution* 12:e8468. https://doi.org/10.1002/ece3.8468

Table 1: Estimates of habitat-specific density of the skink *Oligosoma infrapunctatum* from model with no individual covariate of detection hazard λ_0, and a model with spline smooth of snout-vent length $\lambda_0 \sim s(SVL, k = 3)$.

Model	Habitat class	Density (CI) ha^{-1}	
$\lambda_0 \sim 1$	Low	10.7	(1.4, 81.9)
	High	524.4	(455.3, 604.0)
$\lambda_0 \sim s(SVL, k = 3)$	Low	18.8	(2.7, 129.7)
	High	815.6	(432.0, 1539.6)

Figure captions

Figure 1. Example of simulated spatial covariate c_1 and activity centres (black dots) generated from inhomogeneous Poisson model for intensity as a log-linear function of the covariate. Detector sites are shown as red crosses.

Figure 2. Bias of estimated coefficients of SECR model for density in relation to unmodeled individual heterogeneity in baseline detection parameter λ_0. The log-linear spatial density model had intercept β_0 and slope β_1 with respect to covariate c_1 (Fig. 1). Simulated data: average of 2000 replicates with 95% CI.

Figure 3. Fitted relationship between the baseline detection hazard λ_0 and the individual covariate snout-vent length (SVL) of pitfall-trapped skinks *Oligisoma infrapunctatum*. Estimates from regression splines fitted in a conditional-likelihood SECR model. Rug plot indicates distribution of observed SVL.

Figure 4. Annual population growth rate λ of brushtail possums *Trichosurus vulpecula*. The model for λ used either a 9-level factor for year-specific estimates (blue) or a spline smooth with $k = 7$ (orange, joined). Bars are 95% confidence intervals. The smooth model had lower AIC and shorter intervals, similar to those from an open population model (Appendix S3). Estimates of λ as the ratio of successive session-specific estimates of density from a full-likelihood model exactly overlaid the session-specific conditional-likelihood $\hat{\lambda}$ and are not shown.

 # Figures

Figure 1

Figure 1

Figure 2

Figure 2

Figure 3

Figure 3

Figure 4

Figure 4