History shapes regulatory and evolutionary responses to tigecycline in strains of *Acinetobacter baumannii* from the pre- and post-antibiotic eras

Alecia B Rokes[1,2], Alfonso Santos-Lopez[3], Vaughn S Cooper[1,2]

[1]University of Pittsburgh, Department of Microbiology and Molecular Genetics, Pittsburgh, PA, USA

[2]Center for Evolutionary Biology and Medicine, University of Pittsburgh, Pittsburgh, PA, USA

[3]Department of Molecular Biology, Autonomous University of Madrid, Madrid, Spain

ASL
ORCID: 0000-0002-9163-9947
ABR
ORCID: 0000-0001-9496-0296
VSC
ORCID: 0000-0001-7726-0765

1 History shapes regulatory and evolutionary responses to tigecycline in strains of *Acinetobacter*
2 *baumannii* from the pre- and post-antibiotic eras
3

ABSTRACT

Evolutionary history encompasses genetic and phenotypic bacterial differences, but the
extent to which history influences drug response and antimicrobial resistance (AMR) adaptation
is unclear. Historical contingencies arise when elements from an organism's past leave lasting
effects on the genome, altering the paths available for adaptation. We utilize strains isolated
before and after widespread antibiotic use to study the impact of deep historical differences
shaped by decades of evolution in varying antibiotic and host pressures. We evaluated these
effects by comparing immediate and adaptive responses of two strains of *Acinetobacter*
baumannii to the last-resort antibiotic, tigecycline (TGC). When grown in subinhibitory TGC, the
two strains demonstrated divergent transcriptional responses suggesting that baseline
transcript levels may dictate global responses to drug and their subsequent evolutionary
trajectories. Experimental evolution in TGC revealed clear differences in population-genetic
dynamics – with hard sweeps in populations founded by one strain and no mutations reaching
fixation in the other strain. Transcriptomes of evolved populations no longer showed signatures
of drug response, as was seen in the ancestors, suggesting that genetic adaptation may
outweigh preexisting differences in transcriptional networks. Genetically, AMR was acquired
through predictable mechanisms of increased efflux and drug target modification; however, the
two strains adapted by mutations in different efflux regulators. Fitness tradeoffs of AMR were
only observed in lineages evolved from the pre-antibiotic era strain, suggesting that decades of
adaptation to antibiotics resulted in preexisting compensatory mechanisms in the more
contemporary isolate, an important example of a beneficial effect of historical contingencies.

SIGNIFICANCE STATEMENT
Acinetobacter baumannii is a high priority pathogen often causing multidrug resistant
nosocomial infections. Many healthcare systems experience clonal outbreaks of *A. baumannii*
infections, yet treatment strategies are often strain-agnostic, ignoring the importance of strain
differences. We show that historical differences between two strains, one isolated prior to
widespread antibiotic use and the other following decades of selection to clinical conditions,
dictate transcriptional patterns and response to a last-resort antibiotic and influence the
genetic and phenotypic routes of resistance adaptation. While our study focuses on two
reference strains of *A. baumannii,* these findings can be more broadly applicable to other
pathogenic organisms in which a better understanding of the forces influencing resistance
adaptation is essential for combating the antimicrobial resistance crisis.
37
38

INTRODUCTION

<u>INTRODUCTION</u>

Evolution is an inescapable process, as the worsening antimicrobial resistance (AMR)

crisis demonstrates (1). However, the AMR crisis will be less daunting if we are better able to

understand, predict, and direct the evolutionary outcomes of antibiotic treatment (2, 3). With

better resolution of factors affecting the routes to resistance, we may be able to work toward

directing evolution to avenues with exploitable resistance tradeoffs (4, 5). However, genotypic

and phenotypic routes to resistance, as well as the resulting fitness tradeoffs, may be highly

influenced by strain history, decreasing the range and applicability of evolutionary strategies to

counteract the evolution of AMR (6, 7).

There are three primary forces, often intertwined, that influence evolution: history,

chance, and selection (5, 8, 9). Natural selection causes the most fit alleles in a population to

rise in frequency while less fit mutants are lost from the population. Selection needs heritable

phenotypic variance to act upon that is produced by mutations arising randomly, a factor of

chance. Chance events also occur in the demography of microbial populations, such as

population bottlenecks arising during infections (10). The outcome of a mutation is also

affected by the genetic background in which it arises (11, 12), combining chance and the final

force of evolutionary history, which can influence adaptation in diverse ways but remains

understudied.

Evolutionary history encompasses the genetic and phenotypic remnants from a

bacterium's past including biotic and abiotic factors, such as interactions with other cells and

previous environmental exposures like antibiotics, respectively (13). The phylogenetic origin of

the bacterium is also a major component of evolutionary history. These differences can impose

historical contingencies through which preexisting elements in the genome, transcriptome, or

phenome alter or constrain evolutionary paths available for adaptation (13, 14). History may

dictate the type, identity, rate, and order of mutations acquired (15–18). Additionally, history

likely influences fitness and phenotype associated with a mutation, thereby dictating

evolutionary trajectories and outcomes (19–21). This study seeks to understand how historical

contingencies influence evolutionary trajectories under the strong selective pressure of

antibiotic treatment (22).

Contingencies imposed by history exacerbate our inability to accurately predict bacterial

responses to antibiotics and AMR evolution. *Acinetobacter baumannii* is a nosocomial pathogen

of increasing concern due to limited and often ineffective treatment options (23). With time

and increased antibiotic use, strains of *A. baumannii* have acquired mutations and mobile

genetic elements, on top of many conserved genes, that endow desiccation resistance, stress

tolerance, and drug resistance (24, 25). Such historical differences could cause archaic strains,

which are often represented by laboratory reference strains, to respond and evolve differently

than contemporary infectious strains (26, 27). Importantly, hospitals typically experience clonal

outbreaks of *A. baumannii* caused by closely related isolates of the same sequence type (28,

29). Across hospital systems, *A. baumannii* infections exhibit high diversity (30, 31), but only a

handful of treatments are used. This practice ignores the importance of genetic background

and evolutionary history in the treatment process, likely leading to differences in treatment

response (32, 33). A better understanding of how various strains respond and adapt to

antibiotics would result in improved patient outcomes.

The widespread use of antibiotics in clinical practice has dramatically affected pathogen

evolution (7, 34). Comparing strains isolated before and after the clinical introduction of

antibiotics provides an excellent framework to study how deep historical differences influences

the evolution of resistance (35). To this end, we used two strains of *A. baumannii,* one that was

isolated early in the antibiotic era and has not been exposed to many contemporary antibiotics,

and one that is better representative of current multi-drug-resistant (MDR) clinical infections

having been isolated much more recently. Comparing these strains enables assessment of how

a history of evolving with the human host and novel antimicrobial pressures impacts resistance

evolvability.

Resistance adaptation can be investigated at multiple levels: from the narrowest, the

site-specificity of individual mutations, to more systematic resistance pathways, and even more

broadly, at the level of population-wide responses and adaptive dynamics (36, 37). This study

looks within and across these levels to determine the areas in which evolutionary history is

most influential. We find that deep history influences the basal transcriptome, driving

differences in the response to TGC. The strains also differed in their evolved responses to

prolonged TGC stress at both narrow molecular-genetic and broader population-genetic levels,

despite superficial similarities. This work highlights the importance of evolutionary history in

the response and adaptation of a high-priority pathogen to a last-resort antibiotic, showing that

strain differences can reduce the predictability of treatment outcomes.

<u>RESULTS</u>

<u>Genomic and phenotypic classifications of the strains</u>

Previous studies of historical differences in strain response to antibiotics have used

strains separated by relatively few mutations (9). Here, we study effects of deeper history on

the evolution of AMR in two different *A. baumannii* strains isolated nearly 60 years apart, *i.e.*

before and after the widespread clinical use of antibiotics. The first strain, 17978UN, is a variant

of the commonly used laboratory reference strain ATCC 17978 (38). This strain was isolated in

1951 in the early stages of the introduction of antibiotics in clinical practice (39). In contrast,

the second strain, AB5075-UW, is a variant of AB5075, which was isolated in 2008 from a MDR

infection representative of the current global clinical burden of *A. baumannii* (40, 41).

Therefore, the evolutionary histories of both strains have likely been shaped by different

exposures to antibiotics, hospitals, and the human host. In many ways 17978UN represents

archaic *A. baumannii* infections in comparison to contemporary MDR infections represented by
AB5075-UW.
We first characterized genomic and phenotypic differences between the two strains to
evaluate their historical distinctions. 17978UN is a member of sequence type ST437 whereas
AB5075-UW is assigned to ST1, part of the clinically dominant clonal complex 1 (40). Their
genomes share an average nucleotide identity of 97.5% and 3265 homologous core genes.
17978UN contains an additional 694 accessory genes whereas AB5075-UW encodes 751
accessory genes (Data S1), with no significant differences in gene content grouped by Clusters
of Orthologous Genes (COG) categories (Table S1). However, as expected by its lineage and
history of antibiotic exposure, AB5075-UW has significantly more resistance-associated
elements than 17978UN (Table S1) and is more resistant to a variety of antibiotics (Table S2).
Notably, both strains were susceptible to at least two different classes of antibiotics.
These significant differences in genome content and existing AMR led to the hypothesis
that the strains would respond differently to a contemporary, clinically relevant antibiotic.
Tigecycline (TGC) is a translation-inhibiting antimicrobial compound that was approved for
clinical use in 2005 (42). Resistance to TGC in clinical isolates of *A. baumannii* is typically
attributed to overexpression of efflux pumps or ribosomal modifications (43, 44). The minimum
inhibitory concentration (MIC) of TGC differed only slightly between the strains (17978UN
MIC_{TGC} = 0.125 µg/mL, AB5075-UW MIC_{TGC} = 0.25 µg/mL). Further, subinhibitory TGC (0.06
µg/mL) imposed approximately 75% and 60% fitness defects in 17978UN and AB5075-UW,
respectively (Fig. S1). This comparable level of growth inhibition by TGC enables the study of
influences of deep evolutionary history on both transcriptional response and subsequent
evolutionary adaptation.
138
<u>Strain-dependent transcriptional response acts on historically differentially utilized genes</u>
Given the stress imposed on growth by TGC, even at subinhibitory levels, we predicted
that transcriptional responses to this drug would be largely conserved between strains, likely
consisting of stress-response genes, drug resistance mechanisms, or ribosomal genes specific to
the mechanism of action of TGC (45, 46). We compared transcript levels following growth in
subinhibitory TGC to growth in minimal media lacking antibiotic. The defined minimal media
used here and throughout was M9 salts buffered supplemented with glucose, amino acids, and
other elements as described in the methods (47). As expected, TGC stress caused large
transcriptional changes biased toward gene downregulation in both strains, but there was
surprisingly minimal overlap among differentially expressed genes between strains (Fig. 1 *A* and
B). Only 144 genes, representing 24% of the total downregulated genes, and 53 genes,
representing 18% of upregulated genes, were differentially expressed in the same direction in
both strains. One might intuit that the limited shared response to drug may have resulted from
transcriptional differences among strain-specific genes, but this was not the case. Rather, most

genes that were differentially expressed were within the shared genome (Fig. 1 *A* and *B*; Data
S1), including many genes with the greatest changes in expression that shifted in opposite
directions between strains (Fig. S2 and Fig. S3). Together these results show that the response
to TGC stress is historically contingent.
To test whether baseline transcriptional differences explained varied strain response to
TGC, we compared transcript abundance of conserved genes in the absence of drug (Fig. 1*C*,
left). This comparison revealed differences in baseline transcriptional state that, upon TGC
pressure, caused expression to converge upon a common pattern of gene usage (Fig. 1*C*, *right*).
Genes with higher basal transcription in each strain were preferentially downregulated within
that strain in response to TGC treatment (Fig. 1*C*; Fig. S4), producing a more conserved
transcriptional state under TGC pressure (Animation S1). These historical differences in basal
gene expression explain the strain-dependent responses to TGC pressure.
165
Experimental evolution for tigecycline resistance results in strain-dependent tradeoffs between
fitness and resistance
Given the genomic signatures of history seen in the transcriptional response to TGC, we
hypothesized that evolutionary adaptation to this antibiotic would proceed along different
paths in each strain. We propagated three replicate populations of both strains for 12 days in
increasing concentrations of TGC (methods and Fig. S5). All populations evolved excess
resistance beyond the drug concentration in the media, which was maintained throughout the
experiment, such that all populations became nearly five times more resistant to TGC than their
respective ancestors (Fig. 2).
Antibiotic resistance is often associated with a fitness cost and these costs can depend
on the strain background (17, 48–50). We expected to find fitness deficits of TGC-evolved
populations when grown in the absence of drug, whereas we expected higher fitness of TGC-
evolved populations than their respective ancestors when grown in subinhibitory TGC. As
expected, both sets of evolved populations exhibited substantial fitness increases in
subinhibitory TGC. However, evolved fitness responses in the absence of drug differed.
Populations derived from the laboratory reference strain, 17978UN, incurred a 25-50%
reduction in fitness, whereas no measurable fitness cost was observed for populations derived
from the more recently isolated clinical strain, AB5075-UW (Fig. 2). These results show that
evolutionary tradeoffs between fitness and resistance to TGC differ by founding strain; and
suggest that contemporary MDR strains may adapt to new drugs with limited collateral fitness
costs.
187
Varying population dynamics within conserved mechanisms of TGC resistance
The divergent relationships between fitness and resistance indicated that the genetic
pathways to TGC resistance may differ between the two strains. We used longitudinal whole-

population, whole-genome sequencing to investigate the population-genetic dynamics of
adaptation. Following prior evolution experiments in antibiotics (47, 51), we predicted that
antibiotic pressure in well-mixed populations would be sufficiently strong that mutations
providing resistance should rapidly sweep within the population. Indeed, mutations in genes
associated with drug resistance rapidly fixed in all populations of AB5075-UW (Fig. 3; Data S2),
but surprisingly, no mutations reached fixation in populations of 17978UN. A closer inspection
of the mutations in each 17978UN line suggests that in total, they represent a soft sweep of
resistance-associated mutations. Soft sweeps occur when multiple independent sub-lineages
acquire mutations in the same genes; individual mutations never reach fixation, but the sum of
all co-existing analogous mutations in a gene or pathway approach fixation within the
population (52).
The evolved mutations pointed to two primary mechanisms of resistance to TGC:
modification of the enzymatic target of the drug, the ribosome, and decreasing the intracellular
concentration of antibiotic, primarily through efflux pumps. All replicate populations of both
strains featured high-frequency mutations affecting the same pathways for drug resistance and
even in the same gene. This gene encodes a *S*-Adenosyl-methionine-dependent
methyltransferase (SAM-MT; Fig. 3*A*) that has been associated with creating methylation
patterns required for TGC binding to the ribosome (53). Interestingly, all SAM-MT mutations
that reached high frequency under TGC stress were gene disruptions caused by mobilized
insertion sequences that are predicted to eliminate function (Data S2). Although each
population acquired mutations in the same SAM-MT, the specific insertion sequence causing
the loss of function as well as the site of insertion varied among lineages, with more variability
within populations of AB5075-UW. In addition to the SAM-MT, other genes related to the target
of TGC acquired mutations, such as ribosome recycling factors (Fig. 3*A*; Data S2) that likely aid
in rescuing stalled ribosomes (43).
All populations also evolved mutations in regulators of efflux pumps, but the specific
resistance-nodulation-division (RND) pump that was affected differed between strains.
Populations of 17978UN also acquired mutations affecting other systems predicted to reduce
intracellular concentrations of TGC. For instance, two 17978UN populations acquired mutations
in major facilitator superfamily (MFS) transporters that rise to high frequency (Fig. 3*B;* Data S2)
(54). In summary, the identities, timing, and frequencies of evolved mutations differed between
strains, whereas replicate populations founded by the same strain exhibited high gene-level
parallelism.
224
<u>Strain-dependent efflux pump preferences</u>
All populations adapted to TGC through mutations in drug efflux systems (Fig. 3*B*), yet
different strains evolved with different genetic signatures. *A. baumannii* encodes three RND-
efflux pumps known to be associated with drug resistance adaptation (55, 56). Two-component

system *adeRS* regulates efflux operon *adeABC,* while the other two pumps, *adeFGH* and *adeIJK*
are regulated by negative regulators *adeL* and *adeN,* respectively (Fig. 4*A*) (57). Both strains
encode all three pumps with a high level (97-100%) of amino acid conservation (Table S2), with
the only exception that 17978UN does not encode the outer membrane protein, AdeC, but
AdeK can complement function (55). Populations of AB5075-UW acquired nonsynonymous
mutations throughout the *adeRSABC* efflux system, altering *adeA*, *adeB*, and *adeS*. One of two
predominant single nucleotide polymorphisms (SNPs) in *adeS* (R152K and G160S) fixed in all
AB5075-UW populations, and an additional SNP in *adeB* (E939Q) was nearly fixed in one
population (Data S2). In contrast, each evolved 17978UN population acquired mutations in
adeL caused by an insertion sequence (IS) disruption in this gene that rose to high frequency
(Fig. 3*B*; Data S2). This disruption exhibited site-specific parallelism in which the mutation was
identical within all three TGC-evolved populations, but never reached 100% frequency. As an
example of the soft sweep dynamics described previously, in evolved population 3 we observe
an additional mutation in *adeL* and the two mutations in this gene, cumulatively, reach fixation.
Interestingly, a nonsynonymous SNP in *adeR* also arose in 17978UN populations 1 and 3 on day
three but it was quickly outcompeted by *adeL* mutations (Fig. 3*B*), further emphasizing the
strain-dependence of efflux mutations*.*
There are many ways in which historical contingencies could cause such strain-specific
preferences of efflux targets. One possibility is that the sequences of these target genes had
already differentiated, but all sites that acquired adaptive mutations were initially identical in
the ancestral genomes (Table S3). Another possibility is that the ancestral strains had previously
evolved differential regulation of these pumps by other mechanisms. To test this notion, we
measured expression of efflux genes in the presence and absence of subinhibitory TGC but
found comparable levels of expression of *adeABC* and *adeFGH* (Fig. 4*B*), discounting this
explanation. We found that 17978UN exhibits relatively greater expression of the *adeIJK* pump
in both growth conditions, but this pump was not the target of selection in either strain. In the
absence of TGC, 17978UN had increased efflux activity compared to AB5075-UW (Fig. 4*C*),
possibly due to the differences in AdeIJK expression. Interestingly, drug pressure induced
significantly different states of efflux activity, increasing efflux activity in AB5075-UW but
decreasing it in 17978UN (Fig. 4*C*). As expected, there were significant differences in efflux
activity and expression in the evolved populations, confirming that the mutations in efflux
regulators produced a meaningful phenotype. All evolved populations, except 17978UN
evolved population 3 which still trended higher, evolved significantly increased efflux activity
than their respective ancestors (Fig. 4*D*). Lastly, we show that the mutated regulators
specifically increase expression of their associated pump (Fig. 4*E*). In summary, neither
sequence divergence nor functional differences in the primary targets of selection explained
the alternative evolutionary pathways to increased drug efflux. These results strongly suggest

that more complex interactions between efflux systems and the genome favor different routes

to drug adaptation.

<u>Diminished transcriptional response and historical influences following TGC adaptation</u>

The ancestral strains exhibited divergent transcriptomic changes in response to TGC

stress (Fig. 1), but both adapted to TGC via broadly similar mechanisms. To evaluate the

transcriptional signature of evolved drug tolerance, we conducted RNA sequencing of evolved

populations grown with or without TGC treatment and found remarkably few differences (Fig.

5*A*). Compared to the ancestral response, where many genes changed in abundance upon TGC

treatment, all evolved populations exhibited nearly identical transcriptomes in the two

conditions. This trend is more conspicuous in populations of AB5075-UW, where no genes are

significantly different (p-value < 0.05 and absolute value log fold change > 1) between the two

conditions, while a few genes remain significantly differently expressed upon TGC stress in

populations of 17978UN (Fig. S6, Data S3).

While the muted transcriptional response to drug is observed in all evolved populations,

it does not tell us if evolved transcriptional responses are conserved between strains. Principal

component analysis of transcript abundance confirmed that TGC treatment shifts the initially

diverging ancestral transcriptomes in the same direction while still preserving strain differences

(Fig. 5*B*). Transcriptomes of evolved populations, either treated or untreated, strongly grouped

together. This cluster of evolved populations groups more closely with the untreated ancestors,

further supporting limited transcriptional response to TGC stress by evolved populations (Fig.

5*B*; Fig. S6). Taken together, these convergent evolved transcriptomes show that genomic drug

adaptations alleviate the need to transcriptionally respond to TGC pressure and result in

reduced historical signatures on the transcriptome compared to the divergent ancestral

transcriptomes.

<u>DISCUSSION</u>

Antimicrobial resistance (AMR) is a problem of evolution that is most often observed in

retrospect, when treatment fails. We still have much to learn about how AMR evolves,

including the diversity of genetic pathways to resistance (22), the extent to which these

pathways depend on the environment (47) including the host immune status (58), and whether

these pathways differ among historically divergent strains (18, 20). In this study, we focused on

this last factor by evaluating how reference strains isolated nearly 60 years apart, spanning eras

of antibiotic usage, differ in their adaptation to a clinically significant drug. Although both

strains of *A. baumannii* adapted to grow in media containing tigecycline (TGC) via common

pathways, the specific genes and mutations differed with the founding strain. On the other

hand, replicate populations derived from the same strain evolved in parallel, indicating a level

of genetic predictability within but not between strains.

Perhaps the most striking difference between strains was the fitness consequence of

TGC adaptation: lineages derived from the modern clinical strain experienced no measurable

fitness costs whereas those derived from the historical strain, 17978UN, incurred deficits of

25% or more (Fig. 2). Fitness effects are often related to the stability or persistence of

resistance. Costly resistance has long been proposed as a beneficial trait for humanity that can

preserve drug efficacy in the face of selection (3, 59, 60). Here we find an alarming lack of a

fitness cost of evolved resistance in lineages derived from the contemporary clinical isolate,

AB5075-UW. The evolutionary history of AB5075-UW includes frequent antibiotic exposure

prior to our studies that could have selected for compensatory adaptations that alleviate costs

of TGC resistance (61). The lack of a fitness-resistance tradeoff in AB5075-UW suggests that

passive loss of resistance once drug pressure is alleviated will be rare in related strains,

increasing the need for alternative treatment options.

Differences in the genetic causes of resistance between strains offer evidence that

historical contingency influences routes to AMR. Many bacterial genomes including *A.*

baumannii encode multiple RND efflux pumps and to explain this apparent redundancy, a

hierarchical model of sequential activation has been proposed (54, 57, 62). We find that the

first, dominant mutations affecting drug efflux in each strain affected regulators of different

pumps, with SNPs in *adeS* sweeping to fixation in AB5075-UW populations and gene disruptions

caused by IS in *adeL* reaching high frequencies in 17978UN populations (Fig. 3). This suggests

that the hierarchy of these pumps differs between strains in the face of TGC pressure and raises

the possibility that variation in their regulatory network or gene content evolved in the past.

Perhaps the diminished priority of the AdeRSABC pump in 17978UN stems to the lack of *adeC* in

this strain (Fig. 4*A*; Data S1). These lineages would need to accumulate two mutations to

overexpress one functional pump: one that upregulates AdeAB and one that upregulates AdeK

to serve as the outer membrane protein (55). The historical contingency imposed from the loss

of *adeC* sometime in the history of 17978UN may be driving the strain-dependent efflux

preferences for drug adaptation.

Both the spectra of evolved mutations and their dynamics differed between strains,

suggesting intrinsic biases that influence genetic diversity. Despite having the same number of

encoded insertion sequences (IS) as AB5075-UW (Table S1), nearly all high-frequency mutations

in populations of 17978UN were caused by IS interrupting open reading frames, whereas

populations of AB5075-UW experienced a greater diversity of mutation types. IS elements are

known to play an important role in *A. baumannii* mutagenesis (63). The strain-dependent

historical contingencies on insertion sequence mobilization, repertoire, and insertion site could

have lasting impacts on the future stability of these mutations and, therefore, the stability of

resistance stemming from IS mediated mechanisms (63, 64). Further, resistance-associated

mutations fixed in populations of AB5075-UW, but no mutations fixed in populations of

17978UN, indicating coexisting lineages. The maintained diversity in 17978UN lineages may be

due to relatively greater mutation availability causing the coincident rise of multiple adapted
lineages. When multiple mutations in the same gene collectively reach fixation, this
phenomenon is known as a soft sweep (65, 66). At the population level, the diversity
maintained in a population undergoing selection from antibiotics is an important indicator of
future paths available for additional adaptation in new environments. In this regard,
populations of 17978UN may be more evolvable when challenged with a new stress.
It would be ideal to be able to predict evolutionary trajectories of AMR from patterns of
transient response to an antibiotic. While we do not see clear signatures that may enable
predictions of drug adaptation at the gene level (45), our work shows that strain background
greatly influences transcriptomic response to antibiotics. The large transcriptomic differences
between strains in the absence of drug, which in turn influenced their responses to TGC (Fig. 1),
was remarkable and could be used to map functional contingencies between strains. Many
more transcriptomes of additional strains would be needed to evaluate the power of
transcriptomes as markers of evolutionary history and to anticipate drug responses.
Nonetheless, our studies suggest the existence of a conserved transcriptional state under TGC
stress that different unexposed cell-states migrate toward. The conserved transcriptomes of
evolved populations harboring several new AMR mutations further support a homogeneous
state associated with drug adaptation (Fig. 5). We noted that the evolved transcriptomes more
closely represent the untreated ancestral state, suggesting that the evolved populations are,
essentially, not feeling the pressure of the antibiotic, likely due to the adaptations in drug efflux
decreasing the intracellular concentrations of TGC (Fig. 4). It would be valuable to test if the
collapse of transcriptional heterogeneity upon transient antibiotic pressure as well as
prolonged adaptation is a general pattern or is contingent on the class or concentration of
antibiotic applied. A lack of transcriptional response to antibiotics in clinically adapted samples
could be used as another biomarker upon which to predict the extent of genetic resistance
adaptation.
In a world shifting toward personalized medicine, we should also shift treatment of
infections to be tailored to the specific strain or sequence type causing the infection. This work
supports the idea that initial responses as well as evolutionary trajectories to antibiotic stress
can be highly strain-dependent, and therefore, treatments should be as well. If we are better
able to predict the impact of historical contingencies in the transcriptional response and in
fitness tradeoffs associated with AMR evolution, we will be one step closer to personalized
treatment of MDR infections.
375
METHODS
Bacterial strains and growth conditions
Laboratory reference strain ATCC 17978UN (38, 39) and clinical reference strain
AB5075-UW (40, 41) were used throughout the study and grown under the same conditions.

Unless otherwise stated, bacterial cultures were grown in 5ml M9+ media, as previously
described (47). Briefly, M9+ media is a salts-buffered media with glucose (11.1mM) as the
primary carbon source. It contains 0.1 mM $CaCl_2$, 1 mM $MgSO_4$, 42.2 mM Na_2HPO_4, 22 mM
KH2PO$_4$, 21.7 mM NaCl, 18.7 mM NH_4Cl and is supplemented with 20 mL/L MEM essential
amino acids (Gibco 11130051), 10 mL/L MEM nonessential amino acids (Gibco 11140050), and
1 mL each of trace mineral solutions A, B, and C (Corning 25021–3 Cl). Cultures were incubated
at 37°C with shaking or on a roller-drum at approximately 250rpm.
387
Genomic comparison of the two strains
Genomes were annotated with bakta (v1.6.1, database v4.0) (67) and input to Panaroo
(v1.3) in the sensitive clean mode to obtain a gene presence/absence list that included plasmid
encoded genes (68). We assessed pairwise nucleotide similarity of the two strains with pyANI
(v0.2.12; (69). Multilocus sequence types were confirmed using mlst (v2.11;
https://github.com/tseemann/mlst) (70). Elements associated with resistance, including known
single nucleotide polymorphisms (SNPs), were detected with AMRfinderPlus (version 3.11.26
with database version 2023-11-15) (71). Two-proportion z-tests on two-tailed hypotheses were
used to determine if gene content was significantly different in the two strains, accounting for
the difference in genome size (Table S1). The annotated reference genomes used in this study
can be found in the NCBI BioProject ID PRJNA1214285.
399
Measuring resistance
Broad resistance profiles of the two strains were determined using Sensitire plates
(ThermoFisher, GN3F) following the manufactures protocols. Minimum inhibitory concentration
(MIC) assays were performed to measure susceptibility levels more accurately to Tigecycline
(TGC, Sigma 220620-09-7) following modified CLSI methods (9, 72). More detailed methods can
be found in the supplement.
406
RNA extraction and purification
Cultures for transcriptomic analysis of the ancestor strains were seeded from individual
colonies in biological triplicate into 5mL M9plus media. Population cultures were started
directly from the freezer stock into 5mL M9plus to avoid unnecessary outgrowths. Dilutions of
the cultures into M9plus with or without TGC treatment at 0.06 µg/mL were grown to mid-late
exponential phase for RNA extraction. RNA was extracted following a modified TRIzol
(Invitrogen Cat. No. 15596026) protocol, purified using the Invitrogen PureLink® RNA Mini Kit
(Thermo Cat. No. 12183025) and treated twice with DNase. See supplemental methods for
more details.
416
RNA sequencing and analysis

Full details for RNA sequencing and analysis can be found in the supplemental methods.
Briefly, once RNA was purified and satisfied quality control measures, we performed ribosomal
RNA depletion using the RiBO-COP rRNA Depletion Kit for Gram-Negative Bacteria (Lexogen
Cat. No. 126.96). Library generation was done using Lexogen RNA sequencing kits, following
manufacturers protocols. Sequencing was performed in house on an Illumina NextSeq550 with
the corresponding High-Output Kit v2.5 75 cycles (Illumina 20024906). Raw reads are uploaded
to the NCBI BioProject ID PRJNA1214285. Reads were assessed for quality, processed, and
aligned to the reference genomes using kallisto (73). This analysis provided the gene abundance
metrics, measured in transcripts per million transcripts (TPM), which normalizes by both gene
length and sequencing depth. Differential expression analysis was done with DESeq2 (74).
Unless otherwise stated, cutoffs for significance were set to a false discovery rate adjusted p-
value < 0.05 and a magnitude of $\log_2$FoldChange > 1. RNA sequencing performed on three
biological replicates for each sample by treatment combination except for removal of two
replicates due to low sequencing coverage or outlier status (17978UN population 1 untreated
replicate A and AB5075-UW population 2 untreated replicate C), and averages of the replicates
are presented.
434
Experimental evolution with TGC
Our experimental design was adapted from previous studies (47, 51). Lineages were
inoculated from overnight culture with a 1:100 dilution into fresh media with or without TGC,
resulting in approx. 6.6 generations per day (75). The experiment was designed such that the
populations were not mutation limited (see supplemental methods). In the TGC treated
condition, the TGC level in the media was tailored to the initial susceptibility level of each
strain, starting at 0.5x the ancestral MIC (subinhibitory). Every three days, the concentration of
TGC in the media was doubled (Fig. S5). We propagated the lineages for 12 days, with the final
TGC media concentration at 4x the ancestral MIC, which crossed the clinical breakpoint for
resistance in both strains. Populations were periodically sampled and frozen in 9% DMSO at -
80°C, as well as pelleted and frozen at -20°C for DNA extraction.
446
Whole-population, whole-genome sequencing, and analysis
DNA was extracted from frozen cell pellets using the DNeasy blood and tissue kit for the
QIAcube (Qiagen, Hilden, Germany) with a 10-minute elution into nuclease-free water. Libraries
were prepared in-house as previously described (76) or using the plexWell™ kit following
manufacturer's directions (SeqWell PW096). Libraries were sequenced using an Illumina
NextSeq550 sequencer with a 300 cycle mid-output kit (Illumina 20024905). Raw reads are
uploaded to the NCBI BioProject ID PRJNA1214285. Reads were demultiplexed, trimmed, and
quality checked as described in the supplemental methods for the RNAseq reads. Breseq
(v0.35.0) was used for read mapping and variant calling (77) with subsequent filtering following

previously published rationale (51) with special attention to new junction calls. Filtering,
consolidating for allele frequencies, and plotting were done in RStudio (R v4.2.1) with the
packages ggplot2 (v3.4.2; https://CRAN.R-project.org/package=ggplot2) and tidyr (v1.3.0;
https://CRAN.R-project.org/package=tidyr).
460
Growth curves as measure of fitness
We use bacterial growth curves to measure absolute fitness of ancestral clonal samples
as well as to measure aggregate absolute fitness of evolved populations (78). Growth curves
were seeded to mimic the transfers of the evolution experiment. A large sample of the frozen
population (or ancestor) stock was inoculated, in biological triplicate, into M9plus media and
grown for 24h. Overnight cultures were then diluted 1:100 into fresh media and OD600
measured every 10 minutes for 24h. Fitness was measured, in technical triplicate, in plain
M9plus as well as in M9plus containing 0.06 µg/mL TGC. We use area under the curve (AUC) to
measure fitness of populations normalized by the AUC of their respective ancestor ($AUC_{evolved}$
$_{population}$/$AUC_{ancestor\ average}$). Analyses were done in RStudio (R v4.2.1) utilizing previously
published pipelines (79) (https://github.com/mjfritz/Growth_Curves_in_R).
472
Measurements of efflux expression and activity
Efflux activity of late exponential phase cultures was assessed using the ethidium
bromide efflux activity assay. Fluorescence intensity was measured at three minutes post
addition of ethidium bromide to washed cells with an excitation of 530 nm and emission of 600
nm. Quantitative reverse transcriptive PCR was used to measure expression of efflux pump
components. RNA extraction was done as previously described (supplemental methods). The
Power SYBR Green RNA-to-Ct 1-Step Kit (Applied Biosystems) was used for cDNA synthesis and
qPCR with custom primers (Table S2). Data were analyzed for ΔCt and ΔΔCt, normalized by rpoB
and ancestor, respectively, within strain background. Expression values are presented as 2^{-}
$^{normalized\ expression}$ such that higher values indicate increased expression. We used the main efflux
pump gene (adeB, G, and J) as a representative for pump expression (57, 80). More information
on assays of efflux activity and expression can be found in supplemental methods.
485

ACKNOWLEDGEMENTS
We thank all members of the Cooper Lab as well as our collaborators in the Combating
Antibiotic-Resistant Bacteria Interdisciplinary Research Units (NIH CARBIRU) for thoughtful
discussions and feedback. We are especially grateful for bioinformatic guidance from Emma G.
Mills and Nanami Kubota. We thank Michelle R. Scribner for direction on the initial project
design and assistance in uploading data to the NCBI BioProject. This work was supported by NIH
U19AI158076 and, in part, by PA CURES Grant #4100085725. ABR is also supported by NIH
1F31AI172279-01A1 as well as a Catalyst Award from the Pittsburgh Center for Evolutionary

Biology and Medicine (CEBaM). ASL is currently supported by the Ministry of Science,

Innovation and Universities through a Ramon y Cajal fellowship (RYC2022-037765-I).

<u>**FIGURES**</u>

500
501
Fig. 1: Transcriptional response to TGC is highly strain dependent and results in a shift toward
a more conserved transcriptional state. Counts of differentially expressed genes, upregulated
(*A*) or downregulated (*B*), in each strain in response to TGC treatment. Overlapping regions
represent genes that were similarly differentially expressed in both strains. Percentage of genes
that were significantly differentially expressed in one strain but are still encoded by both strains
(shared genome) is indicated, showing minimal involvement of accessory genes. (*C*) Each point
represents the abundance of one gene in each strain in that condition (left: untreated plain
M9plus media; right: TGC treated with 0.06 µg/mL TGC). Diagonal y = x line represents equal
gene abundance in both strains and R^2 linear regression fits to this line are indicated. Points are
colored based on within-strain significant differential expression upon TGC treatment. See
Animation S1 for an animation of these transcriptional responses, and Fig. S4 for accessory
gene transcriptomic responses and a version of this plot with colors separated.
514

515
Fig. 2: Strain specific patterns in fitness-resistance tradeoffs. Minimum inhibitory
concentration (MIC) to TGC was measured for ancestors and for day-12 TGC-evolved
populations. Relative fitness was calculated from growth curves as described in the methods.
Fitness of 1.00 (gray horizontal dotted line) indicates the population grew equally well to its
respective ancestor in that environment (untreated M9plus or TGC treatment at 0.06 µg/mL
TGC). Points represent mean across replicates and error bars show standard deviation of the
mean.

523
Fig. 3: **Whole-population genome sequencing reveals similar resistance mechanisms but**
different population dynamics between strains. Evolved mutation frequencies over time, with
line colors corresponding to affected gene and line patterns indicating mutation type.
Abbreviations: SNP = single nucleotide polymorphism; IS = insertion sequence, indel = insertion
or deletion, and junction = structural variant indicated by a new junction call. Multiple lines of
the same color indicate more than one unique mutation arose in that gene. (*A*) Mutations in
genes associated with the ribosome. (*B*) Mutations in genes associated with efflux pumps.

Fig. 4: Genetic distinctions of evolved efflux expression and activity (*A*) Representation of the RND efflux pump operons/regulators. All RND genes are highly conserved between the two strains (supplemental table 2), except that 17978UN does not encode *adeC.* Arrow size is scaled to the length of the gene. Inverted triangles depict mutations present on day 12, colored by the population in which they are found. Ancestral efflux activity (*B*) and expression (*C*) in treated (0.06 μg/mL TGC) or untreated states. (*D*) Efflux activity of the evolved populations when grown in the untreated state relative to respective ancestor. (*E*) Efflux expression in evolved populations normalized by the respective untreated ancestor. Points represent mean and error bars depict standard deviation. Statistically significant differences were determined with t-tests in B and C with the comparisons depicted by lines, and with Wilcoxon tests compared to an ancestral value set at 1 in D and E. Significance criteria: ns for $p > 0.05$, * for $p \leq 0.05$, ** for $p \leq 0.01$, *** for $p \leq 0.001$, and **** for $p \leq 0.0001$.

547
Fig. 5: Transcriptional response evolves to minimally react to TGC pressure. (*A*) Conserved
transcript abundance regardless of TGC treatment in all evolved populations. Each point
represents the abundance of one gene in each sample. Point position depicts expression in
each condition (untreated M9plus, or 0.06 µg/mL TGC). Top row shows ancestor and TGC-
evolved populations for 17978UN, and the bottom row is the same for AB5075-UW. Diagonal y
= x line represents equal gene abundance in both conditions and R^2 linear regression fits to this
line are indicated. (*B*) Principal component analysis showing that ancestral transcriptomes are
strain dependent, but TGC treatment produces a conserved shift. Evolved population
transcriptional adaptation is conserved between strains and indifferent to TGC treatment. Point
color indicates sample, shape indicates culture condition either untreated or 0.06 µg/mL TGC
treated, fill indicates either ancestral or evolved populations.
559
560
561
562

SUPPLEMENTAL ANIMATION

Animation S1: Tigecycline treatment induces a shift toward a more conserved transcriptional

response. Animation of main text figure 1, showing movement of genes from the untreated

state to the TGC treated state. See figure 1 for color key and legend.

<u>**SUPPLEMENTAL FIGURES**</u>

Fig. S1: Ancestral growth in treatment conditions. Growth was measured in minimal M9plus
media (Untreated) or M9plus with addition of 0.06 µg/mL TGC (TGC treated) with three
biological replicates (batches), each with three technical replicates. (*A*) Growth curves for
ancestors grown in different conditions. Solid line represents mean, and ribbons depict
standard deviation. (*B*) Area under the curve (AUC) quantifications of curves in A (used to
normalize evolved population growth in Fig. 2). (*C*) Treatment effect on ancestral growth. Y-axis
shows the percent growth defect caused by TGC treatment on each strain, normalized by
within-batch growth in plain media. For B and C, mean and standard deviation depicted by
point and error bars, p-values calculated from pairwise t-test.

586
587

Fig. S2: Transcriptional response to TGC is strain dependent. Volcano plots of differential
expression in 17978UN (*A*) and AB5075-UW (*B*). Points are colored based on significance status
and fold change; Labels are colored based on if the gene is significantly differentially expressed
in only 17978UN (red), in only AB5075-UW (blue), or significant in both strains (gray). Labels
were manually curated to label the strongest hits or hits that showcase differences between
the strains. RNAseq was done in triplicate for each strain x treatment. Genes were included as
significant in the counts/plots if they had an absolute value of log 2-fold change > 1 and a p-
value ≤ 0.05.

|log2FoldChange| > 1   **|log2FoldChange| > 2**

597
Fig. S3: Significance cut-off criteria does not affect amount of differential expression overlap between strains. All differentially expressed genes (both upregulated and downregulated) in response to TGC treatment were summed. Varying cut-offs for calling significance in p-value and in log 2-fold change were tested to determine if the patterns of differential expression overlap between strains would differ. The overlap in differentially expressed genes between strains slightly decreases as increasingly strict significance criteria are applied.

605
606
Fig. S4: Strain-dependent transcriptional response to TGC results in a shift toward a more
conserved transcriptional state. (*A*) Specific directional gene responses with genes colored
based on how that gene responds within-strain upon TGC pressure. The top panels show gene
abundances in untreated media and bottom panels show gene abundance in TGC treatment
(0.06 µg/mL). Only genes falling under the bin of directional response noted above the plots are
colored. All other genes, even those with significant TGC responses, are gray in these plots. (*B*)
Abundance plots of the accessory genes, using the same colors as in A and in main text figure 1.
See figure 1 legend for more details.
615

616
617 **Fig. S5: Experimental design and post-experiment assay timings.** Schematic for the design of
618 the experimental evolution. For in-depth design, see the methods section. Bottom panels
619 highlight when various phenotype/genotype assays were performed. Dark steps indicate TGC
620 concentration doubling every three days of the experiment. Faint bar on bottom represents
621 lineages propagated in M9plus media lacking antibiotic to assess media adaptations.
622

624
Fig. S6: Transcriptional response to TGC in evolved populations is minimal, especially in
lineages of AB5075-UW. A handful of genes significantly respond to TGC treatment (0.06
µg/mL), compared to growth in untreated media, in evolved populations of 17978UN (*top*), but
no genes significantly respond to TGC in any evolved population of AB5075-UW (*bottom*). All
samples are from day 12 of the TGC evolution experiment. Data is from averages of three
biological replicates for all sample x treatment combinations except for removal of two
replicates due to low sequencing coverage or outlier status (17978UN population 1 untreated
replicate A and AB5075-UW population 2 untreated replicate C). Differential expression
significance cutoffs: |LFC|>1 and p-value ≤ 0.05.
634
635

636 <u>**SUPPLEMENTAL TABLES**</u>
637 Tables S1-S4 can also be found at https://github.com/vscooper/tigecycline
638
639 **Table S1: Genomic comparisons between strains.** Summary of strain similarities and
640 differences in various genomic categorizations, including gene count in clusters of orthologous
641 group (COG) categories, total genes assigned COG categories, AMR finder found resistance
642 elements, and annotated insertion sequences. A proportional z-test was performed for all
643 appropriate comparisons with the relevant gene count as the proportion denominator.

		AB5075-UW	17978UN	p-value (proportional z-test)
	Cell cycle control, cell division, chromosome partitioning	34	37	1
	Cell motility	40	44	1
	Cell wall, membrane and envelope biogenesis	170	188	0.957
	Cytoskeleton	2	2	1
	Defense mechanisms	61	75	0.547
	Extracellular structures	36	39	1
	Intracellular trafficking, secretion, and vesicular transport	58	60	0.832
	Posttranslational modification, protein turnover, chaperones	115	128	0.94
	Signal transduction mechanisms	108	122	0.855
	Replication, recombination and repair	87	104	0.583
COG Categories	RNA processing and modification	1	2	1
	Transcription	251	287	0.619
	Translation, ribosomal structure and biogenesis	217	224	0.561
	Amino acid transport and metabolism	276	278	0.33
	Carbohydrate transport and metabolism	126	126	0.508
	Coenzyme transport and metabolism	155	170	1
	Energy production and conversion	164	170	0.654
	Inorganic ion transport and metabolism	165	191	0.619
	Lipid transport and metabolism	197	219	0.902
	Nucleotide transport and metabolism	84	84	0.61
	Secondary metabolites biosynthesis, transport and catabolism	55	68	0.553
	Function unknown	113	130	0.736
	General function prediction only	210	232	0.951
	Total number of genes with COG assignments	2725	2980	**1.55E-12**
	AMRfinder Plus found elements	19	6	**0.017**
	Annotated IS elements	14	14	1

Antibiotic	17978UN	AB5075-UW
Amikacin	≤ 8 µg/mL / S	32 µg/mL / I
Ampicillin	≤ 4 µg/mL / S	> 32 µg/mL / R
Ampicillin/sulbactam	≤ 4/2 µg/mL / S	16/8 µg/mL / I
Aztreonam	16 µg/mL / I	> 32 µg/mL / R
Cefazolin	> 32 µg/mL / R	> 32 µg/mL / R
Cefepime	≤ 4 µg/mL / S	> 32 µg/mL / R
Cephalothin	> 16 µg/mL / R	> 16 µg/mL / R
Meropenem	≤ 1 µg/mL / S	> 8 µg/mL / R
Ertapenem	≤ 2 µg/mL / S	> 16 µg/mL / R
Cefuroxime	16 µg/mL / R	> 32 µg/mL / R
Gentamicin	≤ 2 µg/mL / S	≤ 2 µg/mL / S
Ciprofloxacin	≤ 0.5 µg/mL / S	> 4 µg/mL / R
Piperacillin / tazobactam	≤ 16/4 µg/mL / S	> 128/4 µg/mL / R
Cefoxitin	32 µg/mL / R	> 32 µg/mL / R
Trimethoprim / sulfamethoxazole	> 4/76 µg/mL / R	≤ 0.5 µg/mL / S
Cefpodoxime	4 µg/mL / I	> 16 µg/mL / R
Ceftazidime	2 µg/mL / S	4 µg/mL / S
Tobramycin	≤ 4 µg/mL / S	≤ 4 µg/mL / S
Tigecycline*	**≤ 1 µg/mL / S**	**≤ 1 µg/mL / S**
Ticarcillin / clavulanic acid	≤ 16/2 µg/mL / S	> 64/2 µg/mL / R
Ceftriaxone	8 µg/mL / I	16 µg/mL / I
Tetracycline	≤ 0.5 µg/mL / S	≤ 0.5 µg/mL / S

Table S2: Resistance profiles for the laboratory reference strain, 17978UN, and the multi-drug resistant clinical reference strain, AB5075-UW. Susceptibility/resistance state was measured in duplicate for each strain to a variety of Gram-negative acting antibiotics via Sensititre plates #GN3F (ThermoFisher). Sensitive (S), intermediate (I), and Resistant (R) classifications were assigned using CLSI standards for Gram-negative bacteria. *Tigecycline was chosen for further use in this study and broth-dilution MICs were determined to differ 1-fold between the strains.

Protein	Length	Amino Acid Pairwise Identity (%)
AdeS	362	97.2
AdeR	248	100
AdeA	397	98.7
AdeB	1037	99.5
AdeC	466	*
AdeN	218	100
AdeI	417	100
AdeJ	1059	99.9
AdeK	485	100
AdeL	344	99.7
AdeF	407	99.5
AdeG	1060	99.9
AdeH	483	99.8

Table S3: Conservation of efflux proteins between 17978UN and AB5075-UW. Pairwise identity of amino acid sequences. * *adeC* is only encoded by AB5075-UW, but it is suggested that *adeK* can assume the function of *adeC* in 17978UN (Leus et. al., 2018). Importantly, all sites that will be mutated upon TGC adaptation are initially identical between the strains.

717 **Table S4: List of primers used for Q-RT-PCR.** Quantative, reverse-transcriptase PCR was
718 performed on the inner membrane pump of the three RND efflux pumps as documented in the
719 methods using primers with the characteristics listed here. The housekeeping gene *rpoB* was
720 used as a control.

Target	Direction	Sequence (5'-3')	Melt Temp (ºC)	Product Size	Source
rpoB	forward	TCCGCACGTAAAGTAGGAAC	57.6	155	Coyne et al. Antimicrobial Agents and Chemotherapy.
	reverse	ATGCCGCCTGAAAAAGTAAC	57		
adeB	forward	AAAGGTATTGGCTACGAGTGG	57.7	132	Lin et al. Frontiers in Microbiology. 2017
	reverse	TGCCCAGCTTTCATAGAGTG	57.6		
adeG	forward	CAAAGTCCAGCATTACCAGC	57.1	136	This study, geneious primer3
	reverse	CTGCCCCAGGTTATTCATCT	57		
adeJ	forward	TGACCAACACCTGTGAATGA	57	164	This study, geneious primer3
	reverse	GAAGATCAGGGTGTGGTCAT	56.9		

SUPPLEMENTAL DATA LEGENDS
There are three submissions for supplemental data, each a separate file that can be accessed
through https://github.com/vscooper/tigecycline
Please contact authors for additional information about supplemental data.
Data S1: Ancestral gene annotations and RNA sequencing results.
Data S2: Filtered results of whole-population, whole-genome sequencing throughout the
evolution experiment.
Data S3: RNA sequencing results of the evolved populations.
731
732

SUPPLEMENTAL METHODS

Supplemental methods citations found after main text citations.

Bacterial strains and growth conditions

Laboratory reference strain ATCC 17978UN (1, 2) and clinical reference strain AB5075-

UW (3, 4) were used throughout the study and grown under the same conditions. Unless

otherwise stated, bacterial cultures were grown in 5ml M9+ media, as previously described (5).

Briefly, M9+ media is a salts-buffered media with glucose (11.1mM) as the primary carbon

source. It contains 0.1 mM $CaCl_2$, 1 mM $MgSO_4$, 42.2 mM Na_2HPO_4, 22 mM KH2PO$_4$, 21.7 mM

NaCl, 18.7 mM NH_4Cl and is supplemented with 20 mL/L MEM essential amino acids (Gibco

11130051), 10 mL/L MEM nonessential amino acids (Gibco 11140050), and 1 mL each of trace

mineral solutions A, B, and C (Corning 25021–3 Cl). Cultures were incubated at 37°C with

shaking or on a roller-drum at approximately 250rpm.

Genomic comparison of the two strains

Genomes were annotated with bakta (v1.6.1, database v4.0) (6) and input to Panaroo

(v1.3) in the sensitive clean mode to obtain a gene presence/absence list that included plasmid

encoded genes (Data S1) (7). We assessed pairwise nucleotide similarity of the two strains with

pyANI (v0.2.12; (8). Multilocus sequence types were confirmed using mlst (v2.11;

https://github.com/tseemann/mlst) (9). Elements associated with resistance, including known

single nucleotide polymorphisms (SNPs), were detected with AMRfinderPlus (version 3.11.26

with database version 2023-11-15) (10). Two-proportion z-tests on two-tailed hypotheses were

used to determine if gene content in clusters of orthologous group (COGs) categories was

significantly different in the two strains, accounting for the difference in genome size (Table S1)

(11). The annotated reference genomes used in this study can be found in the NCBI BioProject

ID PRJNA1214285.

Measuring resistance

Broad resistance profiles of the two strains were determined using Sensititre plates

(ThermoFisher, GN3F), commercially available panels of 23 antibiotics relevant to Gram-

negative pathogens at dilutions spanning the clinical breakpoint for resistance as set by the

Clinical and Laboratory Standards Institute (CLSI) (12). Broth microdilution and inoculation of

the plates was performed following the manufacturer's instructions. Sensititre plates were

performed in biological triplicate for each strain.

Minimum inhibitory concentration (MIC) assays were performed to measure

susceptibility levels more accurately to tigecycline (TGC, Sigma 220620-09-7). MIC assays were

performed following modified CLSI methods (12). MIC results presented here are from MIC

assays performed in M9plus media. Reference MIC assays were performed in the standard

Mueller Hinton broth and showed little-to-no (maximum difference of 1-fold) difference to

those performed in M9plus media. Biological triplicate (multiple samples from the frozen

ancestors or populations) with technical triplicates for each assay were performed. We present

the concentration of antibiotic that inhibits 90% growth (IC$_{90}$), measured via optical density at

600nm, as the MIC for each antibiotic.

RNA extraction and purification

For transcriptome analysis of the ancestor strains, cultures were seeded from individual

colonies in biological triplicate into 5mL M9plus media. Population cultures were started

directly by inoculating a large sampling of freezer stock into 5mL M9plus, also in triplicate. All

cultures were grown over-night at 37°C and back diluted 1:100 into 25 mL pre-warmed M9plus

with or without 0.06 µg/mL TGC. In this way, each biological replicate over-night culture results

in a paired set of treated and untreated TGC culture for analysis, thus limiting the effect of

sampling bias for the populations. Cultures were grown until reaching mid-log phase (OD600 =

0.6 +/- 0.05), at which time cultures were pelleted, resuspended in TRIzol (Invitrogen Cat. No.

15596026) and stored at -80°C for subsequent extraction and purification.

RNA was extracted following a modified protocol (13) for TRIzol extraction with

purification using the Invitrogen PureLink® RNA Mini Kit. Extraction included one bead beating

step followed by TRIzol-chloroform extraction with ethanol precipitation. Purification was done

using RNA columns from the Invitrogen PureLink system with final elution into nuclease free

water. RNA samples were treated with DNase 1 off-column with additional DNase spike-in

during incubation (NEB DNase and buffer Cat. No. M0303S) as well as treated with DNase 1 on-

column (Qiagen Cat. No. 79254) during final purification steps on the Invitrogen columns. RNA

concentration and quality were assessed using nanodrop (all 260/280 ratios > 2). Pure RNA was

stored at -20°C for less than a week prior to library preparation and at -80°C for longer-term

storage. Step-by-step RNA protocols are available upon request.

RNA sequencing and analysis

RNA integrity, fragment distribution, and RIN[e] scores were assessed on Tapestation

using the High Sensitivity RNA ScreenTape, sample buffer, and ladder from Agilent Technologies

(Part Numbers: 5067 -5579, -5580, -5581). All RIN[e] scores satisfied subsequent ribo-depeletion

requirements for quality (the majority of RIN[e] scores were >8). rRNA depletion was performed

using the RiBO-COP rRNA Depletion Kit for Gram-Negative Bacteria (Lexogen Cat. No. 126.96)

and directly following the manufacturer's protocols with an initial RNA input of approximately

200ng. RNA libraries including ancestral strain RNAs were prepared using the CORALL Total

RNA-Seq Library Prep Kit (Lexogen Cat. No. 095) while the evolved populations were sequenced

with the RNA-Seq V2 Library Prep Kit with UDIs (Lexogen Cat. No. 171.96). Both library

preparations included the PCR add-on step to determine ideal cycle number (11 cycles) for

library amplification (Lexogen Cat. No. 020). The uniquely indexed samples for each library were

pooled based on fragment size and concentration and the pool was diluted to 4nM for

sequencing. Each library was denatured and loaded at a final concentration of 2pM with a 5%

phiX spike-in. Sequencing was performed on the Illumina NextSeq550 with the corresponding

High-Output Kit v2.5 75 cycles (Illumina 20024906). Raw reads are uploaded to the NCBI

BioProject ID PRJNA1214285.

Reads were demultiplexed using bcl2fastq (v2.17.1) and trimmed with trimmomatic

(v0.36, custom criteria as follows: LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15 MINLEN:36).

Read quality was confirmed with fastQC (v0.11.5) and kallisto (v0.48.0, flags as follows: --single

-l 315 -s 26) was used for pseudoalignment with custom made index files from the

concatenated bakta annotated fasta files previously described for each reference strain (14).

Sequencing coverage was satisfactory (>10^7 reads/sample, resulting in >200x coverage) for all

samples apart from one replicate of AB5075-UW evolved population 2 in the untreated

condition, which had unusually low coverage and was removed from further analysis. We also

removed one replicate of 17978UN evolved population 1 in the untreated condition. DESeq2

(v1.38.3; (15)) was used in RStudio (R v4.2.1) for differential expression analysis supplemented

with EnhancedVolcano (v1.16.0) for plotting. Transcript abundance is reported as transcripts

per million transcripts (TPM) which normalizes by both gene length and sequencing depth. TGC

imposed differential expression analyses were performed with guidance from established

tutorials (15, 16). In the results function for analysis of the DESeq Data Set, we specified to turn

off independent filtering and cooksCutoff to keep outliers and calls with low coverage in the

analyses, then the apeglm method was applied for log fold change shrinkage (17). Unless

otherwise stated, cutoffs for significance were set to a false discovery rate adjusted p-value <

0.05, with the Benjamini and Hochberg correction for multiple comparisons, and a magnitude

of log_2FoldChange > 1.

Experimental evolution with TGC

Our experimental design was adapted from previous studies (5, 18). Both strains were

resurrected from the freezer via streaking on LB agar plates. A single colony of each strain was

used to inoculate a 5mL overnight culture in M9+. After approximately 18 hours, the saturated

culture was used to inoculate multiple independent lineages for the evolution experiment. We

propagated five replicate lineages of each strain in each condition, either with or without TGC.

Lineages were inoculated from overnight culture (either the starting overnight, or the prior day)

with a 1:100 dilution into fresh media with or without TGC. 1:100 dilutions from saturated

overnight cultures results in approximately 6.6 generations per day (19). The large population

size combined with the 1% bottleneck, mutation rate, and doubling time of *A. baumannii*

provides a large mutation supply, with a probability >1 that each site can see a mutation (19).

Lineages were propagated in the presence and absence of TGC, resulting in two

conditions. The no drug lineages were used to control for adaptations to the media and

experimental conditions. In the TGC treated condition, the TGC level in the media was tailored

to the initial susceptibility level of each strain, starting at 0.5x the ancestral MIC (subinhibitory).

Every three days, the concentration of TGC in the media was doubled (Fig. S5). We propagated

the lineages for 12 days, with the final TGC media concentration at 4x the ancestral MIC, which

crossed the clinical breakpoint for resistance in both strains. On days 0, 1, 3, 4, 6, 7, 9, 10, 12,

corresponding to days before and after antibiotic level increases, the populations were stocked

for later assays and sequencing (frozen in 9% DMSO at -80°C, and a cell pellet from 1mL of

culture was frozen at -20°C).

Whole-population, whole-genome sequencing, and analysis

DNA was extracted from frozen cell pellets using the DNeasy blood and tissue kit for the

QIAcube (Qiagen, Hilden, Germany) with a 10-minute elution into nuclease-free water. Libraries

were prepared in-house as previously described (20, 21) or using the plexWell™ kit following

manufacturer's directions (SeqWell PW096). Libraries were sequenced using an Illumina

NextSeq550 sequencer with a 300 cycle mid-output kit (Illumina 20024905). Reads were

demultiplexed, trimmed, and quality checked as described in the supplemental methods for the

RNAseq reads. Raw reads are uploaded to NCBI BioProject ID PRJNA1214285. Breseq (v0.35.0)

was used for read mapping and variant calling (22) with subsequent filtering following

previously published rationale (18) with special attention to new junction calls. Filtering,

consolidating for allele frequencies, and plotting were done in RStudio (R v4.2.1) with the

packages ggplot2 (v3.4.2; https://CRAN.R-project.org/package=ggplot2) and

tidyr (v1.3.0; https://CRAN.R-project.org/package=tidyr).

Growth curves as measure of fitness

We use bacterial growth curves to measure absolute fitness of ancestral clonal samples

as well as to measure aggregate absolute fitness of evolved populations (23). Growth curves

were seeded to mimic the transfers of the evolution experiment. A large sample of the frozen

population (or ancestor) stock was inoculated, in biological triplicate, into M9plus media and

grown for 24h. Overnight cultures were then diluted 1:100 into fresh media and OD600

measured every 10 minutes for 24h. Fitness was measured, in technical triplicate, in plain

M9plus as well as in M9plus containing 0.06 µg/mL TGC. We use area under the curve (AUC) to

measure fitness of populations normalized by the AUC of their respective ancestor ($AUC_{evolved}$

$_{population}$/$AUC_{ancestor\ average}$). Analyses were done in RStudio (R v4.2.1) utilizing previously

published pipelines (24) (https://github.com/mjfritz/Growth_Curves_in_R).

Quantitative reverse transcriptive PCR

To measure evolved efflux pump expression, we extracted RNA as described above for

ancestors and day-12 TGC evolved populations grown in the presence (0.06 µg/mL TGC,

ancestral cultures only) or absence of TGC (for ancestral and evolved populations). 50ng of
purified RNA was added per reaction for Q-RT-PCR according to manufacturer's instructions for
the Power SYBR Green RNA-to-Ct 1-Step Kit (Applied Biosystems). Reactions were performed in
biological and technical triplicate and run on a QuantStudio3 thermocycler using the default
threshold detection. Reactions lacking reverse transcriptase mix were included as controls for
DNA contamination. Data were analyzed for ΔCt and ΔΔCt, normalized by *rpoB* and ancestor,
respectively, within strain background. Expression values are presented as $2^{-normalized\ expression}$
such that higher values indicate increased expression. We used the main efflux pump gene
(*adeB, G, and J)* as a representative for pump expression (25, 26). Primers were designed with
Geneious to amplify approximately 150 base pair regions and are listed in Table S4.
895
<u>Ethidium bromide efflux activity assay</u>
Efflux activity was measured with the ethidium bromide uptake assay loosely based on
previously published methods (27). Ancestral cultures were grown overnight in M9plus, either
with or without addition of 0.06 μg/mL TGC, but not until saturation, and 500 μL was pelleted,
washed twice, and resuspended with PBS. Efflux activity of day-12 TGC-evolved populations
was measured only in untreated M9plus. A black flat-bottom 96-well plate (Corning ref. 3925)
was seeded with 90 μL of resuspended cells and 10 μL of 10 μg/mL ethidium bromide
(Invitrogen by Thermo Fisher Scientific ref. 15585-011). Fluorescence intensity was measured at
three minutes post addition of ethidium bromide with an excitation of 530 nm and emission of
600 nm. High efflux activity would result in a low fluorescence intensity as less ethidium
bromide is left within the cells binding to DNA. We present efflux activity as 1/RFU such that
higher values indicate more efflux. Efflux activity of the day-12 TGC-evolved populations when
grown in the untreated state relative to respective ancestor was calculated as
$1/(RFU_{query}/RFU_{untreated\ ancestor})$. Efflux activity was measured in biological triplicate with at least
three technical replicates each.
911
912
913

Main Text Citations

1.   B. Aslam, *et al.*, Antibiotic resistance: a rundown of a global crisis. *Infect. Drug Resist.* **11**, 1645–1658 (2018).

2.   M. Lässig, V. Mustonen, A. M. Walczak, Predicting evolution. *Nat. Ecol. Evol.* **1**, 77 (2017).

3.   T. Gabaldón, Nothing makes sense in drug resistance except in the light of evolution. *Curr. Opin. Microbiol.* **75**, 102350 (2023).

4.   A. C. Palmer, R. Kishony, Understanding, predicting and manipulating the genotypic evolution of antibiotic resistance. *Nat. Rev. Genet.* **14**, 243–248 (2013).

5.   M. Travisano, J. A. Mongold, A. F. Bennett, R. E. Lenski, Experimental tests of the roles of adaptation, chance, and history in evolution. *Science* **267**, 87–90 (1995).

6.   J. Plucain, *et al.*, Contrasting effects of historical contingency on phenotypic and genomic trajectories during a two-step evolution experiment with bacteria. *BMC Evol. Biol.* **16**, 86 (2016).

7.   R. C. MacLean, A. San Millan, The evolution of antibiotic resistance. *Science* **365**, 1082–1083 (2019).

8.   Z. D. Blount, R. E. Lenski, J. B. Losos, Contingency and determinism in evolution: Replaying life's tape. *Science* **362** (2018).

9.   A. Santos-Lopez, *et al.*, The roles of history, chance, and natural selection in the evolution of antibiotic resistance. *eLife* **10** (2021).

10.   L. Garoff, *et al.*, Population Bottlenecks Strongly Influence the Evolutionary Trajectory to Fluoroquinolone Resistance in Escherichia coli. *Mol. Biol. Evol.* **37**, 1637–1646 (2020).

11.   A. Wong, Epistasis and the evolution of antimicrobial resistance. *Front. Microbiol.* **8**, 246 (2017).

12.   R. C. MacLean, Predicting epistasis: an experimental test of metabolic control theory with bacterial transcription and translation. *J. Evol. Biol.* **23**, 488–493 (2010).

13.   Z. D. Blount, A case study in evolutionary contingency. *Stud. Hist. Philos. Biol. Biomed. Sci.* **58**, 82–92 (2016).

14.   G. J. Vermeij, Historical contingency and the purported uniqueness of evolutionary innovations. *Proc Natl Acad Sci USA* **103**, 1804–1809 (2006).

15.　Y. Wang, *et al.*, Benefit of transferred mutations is better predicted by the fitness of
recipients than by their ecological or genetic relatedness. *Proc Natl Acad Sci USA* **113**,
5047–5052 (2016).

16.　S. Dionisi, J. S. Huisman, Digest: The importance of genetic background in bacterial
evolution. *Evolution* **78**, 593–594 (2024).

17.　N. Filipow, S. Mallon, S. Shewaramani, R. Kassen, A. Wong, The impact of genetic
background during laboratory evolution of Pseudomonas aeruginosa in a cystic fibrosis-
like environment. *Evolution* **78**, 566–578 (2024).

18.　S. Cisneros-Mayoral, L. Graña-Miraglia, D. Pérez-Morales, R. Peña-Miller, A. Fuentes-
Hernández, Evolutionary history and strength of selection determine the rate of
antibiotic resistance adaptation. *Mol. Biol. Evol.* **39** (2022).

19.　K. J. Card, T. LaBar, J. B. Gomez, R. E. Lenski, Historical contingency in the evolution of
antibiotic resistance after decades of relaxed selection. *PLoS Biol.* **17**, e3000397 (2019).

20.　T. N. Batarseh, S. N. Batarseh, A. Rodríguez-Verdugo, B. S. Gaut, Phenotypic and
Genotypic Adaptation of Escherichia coli to Thermal Stress is Contingent on Genetic
Background. *Mol. Biol. Evol.* **40** (2023).

21.　K. M. Flynn, T. F. Cooper, F. B.-G. Moore, V. S. Cooper, The environment affects epistatic
interactions to alter the topology of an empirical fitness landscape. *PLoS Genet.* **9**,
e1003426 (2013).

22.　F. Baquero, *et al.*, Evolutionary pathways and trajectories in antibiotic resistance. *Clin.*
Microbiol. Rev. **34**, e0005019 (2021).

23.　C.-R. Lee, *et al.*, Biology of Acinetobacter baumannii: Pathogenesis, Antibiotic Resistance
Mechanisms, and Prospective Treatment Options. *Front. Cell. Infect. Microbiol.* **7**, 55
(2017).

24.　M. Touchon, *et al.*, The genomic diversification of the whole Acinetobacter genus:
origins, mechanisms, and consequences. *Genome Biol. Evol.* **6**, 2866–2882 (2014).

25.　A. Iovleva, *et al.*, Carbapenem-Resistant Acinetobacter baumannii in U.S. Hospitals:
Diversification of Circulating Lineages and Antimicrobial Resistance. *MBio* **13**, e0275921
(2022).

26.　M. J. Dorman, N. R. Thomson, "Community evolution" - laboratory strains and pedigrees
in the age of genomics. *Microbiology (Reading, Engl)* **166**, 233–238 (2020).

27.   M. Del Mar Cendra, E. Torrents, Differential adaptability between reference strains and
976   clinical isolates of Pseudomonas aeruginosa into the lung epithelium intracellular
977   lifestyle. *Virulence* **11**, 862–876 (2020).

978   28.   G. Makke, *et al.*, Whole-Genome-Sequence-Based Characterization of Extensively Drug-
979   Resistant Acinetobacter baumannii Hospital Outbreak. *mSphere* **5** (2020).

980   29.   C. H. Camargo, *et al.*, Clonal spread of ArmA- and OXA-23-coproducing Acinetobacter
981   baumannii International Clone 2 in Brazil during the first wave of the COVID-19
982   pandemic. *J. Med. Microbiol.* **71** (2022).

983   30.   M. Marazzato, *et al.*, Genetic Diversity of Antimicrobial Resistance and Key Virulence
984   Features in Two Extensively Drug-Resistant Acinetobacter baumannii Isolates. *Int. J.
985   Environ. Res. Public Health* **19** (2022).

986   31.   M. R. Galac, *et al.*, A Diverse Panel of Clinical Acinetobacter baumannii for Research and
987   Development. *Antimicrob. Agents Chemother.* **64** (2020).

988   32.   T. van Opijnen, S. Dedrick, J. Bento, Strain Dependent Genetic Networks for Antibiotic-
989   Sensitivity in a Bacterial Pathogen with a Large Pan-Genome. *PLoS Pathog.* **12**, e1005869
990   (2016).

991   33.   L. Rizzetto, *et al.*, Strain dependent variation of immune responses to A. fumigatus:
992   definition of pathogenic species. *PLoS ONE* **8**, e56651 (2013).

993   34.   J. Davies, D. Davies, Origins and evolution of antibiotic resistance. *Microbiol. Mol. Biol.
994   Rev.* **74**, 417–433 (2010).

995   35.   A. Kaul, C. Souque, M. Holland, M. Baym, Genomic resistance in historical clinical isolates
996   increased in frequency and mobility after the age of antibiotics. *BioRxiv* (2025)
997   https://doi.org/10.1101/2025.01.16.633422.

998   36.   A. P. Hendry, Prediction in ecology and evolution. *Bioscience* (2023)
999   https://doi.org/10.1093/biosci/biad083.

37.   K. E. Erickson, P. B. Otoupal, A. Chatterjee, Transcriptome-Level Signatures in Gene
Expression and Gene Expression Variability during Bacterial Adaptive Evolution. *mSphere*
2 (2017).

38.   C. D. M. Wijers, *et al.*, Identification of Two Variants of Acinetobacter baumannii Strain
ATCC 17978 with Distinct Genotypes and Phenotypes. *Infect. Immun.* **89**, e0045421
(2021).

39.   P. Baumann, M. Doudoroff, R. Y. Stanier, A study of the Moraxella group. II. Oxidative-
negative species (genus Acinetobacter). *J. Bacteriol.* **95**, 1520–1541 (1968).

40.   A. C. Jacobs, *et al.*, AB5075, a Highly Virulent Isolate of Acinetobacter baumannii, as a
Model Strain for the Evaluation of Pathogenesis and Antimicrobial Treatments. *MBio* **5**,
e01076-14 (2014).

41.   L. A. Gallagher, *et al.*, Resources for Genetic and Genomic Analysis of Emerging Pathogen
Acinetobacter baumannii. *J. Bacteriol.* **197**, 2027–2035 (2015).

42.   N. D. Greer, Tigecycline (Tygacil): the first in the glycylcycline class of antibiotics. *Proc*
(Bayl Univ Med Cent) **19**, 155–161 (2006).

43.   X. Hua, *et al.*, Novel tigecycline resistance mechanisms in Acinetobacter baumannii
mediated by mutations in adeS, rpoB and rrf. *Emerg. Microbes Infect.* **10**, 1404–1417
(2021).

44.   K. Lucaßen, *et al.*, Prevalence of RND efflux pump regulator variants associated with
tigecycline resistance in carbapenem-resistant Acinetobacter baumannii from a
worldwide survey. *J. Antimicrob. Chemother.* **76**, 1724–1730 (2021).

45.   X. Hua, Q. Chen, X. Li, Y. Yu, Global transcriptional response of Acinetobacter baumannii
to a subinhibitory concentration of tigecycline. *Int. J. Antimicrob. Agents* **44**, 337–344
(2014).

46.   L. Gao, X. Ma, Transcriptome Analysis of Acinetobacter baumannii in Rapid Response to
Subinhibitory Concentration of Minocycline. *Int. J. Environ. Res. Public Health* **19** (2022).

47.   A. Santos-Lopez, C. W. Marshall, M. R. Scribner, D. J. Snyder, V. S. Cooper, Evolutionary
pathways to antibiotic resistance are dependent upon environmental structure and
bacterial lifestyle. *eLife* **8** (2019).

48.   T. Vogwill, M. Kojadinovic, R. C. MacLean, Epistasis between antibiotic resistance
mutations and genetic background shape the fitness effect of resistance across species
of Pseudomonas. *Proc. Biol. Sci.* **283** (2016).

49.   T. Vogwill, R. C. MacLean, The genetic basis of the fitness costs of antimicrobial
resistance: a meta-analysis approach. *Evol. Appl.* **8**, 284–295 (2015).

50.   K. J. Card, J. A. Jordan, R. E. Lenski, Idiosyncratic variation in the fitness costs of
tetracycline-resistance mutations in Escherichia coli. *Evolution* **75**, 1230–1238 (2021).

51.   M. R. Scribner, A. Santos-Lopez, C. W. Marshall, C. Deitrick, V. S. Cooper, Parallel
Evolution of Tobramycin Resistance across Species and Environments. *MBio* **11** (2020).

52.   K. B. Harris, K. M. Flynn, V. S. Cooper, Polygenic adaptation and clonal interference
1039   enable sustained diversity in experimental Pseudomonas aeruginosa populations. *Mol.*
1040   *Biol. Evol.* (2021) https://doi.org/10.1093/molbev/msab248.

1041   53.   Q. Chen, *et al.*, Decreased susceptibility to tigecycline in Acinetobacter baumannii
1042   mediated by a mutation in trm encoding SAM-dependent methyltransferase. *J.*
1043   *Antimicrob. Chemother.* **69**, 72–76 (2014).

1044   54.   S. Sharma, V. Kaushik, M. Kulshrestha, V. Tiwari, Different Efflux Pump Systems in
1045   Acinetobacter baumannii and Their Role in Multidrug Resistance. *Adv. Exp. Med. Biol.*
1046   **1370**, 155–168 (2023).

1047   55.   I. V. Leus, *et al.*, Substrate Specificities and Efflux Efficiencies of RND Efflux Pumps of
1048   Acinetobacter baumannii. *J. Bacteriol.* **200** (2018).

1049   56.   S. Raza, *et al.*, Molecular Characterization of Resistance-Nodulation-cell Division Efflux
1050   Pump Genes in Multidrug-Resistant Acinetobacter baumannii. *J. Glob. Infect. Dis.* **13**,
1051   177–179 (2021).

1052   57.   S. Coyne, P. Courvalin, B. Périchon, Efflux-mediated antibiotic resistance in
1053   Acinetobacter spp. *Antimicrob. Agents Chemother.* **55**, 947–953 (2011).

1054   58.   W. Huo, *et al.*, Immunosuppression broadens evolutionary pathways to treatment
1055   failure during *Acinetobacter baumannii* pneumonia. *BioRxiv* (2021)
1056   https://doi.org/10.1101/2021.04.07.438861.

1057   59.   N. Harmand, R. Gallet, G. Martin, T. Lenormand, Evolution of bacteria specialization
1058   along an antibiotic dose gradient. *Evol. Lett.* **2**, 221–232 (2018).

1059   60.   L. Praski Alzrigat, D. L. Huseby, G. Brandis, D. Hughes, Resistance/fitness trade-off is a
1060   barrier to the evolution of MarR inactivation mutants in Escherichia coli. *J. Antimicrob.*
1061   *Chemother.* **76**, 77–83 (2021).

1062   61.   B. R. Levin, V. Perrot, N. Walker, Compensatory mutations, antibiotic resistance and the
1063   population genetics of adaptive evolution in bacteria. *Genetics* **154**, 985–997 (2000).

1064   62.   E. M. Darby, V. N. Bavro, S. Dunn, A. McNally, J. M. A. Blair, RND pumps across the genus
1065   Acinetobacter: AdeIJK is the universal efflux pump. *Microb. Genom.* **9** (2023).

1066   63.   S. Gaiarsa, *et al.*, Can Insertion Sequences Proliferation Influence Genomic Plasticity?
1067   Comparative Analysis of Acinetobacter baumannii Sequence Type 78, a Persistent Clone
1068   in Italian Hospitals. *Front. Microbiol.* **10**, 2080 (2019).

1069   64.   M. S. Wright, S. Mountain, K. Beeri, M. D. Adams, Assessment of Insertion Sequence
1070   Mobilization as an Adaptive Response to Oxidative Stress in Acinetobacter baumannii
1071   Using IS-seq. *J. Bacteriol.* **199** (2017).

1072   65.   B. H. Good, I. M. Rouzine, D. J. Balick, O. Hallatschek, M. M. Desai, Distribution of fixed
1073   beneficial mutations and the rate of adaptation in asexual populations. *Proc Natl Acad*
1074   *Sci USA* **109**, 4950–4955 (2012).

1075   66.   M. M. Desai, D. S. Fisher, Beneficial mutation selection balance and the effect of linkage
1076   on positive selection. *Genetics* **176**, 1759–1798 (2007).

1077   67.   O. Schwengers, *et al.*, Bakta: rapid and standardized annotation of bacterial genomes via
1078   alignment-free sequence identification. *Microb. Genom.* **7** (2021).

1079   68.   G. Tonkin-Hill, *et al.*, Producing polished prokaryotic pangenomes with the Panaroo
1080   pipeline. *Genome Biol.* **21**, 180 (2020).

1081   69.   L. Pritchard, R. H. Glover, S. Humphris, J. G. Elphinstone, I. K. Toth, Genomics and
1082   taxonomy in diagnostics for food security: soft-rotting enterobacterial plant pathogens.
1083   *Anal. Methods* **8**, 12–24 (2016).

1084   70.   K. A. Jolley, M. C. J. Maiden, BIGSdb: Scalable analysis of bacterial genome variation at
1085   the population level. *BMC Bioinformatics* **11**, 595 (2010).

1086   71.   M. Feldgarden, *et al.*, AMRFinderPlus and the Reference Gene Catalog facilitate
1087   examination of the genomic links among antimicrobial resistance, stress response, and
1088   virulence. *Sci. Rep.* **11**, 12728 (2021).

1089   72.   C. CLSI, *M100: Performance Standards for Antimicrobial Susceptibility Testing*, 32nd Ed.
1090   (Clinical and Laboratory Standards Institute, 2022).

1091   73.   N. L. Bray, H. Pimentel, P. Melsted, L. Pachter, Near-optimal probabilistic RNA-seq
1092   quantification. *Nat. Biotechnol.* **34**, 525–527 (2016).

1093   74.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
1094   RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

1095   75.   A. Santos-Lopez, *et al.*, Evolved resistance to a novel cationic peptide antibiotic requires
1096   high mutation supply. *Evol. Med. Public Health* **10**, 266–276 (2022).

1097   76.   C. B. Turner, C. W. Marshall, V. S. Cooper, Parallel genetic adaptation across
1098   environments differing in mode of growth or resource availability. *Evol. Lett.* **2**, 355–367
1099   (2018).

77.   D. E. Deatherage, J. E. Barrick, Identification of mutations in laboratory-evolved
microbes from next-generation sequencing data using breseq. *Methods Mol. Biol.* **1151**,
165–188 (2014).

78.   A. Alonso-Del Valle, *et al.*, Variability of plasmid fitness effects contributes to plasmid
persistence in bacterial communities. *Nat. Commun.* **12**, 2653 (2021).

79.   M. J. Muraski, *et al.*, Adaptation to Overflow Metabolism by Mutations That Impair tRNA
Modification in Experimentally Evolved Bacteria. *MBio* **14**, e0028723 (2023).

80.   F. Lin, *et al.*, Molecular Characterization of Reduced Susceptibility to Biocides in Clinical
Isolates of Acinetobacter baumannii. *Front. Microbiol.* **8**, 1836 (2017).
1109
1110

1111   Supplemental Methods Citations

1112   1.   C. D. M. Wijers, *et al.*, Identification of Two Variants of Acinetobacter baumannii
1113   Strain ATCC 17978 with Distinct Genotypes and Phenotypes. *Infect. Immun.* **89**,
1114   e0045421 (2021).

1115   2.   P. Baumann, M. Doudoroff, R. Y. Stanier, A study of the Moraxella group. II.
1116   Oxidative-negative species (genus Acinetobacter). *J. Bacteriol.* **95**, 1520–1541
1117   (1968).

1118   3.   A. C. Jacobs, *et al.*, AB5075, a Highly Virulent Isolate of Acinetobacter baumannii,
1119   as a Model Strain for the Evaluation of Pathogenesis and Antimicrobial Treatments.
1120   *MBio* **5**, e01076-14 (2014).

1121   4.   L. A. Gallagher, *et al.*, Resources for Genetic and Genomic Analysis of Emerging
1122   Pathogen Acinetobacter baumannii. *J. Bacteriol.* **197**, 2027–2035 (2015).

1123   5.   A. Santos-Lopez, C. W. Marshall, M. R. Scribner, D. J. Snyder, V. S. Cooper,
1124   Evolutionary pathways to antibiotic resistance are dependent upon environmental
1125   structure and bacterial lifestyle. *eLife* **8** (2019).

1126   6.   O. Schwengers, *et al.*, Bakta: rapid and standardized annotation of bacterial
1127   genomes via alignment-free sequence identification. *Microb. Genom.* **7** (2021).

1128   7.   G. Tonkin-Hill, *et al.*, Producing polished prokaryotic pangenomes with the Panaroo
1129   pipeline. *Genome Biol.* **21**, 180 (2020).

1130   8.   L. Pritchard, R. H. Glover, S. Humphris, J. G. Elphinstone, I. K. Toth, Genomics and
1131   taxonomy in diagnostics for food security: soft-rotting enterobacterial plant
1132   pathogens. *Anal. Methods* **8**, 12–24 (2016).

1133   9.   K. A. Jolley, M. C. J. Maiden, BIGSdb: Scalable analysis of bacterial genome
1134   variation at the population level. *BMC Bioinformatics* **11**, 595 (2010).

1135   10.   M. Feldgarden, *et al.*, AMRFinderPlus and the Reference Gene Catalog facilitate
1136   examination of the genomic links among antimicrobial resistance, stress response,
1137   and virulence. *Sci. Rep.* **11**, 12728 (2021).

1138   11.   R. L. Tatusov, M. Y. Galperin, D. A. Natale, E. V. Koonin, The COG database: a tool
1139   for genome-scale analysis of protein functions and evolution. *Nucleic Acids Res.*
1140   **28**, 33–36 (2000).

1141   12.   C. CLSI, *M100: Performance Standards for Antimicrobial Susceptibility Testing*,
1142   32nd Ed. (Clinical and Laboratory Standards Institute, 2022).

13.   A. C. Stephens, L. R. Thurlow, A. R. Richardson, Mechanisms Behind the Indirect
Impact of Metabolic Regulators on Virulence Factor Production in Staphylococcus
aureus. *Microbiol. Spectr.* **10**, e0206322 (2022).

14.   N. L. Bray, H. Pimentel, P. Melsted, L. Pachter, Near-optimal probabilistic RNA-seq
quantification. *Nat. Biotechnol.* **34**, 525–527 (2016).

15.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

16.   T. Bliek, J. Chouaref, F. van der Kloet, J. Szkodon, M. Galland, *Introductionto RNA-*
seq lesson (2020) (July 1, 2024).

17.   A. Zhu, J. G. Ibrahim, M. I. Love, Heavy-tailed prior distributions for sequence count
data: removing the noise and preserving large differences. *Bioinformatics* **35**, 2084–
2092 (2019).

18.   M. R. Scribner, A. Santos-Lopez, C. W. Marshall, C. Deitrick, V. S. Cooper, Parallel
Evolution of Tobramycin Resistance across Species and Environments. *MBio* **11**
(2020).

19.   A. Santos-Lopez, *et al.*, Evolved resistance to a novel cationic peptide antibiotic
requires high mutation supply. *Evol. Med. Public Health* **10**, 266–276 (2022).

20.   M. Baym, *et al.*, Inexpensive multiplexed library preparation for megabase-sized
genomes. *PLoS ONE* **10**, e0128036 (2015).

21.   C. B. Turner, C. W. Marshall, V. S. Cooper, Parallel genetic adaptation across
environments differing in mode of growth or resource availability. *Evol. Lett.* **2**, 355–
367 (2018).

22.   D. E. Deatherage, J. E. Barrick, Identification of mutations in laboratory-evolved
microbes from next-generation sequencing data using breseq. *Methods Mol. Biol.*
1151, 165–188 (2014).

23.   A. Alonso-Del Valle, *et al.*, Variability of plasmid fitness effects contributes to
plasmid persistence in bacterial communities. *Nat. Commun.* **12**, 2653 (2021).

24.   M. J. Muraski, *et al.*, Adaptation to Overflow Metabolism by Mutations That Impair
tRNA Modification in Experimentally Evolved Bacteria. *MBio* **14**, e0028723 (2023).

25.   S. Coyne, P. Courvalin, B. Périchon, Efflux-mediated antibiotic resistance in
Acinetobacter spp. *Antimicrob. Agents Chemother.* **55**, 947–953 (2011).

26. F. Lin, *et al.*, Molecular Characterization of Reduced Susceptibility to Biocides in
Clinical Isolates of Acinetobacter baumannii. *Front. Microbiol.* **8,** 1836 (2017).

27. Z. R. Lonergan, *et al.*, An Acinetobacter baumannii, Zinc-Regulated Peptidase
Maintains Cell Wall Integrity during Immune-Mediated Nutrient Sequestration. *Cell*
Rep. **26**, 2009-2018.e6 (2019).

1179