fCUT&Tag-Seq: An Optimized CUT&Tag-based Method for High-Resolution Profiling of Histone Modifications and Chromatin-Binding Proteins in Fungi

Haiting Wang [1^]**, Yongjunlin Tan**[2^]**, Jiayue Ma**[1^]**, Jie Yang**[1,3]**, Mengran Liu**[1,3]**, Peng Jiang**[1,3]**, Shan Lu**[4]**, Haoxue Xia**[5]**, Guangfei Tang**[5]**, Wende Liu**[5]**, Hui-Shan Guo**[1*]**, Chun-Min Shan**[1*]

Affiliations:

1. Department of Agri-microbiomics and Biotechnology, State Key Laboratory of Microbial Diversity and Innovative Utilization, Institute of Microbiology, Chinese Academy of Sciences, Beijing 100101, China

2. Shanxi Agricultural University, Jinzhong 030801, China.

3. University of Chinese Academy of Sciences, Beijing 100049, China

4. College of Agriculture, Guangxi Key Laboratory of Sugarcane Biology, Guangxi University, Nanning 530004, China

5. State Key Laboratory for Biology of Plant Diseases and Insect Pests, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing 100193, China

^: contribute equally.

* E-mail: shancm@im.ac.cn (C.-M.S.); guohs@im.ac.cn (H.-S.G.);

E-mail and ORCID for all authors:

Haiting Wang > wanght@im.ac.cn, 0000-0003-3477-0217

Yongjunlin Tan > t17835090800@163.com, 0009-0007-1288-0147

Jiayue Ma > majiajia0529@gmail.com, 0009-0002-4644-1998

Jie Yang > yangjie@im.ac.cn, 0009-0003-9335-5803

Mengran Liu > liumr@im.ac.cn, 0009-0009-9946-6524

Peng Jiang > jiangp@im.ac.cn, 0009-0008-8151-4682

Shan Lu > lushan@gxu.edu.cn, 0000-0001-5368-896x

Haoxue Xia > xiahaoxue97@163.com, 0009-0000-4077-675X

Guangfei Tang > tangguangfei@caas.cn, 0000-0002-6999-6999

Wende Liu > liuwende@caas.cn, 0000-0002-5570-1395

Hui-Shan Guo > guohs@im.ac.cn, 0000-0002-3057-9303

Chun-Min Shan > shancm@im.ac.cn, 0000-0002-7080-5221

Abstract

Histone modifications and chromatin-binding proteins play crucial roles in

regulating gene expression in eukaryotes, with significant implications for fungal

pathogenicity and development. However, profiling these modifications or proteins

across the genome in fungi remains challenging due to the technical limitations of the

traditional, widely used ChIP-Seq method. Here, we present an optimized

CUT&Tag-Seq protocol (fCUT&Tag-Seq) specifically designed for filamentous fungi

and dimorphic fungi. Our approach involves the preparation of protoplasts and

nuclear extraction to enhance antibody accessibility, along with formaldehyde

crosslinking to improve protein-DNA binding efficiency. We then successfully applied

fCUT&Tag-Seq to accurately profile multiple histone modifications like H3K9me3,

H3K27me3, H3K4me3, and H3K18ac, across different plant pathogenic or model

fungal species, including *Verticillium dahliae*, *Neurospora crassa*, *Fusarium*

graminearum, and *Sporisorium scitamineum*, showing good signal-to-noise ratios,

reproducibility, and detection sensitivity. Furthermore, we extended this method to

profile chromatin-binding proteins, such as the histone acetyltransferase Gcn5. This

study establishes fCUT&Tag-Seq as a robust and useful tool for fungal epigenetic

research, enabling detailed exploration of chromatin dynamics and advancing our

understanding of fungal gene regulation, development, and pathogenicity.

Keywords: fCUT&Tag-Seq; Fungi; Histone modifications; Chromatin-binding

proteins; Plant pathogen

Impact Statement

We developed a faster, low□input method to study how genes are turned on and off in

fungi, even in tough species that are difficult to analyze with standard methods. Our

new approach, named fCUT&Tag-Seq, requires only 10,000 cells and can be

completed in just two days. It delivers clearer, more reliable results and has already

been successfully applied to multiple fungi, including crop pathogens. By revealing

the molecular "control switches" that govern fungal development and virulence, we

expect it will accelerate basic research and help identify new targets for controlling

destructive plant diseases.

Introduction

Chromatin structure and modifications are essential in regulating gene

expression in eukaryotic organisms. The basic unit of chromatin, the nucleosome,

consists of DNA wrapped around histone octamers composed of core histones H2A,

H2B, H3, and H4[1]. The N-terminal tails of these histones can undergo various

post-translational modifications (PTMs), such as methylation, acetylation,

phosphorylation, ubiquitination, sumoylation, and ADP-ribosylation[2, 3]. These

modifications can alter chromatin structure and impact vital biological processes, such

as gene expression, DNA repair, and chromosome organization[4, 5].

Among these modifications, histone methylation is critical in regulating gene

expression. For instance, trimethylation of histone H3 at lysine 9 (H3K9me3) and

lysine 27 (H3K27me3) is highly associated with heterochromatin formation and gene

silencing. In contrast, trimethylation of histone H3 at lysine 4 (H3K4me3), which

typically occurs at transcription initiation sites, is commonly enriched near gene

promoters and is associated with active transcription[6-8]. Furthermore,

chromatin-binding proteins, such as histone acetyltransferases and transcription

factors, play crucial roles in modifying histones or binding to promoters, thereby

regulating gene expression, development, stress responses, and pathogenicity in

eukaryotes, including fungi[9, 10].

Traditional methods for studying histone modifications and chromatin-binding

proteins have relied heavily on Chromatin Immunoprecipitation (ChIP) techniques,

developed in the mid-1980s. The subsequent advent of ChIP, followed by

high-throughput sequencing (ChIP-Seq), enables comprehensive identification of

histone modifications or protein-binding sites across the whole genome. While

ChIP-Seq has become a foundational approach for genome-wide chromatin analysis,

it still has several limitations. The method requires large amounts of input material,

may suffer from epitope masking, and often produces high background noise with low

signal-to-noise ratios[11-14]. These challenges are particularly pronounced when

studying filamentous fungi, where rigid cell walls and secondary metabolites reduce

antibody accessibility and affect the efficiency of immunoprecipitation[15].

The recent development of Cleavage Under Targets and Tagmentation

(CUT&Tag) offers a more efficient method for profiling chromatin features[16-18].

This technique employs a tethered Tn5 transposase, directed by specific antibodies, to

simultaneously cleave DNA at target sites and integrate sequencing adapters.

CUT&Tag-Seq offers several advantages over traditional ChIP-Seq, including

higher-resolution mapping with lower cell input requirements (as few as 60 cells) and

reduced background noise[19, 20]. While ChIP-seq remains the benchmark method

for most chromatin studies, CUT&Tag offers unique advantages for studies that

require live cell analysis or involve non-culturable samples, thereby providing a

complementary approach to address the technical limitations of traditional methods.

While CUT&Tag has been successfully applied to mammalian cells, plant cells, and

yeast, its application in filamentous fungi has been limited due to technical challenges

posed by the fungal cell wall.

In this study, we optimized a fungal CUT&Tag-Seq method (fCUT&Tag-Seq)

for pathogenic/model fungi (e.g., *Verticillium dahliae*, *Neurospora crassa*),

overcoming cell wall challenges via protoplast preparation and nuclear extraction.

Formaldehyde crosslinking improved detection of low-abundance chromatin-binding

proteins. Applied to histone modifications (H3K9me3, H3K4me3, etc.) and *SsGcn5*,

our method yields high-quality data with low cell input, outperforming ChIP-Seq in

signal-to-noise ratio and reproducibility in pathogenic fungi. fCUT&Tag-Seq

streamlines fungal epigenetics research, reducing time and sample requirements while

enhancing chromatin analysis.

Results

Optimization and Implementation of fCUT&Tag-Seq Method for Fungal Systems

We developed an optimized fCUT&Tag-Seq specifically adapted for fungal

systems, building upon the traditional approach using the Vazyme Hyperactive

Universal CUT&Tag Assay Kit for Illumina Pro (Cat. TD904). Our optimization

addresses the unique challenges posed by fungal cell architecture while maintaining

the high sensitivity and specificity characteristic of CUT&Tag methodology. The

modified protocol incorporates several key steps for fungal applications **(Fig. 1).** To

overcome the barrier presented by the fungal cell wall, we added a controlled

enzymatic digestion step to generate protoplasts, followed by gentle nuclear

extraction to preserve chromatin integrity. The isolated nuclei are then specifically

captured by Concanavalin A-coated magnetic beads, which provide a stable platform

for subsequent reactions while minimizing sample loss.

The workflow proceeds through several precisely controlled steps **(Fig. 1)**: (1)

Nuclear membrane permeabilization is enhanced through carefully optimized

digitonin treatment, facilitating antibody access while maintaining nuclear structural

integrity. (2) Primary antibodies specific to the target histone modifications are

introduced, followed by secondary antibodies that serve to amplify the detection

signal. (3) The pA/G-Tn5 transferase complex is recruited to the antibody-bound

locations through protein A/G interactions. (4) Upon Mg^{2+} activation, the tethered Tn5

simultaneously cleaves adjacent DNA and integrates sequencing adapters, enabling

efficient library preparation through subsequent PCR amplification. After library

construction, next-generation sequencing and data analysis can be performed to map

the distribution pattern of histone modifications.

fCUT&Tag-Seq Demonstrates Superior Quality Control Metrics and Target

Enrichment Compared to ChIP-Seq in *V. dahliae*

To evaluate if we could use fCUT&Tag-Seq for profiling histone modifications

in filamentous fungi, we first selected the plant pathogenic fungus *V. dahliae* as a

model system. This pathogenic fungus, known for its broad host range, causes severe

wilt, yellowing, and vascular browning in infected plants, often resulting in significant

crop losses[21]. Given the known roles of H3K9me3 and H3K27me3 in gene

silencing and pathogenicity[2, 22-24], we focused our analysis on these two

modifications. To enhance our understanding of *V. dahliae*'s genetic background, we

sequenced the whole genome of *V. dahliae* V592 strain using the Pacbio HiFi

platform **(Fig. S1)**. Using the genome of this specific strain as the reference for

mapping can help avoid errors introduced by inter-species genomic differences and

also capture species-specific epigenetic modifications of genes, thereby providing

more accurate information for the following analyses.

To evaluate the efficiency of our optimized fCUT&Tag-Seq method for profiling

histone modifications in filamentous fungi, we conducted a comparative analysis in *V.*

dahliae with other established methods. To validate the importance of protoplast

extraction, we initially performed CUT&Tag experiments using spores from both

wild-type and H3K9me3 methyltransferase mutant (Vd*Δkmt1*) strains to assess the

genome-wide distribution of H3K9me3. However, this approach failed to generate a

sufficient enrichment signal **(Fig. S2A)**. When comparing our approach with

previously reported protocols specially developed for fungi (such as CUT&RUN for

Candida albicans and CUT&Tag for *Saccharomyces pombe*), we attempted

spheroplasting *V. dahliae* hyphae using 1.6 mg/mL Zymolyase[15, 25], incubating

samples at 30°C for 30 min. However, this approach was unsuccessful as no

spheroplasting occurred, with mycelial structures remaining intact **(Fig. S2B)**.

Similarly, the application of the standard CUT&Tag protocol, designed for human

cells, without prior protoplast extraction, yielded insufficient data[16]. Importantly,

performing CUT&Tag directly on isolated protoplasts without nuclear extraction also

failed to produce enrichment signals **(Fig. S2A)**, demonstrating that the nuclear

extraction step is essential for successful CUT&Tag profiling in *V. dahliae*. Through

further optimization, we determined that although using as low as 1×10^4 cells was

sufficient to show acceptable enrichment, $2.5 \times 10\square$ cells provided optimal data **(Fig.**

S2C). Sequencing depths of approximately 1 Gb were sufficient to achieve the

desired resolution **(Fig. S2D)**. These data collectively demonstrate the critical

importance of protoplast preparation and nuclear extraction in our optimized

fCUT&Tag-Seq protocol, enabling robust and high-quality profiling of histone

modifications in filamentous fungi.

We then compared traditional ChIP-Seq data from the SRA database (Home -

SRA - NCBI) and fCUT&Tag-Seq datasets based on multiple quality control metrics.

Initial quality control analysis of fCUT&Tag-Seq libraries revealed characteristic

nucleosomal ladder patterns through agarose gel electrophoresis, indicating successful

fragmentation and library preparation **(Fig. 2A)**. Fragment length distribution analysis

further confirmed the expected size distribution pattern typical of nucleosome-sized

fragments **(Fig. 2B)**, validating the technical success of our protocol. Comparative

analysis between fCUT&Tag-Seq and traditional ChIP-Seq demonstrated several key

advantages of our optimized method. Most notably, fCUT&Tag-Seq exhibited a

significantly higher percentage of reads mapped to peak regions and lower duplication

rates (8.73%) compared to ChIP-Seq (88.45%), indicating substantially improved

library complexity and data utilization efficiency **(Fig. 2C and 2D)**. Following

alignment to the genome and subsequent filtering, we conducted a comprehensive

analysis of quality metrics using ChIPQC. The fCUT&Tag-Seq data showed superior

Signal Space Distance (SSD) scores, reflecting enhanced enrichment efficiency.

Furthermore, the Fraction of Reads in Peaks (FRiP) consistently exceeded 7.8% in

fCUT&Tag-Seq samples, with H3K9me3 modification showing particularly improved

signal-to-noise ratios compared to ChIP-Seq **(Fig. 2D)**. And fCUT&Tag-Seq

identified a greater signal intensity, indicating higher signal-to-noise **(Fig. 2E)**. Peak

calling analysis revealed comparable patterns between fCUT&Tag-Seq and ChIP-Seq

data, validating the biological relevance of our results. These comprehensive quality

metrics demonstrate that our fCUT&Tag-Seq protocol not only meets standard quality

benchmarks but also offers superior performance. The improved signal-to-noise ratio

was particularly evident in genome-wide coverage plots, where fCUT&Tag-Seq

showed more distinct enrichment patterns and clearer peak boundaries **(Fig. 2F)**. We

have also performed strictly side-by-side comparisons of fCUT&Tag and ChIP-Seq in

V. dahliae. The results showed that ChIP-Seq failed to show signal enrichment, while

the fCUT&Tag assay yielded a strong H3K9me3 signal at the repeat region **(Fig.**

S2E). These comprehensive quality metrics demonstrate that our fCUT&Tag-Seq

protocol not only meets standard quality benchmarks but also offers superior

performance compared to traditional ChIP-Seq methods in terms of data quality,

enrichment efficiency, and technical reproducibility in *V. dahliae*. The improved

metrics suggest that this method provides a more reliable and efficient approach for

studying histone modifications in *V. dahliae*.

fCUT&Tag-Seq Method Efficiently Profiles Histone Modifications in *V. dahliae*

Following validation of our optimized protocol, we applied fCUT&Tag-Seq to

investigate the genome-wide distribution of two key repressive histone modifications,

H3K9me3 and H3K27me3, in *V. dahliae*. To evaluate the specificity of our method,

we analyzed both wild-type strains and their corresponding methyltransferase mutants

(Vd*Δkmt1* and Vd*Δezh2)*. We incorporated spike-in controls for data normalization

and used IgG antibodies as negative controls to account for background signal.

Genome-wide profiling revealed robust enrichment of both H3K9me3 and

H3K27me3 modifications in wild-type strains **(Fig. 3A and 3B)**. H3K9me3 is a

repressive mark mainly localized in constitutive heterochromatic regions, while

H3K27me3 is a repressive mark responsible for facultative gene silencing, usually

localized in euchromatin. In contrast, control experiments using non-specific IgG

antibodies showed minimal signal across the genome, confirming the specificity of

our approach **(Fig. S3A)**. Importantly, we observed significantly reduced levels of

H3K9me3 and H3K27me3 in their respective methyltransferase mutants, Vd*Δkmt1*

and Vd*Δezh2*, consistent with the loss of these enzymatic activities **(Fig. 3A and 3B)**.

Analysis of the genomic distribution of these modifications revealed distinct

patterns. Genomic feature distribution showed H3K9me3 peaks were predominantly

located in gene promoter regions and distal intergenic spaces in constitutive

heterochromatin, while H3K27me3 was mostly located in the gene promoter region in

euchromatin **(Fig. 3C)**. These distribution patterns align with the known roles of these

modifications in transcriptional regulation. A comparison of signal intensities between

fCUT&Tag-Seq and ChIP-Seq data demonstrated superior enrichment in

fCUT&Tag-Seq experiments, particularly across gene bodies and their 3-kb flanking

regions **(Fig. S3B)**.

To assess the reproducibility of our modified method, we performed correlation

analyses between biological replicates. fCUT&Tag-Seq demonstrated excellent

reproducibility, with Pearson correlation coefficients of 0.99 and 1.0 for H3K9me3

and H3K27me3 modifications, respectively. These values exceeded those obtained

from traditional ChIP-Seq experiments (0.91 and 0.89), highlighting the enhanced

reliability of our optimized method **(Fig. 3D)**.

These results collectively demonstrate that our fCUT&Tag-Seq protocol enables

highly specific and reproducible mapping of histone modifications in *V. dahliae*,

providing a robust tool for investigating chromatin-based regulation in this important

plant pathogen.

fCUT&Tag-Seq Method Shows Broad Applicability Across Model Filamentous

Fungi

To assess whether our fCUT&Tag-Seq protocol could be extended beyond *V.*

dahliae, we tested it on two additional filamentous fungi: *N. crassa* and *F.*

graminearum. Both species serve as valuable model organisms for studying fungal

biology and pathogenicity, making them ideal candidates for validating broad

applicability. Following the same experimental workflow, we focused on histone

modifications H3K9me3 in *N. crassa*, and H3K4me3 in *F. graminearum*.

N. crassa, a member of the Ascomycota phylum, is widely recognized as a model

organism due to its ease of cultivation and haploid genetics, which facilitate

straightforward genetic analysis. Over the past few decades, it has been instrumental

in studying various epigenetic modifications, including DNA methylation and gene

silencing mechanisms[26-28]. We examined the levels of H3K9me3 in both the

wild-type strain and the H3K9me3 methyltransferase mutant Nc*Δdim5*. Quality

control assessments, including agarose gel electrophoresis and sequencing data

analysis, confirmed that the fCUT&Tag-Seq libraries met the required standards **(Fig.**

4A and 4B). The quality metrics indicated excellent performance for the *N. crassa*

wild-type samples **(Fig. 4C)**. Notably, the peak distribution patterns observed were

consistent with those identified through ChIP-Seq, yet fCUT&Tag-Seq demonstrated

higher signal intensity **(Fig. 4D)**. The correlation between fCUT&Tag-Seq and

ChIP-Seq data reached 0.73 **(Fig. 4E)**, further validating the reliability of our method.

Additionally, biological replicates exhibited strong correlation, indicating high

reproducibility **(Fig. 4F)**. Comparative analysis revealed that H3K9me3 was

significantly enriched in the wild-type strain, while the Nc*Δdim5* mutant displayed

markedly lower modification levels **(Fig. 4G)**. As expected, H3K9me3 modifications

were predominantly located in distal intergenic regions and promoter regions **(Fig.**

4H).

Similar results were obtained for *F. graminearum*, the causative agent of

fusarium head blight in wheat[29]. In addition to analyzing H3K9me3 and

H3K27me3 modifications, we also assessed H3K4me3 levels, which are associated

with gene activation. The sequencing quality metrics of fCUT&Tag-Seq for *F.*

graminearum were superior to those of ChIP-Seq, with a greater percentage of reads

mapping to peak regions **(Fig. 5A and 5B)**. The Signal Space Distance (SSD) score

and the percentage of reads in peaks (Rip%) reached 21 and 46, respectively **(Fig. 5B)**.

These findings confirm the robustness of our method, as the wild-type strain exhibited

significant enrichment **(Fig. 5C)**. Notably, peak enrichment was predominantly

observed in the promoter regions (96.51%) **(Fig. 5D)**. The Pearson correlation

coefficient was 0.99, indicating a strong correlation between the sample groups **(Fig.**

5E).

Overall, these results demonstrate that our fCUT&Tag-Seq method is not only

effective for detecting histone modifications in *V. dahliae*, but also applicable to other

model fungi, including *N. crassa* and *F. graminearum*. This versatility underscores the

potential of fCUT&Tag-Seq as a powerful tool for studying epigenetic modifications

across a broad range of fungal species.

fCUT&Tag-Seq Method Enables Profiling of Chromatin-Binding Proteins

To extend the utility of our optimized fCUT&Tag-Seq protocol beyond histone

modifications, we adapted the method to profile chromatin-binding proteins, which

play crucial roles in transcriptional regulation. Given the typically lower abundance of

chromatin-binding proteins compared to histones, we incorporated an additional

formaldehyde crosslinking step before protoplast preparation. This modification

strengthens protein-chromatin interactions, thereby improving detection sensitivity.

Following fragmentation, we performed sequential decrosslinking, protease digestion,

and RNase treatment to ensure efficient DNA purification and library preparation **(Fig.**

6).

To validate this adapted fCUT&Tag-Seq method, we investigated the histone

acetyltransferase SsGcn5 in *S. scitamineum*, a dimorphic pathogenic fungus

responsible for sugarcane smut. Sugarcane smut is one of the most significant

diseases affecting global sugarcane production worldwide. The disease poses

substantial economic threats, highlighting the urgent need for effective prevention and

control strategies against *S. scitamineum*[30-32]. It has been reported that histone

acetylation may play a critical role in fungal pathogenesis[33-36]. To explore the

relationship between histone acetylation and fungal pathogenicity, we screened

possible histone acetyltransferase candidate genes and introduced a Flag epitope tag

into the histone acetyltransferase Gcn5 in two *S. scitamineum* strains with different

mating types (JG35 and JG36), using our previously reported method[37].

Quality control analysis of the resulting fCUT&Tag-Seq libraries revealed proper

DNA fragmentation patterns by agarose gel electrophoresis and expected nucleosomal

size distributions in sequencing reads **(Fig. 7A and 7B)**, confirming successful library

preparation. Comprehensive quality metrics and genome-wide profiling demonstrated

robust enrichment of SsGcn5-3Flag binding sites across the *S. scitamineum* genome

(Fig. 7C and 7D). Notably, comparative analysis revealed a strong correlation

between SsGcn5 binding sites and regions of H3K18 acetylation **(Fig. 7E)**, consistent

with the known catalytic activity of Gcn5. Feature distribution analysis showed that

SsGcn5 binding sites were predominantly located in promoter regions (85.93% of

total peaks) **(Fig. 7F)**, suggesting a primary role in transcriptional regulation. The

high reproducibility of our method was demonstrated by the near-perfect Pearson

correlation coefficient (approximately 1.0) between biological replicates **(Fig. 7G)**.

Our results confirmed that the fCUT&Tag-Seq method not only excels in

detecting histone modifications, but also effectively profiles chromatin-binding

proteins in fungi. By enhancing the detection of low-abundance proteins, this method

also provides a powerful tool for exploring chromatin-binding proteins in regulating

gene expression and chromatin dynamics across a broad range of fungal species.

Discussion

Over prolonged evolutionary periods, plant pathogenic fungi have developed

intricate strategies to penetrate and colonize their hosts. Precise control of pathogenic

gene expression is pivotal to these infection processes, influencing fungal

development, environmental stress responses, and virulence[38]. Emerging evidence

highlights the significance of epigenetic regulation, including histone methylation,

histone acetylation, and chromatin remodeling, in orchestrating transcriptional

reprogramming in plant pathogens[39-41]. However, until recently, the available tools

for epigenetic research in diverse fungal systems were still very limited, making

advanced investigations into fungal chromatin biology challenging.

In this study, we optimized and extended CUT&Tag-Seq to several important

fungal species, including *V. dahliae*, *F. graminearum*, *N. crassa*, and *S. scitamineum*.

Our fCUT&Tag-Seq method effectively addresses the unique obstacles posed by

fungal cell walls by integrating protoplast preparation and gentle nuclear extraction.

Additionally, a formaldehyde crosslinking step was introduced to strengthen

DNA-protein interactions, particularly beneficial for profiling low-abundance

chromatin-binding proteins.

Our data revealed that the fCUT&Tag-Seq method accurately and sensitively

detects histone modifications (H3K9me3, H3K27me3, H3K4me3, and H3K18ac) as

well as chromatin-binding proteins in these fungi. We also performed genome-wide

profiling of H3K9me3 and H3K27me3 distributions, revealing distinct

compartmentalization patterns characteristic of fungal epigenomes. Our analyses

demonstrate that H3K9me3 predominantly marks constitutive heterochromatin

regions, showing strong enrichment at pericentromeric and telomeric domains,

consistent with its role in silencing repetitive elements[42-44]. In contrast, H3K27me3

preferentially marks facultative heterochromatin, regulating transcriptionally dynamic

regions that modulate stress responses and developmental transitions[45, 46]. Its

reversible deposition enables rapid gene expression remodeling in response to

environmental cues, such as host infection or nutrient availability[47].

As epigenetic modifications play crucial functional roles during fungal infection

processes. Taking advantage of the compositional differences between plant and

fungal cell walls, we would be able to isolate fungal protoplasts in planta through

enzymatic digestion directly in the near future. Using the fCUT&Tag-Seq method, we

may profile fungal-specific epigenetic modifications during host infection, thereby

studying their critical associations with fungal pathogenicity. Notably, the minimal

cellular input requirement of this methodology (potentially as few as 10,000 cells)

may offer future applications for studying unculturable fungal species.

While recent studies have established robust ChIP-Seq protocols for model fungi

like *N. crassa* (e.g., datasets SRR1566112, SRR12229310) [48, 49], our comparative

analyses revealed measurable differences in H3K9me3/H3K27me3 detection

sensitivity and reproducibility between established methods and our fCUT&Tag-Seq

approach. More importantly, our method mainly lies in its consistency across

non-model fungi, where traditional ChIP-Seq is particularly challenging.

Although our fCUT&Tag-Seq method has demonstrated significant

improvements in data quality and sensitivity, it is still important to acknowledge the

limitations. Challenges remain in enriching low-abundance proteins and

chromatin-binding proteins located in heterochromatin regions, where accessibility to

antibodies is largely reduced. Future studies should focus on further optimizing

methods for relaxing heterochromatin to enhance the detection of these proteins.

Materials and Methods

Fungal Strains and Culture Conditions

The following fungal strains were used in this study: *V. dahliae* strain V592

(isolated from cotton in Xinjiang, China)[50, 51], *N. crassa* 87-3 and Nc*Δdim5*

(provided by Xiao Liu) [52], *S. scitamineum* strains JG35 and JG36 (provided by

Shan Lu) [53], and *F. graminearum* wild-type strain (provided by Guangfei Tang and

Haoxue Xia)[54]. All strains were maintained and cultured according to standard

protocols specific to each species.

Generation of Mutant and Tagged Fungal Strains

For *V. dahliae*, *VdKMT1* (*VDAG_07826*) and *VdEZH2* (*VDAG_00983*) deletion

strains (Vd*Δkmt1* and Vd*Δezh2*) were constructed as previously reported[55]. 1 kb

genomic sequences upstream and downstream of target genes were amplified from

V592 DNA with the following primer pairs: VdKMT1-A/ VdKMT1-a, VdKMT1-d/

VdKMT1-B, VdEZH2-A/ VdEZH2-a, VdEZH2-d/ VdEZH2-B (listed in Table S1)

using the Phanta Max Super-Fidelity DNA Polymerase (P505, Vazyme) and cloned

into the pGKO-HPT/pGKO-NAT vector *via* homologous recombination.

Sequence-verified vectors were then used to transform *V. dahliae* by *Agrobacterium*

tumefaciens-mediated transformation (ATMT) system. *V. dahliae* transformants were

selected on potato dextrose agar (PDA) plates supplemented with

5-fluoro-2'deoxyuridine (5FU) and hygromycin B / Nourseothricin. Putative

transformants were screened using PCR to confirm the successful deletion of the

target genes with primer pairs V1/Hpt-R and V2/Hpt-F. The primers used for

construction are listed in Table S1.

For *S. scitamineum*, the SsGcn5-Flag strain was constructed using homologous

recombination. The vector pEX1-GAP-Flag-HPT, carrying a Flag tag, was modified

from pEX1-GAP-HPT. Approximately 1.5 kb of upstream (5') and downstream (3')

flanking sequences near the target protein's stop codon (excluding the stop codon)

were amplified from the wildtype strains JG35 and JG36 using primer pairs:

SsGcn5-tg-A/SsGcn5-a and SsGcn5-d/SsGcn5-B, respectively (Table S1). Two

truncated fragments of the hygromycin resistance gene HPT (HPT-up and HPT-down)

were amplified from the pEX1-GAP-Flag-HPT plasmid with the primer pairs

HPT-LB-F/HPT-LB-R and HPT-RB-F/HPT-RB-R. Overlap PCR was employed to

fuse the upstream sequence with HPT-up (primers SsGcn5-tg-A/HPT-LB-R) and the

downstream sequence with HPT-down (primers HPT-RB-F/SsGcn5-B). The products

were introduced into *S. scitamineum* strains JG35 and JG36 via PEG-mediated

protoplast transformation. Transformants were selected on YEPSA medium

containing 150 μg/mL hygromycin B and verified by PCR using primer pairs

SsGcn5-C(V1)/HPT-LB-R and SsGcn5-D(V2)/HPT-RB-F.

fCUT&Tag-Seq for Histone Modification

To profile histone modifications using fCUT&Tag-Seq, protoplasts were

prepared by digesting mycelium (*V. dahliae, N. crassa,* and *F. graminearum*) or spores

(*S. scitamineum*) with 10 mL of enzyme solution (0.25 g VinoTaste®Pro enzyme in

12.5 mL 0.7 M NaCl for *V. dahliae*, *N. crassa,* and *S. scitamineum*; 0.1 g Diselease,

0.2 g Celulase, and 0.2 g Lysozyme were diluted with 0.7 M NaCl to 20 mL for *F.*

graminearum). *F. graminearum* was incubated at 30°C with shaking at 85 rpm, while

the others were incubated at 33°C at 60 rpm. Digestion times were 3 h for *V. dahliae*

mycelium, *F. graminearum* mycelium, and *N. crassa* mycelium, and 30 min for *S.*

scitamineum spores. Protoplasts were filtered through Miracloth (475855-1R; Merck),

and centrifuged for 10 min at 3800 rpm (*V. dahliae* and *N. crassa*), 2200 rpm (S.

scitamineum), or 5000 rpm (*F. graminearum*) at 25□. Protoplasts were resuspended in

10 mL of 0.7 M NaCl and then counted using a hemocytometer.

Nuclei were extracted from 1 million protoplasts per 1.5 mL tube by

centrifugation at 3000 rpm for 5 min at 25 °C, followed by resuspension in 100 μL

nuclear extract buffer (TD904; Vazyme), incubation on ice for 10 min, and a second

centrifugation at 2500 rpm for 6 min at 25 °C. The supernatant was discarded, and

nuclei were resuspended in 100 μL of wash buffer (TD904; Vazyme).

Nuclei were adsorbed by mixing activated Concanavalin A-coated magnetic

beads (TD904; Vazyme) with 250,000 resuspended nuclei and 75 μl of wash buffer

(TD904; Vazyme) in 200μL 8-row tubes, followed by incubation at room temperature

for 10 min with 2-3 inversions.

For antibody binding, tubes were placed on a magnetic rack, the supernatant was

discarded, and 50 μL of antibody buffer (TD904; Vazyme) was added to resuspend the

bead-nucleus complex. Primary antibodies (1 μL each) against H3K9me3 (Cat39161;

Active Motif), H3K27me3(Cat39155; Active Motif), or H3K4me3 (Cat39060; Active

Motif) were added, according to the ratio used. After mixing by inverting the tube,

centrifuge immediately, then place in a 4℃ refrigerator for overnight incubation

under static conditions (no rotation).

The next day, the beads were centrifuged briefly, placed on a magnetic rack, and

the supernatant was discarded. 0.5 μL of secondary rabbit antibody (Ab207; Vazyme)

in 49.5 μL of Dig wash buffer (792 μL Wash Buffer, 8 μL 5% Digitonin) was added

and incubated with rotation (9-11 rpm/min) at room temperature for 1h. Complexes

were washed three times with 200 μL of Dig wash buffer.

For transposase incubation, 2 μL of Hyperactive pA/G-Transposon (TD904;

Vazyme) in 98 μL Dig-300 buffer (100 μL 10 × Dig-300 Buffer, 2 μL 5% Digitonin,

20 μL 50 × protease inhibitor, 878 μL ddH$_2$O) was added to each reaction and

incubated with gentle rotation (9-11 rpm/min) at room temperature for 1 h.

Subsequently, after instantaneous centrifugation and placing on a magnetic rack,

supernatant was discarded and washed three times using 200 μL Dig-300 buffer.

For fragmentation, 10 μL of TruePrep Tagment Buffer L (TD904; Vazyme) in 40

μL Dig-300 buffer was added to each reaction and incubated at 37°C for 1 h.

DNA was extracted by adding 2 µL of 10% SDS (TD904; Vazyme) and 0.5 µL

of DNA spike-in (TD904; Vazyme), incubating at 55°C for 10 min, transferring

supernatant to a new tube, adding 50 µL activated DNA extract beads (TD904;

Vazyme), incubating for 20 min at room temperature with mixing, washing twice with

200 µL 1× B&W buffer (TD904; Vazyme), drying, and resuspending in 30 µL of

RNAase-free ddH$_2$O.

Libraries were constructed by PCR (50 µL reaction: 15µL bead-bound DNA, 25

µL 2× CUT&Tag Amplification Mix [TD904; Vazyme], 5µL N5XX Primer [TD202;

Vazyme], 5µL N7XX Primer [TD202; Vazyme]) following Illumina's adaptor pooling

guide. PCR conditions: 72°C for 3 min, 95°C for 3 min; then 9-20 cycles of 98°C for

10 s, 60°C for 5 s, 72°C for 1 min; hold at 4°C.

PCR products were purified by adding 100 µL VAHTS DNA clean beads (N411;

Vazyme), incubating for 5 min, washing twice with 200 µL 80% ethanol, drying,

resuspending in 22 µL RNase-free ddH□O, and incubating for 5 min before quality

control and sequencing.

fCUT&Tag-Seq for chromatin-binding proteins

To profile lower-abundance chromatin-binding proteins, a formaldehyde

crosslinking step was added. Approximately 1 g of fungal spores was fixed with 9 mL

of 0.1% formaldehyde for 8 min at room temperature, quenched with 1 mL of 2 M

glycine for 5 min at room temperature. The cross-linked cells were washed three

times with 20 mL of 0.7 M NaCl.

Protoplast formation, nuclei capture with ConA-beads, antibody incubation, and

pA/G-Tn5 transposome assembly followed the histone modification protocol. After

fragmentation, crosslinks were reversed by incubating overnight at 65°C with 3 μL 5

M NaCl. RNA was removed by RNase A treatment (37°C for 30 min), proteins by

Proteinase K (53°C for 1 h), and proteases by inhibitors (room temperature for 5 min).

Libraries were amplified by PCR as described.

ChIP-Seq

ChIP assays were performed as previously described[56-58]. Mycelia were

cross-linked with 1% formaldehyde solution (0.4 M sucrose, 10 mM Tris-HCl [pH

8.0], 1 mM EDTA, 1 mM PMSF, 1% formaldehyde) at room temperature for 15

minutes, with the flask inverted every 2–3 minutes. To stop cross-linking, glycine was

added to a final concentration of 0.2 M, followed by incubation for 5 minutes. Fixed

cells were then washed with 1× PBS, cryomilled in liquid nitrogen, and lysed in lysis

buffer (50 mM HEPES [pH 7.5], 137 mM NaCl, 1 mM EDTA, 0.1% SDS, 1% Triton

X-100, 0.1% sodium deoxycholate, 1 mM PMSF, 1 μg/mL leupeptin, 1 μg/mL

pepstatin A). Chromatin was sonicated to an average DNA fragment size of

~500–1000 bp. After sonication, samples were centrifuged at 10,000 ×g and 4 °C for

15 minutes to collect the supernatant. For immunoprecipitation, 2 μL of anti-H3K9ac

antibody (Cat. No. 39161; Active Motif) was incubated with aliquots of clarified cell

lysates (containing 2 mg protein each) at 4 °C overnight with rotation. After overnight

incubation, 50 μL of SS DNA/Protein G Sepharose (pre-treated three times with lysis

buffer) was added, followed by further incubation at 4 °C for 2 hours. Sepharose

beads were collected and washed once with 1 mL of each of the following buffers:

high-salt wash buffer (lysis buffer supplemented with 0.5 M NaCl), 1× LNDET (0.25

M LiCl, 1% NP40, 1% deoxycholate, 1 mM EDTA, 10 mM Tris [pH 8]), and 50 mM

Tris-HCl [pH 8], 1 mM EDTA. Immune complexes were eluted from the beads using

freshly prepared elution buffer (1% SDS, 0.1 M NaHCO₃) via two rounds of elution

at room temperature (250 μL per elution).

fCUT&Tag-Seq Data Analysis

Raw ChIP-Seq data were downloaded from the SRA database using the SRA

Toolkit (version 3.1.1). Raw data of ChIP-Seq and fCUT&Tag-Seq were first

processed using fastp (0.23.4)[59] to remove adapters and low-quality sequences. The

cleaned reads were then aligned to the spike-in sequences using BWA (0.7.18)[60].

After alignment, SAMtools (version 1.21)[61] was used to count the number of

successfully mapped reads. The sample with the fewest mapped reads was selected as

the baseline for spike-in normalization. The normalized data were then re-aligned to

the reference genome using BWA (0.7.18), with reads having a quality score below 20

being filtered out using samtools (1.21). The alignment results were then processed

using Picard (Picard Tools, developed by the Broad Institute) for sorting and

deduplication. Peak calling was performed using MACS2 (2.2.9.1)[62], with IgG

controls used to correct for background. The "-B" flag was employed to generate

histone modifications and control lambda .bdg files, which were subsequently

compared using the bdgcmp function in MACS2. ChIPQC (1.42.0, Bioconductor -

ChIPQC) was applied for quality control of the fCUT&Tag analysis. Utilize Spark.py

(2.6.2)[63] to generate a genomic browser view, incorporating the -gs parameter to

ensure all groups are autoscaled to a unified y-axis. Peak annotation was performed

using ChIPseeker (1.42.0)[64] to assign genomic features to the identified peaks.

Correlation analysis was carried out using the multiBigwigSummary and

plotCorrelation functions of deepTools (3.3.5)[65], while heatmaps were generated

using the computeMatrix and plotHeatmap functions of deepTools (3.3.5).

Figure legends

Figure 1. Schematic overview of the fCUT&Tag-Seq workflow for filamentous

fungi.

Illustration of the key steps in the optimized fCUT&Tag-Seq protocol in filamentous

fungi, covering key steps from protoplast preparation to sequencing. The method

involves digitonin permeabilization to enhance nuclear accessibility.

Figure 2. Comprehensive quality control analysis of fCUT&Tag-Seq compared to

ChIP-Seq in *Verticillum dahliae*.

A. Nucleosomal ladder pattern analysis of fCUT&Tag-Seq libraries by agarose gel

electrophoresis. Libraries were generated from H3K9me3 and H3K27me3 histone

modification profiling experiments in *Verticillum dahliae*. **B.** Fragment length

distribution analysis of sequencing reads from fCUT&Tag-Seq and ChIP-Seq

experiments. Green histograms represent H3K9me3 libraries (replicate 1 and replicate

2), while blue histograms show H3K27me3 libraries (replicate 1 and replicate 2). **C.**

Bar plot showing the fraction of reads in peaks (FRiP) for the fCUT&Tag-Seq and

ChIP-Seq datasets (replicates 1 and 2). **D.** Comparative analysis of key quality metrics

between fCUT&Tag-Seq and ChIP-Seq methods. Quality assessment encompasses

sequencing-level metrics (analyzed by Fastp), including total read counts, filtered read

retention, and PCR duplicate rates, along with enrichment-specific parameters

(evaluated by ChIPQC) measuring alignment efficiency (Map%), post-processing

duplicates (Dup%), the uniformity of reads covering the entire genome (SSD), and

Fraction of Reads in Peaks (Frip). **E.** Signal intensity profiles across peaks in

fCUT&Tag-Seq versus ChIP-Seq datasets: the upper panel displays the H3K9me3

modification, while the lower panel shows the H3K27me3 modification. **F.** Genomic

distribution analysis of identified peaks comparing fCUT&Tag-Seq and ChIP-Seq

methods. The displayed region is randomly selected. The upper panel represents the

H3K9me3 modification, and the lower panel shows the H3K27me3 modification.

Figure 3. Genome-wide profiling of histone modifications in *V. dahliae* using

fCUT&Tag-Seq.

A. Genome browser view of H3K9me3 distribution in wild-type and Vd*Δkmt1* strains.

B. Genome browser view of H3K27me3 distribution in wild-type and Vd*Δezh2*

strains. **C.** Comparative analysis of genomic feature distribution for H3K9me3 and

H3K27me3 modifications between fCUT&Tag-Seq and ChIP-Seq approaches. The

left panel shows the distribution of H3K9me3 peaks across different genomic

elements, and the right panel shows H3K27me3 peak distribution. **D.** Pearson

correlation analysis of signals between biological replicates for H3K9me3 and

H3K27me3 modifications in ChIP-Seq (replicate 1 and replicate 2) and

fCUT&Tag-Seq experiments (replicate 1 and replicate 2). Pearson correlations were

calculated using deepTools, with the multiBamSummary tool followed by

plotCorrelation. Read counts were divided into 1,000-bp bins across the genome for

the analysis.

Figure 4. Application of fCUT&Tag-Seq for histone modifications analysis in *N.*

***crassa*.**

A. Nucleosomal ladder pattern analysis of fCUT&Tag-Seq libraries by agarose gel

electrophoresis. Libraries were generated from two replicates of H3K9me3 profiling

experiments in *N. crassa*. **B.** Fragment length distribution analysis of sequencing

reads from fCUT&Tag-Seq experiments. Histograms display the expected

nucleosomal fragmentation patterns for two replicates. **C.** Comparative analysis

between fCUT&Tag-Seq and ChIP-Seq data for H3K9me3 modification in *N. crassa*.

D. Genome browser view of peak distribution analysis comparing fCUT&Tag-Seq

and ChIP-Seq results. The upper panel shows H3K9me3 enrichment in ChIP-Seq, and

the lower panel displays the modification distribution from fCUT&Tag-Seq. **E.**

Correlation matrix analysis between fCUT&Tag-Seq and ChIP-Seq data, indicating

strong agreement. Pearson correlations were calculated using deepTools

(multiBamSummary followed by plotCorrelation), based on read counts divided into

1000-bp bins across the genome. **F.** Pearson correlation analysis of H3K9me3 signals

between biological replicates in ChIP-Seq and fCUT&Tag-Seq. **G.** Genome browser

view of the H3K9me3 peaks accross representative genomic regions in wild-type and

Nc*Δdim5* strains in *N. crassa*. **H.** Feature distribution comparison of H3K9me3

modification between ChIP-Seq and fCUT&Tag-Seq.

Figure 5. Genome-wide profiling of H3K4me3 modification in *F. graminearum*

using fCUT&Tag-Seq analysis.

A. Fragment length distribution of sequencing libraries for fCUT&Tag-Seq. The left

panel shows the fragment length distribution for the H3K4me3 library in replicate 1,

and the right panel shows the distribution for replicate 2. **B.** Comparative analysis of

key quality metrics for ChIP-Seq and fCUT&Tag-Seq datasets. **C.** Genome browser

view of H3K4me3 modification peak distribution analysis comparing fCUT&Tag-Seq

and ChIP-Seq results. The genome regions were randomly selected. Green peaks

represent ChIP-Seq results, and blue peaks represent fCUT&Tag-Seq results. **D.** The

pie chart and UpSet plot illustrate the distribution of peaks across different genomic

features. The results showing that H3K4me3 is predominantly enriched in promoter

regions. Distinct colors represent different genomic features. **E.** Pearson correlation

coefficients between sample groups for ChIP-Seq and fCUT&Tag-Seq. The scatter

plot displays normalized read counts in 1,000-bp bins across the genome.

Figure 6. Schematic overview of the optimized fCUT&Tag-Seq method for

detecting chromatin-binding proteins.

The workflow diagram illustrates the key steps of the fCUT&Tag-Seq protocol for

analyzing chromatin-binding proteins in fungi, including protoplast preparation,

formaldehyde crosslinking, nuclear extraction, and subsequent fCUT&Tag-Seq

procedures.

Figure 7. Genome-wide profiling of SsGcn5 and its correlation with H3K18ac

modification in *S. scitamineum* using fCUT&Tag-Seq.

A. Quality assessment of fCUT&Tag-Seq libraries by agarose gel electrophoresis.

DNA fragment distributions are shown for two biological replicates of JG35 (lanes

1-2) and JG36 (lanes 3-4) mating types. **B.** Fragment length distribution analysis of

sequencing reads demonstrating expected nucleosomal patterns. Blue and pink

histograms represent Gcn5-Flag distributions in JG35 and JG36 strains, respectively.

C. Comprehensive quality metrics for fCUT&Tag-Seq datasets in DNA-binding

protein enrichment in JG35 and JG36 strains. **D.** Genome browser view of

SsGcn5-Flag binding sites identified by fCUT&Tag-Seq. Genome regions were

selected randomly. **E.** Correlation analysis between SsGcn5-Flag binding sites and

H3K18ac modification regions, showing the functional relationship between histone

acetyltransferase localization and its catalytic activity. **F.** Genomic feature distribution

analysis of SsGcn5-Flag peaks in JG35 and JG36 strains, showing the relative

enrichment across different genomic elements (promoters, gene bodies, intergenic

regions). **G.** Correlation analysis between biological replicates for SsGcn5-Flag

fCUT&Tag-Seq experiments.

Supporting information

Fig.S1 *V. dahliae* V592 genome assembly presentation. (A) The output quality of

sequencing data. Sample: sample name; SeqNum: number of sequences; SumBase:

total number of databases; N50Len: N50 length of sequencing data; N90Len: length

of sequencing data N90; MeanLen: mean length of sequencing sequence; MaxLen:

maximum length of sequencing sequence. (B) Assembly result statistic. Length:

length of the sequence after concatenation; GC (%): GC content of the concatenated

sequence. (C) Coding gene prediction results. (D) The CIRCOS plot of the genome.

Fig.S2 Optimization and comparative evaluation of the fCUT&Tag-Seq protocol

for *V. dahliae*. (A). Genome browser view of H3K9me3 distribution in wild-type and

Vd*Δkmt1* strains using nucleus, spores or protoplast. (B) Comparison of the state of

protoplasts extracted by different methods. The upper panel showed that no

spheroplasting state was observed, extracted by zymolyase described in the

CUT&RUN approach for *Candida albicans* and the CUT&Tag method for

Saccharomyces pombe, and the lower panel showed the protoplast state using the

optimized fCUT&Tag-Seq technique. The images were taken by a confocal

microscope (Leica TCS SP8). Scale bar: 25 µm. (C) Genome browser view of

H3K9me3 distribution in wild-type and Vd*Δkmt1* strains using different cell numbers.

(D) Genome browser view of H3K9me3 distribution in wild-type and Vd*Δkmt1*

strains using different sequencing volumes. (E) Genome browser view of H3K9me3

distribution in wild-type and Vd*Δkmt1* strains using fCUT&Tag-Seq and ChIP-Seq,

respectively. The Y-axis values in ChIP-seq tracks represent normalized read coverage

depth (in reads per million, RPM), calculated by scaling raw read counts to $10⬚$.

Fig.S3 Genome-wide profiling of nonspecific IgG signals and histone

modification patterns in *V. dahliae* using fCUT&Tag-Seq. (A) Genome browser

view of the H3K9me3, H3K27me3, and nonspecific IgG antibody signals across

representative genomic regions in the wild-type strain. Data were presented at the

same scale as specific antibody signals. (B) Heatmap analysis of H3K9me3 and

H3K27me3 signals near protein-coding genes in fCUT&Tag-Seq and ChIP-Seq. Scale

regions were 3,000 bp upstream of the translation starting site (TSS), 3,000 bp

downstream of the translation end site (TES), and a 1,000-bp region on the gene body.

The lengths were plotted using the computeMatrix and plotHeatmap tools in

deepTools.

Fig.S4 Original images of Figure 2A, Figure 4A and Figure 7A. (A) Original

images of Figure 2A. (B) Original images of Figure 4A. (C) Original images of Figure

7A.

Table.S1 Primers used in this study.

Acknowledgments

We thank Xiao Liu for providing *Neurospora crassa* strains. The work was supported

by the China National Key Research and Development Program (Grant no.

2024YFC3406000), the Chinese Academy of Sciences (CAS) Projects for Young

Scientists in Basic Research (Grant no. YSBR-080), the Strategic Priority Research

Program of Chinese Academy of Sciences (Grant no. XDA0450000) to C-M.S., and

the Young Scientists Fund of the National Natural Science Foundation of China

(Grant no. 32402338) to H.W.

The funder has not played any role in the study design, data collection and analysis,

the decision to publish, or the preparation of the manuscript.

Author Contributions:

H.W.: methodology, investigation, data curation, formal analysis, writing-original

draft, and writing—review and editing;

Y.T.: methodology, investigation, data curation, formal analysis, writing-original draft,

and writing—review and editing;

J.M.: biostatistics analysis, data curation, writing-original draft, and writing—review

and editing;

J.Y.: formal analysis, writing—original draft, and writing-review and editing;

M.L.: formal analysis, writing—original draft, and writing-review and editing;

P.J.: biostatistics analysis, writing-original draft, and writing—review and editing;

S.L.: supervision, investigation, resources, data curation;

H.X.: methodology, investigation, data curation, finish the experiments involving

Fusarium graminearum;

G.T.: methodology, investigation, data curation, finish the experiments involving

Fusarium graminearum;

W.L.: supervision, methodology, data curation, finish the experiments involving

Fusarium graminearum;

H-S.G.: supervision, project administration, methodology, resources, data curation,

writing-review and editing.

C-M.S.: conceptualization, supervision, project administration, methodology,

resources, data curation, writing-original draft, writing-review and editing.

All authors have read and agreed to the published version of the manuscript.

Data Availability

All ChIP-Seq data were obtained from the SRA database (Home - SRA - NCBI).

The ChIP-Seq data for H3K9me3 modification in *V. dahliae* were downloaded from

SRR10571946 and SRR10571947, respectively, and data for H3K27me3 modification

were downloaded from SRR10571948 and SRR10571949. For H3K9me3

modification in *N. crassa*, the ChIP-Seq data were downloaded from SRR1566112

and SRR12229310. The ChIP-Seq data for H3K4me3 modification in *F. graminearum*

were downloaded from SRR21677668 and SRR21677669. The reference genomes

used in the mapping process are as follows: PRJNA1212841 for *V. dahliae* V592,

GCF_000182925.2 for *N. crassa*, FgraminearumPH-1 for *F. graminearum* (available

at FungiDB), and GCA_001010845.1 for *S. scitamineum*. The fCUT&Tag-Seq data

and ChIP-Seq data used in this study are available in the NCBI BioProject database

under accession numbers PRJNA1213569, PRJNA1166993, and PRJNA1248591.

Ethics statement

No animal experiments nor human samples were involved in this study.

Conflicts of Interest

The authors declare that there is no conflict of interest.

References

1.   Luger K, Mäder AW, Richmond RK, Sargent DF, Richmond TJ. Crystal

structure of the nucleosome core particle at 2.8 A resolution. *Nature.* 1997;

389:251-260. https://doi.org/10.1038/38444.

2.   Kouzarides T. Chromatin modifications and their function. *Cell.* 2007;

128:693-705. https://doi.org/10.1016/j.cell.2007.02.005.

3.   Allis CD, Jenuwein T. The molecular hallmarks of epigenetic control. *Nat Rev*

Genet. 2016; 17:487-500. https://doi.org/10.1038/nrg.2016.59.

4.   Zhang Y, Sun Z, Jia J, Du T, Zhang N, Tang Y, et al. Overview of histone

modification.   *Adv   Exp   Med   Biol.*   2021;   1283:1-16.

https://doi.org/10.1007/978-981-15-8104-5_1.

5.   Strahl BD, Allis CD. The language of covalent histone modifications. *Nature.*

2000; 403:41-45. https://doi.org/10.1038/47412.

6.   Lachner M, O'Carroll D, Rea S, Mechtler K, Jenuwein T. Methylation of

histone H3 lysine 9 creates a binding site for HP1 proteins. *Nature.* 2001;

410:116-120. https://doi.org/10.1038/35065132.

7.   Schuettengruber B, Bourbon HM, Di Croce L, Cavalli G. Genome regulation

by polycomb and trithorax: 70 Years and Counting. *Cell.* 2017; 171:34-57.

https://doi.org/10.1016/j.cell.2017.08.002.

8.   Wang H, Fan Z, Shliaha PV, Miele M, Hendrickson RC, Jiang X, et al.

H3K4me3 regulates RNA polymerase II promoter-proximal pause-release.

Nature. 2023; 615:339-348. https://doi.org/10.1038/s41586-023-05780-8.

9.   Chen M, Liu Y, Liu Z, Su L, Yan L, Huang Y, et al. Histone acetyltransferase

Gcn5-mediated histone H3 acetylation facilitates cryptococcal morphogenesis

and   sexual   reproduction.   *mSphere.*   2023;   8:e0029923.

https://doi.org/10.1128/msphere.00299-23.

10.   Shivarathri R, Tscherner M, Zwolanek F, Singh NK, Chauhan N, Kuchler K.

The fungal histone acetyl transferase Gcn5 controls virulence of the human

pathogen *Candida albicans* through multiple pathways. *Sci Rep.* 2019; 9:9445.

https://doi.org/10.1038/s41598-019-45817-5.

11.   Park PJ. ChIP-seq: advantages and challenges of a maturing technology. *Nat*

Rev Genet. 2009; 10:669-680. https://doi.org/10.1038/nrg2641.

12.   Furey TS. ChIP-seq and beyond: new and improved methodologies to detect

and characterize protein-DNA interactions. *Nat Rev Genet.* 2012; 13:840-852.

https://doi.org/10.1038/nrg3306.

13.   Locke G, Haberman D, Johnson SM, Morozov AV. Global remodeling of

nucleosome   positions   in   *C. elegans. BMC Genomics.*   2013;   14:284.

https://doi.org/10.1186/1471-2164-14-284.

14.   Dahl JA, Collas P. A rapid micro chromatin immunoprecipitation assay

(microChIP).   *Nat   Protoc.*   2008;   3:1032-1045.

https://doi.org/10.1038/nprot.2008.68.

15.   Torres-Garcia S, Huang Y, Dewornu FS, Tong P, Yeboah R, Allshire R, et al.

Genome-wide   profiling   of   histone   modifications   in   fission   yeast   using

CUT&Tag.   *Methods   Mol   Biol.*   2025;   2862:309-320.

https://doi.org/10.1007/978-1-0716-4168-2_22.

16.   Kaya-Okur HS, Wu SJ, Codomo CA, Pledger ES, Bryson TD, Henikoff JG, et

773   al. CUT&Tag for efficient epigenomic profiling of small samples and single

774   cells.   *Nat   Commun.*   2019;   10:1930.

775   https://doi.org/10.1038/s41467-019-09982-5.

776   17.   Fu Z, Jiang S, Sun Y, Zheng S, Zong L, Li P. Cut&tag: a powerful epigenetic

777   tool   for   chromatin   profiling.   *Epigenetics.*   2024;   19:2293411.

778   https://doi.org/10.1080/15592294.2023.2293411.

779   18.   Bartosovic M, Kabbe M, Castelo-Branco G. Single-cell CUT&Tag profiles

780   histone   modifications   and   transcription   factors   in   complex   tissues. *Nat

781   *Biotechnol.* 2021; 39:825-835. https://doi.org/10.1038/s41587-021-00869-9.

782   19.   Tao XY, Guan XY, Hong GJ, He YQ, Li SJ, Feng SL, et al. Biotinylated Tn5

783   transposase-mediated CUT&Tag efficiently profiles transcription factor-DNA

784   interactions   in   plants.   *Plant   Biotechnol   J.*   2023;   21:1191-1205.

785   https://doi.org/10.1111/pbi.14029.

786   20.   Kaya-Okur HS, Janssens DH, Henikoff JG, Ahmad K, Henikoff S. Efficient

787   low-cost chromatin profiling with CUT&Tag. *Nat Protoc.* 2020; 15:3264-3283.

788   https://doi.org/10.1038/s41596-020-0373-x.

789   21.   Klosterman SJ, Atallah ZK, Vallad GE, Subbarao KV. Diversity, pathogenicity,

790   and   management   of   verticillium   species.   *Annu   Rev   Phytopathol.*   2009;

791   47:39-62. https://doi.org/10.1146/annurev-phyto-080508-081748.

792   22.   Studt-Reinhold   L,   Atanasoff-Kardjalieff   AK,   Berger   H,   Petersen   C,

793   Bachleitner S, Sulyok M, et al. H3K27me3 is vital for fungal development and

secondary metabolite gene silencing, and substitutes for the loss of H3K9me3

in the plant pathogen *Fusarium proliferatum. PLoS Genet.* 2024; 20:e1011075.

https://doi.org/10.1371/journal.pgen.1011075.

23.   Chen YY, Zhu C, Zhao JH, Liu T, Gao F, Zhang YC, et al. DNA

methylation-dependent epigenetic regulation of *Verticillium dahliae* virulence

in plants. *aBIOTECH.* 2023; 4:185-201.

https://doi.org/10.1007/s42994-023-00117-5.

24.   Tang G, Yuan J, Wang J, Zhang YZ, Xie SS, Wang H, et al. Fusarium BP1 is a

reader of H3K27 methylation. *Nucleic Acids Res.* 2021; 49:10448-10464.

https://doi.org/10.1093/nar/gkab844.

25.   Qasim MN, Valle Arevalo A, Paropkari AD, Ennis CL, Sindi SS, Nobile CJ, et

al. Genome-wide profiling of transcription factor-DNA binding interactions in

Candida albicans: A comprehensive CUT&RUN method and data analysis

workflow. *J Vis Exp.* 2022. https://doi.org/10.3791/63655.

26.   Aramayo R, Selker EU. *Neurospora crassa*, a model system for epigenetics

research. *Cold Spring Harb Perspect Biol.* 2013; 5:a017921.

https://doi.org/10.1101/cshperspect.a017921.

27.   Honda S, Lewis ZA, Huarte M, Cho LY, David LL, Shi Y, et al. The DMM

complex prevents spreading of DNA methylation from transposons to nearby

genes in *Neurospora crassa. Genes Dev.* 2010; 24:443-454.

https://doi.org/10.1101/gad.1893210.

28.   Chicas A, Cogoni C, Macino G. RNAi-dependent and RNAi-independent

mechanisms contribute to the silencing of RIPed sequences in *Neurospora*

crassa. *Nucleic* *Acids* *Res.* 2004; 32:4237-4243.

https://doi.org/10.1093/nar/gkh764.

29. Kim JE, Nam H, Park J, Choi GJ, Lee YW, Son H. Characterization of the

CCAAT-binding transcription factor complex in the plant pathogenic fungus

Fusarium *graminearum.* *Sci* *Rep.* 2020; 10:4898.

https://doi.org/10.1038/s41598-020-61885-4.

30. Ling H, Fu X, Huang N, Zhong Z, Su W, Lin W, et al. A sugarcane smut

fungus effector simulates the host endogenous elicitor peptide to suppress

plant immunity. *New* *Phytol.* 2022; 233:919-933.

https://doi.org/10.1111/nph.17835.

31. Rajput MA, Rajput NA, Syed RN, Lodhi AM, Que Y. Sugarcane smut: current

knowledge and the way forward for management. *J Fungi (Basel).* 2021; 7.

https://doi.org/10.3390/jof7121095.

32. Bhuiyan SA, Magarey RC, McNeil MD, Aitken KS. Sugarcane smut, caused

by *Sporisorium scitamineum*, a major disease of sugarcane: A contemporary

review. *Phytopathology.* 2021; 111:1905-1917.

https://doi.org/10.1094/phyto-05-21-0221-rvw.

33. Elías-Villalobos A, Fernández-Álvarez A, Moreno-Sánchez I, Helmlinger D,

Ibeas JI. The Hos2 histone deacetylase controls *Ustilago maydis* virulence

through direct regulation of mating-type genes. *PLoS Pathog*. 2015;

11:e1005134. https://doi.org/10.1371/journal.ppat.1005134.

34.   Hou X, Liu L, Li Y, Wang P, Pan X, Xu D, et al. Regulation of histone

acetylation modification on biosynthesis of secondary metabolites in fungi. *Int*

J Mol Sci. 2024; 26. https://doi.org/10.3390/ijms26010025.

35.   Su C, Lu Y, Liu H. N-acetylglucosamine sensing by a GCN5-related

N-acetyltransferase induces transcription via chromatin histone acetylation in

fungi. *Nat Commun.* 2016; 7:12916. https://doi.org/10.1038/ncomms12916.

36.   Zhang S, Guo Y, Li S, Li H. Histone acetyltransferase CfGcn5-mediated

autophagy governs the pathogenicity of *Colletotrichum fructicola. mBio.* 2022;

13:e0195622. https://doi.org/10.1128/mbio.01956-22.

37.   Yang J, Liu M, Jiao Y, Guo HS, Shan CM, Wang H. An efficient homologous

recombination-based in situ protein-labeling method in *Verticillium dahliae*.

Biology (Basel). 2024; 13. https://doi.org/10.3390/biology13020081.

38.   John E, Singh KB, Oliver RP, Tan KC. Transcription factor control of

virulence in phytopathogenic fungi. *Mol Plant Pathol.* 2021; 22:858-881.

https://doi.org/10.1111/mpp.13056.

39.   Millán-Zambrano G, Burton A, Bannister AJ, Schneider R. Histone

post-translational modifications - cause and consequence of genome function.

Nat   Rev   Genet.   2022;   23:563-580.

https://doi.org/10.1038/s41576-022-00468-7.

40.   Collemare J, Seidl MF. Chromatin-dependent regulation of secondary

metabolite biosynthesis in fungi: is the picture complete? *FEMS Microbiol*

Rev. 2019; 43:591-607. https://doi.org/10.1093/femsre/fuz018.

41.   Kang H, Fan T, Wu J, Zhu Y, Shen WH. Histone modification and chromatin

861   remodeling in plant response to pathogens. *Front Plant Sci*. 2022; 13:986940.

862   https://doi.org/10.3389/fpls.2022.986940.

863   42.   Becker JS, Nicetto D, Zaret KS. H3K9me3-dependent heterochromatin:

864   barrier to cell fate changes. *Trends Genet.* 2016; 32:29-41.

865   https://doi.org/10.1016/j.tig.2015.11.001.

866   43.   Muramatsu D, Singh PB, Kimura H, Tachibana M, Shinkai Y. Pericentric

867   heterochromatin generated by HP1 protein interaction-defective histone

868   methyltransferase Suv39h1. *J Biol Chem.* 2013; 288:25285-25296.

869   https://doi.org/10.1074/jbc.M113.470724.

870   44.   Saksouk N, Simboeck E, Déjardin J. Constitutive heterochromatin formation

871   and transcription in mammals. *Epigenetics Chromatin.* 2015; 8:3.

872   https://doi.org/10.1186/1756-8935-8-3.

873   45.   Cai Y, Zhang Y, Loh YP, Tng JQ, Lim MC, Cao Z, et al. H3K27me3-rich

874   genomic regions can function as silencers to repress gene expression via

875   chromatin interactions. *Nat Commun.* 2021; 12:719.

876   https://doi.org/10.1038/s41467-021-20940-y.

877   46.   Trojer P, Reinberg D. Facultative heterochromatin: is there a distinctive

878   molecular signature? *Mol Cell.* 2007; 28:1-13.

879   https://doi.org/10.1016/j.molcel.2007.09.011.

880   47.   Cheng K, Xu Y, Yang C, Ouellette L, Niu L, Zhou X, et al. Histone tales:

881   lysine methylation, a protagonist in Arabidopsis development. *J Exp Bot.* 2020;

71:793-807. https://doi.org/10.1093/jxb/erz435.

48.   Klocko AD, Rountree MR, Grisafi PL, Hays SM, Adhvaryu KK, Selker EU.

Neurospora importin α is required for normal heterochromatic formation and

DNA   methylation.   *PLoS   Genet.*   2015;   11:e1005083.

https://doi.org/10.1371/journal.pgen.1005083.

49.   Ferraro AR, Ameri AJ, Lu Z, Kamei M, Schmitz RJ, Lewis ZA. Chromatin

accessibility profiling in *Neurospora crassa* reveals molecular features

associated with accessible and inaccessible chromatin. *BMC Genomics.* 2021;

22:459. https://doi.org/10.1186/s12864-021-07774-0.

50.   Gao F, Zhou BJ, Li GY, Jia PS, Li H, Zhao YL, et al. A glutamic acid-rich

protein identified in *Verticillium dahliae* from an insertional mutagenesis

affects microsclerotial formation and pathogenicity. *PLoS One.* 2010;

5:e15319. https://doi.org/10.1371/journal.pone.0015319.

51.   Liu Q, Li Y, Wu H, Zhang B, Liu C, Gao Y, et al. Hyphopodium-specific

signaling is required for plant infection by *Verticillium dahliae*. *J Fungi*

(Basel). 2023; 9. https://doi.org/10.3390/jof9040484.

52.   Liu XL, Yang Y, Hu Y, Wu J, Han C, Lu Q, et al. The nutrient-sensing GCN2

signaling pathway is essential for circadian clock function by regulating

histone acetylation under amino acid starvation. *Elife.* 2023; 12.

https://doi.org/10.7554/eLife.85241.

53.   Lu S, Wang Y, Shen X, Guo F, Zhou C, Li R, et al. SsPEP1, an effector with

essential cellular functions in sugarcane smut fungus. *J Fungi (Basel).* 2021; 7.

https://doi.org/10.3390/jof7110954.

54.   Tang G, Xia H, Huang Y, Guo Y, Chen Y, Ma Z, et al. Liquid-liquid phase

separation of H3K27me3 reader BP1 regulates transcriptional repression.

Genome Biol. 2024; 25:67. https://doi.org/10.1186/s13059-024-03209-7.

55.   Wang S, Xing H, Hua C, Guo HS, Zhang J. An improved single-step cloning

strategy simplifies the agrobacterium tumefaciens-mediated transformation

(ATMT)-based gene-disruption method for *Verticillium dahliae*.

Phytopathology. 2016; 106:645-652.

https://doi.org/10.1094/phyto-10-15-0280-r.

56.   Yang Y, Duan Z, Liu XL, Li Z, Shen Z, Gong S, et al. Checkpoint kinases

regulate the circadian clock after DNA damage by influencing chromatin

dynamics. *Nucleic Acids Res.* 2025; 53. https://doi.org/10.1093/nar/gkaf162.

57.   Zhou Z, Liu X, Hu Q, Zhang N, Sun G, Cha J, et al. Suppression of

WC-independent frequency transcription by RCO-1 is essential for

Neurospora circadian clock. *Proc Natl Acad Sci U S A*. 2013; 110:E4867-4874.

https://doi.org/10.1073/pnas.1315133110.

58.   Cui G, Dong Q, Duan J, Zhang C, Liu X, He Q. NC2 complex is a key factor

for the activation of catalase-3 transcription by regulating H2A.Z deposition.

Nucleic Acids Res. 2020; 48:8332-8348. https://doi.org/10.1093/nar/gkaa552.

59.   Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ

preprocessor. *Bioinformatics.* 2018; 34:i884-i890.

https://doi.org/10.1093/bioinformatics/bty560.

60.   Li H. Aligning sequence reads, clone sequences and assembly contigs with

927   BWA-MEM. *ArXiv.* 2013; 1303. https://doi.org/10.48550/arXiv.1303.3997.

928   61.   Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al.

929   Twelve years of SAMtools and BCFtools. *Gigascience.* 2021; 10.

930   https://doi.org/10.1093/gigascience/giab008.

931   62.   Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, et al.

932   Model-based analysis of ChIP-Seq (MACS). *Genome Biol.* 2008; 9:R137.

933   https://doi.org/10.1186/gb-2008-9-9-r137.

934   63.   Kurtenbach S, William Harbour J. SparK: A Publication-quality NGS

935   Visualization Tool. *bioRxiv*. 2019:845529. https://doi.org/10.1101/845529.

936   64.   Yu G, Wang LG, He QY. ChIPseeker: an R/Bioconductor package for ChIP

937   peak annotation, comparison and visualization. *Bioinformatics*. 2015;

938   31:2382-2383. https://doi.org/10.1093/bioinformatics/btv145.

939   65.   Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, et al.

940   deepTools2: a next generation web server for deep-sequencing data analysis.

941   *Nucleic Acids Res.* 2016; 44:W160-165. https://doi.org/10.1093/nar/gkw257.

DAY 1

DAY 2

(1) Protoplast preparation

(2) Nucleus extraction

(3) Incubation of ConA beads and nuclei

ConA Beads

(4) Incubation of primary antibody

(5) Incubation of secondary antibody

(6) Incubation with pA/G-Tn5

Adapter1 Adapter2

(7) Mg²⁺-activated fragmentation

Mg²⁺

(8) DNA extraction

(9) PCR amplification for library construction

(10) Sequencing

A

Marker | rep1 | rep2

C

Type	Sample	Fastp			ChipQC		
		Raw Reads	After Filter	Dup%	Map%	SSD	Frip
ChIP-Seq	H3K9me3_rep1	6.638665 M	5.382923 M	78.88	95	1.2	3.1
	H3K9me3_rep2	4.179488 M	3.933892 M	9.69	96	2.4	8.5
fCUT&Tag -Seq	H3K9me3_rep1	18.403866 M	17.952596 M	10.19	100	9	6.5
	H3K9me3_rep2	19.364860 M	19.155852 M	12.51	100	9.5	7.2

B

rep1 — rep2
Count — Insert Size — cumulative fraction of reads > insert size

D

ChIP-Seq
38036254 — 100 kb — rep1
2628052 — rep2
fCUT&Tag
331853655 — rep1
rep2
NC_026505.1: 3732166-4637095

23184238 — 100 kb — rep1
1613927 — rep2
331853655 — rep1
rep2
NC_026501.1: 3374994-4711946

E

ChIP-Seq_rep1 / ChIP-Seq_rep2 / fCUT&Tag-rep1 / fCUT&Tag-rep2

0.71	0.73	0.97	1.00
0.71	0.71	1.00	0.97
0.99	1.00	0.71	0.73
1.00	0.99	0.71	0.71

0.0 0.2 0.4 0.6 0.8 1.0

F

ChIP-Seq
rep1 — pearson=0.97 — rep2

fCUT&Tag-Seq
rep1 — pearson=0.99 — rep2

G

174292803 — 1 Mb — WT
NcΔdim5
NC_026503.1: 1-5,274,802

167211163 — 1 Mb — WT
NcΔdim5
NC_026501.1: 1-9,798,893

H

rep1 rep2 rep1 rep2
Percentage(%)

- Promoter (≤1kb)
- Promoter (1-2kb)
- 3' UTR
- 1st Exon
- Other Exon
- 1st Intron
- Other Intron
- Downstream (≤300)
- Distal Intergenic
- 5' UTR

ChIP-seq / fCUT&Tag-seq

A

rep1

rep2

Count

cumulative fraction of reads > insert size

Insert Size

B

Type	Sample	Fastp			ChipQC		
		Raw Reads	After Filter	Dup%	Map%	SSD	Frip
ChIP-Seq	rep1	20.388190M	20.388190M	38.40	98	2.3	18
	rep2	20.190939M	20.190939M	38.72	98	2.4	21
fCUT&Tag-Seq	rep1	39.067562M	38.931750M	18.08	100	21	46
	rep2	56.074418M	55.690382M	23.02	100	24	47

C

ChIP-Seq

58231413

10 kb

rep1

rep2

62074755

10 kb

rep1

rep2

fCUT&Tag

1049355440

rep1

rep2

845477281

rep1

rep2

HG970332: 7,827,549-7,964,530

HG970332: 4,011,619-4,080,109

D

Promoter (≤1kb) (75.96%)
Promoter (1-2kb) (20.55%)
3' UTR (0.18%)
1st Exon (0.21%)
Other Exon (1.99%)
Other Intron (0.05%)
Distal Intergenic (1.06%)

Promoter
genic
Exon
Intron
fiveUTR
Intergenic
threeUTR
downstream
distal_intergenic

E

ChIP-Seq

pearson=0.96

rep1

rep2

fCUT&Tag-Seq

pearson=0.99

rep1

rep2

DAY 1

Formaldehyde crosslinking
of mycelium

Protoplast preparation

Nucleus extraction

ConA Beads

Incubation of
Con A beads and nuclei

Incubation of primary antibody

DAY 2

Incubation of secondary antibody

Adapter1 Adapter2

Incubation with pA/G-Tn5

Mg2+

Mg^{2+}-activated fragmentation

Decrosslinking

DAY 3

RNase and protease digestion

DNA extraction

PCR amplification
for library construction

Sequencing

C

sample	fastp			ChipQC		
	raw reads	after filter	dup%	map%	SSD	Frip
35_Gcn5-Flag_rep1	83.098330M	82.579590M	0.32182209	100	18	12
35_Gcn5-Flag_rep2	28.955856M	28.728874M	0.22725545	100	13	13
36_Gcn5-Flag_rep1	58.758662M	58.445130M	0.24264974	100	15	8.9
36_Gcn5-Flag_rep2	55.960086M	55.311566M	0.19444187	100	14	10