Avoidance of fratricide and improved efficacy of CD33-MSLN CAR-iNK cells in treating acute myeloid leukemia

Yao Wang[1†], Xiujuan Zheng[1,2,3,4†], Zhiqian Wang[2,3,4], Ziyun Xiao[2,3,4], Yunqing Lin[2,3,4], Fan Zhang[2,3,4], Yanhong Liu[2,3], Pengcheng Liu[2,3,4], Qitong Weng[2,3], Leqiang Zhang[2,3,4], Chengxiang Xia[3], Dehao Huang[3], Lijuan Liu[2,3], Yanping Zhu[2,3], Qi Zhang[2,3], Hanmeng Qi[3], Yi Chen[3], Yiyuan Shen[3], Chenyuan Zhang[2,3,4], Jiacheng Xu[2,3,4], Yaoqin Zhao[2,3,4], Jiaxin Wu[2,3,4], Tongjie Wang[2,3], Mengyun Zhang[2,3], Minming Li[5], Wenbin Qian[6], Aibin Liang[7], Xin Du[5], Wenyu Yang[8], Xiaofan Zhu[8], Fangxiao Hu[3*], Jinyong Wang[1,2,3,4*]

[1]Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou 510530, China

[2]State Key Laboratory of Organ Regeneration and Reconstruction, Institute of Zoology, Chinese Academy of Sciences; Beijing, 100101, China

[3]Beijing Institute for Stem Cell and Regenerative Medicine, Beijing 100101, China

[4]University of Chinese Academy of Sciences, Beijing 100049, China

[5]Department of Hematology, Guangdong Provincial People's Hospital, Guangdong Academy of Medical Sciences, Guangzhou 510080, China

[6]Department of Hematology, the Second Affiliated Hospital, College of Medicine, Zhejiang University, Hangzhou, Zhejiang, 310009, China

[7]Department of Hematology, Tongji Hospital of Tongji University, Shanghai 200065, China

[8]State Key Laboratory of Experimental Hematology, National Clinical Research Center for Blood Diseases, Haihe Laboratory of Cell Ecosystem, Institute of Hematology & Blood Diseases Hospital, Chinese Academy of Medical Science & Peking Union Medical College, Tianjin 300020, China

† These authors contributed equally: Yao Wang, Xiujuan Zheng.

*Correspondence: Fangxiao Hu (hufangxiao@biscrm.ac.cn), Jinyong Wang

(wangjinyong@ioz.ac.cn)

One Sentence Summary: Pluripotent stem cell-derived CD33^{KO}-LOOP CAR-iNK cells avoid

fratricide and demonstrate superior cytotoxicity against CD33$^+$MSLN$^+$ leukemia cells.

Abstract

CAR-NK cell therapy has been investigated for treating acute myeloid leukemia (AML). CD33

is a well-established therapeutic target for AML. Conventional CD33 CAR-NK cells are prone

to fratricide due to endogenous expression of CD33 on expanded NK cells. Mesothelin

(MSLN), a tumor differentiation antigen, is expressed in approximately 36% of pediatric AML

cases and 14% of adult AML cases. In this study, we designed a novel CD33-MSLN loop CAR

(Loop CAR) and evaluated its anti-tumor efficacy in human umbilical cord blood-derived NK

(UCB-NK) cells and human pluripotent stem cell-derived NK (hPSC-iNK) cells. The Loop

CAR exhibited superior cytotoxicity against dual-antigen-positive tumor cell lines and primary

AML cells. To further optimize this approach, we established a hPSC-derived cell line via

knockout of endogenous CD33 gene (CD33^{KO}) and integration of the Loop CAR construct.

Using organoid induction technology, we generated mature CD33^{KO}-Loop CAR-iNK cells,

characterized by expression of endogenous CD16. This innovative strategy effectively

mitigated NK cell fratricide and significantly enhanced CD33-mediated specific cytotoxicity.

Moreover, the CD33^{KO}-Loop CAR-iNK cells demonstrated superior tumor-killing activity in

AML xenograft mice and significantly prolonged survival of the treated animals. hPSC-derived

CD33^{KO}-Loop CAR-iNK cells have translational potential for treating AML.

INTRODUCTION

Acute myeloid leukemia (AML) is an aggressive hematologic malignancy that demands urgent therapy (*1*). As new therapeutic targets are being identified, significant efforts have highlighted mesothelin (MSLN) as a promising target in AML. The MSLN is expressed in 36% of pediatric AML cases and 14% of adult AML cases, with 76% of MSLN-positive (MSLN$^+$) patients retaining MSLN expression at relapse. Furthermore, 4% of MSLN-negative (MSLN$^-$) AML patients acquire MSLN expression upon relapse (*2-4*). Preclinical data suggest that MSLN chimeric antigen receptor (CAR)-T cells effectively eliminate MSLN$^+$ AML cells in xenograft models (*5*). Therefore, MSLN is a promising target for AML immunotherapy.

Natural killer (NK) cells have demonstrated considerable potential in enhancing outcomes for AML treatment, particularly in the context of hematopoietic stem cell transplantation (*6*). The adoptive transfer of NK cells for AML therapy has been widely investigated in clinical settings (*7*). CAR-NK cells, which combine the target killing ability of CAR technology with the innate nonspecific cytotoxicity of NK cells, have shown promising results for tumor treatment (*8, 9*). CD33 CAR-NK cells, in particular, have been tested in relapsed/refractory AML therapies, with clinical data demonstrating their efficacy and safety (*10*). Additionally, pluripotent stem cell (PSC)-derived NK or CAR-NK cells have shown therapeutic potential in both preclinical and clinical settings for AML (*11, 12*). PSC-derived NK (iNK) cells provide a homogeneous alternative to NK cells isolated from human tissues (*11, 13, 14*).

Despite these advancements, challenges remain in CAR immune cell therapy, such as tumor antigen escape, on-target off-tumor toxicity, and unintended fratricide due to the shared expression of target antigens between CAR immune cells and tumor cells (*15-17*). To overcome these issues, dual-targeting CARs have been designed to simultaneously target two antigens expressed on tumor cells (*18, 19*). Inactivation of the target antigen in the CAR-engineered immune cells, or in healthy tissue cells, can mitigate fratricide and spare normal tissues. For

example, the disruption of CD7 in CD7 CAR-T cells prevents fratricide and allows for the

expansion of these cells (20). Similarly, genetic inactivation of CD33 in hematopoietic stem

cells enables CD33-deficient hematopoiesis to avoid the killing of CD33 CAR-T cells (21).

Since CD33 is also expressed on NK cells, CD33 CAR-NK cells face similar fratricidal

challenges, which can hinder their expansion (17, 22). Human umbilical cord blood NK (UCB-

NK) cells, which exhibit a 1.4%-25.5% CD33 expression range in their resting state, show

reduced cytotoxicity against K562 cells (23). Moreover, NK cells upregulate CD33 expression

levels to nearly 50% during *in vitro* expansion (17).

In this study, we designed a bivalent loop CAR targeting both CD33 and MSLN antigens

(CD33-MSLN loop CAR, denoted as Loop CAR). We engineered this Loop CAR into UCB-

NK cells and hPSCs to generate Loop CAR-NK and Loop CAR-iNK cells. Both Loop CAR-

NK and Loop CAR-iNK cells exhibited superior cytotoxicity against tumor cells expressing

CD33 and MSLN antigens, including AML cell lines and patient-derived tumor cells.

Transcriptomic analysis confirmed that the Loop CAR enhanced NK cell cytotoxicity.

Additionally, we generated a CD33 gene knockout (CD33^{KO}) hPSC line expressing the Loop

CAR (CD33^{KO}-Loop CAR-hPSCs) and successfully produced CD33^{KO}-Loop CAR-iNK cells.

These CD33^{KO}-Loop CAR-iNK cells exhibited enhanced expansion and anti-tumor efficacy *in*

vitro and effectively suppressed tumor progression in xenograft AML models. Our findings

highlight the clinical potential of Loop CAR-NK and Loop CAR-iNK cells for the treatment

of CD33^{+}MSLN^{+} AML.

RESULTS

CD33-MSLN loop CAR-NK cells exhibit superior cytotoxicity against CD33- and MSLN-

positive tumor cells

To enhance the efficacy of NK cells targeting mesothelin (MSLN)-expressing acute myeloid

leukemia (AML), we employed a bivalent loop CAR strategy, designing a dual-target CAR

targeting both CD33 and MSLN. Specifically, the variable light (V_L) and heavy (V_H) chains of

anti-CD33 single-chain variable fragment (CD33 scFv) and anti-MSLN scFv (MSLN scFv),

the $(G_4S)_3$ linker (Linker 3), and the Whitlow linker (*24*) were sequentially fused to create a

bivalent scFv construct (CD33 V_L-linker 3-MSLN V_H-whitlow linker-MSLN V_L-linker 3-

CD33 V_H, CD33-MSLN loop). The CD33-MSLN loop scFv was then linked to the CD8α hinge,

the CD8 transmembrane domain (TMD), and the CD3 ζ signaling domain (SD) to construct

the final CD33-MSLN loop CAR (denoted as Loop CAR). We applied the MSLN CAR and

CD33 CAR as controls (**Fig. 1A**). To assess the potential of Loop CAR in enhancing tumor-

killing activity in NK cells, we first incorporated the Loop CAR element into the SFG vector,

followed by retroviral packaging as previously reported (*25*). After T-cell depletion, umbilical

cord blood NK (UCB-NK) cells were activated using K562-mbIL-21 feeder cells for 6 days,

followed by transduction with Loop CAR retroviral particles (MOI = 5). The transduced cells

were then expanded for 4 days in the presence of K562-mbIL-21 feeder cells (**Fig. 1B** and **fig.**

S1A). The proportion of Loop CAR$^+$ NK cells (CD56$^+$CD33 CAR$^+$MSLN CAR$^+$) reached 43.4%

two days after transduction, while control groups, including EGFP-NK, CD33 CAR-NK, and

MSLN CAR-NK, achieved transduction rates of 48.2%, 55.6%, and 52.1%, respectively (**Fig.**

1C). Loop CAR-NK cells showed comparable expression of typical NK cell activating and

inhibitory receptors, as well as cytotoxic granules, to the control groups (**fig. S1B**).

We next evaluated the cytotoxicity of Loop CAR-NK cells by sorting CAR$^+$ NK cells and

coculturing them with Nomo-1 (CD33$^+$MSLN$^+$), AGS (CD33$^-$MSLN$^+$), or HL-60

(CD33$^+$MSLN$^-$) tumor cells (**Fig. 1**, **D** and **E**, **fig. S1C**). Loop CAR-NK cells exhibited

significantly superior cytotoxicity against Nomo-1 cells compared to EGFP-NK, CD33 CAR-

NK, and MSLN CAR-NK cells across all effector-to-target (E:T) ratios (E:T = 0.2:1, 0.4:1,

0.8:1, 1.6:1) (**Fig. 1F**). In addition, Loop CAR-NK cells also demonstrated potent killing

activity against MSLN- or CD33-expressing tumor cells, such as AGS (CD33$^-$MSLN$^+$) and

HL-60 ($CD33^{+}MSLN^{-}$), with cytotoxicity comparable to MSLN CAR-NK or CD33 CAR-NK cells (**Fig. 1**, **G** and **H**). Mechanistically, Loop CAR-NK cells exhibited the highest expression of CD107a, a marker associated with NK cell functional activity, when cocultured with Nomo-1 cells (**fig. S1**, **D-F**). Furthermore, Loop CAR molecules accumulated at the immunological synapse (IS) formed between Loop CAR-NK cells and Nomo-1 cells, confirming that the Loop CAR participates in IS formation in a CD33-MSLN-dependent manner (**Fig. 1I**). Notably, Loop CAR-NK cells also demonstrated superior cytotoxicity against primary AML cells expressing both CD33 and MSLN. Mononuclear cells (MNCs) were isolated from the bone marrow (BM) or peripheral blood (PB) of three AML patients (**Fig. 1J**). All three patients' MNCs expressed CD33 (Patient #1: 87.5%, Patient #2: 95.6%, Patient #3: 98.0%) and MSLN (Patient #1: 41.3%, Patient #2: 23.5%, Patient #3: 57.8%) (**Fig. 1K**). The expression of MSLN was further confirmed by qPCR, which showed consistency with the protein expression levels (**Fig. 1L** and **table S1**). Loop CAR-NK cells (Effectors, E) were cocultured with these primary AML cells (Targets, T) for 20 hours at various E:T ratios (0:1, 0.5:1, 1:1, 2:1, and 5:1) (**Fig. 1M**). Loop CAR-NK cells exhibited significantly higher cytotoxicity against all patient-derived targets compared to CD33 CAR-NK and MSLN CAR-NK cells. In addition, both CD33 CAR-NK and MSLN CAR-NK cells killed more targets than EGFP-NK cells (**Fig. 1**, **N-P**). Collectively, these data demonstrate that Loop CAR-NK cells exhibit superior specific cytotoxicity against tumor cells expressing both CD33 and MSLN compared to CD33 CAR-NK or MSLN CAR-NK cells.

Loop CAR-NK cells exhibit a more pronounced activation status after incubation with Nomo-1 cells

To investigate the superior cytotoxicity of Loop CAR-NK cells, we performed single-cell RNA sequencing (scRNA-seq) on sorted Loop CAR-NK cells after incubation with Nomo-1 cells for 72 hours (**Fig. 2A**). UMAP analysis revealed that the transcriptional profiles of Loop CAR-

NK cells were similar to those of CD33 CAR-NK or MSLN CAR-NK cells. However, all CAR-

NK cells exhibited distinct transcriptional profiles compared to EGFP-NK cells (**Fig. 2B**). We

further examined the differentially expressed genes (DEGs) between Loop CAR-NK cells and

CD33 CAR-NK or MSLN CAR-NK cells. Gene Ontology (GO) enrichment analysis indicated

that genes upregulated in Loop CAR-NK cells relative to MSLN CAR-NK cells were primarily

involved in cell division, positive regulation of the Fc receptor-mediated stimulatory signaling

pathway, and positive regulation of the RIG-1 signaling pathway (**Fig. 2C** and **fig. S2**, **A** and

B). Similarly, genes upregulated in Loop CAR-NK cells compared to CD33 CAR-NK cells

were enriched in lymphoid cell differentiation, positive regulation of the RIG-1 signaling

pathway, and formation of immunological synapses. (**Fig 2D** and **fig. S2**, **C** and **D**). Among the

activation-related genes, several hallmarks of NK cell activation and cytotoxic function,

including *CD244*, *SLAMF7*, and *ITGAL* (LFA-1), were upregulated (*26-28*). Additionally,

genes involved in the RIG-I signaling pathway, such as *ANKRD17*, *PUM1*, and *PUM2* (*29, 30*),

and those associated with immunological synapse (IS) formation, including *NCAM1*, *PTK2B*,

and *DOCK8* (*31-34*), were also significantly elevated (**Fig. 2**, **E-G**). These findings suggest

that Loop CAR-NK cells exhibit enhanced activation following incubation with Nomo-1 cells,

which likely contributes to their superior cytotoxicity compared to CD33 CAR-NK or MSLN

CAR-NK cells.

The fratricide of Loop CAR-NK and Loop CAR-iNK cells precludes their expansion

The expansion ability of CD33 CAR-NK cells has been impaired due to fratricide mediated by

the CD33 CAR, as CD33$^+$ NK cells are eliminated by CAR-expressing NK cells (*17*). Here,

the expression of CD33 in UCB-derived Loop CAR-NK cells and hPSC-derived Loop CAR-

iNK cells also led to fratricide. In expanded UCB-NK cells, the expression of CD33 was

observed at 26.9% (**Fig. 3A**). To assess fratricide, we sorted CD33$^-$ and CD33$^+$ NK cells as

targets and co-incubated them with Loop CAR-NK cells (**Fig. 3**, **B** and **C**). As expected, Loop

CAR-NK cells exhibited specific cytotoxicity against CD33$^+$ NK cells at the indicated effector-

to-target (E:T) ratios, while no cytotoxicity was observed against CD33$^-$ NK cells (**Fig. 3D).**

We then sorted an equal number of EGFP-NK and Loop CAR-NK cells and cultured them

consecutively without K562-mbIL-21 stimulation. Over a 12-day period, the number of Loop

CAR-NK cells was lower than EGFP-NK cells, and the gap gradually widened, confirming

that fratricide impaired the expansion of Loop CAR-NK cells (**Fig. 3E**). In our previous work,

we established a method for inducing iNK and CAR-iNK cells from pluripotent stem cells (*11,*

35). Here, we generated a Loop CAR-expressing hPSC line (Loop CAR-hPSCs) and then

conducted the induction of Loop CAR-iNK cells (**Fig. 3F**). Briefly, Loop CAR-hPSCs were

induced for two days to differentiate into Loop CAR-expressing lateral plate mesoderm cells

(Loop CAR-iLPM), which were then combined with OP9 cells to form organoid aggregates.

After 25 days of induction, Loop CAR-iNK cells were generated (**Fig. 3G**). The resulting Loop

CAR-iNK cells expressed similar levels of CD56 and CD16 as iNK cells, with 78.6% of the

cells expressing the Loop CAR (**Fig. 3H**). Moreover, Loop CAR-iNK cells showed comparable

expression of typical NK cell cytotoxic effectors to the iNK cells (**fig. S3A**). Notably, the CD33

expression level in iNK cells was 69.7%, which was significantly higher than that observed in

tissue-derived NK cells (**Fig. 3I**). Next, we sorted CD33$^-$ and CD33$^+$ iNK cells and co-

incubated them with Loop CAR-iNK cells (**Fig. 3, J** and **K**). The Loop CAR-iNK cells showed

significant cytotoxicity against CD33$^+$ iNK cells compared to CD33$^-$ iNK cells (**Fig. 3L**). As

observed with the Loop CAR-NK cells, the number of Loop CAR-iNK cells gradually declined

during a 12-day expansion period, reinforcing that fratricide significantly impaired the

expansion of Loop CAR-iNK cells (**Fig. 3M**). Taken together, the expression of CD33 in Loop

CAR-NK and Loop CAR-iNK cells led to fratricide, which precluded their expansion.

CD33 inactivation in iNK cells preserves their broad-spectrum cytotoxicity and facilitates

the expansion of Loop CAR-iNK cells

To address the fratricide caused by CD33 expression in Loop CAR-iNK cells, we generated a CD33 gene knockout hPSC (CD33^{KO}-hPSC) line using the CRISPR/Cas12i system. Vectors carrying Cas12i, sgRNA1 (5'-TTCAGAAGTGGCCGCAAGGGAAG-3'), sgRNA2 (5'-TTGCTAGTCATCCCCAAATATCC-3'), and a neomycin resistance gene were transduced into hPSCs (**Fig. 4A**). Following neomycin selection, single clones were isolated and genotyped using four pairs of PCR primers (**fig. S4**, **A** and **B**). Ultimately, we identified a CD33^{KO}-hPSC clone with the deletion of exon 3 and exon 4 in one allele of CD33 and the insertion of a loxp-NeoR-loxp sequence in the other (**Fig. 4B** and **fig. S4**, **C** and **D**). Subsequently, CD33^{KO}-hPSCs successfully generated CD33-deficient iNK cells (CD33^{KO}-iNK) with immunophenotype similar to iNK cells (**Fig. 4C**). Moreover, the yields of CD33^{KO}-iNK cells were comparable to those of iNK cells (**Fig. 4D**). Importantly, CD33^{KO}-iNK cells exhibited cytotoxicity against K562 cells equivalent to that of iNK cells, demonstrating that the loss of CD33 did not compromise their broad-spectrum cytotoxicity (**Fig. 4E**). We then introduced the Loop CAR into CD33^{KO}-hPSCs to establish the CD33^{KO}-Loop CAR-hPSC line. As anticipated, these cells successfully differentiated into CD33^{KO}-Loop CAR-iNK cells, which exhibited expression levels of CD56 and CD16 comparable to those of iNK cells, with a CAR expression rate of 89.3% (**Fig. 4**, **F** and **G**). Furthermore, the expression of typical activating and inhibitory receptors, as well as cytotoxic granules, was consistent with that of iNK cells (**fig. S4E**). Most notably, CD33^{KO}-Loop CAR-iNK cells demonstrated a similar expansion capacity to that of iNK cells, indicating that CD33 disruption effectively resolved the fratricide issue and restored the proliferation of Loop CAR-iNK cells (**Fig. 4H**). In conclusion, we successfully mitigated the fratricide issue in Loop CAR-iNK cells by knocking out CD33 in hPSCs, thereby preserving their broad-spectrum cytotoxicity and restoring their expansion.

Genetic disruption of CD33 in Loop CAR-iNK cells enhances their specific cytotoxicity

To evaluate the specific cytotoxicity of CD33^{KO}-Loop CAR-iNK cells, we incubated them with

Nomo-1 cells or MSLN-expressing HL-60 cells (MSLN$^+$ HL-60) (**Fig. 5**, **A** and **B**). Notably,

CD33^{KO}-Loop CAR-iNK cells exhibited significantly enhanced cytotoxicity compared to Loop

CAR-iNK cells, suggesting that fratricide in the latter compromised their specific cytotoxicity

(**Fig. 5**, **C** and **D**). To further evaluate the durability of the cytotoxic response, we performed

serial killing assays by analyzing the cytotoxicity of CD33^{KO}-Loop CAR-iNK cells (**Fig. 5E**).

Compared to Loop CAR-iNK cells, CD33^{KO}-Loop CAR-iNK cells consistently demonstrated

superior specific cytotoxicity against Nomo-1 cells in each round killing assay (**Fig. 5F**).

Additionally, we assessed the expression levels of key effector molecules, including TNF-α,

IFN-γ, and CD107a, in CD33^{KO}-Loop CAR-iNK cells following incubation with Nomo-1 and

MSLN$^+$ HL-60 cells. The results revealed that CD33^{KO}-Loop CAR-iNK cells expressed

significantly higher levels of these molecules compared to Loop CAR-iNK cells, further

corroborating their enhanced effector function (**Fig. 5**, **G-O**). Collectively, these findings

underscore the critical role of CD33 disruption in restoring and strengthening the specific

cytotoxicity of Loop CAR-iNK cells, demonstrating its importance in overcoming fratricide-

induced functional impairments.

CD33^{KO}-Loop CAR-iNK cells effectively suppress AML tumor development in xenograft

animals

To further evaluate the anti-tumor efficacy of CD33^{KO}-Loop CAR-iNK cells, we employed an

AML xenograft mouse model, using iNK cells as control effectors (**Fig. 6A**). MSLN$^+$ HL-60

cells were intravenously injected into NCG mice (5×10^5 cells/mouse) on day -1 to establish

the AML xenograft models. Mice with comparable tumor burdens were randomly assigned to

three groups (Tumor only, iNK, and CD33^{KO}-Loop CAR-iNK). CD33^{KO}-Loop CAR-iNK cells

were administered intravenously at a dose of 1×10^7 cells/mouse on days 0 and 3 (**Fig. 6B**).

Tumor progression was monitored weekly using bioluminescence imaging (BLI). The results

showed that the tumor burden in mice treated with CD33^{KO}-Loop CAR-iNK cells was

significantly lower than in those treated with iNK cells, demonstrating the potent anti-tumor

activity of CD33^{KO}-Loop CAR-iNK cells (**Fig. 6**, **C** and **D**). Moreover, the enhanced tumor

suppression translated into a significant survival benefit, with CD33^{KO}-Loop CAR-iNK-treated

mice achieving a median survival of 42 days, compared to shorter survival in the control groups

(**Fig. 6E**). Collectively, these findings demonstrate that CD33^{KO}-Loop CAR-iNK cells

effectively eliminate AML tumor cells *in vivo* and prolong the survival of xenograft models,

underscoring their therapeutic potential for clinical application.

DISCUSSION

The dual CAR strategy is an effective approach to prevent tumor escape caused by antigen

heterogeneity or loss. Bivalent CAR, one CAR containing two scFvs, which can result in higher

transduction efficiency than bicistronic CAR, two single-targeted CARs. Bivalent CAR can be

categorized into tandem CAR and loop CAR (*36*). Previous studies have reported that loop

CAR exhibited superior anti-tumor potency than tandem CAR (*37, 38*). In this study, we

developed a bivalent loop CAR targeting CD33 and mesothelin (MSLN) (CD33-MSLN loop

CAR, denoted as Loop CAR). When the Loop CAR expression cassette was introduced into

umbilical cord blood NK (UCB-NK) cells, the resulting Loop CAR-NK cells exhibited superior

cytotoxicity against CD33 and MSLN-expressing tumor cells, including Nomo-1 and primary

CD33^{+}MSLN^{+} AML cells. Furthermore, Loop CAR-NK cells demonstrated comparable

cytotoxicity against CD33^{+}MSLN^{-} and CD33^{-}MSLN^{+} tumor cells compared to CD33 CAR-

NK and MSLN CAR-NK cells. The accumulation of Loop CAR proteins at the immunological

synapse between Loop CAR-NK cells and Nomo-1 cells likely contributes to their enhanced

specific cytotoxicity (*39, 40*). Additionally, Nomo-1 cells stimulated Loop CAR-NK cells to

express higher levels of CD107a, a membrane protein associated with NK cell cytotoxic

activity (*41*). Transcriptome analysis revealed that Loop CAR-NK cells exhibited upregulated

signaling pathways related to cell division and activation after incubation with Nomo-1 cells.

Genes such as *NCAM1*, *PTK2B*, and *DOCK8*, which are involved in immunological synapse

formation and cytotoxic function, were significantly upregulated (*31-34, 42*).

Consistent with the previous report (*23*), a fraction of UCB-NK cells naturally express CD33.

Our findings confirmed that hPSC-derived iNK cells exhibit even higher levels of CD33

expression compared to UCB-NK cells. This led to fratricide in both Loop CAR-NK and Loop

CAR-iNK cells, ultimately impairing their expansion. Similar fratricide-induced expansion

impairments have been previously reported in human tissue-derived CD33 CAR-NK cells (*17*),

highlighting the necessity of addressing this challenge. To overcome the fratricide issue, we

genetically disrupted CD33 in hPSCs (CD33^{KO}-hPSCs) using CRISPR/Cas technology.

Previous studies have shown that CD33 knockout in hematopoietic stem cells allows for the

development of a functional hematopoietic system without impairing differentiation potential

(*21*). Our results demonstrated that CD33^{KO}-hPSCs successfully differentiated into iNK cells

without affecting their induction efficiency or anti-tumor activity. Although CD33 has been

considered an inhibitory receptor due to its cytoplasmic tyrosine-based inhibitory motifs

(ITIMs) (*43*). We observed no significant differences in cytotoxicity against K562 cells

between CD33^{KO}-iNK and iNK cells.

As expected, CD33^{KO}-Loop CAR-hPSCs efficiently generated CD33^{KO}-Loop CAR-iNK cells.

The elimination of CD33 expression successfully prevented fratricide and restored the

expansion capacity of Loop CAR-iNK cells. Interestingly, CD33^{KO}-Loop CAR-iNK cells

demonstrated superior specific cytotoxicity against Nomo-1 and MSLN$^+$ HL-60 cells

compared to Loop CAR-iNK cells, indicating that fratricide in the latter not only impaired their

expansion but also reduced their specific cytotoxicity. Moreover, CD33^{KO}-Loop CAR-iNK

cells significantly suppressed tumor progression and prolonged the survival of AML xenograft

mice. We used MSLN$^+$ HL-60 cells, an M2 subtype of acute myeloid leukemia (AML), to

construct the tumor models, as these cells exhibit high tumorigenic potential *in vivo*. Additionally, we attempted to use Nomo-1 cells, an M5a subtype of acute monocytic leukemia, to establish a tumor model. However, the tumorigenic efficiency was suboptimal with Nomo-1 cells. Given NK cells' relatively lower genome editing efficiency compared to T cells (*44*), targeting endogenous CD33 at the pluripotent stem cell (PSC) stage represents a more practical approach to avoid potential fratricide. With their amenability to genetic engineering, clonal selection capabilities, and lack of T cell contamination, PSC-derived NK cells hold promise as standardized, off-the-shelf immunotherapy products (*45, 46*).

Despite significant progress in identifying AML-specific antigens as therapeutic targets, challenges remain due to their genomic and phenotypic heterogeneity and overlapping expression with normal hematopoietic cells (*5*). CD33, expressed in over 80% of AML cells, is absent from primitive stem cells and multipotent progenitor cells, making it an attractive target for AML immunotherapy (*47, 48*). Recent clinical data have demonstrated the safety and efficacy of functionally enhanced CD33 CAR-NK cells (*10*). MSLN, a well-established target in solid tumors, is expressed in approximately 36% of pediatric AML cases and 14% of adult AML cases, but is virtually absent in normal hematopoietic cells (*4*). MSLN CAR-T cells can efficiently eliminate MSLN-positive (MSLN$^+$) AML cells in xenograft models derived from cell lines and patients (*5*). However, MSLN is actively cleaved from the cell membrane and is not highly expressed in AML cells. Strategies such as small molecules (e.g., INCB7839, GM6001) and immunotoxins (e.g., RG7787, SS1P) have been developed to upregulate MSLN expression and reduce shedding, thereby enhancing its viability as a therapeutic target (*5, 49, 50*). Our findings demonstrated that CD33-MSLN loop CAR-NK cells exhibited superior cytotoxicity against primary AML cells expressing both CD33 and MSLN (23.5%-57.8%). Moreover, CD33^{KO}-CD33-MSLN loop CAR-iNK cells displayed robust anti-tumor efficacy both *in vitro* and in xenograft models. These results indicate that the CD33-MSLN loop CAR

design effectively enhances the tumor-killing potential of NK cells and supports the successful

development of a dual-targeting CAR strategy. Additionally, by genetically inactivating CD33,

we effectively addressed the fratricide issue, further improving the function of CD33-MSLN

loop CAR-iNK cells. In conclusion, our study provides compelling preclinical evidence

supporting the further evaluation of CD33-MSLN loop CAR-NK and CD33^{KO}-CD33-MSLN

loop CAR-iNK cells as promising candidates for clinical trials in the treatment of MSLN^{+} AML.

MATERIALS AND METHODS

Study design

The objective of this study was to develop a more efficient therapeutic CAR-NK cell for the treatment of acute myeloid leukemia (AML). This was achieved by designing a bivalent CD33-MSLN loop CAR (Loop CAR) and inactivating CD33, thereby generating Loop CAR-iNK cells that avoid fratricide and effectively target and kill $MSLN^+CD33^+$ AML cells. The construction of Loop CAR was described in the results. The functionality of the Loop CAR was validated by cloning its expression elements into the SFG vector and employing the RD114 retroviral system in conjunction with umbilical cord blood NK cells to generate Loop CAR-NK cells. The superior *in vitro* cytotoxicity of Loop CAR-NK cells compared to CD33 CAR-NK and MSLN CAR-NK cells was assessed through tumor killing assays against $CD33^+MSLN^+$ tumor cell lines (Nomo-1 and $MSLN^+$ HL-60). Furthermore, the expression of CD107a in CAR-NK cells stimulated by Nomo-1 cells was analyzed to evaluate degranulation capacity. Single-cell RNA sequencing (scRNA-seq) using the 10× Genomics platform further demonstrated the enhanced effector functions of loop CAR-NK cells at the transcriptomic level following co-incubation with Nomo-1 cells.

To extend this strategy, the Loop CAR was introduced into hPSCs to generate Loop CAR-iNK cells. However, functional assays revealed that both Loop CAR-NK and Loop CAR-NK iNK cells exhibited fratricide by targeting $CD33^+$ NK cells or iNK cells, leading to limited expansion. To address this challenge, the CRISPR/Cas12i system was employed to generate a CD33 knockout ($CD33^{KO}$) hPSC line. We first confirmed that CD33 disruption did not affect the immunophenotype or cytotoxic function of iNK cells through flow cytometric immunophenotypic analysis and tumor-killing assays against K562 cells. Subsequently, the Loop CAR was introduced into the $CD33^{KO}$-hPSCs to produce $CD33^{KO}$-Loop CAR-iNK cells. Flow cytometric analysis and *in vitro* expansion experiments confirmed that the loss of CD33

did not alter the immunophenotypic characteristics of Loop CAR-iNK cells but significantly enhanced their expansion potential. To validate the improved *in vitro* effector functions of $CD33^{KO}$-Loop CAR-iNK cells, tumor-killing assays and flow cytometry analysis of CD107a expression and effector molecule secretion (IFN-γ and TNF-α) were conducted following stimulation with Nomo-1 and $MSLN^+$ HL-60 cells. To assess the *in vivo* anti-tumor efficacy of $CD33^{KO}$-Loop CAR-iNK cells, AML xenograft models were established using $MSLN^+$ HL-60 cells in NCG mice. Tumor-bearing mice were treated with either iNK or $CD33^{KO}$-Loop CAR-iNK cells, and therapeutic efficacy was monitored through weekly bioluminescence imaging. Blinding was not employed in the experiments. Each experiment was conducted independently on different days using different batches of reagents. Experimental data were repeated to ensure an adequate number of biological replicates. Details regarding sample sizes, replicates, and statistical analyses are provided in the figures, figure legends, and supplementary materials.

Cell culture

Human embryonic kidney (HEK) 293T cells (ATCC) were cultured in Dulbecco's Modified Eagle Medium (DMEM, Gibco) supplemented with 10% fetal bovine serum (FBS, HUANGKE). Human cervical cancer HeLa cells (Procell Life Science & Technology Co., Ltd) were maintained in Minimum Essential Medium α (MEMα, Gibco) with 10% FBS. Mesothelin-expressing ($MSLN^+$) AML cell lines, Nomo-1 (COBIOER), and luciferase-expressing Nomo-1 were maintained in RPMI 1640 (Gibco) supplemented with 10% FBS. Human gastric adenocarcinoma AGS cells (Procell Life Science & Technology Co., Ltd) were cultured in Ham's F12 (Gibco) supplemented with 10% FBS. HL-60 (Procell Life Science & Technology Co., Ltd) and luciferase-expressing $MSLN^+$ HL-60 （$MSLN^+$ HL-60 $luci^+$） cells were cultured in Iscove's Modified Dulbecco's Medium (IMDM, Gibco) supplemented with 20% FBS. To generate MSLN-expressing HL-60 cell line, the cDNA encoding MSLN was inserted into the PiggyBac expression vector (SBI) to generate a recombinant vector. The

MSLN-expressing PiggyBac vector and the transposase expression vector were electroporated

into HL-60 cells using Electroporator EX+ (Celetrix). Seven days after electroporation, HL-60

cells were sorted for two rounds to establish the MSLN-expressing HL-60 cell line. The PSC

line (Q380-ESC) was provided by the National Stem Cell Resource Center, Institute of Zoology,

Chinese Academy of Sciences. All hPSCs were kept in Essential 8 Medium (Gibco) on

Vitronectin-Recombinant Human Protein (rhVTN-N, Gibco) coated plates. The hPSCs

differentiation toward iNK and the related anti-tumor activity assessments of iNK cells in

animals were approved by the Biomedical Research Ethics Committee of the Institute of

Zoology, Chinese Academy of Sciences.

Generation of CAR constructs and CAR-modified NK cells

The monospecific CAR constructs included a signal peptide (SP), the anti-CD33 or anti-

mesothelin (MSLN) single-chain variable fragment (scFv) (*51, 52*), the CD8α hinge domain, a

transmembrane domain (TMD) of CD8, and a signaling domain (SD) of CD3ζ. The Loop CAR

construct was designed using the same backbone, with the scFv arrangement as follows: CD33

(V_L)-MSLN (V_H)-MSLN (V_L)-CD33 (V_H) were fused with the G4S linker and the whitlow

linker (*53*). The EGFP coding sequence or CAR constructs were cloned into SFG recombinant

retroviral vectors (*54*). Retroviruses carrying the CAR vectors were generated as previously

described (*25*). Umbilical cord blood (UCB) units were provided by Guangdong Cord Blood

Bank (Guangzhou, China). CD3$^-$ umbilical cord blood mononuclear cells (UCBMCs) were

isolated, and activated NK cells were produced as previously described (*25*). On day 0,

activated NK cells were transduced with retroviral supernatants at a multiplicity of infection

(MOI) of 5, following the Vectofusin-1 (Miltenyi Biotec) based transduction protocol. Infection

rates were assessed by flow cytometry on day 2. CAR-NK cells were harvested for use after 4

days of expansion. The CAR-NK cells were sorted and expanded for 12 days without

stimulation by K562-mbIL-21 cells (Hangzhou Zhongying Biomedical Technology Co., Ltd).

Flow cytometry

Transduced UCB-NK cells were stained with fluorochrome-conjugated antibodies against CD3

(UCHT1), CD56 (HCD56) (Biolegend), CD33 scFv, and MSLN scFv (ACROBiosystems) for

analysis of CAR expression using a BD LSRFortessa flow cytometer (BD Biosciences). Anti-

CD56, anti-CD33 scFv, and anti-MSLN scFv antibodies were used to sort CAR-NK cells using

Sony MA900 (SONY). Human anti-CD33 antibody (P67.6, Biolegend) and human anti-MSLN

antibody (Abcam) were used to determine the expression levels of CD33 and MSLN in tumor

cell lines and primary AML cells. The immunophenotypes of iNK cells were analyzed using

antibodies against CD45 (HI30, Biolegend), CD3, CD56, CD16 (3G8, Biolegend), CD33 scFv

and MSLN scFv. To evaluate the expression of NK receptors and effector molecules, we used

antibodies against CD56, NKp44 (P44-8), NKp30 (P30-15), NKG2D (1D11), CD69 (FN50),

CD16, CD319 (162.1), CD226 (11A8), CD96 (NK92.39), CD94 (HP-3D9), NKG2A

(S19004C), Perforin (dG9), GzmB (QA18A28) (all from Biolegend), CD33 scFv and MSLN

scFv. Cells were resuspended in DAPI (Sigma-Aldrich) or Dulbecco's phosphate-buffered

saline (BasalMedia Technologies) for analysis with the BD LSRFortessa cytometer (BD

Biosciences). Data were analyzed with FlowJo software (FlowJo_v10.8.1).

Cytotoxicity assays

For cytotoxicity assessment, target cells, labeled with eBioscience™ Cell Proliferation Dye

eFluor™ 670 (eFluor 670) (Invitrogen), were co-cultured with NK cells in the indicated

effector (E) to target (T) ratios (E:T). Target cell death was quantified using the BD

LSRFortessa flow cytometer or BD LSRFortessa X-20 cytometer (BD Biosciences) by

measuring the percentage of DAPI$^+$ among eFluor 670-labeled cells. In serial killing assays,

iNK cells received three rounds tumor challenge. Round 1: eFluor 670-labeled (eFluor 670$^+$)

Nomo-1 cells were mixed with relevant iNK cells and co-incubated for 12 h (E:T = 1:1).

Analysis of the cell lysis of eFluor 670$^+$ Nomo-1 cells at 12 h represents the Round 1

cytotoxicity of the relevant iNK cells. Round 2: Unlabeled Nomo-1 cells were mixed with

relevant iNK cells and co-incubated for 12 h (E:T = 1:1). Equal numbers of eFluor 670-labeled

Nomo-1 cells were added and co-incubated for the next 12 h. Analysis of the cell lysis of eFluor

670$^+$ Nomo-1 cells at 24 h represents the Round 2 cytotoxicity of the relevant iNK cells. Round

3: Unlabeled Nomo-1 cells were mixed with relevant iNK cells (E:T = 1:1). Equal numbers of

unlabeled Nomo-1 cells and eFluor 670-labeled Nomo-1 cells were added at 12 h and 24 h

respectively. Analysis of the cell lysis of eFluor 670$^+$ Nomo-1 cells at 36 h represents the Round

3 cytotoxicity of the relevant iNK cells. Specific cytotoxicity was calculated using the formula:

(percentage of specific death – percentage of spontaneous death) / (1 – percentage of

spontaneous death) × 100%.

Assessment of CD107a, TNF-α, and IFN-γ by flow cytometry

To evaluate CD107a expression, NK or iNK cells were co-cultured with or without Nomo-1 or

MSLN$^+$ HL-60 cells at an E:T ratio of 0.5:1 for 4 h. After incubation, cells were stained with

antibodies against CD56, CD107a (H4A3, Biolegend), CD33 scFv, and MSLN scFv. To

evaluate TNF-α and IFN-γ expression, iNK cells were co-cultured with or without Nomo-1 or

MSLN$^+$ HL-60 cells at an E:T ratio of 0.5:1 for 2 h, followed by adding BFA/Monensin

(MULTISCIENCES) for additional 2-h incubation. After incubation, cells were stained with

antibodies against CD56, CD33 scFv, and MSLN scFv. FIX &PERM Kit (MULTISCIENCES)

was used for fixation and permeabilization, followed by intracellular staining for TNF-α

(Biolegend, MAb11) or IFN-γ (Biolegend, 4S.B3). The cells were analyzed with BD

LSRFortessa X-20 cytometer (BD Biosciences). Data were analyzed with FlowJo software

(FlowJo_v10.8.1).

Immunofluorescence imaging

Immunofluorescence imaging of immune synapses of Loop CAR-NK cells by confocal

microscopy as previously described (*55*). Briefly, immune synapses of Loop CAR-NK cells

were formed by suspending NK cells at 5×10^6/mL and Nomo-1 target cells at 2.5×10^6/mL.

Equal volumes of Loop CAR-NK cell and Nomo-1 target cell suspensions were mixed and

incubated to form conjugates. Loop CAR-NK and Nomo-1 cell conjugates were permeabilized

using the FIX & PERM kit (MULTISCIENCES) and stained with antibodies against phalloidin

(Thermo Fisher Scientific), CD33 scFv, MSLN scFv, pericentrin (Abcam), and perforin

(Biolegend). Sequential scanning with a Leica TCS SP8 laser scanning confocal microscope

imaged NK cell immune synapses. The images were acquired using LASAF software (Leica).

Quantitative real-time PCR

The *MSLN* transcripts were reverse transcribed into cDNA using HiScript III All-in-One RT

SuperMix Perfect for qPCR (Vazyme) and quantified by quantitative real-time PCR using

Hieff® qPCR SYBR® Green Master Mix (Yeasen Biotechnology). *GAPDH* was used for

normalization. The relative gene expression levels were calculated as $2^{-\Delta Ct}$. All primers used in

this study were listed in **table S1**.

Primary samples

Primary AML cells were isolated from AML diagnostic bone marrow or peripheral blood

samples obtained from the State Key Laboratory of Experimental Hematology, National

Clinical Research Center for Blood Diseases, Institute of Hematology & Blood Diseases

Hospital, Chinese Academy of Medical Science & Peking Union Medical College, and Department of Hematology, Guangdong Provincial People's Hospital. Primary AML cells were thawed for immunophenotype analysis of MSLN expression and cultured in IMDM supplemented with 20% FBS, 10 ng/ml GM-CSF (MedChemExpress), 100 U/ml DNase I (Sigma Aldrich), and 50 uM GM6001 (Selleck) (a matrix metalloprotease inhibitor that stabilizes MSLN expression on the cell surface (5) for cytotoxicity assays. *In vitro* killing assays were performed as mentioned above. The use of patient samples was carried out by the provisions of the Declaration of Helsinki. All patient samples were collected with patient consent signatures and reviewed and approved by the Ethics Committee of the Hospital for Hematology & Blood Diseases and the Guangdong Provincial People's Hospital, respectively.

Single-cell RNA-sequencing and data analysis

FACS-sorted EGFP-NK ($CD56^+EGFP^+$) and CAR-NK cells ($CD56^+CAR^+$) were activated with eFluor 670 labeled (eFluor 670^+) Nomo-1 cells for 72 h. Fifty thousand cells sorted (eFluor 670^- and $CD56^+$) from activated NK populations underwent RNA-sequencing (10× Genomics). Droplet-based scRNA-seq datasets were generated using a Chromium system (10× Genomics, PN120263) following the manufacturer's instructions. The raw data were uploaded to the public database of Genome Sequence Archive (HRA008163). Subsequently, these data sets were aligned and quantified using the CellRanger software package (version 6.0.0) and subjected to Seurat (version 3.2.3) for further analysis. To pass quality control, cells were required to have at least 200 genes detected, less than 7,500 genes detected, and less than 20% of aligned reads mapped to mitochondrial genes. 15,408, 17269, 15273, and 16734 cells from EGFP-NK, CD33 CAR-NK, MSLN CAR-NK, and Loop CAR-NK scRNA-seq datasets that passed quality control were merged and used for downstream analysis. The Unique Molecular Identifier (UMI) value was normalized using Seurat's 'LogNormalize' and 'ScaleData'

methods. The Principal Component Analysis (PCA) used 2,000 highly variable genes (as defined by Seurat), and clustering was performed on the most variable principal components (PCs). The most variable PCs were determined by the Jackstraw method. Then, the 36 most variable PCs were used for clustering and Uniform Manifold Approximation and Projection (UMAP) analysis with resolution 0.2 and other default parameters. The median and dot plots for gene expression were plotted using the Seurat VlnPlot function and the ggplot2 package. Differential expression genes (DEGs) were identified using the neurinoma test with the parameter 'min.pct = 0.25, logfc threshold = 0.25'. Gene ontology (GO) enrichment analysis for DEGs was performed by the clusterProfiler package (version 4.2.2) with BH adjusted cutoff value = 0.05. Box plots of the normalized expression of DEGs were plotted using the ggplot2 package. Volcano plots were plotted using ggplot2 and the ggrepel package.

Regeneration of CAR-iNK cells

CAR-expressing hPSCs were engineered using the piggyBac transposon system, as mentioned above. CAR-iNK cells were regenerated from CAR-transfected hPSCs and harvested for use as previously described (*11*). Briefly, CAR-hPSCs were performed a two-day induction to produce CAR lateral plate mesoderm cells (CAR-iLPM). Then 2×10^4 CAR-iLPM were combined with 5×10^5 OP9 cells to form organoid aggregates and seeded in the transwell (Corning, 3450). Subsequently, the medium was changed every 2-3 days to induce NK cell regeneration. After 25 days of induction, mature CAR-iNK cells were harvested for use.

CRISPR/Cas12i-mediated CD33 knockout in hPSCs

Two sgRNA sequences, sgRNA1 (TTCAGAAGTGGCCGCAAGGGAAG) and sgRNA2 (TTGCTAGTCATCCCCAAATATCC), targeting exon 3 and exon 4 of CD33 respectively, were designed using a web-based guide-RNA designer (http://www.rgenome.net/) and

subsequently cloned into the Cas12i expression vector (a gift from Dr. Wei Li of the Institute

of Zoology, Chinese Academy of Sciences) (*56*). To create CD33 knockout hPSCs (CD33^{KO}-

hPSCs), the two Cas12i-sgRNA expression vectors and a donor vector carrying the neomycin

resistance gene (NeoR) were co-transfected into hPSCs using electroporation. Five days post-

transfection, G418 (InvivoGen) was introduced to select the targeted hPSCs for one week.

Following G418 selection, the cells were digested and single cells were seeded into 96-well

plates to form single-cell clones. PCR and sequencing were used to verify the mutation sites,

thereby establishing CD33 knockout clones.

Genome PCR

Single-cell clones of hPSCs resistant to G418 were harvested for DNA extraction using the

TIANamp Genomic DNA kit (TIANGEN). The CD33 locus target sequences were amplified

with the 2 × Rapid Taq Master Mix (Vazyme) and the primer pairs depicted in **fig. S4**.

AML xenograft models with CAR-iNK cell treatment

The MSLN^{+} HL-60 luci^{+} cells (5×10^5 cells/mouse) were injected intravenously into NCG

mice (NOD/ShiLtJGpt-*Prkdc*^{em26Cd52}*Il2rg*^{em26Cd22}/Gpt strain, GemPharmatech Co., Ltd.) to

establish human AML xenograft models on day -1. On day 0, bioluminescence imaging (BLI,

IVIS Spectrum PerkinElmer) was used to quantify the tumor graft burden in each group. After

irradiation (1.0 Gy, Rad Source RS2000), the animals were randomly assigned to treatment

groups and received an intravenous injection of the equivalent of 1×10^7 fresh CAR^{+} iNK or

unmodified iNK cells. Three days later (day 3), the animals received the second intravenous

injection with the same cell count. Tumor aggressiveness was determined weekly by BLI. Mice

with a significant tumor burden were humanely euthanized. All animal experiments were

conducted under the Institutional Animal Care and Use Committee (IACUC) of the Institute of

Zoology of the Chinese Academy of Sciences (IOZ-2021-182).

Statistical analysis

All quantitative analysis were performed with SPSS (IBM SPSS Statistics 25). The Shapiro-Wilk normality test assessed the normal distribution of the data. Quantitative differences (mean ± SD) between samples were compared using two-tailed Student's t-test, Mann-Whitney U test, one-way ANOVA test, and two-way ANOVA test. The *P values* were two-sided, and $P < 0.05$ was considered statistically significant. Survival curves for tumor-bearing models were plotted using the Kaplan-Meier method and compared to the treatment group using logarithmic rank (Mantel-Cox) tests. Statistical analysis was performed using the GraphPad Prism 9 software.

List of Supplementary Materials

Fig. S1 to S4 for multiple supplementary figures.

Table S1 for multiple supplementary table.

References and Notes

1. H. Kantarjian, G. Borthakur, N. Daver, C. D. DiNardo, G. Issa, E. Jabbour, T. Kadia, K. Sasaki, N. J. Short, M. Yilmaz, F. Ravandi, Current status and research directions in acute myeloid leukemia. *Blood Cancer J* **14**, 163 (2024).
2. D. Steinbach, A. Schramm, A. Eggert, M. Onda, K. Dawczynski, A. Rump, I. Pastan, S. Wittig, N. Pfaffendorf, A. Voigt, F. Zintl, B. Gruhn, Identification of a set of seven genes for the monitoring of minimal residual disease in pediatric acute myeloid leukemia. *Clin Cancer Res* **12**, 2434-2441 (2006).
3. D. Steinbach, M. Onda, A. Voigt, K. Dawczynski, S. Wittig, R. Hassan, B. Gruhn, I. Pastan, Mesothelin, a possible target for immunotherapy, is expressed in primary AML cells. *Eur J Haematol* **79**, 281-286 (2007).
4. A. J. Kaeding, S. P. Barwe, A. Gopalakrishnapillai, R. E. Ries, T. A. Alonzo, R. B. Gerbing, C. Correnti, M. R. Loken, L. E. Broderson, L. Pardo, Q. H. Le, T. Tang, A. R. Leonti, J. L. Smith, C. K. Chou, M. Xu, T. Triche, S. M. Kornblau, E. A. Kolb, K. Tarlock, S. Meshinchi, Mesothelin is a novel cell surface disease marker and potential therapeutic target in acute myeloid leukemia. *Blood Adv* **5**, 2350-2361 (2021).
5. Q. Le, S. Castro, T. Tang, A. M. Loeb, T. Hylkema, C. N. McKay, L. Perkins, S. Srivastava, L. Call, J. Smith, A. Leonti, R. Ries, L. Pardo, M. R. Loken, C. Correnti, S. Fiorenza, C. J. Turtle, S. Riddell, K. Tarlock, S. Meshinchi, Therapeutic Targeting of Mesothelin with Chimeric Antigen Receptor T Cells

in Acute Myeloid Leukemia. *Clin Cancer Res* **27**, 5718-5730 (2021).

6. K. C. Hsu, C. A. Keever-Taylor, A. Wilton, C. Pinto, G. Heller, K. Arkun, R. J. O'Reilly, M. M. Horowitz, B. Dupont, Improved outcome in HLA-identical sibling hematopoietic stem-cell transplantation for acute myelogenous leukemia predicted by KIR and HLA genotypes. *Blood* **105**, 4878-4884 (2005).

7. S. Z. D'Silva, M. Singh, A. S. Pinto, NK cell defects: implication in acute myeloid leukemia. *Front Immunol* **14**, 1112059 (2023).

8. Y. Kaito, Y. Imai, Evolution of natural killer cell-targeted therapy for acute myeloid leukemia. *Int J Hematol* **120**, 34-43 (2024).

9. Y. Masamoto, M. Kurokawa, A key to engineering natural killer cells to attack acute myeloid leukemia. *Haematologica* **109**, 1032-1034 (2024).

10. R. Huang, X. Wang, H. Yan, X. Tan, Y. Ma, M. Wang, X. Han, J. Liu, L. Gao, L. Gao, G. Jing, C. Zhang, Q. Wen, X. Zhang, Safety and efficacy of CD33-targeted CAR-NK cell therapy for relapsed/refractory AML: preclinical evaluation and phase I trial. *Exp Hematol Oncol* **14**, 1 (2025).

11. D. Huang, J. Li, F. Hu, C. Xia, Q. Weng, T. Wang, H. Peng, B. Wu, H. Wu, J. Xiong, Y. Lin, Y. Wang, Q. Zhang, X. Liu, L. Liu, X. Zheng, Y. Geng, X. Du, X. Zhu, L. Wang, J. Hao, J. Wang, Lateral plate mesoderm cell-based organoid system for NK cell regeneration from human pluripotent stem cells. *Cell Discov* **8**, 121 (2022).

12. A. Eckstrom, A. Tyagi, S. Mahmood, L. Wong, B. Valamehr, A. Rao, A. Agrawal, M. Siddiqui, V. L. Battula, FT538, iPSC-derived NK cells, enhance AML cell killing when combined with chemotherapy. *J Cell Mol Med* **29**, e70169 (2025).

13. D. A. Knorr, Z. Ni, D. Hermanson, M. K. Hexum, L. Bendzick, L. J. Cooper, D. A. Lee, D. S. Kaufman, Clinical-scale derivation of natural killer cells from human pluripotent stem cells for cancer therapy. *Stem Cells Transl Med* **2**, 274-283 (2013).

14. Y. Li, D. L. Hermanson, B. S. Moriarity, D. S. Kaufman, Human iPSC-Derived Natural Killer Cells Engineered with Chimeric Antigen Receptors Enhance Anti-tumor Activity. *Cell Stem Cell* **23**, 181-192 e185 (2018).

15. E. J. Orlando, X. Han, C. Tribouley, P. A. Wood, R. J. Leary, M. Riester, J. E. Levine, M. Qayed, S. A. Grupp, M. Boyer, B. De Moerloose, E. R. Nemecek, H. Bittencourt, H. Hiramatsu, J. Buechner, S. M. Davies, M. R. Verneris, K. Nguyen, J. L. Brogdon, H. Bitter, M. Morrissey, P. Pierog, S. Pantano, J. A. Engelman, W. Winckler, Genetic mechanisms of target antigen loss in CAR19 therapy of acute lymphoblastic leukemia. *Nat Med* **24**, 1504-1506 (2018).

16. M. L. Cooper, J. Choi, K. Staser, J. K. Ritchey, J. M. Devenport, K. Eckardt, M. P. Rettig, B. Wang, L. G. Eissenberg, A. Ghobadi, L. N. Gehrs, J. L. Prior, S. Achilefu, C. A. Miller, C. C. Fronick, J. O'Neal, F. Gao, D. M. Weinstock, A. Gutierrez, R. S. Fulton, J. F. DiPersio, An "off-the-shelf" fratricide-resistant CAR-T for the treatment of T cell hematologic malignancies. *Leukemia* **32**, 1970-1983 (2018).

17. M. Hejazi, C. Zhang, S. B. Bennstein, V. Balz, S. B. Reusing, M. Quadflieg, K. Hoerster, S. Heinrichs, H. Hanenberg, S. Oberbeck, M. Nitsche, S. Cramer, R. Pfeifer, P. Oberoi, H. Ruhl, J. Oldenburg, P. Brossart, P. A. Horn, F. Babor, W. S. Wels, J. C. Fischer, N. Moker, M. Uhrberg, CD33 Delineates Two Functionally Distinct NK Cell Populations Divergent in Cytokine Production and Antibody-Mediated Cellular Cytotoxicity. *Front Immunol* **12**, 798087 (2021).

18. C. C. Kloss, M. Condomines, M. Cartellieri, M. Bachmann, M. Sadelain, Combinatorial antigen recognition with balanced signaling promotes selective tumor eradication by engineered T cells. *Nat Biotechnol* **31**, 71-75 (2013).

19. D. Abate-Daga, M. L. Davila, CAR models: next-generation CAR modifications for enhanced T-cell function. *Mol Ther Oncolytics* **3**, 16014 (2016).

20. D. Gomes-Silva, M. Srinivasan, S. Sharma, C. M. Lee, D. L. Wagner, T. H. Davis, R. H. Rouce, G. Bao, M. K. Brenner, M. Mamonkin, CD7-edited T cells expressing a CD7-specific CAR for the therapy of T-cell malignancies. *Blood* **130**, 285-296 (2017).

21. M. Y. Kim, K. R. Yu, S. S. Kenderian, M. Ruella, S. Chen, T. H. Shin, A. A. Aljanahi, D. Schreeder, M. Klichinsky, O. Shestova, M. S. Kozlowski, K. D. Cummins, X. Shan, M. Shestov, A. Bagg, J. J. D.

Morrissette, P. Sekhri, C. R. Lazzarotto, K. R. Calvo, D. B. Kuhns, R. E. Donahue, G. K. Behbehani, S. Q. Tsai, C. E. Dunbar, S. Gill, Genetic Inactivation of CD33 in Hematopoietic Stem Cells to Enable CAR T Cell Immunotherapy for Acute Myeloid Leukemia. *Cell* **173**, 1439-1453 e1419 (2018).

22. N. Albinger, R. Pfeifer, M. Nitsche, S. Mertlitz, J. Campe, K. Stein, H. Kreyenberg, R. Schubert, M. Quadflieg, D. Schneider, M. W. M. Kuhn, O. Penack, C. Zhang, N. Moker, E. Ullrich, Primary CD33-targeting CAR-NK cells for the treatment of acute myeloid leukemia. *Blood Cancer J* **12**, 61 (2022).

23. R. Handgretinger, H. J. Schafer, F. Baur, D. Frank, C. Ottenlinger, H. J. Buhring, D. Niethammer, Expression of an early myelopoietic antigen (CD33) on a subset of human umbilical cord blood-derived natural killer cells. *Immunol Lett* **37**, 223-228 (1993).

24. M. Whitlow, B. A. Bell, S. L. Feng, D. Filpula, K. D. Hardman, S. L. Hubert, M. L. Rollence, J. F. Wood, M. E. Schott, D. E. Milenic, et al., An improved linker for single-chain Fv with reduced aggregation and enhanced proteolytic stability. *Protein Eng* **6**, 989-995 (1993).

25. Y. Wang, J. Li, Z. Wang, Y. Liu, T. Wang, M. Zhang, C. Xia, F. Zhang, D. Huang, L. Zhang, Y. Zhao, L. Liu, Y. Zhu, H. Qi, X. Zhu, W. Qian, F. Hu, J. Wang, Comparison of seven CD19 CAR designs in engineering NK cells for enhancing anti-tumour activity. *Cell Prolif* **57**, e13683 (2024).

26. S. O. Mathew, K. K. Rao, J. R. Kim, N. D. Bambard, P. A. Mathew, Functional role of human NK cell receptor 2B4 (CD244) isoforms. *Eur J Immunol* **39**, 1632-1641 (2009).

27. A. Gutierrez-Guerrero, I. Mancilla-Herrera, J. L. Maravillas-Montero, I. Martinez-Duncker, A. Veillette, M. E. Cruz-Munoz, SLAMF7 selectively favors degranulation to promote cytotoxicity in human NK cells. *Eur J Immunol* **52**, 62-74 (2022).

28. D. F. Barber, M. Faure, E. O. Long, LFA-1 contributes an early signal for NK cell cytotoxicity. *J Immunol* **173**, 3653-3659 (2004).

29. Y. Wang, X. Tong, G. Li, J. Li, M. Deng, X. Ye, Ankrd17 positively regulates RIG-I-like receptor (RLR)-mediated immune signaling. *Eur J Immunol* **42**, 1304-1315 (2012).

30. R. Narita, K. Takahasi, E. Murakami, E. Hirano, S. P. Yamamoto, M. Yoneyama, H. Kato, T. Fujita, A novel function of human Pumilio proteins in cytoplasmic sensing of viral infection. *PLoS Pathog* **10**, e1004417 (2014).

31. J. T. Gunesch, A. L. Dixon, T. A. Ebrahim, M. M. Berrien-Elliott, S. Tatineni, T. Kumar, E. Hegewisch-Solloa, T. A. Fehniger, E. M. Mace, CD56 regulates human NK cell cytotoxicity through Pyk2. *Elife* **9**, (2020).

32. D. Sancho, M. Nieto, M. Llano, J. L. Rodriguez-Fernandez, R. Tejedor, S. Avraham, C. Cabanas, M. Lopez-Botet, F. Sanchez-Madrid, The tyrosine kinase PYK-2/RAFTK regulates natural killer (NK) cell cytotoxic response, and is translocated and activated upon specific target cell recognition and killing. *J Cell Biol* **149**, 1249-1262 (2000).

33. H. Ham, W. Huynh, R. A. Schoon, R. D. Vale, D. D. Billadeau, HkRP3 is a microtubule-binding protein regulating lytic granule clustering and NK cell killing. *J Immunol* **194**, 3984-3996 (2015).

34. H. Ham, S. Guerrier, J. Kim, R. A. Schoon, E. L. Anderson, M. J. Hamann, Z. Lou, D. D. Billadeau, Dedicator of cytokinesis 8 interacts with talin and Wiskott-Aldrich syndrome protein to regulate NK cell cytotoxicity. *J Immunol* **190**, 3661-3669 (2013).

35. Y. Liu, M. Zhang, X. Shen, C. Xia, F. Hu, D. Huang, Q. Weng, Q. Zhang, L. Liu, Y. Zhu, L. Wang, J. Hao, M. Zhang, T. Wang, J. Wang, Mesothelin CAR-engineered NK cells derived from human embryonic stem cells suppress the progression of human ovarian cancer in animals. *Cell Prolif* **57**, e13727 (2024).

36. B. Xie, Z. Li, J. Zhou, W. Wang, Current Status and Perspectives of Dual-Targeting Chimeric Antigen Receptor T-Cell Therapy for the Treatment of Hematological Malignancies. *Cancers (Basel)* **14**, (2022).

37. X. Wang, Z. Dong, D. Awuah, W. C. Chang, W. A. Cheng, V. Vyas, S. C. Cha, A. J. Anderson, T. Zhang, Z. Wang, S. J. Szymura, B. Z. Kuang, M. C. Clark, I. Aldoss, S. J. Forman, L. W. Kwak, H. Qin, CD19/BAFF-R dual-targeted CAR T cells for the treatment of mixed antigen-negative variants of acute lymphoblastic leukemia. *Leukemia* **36**, 1015-1024 (2022).

38. Z. Chen, Y. Liu, N. Chen, H. Xing, Z. Tian, K. Tang, Q. Rao, Y. Xu, Y. Wang, M. Wang, J. Wang, Loop

CD20/CD19 CAR-T cells eradicate B-cell malignancies efficiently. *Sci China Life Sci* **66**, 754-770
(2023).

39. J. S. Orange, K. E. Harris, M. M. Andzelm, M. M. Valter, R. S. Geha, J. L. Strominger, The mature
activating natural killer cell immunologic synapse is formed in distinct stages. *Proc Natl Acad Sci*
U S A **100**, 14151-14156 (2003).

40. E. Liu, Y. Tong, G. Dotti, H. Shaim, B. Savoldo, M. Mukherjee, J. Orange, X. Wan, X. Lu, A. Reynolds,
M. Gagea, P. Banerjee, R. Cai, M. H. Bdaiwi, R. Basar, M. Muftuoglu, L. Li, D. Marin, W. Wierda, M.
Keating, R. Champlin, E. Shpall, K. Rezvani, Cord blood NK cells engineered to express IL-15 and
a CD19-targeted CAR show long-term persistence and potent antitumor activity. *Leukemia* **32**,
520-531 (2018).

41. G. Alter, J. M. Malenfant, M. Altfeld, CD107a as a functional marker for the identification of natural
killer cell activity. *J Immunol Methods* **294**, 15-22 (2004).

42. M. Zhang, M. E. March, W. S. Lane, E. O. Long, A signaling network stimulated by beta2 integrin
promotes the polarization of lytic granules in cytotoxic cells. *Sci Signal* **7**, ra96 (2014).

43. T. Ulyanova, J. Blasioli, T. A. Woodford-Thomas, M. L. Thomas, The sialoadhesin CD33 is a myeloid-
specific inhibitory receptor. *Eur J Immunol* **29**, 3440-3449 (1999).

44. G. M. Robbins, M. Wang, E. J. Pomeroy, B. S. Moriarity, Nonviral genome engineering of natural
killer cells. *Stem Cell Res Ther* **12**, 350 (2021).

45. F. Cichocki, S. J. C. van der Stegen, J. S. Miller, Engineered and banked iPSCs for advanced NK-
and T-cell immunotherapies. *Blood* **141**, 846-855 (2023).

46. B. H. Goldenson, P. Hor, D. S. Kaufman, iPSC-Derived Natural Killer Cell Therapies - Expansion and
Targeting. *Front Immunol* **13**, 841107 (2022).

47. J. Liu, J. Tong, H. Yang, Targeting CD33 for acute myeloid leukemia therapy. *BMC Cancer* **22**, 24
(2022).

48. N. N. Shah, J. Azzi, B. W. Cooper, A. Deol, J. DiPersio, D. Koura, B. McClune, L. S. Muffly, M. U.
Mushtaq, R. Narayan, H. C. Suh, G. Yanik, S. Nath, J. Whangbo, G. Koehne, Phase 1/2 Study of
Donor-Derived Anti-CD33 Chimeric Antigen Receptor Expressing T Cells (VCAR33) in Patients with
Relapsed or Refractory Acute Myeloid Leukemia after Allogeneic Hematopoietic Cell
Transplantation. *Blood* **142**, 4862-4862 (2023).

49. C. Alewine, L. Xiang, T. Yamori, G. Niederfellner, K. Bosslet, I. Pastan, Efficacy of RG7787, a next-
generation mesothelin-targeted immunotoxin, against triple-negative breast and gastric cancers.
Mol Cancer Ther **13**, 2653-2661 (2014).

50. P. Awuah, T. K. Bera, M. Folivi, O. Chertov, I. Pastan, Reduced Shedding of Surface Mesothelin
Improves Efficacy of Mesothelin-Targeting Recombinant Immunotoxins. *Mol Cancer Ther* **15**,
1648-1655 (2016).

51. P. R. Hamann, L. M. Hinman, I. Hollander, C. F. Beyer, D. Lindh, R. Holcomb, W. Hallett, H. R. Tsou,
J. Upeslacis, D. Shochat, A. Mountain, D. A. Flowers, I. Bernstein, Gemtuzumab ozogamicin, a
potent and selective anti-CD33 antibody-calicheamicin conjugate for treatment of acute myeloid
leukemia. *Bioconjug Chem* **13**, 47-58 (2002).

52. P. S. Chowdhury, J. L. Viner, R. Beers, I. Pastan, Isolation of a high-affinity stable single-chain Fv
specific for mesothelin from DNA-immunized mice by phage display and construction of a
recombinant immunotoxin with anti-tumor activity. *Proc Natl Acad Sci U S A* **95**, 669-674 (1998).

53. T. J. Fry, N. N. Shah, R. J. Orentas, M. Stetler-Stevenson, C. M. Yuan, S. Ramakrishna, P. Wolters, S.
Martin, C. Delbrook, B. Yates, H. Shalabi, T. J. Fountaine, J. F. Shern, R. G. Majzner, D. F. Stroncek,
M. Sabatino, Y. Feng, D. S. Dimitrov, L. Zhang, S. Nguyen, H. Qin, B. Dropulic, D. W. Lee, C. L.
Mackall, CD22-targeted CAR T cells induce remission in B-ALL that is naive or resistant to CD19-
targeted CAR immunotherapy. *Nat Med* **24**, 20-28 (2018).

54. I. Riviere, K. Brose, R. C. Mulligan, Effects of retroviral vector design on expression of human
adenosine deaminase in murine bone marrow transplant recipients engrafted with genetically
modified cells. *Proc Natl Acad Sci U S A* **92**, 6733-6737 (1995).

55. K. B. Sanborn, G. D. Rak, A. N. Mentlik, P. P. Banerjee, J. S. Orange, Analysis of the NK cell

immunological synapse. *Methods Mol Biol* **612**, 127-148 (2010).

56.   Y. Chen, Y. Hu, X. Wang, S. Luo, N. Yang, Y. Chen, Z. Li, Q. Zhou, W. Li, Synergistic engineering of
CRISPR-Cas nucleases enables robust mammalian genome editing. *Innovation (Camb)* **3**, 100264
(2022).

Acknowledgments: We thank the staffs of SPF-grade animal facilities at the Institute of

Zoology, Chinese Academy of Sciences for supplying animal care service. **Funding:** This

work was supported by grants from the National Key Research and Development Program of

China (2021YFA1100701, 2024YFA1108302, 2021YFA1100800), the National Natural

Science Foundation of China (82450001, 82470120, 82300132, 32300676, 82350104), and

the Noncommunicable Chronic Diseases-National Science and Technology Major Project

(2023ZD0501300).

Author contributions

Y.W. and X.J.Z. designed and conducted all experiments, performed data analysis, and wrote

the manuscript. Z.Q.W., Z.Y.X., F.Z., Y.H.L., and P.C.L. conducted some experiments and

performed data analysis. Y.Q.L. and Q.T.W. performed scRNA-seq and data analysis. L.Q.Z.,

C.X.X., D.H.H., L.J.L., Y.P.Z., T.J.W., M.Y.Z., W.Y.Y., and M.M.L. participated in multiple

experiments. J.Y.W., X.F.Z., X.D., W.B.Q., and A.B.L. discussed the data and manuscript.

F.X.H. designed the project, designed and guided all experiments, scrutinized all experimental

data, and wrote the manuscript. J.Y.W. revised the manuscript and provided the final approval

of the manuscript.

Competing interests

The authors declare that they have no conflict of interest.

Data and materials availability

The scRNA-seq data of EGFP-NK, CD33 CAR-NK, MSLN CAR-NK, and Loop CAR-NK

cells stimulated by Nomo-1 cells have been uploaded to the Genome Sequence Archive public

database (HRA008163). Other relevant information or data is available from the corresponding

authors upon reasonable request.

Figures

Fig. 1. Development and functional evaluation of Loop CAR-NK cells. (**A**) Schematic

diagram of the design of Loop CAR. (**B**) Schematic diagram showing the generation of Loop

CAR-NK cells procedures and evaluation their cytotoxicity. (**C**) Flow cytometric analysis of

the transduction efficiencies of each group of CAR-NK cells. EGFP virus-infected NK cells

were used as control. (**D**) Flow cytometry histograms showing the expression levels of CD33

and MSLN in target cell lines, including CD33$^+$MSLN$^+$ Nomo-1 AML cells, CD33$^-$MSLN$^+$ AGS gastric adenocarcinoma cells, and CD33$^+$MSLN$^-$ HL-60 AML cells. (**E**) Tumor killing assay design. (**F-H**) Statistics of CAR-NK cells cytotoxicity against Nomo-1 tumor cells (F), AGS cells (G), and HL-60 cells (**H**) at indicated E:T ratios. Nomo-1 cells and HL-60 cells were incubated with effectors for 4 hours (n = 4 per group). AGS cells were incubated with effectors for 6 hours (n = 4 per group). (**I**) Confocal microscopy representative images of immunological synapse formed between loop CAR-NK cells and Nomo-1 cells. The conjugates were stained with antibodies against perforin (Violet), phalloidin-F-actin (Blue), CD33 CAR (Green) and MSLN CAR (Red). (**J**) Schematic diagram showing the collection of peripheral blood or bone marrow cells from patients with AML. (**K**) Flow cytometric analysis of the expression of CD33 and MSLN in primary AML cells. (**L**) Real-time quantitative PCR analysis of MSLN transcript expression levels in primary AML cells. The *GAPDH* gene was used as an internal control. Relative gene expression levels were calculated as $2^{-\Delta Ct}$. (**M**) Schematic diagram showing primary AML cells killed by CAR-NK cells *in vitro*. Each group of NK cells was co-cultured with primary AML cells for 20 hours at indicated E:T ratios (*n* = 4 per group). (**N-P**) Specific cytotoxicity of CAR-NK cells derived from UCB-NK against primary AML cells from three patients (Patient #1: N, Patient #2: O, Patient #3: P). Statistical analysis: one-way ANOVA. NS: not significant; *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

Fig. 2. Molecular characterization of Loop CAR-NK cells. (**A**) Schematic diagram of sorting CAR-NK cells for 10× Genomics' single-cell RNA-seq assay. CAR-NK cells were incubated with Nomo-1 cells (E:T = 1:1) for 72 h before sorting for RNA-seq. (**B**) UMAP visualization contrasting EGFP-NK, CD33 CAR-NK, MSLN CAR-NK, and Loop CAR-NK cells. EGFP-NK cells were used as control. (**C** and **D**) Gene Ontology (GO) analysis of differentially highly expressed genes in Loop CAR-NK cells compared with MSLN CAR-NK (C) and CD33 CAR-NK (D) cells, respectively. Each data point represented a distinct GO set. The color scale corresponded to $-\log_{10}$ transformed false discovery rate (FDR)– adjusted P values. The size of the dot was the number of genes enriched in one term. The enrichment factor was GeneRatio/BgRatio. (**E-G**) Box plots showed the expression profiles of differentially highly expressed genes related to activation and cytotoxicity of NK cells (E), participating in the RIG-1 signaling pathway (F), and involving in immunological synapse formation (G) in Loop CAR-NK cells compared with CD33 CAR-NK and MSLN CAR-NK cells.

Fig. 3

Fig. 3. Expression of CD33 in NK or iNK cells led to fratricide and precluded expansion of Loop CAR-NK or Loop CAR-iNK cells. (**A**) Flow cytometry histograms showing the expression of CD33 in UCB-NK cells. (**B**) Representative FACS plots showing the purity of sorted CD33$^-$ and CD33$^+$ NK cells. (**C**) Schematic diagram of measuring the cytotoxicity of Loop CAR-NK cells against CD33$^+$/CD33$^-$ NK cells *in vitro*. The sorted NK cells were co-cultured with Loop CAR-NK cells for 4 h. (**D**) Statistics of the fratricidal lysis of CAR-NK cells Loop CAR-NK cells (Effectors, E) against CD33$^+$ and CD33$^-$ NK cells (Targets, T) at the indicated E:T ratios (*n* = 4 per group). (**E**) Statistical analysis of the cell number for EGFP-NK and Loop CAR-NK cells at indicated time points. An equal number of EGFP-NK and Loop CAR-NK cells were sorted for expansion *in vitro* (*n* = 3 per group). (**F**) Flow cytometric analysis of Loop-CAR expression in Loop CAR-hPSCs. (**G**) Schematic diagram of Loop CAR-iNK cells generated from Loop CAR-hPSCs. (**H**) Immunophenotypes (CD45$^+$CD3$^-$CD56$^+$CD16$^{+/-}$) and Loop-CAR expression (CD33 CAR$^+$MSLN CAR$^+$) of Loop-CAR-iNK cells. (**I**) Flow cytometry histograms showing the expression levels of CD33 in iNK cells derived from hPSCs. (**J**) Representative FACS plots showing the expession levels of CD33 in sorted CD33$^-$ and CD33$^+$ iNK cells. (**K**) Schematic diagram of measuring the cytotoxicity of Loop CAR-iNK cells against CD33$^+$ and CD33$^-$ iNK cells *in vitro*. (**L**) Statistics of fratricidal lysis of Loop CAR-iNK cells (Effectors, E) against CD33$^+$ and CD33$^-$ iNK cells (Targets, T) at indicated E:T ratios (*n* = 4 per group). (**M**) Statistical analysis of the cell number for iNK and Loop CAR-iNK cells at indicated time points (*n* = 3 per group). An equal number of iNK and Loop CAR-iNK cells were started for culture. Data represent the mean ± SEM. Statistics analysis: one-way ANOVA test (D, L) and two-way ANOVA test (E, M). NS: not significant; **$P < 0.01$, ***$P < 0.001$.

Fig. 4. Inactivation of CD33 restored the expansion of Loop CAR-iNK cells. (**A**) Schematic

diagram of CRISPR/Cas12i-targeted CD33 gene strategy. sgRNA: single guide RNA; HR:

homologous region; NeoR: neomycin resistance gene. (**B**) Schematic diagram showing the

genotype of the established CD33^{KO}-hPSCs. (**C**) Flow cytometric analysis of iNK and

CD33^{KO}-iNK cells. (**D**) Statistics of the yields of iNK and CD33^{KO}-iNK cell (Day 27) induced

from hPSCs and CD33^{KO}-hPSCs (*n* = 3 repeats). (**E**) Statistics of cytotoxicity of iNK and

CD33^{KO}-iNK cells (Effectors, E) against K562 cells (Targets, T) at the indicated E:T ratios (*n*

= 4 per group). (**F**) Schematic diagram of CD33^{KO}-Loop CAR-iNK cells induced from

CD33^{KO}-Loop CAR-hPSCs. (**G**) Immunophenotypes (CD45$^+$CD3$^-$CD56$^+$CD16$^{+/-}$) and Loop-

CAR expression in iNK and CD33^{KO}-Loop CAR-iNK cells. (**H**) Statistical analysis of the cell

number for iNK and CD33^{KO}-Loop CAR-iNK cells at indicated time points. An equal number

of iNK and Loop CAR-iNK cells were started for culture (*n* = 3 repeats). Data represent the

mean ± SEM. Statistics analysis: two-tailed Student's t test (D), one-way ANOVA test (E) and

two-way ANOVA test (H). NS: not significant.

Fig. 5

Fig. 5. CD33^{KO}-Loop CAR-iNK cells exhibited superior tumor-killing abilities than Loop CAR-iNK cells. (**A**) Flow cytometry histograms showing the expression of CD33 and MSLN in HL-60 cells engineered with MSLN (MSLN$^+$HL-60). (**B**) Schematic diagram showing Nomo-1 or MSLN$^+$ HL-60 cells killed by Loop CAR-iNK or CD33^{KO}-Loop-CAR-iNK cells *in vitro*. (**C** and **D**) Statistics of specific cytotoxicities analysis of iNK, Loop CAR-iNK and CD33^{KO}-Loop CAR-iNK cells (Effectors, E) against Nomo-1 (C) and CD33$^+$MSLN$^+$ HL-60 cells (D) (Targets, T) at the indicated E:T ratios (*n* = 4 per group). (**E**) Schematic diagram of the serial killing assays of iNK, Loop CAR-iNK, and CD33^{KO}-Loop CAR-iNK cells against Nomo-1 cells. Each group of iNK cells was co-cultured with Nomo-1 cells for 12 h per round at an E:T ratio of 1:1. The fresh Nomo-1 cells were added to the NK cell residues and incubated for every other 12 h. The lysis of Cell Proliferation Dye eFluor 670-labeled (eFluor 670$^+$) Nomo-1 cells in each round represented the cytotoxicity. (**F**) Statistics of specific cytotoxicity of iNK, Loop CAR-iNK, and CD33^{KO}-Loop CAR-iNK cells in serial killing assays (*n* = 4 per group). (**G, J, M**) Representative flow cytometry plots showing the expression of TNF-α (G), IFN-γ (J), and CD107a (M) in each group of iNK cells stimulated by MLSN$^+$ HL-60 and Nomo-1 cells. (**H** and **I**) Statistical analysis of the expression levels of TNF-α (*n* = 3 repeats). (**K** and **L**) Statistical analysis of the expression levels of IFN-γ (*n* = 3 repeats). (**N** and **O**) Statistical analysis of the expression levels of CD107a (*n* = 3 repeats). Data represent the mean ± SEM. Statistics analysis: one-way ANOVA test. NS: not significant; *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

Fig. 6

Fig. 6. CD33^{KO}-Loop CAR-iNK cells efficiently suppressed tumor progression *in vivo*. (**A**)

Schematic diagram of the tumor cells and therapeutic cells used for *in vivo* assay. (**B**) Schematic

diagram of *in vivo* studies. Luciferase-expressing (luci$^+$) MSLN$^+$ HL-60 cells were used to

generate mouse xenograft models (5 × 10^5 cells/mouse, i.v.). Two doses of CD33^{KO}-Loop

CAR-iNK cells (1× 10^7 cells/mouse per dose) were injected into tumor-bearing mice for

therapy. (**C**) Bioluminescence imaging (BLI) of xenograft models of each group (*n* = 5 per

group). (**D**) Statistics of each group's total flux (p/s) (CD33^{KO}-Loop CAR-iNK vs. iNK).

Statistics analysis: two-tailed Student's *t*-test. NS: not significant; **P < 0.01, ***P < 0.001.

(**E**) Kaplan-Meier survival curves of the experimental groups. Statistics analysis: two-tailed

log-rank test. Tumor only vs. iNK, **P < 0.01, CD33^{KO}-Loop CAR-iNK vs. iNK, **P < 0.01.

Supplementary Materials

Fig. S1

Fig. S1. Phenotypes and effector function of Loop CAR-NK cells. (A) Flow cytometric

analysis of the T cells (CD3$^+$) residual in UCBMCs after CD3$^+$ cells depletion on day -6. (B)

Flow cytometry histograms showing the expression of typical NK cell receptors (NKp44,

NKp30, NKG2D, CD69, CD16, CD96, CD94, and NKG2A) and cytotoxic granules (Perforin

and GzmB) in respective NK cells. Control histograms represent FMO control. (C) The

analysis of sorting purity for CD3$^-$CD56$^+$EGFP$^+$ (EGFP-NK) and CD3$^-$CD56$^+$CAR$^+$ (CAR-

NK) cells post 24 h culturing. (D) Experimental design of the analysis of CD107a expression.

EGFP-NK and CAR-NK cells were stimulated at an E:T ratio of 0.5:1 for 4 h. (E)

Representative flow cytometry plots showing the expression of CD107a in EGFP-NK and

CAR-NK cells in response to Nomo-1 cells. (F) Statistic analysis of the mean fluorescence

intensity (MFI) of CD107a in (E), n = 3 repeats. Statistical analysis: one-way ANOVA, ***P

< 0.001.

Fig. S2

Fig. S2. Differentially expressed genes of Loop CAR-NK cells compared with CD33 CAR-

NK and MSLN CAR-NK cells. (A) Volcano plot of differentially expressed genes between

Loop CAR-NK and MSLN CAR-NK cells. (B) GO analysis of differentially down-regulated

genes in Loop CAR-NK cells compared with MSLN CAR-NK. Each data point represents a

distinct GO set. Color scale corresponds to −log10 FDR (adjusted P values). Dot size was the

number of genes enriched in one term. Enrichment Factor was the GeneRatio/BgRatio. (C)

Volcano plot of differentially expressed genes between Loop CAR-NK and CD33 CAR-NK

cells. (D) GO analysis of differentially down-regulated genes in Loop CAR-NK cells compared

with CD33 CAR-NK cells. Each data point represents a distinct GO set. Color scale

corresponds to −log10 FDR (adjusted P values). Dot size was the number of genes enriched in

one term. Enrichment Factor was the GeneRatio/BgRatio.

Fig. S3

A

891 **Fig. S3. Analysis of typical NK molecules of Loop CAR-iNK cells.** (A) Flow cytometry

892 histograms showing the expression of typical NK cell receptors (NKp44, NKp30, NKG2D,

893 CD69, CD96, CD94, and NKG2A) and cytotoxic granules (Perforin and GzmB) in respective

894 iNK cells. Control histograms represent FMO control.

Fig. S4

A

Primer No.	Forword primer sequence	Reverse primer sequence
1	5'-TCCCAGTAGGAACAACTGCTCCCTGAGCATCGTAGACGCC-3'	5'-TAAATGTTCAAGATGATGGATATGCTATTACCCTGATCTG-3'
2	5'-TCCCAGTAGGAACAACTGCTCCCTGAGCATCGTAGACGCC-3'	5'-GATGGATTGCACGCAGGTTC-3'
3	5'-CATGAGGCTAGAAAGGAAATTGGAACTTAAAGAGGCAGTG-3'	5'-GCAGCTGACAACCAGGAGAAGATC-3'
4	5'-CTCGGTGCTCATAATCACCCCACGG-3'	5'-CTATTCATCTGACAAGTGATTCATATCCAGAACATACGAG-3'

Fig. S4. Identification of CD33 genotypes of CD33^{KO}-hPSCs and molecular features of

CD33^{KO}-Loop CAR-iNK cells. (A) The sequences of four pairs of primers used for

genotyping CD33 knock-out PSC clones. (B) Schematic diagram showing the location of

primer pairs used for identifying CD33 genotypes. (C, D) PCR amplification results of each

fragment using the indicated primer pairs. (E) Flow cytometry analysis of typical NK cell

receptors (CD226, CD319, NKp30, CD69, NKG2D, NKp44, NKG2A, CD94, CD96, TARIL)

and cytotoxic granules (Perforin and GzmB) in respective iNK cells. Control histograms

represent FMO control.

Primer name	Primer sequence (5'→3')
GAPDH-F	CAAGAAGTGGGAGCTGGAAG
GAPDH-R	GTCTCCAGGGACGTCACATT
MSLN-F	ACCCAGAAGACTGTGGATGG
MSLN-R	TGCTGTAGCCAAATTCGTTG

904 **Table S1. Primer sequences for real-time quantitative PCR of selected genes.**