# SOX2 empowers a rapid tumorigenic programme from the tumour-resistant population in the skin

Patricia P. Centeno[1,2], Christopher Chester[2,4], Catriona A. Ford[1], Patrizia Cammareri[1,5], Gareth J. Inman[1,3], Thomas Jamieson[1], Rachel A Ridgway[1], Richard Marais[2,4], Andrew D. Campbell[1], Owen J. Sansom[1,3].

[1] Cancer Research UK Scotland Institute, Glasgow, UK.

[2] Cancer Research UK Manchester Institute, Manchester, UK.

[3] School of Cancer Sciences, University of Glasgow, UK.

[4] Present address: Oncodrug Ltd, Alderley Park, Macclesfield, UK.

[5] Present address: Cancer Research UK Scotland Centre, Institute of Genetics and Cancer, University of Edinburgh, Edinburgh, UK.

## Abstract

The skin undergoes continuous renewal, relying on stem and progenitor cells to maintain its barrier function. Over time, multiple oncogenic and tumour suppressor events accumulate. How the skin tolerates these mutations in different populations and how cutaneous squamous cell carcinoma (cSCC) emerges from mutant clones is poorly understood. Rapid cSCC development can ensue in melanoma patients treated with BRAF inhibitors (BRAFi) through paradoxical activation of RAF signalling. To model this in mice, we induced MAPK hyperactivation in two epidermal basal populations: stem-cell-like (K14/K5+) and differentiation-committed (IVL+) cells. Remarkably rapid tumorigenesis ensued from the K14/K5+ basal population defining them as a tumour primed. In contrast, despite mutations being retained and populating the skin, IVL+ cells exhibited tumour resistance with long latency. Nevertheless, both populations can act as cSCC cells of origin. Once transformed, they share histological and transcriptomic hallmarks of epidermal transformation independently of the cell-of-origin or the oncogenic driving mutations. Critically, the pioneer factor SOX2 was uniquely upregulated in tumours arising from the resistant population and was both necessary and sufficient for their transformation. Notably, SOX2 bestow an embryonic profile to human keratinocytes and is expressed in ~20% of human cSCC, suggesting that it could mark tumours originating from committed progenitors.

## Highlights

- **Paradoxical MAPK signalling activation is a tumour promoter in cSCC**
- **K14/K5+ represent a tumour-primed population and IVL+ represents a tumour-resistant population, but both can initiate skin cancer**
- **Oncogene-expressing tumour-resistant population takes over the entire skin epidermis but preserves homeostasis**
- **SOX2 is sufficient to initiate cSCC from the tumour-resistant population by preventing delamination and differentiation and promoting stemness**

Adult stem cells and progenitors continuously regenerate and maintain the mammalian epidermis,

ensuring its structural integrity and protection against the external environment. However, the ageing

skin accumulates a high mutational burden, including mutations in known drivers of cutaneous

squamous cell carcinoma (cSCC) as well as passenger mutations [1]. The proliferation of these mutated

cells is confined under normal conditions by the skin hierarchy and spatial compartmentalisation, but

stressors and promoter events such as wounding, inflammation, UV exposure, and certain oncogenic

mutations can overcome these constraints leading to tumorigenesis [2–7]. cSCC originates from the

keratinocytes, and data from murine model systems have illustrated the tumour's ability to form from

different origins, including the hair follicle, the sebaceous glands, and the interfollicular epidermis

compartments [2,8–10].

The most widely used cSCC mouse model is the well-established two-stage topical application of the

chemical carcinogen 7,12-dimethylbenz[a]anthracene (DMBA), followed by the promoting agent 12-

O-tetradecanoylphorbol-13-acetate (TPA) [2,11]. These tumours develop rapidly from the *Lgr6+* stem cells

residing in the hair follicle and are predominantly driven by *Hras*$^{Q61L}$ and *Trp53* mutations [2,12].

Nevertheless, mutations in *HRAS* are relatively uncommon in spontaneous human cSCC (9-16%) but

exist in healthy skin as passenger mutations [1,10,13]. However, HRAS-driven cSCC is commonly found in

melanoma patients arising within weeks of treatment with BRAF$^{V600E}$ inhibitors (BRAFi) [13–15]. This has

been attributed to the paradoxical activation of the MAPK signalling pathway by BRAFi, due to the

*HRAS* mutations already present in the skin [15,16]. Spontaneous human cSCC exhibits high mutational

burden, mutational heterogeneity and copy number alterations (Bailey et al., 2023), with the main

driver mutations concentrated in the Notch (79.5%) and TP53 (71%) signalling pathways [3]. However,

single mutations in these pathways are insufficient to induce cSCC in mice, and additional

combinations of genetic driver alterations involving the TGFβ and MAPK signalling pathways are

required for the transformation and modelling of disease progression [8,10,17,18]. The complexity and slow

tumour progression of these models have hindered efforts in the field to understand the biology of

skin carcinogenesis.

Two models have been proposed in the interfollicular epidermis to explain the basal layer

heterogeneity. The first model described the existence of two distinct basal progenitors that contribute

differently during homeostasis, wound healing [19] and mechanical stretching [6]. Upon wounding, these

progenitors sense reduced cell density and enhance proliferation by undergoing symmetric self-

renewing division, whilst the skin adopts a transient fluid-like state to facilitate progenitor migration

for regeneration and remodelling [20]. An alternative second model proposes that the heterogeneity of

the basal layer is caused by different levels of commitment to differentiation, suggesting that basal

cells continuously transition towards differentiation following cues from the surrounding

environment[21]. Nevertheless, the comparison between these two different basal populations in

response to oncogenic stimuli remains unexplored.

Here, we assessed the mechanisms that confer competence to initiate cSCC from the two basal cell

populations. We generated a suite of mouse models expressing different oncogenes targeted to the

basal populations of the interfollicular epidermis, marked by the expression of KRT14+/KRT5+ and

IVL+, that rapidly model and resemble human cSCC. Particularly, we modelled the aetiology of fast-

growing cSCC in melanoma patients driven by the hyperactivation of the MAPK signalling pathway. We

defined a tumour-resistant and a tumour-primed population and uncovered a shared transcriptional

programme responsible for transformation. Importantly, we identified SOX2 as a critical factor for the

transformation of the tumour-resistant population. SOX2, highly expressed in ~25% of human cSCC, is

known to induce stem cell characteristics as a super pioneer factor [22] and is recognised for its ability

to reprogram terminally differentiated somatic cells into induced pluripotent stem cells. Our findings

demonstrate the different susceptibility of basal populations to transformation revealing that tumours

retain the transcriptional priming of their cell of origin.

# 86    Results

## 87    Paradoxical MAPK signalling activation is a tumour promoter in cSCC

To study the susceptibility of the two different basal populations to transformation and tumour

formation, we set to generate novel genetically engineered moused models of cSCC that resembled

human disease. Patients of cSCC often carry mutations in the *HRAS* gene causing hyperactivation of

MAPK signalling. Thus, we expressed $Hras^{G12V}$ allele under its endogenous promoter in the basal stem-

like population and the basal committed population by using tamoxifen-inducible Cre recombinase

expressed from the *Krt14* promoter (*Krt14*-CreERT2; hereafter K14) and the *Ivl* promoter (Ivl-CreERT;

hereafter Ivl) respectively (**Figure 1A**).

Expression of one copy of the $Hras^{G12V}$ allele (hereafter HRAS$^{G12V}$) in the K14+ or the IVL+ population

did not result in tumorigenesis (**Figures 1B and 1C**). This result is consistent with human studies

demonstrating that normal-looking skin carries a battery of mutations, including oncogenes, and that

a promotion event is required to trigger transformation[1,2]. Therefore, we topically treated the models

with the classical inflammation and tumour-promoting agent known to stimulate epidermal

proliferation, TPA, (three times per week) but was insufficient for transformation.

Interestingly, melanoma patients commonly treated with BRAFi, such as Dabrafenib, have an increased

risk of developing cSCC due to the presence of dormant *HRAS* mutations in the skin (Hauschild et al.,

2012; Oberholzer et al., 2012; Su et al., 2012). Consequently, these patients develop fast-growing

tumours within weeks of treatment, driven by the paradoxical activation of the MAPK signalling

pathway. Thus, we treated Ivl:HRAS$^{G12V/+}$ and K14:HRAS$^{G12V/+}$ mice with the BRAFi Dabrafenib three

times a week to mimic the patient dosing regimen and promote tumorigenesis. Indeed, this resulted

in tumorigenesis with a tumour-free survival rate of 22 days for the K14:HRAS$^{G12V/+}$ model and 58 days

for the Ivl:HRAS$^{G12V/+}$ model (**Figures 1B and 1C**). BRAFi-promoted tumours exhibited the classic cSCC

hallmarks, including keratin pearls, parakeratosis, and nuclear dysplasia (**Figure 1D**). Moreover, we also

treated with BRAFi healthy-looking mice carrying one copy of the HRAS$^{G12V}$ in the K14+ and IVL+

mutations at 180±5 days post-oncogene induction. These aged-treated mice rapidly developed

tumours within days from the BRAFi treatment, which acts as a potent tumour-promoting agent and

mimics the onset of cSCC from melanoma patients carrying *HRAS* mutations (**Figure 1E**).

Two copies of the *Hras*$^{G12V}$ were sufficient to drive tumorigenesis without the need for a promoting

agent due to higher levels of MAPK activation. Specifically, two copies of *Hras*$^{G12V}$ expressed in the

K14+ population resulted in mice becoming unwell, probably due to the critical function of HRAS in

the gastrointestinal epithelium before any skin-related malignancies appeared. Nevertheless,

K14:HRAS$^{G12V/G12V}$ mice showed hyperproliferation of the skin basal layer, evidenced by increased Ki67

expression (**Figure S1**). In the IVL+ basal population, two copies of the allele drove tumorigenesis with

a tumour-free survival of 24 days (**Figures 1C and S1**). The resulting tumours displayed similar

histopathological features (**Figure S1**).

## Tumour-primed and tumour-resistant populations coexist in the basal layer

To gain further insights into the tumorigenic capacity of these distinct pools of progenitors, we used

the potent oncogene BRAF$^{V600E/+}$ (hereafter BRAF$^{V600E}$), which encodes a different downstream

oncogenic effector of the pro-proliferative MAPK signalling pathway. BRAF$^{V600E}$ mimics the MAPK

hyperactivation seen by the combination of BRAFi and *Hras* mutations. We generated a suite of mouse

models expressing BRAF$^{V600E}$ under the control of K14 and *Krt5* (K5-CreERT; hereafter K5) promoters

targeting the stem-cell-like population and the IVL promoter targeting the committed basal

populations (**Figures  2A and S2A**).

Upon BRAF$^{V600E}$ oncogene induction, both K5+ and K14+ cells rapidly formed tumours (9 and 10 days

to tumour onset), without the need for further promotion, suggesting that these populations are

permissive to tumour development and primed for oncogenic transformation (**Figures 2B and 2C**). In

contrast, BRAF$^{V600E}$ driven from the IVL model required a much longer time to develop tumours (125

days to tumour onset) and tumours grew slowly (**Figures 2B and 2D**).  We also observed that the

Ivl:BRAF$^{V600E}$ model tends to develop multiple tumours at different body sites, including the back, paws,

ears and lips, while the K5- and K14-driven BRAF$^{V600E}$ models develop one tumour on their back at the

precise area of tamoxifen application (**Figure 2E**).

Reducing the dose and duration of the tamoxifen-induction regimen in the K5:BRAF$^{V600E}$ and

K14:BRAF$^{V600E}$ models delayed tumour onset, but tumours still grew rapidly (**Figure S2B**).

Transplantation of subcutaneous pieces from K5:BRAF$^{V600E}$-derived tumours into C57BL/6J and NSG-II2

immunodeficient mice revealed cancer stem cell potential and an ability to form secondary tumours

**(Figure S2C)**. Thus, we termed this population tumour-primed.

The longer latency period for tumours to develop in the Ivl:BRAF$^{V600E}$ model suggests that the IVL+

population requires additional cooperating events to facilitate tumour onset, so we termed this

population tumour-resistant. To further promote cSCC in this population, we exposed the back of the

Ivl:BRAF$^{V600E}$ mice to mild UV [four weekly doses 6 standard erythema doses (SED)], but it did not

accelerate tumour onset despite UV being a risk factor for cSCC due to its increase in mutation burden

(**Figure 2B**).  Similarly, topical treatment with the inflammation-promoting agent TPA (three times per

week) did not accelerate tumorigenesis (**Figure S2D**). No statistical differences in tumour onset were

seen between K14:BRAF$^{V600E}$ cohorts housed in two different establishments or by sex (**Figures S2E and**

**S2F**). However, the Ivl:BRAF$^{V600E}$ model showed a slightly shorter tumour onset in the cohort housed at

the CRUK SI compared to CRUK MI, while no differences were seen between sexes (**Figures 2G and**

**2H**).

We characterised the histopathological features of tumours originating in the K5:BRAF$^{V600E}$,

K14:BRAF$^{V600E}$and Ivl:BRAF$^{V600E}$ models and found remarkable similarities among them. All tumours

were formed by vertical columns of keratinocytes with clearly demarcated limits (**Figure S3A**). Detailed

immunohistochemistry showed distinctive pathological features of cSCC. These included enlarged

interfollicular epidermis and cornified layers with an aberrant accumulation of extracellular keratin

(keratin pearls), nuclear atypia, and incomplete maturation of keratinocytes (parakeratosis) as the

keratinocytes in the cornified layer retain their nuclei (**Figures  2F, 2G and S3B**).

The tumours presented the hallmarks of papilloma with signs of squamous differentiation, as

demonstrated by the increase in the number of layers, expressing the basal markers KRT5 and KRT14,

but also the suprabasal marker KRT1 (**Figures S3C and S3D**). In the most advanced cases, these markers

were expressed throughout the entire epidermal layer, indicating a complete loss of cellular hierarchy

and disruption of multi-layer organisation. Moreover, the increase in Ki67 denotes epidermal

hyperproliferation and hyperplasia of the interfollicular epidermis (**Figure S3E**). These lesions were

highly proliferative, recapitulating the rapid onset and histopathology observed in human cSCC.

Overall, despite both populations residing in the basal layer and sharing the microenvironment, they
show profound differences in susceptibility to transformation upon oncogene induction.

## 171 Oncogene-expressing tumour-resistant population takes over the entire skin
## 172 epidermis but remains tumour-restrictive.

We next aimed to understand how oncogene expression in the tumour-resistant population
contributes to the clonal dynamics in the basal layer. In homeostasis, IVL+ marks a minority population
in the basal layer that is committed to differentiation, although its differentiation does not prevent cell
cycle entry [21]. We followed in IVL+ or KRT14+ expressing populations by coexpressing the *Rosa26* LSL-
tdTomato fluorescence protein (tdRFP) reporter in the presence and absence of BRAF$^{V600E}$ (**Figure 3A**).
Lineage-tracing experiments were conducted at different time points, ranging from before
macroscopical changes appear in the skin up to day 160 or clinical endpoint, post tdRFP/oncogene
induction (**Figures 3B**).

First, co-immunofluorescence analysis with confocal microscopy of control skin (no oncogene)
revealed that by day 8 in the K14:tdRFP model, all cells in the basal layer are labelled by RFP and remain
labelled at day 28, marking a stable population as previously shown[19,21]. In the Ivl:tdRFP model at day
8, RFP labels only a small population of basal cells that co-expressed KRT14 and IVL transcripts, and
some of those clones are maintained at day 28 (**Figure 3C**).

Then, we compared the RFP-labelled populations between control and oncogene-expressing skin. In
the K14:tdRFP and K14:BRAF$^{V600E}$-tdRFP models, all cells in the basal layer and their progeny in the
suprabasal layer were labelled to saturation. Thus, there was no difference in clonal dynamics between
groups (**Figure 3D**). An initial ~10% drop in label-retaining cells was observed after day 8, when the
induction regimen finished, but the population remained stable, renewing itself after that (**Figure 3E**).
There were no differences in epidermis thickness between groups (**Figure 3F**).

In the Ivl:tdRFP model, the labelled cells were mainly located in the suprabasal layer with a few labelled
cells in the basal layer at day 8 (**Figure 3G**). A ~50% drop in the label-retaining population in the basal
layer was seen after day 8, consistent with committed cell differentiating and delaminating (**Figure
3H**). Suprabasal cells that were not located under a label retainer IVL+ basal cluster turned over, and
those regions completely lost the tdRFP label. Nevertheless, the labelled population remained stable
for up to 160 days, marking a long-residing stable population of the basal layer able to self-renew and
contribute with their progeny to the suprabasal layers (**Figures 3G and 3H**).

In contrast, upon oncogene activation in the Ivl:BRAF$^{V600E}$-tdRFP model, not only did we see a decrease
in the label-retaining population after day 8, but we saw a ~40% increase in the clonal persistence of

labelled basal cells, which overtook the unlabelled population (**Figures 3G and 3H**). At day 15, the

entire basal layer was labelled, similar to what we see in the K14:BRAF[V600E]tdRFP. Moreover, in the

Ivl:BRAF[V600E], there was a sharp increase in epidermal thickness, compared to control, due to the

increased proliferation in differentiating cells that became smaller with time but remained significant

during the study (**Figure 3I**). Thus, despite most basal cells carrying the oncogenic mutation,

constraints remained in place resisting transformation.

These data show that oncogene expression in the committed IVL+ population led to a clonal

proliferation dynamic similar to the stem-cell K14+ population but remained tumour-restrictive.

## cSCC shared transcriptional profile

Given the similarities in histology, we next conducted transcriptional profiling to identify potential

vulnerabilities in these tumours. After batch correction to correct for any bias arising from batches

sequenced at different times (**Figure S4A**), we merged the K5:BRAF[V600E] and K14:BRAF[V600E] tumour

samples as they clustered independently of the model, and our analysis returned no differentially

expressed genes (**Figures S4B and S4C**). Note that these tumours arose from the same cell populations

simultaneously expressing KRT5 and KRT14. Then, we compared K14/K5:BRAF[V600E]- and Ivl:BRAF[V600E]-

derived tumours against normal skin from control littermates. Unsupervised clustering and principal

component analysis (PCA) showed that the tumours separated from the normal skin samples (**Figure**

**4A**).

Both tumour groups revealed a large number of transcriptionally upregulated genes when compared

to normal skin (**Figures 4B and 4C**). Nevertheless, the vast majority of upregulated genes (82.8%) were

shared between the K14/K5:BRAF[V600E] and Ivl:BRAF[V600E] tumours (**Figure 4D**). Importantly, many

transcriptional pathways we would expect to induce by promoters such as TPA are activated by the

oncogene alone. Gene set enrichment analysis highlighted commonly upregulated signatures,

including those associated with the *Myc*, *Lgr5* and *Lgr6* wound healing response [23], embryonic

epidermis [24], and the human-derived tumour-specific keratinocyte population [25] among others (**Figure**

**4E and S5A**). *Myc* has been identified in wound healing as a crucial transcription factor during

sebaceous duct dedifferentiation, generating a broad range of interfollicular epidermis cells that

contribute to re-epithelialisation [5].

We also examined the expression of selected differentially expressed genes enriched in the tumours

(**Figure 4F**). We observed high expression of *Myc*, the known cSCC markers *Krt6a* and *Krt6b* [26], and

*Anxa1* ligand, which is secreted by wounded epithelia during regeneration and re-epithelisation [23].

ANXA1 is also enriched in the tumour-specific keratinocyte population signature [25], and it has been

shown to play a tumour-promoting role in other cancers of epithelial origin, such as cancers of the

prostate [27] and breast [28]. We also found increased expression of the basal markers *Ly6a* [23] and *Anxa1*,

which have been proposed as a marker of foetal reprogramming and regenerative stem cells in the

regenerating colonic epithelium and colorectal tumours (Gil Vazquez et al., 2022; Yui et al., 2018).

We also observed increased expression of the interleukins *Il24* and *Il6*. Interestingly, a recent model

for tissue injury-sensing uncovered the role of IL24, produced at wound sites in response to hypoxia,

in promoting epithelial proliferation and re-epithelialisation via pSTAT3 activation [31]. This pathway may

also extend to cSCC, as we see increased expression of the hypoxia sensor HIF1α and IL24 and pSTAT3

in our models (**Figure 4F)**. In addition, IL6 stimulates the release of proinflammatory cytokines from

the cutaneous microenvironment in response to wounding, activating TGFβ receptor signalling and the

STAT3 signal transduction pathway [31,32].

These transcriptomic results were validated by RNAscope for *Ly6a* and immunohistochemistry for

MYC, ANXA1 and pSTAT3 in our models (**Figure 4G and 4H**). Similar findings were seen in tumours

originating from the hair follicle stem cells upon MAPK signalling hyperactivation and deletion of the

TGFβ receptor 1 (*Alk5*) (Lgr5:BRAF$^{V600E}$-ALK5$^{fl/fl}$, Lgr5:HRAS$^{G12V/G12V}$-ALK5$^{fl/fl}$, Lgr5:KRAS$^{G12D/+}$-ALK5$^{fl/fl}$)[10]

and deletion of *Trp53* and *Notch2* (Lgr5:TRP53$^{fl/fl}$-NOTCH2$^{fl/fl}$) (**Figure S5B**).

Overall, these results reveal shared molecular and transcriptional hallmarks of epidermal

transformation independently of the cell-of-origin, tumour latency, or the oncogenic driver mutations.

## SOX2 is expressed in the tumour-resistant population

We next assessed whether transcriptional profiling could identify potential long-lasting markers of the

cell-of-origin of these tumours. Thus, we directly compared the transcriptomic profile of tumours from

K5/K14:BRAF$^{V600E}$ and Ivl:BRAF$^{V600E}$ model, which revealed only a handful of differentially expressed

genes (**Figure 5A**). Notably, SOX2 emerged as a key transcriptional difference, being highly enriched in

Ivl:BRAF$^{V600E}$ tumours, relative to K5/K14:BRAF$^{V600E}$.

SOX2 has been identified as a super pioneer transcription factor able to bind to close chromatin

through its ability to interfere with the maintenance of DNA methylation [22]. Furthermore, SOX2 is

commonly used as a marker of cancer stem cells and is upregulated in cSCC patients, where it rewires

cells for tumour initiation and growth [33,34]. We confirmed SOX2 enriched expression in Ivl:BRAF$^{V600E}$

compared to K5:BRAF$^{V600E}$, K14:BRAF$^{V600E}$ profile by immunohistochemistry (**Figure 5B**). Similar results

were obtained in HRAS-driven models where SOX2 expression was only seen in those arising from the

IVL+ population (**Figure 5C**). Meanwhile, SOX2 expression was negligible in normal skin during

homeostasis or in TPA-treated skin, suggesting that its expression is not linked to inflammation **(Figure**

**5D**). We then extended our investigation to a broader suite of models. We found SOX2 expressed in DMBA/TPA-induced tumours but not in tumours originating from *Lgr5+* cells (hair follicle bulge) driven by different oncogene combinations (**Figures S6A and S6B)**.

To highlight the cell-specific role of SOX2, we assessed the expression of SOX9, a different master regulator of chromatin accessibility that switches embryonic epidermal cells to a hair follicle stem cell fate and is constitutively active during tumorigenesis [35]. As reported, we found that SOX9 was expressed mainly in the hair follicle cells in healthy skin homeostasis, and its expression was increased in the basal layer of all the tumours irrespective of the cell of origin (**Figure S6C**). The suprabasal markers KRT4 and KRT13, commonly expressed in skin homeostasis, were also differentially expressed between the K5/K14:BRAF$^{V600E}$ and Ivl:BRAF$^{V600E}$ models (**Figure 5A**). However, mRNA expression levels did not correlate with increased protein levels, as we found no significant differences in immunostaining intensity across our models (**Figure S6D**).

These data indicate that once the tumour-resistant population expresses SOX2, it drives and mimics the transcriptional profile of the tumour-prime stem population required cell status for tumorigenesis.

## SOX2 overexpression is sufficient to drive tumorigenesis from the tumour-resistant population

Given the potential aetiological activation of SOX2 in skin tumorigenesis, we assessed whether it is required for cSCC development in a cell-specific manner. We crossed the K14:BRAF$^{V600E}$ and Ivl:BRAF$^{V600E}$ models with mice harbouring a conditional *Sox2* knockout allele (SOX2$^{fl}$) (**Figure 6A**). While conditional *Sox2* deletion did not affect tumorigenesis or tumour growth in the K14:BRAF$^{V600E-}$ SOX2$^{fl/fl}$ model (**Figure 6B**), it significantly delayed tumour onset and tumour growth in the Ivl:BRAF$^{V600E-}$Sox2$^{fl/fl}$ model (**Figure 6C**). Loss of *Sox2* had no impact on skin homeostasis in either population, consistent with its lack of expression in normal skin. Similarly, the deletion of *Sox2* had no effect on tumour growth in the K14 model but delayed tumour growth in the IVL model (**Figures 6D and 6E**)

The histopathological features of K14:BRAF$^{V600E-}$SOX2$^{fl/fl}$ and Ivl:BRAF$^{V600E-}$ SOX2$^{fl/fl}$ tumours were similar to those expressing the wild-type *Sox2* allele, including an enlarged epidermis displaying vertical columns of keratinocytes, incomplete maturation of keratinocytes, increased proliferation indicated by Ki67 immunoreactivity mainly in the basal layer, and MYC and pSTAT3 activation (**Figure S7A**). Interestingly, the rare Ivl:BRAF$^{V600E-}$ SOX2$^{fl/fl}$ tumours show patches escaping recombination and clones that retain SOX2 expression, as demonstrated by immunohistochemistry, which highlights its critical role in tumorigenesis (**Figure S7B**). SOX2+ escapee patches were not observed in the K14:BRAF$^{V600E-}$SOX2$^{fl/fl}$ tumours (**Figure S7B**).

Then, we assessed whether the gain of function of SOX2 could render IVL+ cells competent to
tumorigenesis when coupled to a BRAF mutation. To this end, we interbred an allele which allows the
inducible overexpression of SOX2 from the Rosa26 locus [*Rosa26*LSL-SOX2-IRES-eGFP (hereafter
SOX2$^{LSL}$)] to the tumour-resistant population in combination with BRAF$^{V600E}$ in a tamoxifen-inducible
manner. Upon SOX2 overexpression, the oncogene-targeted IVL+ population gave rise to fast-growing
tumours. SOX2 significantly accelerated tumour onset from 65 to 33 days (**Figure 6F)**, while SOX2
overexpression on its own showed no phenotype (**Figure 6C**).

The Ivl:BRAF$^{V600E}$-SOX2$^{LSL}$ tumours show advanced histological features of cSCC, including poorly
differentiated epithelial, keratin pearls, parakeratosis and invasion. These tumours show high and
disorganised proliferation beyond the basal layer and activation of the previously defined shared
transcriptional profile including pSTAT3, ANXA1 and MYC together with high expression of SOX2 (**Figure
6G**)

Overall, these data indicate that SOX2 is a cell-specific requirement for the IVL+ tumour-resistant
population to acquire the competence to initiate cSCC. Thus, SOX2 gain of function in this population
in combination with an oncogene leads to a clonal behaviour similar to the tumour-primed population
and mediates the re-acquisition of stemness in the skin basal compartment.

## SOX2 drives developmental and juvenile epidermal programs and hints at the cell-of-origin

To determine whether SOX2 expression controls stemness in the skin in the absence of oncogene
expression, we first immortalised two healthy human epidermal keratinocyte cell lines, derived from
a juvenile and an adult donor, and subsequently overexpressed SOX2 in the derivative cell lines (**Figures
7A and S8A**). Using these human cell lines, we study the transcriptomic changes induced by SOX2
overexpression in homeostatic conditions.

Interestingly, while SOX2 overexpression triggered transcriptomic changes in the adult cell line,
separating it from the parental cell line, only small transcriptomic changes were seen in the juvenile
SOX2-expressing cell line that clustered with its parental line (**Figures 7B-7D**). The profound SOX2-
driven changes in the adult transcriptome included significant enrichment for the embryonic and
juvenile epidermal signatures (**Figure 7E**). However, this enrichment was not observed in the SOX2
overexpressing juvenile cell line (**Figure 8B**). This could be due to the juvenile cell line resembling a
stem-like population already expressing an embryonic/juvenile-like transcriptome, hence the lack of
impact of SOX2 in these cells. These results resembled what we had seen in the models above; that
the tumour-prime stem-cell population is not altered by the expression of SOX2, while in the
differentiation committed tumour-resistant population, it drives the re-expression of stemness.

Finally, we sought to study how these data translate to human patients. We reanalysed a human dataset containing transcriptional data from different stages of disease progression [17]. We observed that *SOX2* levels were increased in 27% of premalignant actinic keratosis (AK) and 35% of cSCC lesions, whereas *SOX2* expression levels were low in normal skin (NS) (**Figure 7F and 7G**). SOX2 levels did not correlate with the age of the patient or tumour aggressiveness (tumour depth or diameter) (**Figure S8C**). Furthermore, we divided these samples into SOX2 high and low groups and conducted further transcriptional comparisons, however, we did not see any major transcriptional changes either in the AK or cSCC groups (**Figures S8D and S8E**). Moreover, we looked at SOX2 protein expression in a tissue microarray containing 250 human cSCC[36] , which ranged from no expression, in the majority of the samples, to mid and high in 21.7% of the histocores (**Figure 7H**). Those results are in line with the ~25% of human cSCC expressing high or medium levels of SOX2 reported previously [33,34].

This data aligns with the hypothesis that SOX2 is required to drive the re-expression of the stemness program needed to transform the tumour-resistant population, whereas it is not required for the transformation of the already stem-like basal population. Once SOX2 has rewired the epigenome and driven the embryonic and stem-cell pathways needed to trigger tumorigenesis, the tumours that result are indistinguishable from those originating from a stem population (**Figure 7I**).

These results underscore a diverse cell origin for cSCC, where SOX2 marks a subset of tumours emerging from the committed tumour-resistant population.

# Discussion

The heterogeneity of the interfollicular epidermis basal layer is well established. However, considerable debate remains as to whether it comprises two distinct progenitors or two populations that relate to different levels of commitment towards differentiation[19–21,37]. Our research uncovers the mechanism underlying the different tumour-initiating capacities following oncogenic mutations of both populations and identifies SOX2 as a critical enabler that renders committed progenitors susceptible to cSCC initiation.

Importantly, we generated novel genetically engineered mouse models that resemble the rapid cSCC development seen in melanoma patients treated with BRAFi, due to paradoxical MAPK signalling activation [13–15]. BRAFi promoted the development of tumours from cells carrying HRAS$^{G12V}$ mutations that had been sitting in the skin for up to 180 days without driving transformation. This shows that the skin can tolerate mutations in oncogenic pathways with little histological change but can rapidly progress to tumorigenesis given a specific stimulus or promoter.

We uncovered the mechanisms driving the difference in oncogene permissiveness by modelling MAPK

signalling hyperactivation in the stem-cell K14/K5+ population and the committed progenitor IVL+

population. In particular, we obtained rapidly developing tumours within days of oncogene induction

in the K14/K5+; hence, we called this population tumour-primed. However, oncogene induction in the

IVL+ differentiation committed basal population resulted in homeostasis and normal-looking skin

despite the oncogene-bearing cells colonising the entire epithelium within a week. Thus, we termed

this population tumour-resistant. Expression of HRAS$^{G12V}$ or KRAS$^{G12D}$ oncogenes in the skin epithelium

also outcompeted the wild-type population in homeostasis [38]. Furthermore, oncogene expression in

the IVL+ cells increased their residence in the basal layer, preventing delamination and allowing for a

longer latency in the progression to tumorigenesis. Once tumours developed, they resembled those

originating from K5+/K14+ cells.

Transcriptional profiling of the tumours arising from an IVL+ or a K5+/K14+ revealed a common

transcriptional landscape that is enriched for an activated wound-healing response [23], embryonic

developmental pathways [24] and a tumour-specific keratinocyte population seen in human cSCC [25]. In

addition, we observed strong MYC activation across all tumours, which has been proposed as a

mechanism to confer stem cell properties and dedifferentiation in the *Gata6*+ epidermal populations

[5]. At the same time, we revealed evidence of hijacking the tissue injury-sensing and repair pathway,

triggered by hypoxia and driven by IL24 and STAT3 activation that resolves in a coordinated tissue

repair after wounding [31]. Similar to wounds, the core of a tumour is hypoxic, and we observed

increased transcription of *Hif1a* and *Il24*, as well as STAT3 phosphorylation that enhanced oncogene-

driven proliferation to generate the critical tumour mass required for progression. These results

highlight the close interplay between proliferation during tumorigenesis and re-epithelisation during

wound sealing.

Much of our understanding of cSCC in the mouse has come from seminal work using DMBA/TPA[2,11] or

more lately mouse models that have focussed on using stem cell /progenitor to drive KRAS or other

MAPK mutations in combination with TP53[8,10,17]. These models typically result in widespread

oncogenic mutations across the skin, yet only a single tumour often emerges, mirroring human studies

that highlight the skin's ability to tolerate oncogenic mutations. Despite the diversity of oncogenic

events across these models, they converge on a core set of skin transformation pathways, as shown

here. These pathways can be rapidly triggered by BRAF mutation or by paradoxical MAPK activation

via BRAFi in HRAS-mutant cells, leading to tumorigenesis within days or over months to years in models

with longer latency. We propose that the acquisition of this oncogenic programme —also seen in

human cSCC[25] and inducible by inflammation, injury or paradoxical activation of MAPK— allows the

transformation of "normal" epithelial-containing oncogenes into a tumour without requiring
additional oncogenic events.

One of our key findings was that SOX2 activation in the tumour-resistant population is sufficient to
render this population susceptible to transformation. Knocking out this gene *in vivo* delayed tumour
onset and growth but did not affect the tumour-primed population. Conversely, SOX2 overexpression
in combination with oncogene expression rapidly accelerated tumorigenesis *in vivo,* showing that SOX2
is both necessary and sufficient for transforming the IVL+ lineage.

The limited capacity of the IVL+ basal population to switch to symmetric division and its tendency to
preferentially undergo asymmetric division is part of its transcriptome-engrained commitment
towards differentiation[19,21] and could explain its tumour-resistant phenotype also seen in basal cell
carcinoma[37]. Indeed, in basal cell carcinoma, only the stem-cell population and not the progenitors
have been proposed to initiate tumorigenesis[37]. However, we argue that both populations (stem-like
and committed) can trigger skin tumorigenesis.

There is evidence that SOX2 activation happens in ~20% of cSCC patients[33,34]. This suggests that the
IVL+ basal population acts as a cell-of-origin for these tumours and that SOX2 may serve as a marker
for classifying human cSCC according to the cell of origin. Still, the committed progenitors are tumour-
resistant, and they need to re-acquire the stemness features to stop differentiation and apoptosis and
allow transformation. Thus, this requirement of a stemness transcriptome, acquired by SURVIVIN
expression in basal cell carcinoma[37] or by SOX2 in cSCC reconciliates our results. The convergent
"stemness" state required for tumour initiation is also found in tumours developed from the
DMBA/TPA model [2].

Although the role of SOX2 in cSCC initiation has been previously described[33,34], our study shows that
its expression is only required and essential in the tumour-resistant population, where it confers the
stemness required to overcome their inability to initiate cSCC. SOX2 is a super pioneer factor able to
bind closed chromatin and rewire the epigenome to drive stem cell features [22,39]. Indeed, we show that
SOX2 overexpression in homeostasis can drive embryonic and juvenile transcriptomes. Thus, SOX2
would allow the tumour-resistant population to suppress, freeze, or completely abolish its
commitment towards differentiation by activating the stemness profile required for transformation.
IVL+ cells may dedifferentiate and become IVL- to drive tumorigenesis and to remain in the basal layer,
where the proliferative environment is confined. Indeed, cancer stemness is not restricted to the

unidirectional route (stem cell to progenitors) that applies in normal tissues, as shown by therapeutic

agents inducing different differentiation states[2].

We hypothesise that the more fluid and gradual differentiation mechanism recently proposed[21] could

be reverted towards dedifferentiation as the committed cells residing in the basal layer have not

necessarily exited the cell cycle. This model could explain the almost identical transcriptomic profile

obtained from tumours derived from the K14/K5+ population and the Ivl+ population, as both

populations will go on to delaminate, which argues against the presence of two distinct epidermal

populations.

How SOX2 is activated by oncogenic mutations is still elusive. It may be directly or indirectly

downstream of the commonly activated pERK or MYC signalling pathways. A deep understanding not

only of the tissue-specific mechanisms that drive transformation but also of the cell-specific

vulnerabilities that govern epithelial maintenance in homeostasis and tumorigenesis will allow the

development of rational cancer therapies to contain tumours at their source.

Collectively, we have generated new genetically driven *in vivo* models to study cSCC and identified

SOX2 as a key regulator of the different susceptibility of the tumour-primed and tumour-resistant skin

populations. SOX2 expression in oncogene-carrying tumour-resistant clones increases their residence

time in the basal layer and prevents delamination by driving a stem-like transcriptomic profile that

enables this population to initiate tumorigenesis at a longer latency.

# Online Methods

## Genetically engineered mouse models and husbandry

Mice were housed in accordance with UK Home Office Regulations, maintained in a pathogen-free

facility under a 12-h light-dark cycle at a constant temperature between 19–23°C, and 55 ± 10%

humidity, and given drinking water and fed standard chow diet *ad libitum*. Experiments conducted at

the CRUK Scotland Institute Animal Facility were in accordance with the UK Home Office guidelines,

under project licence PP3908577, and were reviewed and approved by the University of Glasgow

Animal Welfare and Ethical Review Board (AWERB). Experiments conducted at the CRUK Manchester

Institute were performed at Alderley Park Animal Research Unit, with breeding at the University of

Manchester incubator breeding facility, in accordance with UK Home Office regulations, under project

licences PE4369EDB and P671A5B06, and reviewed and approved by the CRUK Manchester Institute's

AWERB. Animals were monitored regularly until terminally sacrificed by schedule 1 at clinical endpoint:

either when the cumulative tumour burden had reached a maximum volume of 1500 mm$^3$ or when a

single tumour exceeded 15 mm in diameter, weight loss >20%, or any other signs of ill health and

distress. For all mouse studies, no formal randomisation was performed, and researchers were not

blinded to the mouse genotypes. No exclusions were performed and all mice included in experimental

cohorts were included in the analysis.

Genotyping was performed by Transnetyx according to the previously published protocols provided in

the references for each allele. Mice were maintained on a C57BL/6J background, carrying the following

alleles/transgenes: *Ivl*-CreERT2 [8], *Krt5*-CreERT2 [40], *Krt14*-CreERT2 [41], BRAF$^{V600E}$ [42,43], HRAS$^{G12V}$ [44],

*Rosa26LSL-tdRFP* [45], *Sox2$^{fl}$* [46] and *Rosa26LSL-SOX2-IRES-eGFP* [47]. Archived tumour blocks [from[10] ]

included the additional following alleles: *Lgr5*-CreERT2 [48], *Kras*$^{G12D}$ [49], *Notch2$^{fl}$* [50], and *Trp53$^{fl}$* [51].

Genetic recombination was induced in mice of either sex 8-12 weeks of age by topical application to

the shaved backs of the mice of 100 μL tamoxifen (10 mg/ml in ethanol; T5648, Sigma) in models

driven by Ivl-CreERT2, or 4 μl 4-hydroxytamoxifen (20 mg/ml in ethanol; H6278, Sigma) in models

driven by *Krt5*-CreERT2 and *Krt14*-CreERT, unless otherwise stated. This was repeated three more

times over 8 days.

## UV irradiation

UV irradiation experiments were conducted at the CRUK Manchester Institute husbandry facility

following the UK Home Office regulations under project licences PE4369EDB and P671A5B06. In

cohorts exposed to UVR, mice were subjected to UVR exposure once per week for 4 weeks, beginning

4 weeks after the final tamoxifen application. For this process, mice were anaesthetised by

intraperitoneal injection with 1 mg/kg Domitor and 100 mg/kg ketamine, with 5 mg/kg Antisedan

anaesthetic reversal. The backs were shaved, and a black cloth was used to cover regions that were to

remain unexposed. The Waldmann UV181 unit with UV6 broad wavelength (280-380 nm) lamp was

used for UVR exposure and the intensity was tested regularly with a USB2000+ spectroradiometer

(Ocean Optics). The UV dose used was 0.6 kJ/m$^2$, which equated to 3 minutes and 14 seconds of

exposure. Following irradiation, E45 moisturising cream was topically applied to the back.

## Tumour fragment implantation

Tumour implantation experiments were conducted at the CRUK Manchester Institute husbandry

facility following the UK Home Office regulations under project licences PE4369EDB and P671A5B06.

A tumour derived from a K5:BRAF$^{V600E}$ mouse was resected upon clinical endpoint under a laminar flow

hood aseptically. The tumour was cut into 2-3 mm$^3$ pieces for immediate implantation on anesthetised

6-8 week-old C57/6J littermates and NSGII2 mice. Briefly, recipient animals were anesthetised with

isoflurane and kept on a heated stage. A tumour piece was implanted subcutaneously between

the skin and the peritoneal wall. The surgical incision was closed using a surgical clip.

Rymadyl/Carprofen analgesic was given at 4mg/kg. Mice were monitored until consciousness was

regained and then monitored daily. The surgical clip was removed 7 days post-surgery.

## *In vivo* drug treatments

494 *In vivo* drug treatments were conducted at the CRUK Scotland Institute Animal Facility following the

495 UK Home Office guidelines, under project licence PP3908577. Skin inflammation was induced by

496 topically applying 150 µl of TPA (31.25 µg/ml) in acetone (Sigma-Aldrich 16561-29-8) to shaved dorsal

497 skin three times per week until tumour signs appeared. The BRAFi Dabrafenib was administered by

498 daily oral gavage (30 mg/kg in 100 µl) during the experiment.

## Immunohistochemistry and immunofluorescence

Organs collected in 10% neutral buffered formalin were stored at room temperature for 20-28 hours,

followed by transfer to 70% ethanol and storage at 4°C until processing. All haematoxylin and eosin

(H&E), immunohistochemistry, co-immunofluorescence and *in situ* hybridisation staining were

performed on 4-µm formalin-fixed paraffin-embedded sections (FFPE) which had previously been

heated at 60°C for 2 hours.

The following antibodies were used on a Leica Bond Rx autostainer: KRT5 (905501, Biolegend), Ki67

(12202, Cell Signaling), SOX2 (14962, Cell Signaling) and pSTAT3 (9131, Cell Signaling). All FFPE sections

underwent onboard dewaxing (AR9222, Leica) and epitope retrieval using ER2 solution (AR9640, Leica)

for 20 minutes at 95°C. Sections were rinsed with Leica wash buffer (AR9590, Leica) before peroxidase

block was performed using an Intense R kit (DS9263, Leica) for 5 minutes. Sections were rinsed with

wash buffer before primary antibody application at an optimal dilution (KRT5, 1/1500; Ki67, 1/1000;

pSTAT3, 1/100; SOX2, 1/200). The sections were rinsed with wash buffer before the application of anti-

rabbit EnVision HRP-conjugated secondary antibody (K4003, Agilent) for 30 minutes. The sections were

rinsed with wash buffer, visualised using DAB, and counterstained with haematoxylin from the Intense

R kit.

FFPE sections for KRT14 (ab7800, Abcam), c-MYC (ab32072, Abcam), RFP (600-401-379, Rockland) and

SOX9 (AB5535, Millipore) staining were stained on an Agilent autostainer Link48. FFPE sections were

loaded into an Agilent pre-treatment module to be dewaxed and undergo heat-induced epitope

retrieval (HIER) using a High pH target retrieval solution (K8004, Agilent). After HIER, the sections were

rinsed in FLEX wash buffer (K8007, Agilent) before being loaded onto the Agilent autostainer. The

sections underwent peroxidase blocking (S2023, Agilent) for 5 minutes and were rinsed with FLEX

buffer. The primary antibody application was at an optimised dilution (KRT14, 1/300; c-MYC, 1/800;

RFP, 1/1000; SOX9, 1/500). Sections were washed with FLEX buffer before application of anti-rabbit

EnVision secondary antibody for 30 minutes. Sections were rinsed with FLEX wash buffer before

applying Liquid DAB (K3468, Agilent) for 10 minutes. Sections were washed in water and counterstained with haematoxylin 'Z' (RBA-4201-00A, CellPath).

In-situ hybridisation detection for *Anxa1* (509298), *Ivl* (422538), *Ly6a* (427578), PPIB (313918; positive control) and dapB (312038; negative control) (all Bio-Techne) mRNA was performed using RNAScope 2.5 LSx (Brown) detection kit (322700; Bio-Techne) according to the manufacturer's instructions. H&E staining was performed on a Leica autostainer (ST5020). Sections were dewaxed in xylene, taken through graded ethanol solutions and stained with haematoxylin 'Z' (RBA-4201-00A, CellPath) for 13 minutes. Sections were washed in water, differentiated in 1% acid alcohol, washed and the nuclei stained blue in Scott's tap water substitute (in-house). After washing with tap water, sections were placed in Putt's eosin (in-house) for 3 minutes. To complete H&E, IHC & ISH staining, sections were rinsed in tap water, dehydrated through a series of graded alcohols, and placed in xylene. The stained sections were coverslipped in xylene using DPX mountant (SEA-1300-00A, CellPath).

Sections for KRT14 and RFP co-IF staining were loaded onto a Leica Bond Rx autostainer. The FFPE sections underwent on-board dewaxing and epitope retrieval using ER2 solution for 20 minutes at 95°C. Sections were rinsed with Leica wash buffer (AR9590, Leica) before application of 10% normal goat serum (X090710-8, Agilent) for 30 minutes. The sections were rinsed with Leica wash buffer before application of anti-RFP antibody at 1/1000 dilution for 1 hour. Sections were rinsed with Leica wash buffer and goat anti-rabbit IgG 488 (A11034, Invitrogen) secondary antibody diluted 1/250 for 30 minutes. After rinsing with Leica wash buffer KRT14 antibody was applied at 1/750 dilution for 1 hour. Sections were rinsed with Leica wash buffer and goat anti-mouse IgG 647 secondary antibody (A21236, Invitrogen diluted) 1/250 for 30 minutes before application of DAPI (MBD0015, Sigma-Aldrich). To complete the staining, sections were mounted using ProLong Diamond antifade mountant (P36970, Thermo Fisher Scientific).

Immunohistochemistry images were acquired using a SCN400F slide scanner (Leica Microsystems) at ×20 or x40 magnification. Confocal images were collected on a Zeiss 710 point-scanning confocal microscope, built on an inverted Zeiss Axio Imager.Z2 stand. Images were acquired using a EC Plan-Neofluar 40x/1.30 Oil and a confocal pinhole diameter of 107 μm. Multi-channel images were captured sequentially: DAPI (nuclear marker) using 405 nm excitation and 410-481 nm emission bandwidth, IgG 488 using 488 nm excitation and 514-582 nm emission and IgG 647 using 633 nm excitation and 638-747 nm emission. Images were collected with a 1x zoom with an image size of 2320 x 2320 pixels, yielding a pixel size of 92 x 92 nm, and a 1.38 us pixel dwell time. Z-stacks were collected using a step size of 2 um. Images were acquired using the software Zen LSM 2.1 Black (Zeiss).

## RNAseq sequencing and analysis

RNA was extracted from fresh frozen tumour samples or human cell line pellets using an AllPrep DNA/RNA kit (74104, Qiagen) according to the manufacturer's instructions. Indexed poly(A) libraries were prepared using 200 ng of total RNA and 14 cycles of amplification with the Agilent SureSelect Strand-Specific RNA Library Preparation Kit for Illumina Sequencing (G9691B, Agilent). RNA polyA libraries were sequenced using NovaSeq 6000 XP SP, 200 cycles, paired-end reads (2 x 101 bp) 20-30 million reads. RNA preparation and sequencing of mouse samples were performed at the CRUK Manchester Institute histology core and sequencing facility. RNA preparation and sequencing of human cell lines were performed at GENEWIZ Azenta.

For mouse samples raw sequence quality control was performed using FastQC (v0.11.8) before and after removing adapters and low-quality base calls (Phred score <20) using TrimGalore with default options (v0.6.6). Trimmed reads were aligned to GRCm38, release 100 using Hisat2 (v2.1.0). Gene counts were subsequently estimated using featurecounts (subread/1.6.3). For human cell lines, raw sequence quality control and alignment were performed using Nextfow nf-core/rnaseq pipeline using STAR and Salmon [52]. Trimmed reads were aligned to GRCh38, release 113. Sva::ComBat-seq was used for batch correction (v3.52.0). For all samples, after removing transcripts without a minimum of 5 reads in at least one sample, the differential expression analysis between mouse tumours and skin was performed using the R package DESeq2 (v1.44.0). The resultant p-values were corrected for multiple comparisons using the Benjamini-Hochberg approach. The following additional R packages were used for downstream analysis: pheatmap (v1.0.8), fgsea (v1.30.0), enhancedVolcano (v1.22.0), clusterProfiler (v4.12.6), dplyr (v1.1.4), tidyr (v1.3.1), AnnotationHub (v3.12.0), ggvenn (0.1.10), AnnotationDbi (v1.66.0), mixOmics (v6.26.0).

## Culture of human keratinocyte cell lines and introduction of genetic modifications

Human keratinocyte cell lines were maintained in serum-free Keratinocyte Growth Medium 2 (C-20011, PromoCell) until confluent and then split using 0.25% Trypsin-EDTA (T4049-100ML, Merck) and quenched using Defined Trypsin Inhibitor (R007100, Gibco). All cell culture incubators were set to 37°C and 5% CO2.

Normal human primary juvenile keratinocytes (JHEK) (C-12006, PromoCell) and normal human primary adult keratinocytes (AHEK) (102-05a, Cell Applications) were immortalised with in-house produced lentivirus expressing the hTERT-hygromycin resistant pLV-EF1a-IRES-Hygro plasmid and selected with hygromycin (300 µg/ml). The plasmid was a gift from Tobias Meyer (85140, Addgene) [53]). Then, nucleofection-based CRISPR-Cas-9 methodology was used to generate p16 KO using the 4D Nucleofector Core Unit (AAF-1002B with AAF-1002X, Lonza). A gRNA with enhanced stability targeting

*Cdkn2a* (p16) was designed and purchased from Integrated DNA Technologies (IDT). Ribonucleoprotein

(RNP) complexes were formed by diluting 120 pmol sgRNA, 104 pmol Cas9 Nuclease V3 (1081059,

IDT), 1 μl of Alt-R Cas9 Electroporation Enhancer (1075916, IDT) in Nuclease-Free IDTE, pH 7.5 (1X TE

solution) (I11-01-02-02, IDT). RNP complex was added to 20 μl of Primary Cell P3 Nucleofector Solution

(V4XP-3032, Lonza) and nucleofected using programme CM137. The mixture was then transferred to

a 20 μl nucleocuvette strip and nucleofected using programme CM137. The targeted sequence was

PCR-amplified, sequenced at Eurofins and KO-validated using the TIDE algorithm (achieving >90%

efficiency)[54]. *SOX2* overexpression lentivirus with puromycin resistance was purchased from

VectorBuilder ready-to-use (vector id VB900088-3008dcy). SOX2-overexpressing keratinocytes were

selected with 1 μg/ml puromycin. GFP overexpression control lentivirus with puromycin resistance

purchased from VectorBuilder ready-to-use were used as controls. Cell pellets were regularly collected

and submitted for cell line authentication at Eurofins and tested for Mycoplasma in-house.

## Immunoblotting

Proteins were extracted from whole-cell lysates using RIPA buffer (R0278, Sigma-Aldrich) mixed with 1

tablet of cOmplete™ Protease Inhibitor Cocktail (11873580001, Sigma-Aldrich). Cell suspensions were

incubated on ice for 10 minutes before centrifuging at 13,000 rpm for 10 minutes at 4°C. Protein

extracts from the supernatants were quantified using the Pierce™ BCA Protein Assay Kit (23225,

Invitrogen). Protein extracts (25 μg) were denatured in 4x Laemmli sample buffer (1610747, BIO-RAD)

at 95 °C for 5 minutes. The samples were separated on NuPAGE™ 4-12% Bis-Tris Mini Protein Gels

(NP0322BOX, Invitrogen) at 120 V for 1 hour in 1x MOPS SDS running buffer (B0001, Invitrogen)

together with the Chameleon Duo pre-stained Protein Ladder (928-60000, LI-COR). The proteins were

transferred onto a nitrocellulose membrane (170-4270, BIO-RAD), using the BIO-RAD Trans-Blot Turbo

Transfer System. The membrane was blocked with 1x TBST with 5% BSA (A7906, Sigma-Aldrich) for 1

hour at room temperature in the dark. The membrane was then incubated in anti-SOX2 antibody

1:1000 (23064, Cell Signalling) and anti-β-actin 1:10,000 (MA1-140, Invitrogen) diluted in 1x TBST 5%

BSA, at 4°C overnight in the dark. β-actin was used as a loading control. Followed by 3x10 minute

washes in 1x TBST in the dark and subsequently incubated in fluorescence-conjugated secondary

antibodies, diluted in 5% BSA in 1x TBST, at room temperature for 45 minutes in the dark. Secondary

antibodies used were anti-rabbit IgG (A32735, Thermo Fisher Scientific) and anti-mouse IgG

(A21057, Thermo Fisher Scientific). This was followed by 3x10 minute washes in 1x TBST before the

membrane was scanned using the LI-COR Odyssey® CLx Infrared Imaging System.

## Data availability

The data discussed in this publication have been deposited in NCBI's Gene Expression Omnibus. The accession numbers for the RNA-seq data reported in this paper are NCBI GEO GSE280236 (mouse data) and NCBI GEO GSE281000 (human cell lines data). No new code was generated in this publication.

## Acknowledgements

We thank the Core Services and Advanced Technologies at the Cancer Research UK Scotland and Manchester Institutes, particularly the Biological Services Unit, Histology Service and Molecular Technologies. We also thank Claire Mitchell and the BAIR Biological Advance Image Resource service at Cancer Research UK Scotland Institute. We thank Kevin Haigis (Dana-Farber Cancer Institute) for providing the HRAS[G12V] mouse model. We are grateful to Sarah Ressel and Shan Quah for the critical review of the manuscript and to Nathalie Sphyris for expert writing advice (Cancer Research UK Scotland Institute). This work was supported by Cancer Research UK Cancer Grand Challenges and The Mark Foundation for Cancer Research to the SPECIFICANCER team (A29055) and by CRUK core funding to the Cancer Research UK Manchester Institute (C5759/A27412) and the Cancer Research UK Scotland Institute (C5759/A31287). P.P.C. and C.C. were funded by the Cancer Research UK Grand Challenge SPECIFICANCER Consortium (A29055 & A27412). C.A.F, R.A.R., P.C., T.J, O.J.S. were supported by CRUK Beatson Institute core funding (A17196, A31287).

## Author Contributions

P.P.C., A.C., R.M. and O.J.S. designed experiments and interpreted results. P.PC, C.C., C.A.F., R.A.R., T.J.S. and P.C. performed experiments and analysed results. P.P.C. analysed publicly available human cancer data sets. P.P.C. processed and analysed the RNA sequencing data. P.P.C., A.C and O.J.S. wrote the paper, and reviewed and discussed the drafted manuscript. All authors contributed to the manuscript.

## Competing Interest

The O.J.S. laboratory receives funding from Novartis, Cancer Research Technology (Cancer Research Horizons), Boehringer Ingelheim and AstraZeneca for other unrelated projects. R.M. is the founder and CEO of MyT Bioscience Ltd and the founder, Director and CSO of Oncodrug Ltd. The remaining authors declare no competing interests.

# REFERENCES

1. Martincorena I, Roshan A, Gerstung M, et al. High burden and pervasive positive selection of somatic mutations in normal human skin. *Science (1979)*. 2015;348(6237):880-886. doi:10.1126/science.aaa6806

2. Taylor MA, Kandyba E, Halliwill K, et al. Stem-cell states converge in multistage cutaneous squamous cell carcinoma development. *Science*. 2024;384(6699):eadi7453. doi:10.1126/SCIENCE.ADI7453

3. Chang D, Shain AH. The landscape of driver mutations in cutaneous squamous cell carcinoma. *NPJ Genom Med*. 2021;6(1):61. doi:10.1038/s41525-021-00226-4

4. Ge Y, Gomez NC, Adam RC, Yuan S, Elemento O, Fuchs Correspondence E. Stem Cell Lineage Infidelity Drives Wound Repair and Cancer. *Cell*. 2017;169:636-650. doi:10.1016/j.cell.2017.03.042

5. Bernabé-Rubio M, Ali S, Bhosale PG, et al. Myc-dependent dedifferentiation of Gata6+ epidermal cells resembles reversal of terminal differentiation. *Nature Cell Biology 2023*. Published online September 21, 2023:1-13. doi:10.1038/s41556-023-01234-5

6. Aragona M, Sifrim A, Malfait M, et al. Mechanisms of stretch-mediated skin expansion at single-cell resolution. *Nature 2020 584:7820*. 2020;584(7820):268-273. doi:10.1038/s41586-020-2555-7

7. Page ME, Lombard P, Ng F, Göttgens B, Jensen KB. The Epidermis Comprises Autonomous Compartments Maintained by Distinct Stem Cell Populations. *Cell Stem Cell*. 2013;13(4):471-482. doi:10.1016/j.stem.2013.07.010

8. Lapouge G, Youssef KK, Vokaer B, et al. Identifying the cellular origin of squamous skin tumors. *Proceedings of the National Academy of Sciences*. 2011;108(18):7431-7436. doi:10.1073/pnas.1012720108

9. Kretzschmar K, Weber C, Driskell RR, Calonje E, Watt FM. Compartmentalized Epidermal Activation of β-Catenin Differentially Affects Lineage Reprogramming and Underlies Tumor Heterogeneity. *Cell Rep*. 2016;14(2):269-281. doi:10.1016/J.CELREP.2015.12.041

10. Cammareri P, Rose AM, Vincent DF, et al. Inactivation of TGFβ receptors in stem cells drives cutaneous squamous cell carcinoma. *Nat Commun*. 2016;7(1):12493. doi:10.1038/ncomms12493

11. Quintanilla M, Brown K, Ramsden M, Balmain A. Carcinogen-specific mutation and amplification of Ha-ras during mouse skin carcinogenesis. *Nature 1986 322:6074*. 1986;322(6074):78-80. doi:10.1038/322078a0

12. Nassar D, Latil M, Boeckx B, Lambrechts D, Blanpain C. Genomic landscape of carcinogen-induced and genetically induced mouse skin squamous cell carcinoma. *Nat Med*. 2015;21(8):946-954. doi:10.1038/nm.3878

13. Oberholzer PA, Kee D, Dziunycz P, et al. RAS Mutations Are Associated With the Development of Cutaneous Squamous Cell Tumors in Patients Treated With RAF

Inhibitors. *Journal of Clinical Oncology*. 2012;30(3):316-321. doi:10.1200/JCO.2011.36.7680

14. Hauschild A, Grob JJ, Demidov L V., et al. Dabrafenib in BRAF-mutated metastatic melanoma: A multicentre, open-label, phase 3 randomised controlled trial. *The Lancet*. Published online 2012. doi:10.1016/S0140-6736(12)60868-X

15. Su F, Viros A, Milagre C, et al. *RAS Mutations in Cutaneous Squamous-Cell Carcinomas in Patients Treated with BRAF Inhibitors A Bs t r Ac t*. Vol 366.; 2012. doi:2012;366:207-15

16. Poulikakos PI, Zhang C, Bollag G, Shokat KM, Rosen N. RAF inhibitors transactivate RAF dimers and ERK signalling in cells with wild-type BRAF. *Nature*. 2010;464(7287):427-430. doi:10.1038/nature08902

17. Bailey P, Ridgway RA, Cammareri P, et al. Driver gene combinations dictate cutaneous squamous cell carcinoma disease continuum progression. *Nat Commun*. 2023;14(1):5211. doi:10.1038/S41467-023-40822-9

18. White AC, Tran K, Khuu J, et al. Defining the origins of Ras/p53-mediated squamous cell carcinoma. *Proceedings of the National Academy of Sciences*. 2011;108(18):7425-7430. doi:10.1073/pnas.1012670108

19. Mascré G, Dekoninck S, Drogat B, et al. Distinct contribution of stem and progenitor cells to epidermal maintenance. *Nature*. 2012;489(7415):257-262. doi:10.1038/nature11393

20. Sarate RM, Hochstetter J, Valet M, et al. Dynamic regulation of tissue fluidity controls skin repair during wound healing. *Cell*. 2024;187. doi:10.1016/j.cell.2024.07.031

21. Cockburn K, Annusver K, Gonzalez DG, et al. Gradual differentiation uncoupled from cell cycle exit generates heterogeneity in the epidermal stem cell layer. *Nature Cell Biology 2022 24:12*. 2022;24(12):1692-1700. doi:10.1038/s41556-022-01021-8

22. Vanzan L, Soldati H, Ythier V, et al. High throughput screening identifies SOX2 as a super pioneer factor that inhibits DNA methylation maintenance at its binding sites. *Nat Commun*. 2021;12(1):3337. doi:10.1038/s41467-021-23630-x

23. Joost S, Jacob T, Sun X, et al. Single-Cell Transcriptomics of Traced Epidermal and Hair Follicle Stem Cells Reveals Rapid Adaptations during Wound Healing. *Cell Rep*. 2018;25(3):585-597.e7. doi:10.1016/J.CELREP.2018.09.059

24. Sennett R, Wang Z, Rezza A, et al. An Integrated Transcriptome Atlas of Embryonic Hair Follicle Progenitors, Their Niche, and the Developing Skin. *Dev Cell*. 2015;34(5):577-591. doi:10.1016/J.DEVCEL.2015.06.023

25. Ji AL, Rubin AJ, Thrane K, et al. Multimodal Analysis of Composition and Spatial Architecture in Human Squamous Cell Carcinoma. *Cell*. 2020;182(2):497-514.e22. doi:10.1016/j.cell.2020.05.039

26. Zou DD, Sun YZ, Li XJ, et al. Single-cell sequencing highlights heterogeneity and malignant progression in actinic keratosis and cutaneous squamous cell carcinoma. *Elife*. 2023;12. doi:10.7554/eLife.85270

27. Bizzarro V, Belvedere R, Milone MR, et al. Annexin A1 is involved in the acquisition and maintenance of a stem cell-like/aggressive phenotype in prostate cancer cells with acquired resistance to zoledronic acid. *Oncotarget*. 2015;6(28):25074. doi:10.18632/ONCOTARGET.4725

28. Al-Ali HN, Crichton SJ, Fabian C, et al. A therapeutic antibody targeting annexin-A1 inhibits cancer cell growth in vitro and in vivo. *Oncogene 2024 43:8*. 2024;43(8):608-614. doi:10.1038/s41388-023-02919-9

29. Yui S, Azzolin L, Maimets M, et al. YAP/TAZ-Dependent Reprogramming of Colonic Epithelium Links ECM Remodeling to Tissue Regeneration. *Cell Stem Cell*. 2018;22(1):35-49.e7. doi:10.1016/J.STEM.2017.11.001

30. Gil Vazquez E, Nasreddin N, Valbuena GN, et al. Dynamic and adaptive cancer stem cell population admixture in colorectal neoplasia. *Cell Stem Cell*. 2022;29(8):1213-1228.e8. doi:10.1016/j.stem.2022.07.008

31. Liu S, Hur YH, Cai X, et al. A tissue injury sensing and repair pathway distinct from host pathogen defense. *Cell*. 2023;186(10):2127-2143.e22. doi:10.1016/j.cell.2023.03.031

32. Johnson BZ, Stevenson AW, Prêle CM, Fear MW, Wood FM. The Role of IL-6 in Skin Fibrosis and Cutaneous Wound Healing. *Biomedicines 2020, Vol 8, Page 101*. 2020;8(5):101. doi:10.3390/BIOMEDICINES8050101

33. Siegle JM, Basin A, Sastre-Perona A, et al. SOX2 is a cancer-specific regulator of tumour initiating potential in cutaneous squamous cell carcinoma. *Nat Commun*. 2014;5. doi:10.1038/NCOMMS5511

34. Boumahdi S, Driessens G, Lapouge G, et al. SOX2 controls tumour initiation and cancer stem-cell functions in squamous-cell carcinoma. *Nature*. 2014;511(7508):246-250. doi:10.1038/NATURE13305

35. Yang Y, Gomez N, Infarinato N, et al. The pioneer factor SOX9 competes for epigenetic factors to switch stem cell fates. *Nature Cell Biology 2023*. Published online July 24, 2023:1-11. doi:10.1038/s41556-023-01184-y

36. Rose AM, Spender LC, Stephen C, et al. Reduced SMAD2/3 activation independently predicts increased depth of human cutaneous squamous cell carcinoma. *Oncotarget*. 2018;9(18):14552-14566. doi:10.18632/ONCOTARGET.24545

37. Canato S, Sarate R, Carvalho-Marques S, et al. Survivin Promotes Stem Cell Competence for Skin Cancer Initiation. *Cancer Discov*. 2025;15(2):427-443. doi:10.1158/2159-8290.CD-24-0263

38. Gallini S, Annusver K, Rahman NT, et al. Injury prevents Ras mutant cell expansion in mosaic skin. *Nature 2023*. Published online June 21, 2023:1-9. doi:10.1038/s41586-023-06198-y

39. Schaefer T, Lengerke C. SOX2 protein biochemistry in stemness, reprogramming, and cancer: the PI3K/AKT/SOX2 axis and beyond. *Oncogene 2019 39:2*. 2019;39(2):278-292. doi:10.1038/s41388-019-0997-x

765    40.    Van Keymeulen A, Rocha AS, Ousset M, et al. Distinct stem cells contribute to
766            mammary gland development and maintenance. *Nature*. 2011;479(7372):189-193.
767            doi:10.1038/nature10573

768    41.    Vasioukhin V, Degenstein L, Wise B, Fuchs E. The magical touch: Genome targeting in
769            epidermal stem cells induced by tamoxifen application to mouse skin. *Proc Natl Acad
770            Sci U S A*. 1999;96(15):8551-8556. doi:10.1073/pnas.96.15.8551

771    42.    Mercer K, Giblett S, Green S, et al. Expression of endogenous oncogenic V600EB-raf
772            induces proliferation and developmental defects in mice and transformation of
773            primary fibroblasts. *Cancer Res*. 2005;65(24):11493-11500. doi:10.1158/0008-
774            5472.CAN-05-2211

775    43.    Dhomen N, Reis-Filho JS, da Rocha Dias S, et al. Oncogenic Braf Induces Melanocyte
776            Senescence and Melanoma in Mice. *Cancer Cell*. 2009;15(4):294-303.
777            doi:10.1016/j.ccr.2009.02.022

778    44.    Bender RHF, Haigis KM, Gutmann DH. Activated k-ras, but not h-ras or N-ras, regulates
779            brain neural stem cell proliferation in a raf/rb-dependent manner. *Stem Cells*.
780            2015;33(6):1998-2010. doi:10.1002/STEM.1990

781    45.    Madisen L, Zwingman TA, Sunkin SM, et al. A robust and high-throughput Cre reporting
782            and characterization system for the whole mouse brain. *Nat Neurosci*. 2010;13(1):133-
783            140. doi:10.1038/NN.2467

784    46.    Shaham O, Smith AN, Robinson ML, Taketo MM, Lang RA, Ashery-Padan R. Pax6 is
785            essential for lens fiber cell differentiation. *Development*. 2009;136(15):2567-2578.
786            doi:10.1242/DEV.032888

787    47.    Lu Y, Futtner C, Rock JR, et al. Evidence That SOX2 Overexpression Is Oncogenic in the
788            Lung. *PLoS One*. 2010;5(6):e11022. doi:10.1371/JOURNAL.PONE.0011022

789    48.    Barker N, Van Es JH, Kuipers J, et al. Identification of stem cells in small intestine and
790            colon by marker gene Lgr5. *Nature 2007 449:7165*. 2007;449(7165):1003-1007.
791            doi:10.1038/nature06196

792    49.    Jackson EL, Willis N, Mercer K, et al. Analysis of lung tumor initiation and progression
793            using conditional expression of oncogenic K-ras. *Genes Dev*. 2001;15(24):3243-3248.
794            doi:10.1101/GAD.943001

795    50.    McCright B, Lozier J, Gridley T. Generation of new Notch2 mutant alleles. *Genesis*.
796            2006;44(1):29-33. doi:10.1002/GENE.20181

797    51.    Jonkers J, Meuwissen R, Van der Gulden H, Peterse H, Van der Valk M, Berns A.
798            Synergistic tumor suppressor activity of BRCA2 and p53 in a conditional mouse model
799            for breast cancer. *Nat Genet*. 2001;29(4):418-425. doi:10.1038/NG747

800    52.    Ewels PA, Peltzer A, Fillinger S, et al. The nf-core framework for community-curated
801            bioinformatics pipelines. *Nature Biotechnology 2020 38:3*. 2020;38(3):276-278.
802            doi:10.1038/s41587-020-0439-x

803    53.    Hayer A, Shao L, Chung M, et al. Engulfed cadherin fingers are polarized junctional
804            structures between collectively migrating endothelial cells. *Nat Cell Biol*.
805            2016;18(12):1311. doi:10.1038/NCB3438

806    54.    Brinkman EK, Chen T, Amendola M, Van Steensel B. Easy quantitative assessment of
807            genome editing by sequence trace decomposition. *Nucleic Acids Res*.
808            2014;42(22):e168-e168. doi:10.1093/NAR/GKU936

811 FIGURES

## Figure 1

**Figure 1. Paradoxical MAPK signalling activation is a tumour promoter in cSCC.**

(A) Schematic representation of the mouse models used (top) and the epidermis (bottom) formed by the suprabasal, formed by IVL+ cells, and the basal layer, which includes a mixture of K14+/K5+ and IVL+ cells. Created with BioRender.

(B-C) Kaplan-Meier tumour-free survival plot for K14:HRAS$^{G12V/+}$ and those treated with the BRAFi Dabrafenib and TPA (B), and Ivl:HRAS$^{G12V/+}$ , Ivl:HRAS$^{G12V/G12V}$ and those treated with BRAFi Dabrafenib and TPA (C).

(D) Representative histological images including H&E and Ki67 immunohistochemistry (IHC) in tumours derived from K14:HRAS$^{G12V/+}$ and Ivl:HRAS$^{G12V/+}$ mice treated with the BRAFi Dabrafenib at clinical endpoint.

(E) Kaplan-Meier tumour-free survival plots for K14:HRAS$^{G12V/+}$ treated with BRAFi Dabrafenib 182 days after oncogene induction (left) and Ivl:HRAS$^{G12V/+}$ treated with BRAFi Dabrafenib 189 days after oncogene induction (right).

# Figure 2

**Figure 2: Tumour-primed and tumour-resistant populations coexist in the basal layer**

(A) Schematic representation of the mouse models used for skin tumorigenesis.

(B) Kaplan-Meier plot displaying tumour-free survival for K5:BRAF$^{V600E}$, K14:BRAF$^{V600E}$, Ivl:BRAF$^{V600E}$, and Ivl:BRAF$^{V600E}$ UV-treated female mice aged until clinical endpoint.

(C-E) Total tumour growth in K5:BRAF$^{V600E}$ and K14:BRAF$^{V600E}$ (C) and in Ivl:BRAF$^{V600E}$ and UV-treated Ivl:BRAF$^{V600E}$

(D) female mice and total tumour number (Mean and SEM) (E) in female mice aged until clinical endpoint.

(F-G) Representative H&E of K5:BRAF$^{V600E}$ (F) and Ivl:BRAF$^{V600E}$ (G) tumours at clinical endpoint showing keratin pearls are indicated (KP) and arrows indicate nuclear atypia.

# Figure 3

**Figure 3: Oncogene-expressing tumour-resistant population takes over the entire skin but remains tumour-restrictive.**

(A) Schematic representation of the mouse models used for lineage tracing.

(B) Schematic representation of the experimental design used for lineage tracing.

(C) Representative Z-stack co-immunofluorescence (IF) of KRT14 and RFP at day 8 and day 28 post tdRFP induction in K14:tdRFP and Ivl:tdRFP normal skin.

(D,G) Representative RFP IHC at different time points post-induction in K14:tdRFP and K14:BRAF$^{V600E}$-tdRFP (D) and Ivl:tdRFP and Ivl:BRAF$^{V600E}$-tdRFP (G).

(E,H) Quantification of the clonal persistence in the basal epidermis region of K14:tdRFP and

K14:BRAF$^{V600E}$-tdRFP (E), and Ivl:tdRFP and Ivl:BRAF$^{V600E}$-tdRFP (H) mice at different times post-

induction.

(F,I) Skin thickness quantification at different time points post-induction in K14:tdRFP and

K14:BRAF$^{V600E}$-tdRFP (F) and Ivl:tdRFP and Ivl:BRAF$^{V600E}$-tdRFP (I). At least 10 measurements from three

different skin stripes per animal. Points represent individual animals.  Unpaired T-test. NS not

significant P>0.05, ** p<0.01 and *** p<0.001.

# Figure 4

**Figure 4: Shared transcriptional profile independent of the cell-of-origin or oncogene**

(A) Principal component analysis (PCA) of normalised expression values showing the correlation between the transcriptional profiles of normal skin (NS, n=4), Ivl:BRAF[V600E] (IVL, n=6) and K5/K14:BRAF[V600E] (K5_K14, n=9).

(B, C) Volcano plots showing differentially expressed genes in IVL tumours (B) and K5_K14 tumours (C) compared with normal skin using the Wald test (two-tailed).

(D) Venn diagram showing upregulated genes shared between Ivl:BRAF[V600E] (Ivl) and K5/K14:BRAF[V600E] (K5_K14) tumours when compared with normal skin.

(E) Gene set enrichment analysis of Hallmarks and indicated pathways in K5_K14 tumours and IVL tumours.

Showing pathways significantly enriched (padj<0.01 and NES>2 based on an adaptive multi-level split Monte-

Carlo).

(F) Hierarchical clustering heatmap of selected genes showing normalised expression and correlation between

normal skin (NS), Ivl:BRAF$^{V600E}$ (Ivl) and K5/K14:BRAF$^{V600E}$ (K5_K14).

(G,H) Representative images of IHC validation of selected targets in K14:BRAF$^{V600E}$ and Ivl:BRAF$^{V600E}$ (G) and BRAFi

treated K14:HRAS$^{G12D/+}$ and Ivl:HRAS$^{G12D/+}$ tumours (H) at clinical endpoint.

**Figure 5: SOX2 is expressed in the tumour-resistant population**

871    (A) Volcano plot showing differentially expressed genes between K5/K14:BRAF$^{V600E}$ and Ivl:BRAF$^{V600E}$ tumours

872    using Wald test (two-tailed).

873    (B,C) Representative images of IHC validation of SOX2 in Ivl:BRAF$^{V600E}$ , K5:BRAF$^{V600E}$, and K14:BRAF$^{V600E}$ (B) and

874    BRAFi treated K14:HRAS$^{G12D/+}$ and Ivl:HRAS$^{G12D/+}$ tumours (C) at clinical endpoint.

875    (D) Representative images of IHC of SOX2 in normal skin (left) and normal skin treated with TPA  (right).

# Figure 6

**Figure 6: SOX2 overexpression is sufficient to initiate cSCC from the tumour-resistant population.**

(A) Schematic representation of the mouse models used for tumorigenesis.

(B-E) Kaplan-Meier tumour-free survival plot and tumour growth for K14:BRAF$^{V600E}$, K14:BRAF$^{V600E}$-SOX2$^{fl/fl}$ and K14:SOX2$^{fl/fl}$ (B,D), and for Ivl:BRAF$^{V600E}$, Ivl:BRAF$^{V600E}$-SOX2$^{fl/fl}$ , Ivl:SOX2$^{fl/fl}$ ,Ivl:BRAF$^{V600E}$-SOX2$^{LSL}$, and Ivl:SOX2$^{LSL}$ (C,E). P-values were determined using the log-rank (Mantel-Cox) test. NS not significant P>0.05, ** p<0.01. Note that K14:BRAF$^{V600E}$ and Ivl:BRAF$^{V600E}$ cohorts correspond to CRUK-SI cohorts also shown in  Figure S2 and are shown here for comparison.

(G) Representative H&E and IHC of Ki67, SOX2, MYC, pSTAT3 and ANXA1 in Ivl:BRAF$^{V600E}$-SOX2$^{LSL}$ tumour at clinical endpoint.

# Figure 7

(A) Western-blot validation of SOX2 overexpression in adult (AHEK) and juvenile (JHEK) human keratinocytes.

**Figure 7: SOX2 drives developmental and juvenile epidermal signatures and hints at the cell-of-origin for cSCC patients.**

(A) Western-blot validation of SOX2 overexpression in adult (AHEK) and juvenile (JHEK) human keratinocytes.

(B) Principal component analysis (PCA) of normalised expression values showing the correlation between the transcriptional profiles of JHEK, JHEK-SOX2, AHEK, and AHEK-SOX2. N=3, repeated independent experiments per group.

(C,D) Volcano plot showing differentially expressed genes between JHEK and JHEK-SOX2 (C) and AHEK and AHEK-SOX2 (D) using the Wald test (two-tailed).

(E) Gene set enrichment analysis of Embryonic epidermis and juvenile transcriptional signatures [24] in AHEK and AHEK-SOX2. Pathways are significantly enriched (padj<0.01 based on an adaptive multi-level split Monte-Carlo).

(F) Normalised RNA expression levels (counts) of SOX2 from a cSCC human dataset [17] including normal skin (NS, n=17), and different levels of disease progression including actinic keratosis (AK, n=14) and cutaneous squamous

cell carcinoma (SCC, n=66). Dotted lines mark the mean *SOX2* normalised expression in normal skin and two

standard deviations from the mean. Pie charts showing the percentage of samples with high and low *SOX2*

normalised expression for cSCC and AK groups. Two standard deviations from normal skin mean were used as a

threshold.

(G) Heatmap showing individual *SOX2* normalised expression levels in AK and cSCC groups.

(H) Representative IHC of SOX2 conducted in human tissue microarray containing 250 cSCC histocores described

in[36]. 196 (78.4%) of the samples were negative, and 54 (21.6%) were positive for SOX2 protein expression. The

scale bar is 50 μm.

(I) Graphic abstract of the model arising from this study. cSCC can be initiated from K14/K5+ stem-like cell or

differentiation committed IVL+, the latter will require SOX2 activation and will take longer to escape constraints

and trigger tumorigenesis. SOX2 reprogramming will activate embryonic and stem-like features critical for

tumorigenesis. During this process, actively dividing IVL+ expand through the entire epithelium with little

histological change, preparing for tumour initiation and progression and leading to an alternative route for

tumorigenesis. We propose a common oncogenic programme, seen in human cSCC[25] and inducible by

inflammation, injury or paradoxical activation of MAPK, that allows the transformation of "normal" epithelial-

containing oncogenes into a tumour without additional oncogenic events.