**Comparative proteomics uncovers distinct biomarkers, protein networks and defense**

**responses in tomato during beneficial and pathogenic microbial interactions**

Dhananjaya Pratap Singh[1,*], Sudarshan Maurya[1], Suresh Reddy Yerasu[1], Ratna Prabha[2], Renu

S[3], Birinchi K. Sarma[4], Lovkush Satnami[1], Nagendra Rai[1]

[1]ICAR-Indian Institute of Vegetable Research, Shahanshahpur, Varanasi 201305, India

[2]ICAR-Indian Agricultural Research Institute, Agricultural Knowledge Management Unit,

110012 New Delhi, India

[3]Indian Council of Agricultural Research, Krishi Bhawan, 110001 New Delhi, India

[4]Department of Mycology and Plant Pathology, Institute of Agricultural Sciences, Banaras

Hindu University, Varanasi 221005 India

*Corresponding author:

Dhananjaya Pratap Singh, Principal Scientist (Biotechnology)

email: Dhananjaya.Singh@icar.gov.in

**Abstract**

The molecular mechanisms underlying plant responses to beneficial and pathogenic microbial

interactions has been uncovered in tomato at the proteomic level. Our study employed LC-MS-

based proteomics to investigate differential protein regulation in tomato plants during

interactions with beneficial bacteria (*Bacillus* subtilis BV7) versus pathogenic fungus (*Alternaria*

*solani*). Comparative analysis revealed distinct protein signatures characterizing each interaction:

232 unique proteins in BV7-treated plants versus 96 in pathogen-infected plants, with 54 proteins

shared between treatments. BV7 inoculation enhanced proteins involved in photosynthesis and

primary metabolism, with PSI-K emerging as the top biomarker (score 2.53), while pathogen

infection triggered focused defense responses with Cytochrome b559 (score 2.38) as the key

biomarker. Metabolic pathway analysis demonstrated that BV7 uniquely enhanced vitamin

metabolism (thiamine, riboflavin, folate) and energy production pathways, while pathogen

infection activated defense-related phenylpropanoid biosynthesis. Analysis of defense enzymes

showed pathogen infection induced highest activities of PAL ($39.4\pm0.46$ U $h^{-1}$ $g^{-1}$ fw), SOD

(29.9±0.94 U mg$^{-1}$ protein), POD (3.18±0.07 μg g$^{-1}$ fw min$^{-1}$), APx (15.58±0.23 U mg$^{-1}$ protein),

and GPx (71.51±0.7 U mg$^{-1}$ protein), while BV7 maintained moderate enzyme levels, suggesting

balanced growth-defense responses. The shared proteins between treatments indicate a common

molecular framework potentially contributing to induced systemic resistance. These findings

provide novel insights into plant-microbe interactions at the molecular level, identifying specific

protein biomarkers and metabolic pathways that could be targeted for enhancing crop

productivity and disease resistance. The results have significant implications for developing

biological control strategies and improving sustainable agricultural practices.

**Keywords**

39   Proteomics, LC-MS, Plant-microbe interactions, Tomato, *Bacillus subtilis, Alternaria solani,*

40   Biomarkers

## 1. Introduction

The intricate interactions between plants and microorganisms play a pivotal role in shaping plant

health, productivity, and overall ecosystem dynamics. Plant-microbe interactions can be broadly

categorized as beneficial, neutral, or pathogenic, each eliciting distinct molecular responses

within the host plant (Jones et al, 2019). Understanding the molecular mechanisms underlying

diverse microbial interactions with plants is crucial for developing sustainable agricultural

practices and enhancing crop resilience in the face of biotic and abiotic stresses (Agrahari et al,

2020). Tomato (*Solanum lycopersicum*), a globally important vegetable crop, serves as an

excellent model system for studying plant-microbe interactions due to its economic significance,

genetic tractability, and susceptibility to various pathogens (Krishna et al, 2022, Liu et al, 2022).

The plant's responses to beneficial and pathogenic microbes provide valuable insights into the

complex molecular networks that govern plant immunity, growth promotion, and stress

tolerance.

Beneficial gram-positive bacterium *Bacillus subtilis* has widely recognized as a plant growth-

promoting rhizobacterium (PGPR) that colonizes plant rhizosphere and confers multifaceted

benefits to the host plant, including enhanced nutrient uptake, phytohormone production, and
induced systemic resistance (ISR) against pathogens (Vejan et al, 2016). The mechanisms by
which *B. subtilis* promotes plant growth and health involve complex signaling cascades and
metabolic reprogramming within the host plant (Hashem et al, 2019). On the other hand,
*Alternaria solani* causes a devastating early blight disease in tomato and significantly impacts
production of this crop worldwide. This necrotrophic fungus induces cell death and tissue
necrosis, leading to substantial yield losses. The interaction between tomato and *A. solani*
triggers a distinct set of biochemical and molecular mechanisms leading to defense responses,
including the activation of pathogenesis-related (PR) proteins, phytoalexin production, and
hormonal signaling pathways (Pandey et al, 2023). The contrasting nature of beneficial and
pathogenic interactions provides a unique opportunity to elucidate the molecular signatures that
differentiate plant responses to diverse microbial stimuli. While previous studies have focused on
individual aspects of these interactions, a comprehensive understanding of the global proteomic
changes and associated pathway shifts remains elusive.

Recent advances in high-throughput proteomics technologies have revolutionized our ability to
study complex biological systems at unprecedented depth and resolution (Yan et al, 2022). These
approaches enable the identification and quantification of thousands of proteins simultaneously,
offering a holistic view of cellular processes and regulatory networks (Hu et al, 2015). To
provide a comprehensive understanding of the differential responses of tomato to beneficial and
pathogenic interactions, we have applied state-of-the-art proteomics techniques to study *B.*
*subtilis* and *A. solani* interaction at proteimic level and unravel the intricate molecular landscape
that defines contrasting relationships. The objective of this study was to decipher the differential
proteomic signatures and pathway shifts in tomato plants in response to colonization by the
beneficial *B. subtilis* and infection by the pathogenic *A. solani*. By comparing distinct proteomic
profiles, we have identified key biomarker proteins, metabolic pathways, and regulatory
networks that are specifically modulated in each interaction. This comparative approach will
provide valuable insights into the molecular mechanisms underlying plant growth promotion,
induced resistance, and pathogen defense. Our findings not only advance fundamental
knowledge of plant-microbe interactions at the proteomic level but also pave the way for
innovative strategies to enhance crop resilience and productivity in the face of diverse microbial
challenges. The insights gained from the findings may help in targeting important pathways for

manipulation through microbial inoculations for reducing chemical farm inputs and promoting
environment-friendly crop protection methods.

**2. Material and Methods**

**2.1. Beneficial microbe, pathogen and plant treatments**

Seeds of the tomato plants (*S. lycopersicum* var. Kashi Aman) were sown in pots (20x20x14cm)
filled with sterilized field soil, cocopeat, perlite and vermiculite in a ratio of 3:1:1:1; w/w. Potted
seeds were kept in greenhouse conditions (Singh et al, 2023, 2024). After emergence, seedlings
were allowed to grow and develop at 25°C/20°C (day/night; D/N) thermal condition and 14h/10h
(D/N) photoperiod at a relative humidity of 70%. An artificial light source illuminating
600 $\mu$mol m$^{-2}$ s$^{-1}$ of light was used to keep plants illuminated. Uniformly growing 3 weeks old
seedlings having similar height and growth were used for the bacterial inoculation treatment. The
beneficial bacterium, *Bacillus subtilis* (BV7) was grown in LB broth media (tryptone 10g, yeast
extract 5g, NaCl 10g and distilled water make up to 1L) at 37°C. The microbial inoculant using
*B. subtilis* BV7 was prepared using the spores of the bacterium (2.7x10$^{8}$ mL$^{-1}$ colony-forming
units (CFU) in which the pre-germinated tomato seedlings were dipped for 15 min and then
transplanted to the experimental pots. Separate sets of seedlings were transplanted in separate
pots having control and bacteria-inoculated plants were allowed to grow under similar growth
conditions till 60 days. One set of control plants was inoculated with *A. solani* culture (2.1×10$^{4}$
conidia mL$^{-1}$) on 61$^{th}$ day at their lower leaves. Altogether, three sets of plants were available:
control (non-treated), *B. subtilis* BV7 inoculated and *A. solani* infected plants. Plants were
maintained at 86-90% relative humidity to facilitate disease progression and sampling was done
after 5 days of pathogen inoculation. The disease incidence (%) was noted at the time of
sampling. Each treatment was replicated three times.

**2.2 Enzyme extraction and antioxidant activity**

Enzyme extracts were prepared from the same leaf samples as were used for the extraction of
global plant proteome. Freshly harvested leaf tissues (0.5 g) from control, BV7-inoculated and
pathogen-infected plants were homogenized separately in extraction buffer containing 10 ml
sodium phosphate buffer (0.1M), 0.2 mL mercaptoethanol and 1.0 g of polyvinyl
pyrrolidone (PVPP). The extracted suspension was kept for 1 h after centrifugation (12000 rpm,

30 min, 4°C) and precipitated with 5 volumes of ammonium acetate (0.1M) in methanol. The

enzyme (protein) mixture was stored 1h and then centrifuged (10000 rpm, 10min) to obtain

pellets that were finally re-suspended in extraction buffer (3ml). Protein content was determined

(Lowry et al., 1951). Assays for the analysis of antioxidant activity related enzymes were

performed for phenylalanine ammonia-lyase (PAL), peroxidase (POD), superoxide dismutase

(SOD), ascorbate peroxidase (APx) and guaiacol peroxidase (GPx) as per the methods described

in Singh et al (2020) and Patani et al, (2023) with minor and conditionally suitable

modifications. All the experiments were performed in 3 replications and the data were subjected

to the test of statistical significance using SPSS 16.0.

**2.3. Proteome extraction from plant leaves**

Global proteome extraction was performed using the phenol extraction method (Faurobert et al,

2007). The procedure for sample preparation included grinding of 1g of fresh leaf tissues kept in

frozen liquid nitrogen in a pre-cooled pestle-mortar. The sample was the suspended in ice-cooled

methanol for 1h to remove pigments and phenolic contaminants that interfere with the total

proteome content. After centrifugation in cold condition, the residue was re-suspended in 3ml

extraction buffer (500mM Tris HCl, 50 mM EDTA, 700 mM sucrose, 100 mM KCl, pH 8.0).

Mercaptoethanol (2%) and phenylmethylsulphonyl fluoride (PMSF) (1mM) were added before

re-suspension. The mixture was the vortexed and incubated on ice for 10 minutes with shaking.

An equal volume of Tris-buffered phenol (phenol + tris-HCl saturated buffer, pH 6.6/7.9) was

added. After 10 minutes of incubation, the solution was centrifuged at 12000$g$. The phenolic

phase was removed thrice. Extracted protein was precipitated using 0.1M ammonium acetate in

cold methanol. After centrifugation, the resulting pellet was stored at -20°C. Protein

concentration was determined using Bradford protein assay (Bradford 1976).  The precipitated

protein was dissolved in G-buffer (guanidine hydrochloride) and the protein solution was filtered

through a 0.22µM syringe filter prior to proteomic data generation.

**2.4 Trypsin digestion and tandem mass tag (TMT) labeling**

Protein samples (100µg) were combined with 200µL of 50 mM $NH_4HCO_3$ and digested with

trypsin following López-Ferrer et al, (2008). Samples were treated with 5 mM dithiothreitol

(DTT) and incubated for 1h at 55°C, followed by the addition of 11 mM iodoacetamide and a

15-minute incubation in dark at room temperature. Urea concentration was reduced to <2M.

Trypsin was added at a 1:50 (trypsin:protein) ratio, and samples were digested overnight at 37°C.

A second enzymatic hydrolysis was performed with trypsin at 1:100 (trypsin:protein) for 4h. The

digested peptides were desalted, freeze-dried, and then dissolved in 0.5M triethylammonium

bicarbonate. Finally, the peptides were labeled using a tandem mass tag (TMG) kit according to

the manufacturer's instructions.

**2.5 LC-MS analysis**

Peptide separation was performed using reverse phase nano-liquid chromatography on a Thermo

Scientific Easy nLC 1200 instrument (Thermo scientific). The column used was a PepMap

RSLC C18 2µm, 75µm x 50cm, maintained at 60°C. A 2µg peptide sample was loaded onto the

column. For separation, the solvents were, A: 98% water, 2% acetonitrile, 0.1% formic acid, and

B: 20% water, 80% acetonitrile, 0.1% formic acid with a flow rate constant at 300nL min$^{-1}$. The

gradient profile was 1. 0-2 min: 5% to 10% solvent B, 2. 2-104 min: 10% to 45% solvent B, 3.

104-105 min: 45% to 90% solvent B, 4. 105-115 min: hold at 90% solvent B, 5. 115-116 min:

decrease to 5% solvent B and 6. 116-118 min: hold at 5% solvent B. All the solvents used were

LC-MS grade (Fischer Scientific). Internal calibration utilized a lock mass of 445.12003 Da.

MS data was generated using a Q Exactive Orbitrap mass spectrometer (Thermo Scientific)

coupled to an Easy-nLC 1200 liquid chromatograph. The run time was 0 to 120 min in positive

polarity with a default charge state of 2.0. For MS1, settings included 1 microscan, 70000

resolution, 3e6 AGC, 60ms maximum IT time, and a 350 to 2000 m/z scan range. MS2 settings

were 17500 resolution, 200 to 2000 m/z full scan range, 1e5 AGC, 12ms ion transfer, 100 m/z

fixed first mass, and 27 normalized collision energy. Data dependent analysis (DDA) settings

included 1.00e2 AGC target max, 50.0 S dynamic exclusion time, and unassigned charge

exclusion.

**2.6 Database search**

The annotation of proteins in the samples was performed using Proteome Discoverer 2.5 in

conjunction with the UniProt protein database of *Solanum lycopersicum* (UniProt). Validation

parameters based on q-values included peptide length 6 minimum, 144 amino acid residues

maximum, precursor mass tolerance 10 ppm, secondary fragment ion mass error tolerance 0.02

Da and restriction method trypsin/p with maximum allowed missed cleavage up to 2. The false

discovery rate (FDR) was restricted to 0.01 for both protein and peptide-spectrum match (PSM)

identification.

**2.7 Functional annotation and enrichment analysis**

GO based annotation was done using UniProt-GOA database (http://www.ebi.ac.uk/GOA).

Protein IDs were converted to UniProt IDs and then GO IDs were matched to retrieve

corresponding information from the UniProt-GOA database. InterProScan was used to predict

the GO function of the protein if there was no protein information query on the UniProt-GOA

database. Venny 2.1 (Venny 2.1.0 (csic.es) was used to generate a Venn diagram to classify

unique proteins in the samples. MetaboAnalyst 6.0 and SIMCA 18 (Sartorius) was implied to

perform univariate (volcano plot) analysis and SS Plot was used for data visualization. Data was

normalized taking median and log transformation values and Pareto scaling prior to multivariate

analysis and preparation of correlation heatmap. Panther classification system (pantherdb.org)

database was used to classify GO-Slim biological and molecular functions and categorize protein

class. InterPro (a resource for functional analysis of protein sequence family classification,

prediction of domains and special sites) database was used to analyze the functional domains of

proteins.

The Kyoto Encyclopedia of Genes and Genomes (KEGG) (KEGG: Kyoto Encyclopedia of

Genes and Genomes) pathway database of *S. lycopersicum* (*sly*) was used to annotate protein

pathways. Pathway mapping of proteins was facilitated by an interactive pathway explorer tool

iPATH3.0 (iPath 3: interactive Pathways Explorer (embl.de). The functional protein association

network web-tool STRING (STRING: functional protein association networks (string-db.org)

was used to perform protein-protein interaction (PPI) networks of the up- and down-regulated

proteins identified in the samples. ROC curve analysis module of MetaboAnalyst 6.0 was used

for the identification of protein biomarkers. A *t*-test was employed to test the enrichment of the

differentially expressed proteins against all identified proteins. The GO/KEGG/protein domains

with a corrected *p*-value <0.05 were considered significant.

**3. Results**

The study evaluated how the protein composition changed in the individual set of plant leaves in

control (non-treated), beneficial bacteria inoculation and pathogen (*A. solani*) infection condition

and analyzed comparative proteome profile to evaluate differential impact of beneficial and

pathogenic interactions. We evaluated the activity of defense related antioxidative enzymes PAL,

POD, SOD, APx and GPx in the plant leaves for the assessment of biochemical impact of

beneficial microbe inoculation and pathogen infection at enzymatic level (Fig 1). Comprehensive

biochemical analyses of plant leaf under different treatment conditions, including control (C),

beneficial *Bacillus* BV7 (B), and pathogenic *A. solani* (P) treatments were presented in Fig 1. At

the time of the sampling for proteome extraction, infected plants showed a disease incidence of

72.4%±1.2 showing high degree of pathogen infection (Fig 1a). Determination of total protein

content revealed that *Bacillus* BV7 treatment induced highest protein levels (3.13±0.21mg $g^{-1}$

fw) compared to both the control and pathogen-infected samples, while control and pathogen

treatments showed similar, relatively lower protein contents (Fig 1b). The analysis of the

activities of several defense enzymes across these treatments showed varying response. PAL

activity demonstrated a progressive increase from control to BV7 to pathogen treatments, with

the pathogen treatment showing the highest activity (39.4±0.46 U $h^{-1} g^{-1}$ fw) (Fig 1c). Similarly,

SOD activity exhibited a stepwise increase across treatments, with control showing the lowest

activity, followed by intermediate activity in BV7-treated samples, and the highest activity in

pathogen-infected tissues (29.9±0.94 U $mg^{-1}$ protein) (Fig 1d). Regular POD activity followed a

similar trend, with control samples showing minimal activity, BV7 treatment inducing moderate

activity, and pathogen infection resulting in the highest activity levels (3.18±0.07 μg $g^{-1}$ fw $min^{-1}$

$^{1}$) (Fig 1e).

Analysis of peroxidase enzymes including APx and GPx revealed consistent patterns. APx

activity showed the lowest levels in control samples, with a significant increase in BV7-treated

samples and the highest activity in pathogen-infected tissues (15.58±0.23 U $mg^{-1}$ protein) (Fig

1f). GPx activity also followed mirrored this pattern, demonstrating progressive increases from

control to BV7 to pathogen treatments (71.51±0.7 U $mg^{-1}$ protein) (Fig 1g). The activities of

defense enzymes showed remarkable consistency in their response patterns, with control samples

consistently showing the lowest activity, BV7 treatment inducing intermediate levels, and

pathogen infection triggering the highest enzymatic responses. The only exception to this pattern

was observed in total protein content, where BV7 treatment induced the highest content in plant

leaves. The results indicated that both beneficial (BV7) and pathogenic treatments trigger plant

defensive responses, but at different intensities, with pathogenic infection generally eliciting the

strongest enzymatic responses while the beneficial bacteria induces moderate levels of these

defense-related enzymes. These findings therefore, suggested the need to explore the molecular

basis of the interaction conditions in tomato plants at the proteomic level.

Figure 1.

Figure 1: Biochemical and disease responses in plant leaves under different treatment conditions

including a. inoculation with the beneficial *Bacillus* BV7 (B), infection with the pathogen *A.*

*solani* (P) and Control (C). The disease incidence was reported to be 72.4%±1.2. Quantitative

analyses of b. total protein content, and various biochemical markers including enzymes c.

phenylalanine ammonia-lyase (PAL) activity, d. superoxide dismutase (SOD) activity, e.

peroxidase activity, f. ascorbate peroxidase activity, and g. guaiacol peroxidase activity. Error

bars represent standard error (SE) of the mean.

The LC-MS-based proteomic analysis revealed distinct protein expression patterns across

control, beneficial bacteria (BV7)-inoculated, and pathogen-infected tomato leaves, as illustrated

in the Venn diagram (Fig 2a). The analysis identified unique protein signatures for each
treatment condition, with 316 proteins exclusively present in control plants, 232 proteins unique
to BV7-inoculated plants, and 96 proteins specifically associated with pathogen infection.
Additionally, the diagram showed overlapping proteins, with 202 proteins shared between
control and BV7 inoculation, 54 proteins common between BV7 and pathogen infection, and 26
proteins shared across all the three conditions. In pathogen-infected leaves (Fig 2b), the 96
unique proteins were primarily associated with metabolic processes, cellular metabolism, and
organic substance metabolic processes, as indicated by the GO term analysis. This suggested
significant metabolic reprogramming in response to pathogen infection, likely representing the
plant's defense mechanisms against the pathogen. The control condition (Fig 2c) showed the
highest number of unique proteins (316), with significant enrichment in cellular processes,
metabolic pathways, and macromolecule metabolic processes. The presence of these proteins
indicated normal physiological processes occurring in healthy tomato plants, establishing a
baseline for comparison with treated conditions. In BV7-inoculated plants (Fig 2d), 232 unique
proteins showed enrichment in metabolic processes, cellular processes, and various binding
activities. Notably, the GO analysis revealed significant representation of proteins involved in
cellular component organization, biosynthetic processes, and intracellular membrane-bounded
organelle functions. This protein profile suggested that BV7 inoculation triggers specific cellular
responses that may contribute to enhanced plant growth and defense preparedness.

A detailed analysis on how the unique proteins identified in the control, BV4 inoculated and
pathogen infected plants enriched metabolic pathways, all the GO terms are presented for
'biological process', 'molecular function', 'cellular component', 'subcellular localization',
'protein domain features' and 'annotated key words' in Supplementary Tables S1, S2 and S3.
STRING assigned GO terms for 'biological process' to the uniquely identified proteins in all the
treatment conditions revealed distinct biological responses reflecting the impact of each
conditions on plant health and resilience (Table 1). Majority of unique proteins in the control
plants were primarily associated with cellular processes (244), cell metabolism (145), metabolic
process (164), primary metabolic process (135), nitrogen compound metabolism (122),
macromolecule biosynthesis (109), and organic substance metabolism (143) (Table S1). It
suggested a normal plant physiological process indicating majorly protein enrichment in
foundational cellular mechanism, gene expression, nitrogen metabolism, and macromolecule

biosynthesis. The compartmentalization of proteins is a crucial for protein function as the

specific location of proteins within cells. In control plants, unique proteins were predominantly

localized to intracellular compartments, including cytoplasm, ribosomes, chloroplasts, and the

mitochondria (Table S1). In BV7-inoculated plants, it was seen that besides maintaining

metabolic (155), cellular (176) and cellular metabolic process (121 proteins), the biological

processes for the unique proteins shifted towards enhanced small molecule biosynthesis, nitrogen

metabolism, amino acid metabolism, carboxylic acid and vitamin biosynthetic processes (Table

S2). Increased enrichment of proteins in vitamin biosynthesis (e.g. riboflavin) and protein

refolding indicated beneficial impact of BV7 inoculation, which further caused unique protein

enrichment in subcellular compartments such as the cytoplasm, chloroplasts, peroxisomes, and

nucleosomes, supposedly enhancing cellular processes like metabolic activity, antioxidant

defense (*via* peroxisomes), and photosynthesis (chloroplast proteins) (Table S2). In contrast, in

plants infected with *A. solani,* there were unprecedented drop in the number of GO terms to

which the 96 uniquely present proteins enriched (Table S3) and the enrichment showed a distinct

profile dominated by catabolic and small molecule metabolic processes, indicative of stress-

related responses. In pathogen-infected plants, protein localization was shown in cytoplasmic

and cytosolic compartments. These proteins are involved in stress responses and metabolic

disruptions, indicating a shift from constructive cellular activities to defensive mechanisms

(Table S3).

Figure 2

Figure 2. Plant leaves of tomato showing a. classification of unique proteins using venn diagram in control, BV7 beneficial bacteria inoculated and pathogen infected plant leaves as identified by LC-MS-based proteomics analysis, b. Gene Ontology (GO) terms functional classification (FC) for 96 unique proteins enriching different metabolic pathways in pathogen infected tomato plant leaves, c. FC for 316 proteins unique in control leaves, d. FC for 232 proteins unique in BV7 inoculated plant leaves. GO terms are grouped in different color bars including 'red' for 'biological process', 'blue' for 'molecular function' and 'green' for 'cellular component'

Table 1.

Supplementary Table S1, S2, S3

The proteomes of two plant groups involving control *versus* BV7-inoculated (C*vs*BV7) and control *versus* pathogen infected (C*vs*Pathogen) plants were mutually compared to understand changes at the level of protein profiles. Univariate analysis using volcano plot indicated regulation (up- or -down) of proteins in both the plant groups in terms of fold change (FC>1.5) and *p* values (<0.05) (Fig 3a,b). C*vs*BV7 plant group showed 101 differentially up-regulated and

106 down-regulated proteins (Fig 3a). In C*vs*Pathogen plant group, there were 75 up- and 108

down-regulated proteins (Fig. 3b). The heatmap generated a correlation between overall proteins

in the C*vs*BV7 (Fig. 3c) and in C*vs*Pathogen group (Fig 3d). Heatmap for CvsBV7 showed a

clear segregation into four distinct clusters (represented by the red and blue blocks). The sharp

contrast between these blocks indicated that BV7 treatment leads to consistent and substantial

changes in the overall protein composition compared to control conditions. The presence of both

positive (red) and negative (blue) correlations suggested a complex protein regulation networks

are activated during beneficial bacterial colonization. The heatmap for CvsPathogen displayed a

more complex correlation pattern with multiple smaller clusters, indicating that pathogen

infection triggers a more diverse and perhaps more specialized protein response. The heatmap

showed more gradual transitions between correlation clusters, suggesting interconnected protein

regulation networks responding to pathogen stress. The pattern further differentiated the samples

based on protein composition and indicated major changes in the overall protein profile in BV7

inoculated and pathogen infected plants, indicating distinct changes at the protein regulation

level in tomato in response to both the beneficial and pathogen interaction.

Figure 3

Figure 3. Identification of proteins that differ significantly in terms of their fold change (FC) values at p<0.05 to describe protein regulation (up- and down) in the samples, Volcano plot for a. CvsBV7 and b. CvsPathogen plant group; and correlation heatmap showing sample differentiation based on overall protein composition in c. CvsBV7 and d. CvsPathogen plant group

A distinct but complex enrichment of unique proteins identified in C*vs*BV7 (red lines) and C*vs*Pathogen (blue lines) plant groups in various metabolic pathways was observed in tomato leaves in the pathway map generated by iPath3.0 highlighting unique protein engagement patterns (Fig. 4). Contrast biosynthetic and metabolic pathways were mapped in iPATH by unique proteins found in the plant group Cvs BV7 and CvsPathogen. In CvsBV7-inoculated plant group, there is notable activation of pathways related to primary metabolism, including enhanced photosynthetic processes and energy metabolism. BV7-treated plants show enhanced protein engagement in pathways related to beneficial metabolite production, suggesting

improved plant fitness. The positive influence of BV7 interaction on the plant's energy
production and carbon fixation capabilities is supposed to potentially contribute to improved
plant growth and development. The unique proteins in Cvspathogen-infected plants showed
differential engagement in stress response pathways and secondary metabolism. Inoculation of
beneficial bacteria BV7 enriched vitamin metabolism including that of thiamine, riboflavin, and
folate metabolism more prevalently as compared to the unique proteins in CvsPathogen plant
group. Likewise, phenylpropanoid biosynthesis, tyrosine and phenylalanine metabolism was
highly enriched by the proteins unique in pathogen infected plants leaves showing significant
activation of defense-related pathways, including those involved in oxidative stress response.
This has indicated plant's shift towards defense mechanisms rather than growth-promoting
processes under pathogenic interaction condition.

Particularly noteworthy is the differential regulation of amino acid metabolism and secondary
metabolite biosynthesis between the two conditions. The interconnectedness of these pathways
demonstrated complex metabolic reprogramming during plant-microbe interactions. The
beneficial bacterium BV7 appeared to promote pathways that enhance plant growth and
development, while pathogen infection triggered extensive metabolic changes related to stress
response and defense. This differential pathway engagement has provided insights into how
plants modulate their metabolism in response to beneficial versus pathogenic interactions and
how the treatments underline plant's ability to distinguish between beneficial and harmful
microorganisms, leading to appropriate metabolic responses that either promote growth or
activate defense mechanisms.

**Figure 4.** Comparative pathway analysis of differentially expressed proteins in plants under different treatments. The metabolic pathway map illustrates unique proteins identified in BV7-inoculated plants (red lines, CvsBV7) and pathogen-infected plants (blue lines, CvsPathogen) relative to control plants. The network demonstrates the complex interconnections between various metabolic pathways (shown in blue text), including but not limited to photosynthesis, fatty acid metabolism, amino acid metabolism, and secondary metabolite biosynthesis. The intricate pathway mapping reveals distinct metabolic responses triggered by beneficial (BV7) and pathogenic interactions in plants, highlighting the differential regulation of key biological processes under these conditions.

Figure 5

Figure 5. Identification of biomarker proteins (using ROC curve analysis classification method PLS-DA and feature ranking method PLS-DA built-in in MetaboAnalyst 6.0) and their interactions using String database in CvsBV7 and CvsPathogen plant groups. Top 25 biomarker proteins ranked by average importance along with their functional groups were identified in a. BV7-inoculated and b. pathogen infected plants compared to control. Red and green bars indicate the average importance scores of proteins in CvsBV7 and CvsPathogen plant groups, respectively; c. Protein-protein interaction (PPI) network of key biomarker proteins in CvsBV7 group, with nodes colored according to functional clusters and edge thickness representing interaction strength, d. PPI in CvsPathogen plant group showing interactions and functional clustering representing differential molecular responses to beneficial and pathogenic interactions

Identification of reliable biomarkers has remained central among the major aims of any proteomic analysis (Alharbi et al, 2020). Results based on ROC analysis identified potential protein biomarkers distinguishing between C*vs*BV7 inoculation (Fig 5a) and C*vs*Pathogen (Fig 5b). The comparative analysis of biomarker proteins revealed distinct molecular signatures in plants responding to beneficial bacteria (BV7) versus pathogen infection. In BV7-inoculated

plants (Fig. 5a), PSI-K (Photosystem I subunit K) emerged as the most significant biomarker
with the highest average importance score (2.53), followed by ribosomal protein L35, Glycine-
rich protein, Alcohol acyl transferase, alanine transaminase, Glycine-rich protein, Fruit-ripening
protein, Glucan endo-1,3-beta-D-glucosidase. The predominance of photosynthesis-related and
metabolic proteins among the top biomarkers suggested that BV7 inoculation enhances primary
metabolism and energy production processes. Notably, several stress-response proteins,
including UTP-glucose-1-phosphate uridylyltransferase and NADP-binding domain-containing
proteins, also showed high importance scores (>1.8), indicating an induced balanced regulation
of defense responses along with the growth and development. In contrast, pathogen-infected
plants (Fig. 5b) displayed a distinct set of biomarker proteins primarily associated with defense
responses. The identified biomarker proteins Cytochrome b559 subunit alpha ranked highest
(2.38), followed by the proteins ethylene-responsive proteinase inhibitor 1, thiamine thiazole
synthase, glutamate synthase (ferredoxin), carboxypeptidase A inhibitor-like domain-containing
protein, and 14-3-3 protein 7 in CvsPathogen plant group (Fig. 5b) indicated the protein profile
shift towards signal, pathogenesis-related stress response, catabolic processes and defense
mechanisms during pathogen infection. The functional distribution of proteins in different groups
viz. protein synthesis and ribosomal proteins,metabolic enzymes, proteins related to plant
growth, photosynthesis, chloroplast, cell wall modification, energy metabolism, protein
modification and regulation, stress response, defense and signal transduction highlighted the
sophisticated molecular machinery plants employ to distinguish between beneficial and
pathogenic interactions, appropriately allocating resources either toward growth promotion or
defense responses. The differential regulation of these protein groups demonstrated the plant's
ability to mount specific and appropriate responses to different microbial encounters.

The protein-protein interaction networks provided further insights into the functional
organization of these responses. In BV7-treated plants (Fig. 5c), the network showed
concentrated clustering around metabolic processes, with strong interactions between key
proteins (indicated by edge thickness). The functional clusters, represented by differently colored
nodes, suggest coordinated regulation of growth-promoting pathways. The pathogen-response
network (Fig. 5d) exhibited a more dispersed interaction pattern with multiple functional
clusters, reflecting the complex nature of defense responses. This network organization
demonstrates the activation of diverse defense mechanisms and stress response pathways during

pathogen infection. These findings highlighted how plants engage different protein networks and
metabolic pathways when interacting with beneficial versus pathogenic microorganisms,
providing molecular insights into the plant's ability to respond to different microbial interactions.
Overall, our results have pointed out the complex and specific protein-level responses of plants
to varied microbial interactions, providing valuable insights into the molecular mechanisms
underlying plant-microbe interactions.

## 4. Discussion

The study evaluated proteome level biochemical dynamic changes in the leaves of tomato (var.
Kashi Aman) plants inoculated with *Bacillus* BV7 (beneficial) and infected with *A. solani*
(pathogenic) interaction using comparative proteomics. The findings revealed critical insights
into how beneficial and pathogenic interactions shape the plant's proteome, influencing its
metabolic, biosynthetic, and defense-related processes. Both the beneficial bacterial inoculation
and pathogen infection in tomato affected defense related antioxidative enzyme activity and
presented proteome level distinct changes that reflected molecular mechanisms underlying plant-
microbe interactions. An exclusive data analysis pipeline, including proteome extraction, data
generation, enzyme assay, protein annotation, pathway enrichment analyses and statistical
models was employed to identify significantly regulated proteins and biomarkers. Mapped
altered pathways and associated biological processes were implied to generate a comprehensive
view of the cellular reprogramming events in the plants (Khan et al, 2024).

The antioxidative enzymes such as POD, SOD, APx, and GPx play pivotal roles in plant defense
by mitigating oxidative stress caused by biotic interactions (Rajput et al, 2021). Significant
increase in enzyme activities observed in BV7-inoculated plants highlighted their role in priming
plant defense mechanisms. Significantly higher protein content in BV7-treated plants aligns with
findings by Hashem et al (2019), who demonstrated that beneficial *Bacillus* strains enhance
protein synthesis machinery in host plants. The lower protein content in pathogen-infected
tissues may indicate protein degradation or metabolic disruption during pathogenic invasion,
consistent with observations by Schmey et al. (2024). PAL activity showed progressive elevation
from control to BV7 to pathogen treatments, with pathogen infection inducing the highest
activity. This pattern is of particular significance as the enzyme PAL represents the first
committed step in the phenylpropanoid pathway, crucial for defense compound synthesis (Vogt

2010). The moderate induction of PAL enzyme activity by the *Bacillus* BV7 inoculation

suggested a balanced immune response that may have contributed to improved plant health

(Bhattacharyya et al 2020).

The activity of SOD was highest in pathogen-infected tissues indicating intense oxidative stress

management level, as previously noted by Torres et al (2006) during plant-pathogen interactions

(Kunos et al 2022). The moderate SOD induction in *Bacillus* BV7-treated plants suggested a

primed state of oxidative stress response without the metabolic burden of full activation,

supporting the concept of beneficial microbe-induced systemic resistance described by Pieterse

et al. (2014). The elevated GPx activity in pathogen-infected plants reflected deployment of

ROS-scavenging mechanisms, crucial for managing oxidative stress during pathogen attack

(Wang et al 2019). The intermediate levels of peroxidase activities in BV7-treated plants have

led to presume a calibrated defense response that may contribute to enhanced stress tolerance

without compromising growth in beneficial microbe-plant interactions (Meena et al. 2017,

Tharanath et al, 2024).

The comparative proteomic analysis has revealed distinctive proteomic responses in tomato

under different microbial interactions, highlighting the proteome-level mechanisms plants

employ to differentiate between beneficial and pathogenic interactions. In control plants,

presence of 316 unique proteins enriched in fundamental cellular and metabolic processes

represented a robust baseline of physiological activities. This protein profile indicated an active

primary metabolism regulating photosynthesis and protein synthesis and the enzymes essential

for normal plant growth and development (Cheng et al, 2010, Tcherkez et al, 2020, Ji et al,

2022).

The beneficial bacteria (BV7) inoculation induced a distinct protein regulation pattern with 232

unique proteins, demonstrating proteins related to photosynthesis, primary metabolism, vitamin

biosynthesis (particularly riboflavin), and protein refolding mechanisms. The activation of these

growth-promoting pathways, coupled with maintained basal defense readiness, has shown that

beneficial microbial interaction optimally regulated plant development while ensuring protective

mechanisms to remain functional (Mhlongo et al 2020). In contrast, pathogen-infected plants

exhibited a more focused response with 96 unique proteins, predominantly associated with stress

response and defense-related enzymes. The increased catabolic processes suggested significant

metabolic reprogramming towards defense at the expense of growth (Bentham et al, 2020). This

protein level shift represented a resource allocation strategy where plants prioritize survival over
growth under pathogenic stress (Huot et al 2014). The protein-protein interaction networks
further demonstrated differential responses (Hollander et al 2021). BV7-induced networks
showed concentrated clustering with strong metabolic interactions, indicating systematic
enhancement of growth and development processes. Conversely, pathogen-induced networks
displayed more dispersed clustering with multiple functional groups, reflecting the activation of
diverse stress response mechanisms (Zhang et al., 2022). Interestingly, the common elements of
54 shared proteins in beneficial and pathogenic response pathways, potentially indicated basal
defense priming by beneficial bacteria (Mhlongo et al., 2018).

The results of volcano plot analysis effectively illustrated the magnitude and significance of
protein regulation in both the plant groups. In the CvsBV7 comparison, relatively balanced
distribution of up- and down-regulated proteins (fold change >1.5, p<0.05) suggested a
coordinated metabolic reprogramming in response to beneficial bacterial colonization (Li et al,
2023). The comparable numbers of proteins regulated in both directions indicated that the
inoculation of *Bacillus* BV7 induced both activation and suppression of specific cellular
processes rather than a unidirectional response (Salehin et al, 2021). In contrast, the CvsPathogen
group exhibited higher number of down-regulated proteins suggesting that pathogen infection
has caused broader suppression of normal cellular processes, possibly reflecting the pathogen's
strategy to compromise host defense mechanisms (Huot et al, 2014).

The heatmap of CvsBV7 demonstrated clear segregation into four distinct clusters indicating that
*Bacillus* BV7 treatment leads to well-defined and consistent changes in protein composition
compared to control condition. The presence of both positive and negative correlations suggested
the activation of complex protein regulation networks during beneficial bacterial colonization,
likely reflecting the simultaneous promotion of growth and basal defense mechanisms (Karasov
et al, 2017). The heatmap of CvsPathogen revealed a complex correlation with multiple smaller
clusters. This intricate organization suggested that pathogen infection triggered more diverse and
specialized protein response compared to beneficial bacterial interaction. The gradual transitions
between correlation clusters indicated interconnected protein regulation networks responding to
pathogen stress, highlighting the complexity of defense responses (Ding et al., 2024). Distinct
patterns observed in both heatmaps effectively differentiated the samples based on protein
composition, indicating major changes in the overall protein profile in both BV7-inoculated and

pathogen-infected plants. This differentiation at the protein level suggested proteome-based
distinguished machinery in plants for beneficial and pathogenic interactions, leading to
appropriate physiological responses (Hu et al., 2015).

The metabolic pathway maps by iPath 3.0 revealed that BV7 treatment activated pathways
related to enhanced photosynthesis, primary metabolism, energy production, and vitamin
synthesis, while pathogen infection triggered pathways associated with stress response, defense
mechanisms, and secondary metabolism (Kong et al., 2023). The enhanced engagement of
photosynthetic processes and energy metabolism pathways in BV7-inoculated plants suggested
that beneficial bacteria promoted plant's energy and metabolic efficiency (Stringlis et al., 2018).
The BV7 interaction showed enhanced activity in carbon fixation and energy production
pathways that potentially contribute to improved plant growth and development (Tiwari et al,
2022). Such observations correlate with the positive influence of beneficial bacteria on plant
fitness and productivity (Harman et al, 2021). The differential pathway engagement
demonstrated plant's modulation of their metabolism in response to different microbial
interactions (Nishad et al., 2020).

Particularly striking is the activation of pathways related to beneficial vitamin metabolism
pathways. Increased engagement of proteins in thiamine, riboflavin, and folate metabolism
pathways during BV7 interaction compared to pathogen-infected plants suggested positive
influence of the beneficial interaction on plant's nutritional and metabolic status (Vassilev and
Malusà, 2021). Contrary to this, distinct metabolic profile dominated by pathways involved in
stress response and secondary metabolism was observed in the CvsPathogen group. The
enhanced engagement of proteins in phenylpropanoid biosynthesis, tyrosine, and phenylalanine
metabolism pathways indicated significant activation of defense-related mechanisms. This
metabolic shift is presumed to be due to the plant's reallocation of resources from growth-
promoting processes to defense mechanisms under pathogenic stress (Schultz et al., 2013).
Differential regulation of amino acid metabolism and secondary metabolite biosynthesis between
the two conditions is particularly noteworthy. The interconnectedness of these pathways, as
visualized in the network map, demonstrates the complex metabolic reprogramming that occurs
during plant-microbe interactions. The beneficial bacterium BV7 has appeared to promote
pathways that enhance plant growth and development, while pathogen infection triggers
extensive metabolic changes related to stress response and defense (Shan et al, 2024).

The analysis of protein biomarkers revealed crucial insights into the distinctive molecular
signatures characterizing plant responses to beneficial bacterial inoculation (BV7) and pathogen
infection. In BV7-inoculated plants, PSI-K (Photosystem I subunit K) emerged as the most
significant biomarker, followed by key proteins ribosomal protein L35, glycine-rich protein,
alcohol acyl transferase, and alanine transaminase. The identified biomarker proteins were
majorly linked to photosynthetic efficiency and energy production,protein synthesis and cellular
growth,light harvesting and photosynthesis,carbohydrate metabolism, cell wall modification and
fruit development processes. The predominance of photosynthesis-related and metabolic proteins
among the biomarkers suggested activated primary metabolism and energy production processes
(Stringlis et al., 2018). Several stress-response proteins, including UTP-glucose-1-phosphate
uridylyltransferase and NADP-binding domain-containing proteins, showed high importance
scores (>1.8), indicating that beneficial bacteria help maintain a balanced regulation between
growth promotion and defense preparedness (Timofeeva et al., 2024).

In contrast, pathogen-infected plants showed a distinct set of biomarker proteins primarily
associated with plant defense and stress response. The cytochrome b559 subunit alpha is
associated with photosynthetic electron transport and ROS signaling during stress (Didaran et al,
2024), ethylene-responsive proteinase inhibitor 1+ integrates signaling between ethylene and
jasmonate pathway for plant defense (Lorenzo et al, 2003), thiamine thiazole synthase interacts
with abscisic acid (ABA) signaling (Ki et al, 2016), 14-3-3 protein 7 is associated with
immunity-linked programmed cell death pathway in tomato (Oh et al 2010), expansin-like B1
protein regulate developmental processes (Sun et al, 2021) and thioredoxin-like proteins mediate
redox signaling in plant immunity (Jedelská et al, 2020). Dominance of such proteins upon
pathogenic interaction indicates a shift towards stress response, catabolic processes, and defense
mechanisms during infection (Jones et al., 2019).

Protein-protein interaction (PPI) networks provided further insights into the functional
organization of these responses. In BV7-treated plants, the network exhibits concentrated
clustering around metabolic processes, with strong interactions between key proteins as indicated
by edge thickness. The functional clusters, represented by differently colored nodes, suggest
coordinated regulation of growth-promoting pathways. The pathogen-response network showed
a more dispersed interaction pattern with multiple functional clusters, reflecting the complex
nature of defense responses and indicating the activation of diverse defense mechanisms.

Functional distribution of proteins across different groups (protein synthesis and ribosomal

proteins, metabolic enzymes, proteins related to plant growth, etc.) highlighted the sophisticated

molecular machinery plants employ to distinguish between beneficial and pathogenic

interactions. This differential regulation demonstrated the plant's ability to mount specific and

appropriate responses to different microbial encounters.

**5. Conclusion**

This comprehensive proteomics-based investigation has provided insights into how plants

orchestrate distinct molecular responses to beneficial and pathogenic interactions. The study

demonstrated that beneficial bacteria (BV7) induces a balanced protein expression profile that

promotes growth while maintaining defense readiness, evidenced by 232 unique proteins

predominantly involved in primary metabolism and energy production. In contrast, pathogen

infection has triggered a more focused defensive response with 96 unique proteins primarily

associated with stress response and defense mechanisms. The identification of specific biomarker

proteins - PSI-K (score 2.53) for beneficial interactions and Cytochrome b559 (score 2.38) for

pathogenic responses provided valuable molecular signatures for distinguishing between

different plant-microbe interactions. The identification of 54 shared proteins between BV7 and

pathogen treatments suggested a common molecular framework that may contribute to induced

systemic resistance, highlighting the sophisticated nature of plant immune responses. Particularly

noteworthy is the differential regulation of metabolic pathways, which uniquely revealed that

beneficial bacteria (BV7) has enriched vitamin metabolism (thiamine, riboflavin, folate) and

primary metabolic processes while pathogen infection activated defense-related pathways

including phenylpropanoid biosynthesis, amino acid metabolism and stress responses. This has

clearly demonstrated distinct metabolic reprogramming strategies in plants responding to

different microbial interactions that distinguish between beneficial and harmful microorganisms.

The findings have significant implications for agricultural applications, particularly in

developing strategies for enhancing crop productivity and disease resistance. The identified

protein biomarkers and metabolic pathways could serve as potential targets for improving plant

health and developing more effective biological control methods. Future work building on these

insights could lead to innovative approaches in sustainable agriculture and plant protection

strategies.

**References**

Agrahari RK, Singh P, Koyama H, Panda SK (2020) Plant-microbe interactions for sustainable
agriculture in the post-genomic era. Curr Genomics 21:168-178.
https://doi.org/10.2174/1389202921999200505082116

Alharbi RA (2020) Proteomics approach and techniques in identification of reliable biomarkers
for diseases. Saudi J Biol Sci 27:968-974. https://doi.org/10.1016/j.sjbs.2020.01.020

Bentham AR, De la Concepcion JC, Mukhi N, Zdrzałek R, Draeger M, Gorenkin D, Hughes RK,
Banfield MJ (2020) A molecular roadmap to the plant immune system. J Biol Chem 295:14916-
14935. https://doi.org/10.1074/jbc.REV120.010852

Bhattacharyya C, Banerjee S, Acharya U, Singh R (2020) Evaluation of plant growth promotion
properties and induction of antioxidative defense mechanism by tea rhizobacteria of Darjeeling,
India. Sci Rep 10:15536. https://doi.org/10.1038/s41598-020-72439-z

Cheng Z, McConkey BJ, Glick BR (2010) Proteomic studies of plant-bacterial interactions. Soil
Biol Biochem 42:1673-1684. https://doi.org/10.1016/j.soilbio.2010.05.033

Ding Y, Shi Y, Yang S (2024) Regulatory networks underlying plant responses and adaptation to
cold stress. Annu Rev Genet 58:43-65. https://doi.org/10.1146/annurev-genet-111523-102226

Harman G, Khadka R, Doni F, Uphoff N (2021) Benefits to plant health and productivity from
enhancing plant microbial symbionts. Front Plant Sci 11:610065.
https://doi.org/10.3389/fpls.2020.610065

Hashem A, Tabassum B, Fathi Abd Allah E (2019) *Bacillus subtilis*: A plant-growth promoting
rhizobacterium that also impacts biotic stress. Saudi J Biol Sci 26:1291-1297.
https://doi.org/10.1016/j.sjbs.2019.05.004

Hollander M, Do T, Will T, Helms V (2021) Detecting rewiring events in protein-protein
interaction networks based on transcriptomic data. Front Bioinform 1:724297.
https://doi.org/10.3389/fbinf.2021.724297

Hu J, Rampitsch C, Bykova NV (2015) Advances in plant proteomics toward improvement of
crop productivity and stress resistance. Front Plant Sci 6:209.
https://doi.org/10.3389/fpls.2015.00209

Huot B, Yao J, Montgomery BL, He SY (2014) Growth-defense tradeoffs in plants: a balancing
act to optimize fitness. Mol Plant 7:1267-1287. https://doi.org/10.1093/mp/ssu049

Jedelská T, Luhová L, Petřivalský M (2020) Thioredoxins: emerging players in the regulation of
protein S-nitrosation in plants. Plants 9:1426. https://doi.org/10.3390/plants9111426

Ji L, Yang X, Qi F (2022) Distinct responses to pathogenic and symbionic microorganisms: the
role of plant immunity. Int J Mol Sci 23:10427. https://doi.org/10.3390/ijms231810427

Jones P, Garcia BJ, Furches A, Tuskan GA, Jacobson D (2019) Plant host-associated
mechanisms for microbial selection. Front Plant Sci 10:862.
https://doi.org/10.3389/fpls.2019.00862

Karasov TL, Chae E, Herman JJ, Bergelson J (2017) Mechanisms to mitigate the trade-off
between growth and defense. Plant Cell 29:666-680. https://doi.org/10.1105/tpc.16.00931

Khan Q, Huang X, He Z, Yang Y, Abbas F (2024) An insight into conflict and collaboration
between plants and microorganisms. Chem Biol Technol Agric 11:161.
https://doi.org/10.1186/s40538-024-00684-9

Li CL, Wang M, Wu XM, Chen DH, Lv HJ, Shen JL, Qiao Z, Zhang W (2016) THI1, a thiamine
thiazole synthase, interacts with Ca2+-dependent protein kinase CPK33 and modulates the S-
type anion channels and stomatal closure in *Arabidopsis*. Plant Physiol 170:1090-104.
https://doi.org/10.1104/pp.15.01649

López-Ferrer D, Heibeck TH, Petritis K, Hixson KK, Qian W, Monroe ME, Mayampurath A,
Moore RJ, Belov ME, Camp DG, Smith RD (2008) Rapid sample processing for LC-MS-based
quantitative proteomics using high intensity focused ultrasound. J Proteome Res 7:3860-7.
https://doi.org/10.1021/pr800161x

Lorenzo O, Piqueras R, Sánchez-Serrano JJ, Solano R (2003) ETHYLENE RESPONSE
FACTOR1 integrates signals from ethylene and jasmonate pathways in plant defense. Plant Cell
15:165-78. https://doi.org/10.1105/tpc.007468

Kong F, Yang L (2023) Pathogen-triggered changes in plant development: Virulence strategies
or host defense mechanism? Front Microbiol 14:1122947.
https://doi.org/10.3389/fmicb.2023.1122947

Krishna R, Ansari WA, Soumia PS, Singh M, Singh B (2022) Biotechnological interventions in
tomato (*Solanum lycopersicum*) for drought stress tolerance: achievements and future prospects.
BioTech 11:48. https://doi.org/10.3390/biotech11040048

Kunos V, Cséplő M, Seress D, Eser A, Kende Z, Uhrin A, Bányai J, Bakonyi J, Pál M, Mészáros
K (2022) The stimulation of superoxide dismutase enzyme activity and its relation with the
*Pyrenophora teres* f. teres infection in different barley genotypes. Sustainability 14:2597.
https://doi.org/10.3390/su14052597

Li XY, Zeng ZX, Cheng ZX, Wang YL, Yuan LJ, Zhai ZY, Gong W (2023) Common
pathogenic bacteria-induced reprogramming of the host proteinogenic amino acids metabolism.
Amino Acids 55:1487-1499. https://doi.org/10.1007/s00726-023-03334-w

Liu W, Liu K, Chen D, Liu Y, Mao Y, Liu H (2022) *Solanum lycopersicum*, a model plant for
the studies in developmental biology, stress biology and food science. Foods 11:2402.
https://doi.org/10.3390/foods11162402

Meena KK, Sorty AM, Bitla UM, Choudhary K, Gupta P, Pareek A, Singh DP, Prabha R, Sahu
PK, Gupta VK, Singh HB, Krishanani KK, Minhas PS (2017) Abiotic stress responses and
microbe-mediated mitigation in plants: the omics strategies. Front Plant Sci 8:172.
https://doi.org/10.3389/fpls.2017.00172

Mhlongo MI, Piater LA, Madala NE, Labuschagne N, Dubery IA (2018) The chemistry of plant-
microbe interactions in the rhizosphere and the potential for metabolomics to reveal signaling
related to defense priming and induced systemic resistance. Front Plant Sci 9:112.
https://doi.org/10.3389/fpls.2018.00112

Mhlongo MI, Piater LA, Steenkamp PA, Labuschagne N, Dubery IA (2020) Metabolic profiling
of PGPR-treated tomato plants reveal priming-related adaptations of secondary metabolites and
aromatic amino acids. Metabolites 10:210. https://doi.org/10.3390/metabo10050210

Nishad R, Ahmed T, Rahman VJ, Kareem A (2020) Modulation of plant defense system in
response to microbial interactions. Front Microbiol 11:1298.
https://doi.org/10.3389/fmicb.2020.01298

Patani A, Prajapati D, Ali D, Kalasariya H, Yadav VK, Tank J (2023) Evaluation of the growth-
inducing efficacy of various *Bacillus* species on the salt-stressed tomato (*Lycopersicon*
*esculentum* Mill.). Front Plant Sci 14:1168155. https://doi.org/10.3389/fpls.2023.1168155

Pandey AK, Dinesh K, Nirmala NS, Prajapati GK, Kumar M, Singh A (2023) Insight into tomato
plant immunity to necrotrophic fungi. Curr Res Biotechnol 6:100144.
https://doi.org/10.1016/j.crbiot.2023.100144

Pieterse CM, Zamioudis C, Berendsen RL, Weller DM, Van Wees SC, Bakker PA (2014)
Induced systemic resistance by beneficial microbes. Annu Rev Phytopathol 52:347-375.
https://doi.org/10.1146/annurev-phyto-082712-102340

Rajput VD, Harish, Singh RK, Verma KK, Sharma L, Quiroz-Figueroa FR, Meena M, Gour VS,
Minkina T, Sushkova S, Mandzhieva S (2021) Recent developments in enzymatic antioxidant
defence mechanism in plants with special reference to abiotic stress. Biology (Basel) 10:267.
https://doi.org/10.3390/biology10040267

Salehin A, Puri RR, Hafiz MHR, Itoh K (2021) Effect of co-inoculation of Bacillus sp. strain
with bacterial endophytes on plant growth and colonization in tomato plant (*Solanum*
*lycopersicum*). Microbiol Res 12:480-490. https://doi.org/10.3390/microbiolres12020032

Schmey T, Tominello-Ramirez CS, Brune C, Stam R (2024) *Alternaria* diseases on potato and
tomato. Mol Plant Pathol 25:e13435. https://doi.org/10.1111/mpp.13435

Schultz JC, Appel HM, Ferrieri AP, Arnold TM (2013) Flexible resource allocation during plant
defense responses. Front Plant Sci 4:324. https://doi.org/10.3389/fpls.2013.00324

Shan Y, Wang D, Zhao FH, Chen WL, Wang MJ, Sun FL, Wang HB, Li CY, Wu XF, Qiu DW
(2024) Insights into the biocontrol and plant growth promotion functions of *Bacillus altitudinis*
strain KRS010 against *Verticillium dahliae*. BMC Biol 22:116. https://doi.org/10.1186/s12915-
024-01913-1

Singh DP, Bisen MS, Prabha R, et al (2023) Untargeted metabolomics of *Alternaria solani*-
challenged wild tomato species *Solanum cheesmaniae* revealed key metabolite biomarkers and
insight into altered metabolic pathways. Metabolites 13:585.
https://doi.org/10.3390/metabo13050585

Singh DP, Rai N, Farag MA, Maurya S, Yerasu SR, Bisen MS, Prabha R, Shukla R, Behera TK (2024) Metabolic diversity, biosynthetic pathways, and metabolite biomarkers analysed via untargeted metabolomics and the antioxidant potential reveal for high temperature tolerance in tomato hybrid. Plant Stress 11:100420. https://doi.org/10.1016/j.stress.2024.100420

Stringlis IA, Yu K, Feussner K, de Jonge R, Van Bentum S, Van Verk MC, Berendsen RL, Bakker PA, Feussner I, Pieterse CM (2018) Root transcriptional dynamics induced by beneficial rhizobacteria. Nature Communications 9:1-10. https://doi.org/10.1038/s41467-018-07204-y

Sun W, Yu H, Liu M, Tang Z, Chen Y, Lu T, Ji S (2021) Evolutionary research on the expansin protein family during the plant transition to land provides new insights into the development of Tartary buckwheat fruit. BMC Genomics 22:252. https://doi.org/10.1186/s12864-021-07562-w

Tcherkez G, Carroll A, Abadie C, Mainguet S, Davanture M, Zivy M (2020) Protein synthesis increases with photosynthesis via the stimulation of translation initiation. Plant Science 291:110352. https://doi.org/10.1016/j.plantsci.2019.110352

Tharanath AC, Upendra RS, Rajendra K (2024) Soil Symphony: a comprehensive overview of plant–microbe interactions in agricultural systems. Applied Microbiology 4:1549-1567. https://doi.org/10.3390/applmicrobiol4040106

Timofeeva AM, Galyamova MR, Sedykh SE (2024) How Do Plant Growth-Promoting Bacteria Use Plant Hormones to Regulate Stress Reactions? Plants 13:2371. https://doi.org/10.3390/plants13172371

Tiwari M, Pati D, Mohapatra R, Sahu BB, Singh P (2022) The impact of microbes in plant immunity and priming induced inheritance: a sustainable approach for crop protection. Plant Stress 4:100072. https://doi.org/10.1016/j.stress.2022.100072

Torres MA, Jones JD, Dangl JL (2006) Reactive oxygen species signaling in response to pathogens. Plant Physiol 141:373-8. https://doi.org/10.1104/pp.106.079467

Vassilev N, Malusà E (2021) Special Issue: Microorganisms and Plant Nutrition. Microorganisms 9:2571. https://doi.org/10.3390/microorganisms9122571

Vejan P, Abdullah R, Khadiran T, Ismail S, Nasrulhaq Boyce A (2016) Role of plant growth promoting rhizobacteria in agricultural sustainability—a review. Molecules 21:573. https://doi.org/10.3390/molecules21050573

Vogt T (2010) Phenylpropanoid biosynthesis. Molecular Plant 3:2-20.
https://doi.org/10.1093/mp/ssp106

Wang Y, Ji D, Chen T, Li B, Zhang Z, Qin G, Tian S (2019) Production, Signaling, and
Scavenging Mechanisms of Reactive Oxygen Species in Fruit-Pathogen Interactions. Int J Mol
Sci 20:2994. https://doi.org/10.3390/ijms20122994

Yan S, Bhawal R, Yin Z, Zhou Y, Liu J, An Y, Zhang Z, Li J (2022) Recent advances in
proteomics and metabolomics in plants. Mol Hortic 2:17. https://doi.org/10.1186/s43897-022-
00038-9

Zhang S, Li C, Si J, Han Z, Chen D (2022) Action mechanisms of effectors in plant-pathogen
interaction. Int J Mol Sci 23:6758. https://doi.org/10.3390/ijms23126758

784   Table 1.

785   Table 1. Gene Ontology (GO) terms for the unique proteins found in Control, BV7 inoculated and Pathogen

786   infected tomato plants having distinguished changes in terms of the 'biological processes' due to interaction

787   with *Bacillus subtilis* (BV7) and the Pathogen

788
789   a.  GO enrichment terms (41) for the 'biological process' of unique proteins in Control
790

| Sr. No. | GO term description | GO term ID | Observed gene count | Strength | False discovery rate (FDR) |
|---|---|---|---|---|---|
| 1. | Cellular process | GO:0009987 | 244 | 0.17 | 9.59E-16 |
| 2. | Translation | GO:0006412 | 31 | 0.67 | 9.66E-09 |
| 3. | Amide biosynthetic process | GO:0043604 | 32 | 0.63 | 1.41E-08 |
| 4. | Gene expression | GO:0010467 | 56 | 0.4 | 2.38E-07 |
| 5. | Cellular amide metabolic process | GO:0043603 | 33 | 0.56 | 2.38E-07 |
| 6. | Cellular nitrogen compound biosynthetic process | GO:0044271 | 43 | 0.45 | 1.13E-06 |
| 7. | Cellular macromolecule metabolic process | GO:0044260 | 61 | 0.35 | 1.36E-06 |
| 8. | Cellular metabolic process | GO:0044237 | 145 | 0.18 | 1.79E-06 |
| 9. | Organonitrogen compound biosynthetic process | GO:1901566 | 44 | 0.43 | 2.01E-06 |
| 10. | Cellular component organization or biogenesis | GO:0071840 | 67 | 0.32 | 2.39E-06 |
| 11. | Ribonucleoprotein complex biogenesis | GO:0022613 | 22 | 0.64 | 5.94E-06 |

| 12. | Cellular nitrogen compound metabolic process | GO:0034641 | 75 | 0.28 | 9.81E-06 |
|---|---|---|---|---|---|
| 13. | Cellular macromolecule biosynthetic process | GO:0034645 | 33 | 0.48 | 1.05E-05 |
| 14. | Ribosome biogenesis | GO:0042254 | 18 | 0.64 | 8.78E-05 |
| 15. | Nitrogen compound metabolic process | GO:0006807 | 122 | 0.18 | 0.0001 |
| 16. | Cellular component organization | GO:0016043 | 56 | 0.3 | 0.00021 |
| 17. | Cellular component biogenesis | GO:0044085 | 35 | 0.4 | 0.00021 |
| 18. | Organelle organization | GO:0006996 | 42 | 0.35 | 0.00024 |
| 19. | Macromolecule biosynthetic process | GO:0009059 | 37 | 0.37 | 0.00039 |
| 20. | Metabolic process | GO:0008152 | 164 | 0.13 | 0.00041 |
| 21. | Macromolecule metabolic process | GO:0043170 | 109 | 0.17 | 0.00063 |
| 22. | Proteasome-mediated ubiquitin-dependent protein catabolic process | GO:0043161 | 12 | 0.73 | 0.00087 |
| 23. | Organonitrogen compound metabolic process | GO:1901564 | 90 | 0.19 | 0.001 |
| 24. | Establishment of localization in cell | GO:0051649 | 23 | 0.47 | 0.0011 |
| 25. | Organic substance biosynthetic process | GO:1901576 | 59 | 0.25 | 0.0016 |
| 26. | Cellular biosynthetic process | GO:0044249 | 57 | 0.25 | 0.0018 |
| 27. | Protein metabolic process | GO:0019538 | 73 | 0.22 | 0.0019 |
| 28. | Primary metabolic process | GO:0044238 | 135 | 0.13 | 0.0026 |
| 29. | Cellular localization | GO:0051641 | 29 | 0.38 | 0.0027 |
| 30. | Generation of precursor metabolites and energy | GO:0006091 | 19 | 0.46 | 0.0076 |
| 31. | Aerobic electron transport chain | GO:0019646 | 5 | 1.09 | 0.0109 |
| 32. | Mitochondrial ATP synthesis coupled electron transport | GO:0042775 | 5 | 1.09 | 0.0109 |
| 33. | Intracellular transport | GO:0046907 | 20 | 0.43 | 0.011 |
| 34. | Organic substance metabolic process | GO:0071704 | 143 | 0.11 | 0.0126 |
| 35. | rRNA processing | GO:0006364 | 11 | 0.61 | 0.0141 |
| 36. | Ubiquitin-dependent protein catabolic process | GO:0006511 | 14 | 0.5 | 0.0239 |
| 37. | Microtubule-based process | GO:0007017 | 10 | 0.61 | 0.0251 |
| 38. | Chloroplast localization | GO:0019750 | 4 | 1.16 | 0.0251 |

| 39. | Mitochondrial electron transport, NADH to ubiquinone | GO:0006120 | 3 | 1.37 | 0.0413 |
| 40. | Electron transport chain | GO:0022900 | 12 | 0.51 | 0.0414 |
| 41. | Proteolysis involved in protein catabolic process | GO:0051603 | 15 | 0.44 | 0.0476 |

792    b. GO enrichment terms (41) for the 'biological process' of unique proteins in BV7 inoculated plants

| Sr. No. | GO term description | GO term ID | Observed gene count | Strength | False discovery rate (FDR) |
|---|---|---|---|---|---|
| 1. | Metabolic process | GO:0008152 | 155 | 0.24 | 4.33E-14 |
| 2. | Small molecule metabolic process | GO:0044281 | 51 | 0.58 | 6.27E-13 |
| 3. | Cellular process | GO:0009987 | 176 | 0.16 | 4.70E-10 |
| 4. | Cellular metabolic process | GO:0044237 | 121 | 0.24 | 3.70E-09 |
| 5. | Organic substance metabolic process | GO:0071704 | 132 | 0.21 | 8.17E-09 |
| 6. | Carboxylic acid metabolic process | GO:0019752 | 32 | 0.62 | 1.17E-08 |
| 7. | Cellular amino acid metabolic process | GO:0006520 | 21 | 0.77 | 8.86E-08 |
| 8. | Small molecule biosynthetic process | GO:0044283 | 24 | 0.68 | 1.88E-07 |
| 9. | Organonitrogen compound biosynthetic process | GO:1901566 | 37 | 0.49 | 7.80E-07 |
| 10. | Organonitrogen compound metabolic process | GO:1901564 | 80 | 0.28 | 1.15E-06 |
| 11. | Biosynthetic process | GO:0009058 | 54 | 0.32 | 3.53E-05 |
| 12. | Cellular amino acid biosynthetic process | GO:0008652 | 12 | 0.87 | 5.31E-05 |
| 13. | Cellular biosynthetic process | GO:0044249 | 50 | 0.33 | 5.88E-05 |
| 14. | Carboxylic acid biosynthetic process | GO:0046394 | 17 | 0.67 | 6.47E-05 |
| 15. | Organic substance biosynthetic process | GO:1901576 | 51 | 0.32 | 7.14E-05 |
| 16. | Catabolic process | GO:0009056 | 34 | 0.42 | 8.83E-05 |
| 17. | Alpha-amino acid metabolic process | GO:1901605 | 14 | 0.74 | 0.00011 |
| 18. | Nitrogen compound metabolic process | GO:0006807 | 93 | 0.19 | 0.00028 |
| 19. | Alpha-amino acid biosynthetic process | GO:1901607 | 10 | 0.87 | 0.00036 |
| 20. | Primary metabolic process | GO:0044238 | 107 | 0.17 | 0.00043 |
| 21. | Organic substance catabolic process | GO:1901575 | 30 | 0.41 | 0.00043 |
| 22. | Organonitrogen compound catabolic process | GO:1901565 | 19 | 0.5 | 0.0027 |
| 23. | Nucleosome assembly | GO:0006334 | 5 | 1.2 | 0.0042 |
| 24. | Cellular catabolic process | GO:0044248 | 24 | 0.41 | 0.005 |
| 25. | Protein refolding | GO:0042026 | 5 | 1.12 | 0.0081 |
| 26. | Cellular nitrogen compound metabolic process | GO:0034641 | 50 | 0.24 | 0.0115 |
| 27. | Cellular nitrogen compound biosynthetic process | GO:0044271 | 26 | 0.36 | 0.0115 |
| 28. | Hydrocarbon biosynthetic process | GO:0120251 | 5 | 1.04 | 0.0174 |
| 29. | Generation of precursor metabolites and energy | GO:0006091 | 15 | 0.49 | 0.0201 |

| | | | | | |
|---|---|---|---|---|---|
| 30. | Olefinic compound biosynthetic process | GO:0120255 | 5 | 1 | 0.0252 |
| 31. | Riboflavin biosynthetic process | GO:0009231 | 3 | 1.47 | 0.027 |
| 32. | Peptide metabolic process | GO:0006518 | 16 | 0.45 | 0.0291 |
| 33. | Vitamin biosynthetic process | GO:0009110 | 5 | 0.94 | 0.0333 |
| 34. | Ethylene biosynthetic process | GO:0009693 | 4 | 1.13 | 0.0333 |
| 35. | Pigment metabolic process | GO:0042440 | 6 | 0.83 | 0.0333 |
| 36. | Protein-containing complex assembly | GO:0065003 | 13 | 0.5 | 0.0333 |
| 37. | Organophosphate metabolic process | GO:0019637 | 15 | 0.45 | 0.0338 |
| 38. | Cellular amide metabolic process | GO:0043603 | 17 | 0.41 | 0.0374 |
| 39. | NADP metabolic process | GO:0006739 | 4 | 1.08 | 0.0384 |
| 40. | Monocarboxylic acid metabolic process | GO:0032787 | 12 | 0.5 | 0.0443 |
| 41. | Chromatin assembly | GO:0031497 | 6 | 0.78 | 0.0477 |

794 c. GO enrichment terms (7) for the 'biological process' of unique proteins in Pathogen inoculated plants

| Sr. No. | GO term description | GO term ID | Observed gene count | Strength | False discovery rate (FDR) |
|---|---|---|---|---|---|
| 1. | Metabolic process | GO:0008152 | 60 | 0.21 | 0.0055 |
| 2. | Cellular biosynthetic process | GO:0044249 | 26 | 0.43 | 0.0055 |
| 3. | Organic substance biosynthetic process | GO:1901576 | 27 | 0.43 | 0.0055 |
| 4. | Organic substance metabolic process | GO:0071704 | 55 | 0.21 | 0.0059 |
| 5. | Cellular metabolic process | GO:0044237 | 49 | 0.23 | 0.012 |
| 6. | Cellular process | GO:0009987 | 70 | 0.14 | 0.0242 |
| 7. | Small molecule metabolic process | GO:0044281 | 17 | 0.48 | 0.0242 |

795