Homeostatic bidirectional plasticity in Upbound and Downbound micromodules in a model of the olivocerebellar loop

Elías M. Fernández Santoro[1], Lennart P.L. Landsmeer[1,2], Said Hamdioui[2], Christos Strydis[1,2], Chris I. De Zeeuw[1,3], Aleksandra Badura[1*] and Mario Negrello[1*]

[1] Department of Neuroscience, Erasmus MC, Rotterdam, The Netherlands

[2] Quantum & Computer Engineering Department, Delft University of Technology, Delft, The Netherlands

[3] Netherlands Institute for Neuroscience, Royal Academy of Arts and Sciences, Amsterdam, The Netherlands

*Correspondence:

Mario Negrello (m.negrello@erasmusmc.nl)

Aleksandra Badura (a.badura@erasmusmc.nl)

Department of Neuroscience, Erasmus MC

Wytemaweg 80, 3015 CN Rotterdam

tel: 0031-(0)10 7043589

Abstract

Olivocerebellar learning is highly adaptable, unfolding over minutes to weeks depending on

the task. However, the stabilizing mechanisms of the synaptic dynamics necessary for

ongoing learning remain unclear. We constructed a model to examine plasticity dynamics

under stochastic input and investigate the impact of inferior olive (IO) reverberations on

Purkinje cell (PCs) activity and synaptic plasticity. We explored Upbound and Downbound

cerebellar micromodules, which are organized loops of IO neurons, cerebellar nuclei neurons

and microzones of PCs characterized by their unique molecular profiles and different levels

of baseline firing. Our findings show synaptic weight convergence followed by stability of

synaptic weights. In line with their relatively low and high intrinsic firing, we observed that

Upbound and Downbound PCs have a propensity for potentiation and depression,

respectively, with both PC types reaching stability at differential levels of overall strength of

their parallel-fiber (PF) inputs. The oscillations and coupling of IO neurons participating in

the Upbound and Downbound modules determine at which frequency band PFs can be

stabilized optimally. Our results indicate that specific frequency components drive IO

resonance and synchronicity, which, in turn, regulate temporal patterning across Upbound

and Downbound zones, orchestrating their plasticity dynamics.

Author Summary

The olivocerebellar system is a part of the brain that facilitates learning and controlling

movements. To coordinate movements it integrates sensorimotor information with motor

command signals. The resulting behaviour needs to be continuously adjusted during motor

learning. The leading hypothesis is that changes in strength of the synaptic connections

between neurons underlie the learning process. The challenge is to elucidate the factors that

determine the exact timing and precision of the learned movements. To answer this question

we developed a computational model of the two main types of modules of the olivocerebellar

system that can control different types of movements in a bidirectional fashion. We found

that the rhythm and coupling of the olivary neurons play an important role in controlling and

stabilizing plasticity in the cerebellar cortex of both types of modules, together shaping

learning-dependent timing of motor behaviour.

Introduction

Procedural learning requires integration of sensory inputs with motor outputs, demanding circuits capable of both plasticity and stability. Sensorimotor control systems therefore rely on recurrent loops with feedback connections, enabling adaptive changes in synaptic strength based on ongoing activity patterns [1–3]. Among these systems, the olivocerebellar system stands out for its well-characterized plasticity mechanisms that are tightly linked to sensorimotor contingencies [4]. The olivocerebellar loop consists of the cerebellar cortex (CC), cerebellar nuclei (CN) and the inferior olive (IO), which feeds back into the CC via the climbing fibers (CFs) [5]. The sole output of the CC is generated by the inhibitory Purkinje cells (PCs), which receive sensorimotor information from the excitatory mossy fiber (MF) - parallel fiber (PF) pathway as well as the excitatory CF pathway.

The extensive dendritic trees of PCs receive signals from many PFs, all of which originate from granule cells [6,7]. Together with the inhibitory molecular layer interneurons, the excitatory PFs can modulate the simple spike (SSpk) activity of PCs, allowing for precise bidirectional control [8,9]. Individual PFs form relatively weak synaptic connections with PCs, [10], but as a group they can readily double the intrinsic SSpk firing frequency when needed for online sensorimotor control [11]. In contrast, in most mammals, each PC receives a single CF from the IO, which delivers a powerful excitatory input resulting in an all-or-none complex spike [(CSpk); [4]]. Different from the SSpks (~50-100 Hz), the CSpks occur at a relatively low firing frequency (~1 Hz) during spontaneous activity and they can increase their instantaneous firing frequency about ten-fold during modulation [8]. Since the CSpks are triggered by the IO activity [12], they reflect the integration of its excitatory signals from one or more of the ascending or descending afferent systems [13,14] and its inhibitory input signals from the hindbrain [15–18]. Interestingly, the nuclei that form the sources of the

77   excitatory inputs to the IO are largely overlapping with the sources of the MF pathways that

78   provide the sensorimotor signals to the CC via the PFs [19].

79   During learning, PF to PC synapses undergo activity-dependent changes guided by

the CF signals [20]. Classical models describe long-term depression (LTD) of these synapses

when PF and CF are co-activated within a short time window [20,21], and long-term

potentiation (LTP) when PF activity occurs outside this window [22]. However, alternative

plasticity mechanisms have been proposed, where bidirectional and even inverse plasticity

patterns have been observed depending on the pattern of PF input, the intrinsic firing

properties of PCs, and the local microcircuit context [23–27]. Maintaining a balance between

potentiation and depression is crucial to avoid unbounded growth or elimination of synaptic

weights [28–30], enabling the PF-PC synapses to remain responsive while supporting long-

term adaptation to input statistics.

89   Question remains how stability can be achieved in the olivocerebellar loops with

ongoing sensorimotor activity and following procedural learning. Several studies have shown

that activity-dependent synaptic plasticity is generally rather unstable in recurrent circuits

[1,31,32], often leading to difficulties with network dynamics [1,30,33–36]. Solving this

question for the olivocerebellar system becomes particularly challenging when taking into

account the heterogeneity of this network. Specifically, PCs in different cerebellar

microzones display distinct intrinsic properties, with the PCs expressing Aldolase-C (also

known as Zebrin II-positive PCs) typically exhibiting lower spontaneous SSpk firing

frequency (30-90 Hz) compared to Aldolase-C-negative PCs (60-120 Hz) [9,37,38]. These

electrophysiological differences are accompanied by divergent patterns of connectivity and

plasticity: Aldolase-C-positive and -negative zones differ in their afferent and efferent

projections [37,39–42], and they exhibit different propensities for long-term synaptic

plasticity at their PF to PC synapses [43].

Building on these distinctions, the conceptual framework of so-called "Upbound" (Aldolase-C-positive) and "Downbound" (Aldolase-C-negative) modules was introduced to capture the observation that different microzones support distinct plasticity regimes during learning [44–46]. In particular, because of their relatively low baseline firing frequency, the Upbound modules are characterized by a propensity for LTP over the course of training. Conversely the Downbound modules, which show high levels of baseline activity, show a propensity for LTD [9,43,47–49].

Our model explores the existence of common mechanisms in the olivocerebellar system that can stabilize changes in synaptic weights across modules that support distinct behaviours. We address this question by modeling how stability can emerge in a system where plasticity is continuously shaped by the circuit's own reverberating activity. Unlike recurrent excitatory networks, where Hebbian plasticity often leads to runaway instability, the olivocerebellar system forms a closed loop with distinct architecture: feedforward excitation, delayed inhibitory feedback, and intrinsic STOs in the IO. While this system lacks recurrent excitation, it still poses challenges for stability as the PF-PC synaptic plasticity is coupled to internally generated output. In this context, stability does not mean averting unbounded weight growth, but rather the emergence of robust synaptic weight distributions under ongoing plasticity and stationary input statistics. These dynamics reflect a homeostatic baseline regime in which Upbound and Downbound (Up/Downbound) modules emerge as stable configurations, rather than as pre-imposed initial conditions.

We hypothesize that the resonant dynamics of the olivocerebellar loops, particularly those shaped by the IO STOs and inhibitory feedback, are key to stabilizing PF-PC synaptic weights during procedural learning in both Up/Downbound modules. These dynamics may serve to temporally align CF input with relevant PF activity patterns, thereby promoting structural, frequency-dependent synaptic modifications across repeated exposure to the

similar input temporal regularities. In other words, establishing robust encoding of learned

associations between movements and the predictions of their sensory outcomes [44].

To test this hypothesis, we developed a biologically grounded network model of the

olivocerebellar system. The model incorporates reverberating, quasi-periodic activity [50–52]

and simulates micromodules with distinct electrophysiological properties of PCs, reflecting

Up/Downbound zones. A bidirectional, homeostatic plasticity mechanism, a Bienenstock-

Cooper-Munro-type (BCM) rule, jointly with an LTD vs. LTP mechanism, governs learning

at the PF-PC synapse. These rules are not used as a predefined stability constraint, but rather

to investigate how internally generated activity can lead to stable and interpretable synaptic

configurations under learning-like conditions. In our model, synaptic weights represent the

efficacy of the PF input onto the PCs and evolve over time according to the BCM rule. These

weights modulate the postsynaptic current (PSC) generated by each PF bundle, which in turn

shapes the SSpk activity. The model includes the core components of the olivocerebellar loop

(PCs, CN and IO neurons) with distinct parameter regimes corresponding to Up/Downbound

modules.

While the model simulates synaptic plasticity and evolving weight dynamics, it does

not incorporate explicit motor error signals or task-dependent modulation of CF activity.

Instead, it focuses on exploring how internally generated loop dynamics organize synaptic

weights in the absence of explicit motor error feedback. As such, the model captures baseline

synaptic homeostasis under fixed, ongoing input conditions rather than procedural learning

per se. In this context, it examines how the olivocerebellar loop can autonomously stabilize

its synaptic configuration, shaped by the interaction of intrinsic excitability, PF drive, and IO

feedback. We propose that these stable states may serve as a scaffold upon which learning-

related, supervised plasticity mechanisms (driven by behaviorally relevant CF signals) can

subsequently act. To test whether the stable Up/Downbound baseline states generated by our

model could support learning-related plasticity, we simulated a classical eyeblink

conditioning paradigm. Our data show that the stable Up/Downbound states persist and, upon

stimulus pairings, undergo distinct plasticity-driven changes in firing, consistent with

experimental findings.

Our simulations show that the differences between Up/Downbound zones are not

solely due to intrinsic properties of their PCs, rather, they also emerge from the dynamics of

the olivocerebellar loop. Moreover, different PF input frequency bands differentially drive

CSpk responses, suggesting that specific frequency components promote IO resonance and

this frequency selectivity is further shaped by IO gap junction coupling. Together, our data

highlight that temporal pattern selectivity processing across both Up/Downbound

micromodules is enhanced and stabilized by reverberations in the olivocerebellar loop during

procedural learning.

Results

Designing an olivocerebellar loop model to examine plasticity dynamics

Our hypothesis is that PF-PC synaptic weights are stabilized during procedural learning by

the resonant dynamics of the olivocerebellar loops. To test this, we developed a biologically

plausible model of the olivocerebellar system (**Fig. 1A**), which is decoupled from the rest of

the brain to focus on the olivocerebellar resonant network dynamics that drive plasticity. The

questions we tested with this model are centered on exploration of synaptic stability during

ongoing plasticity. We emulated ongoing arbitrary cortical and subcortical inputs to the

cerebellum via 5 uncorrelated PF bundles, which is conveyed by 100 PCs via 40 CN cells to

40 cells in the IO [53–60]. These IO cells have the ability to respond by advancing or

delaying their phase as a function of when inhibition arrives in the oscillation cycle, as

observed in *in vivo* experiments [16,61–65].

As input to our model network we used bundles of PFs as a noisy current input rather

than individual synapses. A bundle of PFs represents the combination of numerous synaptic

sources, as expected during the awake and behaving state [66,67]. To avoid biasing the input,

we modelled background activity as the sum of several independently generated Ornstein-

Uhlenbeck (OU) processes, taken to represent the integrated, continuous current from large

numbers of PF EPSCs [68]. This use of stochastic but temporally correlated input allowed us

to assess whether synaptic weights stabilize under a fixed input structure, enabling us to

isolate the contribution of internal loop dynamics, such as IO resonance and feedback, to

plasticity. This choice was motivated by theoretical considerations and in vivo experimental

data of granule cell and PF activity during learning, at rest and during walking [66,69,70],

which shows complex, high dimensional firing patterns transmitted to PCs in awake animals.

We refer to this temporally varied OU input as "frozen input" because it was generated once

and then held constant across all simulations. The input was not repeated or manipulated

across trials; it remained fixed throughout all simulations, serving as a controlled condition.

This design allows us to compare network behavior across simulation conditions while

ensuring that any observed differences could be attributed exclusively to the effects of

plasticity mechanisms, rather than to variations in the external input. This is a crucial aspect

of our model as we can test the hypothesis that under the stimulus representing sensorimotor

contingencies there would be synaptic adjustment and weight stabilization in the PF

bundles.Our PF bundles can be seen as bundles of correlated input originating via MF

rosettes [10,71]. Given that around 90% of PF synapses are silent [72], we modelled 5

bundles with approximately 200 active PFs each (**Fig. 1A-C**). The PF-PC synapse acts as a

"meta-synapse", representing the average direction of change of the individual synaptic

weights in the PF bundle. To represent plasticity, we added a weight value for each PF

bundle, which could change independently. Model parameters were tuned to match

spontaneous activity statistics reported experimentally — such as PC SSpk and CSpks firing

rates, CN output, and IO STOs. No tuning was performed to enforce the hypothesized

plasticity directions. Rather, the observed phenomena (e.g. synaptic weight dynamics, CF

pause modulation, and frequency-dependent IO entrainment) emerged from the interactions

of BCM and LTD plasticity, IO coupling, and intrinsic PC excitability under fixed input

conditions.

We tuned the PCs firing frequency to represent a mixture of their intrinsic firing

frequency and their response to the PF input. As shown experimentally [9], when the PF

input was completely absent, PCs had an intrinsic firing frequency that was about 20 Hz

lower than when the PF input was present. For simplicity we focussed on the primary drivers

of the olivocerebellar loop dynamics - the GABAergic CN projection cells, and excluded

cells involved in other loops. CN AdEx model cells were primarily designed to have robust

transmission of rate coding [73]. Our CN cells received divergent input from the PC layer,

which modulated CN responses in an anti-correlated fashion [73,74].

It is known that the IO contains a mixture of cells: some of which oscillate robustly,

some display dampened oscillations, and others that do not seem to display any resonant

behavior. Among the oscillating cells, many exhibit a low-frequency rhythm in the 4-10 Hz

range [52]. Our modelled IO cells oscillated robustly around 5.8-6 Hz and were connected

with electrotonic Connexin-36 synapses [52,75]. IO spiking synchrony was disrupted in

absence of gap junction coupling (**Fig. S1**). Though anatomical clusterization of cells is

highly plausible [55,76], here we examined a small set of cells without explicit clustering.

Figure 1 The olivocerebellar loop model and its population response (Downbound,
coupled). (A) Diagram of the olivocerebellar loop model. It consists of 5 PF bundles, 100
PCs, 40 CN neurons and 40 IO neurons. The projection (P) and convergence (C) ratios
between neuronal populations are indicated for each connection between populations.
Specifically, our model has 5 PF bundle inputs into 100 PCs. Each PC projects to 16 CNs.
The CN layer consists of 40 CN neurons, which receive input on average from 40 PCs (from
30 to 52 PCs). Each CN projects to 10 IO cells. There are 40 IO cells, each of them receiving
on average 10 CNs (from 6 to 16 CNs). The IO cells are modelled with gap junctions and
subthreshold oscillations. Half of the IO cells project back to the PCs with each of those IO
cells projecting to on average 5 PCs (from 2 to 9 cells), while each PC only receives signals
from 1 IO. Each IO neuron connects to all other IO neurons in the population via the gap
junctions. (B) From top to bottom: the PF bundle input modeled as Ornstein-Uhlenbeck
processes; PC responses, CN responses and IO responses (membrane potential and raster).
(C) Autocorrelation of each PF input. (D) Heatmap of the correlation of membrane
potentials within each population.
238
239
Homeostatic plasticity and CSpk-triggered plasticity mechanisms

To capture the essential interactions between LTP and LTD we modelled the PF-PC synapse

with a mixture of a BCM learning rule and a PF input dependent CSpk-triggered LTD

mechanism. This CF-induced LTD operates inversely to the classical BCM mechanism and

changes the probability of BCM-LTP [12]. In our model, the synaptic weights were

potentiated when PC activity was low to maintain excitability. This complex combination of

synaptic plasticity mechanisms enabled us to inspect their interactions and dynamics.

The BCM mechanism (**Fig. 2A**) in our model encapsulates the effect of changes in the

PF activity by either potentiating or depressing the synapse independently of CF input. This

mechanism depends on 3 factors: the average of the recent activity of the PC, ρ_{PC}; the BCM

threshold θ_M — which also depends on ρ_{PC}— and the instantaneous activity of the PF, ρ_{PF}

(**Fig. 2B**). After an IO spike (**Fig. 2C**), a pause in SSpks leads to a CF pause, which causes a

decrease in ρ_{PC}. The LTD component of plasticity due to the CSpk gives a negative change in

synaptic weight also proportional to the ρ_{PF} (**Fig. 2D**). The plasticity threshold

θ_M distinguishing depression from potentiation follows ρ_{PC}, that is θ_M lower than ρ_{PC} results

in potentiation. Conversely, θ_M higher than ρ_{PC} leads to depression. Both these changes are

proportional to activity ρ_{PF} for each synapse (see **Fig. 2E**). The total change in synaptic

weight is computed as the sum of two plasticity components: a BCM-based continuous

update and a CSpk-triggered term (**Fig. 2F**). The BCM plasticity mechanism has only one

hard constraint limiting the potentiation of the PC to frequencies of 250 Hz, as firing rate

beyond this rate is considered pathological.

After an initial transient, synaptic weights converged to stable distributions. However,

we observed small residual stabilizing oscillations around this stable mean, which we refer to

as weight fluctuations. These reflect the continuous balancing process of BCM potentiation

and CSpk-triggered LTD under stochastic input. Importantly, these fluctuations were minute

relative to the absolute weight values and did not show cumulative drift. While shaped by the

structure of the input —such as frequency content and temporal correlations— the

fluctuations were not dependent on the input duration, indicating a steady-state regime rather

than input-length dependence.

While the BCM rule used is not fitted to *in vitro* data, it captures key features of

activity-dependent potentiation and depression shaped by CSpk activity. This allowed us to

examine how loop-level dynamics guide synaptic trajectories. Different BCM parameter

choices may alter the speed of convergence but not the main results: weights stabilized, CSpk

frequency remained frequency-dependent, and Up/Downbound modules retained distinct

profiles. For example, we selected the threshold time constant (τ_M) within a range of 10-20

ms, based on the typical duration of CF pauses. This parameter governs the low-pass filtering

of recent PC activity used to compute the threshold θ_M. We did not tune τ_M to produce

specific outcomes; rather, this range was chosen *a priori* based on known physiology, and

only *a posteriori* did we find that it yielded expected plasticity directions —net depression in

Downbound zones and potentiation in Upbound zones.

Synaptic weights are stable across plasticity epochs

We hypothesized that synaptic weights would reach a new level of stability and stay at the

level for subsequent plasticity events. To investigate these changes in synaptic weights across

plasticity events we did the following simulations (see Methods), where for each epoch we

gave the same frozen input as PF input. We started with the first epoch ("no plasticity") with

the plasticity turned off and with static synaptic weights which were constant throughout this

120 s experiment. After this "no plasticity" epoch finished, BCM and LTD plasticity

mechanisms were enabled and the same frozen input was replayed. At the end of this

plasticity epoch, we took the last value of the filtered synaptic weights and turned it into the

new static weights for the final epoch (after plasticity). Again the same input was given one

last time with the new static weights. This configuration enabled us to compare synaptic

weights at the PF-PC synapse before and after plasticity. To evaluate the impact of plasticity

on the weights, each experiment was performed 4 times with the same frozen input (**Fig. 2G-**

L).

The same frozen input was used in each epoch. It provides consistent, varying input

statistics and temporal correlations across epochs ensuring reproducibility. When plasticity is

switched off, the synaptic weights remain fixed, allowing us to compare resonant activity at

stable weight configurations before and after plasticity. In contrast, during plasticity-on

epochs, the BCM and CSpk-triggered LTD mechanisms allowed the weights to settle into a

new homeostatic equilibrium.

For both Up/Downbound zones the direction of change of the 500 synaptic weights

for the 100 PC population (5 synapses per PC) was defined during the first "plasticity" epoch.

The synaptic weights at this epoch were reduced in the Downbound zone and increased in the

Upbound zone (**Fig. 2I-J**), suggesting that the intrinsic PC currents determined the plasticity

direction. From the second epoch after plasticity we found that the weights were mostly

stable, though a few changed in rank (**Fig. 2G-H**), indicating that the interaction between

CSpk and PF inputs may drive these rank changes. In Downbound zones weights fluctuated

around the mean, indicating a faster rate of weight change, due to the higher intrinsic PC

current (**Fig. 2J**). The Upbound zone, with a lower intrinsic drive, exhibited more synaptic

weight stability after relaxation in epochs 3 and 4 (**Fig. 2G-H**).

To test the robustness of these weight dynamics, we repeated the simulation protocol

using 10 different random seeds, each with independently generated OU input and randomly

sampled network parameters. For each seed, we computed the median weight change per

plasticity epoch (**Fig. 2K-L**). The same bidirectional trend was observed across all

simulations: Upbound modules exhibited early potentiation followed by stabilization, while

Downbound modules showed early depression followed by stabilization. These results

support the generalizability of our findings and indicate that the observed stabilization arises

from the intrinsic dynamics of the model rather than from input-specific contingencies.

319
320
Figure 2 The plasticity direction differs between Up/Downbound micromodules. (A) BCM
mechanism: the x-axis represents the activity of the PC, and the y-axis represents the
predicted plasticity change at a PF-PC synapse before scaling by PF activity. Positive values
indicate potentiation; negative values indicate depression. Each gray curve shows the BCM
function for a different value of the plasticity threshold θ. The colormap encodes the
corresponding θ values (in Hz), with lighter gray indicating lower thresholds. As PC activity
increases, the sliding threshold θ shifts rightward, reducing the potentiation range and
increasing the chances of depression (a larger part of the curve is negative). The same
applies for a lower activity in PC, leading to higher chance of potentiation. (B) CSpk-
triggered change in BCM. When there is a CSpk, the activity of the PC (blue) decreases. The
sliding threshold θ (black) follows and goes lower than the PC activity, leading to
potentiation (green). When θ is higher than the activity there is depression (red). This is
shown for both Up/Downbound zones. (C) Trace of the IO membrane potential for

Downbound coupled scenario. This trace was selected to illustrate the short-timescale effects
of different IO spike intervals on plasticity mechanisms (BCM and CSpk-triggered LTD).
While this particular trace appears bursty, it is not representative of the full IO population.
IO neurons in the model exhibit a range of firing modes including tonic firing and
*quiescence. The mean response profile shows a ~1 Hz firing rate (**Fig. 1B**) and the IO has a*
*large distribution of firing rates across the population (**Fig. 3B**). IO burstiness was not*
explicitly tuned and remained within biologically observed ranges (<6Hz), consistent with
*experimental findings [52,77–80]. (**D**) CSpk-triggered LTD component of synaptic plasticity.*
This shows the change in synaptic weight due to a IO spike (CSpk) event, proportional to the
instantaneous activity of the PF. In the absence of a CSpk, this component is zero; following
*a CSpk, it produces a transient depression that decays back to zero. (**E**) BCM component of*
synaptic plasticity. This reflects the change in synaptic weight based on the recent activity of
*the PC and the BCM threshold θ. It evolves continuously, independently of CF input. (**F**)*
Total synaptic weight update, computed as the sum of the BCM component and the CSpk-
triggered LTD component. All values are unitless and represent normalized synaptic efficacy
*changes per time step in the model. (**G**) Relative difference of synaptic weights of all PCs*
connecting to PF 1 before and after each plasticity (AP 1, 2, 3 and 4) between the epochs
*(current weight divided by the one of the previous event) for Upbound zones. (**H**) Same as I*
*but for the Downbound zones. (**I**) Boxplots depict absolute differences between the epochs for*
the whole population of weights (current weight minus previous weight) for Upbound zones.
*(**J**) Same as K but for the Downbound zones. (**K**) Median synaptic weight changes across four*
plasticity epochs for 10 independent Upbound simulations with randomized network and OU
*input per run. (**L**) Same as M, but for Downbound simulations. Synaptic weights are unitless*
and represent normalized efficacy values in the model.
358

Gap junction coupling creates a global phase and leads to synchronized firing

We hypothesize that gap junctions (i.e. "coupling") have an impact on IO STO synchrony in

that strong coupling leads to higher synchrony. To quantify this effect, we looked at the STO

oscillation phases over time. These were decomposed into a network level (global) phase,

defined as the mean Hilbert phase across all IO neurons, and into an individual-level (delta)

phase, defined as the difference between the Hilbert phase of each IO neuron and the global

phase at each time point. This was done to compare the difference between the individual and

global phases (**Fig. S2-1A**). The global phase of STOs, representing average phase

progression, was very stable in both coupled and uncoupled conditions (**Fig. S2-1B,S2-2B**).

In the condition with coupled IO cells, oscillators phase-locked onto the global phase creating

synchrony (**Fig. S2-1C**) not seen in the uncoupled case (**Fig. S2-2.C**), where oscillators

behaved independently. To investigate whether the coupling also led to dynamically

organized clusters in the IO, we calculated the pairwise Kuramoto phase (**Fig. S2-1D**). At the

scale of one second (**Fig. S2-1E**), transient clustering could be observed, however clusters

were not stable (**Fig. S2-1F-G**) and reflected random phase fluctuations driven by

GABAergic CN input. At the scale of 30 s (quarter-time of the full simulation) no clear

clustering was observed. Network synchrony peaked around IO spikes, but only for the

coupled scenario (**Fig. S2-2A**).

Upbound PCs have a propensity for potentiation and Downbound PCs for depression

As it has been shown that PCs adaptively adjust their SSpk firing rate via CSpks to elicit

blinks during eyeblink conditioning [45,81,82], we hypothesize that the direction of change

of the PC firing rate depends on the intrinsic frequency of each PC microzone. Eyeblink

conditioning is a classical associative learning paradigm where a neutral stimulus, such as

light, is paired with an air puff that evokes reflexive eyelid closure so that the initially neutral

stimulus can eventually trigger blinking. As argued previously, in this associative task,

bidirectional changes in the SSpk activity are expressed in the Downbound zone during initial

stages of training, whereas the Upbound zone controls acceleration of movement in

paradigms such as the vestibulo-ocular learning (**Fig. 3A**) [44]. In fact, several studies

support the distinction between these zones, proposing that they constitute complementary

modules that may act in concert to encode prediction and feedback signals during

sensorimotor learning [5,9,43,47–49]. Notably, while eyeblink conditioning involves learning

over many trials with discrete, stimulus-locked inputs and explicit error signals, our model

instead uses fixed, temporally varied, OU input, which was generated once and reused across

simulations to explore whether intrinsic firing properties and internal dynamics alone can

explain the directionality of plasticity. **Figure 3A** provides a schematic summary of

previously published experimental findings [44], illustrating the typical direction of SSpk

modulation in Up/Downbound PCs during eyeblink conditioning and vestibulo-ocular

learning. In contrast, **Figure 3B** shows how the same directionality emerges in our model

when plasticity mechanisms (BCM, LTD, LTP) are activated, but over much longer

timescales (seconds rather than hundreds of milliseconds. For display, we plot the raw 1-s

binned population mean together with a Savitzky-Golay-smoothed trace. The smoothing

applied across the full concatenated simulation produces small boundary effects at the epoch

transition. Nevertheless, the comparison highlights that, despite the different inputs and

timescales, the baseline level of PC activity is sufficient for the BCM mechanism to establish

the direction of synaptic weight change. PCs in our model were tuned to reproduce the

experimental data (**Fig. 3C-D**), whereby Downbound PCs showed relatively high SSpks (~90

Hz) and CSpks (~1.3 Hz) firing frequencies, while Upbound PCs had lower average SSpk

(~60 Hz) and CSpk [(~0.9 Hz), [9,44]]. Tuning PCs intrinsic currents resulted in the CF

pause reproducing experimental observations (**Fig. 3E**).

The direction of plasticity was the same irrespectively of the absence or presence of

IO gap junction coupling (**Fig. S3A-E**). For the Downbound micromodule, there was a small

increase in the mean of the CSpk firing frequency in the uncoupled case with respect to the

coupled case (**Fig. 3D** and **Fig. S3B**), which led to a wider distribution of the CSpk frequency

squared coefficient of variation [(CV2); (**Fig. S3F**)].

Figure 3. Comparison of experimental results and model results for Downbound and
Upbound zones. (A) Schematic summary (adapted from [44]) of SSpk modulation direction
in Up/Downbound modules during eyeblink conditioning and vestibulo-ocular learning. This
is not a direct experimental trace but a schematic summary of findings based on in vivo PC
recordings during eyeblink conditioning (Downbound) and vestibulo-ocular learning
(Upbound). (B) Model results showing population-averaged PC SSpk activity across the
three epochs (no plasticity, plasticity, after plasticity). Thin translucent trace: raw 1-s binned
population mean across PCs. Thick trace: Savitzky-Golay-smoothed version of the same
trace (window length 110 s, polynomial order 9), applied across the entire concatenated time
series. Minor boundary effects occur at the epoch transitions (vertical dashed lines). Note
that the timescales differ: in the model changes emerge over seconds, whereas in experiments
they appear over many trials and are expressed in relation to stimulus onset within hundreds
of milliseconds. (C-E) Comparison of experimental extracellular recordings in awake mice
from Up/Downbound-identified PCs, adapted from [9] and model results of PC firing
frequency, CSpk firing frequency and CF pause duration. Each gray line connects to the
paired value for a single model cell across the two modules. These pairs share all parameters

except for: (1) the intrinsic firing rate of the PCs and (2) the g_{Cal} and $I^{(IO)}{}_{OU}$ parameters of
the IOs, as shown in Table 7. There is a significant difference in means across
Up/Downbound modules for the SSpk (t=-13.78, p<0.001; two-sample Student's t-test;
n=100), CSpk (t=-4.59, p<0.001; two-sample Student's t-test; n=40), and CF pause (t=-
12.52, p<0.001; two-sample Student's t-test; n=40), which also has a smaller variance in the
Downbound zone (F=2.94, p<0.001; F-test using CCDF of F-distribution; n=40).
437

Inhibition followed by disinhibition of CN elicits CSpk

We hypothesized that specific characteristics of the input would elicit CSpks. We started by

analysing the CSpk-triggered PSC (**Fig. 4A**) and we observed that the CSpks were promoted

by specific oscillations present in the input. In our model, the PSC refers to the current that

enters the PC soma,specifically, the sum of the 5 PF currents, each scaled by their

corresponding synaptic weight. CSpks were evoked when the input carried specific phases

and frequency (around 5-6 Hz), with a preference for reduced input just before the CSpk. At

the level of the CN this translated to inhibition followed by disinhibition, which had

previously been empirically observed by [83,84]. The decrease (following an increase) in

PSC happened 60 ms before the bottom of the subthreshold oscillation (STO) of the IO

membrane response, which was closely related to the synaptic delay in the loop (10 ms

between PC and CN and 50 ms between CN and IO). This suggests that CSpks had a

preference for particular frequencies and phases of the input. A clear contrast is observed

after plasticity, as in Upbound zones the sensitivity to the PSC oscillations was increased and

reduced for Downbound zones.

Every CN spike contributed to a reset of the IO STOs (**Fig. 4B**), in line with

experimental results [16]. We found that in the Upbound zone there was a lower CN spike

frequency and lower amplitude of IO oscillations before the CN spike. As there was more CN

inhibition in the Downbound zone, a lower CN spike frequency led to reduced dampening of

the IO oscillation. In addition, we found that increasing CN inhibition lowered the frequency

of IO STOs while increasing IO spike (CSpk) output (**Fig. S4E**). In the Upbound zone, we

observed a lower CN spike frequency and lower amplitude of IO oscillations before the CN

spike, indicating weaker modulation of IO dynamics. In contrast, in the Downbound zone,

stronger CN inhibition slowed the IO STOs and reduced their amplitude, creating conditions

that increased the likelihood of IO spiking. This may be due to reduced oscillation amplitude,

resulting in milder hyperpolarization and thus a greater chance of crossing spike threshold.

Thus, CN activity modulates IO excitability indirectly, through effects on STO phase and

amplitude, rather than through direct excitation. These nonlinear interactions were further

shaped by PF-PC plasticity: IO responses to CN spikes varied depending on prior synaptic

changes, showing that the same input could produce variable IO responses depending on

network history. This is consistent with experimental observations of trial-to-trial variability

in CSpk timing [7,85], and supports the idea that IO STO phase and amplitude influence

CSpk generation [52]. After plasticity, CN spikes no longer significantly affected IO STO

frequency in the Downbound zone. This result suggests that the IO STOs are affected but not

driven by CN inhibition.

After the first "plasticity" epoch, the PC SSpk activity was increased for the Upbound

zone and decreased for Downbound zone, resulting in both PC populations firing at about 60

Hz after plasticity (**Fig. 4C**). As a result, the CN activity was decreased for the Upbound and

increased for the Downbound zone (**Fig. 4D**). Furthermore, the CSpk frequency was

decreased for Upbound PCs and increased for Downbound PCs (**Fig. 4E**). The length of the

CF pause decreased for both Up/Downbound zones (**Fig. 4F**).

Investigating the effect of gap junction coupling, we observed that the uncoupled

Downbound IO had on average a higher variance and higher CSpk firing frequency (**Fig.**

S4B), leading to a higher variance of SSpk firing frequency for Downbound PCs (**Fig. S4A**).

Nevertheless, the CN (**Fig. S4C**) and CF pause (**Fig. S4D**) were the same for both coupled

and uncoupled conditions.

Figure 4. Zone-dependent spiking resonances before and after plasticity. (A) CSpk-
triggered Postsynaptic current (PSC) recorded at the soma of the PC (top), IO membrane

potential (middle), and the PC membrane potential showing the CF pause (bottom). Traces
are aligned to the CSpk time (t=0), but the time window range of the CF pause trace differs
slightly to best illustrate its response. The synaptic delay between the PSC entering the PC,
going through the CN (10 ms) and reaching the IO (50 ms). As the inhibitory CN input
*coincides with the bottom of the IO STO, a CSpk is elicited. (**B**) CN spike triggered*
*membrane response of the IO, before and after plasticity. (**C**) Boxplots of SSpk frequency*
before and after plasticity. There there is a significant difference in the across the
*Up/Downbound zones before and after plasticity (before plasticity: **t=-13.90, p<0.001,** after*
*plasticity: **t=-5.36, p<0.001; two-sample Student's t-test; n=100**) and the variance is also*
*different after plasticity across the zones (**F=1.69, p=0.005; F-test using CCDF of F-***
***distribution; n=100**). There is a significant difference across plasticity epochs for both zones*
*(**Upbound: t=-7.64, p<0.001,** Downbound: **t=12.89, p<0.001; two-sample Student's t-test;***
n=100**). The variance of the SSpks after plasticity is also significantly different (**Upbound:
F=59.88, p<0.001,** Downbound: **F=36.00, p<0.001; F-test using CCDF of F-distribution;
***n=100**). (**D**) Boxplots of CN spiking frequency before and after plasticity (before plasticity:*
***t=-3.26, p=0.002,** after plasticity: **t=9.15, p<0.001,** Upbound: **t=5.05, p<0.001,** Downbound:*
***t=-7.26, p<0.001; two-sample Student's t-test; n=40**). (**E**) CSpk frequency before and after*
*plasticity (before plasticity: **t=3.98, p<0.001,** after plasticity: **t=9.44, p<0.001,** Upbound:*
***t=2.99, p=0.004,** Downbound: **t=-2.81, p=0.006; two-sample Student's t-test; n=40**). The*
*variance of CSpks is also different after plasticity across the zones (**F=2.91, p<0.001; F-test***
***using CCDF of F-distribution; n=40**). (**F**) CF pause length is significantly different before*
*plasticity (**t=-12.52, p<0.001; two-sample Student's t-test; n=40**). The length of the CF pause*
*is significantly modulated by plasticity in both zones (**Upbound: t=6.37, p<0.001,***
***Downbound: t=-13.00, p<0.001; two-sample Student's t-test; n=40**). The CF pause variance*
*was smaller after plasticity (**Upbound: F=20.76, p<0.001,** Downbound: **F=9.34, p<0.001; F-***
***test using CCDF of F-distribution; n=40**).*
513

Specific input frequency bands differentially drive plasticity

Our previous hypothesis was that specific characteristics of the input would directly influence

CSpk generation. Here, we expand our focus to include the full olivocerebellar loop,

proposing that distinct input frequency bands engage different resonant dynamics of the

circuit, thereby shaping both CSpk timing and the resulting plasticity. As a continuation of

the results shown in **Fig. 4**, we examined how different frequency bands shape loop-level

responses under plasticity. Signals from different sensorimotor sources have different spectral

properties, and PC population SSpks activity has been reported to contain bands from low

delta [86] to high gamma [50] frequencies.

To investigate how these spectral differences influence the loop dynamics under

plasticity we filtered the original PF signal into distinct frequency bands. The filtered signals

were normalized to keep the same spectral power (**Fig. 5A**), and were used in exact copies of

the same network to compare the before and after plasticity activity of the cells across the

different frequency bands (**Fig. 5B-D**). We found that CN dynamics were hardly modulated

by the filter classes, indicating high level of signal divergence and mixing (**Fig. 5C**).

As judged from both SSpk (**Fig. 5B**) and CSpk (**Fig. 5D**) distributions,

Up/Downbound PC zones showed marked differences in how their firing rates responded to

different PF input frequency bands. These differences were most pronounced for SSpks in the

upbound zones.

While the direction of plasticity of CSpks activity qualitatively resembled

physiological observations [9,43,44], frequency distributions were sensitive to plasticity

under different bands, with lower input frequencies driving most of the variability. This was

true for low frequencies (5-10 Hz band), where the CSpks activity was higher for

Up/Downbound zones. Furthermore, frequency bands larger than 10 Hz evoked similar

SSpks and CSspks distributions after plasticity, indicating that the PC network response was

robust to frequency variations.

Figure 5. Response of the olivocerebellar model after plasticity for Up/Downbound zones
for filtered inputs. *(**A**) Different types of filters used on the original input (no filter). (**B**)*
When comparing with the no filter input, the distribution of frequency of SSpks is always
different across frequency bands for the Upbound zone and in the case of the Downbound it
*is different between the 25-30 and 50-75 Hz frequency bands (for statistics see **Table S1**).*
The distribution of the SSpk frequency is significantly different between Up/Downbound
*across the different filter inputs (see **Table S1** for statistics). (**C**) The CN spike frequency*
*stays the same for all types of inputs across modules (see **Table S2** for statistics). (**D**) CSpks*
*frequency stays the same for all types of inputs across modules (for statistics see **Table S2**).*
*The response of the IO varies significantly for the 5-10 Hz frequency band (Upbound: **t=-***
7.86, p<0.001** and **D=0.75, p<0.001**, Downbound: **t=-4.04, p<0.001; two-sample Student's t-
***test** and **D=0.5, p<0.001; two-sample ks-test; n=40**).*

Differential CSpk response to PF input changes drives plasticity in 5-10 Hz band

The largest differential plasticity responses were observed for the 5-10 Hz filtered input in the

Downbound zone. Here we observed a large decrease in SSpk frequency after plasticity (**Fig.**

5B), with a twofold increase of CSpks (**Fig. S5A**). Before plasticity, the SSpk frequency was

approximately the same for the 5-10 Hz filtered input and the "no filter" case, as the current

input power spectrum and weights were the same. We assumed that the decrease in SSpk

frequency after plasticity originated from CSpk-triggered LTD. However, this assumption did

not explain the increase in CSpk frequency for the 5-10 Hz filtered PF input.

We looked at the relationship between the STO frequency and CSpk frequency (**Fig.**

S5B). We found that STO frequency and frequency-distribution width was narrower in the

Downbound than in the Upbound zone. For both Up/Downbound the STO frequency and IO

spike frequency were inversely related. This may seem unintuitive but is explained by

channel dynamics of IO cells, as more CN inhibition both increases the driving force and

reduces Ca-T type inactivation, leading to more prompt spiking upon disinhibition from DCN

(**Fig. S4E**) [16,52,87]. For both the Up/Downbound zone, 5-10 Hz input lowered the STO

frequency further, while at the same time increased the IO spike frequency. The lowering of

the STO frequency was more pronounced in the Upbound zone. We examined the IO spike-

triggered PSC to understand further how the input characteristics affected CSpk occurrences

(**Fig. S5C**). In general, for both IO unfiltered and 5-10 Hz input, IO spikes were preceded by

oscillations below or at the lower range of the STO frequency (5.6 Hz for Upbound, 5.4 Hz

for Downbound). For unfiltered noise, IO spikes were triggered by a small decrease in PSC

amplitude, whereas for 5-10 Hz input, IO spikes were triggered by a shortening of the PSC

oscillation period by 10 ms (**Fig. S5C**). Our results suggest that, for the unfiltered input (OU

input), the smaller frequencies, which are similar to the STO frequency, have a bigger impact

on the system.

Gap junction coupling drives frequency selectivity

Following our previous hypothesis that gap junctions affect IO STO synchrony, we

hypothesized that this change in STO synchrony will be reflected in the loop's response to

different frequencies. We observed that the frequency selectivity of the IO was lost when the

gap junction was uncoupled (**Fig. 6A**). The differences in the mean and distribution of CSpk

and SSpk frequency after plasticity were much less pronounced within the zones in the

uncoupled condition than in the coupled case. Furthermore, the higher IO spike frequency

and lower STO frequency was completely lost for the Upbound zone, but was increased for

the Downbound micromodule when the IO neurons were uncoupled (**Fig. 6B**). The loss of the

inverse STO - IO spike frequency relationship in the Upbound zone co-occurred with a loss

of the oscillatory response in the IO-spike triggered PSC (**Fig. 6C**). A higher CSpk frequency

for the Downbound zone (**Fig. S3B**) led to a less pronounced loss of the inverse STO - IO

spike frequency relationship. This suggests that the gap junction coupling has an effect

upstream to the PC-CN and back to the IO as there is less variation in the CN inhibitory input

onto the IO.

594
Figure 6. Gap junction drives frequency selectivity. (A) Firing frequency distributions for
SSpks (left) and CSpks (right) before and after plasticity for different frequency bands when
gap junctions are deleted. (B) STO frequency vs IO spike frequency for Upbound (left) and
Downbound (right) zones for before (top) and after (bottom) plasticity. Downbound IO has a
wider range of STO frequency and as a result a wider range of IO spikes also. (C) IO spike
triggered PSC response. Uncoupled Upbound zones lose or have much shorter entrainment
phase, while the Uncoupled Downbound still shows strong oscillatory response.

Baseline Up/Downbound states support task-related plasticity during conditioning

Our model reproduced bidirectional plasticity in PC activity in response to arbitrary inputs

(**Fig. 3B**). While frozen noise may approximate inputs in the awake and behaving state, it

does not reflect specific motor behaviors. To test whether the stable Up/Downbound baseline

states generated by the model could support learning-related plasticity, we reproduced a

classical eyeblink conditioning paradigm [88]. A conditioned stimulus (CS, 250 ms) was

delivered to one PF bundle and co-terminated with an unconditioned stimulus (US, 30 ms)

delivered to another PF bundle. A copy of the US was sent to the IO to elicit CF input, as in

standard protocols (**Fig. 7A**). Each block consisted of an initial US-only trial, ten CS+US

pairings, and a final CS-only trial, separated by 8-12s intervals (**Fig. 7B**).

The model captured key features of learning-related plasticity. The US elicited CSpks

across the IO population (**Fig. 7C**), and STO synchrony increased in the Upbound zone after

plasticity (**Fig. 7D**). Before learning, SSpks increased in response to the CS; this increase

disappeared after learning (**Fig. 7E**). These dynamics were consistent with the baseline states

in **Figure 4**, as seen in before/after plasticity comparisons of SSpks, CSpks and CF pause

(**Fig. 7F-H**).

Some discrepancies with experimental data remained. We did not observe a post-

learning decrease in SSpks, likely due to the absence of MLIs, which can suppress PC

activity in response to increased PF input [89,90]. Nor did we observe the CS-aligned CSpk

increase reported experimentally, which may depend on modulatory input from the striatum

or amygdala [91], both absent from our model. However, our model did replicate the

experimentally observed decrease in CSpk frequency in response to the US in the

Downbound zone after learning, suggesting that this effect arises from intrinsic

olivocerebellar dynamics.

626
Figure 7. Response of the olivocerebellar model to an eyeblink conditioning experiment
after plasticity for Up/Downbound zones *(**A**) Different PF bundles receive the CS and US*
*inputs. (**B**) A block of the eyeblink experiment consisting of an initial US (30 ms) followed by*
*10 CS+US pairings (220+30 ms), followed by a CS (220 ms). (**C**) CSpk-triggered responses*
of the IO population (40 cells) membrane potential (top) and raster (bottom) for before and
*after eyeblink training for Up and Downbound zones. (**D**) The Kuramoto synchrony between*
*all 40 IO cells. (**E**) CS-triggered frequency density responses of the populations for SSpk*
*(top) and CSpk (bottom). (**F**) Frequency responses for SSpk (before plasticity: **t=13.48,***
p<0.001**, after plasticity: **t=8.72, p<0.001**, Upbound: **-8.66, p<0.001**, Downbound: **t=12.96,
***p<0.001**; two-sample Student's t-test). The variance of SSpk frequency is significantly*
*different after plasticity across both zones (**F=5.87, p<0.001; F-test using CCDF of F-***
***distribution**) and the variance is also different for both zones across plasticity epochs*
*(Upbound: **F=42.01, p<0.001**, Downbound: **F=7.10, p<0.001; F-test using CCDF of F-***
***distribution**). (**G**) CSpk frequency boxplot (before plasticity: **t=3.68, p<0.001**, after plasticity:*
t=7.39, p<0.001; two-sample Student's t-test**). (**H**) CF Pause boxplot (before plasticity: **t=-
12.78, p<0.001**, after plasticity: **t=5.07, p<0.001**, Upbound: **9.68, p<0.001**, Downbound: **t=-
***10.27, p<0.001; two-sample Student's t-test**). The variance of the CF pause is significantly*
*different after plasticity across both zones (**F=1.83, p=0.001; F-test using CCDF of F-***
***distribution**) and the variance is also different for both zones across plasticity epochs*
*(Upbound: **F=15.62, p<0.001**, Downbound: **F=2.96, p<0.001; F-test using CCDF of F-***
***distribution**).

STO frequency decreases after plasticity for CS-US experiment

To better understand the learned response during CS-US trials, we examined PF

weight changes over time. PF1-3 and PF5 (US) exhibited similar weight dynamics (**Fig. S6-**

1A), while the PF4 (CS) weight showed a pronounced decrease during plasticity, likely

explaining the lack of CSpks at CS onset (**Fig. 7E**). Trial-averaged weights (**Fig. S6-1B**)

confirmed this pattern: the PF4 weight declined steadily across CS trials, whereas the other

PF weights remained stable. During the US, PF1-3 weights briefly decreased, then rapidly

returned to baseline, while PF5 remained unaffected. Before plasticity, increased PC firing

transiently accelerated STO frequency and global oscillator phase (**Fig. S6-1C**), an effect

abolished after plasticity.

When gap junctions were uncoupled, IO STO synchrony decreased overall but still

peaked near the US (**Fig. S6-2A**). SSpk responses were similar to the coupled case, but the

CSpk pattern reversed: the Downbound peak increased while the Upbound decreased post-

learning (**Fig. S6-2B**). Uncoupling also led to higher IO firing rates (**Fig. S6-2C**).

Discussion

How brains are able to maintain acquired sensorimotor patterns while continuous plasticity is a function of reverberating activity in all elements of the system is a central problem in computational neuroscience. The olivocerebellar system is a particularly interesting case to study continuous plasticity dynamics given that many of its neurons have a high level of intrinsic activity, organized in repeated feedforward loops [4,6,7,13,19,37]. To examine the plasticity dynamics of the olivocerebellar system, we developed a biologically grounded model representing distinct, so-called Up/Downbound modules within the network [9,43,44]. We tested the model independently in both conditions to allow direct comparison of the dynamics, stability, and plasticity rules that emerge within each module. The model does not simulate the modules interacting within a shared network, but treats them as parallel, anatomically independent compartments. Our aim was to capture and contrast the intrinsic stabilization dynamics of synaptic weights under ongoing plasticity learning behaviors in the isolated olivocerebellar system, thus keeping connectivity identical in both modules. In designing our model, we applied a reverse engineering approach and remained agnostic with respect to the specific functions that the cerebellar modules may compute, focusing on the closed loop dynamics of PF-PC plasticity subjected to resonances of IO spiking. We also excluded input to the IO and CN from cortical and subcortical structures and treated IO activity exclusively as a function of CN feedback and intrinsic oscillatory activity. Using this network model formed by the evolutionarily conserved, essential components, we analysed resonant dynamics of the loop, PC-PF weight distributions and stability under different frequency inputs, as well as the influence of IO gap junctions on frequency responses across populations. In what follows we will discuss the main results obtained by interrogating the model; these include (1), how steady state distributions of PC activity capture directions of plasticity in Up/Downbound modules, (2) which PF input frequencies most likely evoke

CSpks, (3) how synaptic weight distributions of the PFs of both modules stabilize, and (4) to

what extent this stabilization depends on gap junction coupling in the IO.

Steady state distributions of PC activity

In our model, PF-PC synapses were subject to two types of plasticity, BCM and

CSpk-dependent LTD. The BCM rule is a Hebbian activity dependent learning rule, and as

such is expected to reflect statistical regularities in the input [92,93]. However, in a

biophysically plausible setting with a large set of parameters and feedback there is no

guarantee that IO and PC responses remain bounded over time. To examine weight

convergence, we verified the weight values after multiple epochs of repeated, stochastic

input, modeled as OU noise. We found that synaptic weight distributions and firing rates

stabilized across epochs for both Up/Downbound micromodules. After an initial transient,

Upbound zones displayed more weight stability across epochs, while Downbound modules

showed small, bounded fluctuations around a stable mean, and thus were more sensitive to

loop dynamics. In effect, ongoing plasticity tuned PC activities in the same direction as

observed in physiological experiments [9,43,44] (**Fig. 3**); thus our model showed how

module-specific intrinsic PC firing rates shaped PC-PF plasticity direction. Plasticity did not

simply preserve the initial intrinsic firing asymmetry, but instead reconfigured PC activity

through closed-loop dynamics. Preferred activity levels emerged as a result of interactions

between intrinsic excitability, ongoing input, and network feedback. In this sense, the loop

does not enforce a fixed firing rate signature per module, but rather shapes zone-specific

attractors under homeostatic plasticity. The intrinsic current established a rate preference that

acted as a corrective "elastic force" on the weights. In the case of the Downbound module,

this corrective drive was less efficient at damping weight fluctuations, leading to slower

convergence and greater residual variability, whereas in the Upbound zone, convergence was

faster and more stable. This leads us to predict a higher changeability of synaptic weights in

Downbound zones, as the PC intrinsic rate is kept farther from circuit equilibrium in this

case. This model prediction is in line with the biological observation that proteins implicated

in synaptic plasticity are more prominently expressed in the Downbound zones, than those of

the Upbound zones [43,44,94]. Notably, recent transcriptomics studies indicate that the

subdiversity of different PC microzones is even greater than the two main categories of

Up/Downbound modules, particularly in primates [42,95,96]. Hence, future models will need

to consider reflecting the full heterogeneity of all PC populations.

Using our model's simulations we can conclude that while zone-specific PC

characteristics are known to be intrinsic [9,42–44], the olivocerebellar loop mechanisms

enhance this functional baseline difference and drive it to stability, engaging interactions of

BCM and CSpk-dependent long-term plasticity rules.

Input frequency dependent CSpk responses

Changes of weight of specific PCs in our model were strongly determined by their

recent input, both in terms of PF and CF input. Recent firing rate acted as a low pass proxy to

history of inputs, while the CSpk timing was strongly determined by the phase state of the IO.

We observed a strong preference for specific frequencies and phases of the stochastic PF

input and showed that CSpks probabilities were modulated by specific temporal features of

this input (**Fig. 4A-B**). As the state of the IO was subject to CN population drive, we showed

that the plasticity of the overall system was frequency-dependent.

Our choice of using OU PF noise [68] was motivated by emulating arbitrary input that

may exhibit any frequency and any phase combination. In the long interval of our plasticity

epochs (120 s), this OU input contained very different frequency compositions. This

modeling strategy was chosen to reflect *in vivo* experimental data of granule cell and PF

activity during learning, at rest and during walking [66,67,69,70], which show complex, high

dimensional firing patterns transmitted into PCs in awake animals. The variability of the IO

feedback and the resulting jitter in the latency of the CSpk response across trials, which we

show in our model, has also been observed experimentally [7,85] and can be explained by

phase shifts of IO STOs [52]. We analyzed the extent of the input frequency dependence by

filtering inputs to include only specific frequency bands. We discovered significant

differences in SSpk distributions to filtered inputs, which was particularly apparent in

Downbound zones. Specifically, we found that our model responded preferentially to the 5-

10 Hz frequency band, which overlapped with the IO STO frequency (5-6 Hz), indicating that

specific ranges of PF frequency bands led this model closer to physiological responses.

This frequency preference suggests that IO output is maximized when PF input

contains components near the intrinsic STO frequency of the IO (4-6 Hz). When STOs are

aligned with this input band, IO neurons are more likely to spike, thereby increasing the

probability of IO spiking. When we further investigated the relation between IO STO

frequency and IO spike output, we found them to be inversely related in both Up/Downbound

modules. This arose from differences in inhibitory CN feedback: stronger inhibition in the

Upbound zone slowed the STOs, effectively shifting them away from the preferred frequency

range and thus reducing the likelihood of IO spiking. In contrast, the Downbound zone

received weaker CN inhibition, resulting in faster STOs that remained better aligned with the

preferred frequency range, facilitating more frequent IO spikes. This mechanism

demonstrates how CN inhibition indirectly gates plasticity through STO tuning, and aligns

with *in vivo* recordings showing that CSpk rates differ across modules [7,85], with Upbound

zones typically displaying lower spontaneous CSpk frequencies than Downbound zones.

Stability of PF-PC synaptic weights

The overarching question that motivates our model is the apparent dilemma between

stability of acquired sensorimotor patterns and ever changing synaptic strengths. However,

we did not specifically test the loop in a behaviorally-relevant setting, rather we examined the

general case of any arbitrary input, reflecting ongoing sensorimotor behavior. Future research

with our model could study input signals that more closely represent the contingencies of

sensorimotor learning, with their specific frequencies and phases, related to the interaction to

the body and other brain regions. Upcoming effort could focus on adding the excitatory input

to the IO from the mesodiencephalic junction (MDJ) [16] and even the CN [13]; oscillatory

gating of the granular layer via Golgi cells [10,97]; and/or inhibition from the MLIs [89,90].

While we deliberately excluded the excitatory CN-MDJ-IO pathway and MF collaterals to

the CN to focus on the core feedback loop, we acknowledge that these inputs are likely to

modulate IO excitability and contribute to zonal differences. Experimental evidence shows

that excitatory input via the MDJ can evoke CSpks with much shorter latencies (4-8 ms) [15]

than the slower GABAergic inhibition of over 50 ms [79,98,99], and that MF input to the CN

may shift baseline excitability.

Our model was not tuned for specific outcomes but to test whether stable synaptic

states could emerge from internal loop dynamics. Resonance between PF and CF activity

drove frequency-dependent PF-PC synapses stabilization without task-specific tuning. We

interpret this as a baseline regime of the olivocerebellar loop, where Up/Downbound modules

settle into distinct steady states under stationary input and ongoing plasticity.

We extended this framework by incorporating a classical eyeblink conditioning

paradigm. The baseline states established by ongoing plasticity persisted when task-related

CF inputs were introduced. During CS-US pairings, Up/Downbound modules underwent

distinct learning-related adaptations, demonstrating that homeostatic stabilization provides a

functional substrate for supervised, CF-driven plasticity. This reconciles experimental

observations of stable baseline states with the emergence of biased plasticity during learning.

 Impact of gap junction coupling on stabilization

Gap junctions can cause synchronized IO spikes, in both Up/Downbound zones,

leading to a global phase in which many individual IO neurons are coupled [16,100]. When

the gap junctions were abolished in our model, the frequency selectivity of the IO response

was disrupted. In the Upbound module, this disruption was accompanied by a reduction in the

variability of IO spike rates and a loss of oscillatory structure in IO-triggered PSC responses.

However, in the Downbound module, IO spike variability persisted to a larger degree, likely

due to its intrinsically higher IO and PC firing rates and more robust CSpk activity. This

differential effect underscores the zone-specific contributions of IO electrotonic coupling to

timing and frequency selectivity. Moreover, this result suggested that the gap junctions have

a critical influence on the frequency selectivity of the system. Indeed, when tested, we found

that the frequency selectivity due to the difference in strength of the CN input, highlighted

above, was lost in the absence of IO gap junction coupling. As CN inhibition turned out to

not completely shape the IO STOs, this could indicate that the interplay between CN

inhibition and IO gap junction coupling is at the core of this STO - IO frequency relationship.

It should be noted that the interaction between GABAergic CN input to the IO and regulation

of the coupling strength as well as the relevance of coupling in the IO for the prominence of

its STO's has been the subject of many experimental studies. Both relations have been

confirmed at the morphological, electrophysiological and pharmaceutical level, *in vivo* as

well as *in vitro* [15,16,52,77–80,86,101–105]. The relationship between the level of STO's

and IO frequency band is underscored by the finding that Up/Downbound modules showed

different correlation strengths in this respect. Thus, as gap junction coupling facilitates IO

resonance and as there are module-specific interactions, our data imply that the temporal

pattern selectivity processing across Up/Downbound micromodules is enhanced by dynamic

regulation of electrotonic coupling in the IO. This was further confirmed in the eyeblink

conditioning paradigm: uncoupling reduced overall IO STO synchrony but preserved peak

synchrony around the US. Interestingly, CSpk responses reversed post-learning Downbound

increase and Upbound decrease, highlighting how coupling biases plasticity directionality. IO

firing rates also increased without coupling, suggesting that electrotonic interactions

constrain IO excitability and shape zone-specific plasticity. These modeling data align well

with experiments in which olivary coupling is affected either genetically or

pharmacologically; impairing dendrodendritic IO coupling hampers cerebellar sensorimotor

learning [75].

Limitations and biological generalization

Our model captures key internal dynamics of the olivocerebellar loop, however due to

epistemic considerations several biological features were explicitly excluded from the design.

The PF-PC synaptic plasticity was modeled using a BCM-inspired homeostatic rule in

combination with a CSpk-triggered LTD component, while leading to LTP in the absence of

CSpks. While this captures bidirectional plasticity and weight stabilization, it remains a

phenomenological approximation. Experimental data on PF-PC plasticity, particularly under

in vivo or awake conditions, suggest a large set of mechanisms that our simplified rule does

not fully represent [23–27]. For instance, the CF CSpk itself was not modeled with detailed

waveform dynamics, and thus the influence of CSpk duration, calcium influx or burst

structure on plasticity was not explored [106,107]. In terms of input, we employed stochastic

noise (OU process) to test how arbitrary fluctuations at different frequencies shape loop

dynamics and plasticity. This does not capture structured MF activity, such as bursts or

sequences associated with sensory events or behavior. Nor does it reflect modulations by

Golgi-Granule cell-mediated gating oscillations in the 8-12 Hz range [10,97]. However, we

observed that our model responded most strongly to frequencies near 10 Hz — consistent

with the filtering properties of the granular layer [97]. This suggests that Golgi cell-induced

frequency gating may reinforce IO resonance at these frequencies and further constrain

learning   windows.

To assess whether the stable Up/Downbound configurations could support learning,

we incorporated a classical eyeblink conditioning paradigm typically associated with

Downbound zones [5]. We applied the paradigm to both modules and found that CS-induced

SSpk increases were suppressed after learning, and US-triggered CSpks decreased in the

Downbound zone. Notably, these effects reversed when IO gap junctions were removed. A

recent model, incorporating MLIs, Golgi and granule cells replicated CS-induced SSpk

suppression and CSpk increase during conditioning [45], but did not examine the role of IO

reverberations on PC homeostatic plasticity. Our model shows that removing gap junctions

increased Downbound CSpks and reduced Upbound ones, suggesting IO coupling is required

for the timing of acquired eyeblink responses [75].

Our simplifications were designed to probe ongoing learning properties of the

resonant olivocerebellar loop. The model captures homeostatic stabilization under persistent

input, not task-driven synaptic reorganization. Future versions could test how structured

sensorimotor contingencies shape weight homeostasis, clarifying stabilization versus

learning-related plasticity. Overall, our work highlights the dynamic nature of this loop,

essential for understanding how the cerebellum "learns to learn".

Methods

Network architecture

To balance anatomical accuracy with computational tractability, we modeled each

micromodule with 100 PCs, 40 CN neurons and 40 IO neurons. These ratios are grounded in

known anatomical estimates: within a module, approximately 20-30 PCs converge onto a

single CN neuron, while each PC projects to ~30-50 CN neurons [58]. Half of IO neurons

connect to the PCs. Each of those IO neurons project to ~7-10 PCs, and each PC receives CF

input from a single IO neuron [59,60]. We modeled only the inhibitory CN neurons, as these

are the ones that project to the IO [53–57]. For the CN-IO synapse, we assumed a 25%

reciprocal connectivity probability, resulting in each CN neuron connecting ~10 IO neurons

and each IO neuron receiving from ~10 CN neurons. This structured yet simplified

architecture ensures that core convergence and divergence patterns of the circuit are

preserved while enabling efficient simulation (see **Fig. 1**).

Neuron models and justification

PCs and CN neurons were modeled using simplified adaptive exponential integrate-

and-fire neurons. This abstraction was sufficient to capture the key dynamics relevant for our

study: the modulation of PC firing rate by PF and CF input, and the inhibitory feedback from

PCs to CN. Furthermore, IO neurons are endowed with rather unique properties. They are

one of the most densely coupled via dendro-dendritic gap junctions than any other type of

neuron in the mammalian brain [101,102,108] and they express high levels of voltage-gated

and calcium-dependent ion-channels that allow them to oscillate at frequencies of up to 10 Hz

[52,77–80]. The interplay between the excitatory and inhibitory inputs to the IO as well as the

temporal relation with the subthreshold oscillations (STOs) of the coupled olivary cells

determines whether and when an olivary action potential will be generated [16,64,104,109].

Because our primary focus was on IO-driven resonances and loop-level plasticity, we

prioritized a more detailed multi-compartment IO model to accurately reproduce STOs and

phase-resetting phenomena. Modeling PC morphology or PC calcium dynamics was not

necessary for our hypothesis (see Introduction), which depended primarily on CF timing and

population-level PC output.

Parallel fiber bundles

We simulate the PF input, representing the activity of PF bundles, as an Ornstein-Uhlenbeck

(OU) process:

887
$$\frac{dI^{(PF)}{}_{OU}}{dt} = \frac{I^{(PF)}{}_{OU_0} - I^{(PF)}{}_{OU}}{\tau^{(PF)}{}_{OU}} + \frac{\sigma^{(PF)}{}_{OU}}{\sqrt{\tau^{(PF)}{}_{OU}}} \xi$$

888   Eq. (1)

889   where ξ is a Gaussian white noise with mean $\mu = 0$ and standard deviation $\sigma = 1$ that scales

890   with units of $s^{-0.5}$. The mean is $I^{(PF)}{}_{OU_0}$ and the variance —the integral of the power

891   spectral density— is equal to σ^2. The noise intensity is defined as $\sigma^2 \tau$. Parameter values in

892   the model are shown in **Table 1**.

894   **Table 1.** Parameters of the Ornstein-Uhlenbeck processes

Parameters	Baseline current I_{OU_0} (nA)	OU time constant τ_{OU} (ms)	OU standard deviation σ_{OU} (nA)
PF	1.3	50	0.25
IO	-0.3	50	0.3

896   *PC and CN neuronal models*

897   The PC and CN populations are modeled as adaptive exponential integrate-and-fire (AdEX)

898   models [110]. The general cell equations for the PC and CN are described as:

899
$$\frac{dV}{dt} = \frac{1}{C_m}\left(-g_l(V - V_l) + g_l \Delta_T e^{\left(\frac{V - V_T}{\Delta_T}\right)} + \sum_i I^{(i)} - w\right) \qquad \text{Eq. (2)}$$

900
$$\frac{dw}{dt} = \frac{a(V - V_l) - w}{\tau_w} \qquad \text{Eq. (3)}$$

Here, Eq. 2 describes the membrane potential dynamics, while Eq. 3 shows the dynamics of

the adaptation value. The $\sum_X I^{(i)}$ refers to the sum of the currents specific to cell type X

(where $X \in \{PC, CN\}$ and i indexes the current types affecting that cell); Δ_T is the slope

factor; V_T is the threshold potential; w is the adaptation variable; a is the adaptation coupling

factor, and τ_w is the adaptation time constant. The slope factor Δ_T is a quantification of the

sharpness of the spike, which can be seen as the sharpness of the sodium activation curve if

the activation time constant is ignored [111]. The parameters of all AdEx cells (both CN and

PC) were sampled using evenly spaced values across the listed range in **Table 2**. For each

parameter we generated a list of linearly spaced values equal to the population size (e.g. 100

for PCs), and assigned them to individual cells without replacement. This approach ensures a

uniform spread across the specified range and introduces controlled variability in intrinsic

properties across the population. Hence, each cell in the population will display individual

differences in firing properties. When a spike happens both the membrane potential and the

adaptive variables are updated. The actual threshold potential $V_{cut} = V_T + 5\Delta_T$ is used for the

neuron models. When the potential reaches V_{cut} (see **Table 2**), a spike occurs and both the

membrane potential V is reset to V_r, and the adaptation variable w is updated to the reset

adaptation current b and then decays until either reaching 0 or the next spike time [112].

$$@Cell\ spike: V = V_r;\ w += b \qquad\qquad \text{Eq. (4)}$$

Our goal for using the randomization of parameters is to have variability across the cells in

each population. Since the CN is inhibited by the PC we will already have enough variability

in the inhibitory current as each PC is different. Also, the CN cells have their own variability

across the parameters. Hence, we do not randomize the intrinsic current I_{int} of the CN as we

can then see the direct effect of one PC onto the CN.

924
Table 2. Parameters of the AdEx cell types

Parameters	PC	CN
Membrane capacitance C_m (pF)	$75 \pm 1.0[N_{PC}]$	$281 \pm 1.0[N_{CN}]$

Leakage conductance g_l (nS)	$30 \pm 1.0[N_{PC}]$	$30 \pm 1.0[N_{CN}]$
Leakage reversal potential V_l (mV)	$-70.6 \pm 0.5[N_{PC}]$	$-70.6 \pm 0.5[N_{CN}]$
Threshold potential V_T (mV)	$-50.4 \pm 0.5[N_{PC}]$	$-50.4 \pm 0.5[N_{CN}]$
Slope factor Δ_T (mV)	$2 \pm 0.5[N_{PC}]$	$2 \pm 0.5[N_{CN}]$
Adaptation time constant τ_w (ms)	$144 \pm 2.0[N_{PC}]$	$30 \pm 1.0[N_{CN}]$
Adaptation coupling factor a (nS)	$4 \pm 0.5[N_{PC}]$	$4 \pm 0.5[N_{CN}]$
Reset potential V_r (mV)	$-70.6 \pm 0.5[N_{PC}]$	$-65 \pm 0.5[N_{CN}]$
Reset adaptation current b (nA)	$0.0805 \pm 0.001[N_{PC}]$	$0.0805 \pm 0.001[N_{CN}]$
Intrinsic current I_{int} (nA)	$0.35 \pm 0.21[N_{PC}]$	1.2

<u>Purkinje cell Currents</u>

For the PC, there are 2 currents denoted $I^{(PC)}{}_{int}$ and I_{PF}. The former is the activation

(intrinsic) input current given to the cell to tune the desired intrinsic firing frequency

distribution. The latter is the postsynaptic current (PSC) input entering the PC soma from the

PFs (**Fig. 1A**).

$$\sum I_{PC} = I_{PF} + I^{(PC)}{}_{int} \qquad\qquad \text{Eq. (5)}$$

<u>Cerebellar nuclei Synaptic Currents</u>

The CN currents are shown in Eq. 6, where $I^{(CN)}{}_{int}$ is the intrinsic current of the CN

and $I^{(CN)}{}_{PC}$ is the inhibitory current from the PC to the CN. It is a filtered version of the PC

spike train that decays exponentially in the absence of spikes. Upon each PC spike, $I^{(CN)}{}_{PC}$ is

updated by a constant $\gamma^{(CN)}{}_{PC}$ (see **Table 6**). This means that the total current $I_{int} - I^{(CN)}{}_{PC}$

will be smaller and the CN will spike less (inhibition from PC).

$$\sum I_{CN} = I^{(CN)}{}_{int} - I^{(CN)}{}_{PC} \qquad\qquad \text{Eq. (6)}$$

Inferior olive model

The IO model is based on the De Gruijl model [107] —a modification of the

Schweighofer model [87] with an added axonic current. The general cell equations for the

model are shown below:

943
$$C_m \frac{dV_s}{dt} = -\sum I^{(s)} + I^{(s)}{}_{app} + I^{(IO)}{}_{OU} + I^{(IO)}{}_{CN} \qquad \text{Eq. (7)}$$

944
$$C_m \frac{dV_d}{dt} = -\sum I^{(d)} + I^{(d)}{}_{app} \qquad \text{Eq. (8)}$$

945
$$C_m \frac{dV_a}{dt} = -\sum I^{(a)} \qquad \text{Eq. (9)}$$

946
$$\frac{dI^{(IO)}{}_{OU}}{dt} = \frac{I^{(IO)}{}_{OU_0} - I^{(IO)}{}_{OU}}{\tau^{(IO)}{}_{OU}} + \frac{\sigma^{(IO)}{}_{OU}\xi}{\sqrt{\tau^{(IO)}{}_{OU}}} \qquad \text{Eq. (10)}$$

Where, the membrane potential specific to each compartment, somatic s, dendritic d and

axonic a is represented with V_i (where $i = s, d, a$). The equations for the ionic conductances

of the general cell are shown in **Table 3** and their parameters in **Table 4**. All parameters are

randomized in the same way as for the AdEx parameters. The IO receives an a noise input

$I_{IO_{OU}}$ modeled as an OU process (similarly to the PF input, see **Table 1**). With this input we

model the baseline firing rate of the IO as well as modeling inputs from other brain regions.

Furthermore, the inhibitory input from the CN onto the IO is modeled by the current $I^{(IO)}{}_{CN}$,

as described in the Synapses subsection. Finally, two applied currents $I^{(i)}{}_{app}$ (where $i = s, d$)

are in the dendrites and somatic compartments. These are used when adding external current

into the cell. The dynamics of each activation or inactivation variable (see **Table 5**) is given

by the differential equation below. In this equation z indicates the variables

$h, k, l, n, x, q, r, s, h_a$ and x_a. There are 2 exceptions, m and m_a which have a small enough τ

that they are approximately equal to their steady-states (m_∞ and $m_{\infty a}$, respectively).

960
$$\frac{dz}{dt} = \frac{(z_\infty - z)}{\tau_z} \qquad \text{Eq. (11)}$$

962   **Table 3.** Ionic conductances of the IO cell

Somatic currents $\sum I_s$	Dendritic currents $\sum I_d$	Axonic currents $\sum I_a$
$I^{(s)}{}_{Na} = g_{Na} m_\infty{}^3 h (V_s - V_{Na})$	$I^{(d)}{}_{Ca_h} = g_{Ca_h} r^2 (V_d - V_{Ca})$	$I^{(a)}{}_{Na} = g_{Na_a} m_{\infty a}{}^3 h_a (V_a - V_{Na})$
$I^{(s)}{}_{Ca_l} = g_{Ca_l} k^3 l (V_s - V_{Ca})$	$I^{(d)}{}_{KCa} = g_{KCa} s (V_d - V_K)$	$I^{(a)}{}_{Ka} = g_{Ka} x_a{}^4 (V_a - V_K)$
$I^{(s)}{}_{K_{dr}} = g_{K_{dr}} n^4 (V_s - V_K)$	$I^{(d)}{}_c = g_c (V_d - V_{de}) f (V_d - V_{de})$	$I^{(a)}{}_{la} = g_{la} (V_a - V_l)$
$I^{(s)}{}_{K_s} = g_{K_s} x_s{}^4 (V_s - V_K)$	$I^{(d)}{}_{ld} = g_{ld} (V_d - V_l)$	$I^{(a)}{}_{sa} = \dfrac{g_{int}}{p_2} (V_a - V_s)$
$I^{(s)}{}_h = g_h q (V_s - V_h)$	$I^{(d)}{}_{sd} = \dfrac{g_{int}}{(1 - p)} (V_d - V_s)$	
$I^{(s)}{}_{ls} = g_{ls} (V_s - V_l)$		
$I^{(s)}{}_{ds} = \dfrac{g_{int}}{p} (V_s - V_d)$		
$I^{(s)}{}_{as} = \dfrac{g_{int}}{(1 - p_2)} (V_s - V_a)$		

Table 4. Parameters of the IO Standard Cell

Conductances (mS/cm^2)		Reversal Potentials (mV)	
g_{Na}	$150 \pm 1.0[N_{IO}]$	V_{Na}	$55 \pm 1.0[N_{IO}]$
g_{Ca_l}	$1.4 \pm 0.05[N_{IO}]$	V_{Ca}	$120 \pm 1.0[N_{IO}]$
$g_{K_{dr}}$	$9.0 \pm 0.1[N_{IO}]$	V_K	$-75 \pm 1.0[N_{IO}]$
g_{K_s}	$5.0 \pm 0.1[N_{IO}]$	V_h	$-43 \pm 1.0[N_{IO}]$
g_h	$0.12 \pm 0.01[N_{IO}]$	V_l	$10 \pm 1.0[N_{IO}]$
g_{ls}	$0.017 \pm 0.001[N_{IO}]$	Membrane Capacitance (μF/cm^2)	
g_{Ca_h}	$4.5 \pm 0.1[N_{IO}]$	C$_m$	$1.0 \pm 0.1[N_{IO}]^2$
$g_{K_{Ca}}$	$35 \pm 0.5[N_{IO}]$	Cell Morphology	
g_{ld}	$0.016 \pm 0.001[N_{IO}]$	g$_{int}$	$0.13 \pm 0.001[N_{IO}]$
g_{Na_a}	$240 \pm 1.0[N_{IO}]$	p	$0.25 \pm 0.01[N_{IO}]$
g_{K_a}	$240 \pm 0.5[N_{IO}]$	p$_2$	$0.15 \pm 0.01[N_{IO}]$
g_{la}	$0.017 \pm 0.001[N_{IO}]$		

 Table 5. Steady-state activation/inactivation and time constants of IO ionic conductances

I	Steady-state Activation/Inactivation	Time constant τ (ms)	Forward rate function α	Backward rate function β
I_{Na}	$m_\infty = \dfrac{1}{1 + e^{-\left(\frac{30+V_s}{5.5}\right)}}$			
	$h_\infty = \dfrac{1}{1 + e^{\frac{70+V_s}{5.8}}}$	$\tau_h = 3e^{-\left(\frac{40+V_s}{33}\right)}$		
I_{Ca_l}	$k_\infty = \dfrac{1}{1 + e^{-\left(\frac{61+V_s}{4.2}\right)}}$	$\tau_k = 5$		
	$l_\infty = \dfrac{1}{1 + e^{\frac{85.5+V_s}{8.5}}}$	$\tau_l = 35 + \dfrac{20e^{\left(\frac{160+V_s}{30}\right)}}{1 + e^{\left(\frac{84+V_s}{7.3}\right)}}$		
$I_{K_{dr}}$	$n_\infty = \dfrac{1}{1 + e^{-\left(\frac{3+V_s}{10}\right)}}$	$\tau_n = 5 + 47e^{\left(\frac{50+V_s}{900}\right)}$		
I_{K_s}	$x_{\infty s} = \dfrac{\alpha_{x_s}}{\alpha_{x_s} + \beta_{x_s}}$	$\tau_{x_s} = \dfrac{1}{\alpha_{x_s} + \beta_{x_s}}$	$\alpha_{x_s} = \dfrac{0.13(V_s + 25)}{1 - e^{-\left(\frac{25+V_s}{10}\right)}}$	$\beta_{x_s} = 1.69e^{-0.0125(35+V_s)}$
I_h	$q_\infty = \dfrac{1}{1 + e^{\frac{80+V_s}{4}}}$	$\tau_q = \dfrac{1}{e^{-(14.6+0.086V_s)} + e^{-(1.87+0.0}}$		
I_{Ca_h}	$r_\infty = \dfrac{\alpha_r}{\alpha_r + \beta_r}$	$\tau_r = \dfrac{5}{\alpha_r + \beta_r}$	$\alpha_r = \dfrac{1.7}{1 + e^{-\left(\frac{5+V_d}{13.9}\right)}}$	$\beta_r = \dfrac{0.02(V_d + 8.5)}{e^{\left(\frac{8.5+V_d}{5}\right)} - 1}$
$I_{K_{Ca}}$	$s_\infty = \dfrac{\alpha_s}{\alpha_s + \beta_s}$	$\tau_s = \dfrac{1}{\alpha_s + \beta_s}$	$\alpha_s = min([Ca^{2+}]_0, 0.01)$ $[Ca^{2+}]_0 = 2 \cdot 10^{-5}[Ca^{2+}]$	$\beta_s = 0.015$
I_{Na_a}	$m_{\infty a} = \dfrac{1}{1 + e^{-\left(\frac{30+V_a}{5.5}\right)}}$			
	$h_{\infty a} = \dfrac{1}{1 + e^{-\left(\frac{60+V_a}{5.8}\right)}}$	$\tau_{h_a} = 1.5e^{-\left(\frac{40+V_a}{33}\right)}$		
I_{K_a}	$x_{\infty a} = \dfrac{\alpha_{x_a}}{\alpha_{x_a} + \beta_{x_a}}$	$\tau_{x_a} = \dfrac{1}{\alpha_{x_a} + \beta_{x_a}}$	$\alpha_{x_a} = \dfrac{0.13(V_a + 25)}{1 - e^{-\left(\frac{25+V_a}{10}\right)}}$	$\beta_{x_a} = 1.69e^{-0.0125(35+V_a)}$

Three morphological parameters determine the electrotonic properties of the three-

compartmental model: the ratio of the somatic area to the total somatic and dendritic surface

area p; the ratio of the somatic area to the total somatic and axonic surface area p_2, and the

electrotonic coupling conductance between the 3 compartments. The current flowing from the

somatic into the dendritic compartment I_{sd} and from the dendritic into the somatic

compartments I_{ds}, as well as the leakage currents for the soma I_{ls} and for the dendrites I_{ld} are

represented in this model [87]. We also have the current flowing out from the soma into the

axon I_{sa} and from the axon into the soma I_{as} as well as the leakage for the axonic

compartment I_{la} [107]. Furthermore, the dynamics of the calcium concentration $[Ca^{2+}]$ is

defined as follows:

$$\frac{d[Ca^{2+}]}{dt} = -3.0 I_{Ca_h} - 0.075 [Ca^{2+}] \hspace{2cm} \text{Eq. (12)}$$

Two distinct types of calcium currents exist in the IO neuron: a low-threshold current I_{Ca_l}

located in the soma and a high-threshold current I_{Ca_h} located in the dendrites [77,113]. As the

window of conductance of I_{Ca_l} is around the resting membrane potential, the IO cell is

excited in response to hyperpolarizing currents [77,113]. On the other hand, I_{Ca_h} is non-

inactivating. Thus, a depolarizing dendritic input results in a prolonged plateau potential. As

calcium enters the dendrites, the calcium-activated potassium current I_{KCa} is activated. This

current abruptly terminates the plateau potential after about 30 milliseconds [77,113]. Due to

the long time constant (several hundred milliseconds) of I_{KCa}, there is a long

afterhyperpolarization which, in turn, de-inactivates I_{Ca_l} resulting in a postinhibitory rebound

spike. This current contributes to the subthreshold oscillations as it contributes to amplitude

and frequency [87,114]. Furthermore, somatic sodium spikes are generated with the sodium

current I_{Na}. These are terminated by an outward delayed rectifier potassium current $I_{K_{dr}}$

[77,113]. Finally, the axon hillock includes a sodium current I_{Na_a} (whose inactivation

function was altered to allow for the generation of bursts of spikes) and potassium current I_{K_a}

[107].

 Synapses

        Table 6 shows the parameters and synaptic delays for each synapse in the model. The

 synaptic delay and time constants were chosen from literature for the PC to CN and the CN to

 IO   delays   [99,107,115–118].   Other   parameters   were   selected   to   reproduce   the

 electrophysiological results shown in **Fig. 3**. Synaptic parameters in **Table 6** were held

 constant across simulations to isolate the effects of plasticity mechanisms and network

 architecture. While neuronal parameters in **Tables 2** and **4** were varied to reflect known

 biological heterogeneity, fixed synaptic parameters ensured consistent scaling of synaptic

 inputs, allowing clearer interpretation of plasticity-driven dynamics. When applicable, these

 values are overridden by those in **Table 7** for zone-specific simulations. While the synaptic

 values chosen fall within biologically plausible ranges, we manually verified that the key

 model outcomes —including the frequency preference for CSpk generation and PF-PC

 weight stabilization— were robust to variation in these delay parameters.

1007

Table 6. Baseline synaptic parameters and delays

Synapses	Parameters	Values
PF-PC	Moving threshold θ_{M_0}	60 Hz
	Threshold time constant τ_M	15 ms
	CSpk induced LTD constant A_{CSpk}	-0.01 nA
	CSpk induced LTD time constant τ_{CSpk}	350 ms
PC-CN	Inhibitory current from PC to CN $\gamma^{(CN)}{}_{PC}$	0.004 nA
	Synaptic delay	10 ms
CN-IO	Inhibitory current from CN to IO $\gamma^{(IO)}{}_{CN}$	-1.8 nA
	IO_{CN} time constant $\tau^{(IO)}{}_{CN}$	30 ms
	Synaptic delay	50 ms
IO-IO	Coupling conductance g_c	0.00125 mS/cm^2
IO-PC	Inhibitory current from IO to PC $\gamma^{(PC)}{}_{IO}$	0.22 nA
	$I^{(PC)}{}_{IO}$ time constant $\tau^{(PC)}{}_{IO}$	30 ms
	Synaptic delay	15 ms

1008

PF-PC connectivity

1010   Each PC receives input from 5 PF bundles with different (meta)synaptic weights.

1011 These weights are sampled from the uniform distribution and normalized to a sum of 5. This

1012 resembles a draw from a scaled symmetric Dirichlet distribution with concentration

1013 parameter $\alpha > 1$, i.e. the values are favored to the center of the simplex. The PSC I_{PF} (see

1014 PC model) is calculated as:

1015
$$I_{PF} = \frac{1}{5}\sum_{j}^{N_{PF}} I_j w_j \qquad\qquad \text{Eq. (13)}$$

PF-PC Plasticity

The plasticity mechanism between the PF bundles and PCs consists of an intrinsic

homeostatic mechanism w_{BCM} and a long-term depression (LTD) mechanism w_{CSpk}. The

homeostatic mechanism is based on the Bienenstock-Cooper-Munro-type (BCM) plasticity

mechanism [119]. LTD happens at the IO spike, whereas LTP happens during the absence of

IO spikes. The final weight is as shown in Eq. 14. Where, i is the PC (where $i = 1, \ldots, 100$)

and j is the synapse of that PC (where $j = 1, \ldots, 5$).

1023
$$w_{i,j} = w_{BCM_{i,j}} + w_{CS_{i,j}}$$
     Eq. (14)

BCM

The homeostatic component of synaptic plasticity is governed by a modified BCM

rule. This formulation builds on the original BCM model [119] but introduces additional

damping to match cerebellar PC firing ranges.

Let $\rho_{PF,j}$ denote the effective firing rate (Hz) of the j-th PF bundle, computed from

the amplitude of the OU-driven synaptic current, and let ρ_{PC_i} be the moving average firing

rate (Hz) of the i-th PC over a 200 ms window. The sliding plasticity threshold θ_{M_i} (Hz) for

the i-th PC evolves as shown in Eq. 16.

1032
$$\frac{dw_{BCM_{i,j}}}{dt} = \rho_{PF_j} \phi\left(\rho_{PC_i}, \theta_{M_i}\right)$$
     Eq. (15)

1033
$$\frac{d\theta_{M_i}}{dt} = \left[\left(\rho_{PC_i}\right)^2 - \frac{\theta_{M_i}}{\tau_M}\right]$$
     Eq. (16)

1034 where, τ_M is the threshold time constant. Moreover, ϕ_i represents the general change in

1035 plasticity specific to the i-th PC, as it depends on the firing rate ρ_{PC_i}, and the sliding threshold

1036 θ_{M_i}, both of which vary across PCs. This term is then modulated by ρ_{PF_j} for each PF bundle:

1037
$$\phi_i = \frac{\rho_{PC_i}\left(\rho_{PC_i} - \theta_{M_i}\right)}{\theta_{M_i}}$$
     Eq. (17)

1038   This classical parabolic term (Eq. 17) has zeros at $\rho_{PC} = 0, \theta_M$ and a minimum of $\phi =$

1039 $-\frac{1}{4}\theta_M$ at $\rho_{PC} = \frac{1}{2}\theta_M$. However, left unbounded, this can lead to unrealistically large

1040   potentiation at high firing rates. To limit this, we introduced a damping factors:

$$\frac{dw_{BCM_{i,j}}}{dt} = 5\rho_{PF_j}\, tanh\left(0.01 \frac{\rho_{PC_i}\left(\rho_{PC_i} - \theta_{M_i}\right)}{\theta_{M_i}} \right) \qquad \text{Eq. (18)}$$

where $c = 0.01$ is a shape constant. In implementation (Brian2), we ensure the $tanh$

argument is dimensionless by multiplying the rate term by a time unit (second). This

hyperbolic term flattens the curve at high firing rates, ensuring that potentiation and

depression remain within physiologically plausible bounds.

This formulation yields bidirectional weight changes depending on the position of ρ_{PC}

relative to θ_M, with potentiation when $\rho_{PC} < \theta_M$ and depression when $\rho_{PC} > \theta_M$. The $tanh$

function ensures saturation and boundedness of weight change. The only hard constraint is

imposed on ρ_{PC}, limited to 250 Hz to prevent pathological firing, but no explicit bounds are

placed on w_{BCM_j}.

To assess the robustness of the BCM mechanism, we varied the threshold time

constant, a key parameter controlling the speed at which the PC activity influences synaptic

change. The overall behavior of the model remained robust as the PF-PC weights stabilized

but the Up/Downbound modules diverged in their plasticity outcomes only for a narrow

range of time constant values. Namely, net depression in Downbound modules and net

potentiation in Upbound modules. Outside this range, homeostasis was still achieved, but the

direction of synaptic change was not aligned with the expected biological behavior. These

results suggest that, while the system operates in a resilient homeostatic regime, certain

features of learning dynamics are parameter-sensitive and emerge only within a constrained

space of biologically plausible plasticity kinetics.

LTD vs LTP

LTD is modeled as a negative change of the synaptic weight proportional to the

activity of the PF and with a time decay of 350 ms (**Table 7**) simulating the calcium release

time (**Fig. 2D**). Upon each IO spike, the LTD term was incremented by a constant by a

constant A_{CSpk} is scaled by the absolute deviation from the instantaneous PF input ρ_{PF_j} (Hz)

from its mean level I_{PF_0} (converted to Hz by explicitly multiplying the appropriate unit

scaling factors). As a consequence, LTP occurs in periods without IO spikes.

1068
$$\frac{dw_{CS_{i,j}}}{dt} = \frac{-w_{CS_{i,j}}}{\tau_{CSpk}}$$
Eq. (19)

1069
$$@IO\ spike: w_{CS_{i,j}} += A_{CSpk} \left| \rho_{PF_j} - I_{PF_0} \right|$$
Eq. (20)

1070   As the CSpk-triggered LTD component was incremented by a negative amount and

1071   decays exponentially back toward zero between IO spikes, it always remains negative. This

1072   decay can allow the BCM-driven component $w_{BCM_{i,j}}$ to dominate when its positive

1073   contribution exceeds $w_{CS_{i,j}}$. The total synaptic weight change is the sum of these two

1074   contributions.

1075   **Table 7.** Parameter values for zone-specific simulations of micromodules

Zone	Coupling	Synaptic currents		Conductance	IO OU current parameters	
		CN to IO $\gamma^{(IO)}_{CN}$ (nA)	PC to CN $\gamma^{(CN)}_{PC}$ (nA)	g_{Cal} (mS/cm^2)	$I^{(IO)}_{OU_0}$ (nA)	$\sigma^{(IO)}_{OU}$ (nA)
Upbound	Coupled	-2.0	0.005	1.2	-0.7	0.7
	Uncoupled	-2.0	0.005	1.4	-0.3	0.7
Downbound	Coupled	-1.8	0.004	1.4	-0.3	0.3
	Uncoupled	-1.8	0.004	1.4	-0.6	0.7

1076 <u>PC-CN Synapse</u>

1077   The inhibitory synapse from PC to the CN is modeled by a discrete increase in

1078 $I^{(CN)}{}_{PC}$ at the time of a presynaptic PC spike. After each increase, $I^{(CN)}{}_{PC}$ decays

1079 exponentially back toward zero, resulting in a temporary suppression of CN firing.

1080
$$\frac{dI^{(CN)}{}_{PC}}{dt} = \frac{-I^{(CN)}{}_{PC}}{\tau^{(CN)}{}_{PC}}$$
    Eq. (21)

1081
$$@PC\ spike\colon I^{(CN)}{}_{PC} += \gamma^{(CN)}{}_{PC}$$
    Eq. (22)

1082   Each PC projects to 16 CN cells and each CN cell receives on average from 40 PCs,

1083 ranging from 30 to 52 (**Fig. 1A**). Each PC source indices is repeated for each PC and CN

1084 target indices are sampled without replacement from the CN population.

1085

1086 <u>CN-IO Synapse</u>

1087   Analogously to the PC-CN synapse, the new synaptic current $I^{(IO)}{}_{CN}$ increases with

1088 CN spikes. The decay time of $I^{(IO)}{}_{CN}$ is small (30 ms), and enters the IO at the soma ($I^{(S)}{}_{app}$).

1089 This inhibitory input can be seen as the asynchronous release of GABA onto the IO cell [98].

1090 The update value of $\gamma^{(IO)}{}_{CN}$ is seen in **Table 6** and the synaptic update is done as follows:

1091
$$\frac{dI^{(IO)}{}_{CN}}{dt} = \frac{-I^{(IO)}{}_{CN}}{\tau^{(IO)}{}_{CN}}$$
    Eq. (23)

1092
$$@CN\ spike\colon I^{(IO)}{}_{CN} += \frac{\gamma^{(IO)}{}_{CN}}{N_{CN}}$$
    Eq. (24)

1093   Each CN projects to 10 IO cells and each IO cell receives on average from 10 CN

1094 cells, ranging from 6 to 16 (**Fig. 1A**). The connectivity is generated ensuring these ratios. CN

1095 cell indices are repeated for each CN cell, while IO indices are sampled without replacement

1096 from the IO population, avoiding duplicate connections.

1097

 IO-IO Gap Junction

The electrotonic coupling between the cells is represented by a current I_c (see **Table**

6) in the dendrites and the transjunctional voltage dependence of the gap junction

conductance f [87] is defined below. In the coupled case, each IO neuron connects to all

other neurons in the population (**Fig. 1A**). The transjunctional voltage dependence of the gap

junction conductance (Eq. 26) is defined as $V = V_d - V_{de}$, where both V_d and V_{de} are

dendritic membrane potentials.

1105
$$I_c = g_c(V_d - V_{de})f(V_d - V_{de})$$ Eq. (25)

1106
$$f(V) = 0.6e^{\frac{V^2}{50^2}} + 0.4$$ Eq. (26)

IO-PC Synapse

The modeling of the IO-PC synapse focuses on the pause mechanism following a

CSpk. Each PC receives a CF input from one IO as seen in **Fig. 1A**. As the PC AdEX model

already has an adaptation variable w that increases the interspike interval (ISI), the synaptic

modeling only requires the increase of w following a presynaptic IO spike. The update of the

w adaptation variable is increased as shown in Eq. 27. As w is increased to a higher value, the

PC cannot spike until w decays to a lower value.

1114
$$@IO\ spike\colon w^{PC} \mathrel{+}= \gamma^{(PC)}{}_{IO}$$ Eq. (27)

Only half of the IOs synapse with PCs. Each of these IO CF connects to on average 5

PCs (ranging 2 to 9) and each PC receives from only 1 IO cell (**Fig. 1A**). IO cell indices are

sampled without replacement from the IO population. Each PC is either assigned a unique IO

source, or —when convergence exceeds the IO population size— assigned an IO neuron

randomly selected from a pre-sampled pool, ensuring the intended level of convergence while

avoiding duplicate connections.

Upbound and Downbound zones

To reproduce the physiological results of the different zones we did a grid search over

2 synaptic currents, 1 IO conductance and the IO OU current (see **Table 7**). This grid search

was done in steps of 0.1 for all parameters except for the PC to CN synaptic current $\gamma^{(CN)}_{PC}$

which was done in steps of 0.001. Then we tried to find a set of parameters such that we

could reproduce physiological firing ranges with minimal parameter changes between cases.

Table 7 shows the different parameters that need to be changed to reproduce

electrophysiological results as shown in **Fig. 3**.

The values of **Table 7** override the corresponding baseline parameters listed in **Table**

6 for the indicated zone and coupling condition. These variations reflect biologically

plausible differences in inhibitory strength, dendritic conductance, and IO noise that give rise

to module-specific dynamics. Parameters not listed here retain their default values from

Table 6.

Filtered inputs

Filtered inputs were obtained via bandpass filtering. For each noise input and

frequency range ($f_{c_{low}}$ to $f_{c_{high}}$), a forward 6th butterworth filter was applied on the mean-

offset removed original noise signal, obtaining $y_{filtered}$. Filtering was done as cascaded

second-order sections, taking into account the rate at which the noise was sampled as

$\left(f_s = \frac{1}{dt}\right)$ with dt being the time step of the simulation. To account for power loss due to

filtering, a correction was applied by multiplying the frequency-domain energy:

1141
$$I_{PF_{filtered}} = y_{filtered} \frac{\sqrt{\Sigma|Y_{unfiltered}(f)|^2}}{\sqrt{\Sigma|Y_{filtered}(f)|^2}} + I_{PF_0} \qquad \text{Eq. (28)}$$

1142 *CS-US Experiment*

1143   We used step currents of different amplitude and duration (Table 8) to simulate the

1144 CS and US. The US was copied onto the IO with a 4ms delay and multiplied by 2.8 to have

1145 the desired effect on the IO spike response. The ITI was randomly chosen between 8-12

1146 seconds.

1147
1148
1149
1150   **Table 8.** Parameter values for the different micromodules

Stimulus	Pulse amplitude (nA)	Pulse duration (ms)
CS	1.5	250
US	1.5	30

1151

Simulations

For all simulations the Euler-Maruyama method was used to simulate the frozen

noise. For the rest of the neuron groups, the forward Euler method was used to integrate the

differential equations. Two integration timesteps were used, one for the simulations (0.025

ms) and one for the recording of the neuron output (1 ms). All experiments had a simulation

length of 120 s. We first started by generating a seed which included generating a network

with connectivities, cell parameters and frozen PF input. When modeling the different

micromodules we used the parameters shown in **Table 7**.

The frozen PF input was used for the "no plasticity" experiment. For the subsequent

"plasticity" epoch, we enabled the BCM and LTD mechanisms and applied the same frozen

PF input. At the end of each plasticity epoch, the evolving PF-PC synaptic weights were

smoothed using a Savitzky–Golay filter (window size = 100 samples, polynomial order = 2)

to reduce variability caused by the instantaneous state of the LTD re LTP mechanism.

Without smoothing, the final weight could be biased if the simulation ended immediately

after a CSpk, when the LTD term is transiently active. The final smoothed weight value was

then used as the static weight for the subsequent "after plasticity" simulation with the same

frozen input. This procedure was repeated for the second, third and fourth plasticity epochs

(see **Fig. 2G-L**).

The experiments with filtered signals were done in the same way. The only difference

was that the new PF input was a filtered version of the same frozen input. We still had the

same network seed. We redid the experiments for the following frequency bands: 5-10 Hz,

10-15 Hz, 15-20 Hz, 25-30 Hz, 50-75 Hz, 100-150 Hz and 1-800 Hz (see **Fig. 5A**).

The CS - US experiment (light/puff) was done by adding a step current of 250 ms to

the PF 4 input which represented the CS. The same was done for PF 5 but with a 30 ms step

current input that represented the US. This input was also copied (with a 4ms delay) to the IO

to elicit CSpks. The two signals co-terminated. A block consisted of an initial US followed

by 10 CS-US stimuli with varying ITIs and it finished with a final CS.

For each of the simulations we also did a copy simulation with the uncoupled

scenario. This is the same seed, meaning that we had the same network, same connectivity,

same cell parameters, and same frozen PF input. The only difference was that the coupling

conductance g_c (see Table 3) of the coupling current I_c is lowered to 0 ($g_c = 0$).

Code

All code used for simulations and analysis is available at:

https://github.com/eliasmateo95/CerebellarLoop.

Statistical Analysis

To assess differences across simulation conditions, we used two types of parametric

statistical tests depending on the data distribution. Two-sample Student's *t*-tests were applied

for comparisons of means across groups. F-tests were used to compare variances across

groups. All tests were two-sided and group sized ($n = 40$ for CN and IO cells, $n = 100$ for

PCs) are reported in figure captions.

Acknowledgments

The authors would like to thank Laurens Bosman for his insights and comments on a previous version of the manuscript. We would also like to thank the NVIDIA corporation for supporting our research with the donation of 2 x RTX6000.

Author contributions

EF: Conceptualization, Investigation, Methodology, Writing – original draft, Writing – review & editing. LL: Methodology, Writing – original draft, Writing – review & editing. SH: Funding acquisition, Writing – review. CS: Funding acquisition, Writing – review. CDZ: Conceptualization, Funding acquisition, Investigation, Project administration, Resources, Supervision, Writing – original draft, Writing – review & editing. AB: Conceptualization, Funding acquisition, Investigation, Methodology, Project administration, Resources, Supervision, Writing – original draft, Writing – review & editing. MN: Conceptualization, Funding acquisition, Investigation, Methodology, Project administration, Resources, Supervision, Writing – original draft, Writing – review & editing.

Funding

This work was supported by the Netherlands Organization for Scientific Research (NWO) VIDI/917.18.380,2018/ZonMw (A.B), NWO NWA-ORC 2022 SCANNER (A.B), the Erasmus MC Convergence Health and Technology Integrative Neuromedicine Flagship Program (A.B. and M.N.), the 2021 H2020 funding (n°650003) (M.N). Financial support to CS provided by the European-Union Horizon Europe R&I program through projects SEPTON (no. 101094901) and SECURED (no. 101095717) and through the NWO - Gravitation Programme DBI2 (no. 024.005.022). Financial support to CIDZ was provided by the Netherlands Organization for Scientific Research (NWO-ALW 824.02.001), the Dutch Organization for Medical Sciences (ZonMW 91120067), Medical Neuro-Delta (MD 01092019-31082023), INTENSE LSH-NWO (TTW/00798883), ERC-adv (GA-294775) and ERC-POC (nrs. 737619 and 768914); The NIN Vriendenfonds for Albinism as well as the Dutch NWO Gravitation Program (DBI2).

Conflicts of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

1. Zenke F, Gerstner W, Ganguli S. The temporal paradox of Hebbian learning and homeostatic plasticity. Curr Opin Neurobiol. 2017;43: 166–176.

2. Keck T, Hübener M, Bonhoeffer T. Interactions between synaptic homeostatic mechanisms: an attempt to reconcile BCM theory, synaptic scaling, and changing excitation/inhibition balance. Curr Opin Neurobiol. 2017;43: 87–93.

3. Hölzel RW, Krischer K. Stability and long term behavior of a Hebbian network of Kuramoto oscillators. SIAM J Appl Dyn Syst. 2015;14: 188–201.

4. Fernández Santoro EM, Karim A, Warnaar P, De Zeeuw CI, Badura A, Negrello M. Purkinje cell models: past, present and future. Front Comput Neurosci. 2024;18: 1426653.

5. De Zeeuw CI, Ten Brinke MM. Motor learning and the cerebellum. Cold Spring Harb Perspect Biol. 2015;7: a021683.

6. Heck D, Sultan F. Cerebellar structure and function: making sense of parallel fibers. Hum Mov Sci. 2002;21: 411–421.

7. Chaumont J, Guyon N, Valera AM, Dugué GP, Popa D, Marcaggi P, et al. Clusters of cerebellar Purkinje cells control their afferent climbing fiber discharge. Proc Natl Acad Sci U S A. 2013;110: 16223–16228.

8. De Zeeuw CI, Hoebeek FE, Bosman LWJ, Schonewille M, Witter L, Koekkoek SK. Spatiotemporal firing patterns in the cerebellum. Nat Rev Neurosci. 2011;12: 327–344.

9. Zhou H, Lin Z, Voges K, Ju C, Gao Z, Bosman LWJ, et al. Cerebellar modules operate at different frequencies. Elife. 2014;3: e02536.

10. Sudhakar SK, Hong S, Raikov I, Publio R, Lang C, Close T, et al. Spatiotemporal network coding of physiological mossy fiber inputs by the cerebellar granular layer. PLoS Comput Biol. 2017;13: e1005754.

11. Galliano E, Gao Z, Schonewille M, Todorov B, Simons E, Pop AS, et al. Silencing the majority of cerebellar granule cells uncovers their essential role in motor learning and consolidation. Cell Rep. 2013;3: 1239–1251.

12. Coesmans M, Weber JT, De Zeeuw CI, Hansel C. Bidirectional parallel fiber plasticity in the cerebellum under climbing fiber control. Neuron. 2004;44: 691–700.

13. Wang X, Novello M, Gao Z, Ruigrok TJH, De Zeeuw CI. Input and output organization of the mesodiencephalic junction for cerebro-cerebellar communication. J Neurosci Res. 2022;100: 620–637.

14. Kubo R, Aiba A, Hashimoto K. The anatomical pathway from the mesodiencephalic junction to the inferior olive relays perioral sensory signals to the cerebellum in the mouse. J Physiol. 2018;596: 3775–3791.

15. De Zeeuw CI, Simpson JI, Hoogenraad CC, Galjart N, Koekkoek SK, Ruigrok TJ. Microcircuitry and function of the inferior olive. Trends Neurosci. 1998;21: 391–400.

16. Loyola S, Hoogland TM, Hoedemaker H, Romano V, Negrello M, De Zeeuw CI. How inhibitory and excitatory inputs gate output of the inferior olive. Elife. 2023;12. doi:10.7554/eLife.83239

17. van Hoogstraten WS, Lute MCC, Liu Z, Broersen R, Mangili L, Kros L, et al. Disynaptic

inhibitory cerebellar control over caudal medial accessory Olive. eNeuro. 2024;11:
ENEURO.0262–23.2023.

18. Wang X, Liu Z, Angelov M, Feng Z, Li X, Li A, et al. Excitatory nucleo-olivary pathway shapes
cerebellar outputs for motor control. Nat Neurosci. 2023;26: 1394–1406.

19. Broersen R, De Zeeuw CI. Keeping track of time: An interaction of mossy fibers and climbing
fibers. Neuron. 2024;112: 2664–2666.

20. Gao Z, van Beugen BJ, De Zeeuw CI. Distributed synergistic plasticity and cerebellar learning.
Nat Rev Neurosci. 2012;13: 619–635.

21. Ito M. Mechanisms of motor learning in the cerebellum. Brain Res. 2000;886: 237–245.

22. Gutierrez-Castellanos N, Da Silva-Matos CM, Zhou K, Canto CB, Renner MC, Koene LMC, et
al. Motor Learning Requires Purkinje Cell Synaptic Potentiation through Activation of AMPA-
Receptor Subunit GluA3. Neuron. 2017;93: 409–424.

23. Suvrathan A, Raymond JL. Depressed by learning-heterogeneity of the plasticity rules at parallel
fiber synapses onto Purkinje cells. Cerebellum. 2018;17: 747–755.

24. De Zeeuw CI, Lisberger SG, Raymond JL. Diversity and dynamism in the cerebellum. Nat
Neurosci. 2021;24: 160–167.

25. Zang Y, De Schutter E. Recent data on the cerebellum require new models and theories. Curr
Opin Neurobiol. 2023;82: 102765.

26. Bouvier G, Aljadeff J, Clopath C, Bimbard C, Ranft J, Blot A, et al. Cerebellar learning using
perturbations. Elife. 2018;7: e31599.

27. Titley HK, Kislin M, Simmons DH, Wang SS-H, Hansel C. Complex spike clusters and false-
positive rejection in a cerebellar supervised learning rule. J Physiol. 2019;597: 4387–4406.

28. Lisman J. Long-term potentiation: outstanding questions and attempted synthesis. Long-term
Potentiation: Enhancing Neuroscience for 30 years. Oxford University PressOxford; 2004. pp.
371–392.

29. Allen CB, Celikel T, Feldman DE. Long-term depression induced by sensory deprivation during
cortical map plasticity in vivo. Nat Neurosci. 2003;6: 291–299.

30. Abbott LF, Nelson SB. Synaptic plasticity: taming the beast. Nat Neurosci. 2000;3 Suppl: 1178–
1183.

31. Shimizu G, Yoshida K, Kasai H, Toyoizumi T. Computational roles of intrinsic synaptic
dynamics. bioRxiv. bioRxiv; 2021. doi:10.1101/2021.04.22.441034

32. Sussillo D, Toyoizumi T, Maass W. Self-tuning of neural circuits through short-term synaptic
plasticity. J Neurophysiol. 2007;97: 4079–4095.

33. Canolty RT, Ganguly K, Kennerley SW, Cadieu CF, Koepsell K, Wallis JD, et al. Oscillatory
phase coupling coordinates anatomically dispersed functional cell assemblies. Proc Natl Acad
Sci U S A. 2010;107: 17356–17361.

34. Molter C, Salihoglu U, Bersini H. The road to chaos by time-asymmetric Hebbian learning in
recurrent neural networks. Neural Comput. 2007;19: 80–110.

35. Rabinowitch I, Segev I. Two opposing plasticity mechanisms pulling a single synapse. Trends
Neurosci. 2008;31: 377–383.

36. van Rossum MC, Bi GQ, Turrigiano GG. Stable Hebbian learning from spike timing-dependent
plasticity. J Neurosci. 2000;20: 8812–8821.

37. Voogd J, Ruigrok TJH. The organization of the corticonuclear and olivocerebellar climbing fiber
projections to the rat cerebellar vermis: the congruence of projection zones and the zebrin
pattern. J Neurocytol. 2004;33: 5–21.

38. Long RM, Pakan JMP, Graham DJ, Hurd PL, Gutierrez-Ibañez C, Wylie DR. Modulation of
complex spike activity differs between zebrin-positive and -negative Purkinje cells in the pigeon
cerebellum. J Neurophysiol. 2018;120: 250–262.

39. Voogd J, Glickstein M. The anatomy of the cerebellum. Trends Neurosci. 1998;21: 370–375.

40. Hoang H, Tsutsumi S, Matsuzaki M, Kano M, Kawato M, Kitamura K, et al. Dynamic
organization of cerebellar climbing fiber response and synchrony in multiple functional
components reduces dimensions for reinforcement learning. Elife. 2023;12: e86340.

41. Zhang J, Tran-Anh K, Hirata T, Sugihara I. Striped distribution pattern of Purkinje cells of
different birthdates in the mouse cerebellar cortex studied with the Neurog2-CreER transgenic
line. Neuroscience. 2021;462: 122–140.

42. Chen X, Du Y, Broussard GJ, Kislin M, Yuede CM, Zhang S, et al. Transcriptomic mapping
uncovers Purkinje neuron plasticity driving learning. Nature. 2022;605: 722–727.

43. Voerman S, Urbanus BHA, Schonewille M, White JJ, De Zeeuw CI. Postsynaptic plasticity of
Purkinje cells in mice is determined by molecular identity. Commun Biol. 2022;5: 1328.

44. De Zeeuw CI. Bidirectional learning in upbound and downbound microzones of the cerebellum.
Nat Rev Neurosci. 2021;22: 92–110.

45. Geminiani A, Casellato C, Boele H-J, Pedrocchi A, De Zeeuw CI, D'Angelo E. Mesoscale
simulations predict the role of synergistic cerebellar plasticity during classical eyeblink
conditioning. PLoS Comput Biol. 2024;20: e1011277.

46. De Zeeuw CI, Romano V. Time and tide of cerebellar synchrony. Proc Natl Acad Sci U S A.
2022;119: e2204155119.

47. Wadiche JI, Jahr CE. Patterned expression of Purkinje cell glutamate transporters controls
synaptic plasticity. Nat Neurosci. 2005;8: 1329–1334.

48. Ebner TJ, Wang X, Gao W, Cramer SW, Chen G. Parasagittal zones in the cerebellar cortex
differ in excitability, information processing, and synaptic plasticity. Cerebellum. 2012;11: 418–
419.

49. Suvrathan A, Payne HL, Raymond JL. Timing rules for synaptic plasticity matched to behavioral
function. Neuron. 2016;92: 959–967.

50. de Solages C, Szapiro G, Brunel N, Hakim V, Isope P, Buisseret P, et al. High-frequency
organization and synchrony of activity in the purkinje cell layer of the cerebellum. Neuron.
2008;58: 775–788.

51. Kistler WM. Time-slicing: A model for cerebellar function based on synchronization,
reverberation, and time windows. Neurocomputing. 2001;38-40: 1367–1372.

52. Negrello M, Warnaar P, Romano V, Owens CB, Lindeman S, Iavarone E, et al. Quasiperiodic
rhythms of the inferior olive. PLoS Comput Biol. 2019;15: e1006475.

53. Ruigrok TJ. Cerebellar nuclei: the olivary connection. Prog Brain Res. 1997;114: 167–192.

54. Scheibel ME, Scheibel AB. The inferior olive; a Golgi study. J Comp Neurol. 1955;102: 77–131.

55. Vrieler N, Loyola S, Yarden-Rabinowitz Y, Hoogendorp J, Medvedev N, Hoogland TM, et al.
Variability and directionality of inferior olive neuron dendrites revealed by detailed 3D
characterization of an extensive morphological library. Brain Struct Funct. 2019;224: 1677–
1695.

56. Scheibel ME, Scheibel AB. Neuropil organization in the superior olive of the cat. Exp Neurol.
1974;43: 339–348.

57. Brodal A, Scheibel A, Scheibel M, Walberg F. Areal distribution of axonal and dendritic patterns
in inferior olive. J Comp Neurol. 1956;106: 21–49.

58. Cerebellar dentate nucleus: Organization, cytology and transmitters, by Victoria Chan Palay,
550 pp, illustrated, $98.00, Springer, Berlin and New York, 1977. Ann Neurol. 1978;4: 588–588.

59. Schmolesky MT, Weber JT, De Zeeuw CI, Hansel C. The making of a complex spike: ionic
composition and plasticity. Ann N Y Acad Sci. 2002;978: 359–390.

60. Busch SE, Hansel C. Climbing fiber multi-innervation of mouse Purkinje dendrites with
arborization common to human. Science. 2023;381: 420–427.

61. Ju C, Bosman LWJ, Hoogland TM, Velauthapillai A, Murugesan P, Warnaar P, et al. Neurons of
the inferior olive respond to broad classes of sensory input while subject to homeostatic control.
J Physiol. 2019;597: 2483–2514.

62. Devor A, Yarom Y. Coherence of subthreshold activity in coupled inferior olivary neurons. Ann
N Y Acad Sci. 2002;978: 508.

63. Jacobson GA, Lev I, Yarom Y, Cohen D. Invariant phase structure of olivo-cerebellar
oscillations and its putative role in temporal pattern generation. Proc Natl Acad Sci U S A.
2009;106: 3579–3584.

64. Lefler Y, Torben-Nielsen B, Yarom Y. Oscillatory activity, phase differences, and phase
resetting in the inferior olivary nucleus. Front Syst Neurosci. 2013;7: 22.

65. Torben-Nielsen B, Segev I, Yarom Y. The generation of phase differences and frequency
changes in a network model of inferior olive subthreshold oscillations. PLoS Comput Biol.
2012;8: e1002580.

66. Lanore F, Cayco-Gajic NA, Gurnani H, Coyle D, Silver RA. Cerebellar granule cell axons
support high-dimensional representations. Nat Neurosci. 2021;24: 1142–1150.

67. Badura A, De Zeeuw CI. Cerebellar granule cells: Dense, rich and evolving representations. Curr
Biol. 2017;27: R415–R418.

68. Uhlenbeck GE, Ornstein LS. On the theory of the Brownian motion. Phys Rev. 1930;36: 823–
841.

69. Giovannucci A, Badura A, Deverett B, Najafi F, Pereira TD, Gao Z, et al. Cerebellar granule
cells acquire a widespread predictive feedback signal during motor learning. Nat Neurosci.
2017;20: 727–734.

70. Lee J-H, Khan MM, Stark AP, Seo S, Norton A, Yao Z, et al. Cerebellar granule cell signaling is
indispensable for normal motor performance. Cell Rep. 2023;42: 112429.

71. Sultan F. Distribution of mossy fibre rosettes in the cerebellum of cat and mice: evidence for a
parasagittal organization at the single fibre level. Eur J Neurosci. 2001;13: 2123–2130.

72. Isope P, Barbour B. Properties of unitary granule cell-->Purkinje cell synapses in adult rat
cerebellar slices. J Neurosci. 2002;22: 9668–9678.

73. Abbasi S, Hudson AE, Maran SK, Cao Y, Abbasi A, Heck DH, et al. Robust transmission of rate
coding in the inhibitory Purkinje cell to cerebellar nuclei pathway in awake mice. PLoS Comput
Biol. 2017;13: e1005578.

74. Najac M, Raman IM. Integration of Purkinje cell inhibition by cerebellar nucleo-olivary neurons.
J Neurosci. 2015;35: 544–549.

75. Van Der Giessen RS, Koekkoek SK, van Dorp S, De Gruijl JR, Cupido A, Khosrovani S, et al.
Role of olivary electrical coupling in cerebellar motor learning. Neuron. 2008;58: 599–612.

76. Kølvraa M, Müller FC, Jahnsen H, Rekling JC. Mechanisms contributing to cluster formation in
the inferior olivary nucleus in brainstem slices from postnatal mice. J Physiol. 2014;592: 33–47.

77. Llinás R, Yarom Y. Properties and distribution of ionic conductances generating
electroresponsiveness of mammalian inferior olivary neurones in vitro. J Physiol. 1981;315:
569–584.

78. Leznik E, Llinás R. Role of gap junctions in synchronized neuronal oscillations in the inferior
olive. J Neurophysiol. 2005;94: 2447–2456.

79. Bazzigaluppi P, Ruigrok T, Saisan P, De Zeeuw CI, de Jeu M. Properties of the nucleo-olivary
pathway: an in vivo whole-cell patch clamp study. PLoS One. 2012;7: e46360.

80. Khosrovani S, Van Der Giessen RS, De Zeeuw CI, De Jeu MTG. In vivo mouse inferior olive
neurons exhibit heterogeneous subthreshold oscillations and spiking patterns. Proc Natl Acad Sci
U S A. 2007;104: 15911–15916.

81. ten Brinke MM, Boele H-J, Spanke JK, Potters J-W, Kornysheva K, Wulff P, et al. Evolving
models of Pavlovian conditioning: Cerebellar cortical dynamics in awake behaving mice. Cell
Rep. 2015;13: 1977–1988.

82. Cook AA, Fields E, Watt AJ. Losing the Beat: Contribution of Purkinje Cell Firing Dysfunction
to Disease, and Its Reversal. Neuroscience. 2021;462: 247–261.

83. Witter L, Canto CB, Hoogland TM, de Gruijl JR, De Zeeuw CI. Strength and timing of motor
responses mediated by rebound firing in the cerebellar nuclei after Purkinje cell activation. Front
Neural Circuits. 2013;7: 133.

84. Hoebeek FE, Witter L, Ruigrok TJH, De Zeeuw CI. Differential olivo-cerebellar cortical control
of rebound activity in the cerebellar nuclei. Proc Natl Acad Sci U S A. 2010;107: 8410–8415.

85. Bina L, Romano V, Hoogland TM, Bosman LWJ, De Zeeuw CI. Purkinje cells translate
subjective salience into readiness to act and choice performance. Cell Rep. 2021;37: 110116.

86. Romano V, Reddington AL, Cazzanelli S, Mazza R, Ma Y, Strydis C, et al. Functional
convergence of autonomic and sensorimotor processing in the lateral cerebellum. Cell Rep.
2020;32: 107867.

87. Schweighofer N, Doya K, Kawato M. Electrophysiological properties of inferior olive neurons:
A compartmental model. J Neurophysiol. 1999;82: 804–817.

88. Dijkhuizen S, Van Ginneken LMC, IJpelaar AHC, Koekkoek SKE, De Zeeuw CI, Boele HJ.
Impact of enriched environment on motor performance and learning in mice. Sci Rep. 2024;14:
5962.

89. Wulff P, Schonewille M, Renzi M, Viltono L, Sassoe-Pognetto M, Badura A, et al. Synaptic
inhibition of Purkinje cells mediates consolidation of vestibulo-cerebellar motor learning. Nat
Neurosci. 2009;12: 1042–1049.

90. Badura A, Schonewille M, Voges K, Galliano E, Renier N, Gao Z, et al. Climbing fiber input
shapes reciprocity of Purkinje cell firing. Neuron. 2013;78: 700–713.

91. Oyaga MR, Serra I, Kurup D, Koekkoek SKE, Badura A. Delay eyeblink conditioning
performance and brain-wide c-Fos expression in male and female mice. Open Biol. 2023;13:
220121.

92. Oja E. A simplified neuron model as a principal component analyzer. J Math Biol. 1982;15:
267–273.

93. Brito CSN, Gerstner W. Nonlinear Hebbian learning as a unifying principle in receptive field
formation. PLoS Comput Biol. 2016;12: e1005070.

94. Voerman S, Broersen R, Swagemakers SMA, De Zeeuw CI, van der Spek PJ. Plasticity
mechanisms of genetically distinct Purkinje cells. Bioessays. 2024;46: e2400008.

95. Rodriques SG, Stickels RR, Goeva A, Martin CA, Murray E, Vanderburg CR, et al. Slide-seq: A
scalable technology for measuring genome-wide expression at high spatial resolution. Science.
2019;363: 1463–1467.

96. Hao S, Zhu X, Huang Z, Yang Q, Liu H, Wu Y, et al. Cross-species single-cell spatial
transcriptomic atlases of the cerebellar cortex. Science. 2024;385: eado3927.

97. Solinas S, Nieus T, D'Angelo E. A realistic large-scale model of the cerebellum granular layer
predicts circuit spatio-temporal filtering properties. Front Cell Neurosci. 2010;4: 12.

98. Best AR, Regehr WG. Inhibitory regulation of electrically coupled neurons in the inferior olive
is mediated by asynchronous release of GABA. Neuron. 2009;62: 555–565.

99. Andersson G, Hesslow G. Activity of Purkinje cells and interpositus neurones during and after
periods of high frequency climbing fibre activation in the cat. Exp Brain Res. 1987;67: 533–542.

100.   Lefler Y, Yarom Y, Uusisaari MY. Cerebellar inhibitory input to the inferior olive decreases
electrical coupling and blocks subthreshold oscillations. Neuron. 2014;81: 1389–1400.

101.   Llinas R, Baker R, Sotelo C. Electrotonic coupling between neurons in cat inferior olive. J
Neurophysiol. 1974;37: 560–571.

102.   Sotelo C, Llinas R, Baker R. Structural study of inferior olivary nucleus of the cat:
morphological correlates of electrotonic coupling. J Neurophysiol. 1974;37: 541–559.

103.   de Zeeuw CI, Holstege JC, Ruigrok TJ, Voogd J. Ultrastructural study of the GABAergic,
cerebellar, and mesodiencephalic innervation of the cat medial accessory olive: anterograde
tracing combined with immunocytochemistry. J Comp Neurol. 1989;284: 12–35.

104.   De Gruijl JR, Sokol PA, Negrello M, De Zeeuw CI. Modulation of electrotonic coupling in
the inferior Olive by inhibitory and excitatory inputs: Integration in the glomerulus. bioRxiv.
bioRxiv; 2016. doi:10.1101/072041

105.   Bauer S, van Wingerden N, Jacobs T, van der Horst A, Zhai P, Betting J-HLF, et al. Purkinje
cell activity resonation generates rhythmic behaviors at the preferred frequency of 8 Hz.
Biomedicines. 2022;10: 1831.

106.   Hong S, De Schutter E. Purkinje neurons: What is the signal for complex spikes? Curr Biol.

2008;18: R969–71.

107.   De Gruijl JR, Bazzigaluppi P, de Jeu MTG, De Zeeuw CI. Climbing fiber burst size and
olivary sub-threshold oscillations in a network setting. PLoS Comput Biol. 2012;8: e1002814.

108.   De Zeeuw CI, Lang EJ, Sugihara I, Ruigrok TJ, Eisenman LM, Mugnaini E, et al.
Morphological correlates of bilateral synchrony in the rat cerebellar cortex. J Neurosci. 1996;16:
3412–3426.

109.   Chen X, Kovalchuk Y, Adelsberger H, Henning HA, Sausbier M, Wietzorrek G, et al.
Disruption of the olivo-cerebellar circuit by Purkinje neuron-specific ablation of BK channels.
Proc Natl Acad Sci U S A. 2010;107: 12323–12328.

110.   Brette R, Gerstner W. Adaptive exponential integrate-and-fire model as an effective
description of neuronal activity. J Neurophysiol. 2005;94: 3637–3642.

111.   Goodman D, Brette R. Brian: a simulator for spiking neural networks in python. Front
Neuroinform. 2008;2: 5.

112.   Abbott LF, Dayan P. Theoretical neuroscience. In: MIT Press [Internet]. The MIT Press,
Massachusetts Institute of Technology; 1 Dec 2021 [cited 2 Sep 2024]. Available:
https://mitpress.mit.edu/9780262041997/theoretical-neuroscience/

113.   Llinás R, Yarom Y. Electrophysiology of mammalian inferior olivary neurones in vitro.
Different types of voltage-dependent ionic conductances. J Physiol. 1981;315: 549–567.

114.   Bal T, McCormick DA. Synchronized oscillations in the inferior olive are controlled by the
hyperpolarization-activated cation current I(h). J Neurophysiol. 1997;77: 3145–3156.

115.   Ruigrok TJH, Voogd J. Cerebellar influence on olivary excitability in the cat. Eur J Neurosci.
1995;7: 679–693.

116.   Hesslow G. Inhibition of inferior olivary transmission by mesencephalic stimulation in the
cat. Neurosci Lett. 1986;63: 76–80.

117.   Andersson G, Hesslow G. Inferior olive excitability after high frequency climbing fibre
activation in the cat. Exp Brain Res. 1987;67: 523–532.

118.   Andersson G, Oscarsson O. Projections to lateral vestibular nucleus from cerebellar climbing
fiber zones. Exp Brain Res. 1978;32: 549–564.

119.   Bienenstock EL, Cooper LN, Munro PW. Theory for the development of neuron selectivity:
orientation specificity and binocular interaction in visual cortex. J Neurosci. 1982;2: 32–48.

120.   McInnes L, Healy J, Astels S. hdbscan: Hierarchical density based clustering. J Open Source
Softw. 2017;2: 205.

1501

Supporting Information

Supporting Information Figure S1. *Gap junctions increase spiking and subthreshold synchrony (A) Gap junctions are disabled by setting the gap junction current = 0. For the Downbound micromodule, we compare the IO membrane response (B), raster (C), and correlation matrix of membrane potentials (D) in coupled (top) and uncoupled networks (bottom). Synchrony directly impacts the correlations in weight changes expected at the PF-PC synapse.*

1510

Supporting Information Figure S2-1. ***Synchrony and Clustering.*** *In a single micromodule*
*we observe a few clusters in the short time scale whose stability fades exponentially. (**A**)*
*Kuramoto synchrony metric for STOs is derived from IO Vm in 5 transformation steps. (**B**)*
Stability of global oscillator (i.e. usability as a low dimensional global clock): global phase
(mean Hilbert phase of all IO cells) from start to end of an 80 second period. The deviation
from horizontal indicates phase drifts across phase in radians. Both Upbound and
Downbound IO have small drifts and maintain a robust global phase, equating with a single
*synchronized cluster. (**C**) Delta phase plot. Difference of phase of each IO neuron to the*
global phase over time. Plotted modulo 32pi radians to preserve space. Phase-locked
oscillators are clearly visible as overlapping lines, while lag/leap events are shown as
*vertical jumps. Comparing long (**D,** 30s) and short term (**E,** 1s) clusterization with the*
pairwise kuramoto phase reveals a few clusters but a single global oscillator in the 30s
*frame. (**F**) Voltage trace of the clusters found in **E**. The black bars indicate the 1s used for*
*clustering. (**G**) Cluster stability of 1s clusters. In intervals of 10ms, 1s clusters were found*
with HDBSCAN [120] and compared with each other via NMI (normalized mutual
information). At a distance of 1s, clusters are barely related.

1527
Supporting Information Figure S2-2. ***Uncoupled Synchrony.*** *When uncoupled, the global*
oscillator still exhibits stable clock behavior. In Upbound zones there is less phase
dispersion, but no phase locking appears in uncoupled scenarios. Downbound zone
*oscillators act completely independently. (**A**) IO Spike triggered Kuramoto synchrony with*
average 25/75 percentiles in Downbound, after plasticity. Average STO synchrony is higher
for the coupled case. STO synchrony peaks before and after IO spikes in the coupled case,
but not for the uncoupled case. This indicates that gap junctions not only set a base level of
synchrony, but they also induce synchronous spikes. Note the increase of synchrony before
*the IO spike in the coupled case. (**B**) Stability of the global oscillator (see S2.1B), for the*
*uncoupled case. Mean oscillator phase has smaller variation. (**C**) Delta phase plot (see*
S2.1C) for the uncoupled case (without modulo because phase locking does not occur as in
the coupled case (Figure S2-1). IO neurons in the Upbound zone still keep some synchrony
while for Downbound zones, oscillators behave independently.
1541
1542

Supporting Information Figure S3 Difference between coupled and uncoupled scenarios.
*(**A**) SSpks frequency is significantly different across zones (coupled: **t=13.9, p<0.001**, uncoupled: **t=13.61, p<0.001; two-sample Student's t-test; n = 100**). (**B**) CSpks frequency (coupled: **t=3.98, p<0.001**, uncoupled: **t=2.5, p=0.014; two-sample Student's t-test; n = 40**). The variance of CSpks is significantly different in the uncoupled case across zones(**F=8.65, p<0.001; F-test using CCDF of F-distribution; n = 40**). (**C**) The CN firing frequency (coupled: **t=-3.26, p=0.002**, uncoupled: **t=-2.98, p=0.004; two-sample Student's t-test; n = 40**). (**D**) The length of the CF Pause is significantly different across zones for both coupling cases (coupled: **t=-12.52, p<0.001**, uncoupled: **t=-14.39, p=0.014; two-sample Student's t-test; n = 40**). (**E**) SSpk CV is significantly different for the Downbound zone across coupling cases (**t=5.94, p<0.001; two-sample Student's t-test; n = 100**). For the coupled case, across both zones, the SSpk CV is significantly different (**t=-4.27, p<0.001; two-sample Student's t-test; n = 100**). (**F**) SSpk CV2 is significantly different for both coupled and uncoupled cas, across both zones, (Coupled**: t=-13.06, p<0.001**, uncoupled: **t=-12.69, p<0.001; two-sample Student's t-test; n = 100**). (**G**) CSpk CV is significantly different for the Downbound zone across coupling cases (**t=-3.69, p<0.001; two-sample Student's t-test; n = 100**. For the coupled case, across both zones, the CSpk CV is significantly different (**t=7.16, p<0.001; two-sample Student's t-test; n = 100**). (**H**) CSpk CV2 is significantly different for the coupled case across both zones, (**t=-3.57, p<0.001; two-sample Student's t-test; n = 100**). For the uncoupled case, the variance is significantly different across both zones (**F=6.86, p<0.001; F-test using CCDF of F-distribution; n = 100**).*

Supporting Information Figure S4 More CN inhibition leads to less STO frequency and to more CSpk frequency. (A) *After Plasticity difference for SSpks is significant across coupled cases for the Upbound zone (**t=4.51, p<0.001; two-sample Student's t-test; n=100**) and for coupled case across zones (**t=-5.36, p<0.001; two-sample Student's t-test; n=100**). The variance of SSpk frequency is significantly different for the coupled case across both zones (**F=4.85, p<0.001; F-test using CCDF of F-distribution; n=100**). **(B)** CSpk frequency is significantly different across coupling cases (coupled: **t=9.44, p<0.001**, uncoupled: **t=5.84, p<0.001; two-sample Student's t-test; n=40**) and the variance is also significantly different (coupled: **F=2.91, p<0.001**, uncoupled: **F=12.19, p<0.001; F-test using CCDF of F-distribution; n=40**). **(C)** CN firing frequency is significantly different across coupling cases (coupled: **t=9.15, p<0.001**, uncoupled: **t=8.87, p<0.001; two-sample Student's t-test; n=40**). **(D)** The length of the CF pause is significantly different across coupling cases for Upbound and Downbound zones (Upbound: **t=-6.25, p<0.001**, Downbound: **t=2.5, p=0.013; two-sample Student's t-test; n=40**). It is also significantly different across zones for the uncoupled case (**t=-7.76, p<0.001; two-sample Student's t-test; n=40**). (E) Increase in CN-to-IO inhibitory synapses,, and thus inhibition strength, led to a decrease in IO STO frequency and a concurrent increase in CSpk frequency.*

1585
Supporting Information Figure S5. ***Input Filter 5-10Hz.*** *(**A**) Firing frequency distributions*
*for each population before and after plasticity for different frequency bands. (**B**) STO*
frequency vs IO spike frequency for Upbound (left) and Downbound (right) zones for before
(top) and after (bottom) plasticity. There seems to be an inverse relationship between STO
frequency and IO spike frequency. To reduce the effect of the spike refractory period on STO
frequency, STO frequency is here defined as the mean of the 25-75 percentile of the inverse of
time differences between zero crossings in the Hilbert phase. For the 5-10 Hz filter in the
Upbound zone, STO frequency seems to decrease while STO spikes increase. For the
Downbound zone, STO frequency is lowered after plasticity, correlated to a higher baseline
*CSpk frequency and thus a less pronounced difference to the 5-10 Hz filter. (**C**) IO spike*
triggered PSC for NF (top), 5-10 (middle) and 10-15 Hz (bottom) input filters. In general, IO
spikes for The NF case seem to derive from entrainment followed by a final larger decrease
in PSC. For 5-10 Hz filter, IO spikes seem to derive purely from entrainment to the
oscillatory signal, with also a much clearer (larger amplitude) template before the spike.
1600

1601
Supporting Information Figure S6-1. ***CS-US experiment.*** *(**A**) Average weight shift after*
*plasticity for each parallel fiber. The CS pf sees a large decrease in weight. (**B**) Weight*
changes during plasticity, averages for each CS/US trial. PF4 (CS) weight decreases
continuously during CS. PFs 1,2,3 (noise) decrease during US and return quickly to baseline.
*PF5(US) responds the least to US. (**C**) Mean instantaneous Hilbert frequency of STOs,*
smoothed using 51ms 3rd order Savitzky–Golay filter.
1608
1609

1610

***Supporting Information Figure S6-2**. **Uncoupled CS-US experiment.** (**A**) Kuramoto synchrony for the uncoupled scenario (**B**) CS-triggered frequency density responses of the populations for SSpk (top) and CSpk (bottom). (**C**) SSpk frequency response after plasticity is significantly different across zones for both coupling cases (coupled: **t=-8.72, p<0.001**, uncoupled: **t=-3.02, p=0.003; two-sample Student's t-test**), as well as the variance (coupled: **F=5.87, p<0.001**, uncoupled: **F=1.94, p<0.001; F-test using CCDF of F-distribution**). (**D**) CSpk frequency after plasticity is significantly different across coupling cases in the Downbound zone (**t=-2.28, p=0.026; two-sample Student's t-test**). It is also significantly different across zones for both coupling cases (coupled: **t=7.39, p<0.001**, uncoupled: **t=-5.51, p<0.001; two-sample Student's t-test**), as well as the variance (coupled: **F=1.83, p<0.031**, uncoupled: **F=10.4, p<0.001; F-test using CCDF of F-distribution**).*

Supporting Information Table S1 - PC SSpk frequency statistics (n=100)

	Difference compared to NF Upbound		Difference compared to NF Downbound		Across Upbound and Downbound
	two-sample Student's t-test	two-sample ks-test	two-sample Student's t-test	two-sample ks-test	two-sample ks-test
NF	-	-	-	-	*D=0.48, p<0.001*
5-10	*t=8.18, p<0.001*	*D=0.62, p<0.001*	*t=10.29, p<0.001*	*D=0.72, p<0.001*	*D=0.54, p<0.001*
10-15	*t=11.98, p<0.001*	*D=0.74, p<0.001*	*t=11.81, p<0.001*	*D=0.68, p<0.001*	*D=0.70, p<0.001*
25-30	*t=13.7, p<0.001*	*D=0.77, p<0.001*	*p>0.05*	*p>0.05*	*D=0.55, p<0.001*
50-75	*t=12.56, p<0.001*	*D=0.70, p<0.001*	*p>0.05*	*p>0.05*	*D=0.3, p<0.001*
100-150	*t=11.81, p<0.001*	*D=0.74, p<0.001*	*t=6.5, p<0.001*	*D=0.46, p<0.001*	*D=0.38, p<0.001*
1-800	*t=10.2, p<0.001*	*D=0.67, p<0.001*	*t=5.66, p<0.001*	*D=0.32, p<0.001*	*D=0.21, p=0.024*

1626
1627
1628

Supporting Information Table S2 - CN and IO spike frequency statistics across Upbound and Downbound modules

	CN *(n=40)*	IO *(n=40)*
	two-sample ks-test	two-sample ks-test
NF	*D=0.78, p<0.001*	*D=0.78, p<0.001*
5-10	*D=0.80, p<0.001*	*D=0.53, p<0.001*
10-15	*D=0.78, p<0.001*	*D=0.78, p<0.001*
25-30	*D=0.72, p<0.001*	*D=0.82, p<0.001*
50-75	*D=0.72, p<0.001*	*D=0.80, p<0.001*
100-150	*D=0.75, p<0.001*	*D=0.82, p<0.001*
1-800	*D=0.75, p<0.001*	*D=0.70, p<0.001*

1629
1630
1631
1632

A Olivocerebellar Model

5 PF

100 PC

CF-PC   P: 1:5   PC-CN   P: 1:16
        C: 1:1   C: 40:1

40 CN

CN-IO   P: 1:10
        C: 10:1

40 IO

Gap=F(DeltaV)

B Population Responses

1nA

10mV

200ms

10mV

200ms

10mV

200ms

C PF Autocorrelation

I₁

D Membrane Potential

PC

CN

IO

1.0

0.8

0.2

0.0

A BCM Mechanism

Potentiation

θ (Hz)

θ_M^1 θ_M^2

$\overline{\rho_{PC}}$

Depression

B Upbound   Downbound

CS Triggered BCM Response

CSpk   CSpk

— θ_M
— Recent rate
— Depression
— Potentiation

5Hz

200ms

C IO Membrane Potential

Action Potential
Threshold

20mV

1s

D CSpk component

0.1w

1s

E BCM component

0.1w

1s

F Total weight

0.1w

1s

Synaptic weights for each plasticity epoch (PF1)

G   **H**

Relative difference with previous epoch (PF1)

I   **J**

NP AP1 AP2 AP3 AP4   NP AP1 AP2 AP3 AP4

Absolute difference with previous epoch (all weights)

K   **L**

AP1 AP2 AP3 AP4   AP1 AP2 AP3 AP4

A Upbound   Downbound

Recordings

PC Simple spikes (Hz)

Spontaneous activity   Learning   Spontaneous activity   Learning

De Zeeuw 2021

100ms

B Model

PC Simple spikes (Hz)

No Plasticity   Plasticity   No Plasticity   Plasticity

20s

Recordings   Model

C SSpk Firing Frequency (Hz)

D CSpk Firing Frequency (Hz)

E CF Pause (ms)

Upbound   Downbound   Upbound   Downbound

Zhou 2014

A CSpk Triggered Responses

Upbound   Downbound

B CN Triggered IO Responses

Upbound   Downbound

C SSpk Frequency (Hz)   **D** CN Frequency (Hz)

E CSpk Frequency (Hz)   **F** CF Pause (ms)

Upbound   Downbound

A CS-US Experiment

PF 1
PF 2
PF 3
PF 4
PF 5

PC

CS

US

IO

4ms delay

100ms   1nA

B CS-US Block

Initial   ITI   8 stimuli   ITI

US   CS+US   CS+US   CS

C US-Triggered IO Responses

Upbound   Downbound

CS   US   CS   US   40mV

Before Plasticity

After Plasticity

250ms

D Kuramoto Synchrony
([0,1])

— Before Plasticity
-- After Plasticity

0.8   0.8

0.4   0.4

250ms

E Frequency Density Responses

CS   US

SpkS Frequency Density

0.6

0.4

CSpk Frequency Density

0.25

— Upbound
— Downbound
— Before Plasticity
-- After Plasticity

100ms

F SSpk Frequency (Hz)

Upbound

Before   After
Plasticity   Plasticity

Downbound

Before   After
Plasticity   Plasticity

G CSpk Frequency (Hz)

Upbound

Before   After
Plasticity   Plasticity

Downbound

Before   After
Plasticity   Plasticity

H CF Pause (ms)

Upbound

Before   After
Plasticity   Plasticity

Downbound

Before   After
Plasticity   Plasticity

A IO Coupling

B Membrane Potential

Coupled

Uncoupled

20mV

200ms

20mV

200ms

Gap=0

C Raster

200ms

200ms

D Membrane Potential

A Kuramoto Synchrony

1) IO Membrane Potential
   25mV
   200ms

2) Band Pass 4-10 Hz 6th Order Butterworth
   20mV
   200ms

3) Hilbert Analytical Phase
   2
   200ms

4) |Mean of Complex Phase|
   1
   200ms

5) Low pass 2Hz 6th Order Butterworth
   1
   200ms

B Stability of Global Oscillator

Upbound
Downbound

0.02rad
10s

C Delta Phase

Upbound   Downbound

10π   10π
10s   10s

D Pairwise Kuramoto Phase (30s timeframe)

Upbound

Downbound

NP   AP1   AP2   AP3   AP4

Kuramoto synchrony 1.0 – 0.0

E Pairwise Kuramoto Phase (1s timeframe)

AP1, Clusters = 4

Kuramoto synchrony 1.0 – 0.0

F IO Membrane Potential in Clusters

1s window

Cluster
1 2
3 4

50mV
1s

G Cluster Stability

Upbound
Downbound

200ms

A IO Spike-triggered Kuramoto Synchrony

Downbound, After Plasticity

Coupled

IO Spike

Uncoupled

IO Spike

0.4

1s

B Stability of Global Oscillator

Upbound
Downbound

0.02 rad

10s

C Delta Phase

Upbound

Downbound

50π

10s

50π

10s

A Populations Firing Frequency (Hz)

B STO Frequency vs IO Spike Frequency

C IO Spike-triggered PSC

A Weight Shift After Plasticity

B Weight Change During Plasticity

PF 5 US
PF 2
PF 3
PF 1

PF 4 CS

100ms

C Mean Hilbert STO Frequency (Hz)

Before Plasticity

After Plasticity

200ms

200ms

A Kuramoto Synchrony
([0,1])

Upbound

Downbound

CS US

CS US

— Before Plasticity
--- After Plasticity

0.8

0.8

0.4

0.4

250ms

B Frequency Density Responses

CS US

SpkS Frequency Density

0.6

Upbound
Downbound
Before Plasticity
After Plasticity

CSpk Frequency Densityy

1.0

0.5

100ms

C SSpk Frequency (Hz)

D CSpk Frequency (Hz)

Upbound

Downbound

Coupled Uncoupled

Coupled Uncoupled