Assessing the Evolutionary Trajectory of Arbuscular Mycorrhizal Conserved Genes in Seagrasses and Aquatic Close Relatives

Cassandra L. Ettinger[1,#], Jennifer Arroyo[2,3], Jason E. Stajich[1,4]

[1] Department of Microbiology and Plant Pathology, University of California, Riverside, CA, United States
[2] Department of Biology, California State University Channel Islands, Camarillo, CA, United States
[3] The University of Arizona, Tucson, AZ, United States
[4] Institute for Integrative Genome Biology, University of California, Riverside, Riverside, CA, United States

[#]**Corresponding Author:**
Cassandra L. Ettinger
Email address: cassande@ucr.edu

Keywords: seagrasses, marine plants, aquatic plants, arbuscular mycorrhizal fungi, symbiosis, comparative genomics, plant evolution, Alismatales

Summary

Arbuscular mycorrhizal fungi (AMF) form beneficial associations with plants, and are thought to have been critical to the adaptation of the ancestor of terrestrial plants during the transition onto land. However, the ability of AMF to associate with aquatic plants is unclear. To address this, we used 65 publicly available genomes and transcriptomes (25 freshwater, 23 terrestrial and 17 marine plants) to interrogate the genomic potential to form AMF associations in aquatic plant lineages in the order Alismatales. We explored the presence or absence of homologs of 45 genes, with a a special focus on six critical genes including three that co-evolved with AMF associations (*RAD1*, *STR1*, *STR2*) and three necessary for intracellular symbiosis (*SymRK*, *CCaMK/DMI3*, *CYCLOPS/IDP3*). Our results indicate a pattern likely consistent with independent gene losses (or extreme divergence) of symbiosis genes across aquatic lineages suggesting a possible inability to form AMF associations. However, some of these conserved genes (i.e., *CCaMK/DMI3*) are purported to function in other types of fungal symbioses, such as ectomycorrhizal symbiosis, and were observed here in a subset of aquatic lineages, including seagrasses. Overall, our findings highlight the complex evolutionary trajectories of symbiosis-related genes in aquatic plants, suggesting that while AMF associations may have been lost in certain lineages, others have genes that may allow them to form alternative fungal symbioses which may still play an underappreciated role in their ecology.

Introduction

Arbuscular mycorrhizal fungi (AMF) are obligate mutualistic root symbionts that assist host plants in nutrient uptake in exchange for carbon compounds (Parniske, 2008; Bonfante & Anca, 2009). These fungi are thought to have been essential in facilitating the transition of ancestral plants from an aquatic to terrestrial environment (Redecker *et al.*, 2000; Heckman *et al.*, 2001; Wang *et al.*, 2010; Humphreys *et al.*, 2010). These partnerships are widespread, with the majority of flowering plants (82-85%) maintaining AMF associations, underscoring their ecological importance (Wang & Qiu, 2006; Brundrett, 2009).

AMF were historically thought not to colonize aquatic environments, but have since been observed in wetlands, estuaries, mangrove forests and freshwater ecosystems (Beck-Nielsen & Madsen, 2001; Bohrer *et al.*, 2004; Šraj-Kržič *et al.*, 2006; Kohout *et al.*, 2012). AMF have been reported to form associations with freshwater Alismatales, an order of flowering plants that includes many aquatic lineages, although these associations appear sporadic (Clayton & Bagyaraj, 1984; Wigand & Stevenson, 1994; Khan & Belik, 1995; Beck-Nielsen & Madsen, 2001; Wang & Qiu, 2006; Šraj-Kržič *et al.*, 2006; Radhika & Rodrigues, 2007; Brundrett, 2009). Notably, marine Alismatales (i.e., seagrasses) have not been observed to form AMF associations, which may be due to the additional selective pressures of the marine environment (Nielsen *et al.*, 1999). Instead there have been reports of ectomycorrhizal-like fungal root symbioses in Mediterranean seagrasses, *Posidonia oceanica* and *Thalassodendron ciliatum* (Vohník et al., 2015, 2019; Borovec & Vohník, 2018; Vohník & Josefiová, 2024). Further, a recent study surveying 34 conserved plant genes involved in AMF profiled two Alismatales species, the freshwater duckweed *Spirodela polyrhiza* and the seagrass *Zostera marina*, and concluded both were likely non-mycorrhizal (NM) based on gene loss (Radhakrishnan *et al.*, 2020). However, seagrasses and freshwater lineages in the Alismatales have interesting evolutionary histories which involve multiple independent transitions into aquatic ecosystems (Les *et al.*, 1997; Chen *et al.*, 2022). Therefore, we set out to more robustly profile the Alismatales overall to better understand how aquatic transitions might inform the evolutionary trajectory of genes involved in fungal symbiosis.

To interrogate the genomic potential to form AMF associations in aquatic plant lineages in the order Alismatales, including seagrasses and freshwater relatives, we searched for the presence or absence of key genes implicated in AMF symbiosis in a set of plant genomes and transcriptomes. We used 65 publicly available genomes and transcriptomes downloaded from

NCBI GenBank (Table S1). This included 25 freshwater, 23 terrestrial and 17 marine

representatives, with 49 representing members of the order Alismatales. We performed

reciprocal blast searches of protein sequences to explore the presence or absence of homologs

of 45 genes reported to be related to AMF symbiosis (Table S2) (Delaux *et al.*, 2013, 2014;

Radhakrishnan *et al.*, 2020), with a special focus on six critical genes highlighted by

Radhakrishnan *et al.* (2020) including three that co-evolved with AMF associations (*RAD1*,

STR1, *STR2*) and three necessary for intracellular symbiosis (*SymRK*, *CCaMK/DMI3*,

CYCLOPS/IDP3).

Materials and Methods

Publicly available genomes and transcriptomes from 65 plants were downloaded from NCBI

(Table S1). This included 25 freshwater, 23 terrestrial and 17 marine representatives, with 49

representing members of the order Alismatales ('Darwin Tree of Life'; Ouyang *et al.*, 2006; The

International Brachypodium Initiative, 2010; Massa *et al.*, 2011; Lamesch *et al.*, 2011;

Bennetzen *et al.*, 2012; Ming *et al.*, 2013, 2015; Droc *et al.*, 2013; Regier *et al.*, 2013;

Motamayor *et al.*, 2013; Wang *et al.*, 2014, 2023, 2024; Schmutz *et al.*, 2014; Yang *et al.*, 2015;

Hirsch *et al.*, 2016; D'Esposito *et al.*, 2016; Michael *et al.*, 2017, 2021; Suzuki *et al.*, 2017;

Ruocco *et al.*, 2017; Malandrakis *et al.*, 2017; Zhang *et al.*, 2017; Entrambasaguas *et al.*, 2017;

McCormick *et al.*, 2018; Hoang *et al.*, 2018, 2020; Pecrix *et al.*, 2018; An *et al.*, 2019; Bellinger

et al., 2020; Yin *et al.*, 2021; Ma *et al.*, 2021, 2024; Park *et al.*, 2021; Rowarth *et al.*, 2021;

Abramson *et al.*, 2021; Gao *et al.*, 2022; Frisse *et al.*, 2022; Tang *et al.*, 2022; Bayer *et al.*,

2022; Ndhlovu & von der Heyden, 2022; Li *et al.*, 2023; Pasaribu *et al.*, 2023; Qian *et al.*, 2024).

We obtained the predicted AMF status for each plant family from FungalRoot v. 2

(Soudzilovskaia *et al.*, 2022). FungalRoot classifies plant families as obligate arbuscular

mycorrhizal (AM), non-mycorrhizal (NM), or AM-NM which refers to facultative AMs - plants that

are occasionally colonized by AMF, but are often observed without AMF in other conditions. Of

the 65 plants included in this work: 22 were AM, 23 were NM and 20 were AM-NM.

Using a set of 45 genes reported previously to be involved in AMF symbiosis (Table S2),

including six genes, *SymR/DMI2*, *CCaMK/DMI3*, *CYCLOPS/IPD3*, GRAS transcription factor

RAD1, and two half-ATP-binding cassette (ABC) transporters, *STR* and *STR2*, thought to be

critically important for forming intracellular symbioses and AMF (Delaux *et al.*, 2013, 2014;

Radhakrishnan *et al.*, 2020), we surveyed the genetic landscape of several plants across the

order Alismatales to assess whether they have maintained the potential to form AMF

associations as many species in this order have ancestors that evolutionarily transitioned from
terrestrial to aquatic habitats.

We used command-line NCBI BLAST v. 2.16.0+ (Camacho *et al.*, 2009) to perform reciprocal
blast searches for the set of 45 genes using protein sequences from *Medicago truncatula* (v.4)
as a reference, similar to searches performed in other studies of AMF (Delaux *et al.*, 2014;
Radhakrishnan *et al.*, 2020). For annotated genomes (n=53) we searched with blastp using the
protein sequences from *M. truncatula.* Top matches were then reciprocally searched back
against the annotation for *M. truncatula* using blastp. While, for unannotated genomes (n=2)
and transcriptomes (n=14), we instead searched with tblastn using the protein sequences from
M. truncatula. Top matches were then reciprocally searched against the *M. truncatula*
annotation using blastx. Subsequent analysis and visualization of reciprocal matches was
performed in R v. 4.3.1 (R Core Team, 2021) using the tidyverse v. 2.0.0 R package (Wickham
et al., 2019).

For the purposes of this study, we consider cases where a positive reciprocal match was made
(i.e. where the initially searched *M. truncatula* gene was subsequently the top hit in the
reciprocal search) to be homologous genes. Note, we did not further assess whether positive
reciprocal matches represented genes that were complete or functional. Further we consider
cases where an initial match was made, but the reciprocal match was negative, to potentially
indicate distant homology, such as paralogs, or genes that have undergone significant evolution
or divergence. Finally, we consider cases where no matches were made to most likely indicate
gene loss as the ancestral state of these genes is present. Note that for unannotated genomes
and transcriptomes, there are several additional reasons genes might be missed using this
approach including assembly issues, transcriptional variability, abundance or long introns.
However, including both unannotated genomes and transcriptomes allows us to have a more
robust understanding of the Alismatales gene space using currently available public data.

Results and Discussion

Across the 45 surveyed genes, positive reciprocal matches using protein sequences were found
in 57.27 ± 3.08% of predicted AM plants, 11.78 ± 2.83% of NM-AM plants, and 18.36 ± 4.25% of
NM plants (Figure 1). A similar trend emerged across habitat types: 61.26 ± 3.30% in terrestrial
plants, 10.22 ± 2.97% in freshwater plants, and 14.90 ± 4.32% in marine plants. Recovery of
positive reciprocal matches was highest in annotated genomes, followed by transcriptomes, and

lowest in unannotated genomes (e.g., for AM plants: $82.04 \pm 4.04\%$, $45.56 \pm 4.44\%$, and 23.05

$\pm 3.23\%$, respectively). This suggests that efforts to expand available Alismatales assembled

and annotated genomes, particularly for aquatic Alismatales, would improve detection accuracy,

and likely yield additional insights into the presence and absence of these genes in aquatic

lineages.

Regardless, the lower detection of positive reciprocal matches in freshwater and marine plants

suggests that these plants have lost or have experienced divergence in many of the genes

involved in AMF symbiosis. These findings support the idea that plants transitioning back into

aquatic environments have repeatedly lost the ability to form AMF associations, largely aligning

with the NM and NM-AM predictions for these lineages from the FungalRoot database based on

observational reports. Overall, our results show a pattern of likely independent loss of

conserved AMF associated genes across aquatic and marine Alismatales lineages and are

consistent with previous research showing convergent loss of AMF genes in other NM plants

(Delaux *et al.*, 2014; Radhakrishnan *et al.*, 2020).

Focusing on the six key genes previously identified as vital to AMF symbiosis (Radhakrishnan *et*

al., 2020), we observed similar patterns to the full gene set (Figure S1). Specifically for the three

genes that co-evolved with AMF associations (*RAD1, STR1, STR2*), reciprocal matches using

protein sequences were primarily found in terrestrial AM plants ($83.33 \pm 3.21\%$), with occasional

matches in NM-AM freshwater plants, but none in marine plants. Radhakrishnan *et al. (2020)*

suggested that the observed consistent loss of *RAD1*, *STR1* and *STR2* following the

abandonment of AMF associations indicates there may be selection against these genes once

AMF symbiosis is no longer needed. The absence of these genes in marine lineages reinforces

previous observational reports that seagrasses lack AMF associations (Nielsen *et al.*, 1999).

For the three genes required for intracellular symbiosis (*SymRK*, *CCaMK*/*DMI3*,

CYCLOPS/IDP3) most reciprocal matches using protein sequences were from terrestrial plants

($73.91 \pm 4.35\%$), with only *CCaMK*/*DMI3* detected in both freshwater and marine plants.

Radhakrishnan *et al.* (2020) found that these three genes were not absent from all species that

have lost AMF associations and instead observed that they were maintained in species that

formed other types of fungal and bacterial symbiosis. They also highlighted several additional

genes (*KinF, EPP1, VAPYRIN, LIN, CASTOR*, and *SYN*) which were maintained in species

capable of other types of intracellular symbioses. Of these additional genes, we only recovered

positive matches to *VAPYRIN* in aquatic plants.

Our results suggest that some genes involved in AMF associations (i.e., *CCaMK/DMI3*) are

retained in aquatic lineages, including seagrasses. Many of the 45 surveyed conserved genes

are involved in other microbial symbioses beyond AMF (e.g., orchid mycorrhizas, ericoid

mycorrhizas and legume-rhizobia symbioses) and their retention, loss or divergence may be

important for understanding their interactions with other microorganisms. For example,

CCaMK/DMI3 is thought to play a role in ectomycorrhizal (ECM) symbioses (Li *et al.*, 2024). In

our study, we found positive reciprocal matches for *CCaMK/DMI3* in the seagrass *Posidonia*

oceanica, which may relate to its reported ECM-like symbiotic relationship with the fungal root

endophyte *Posidoniomyces atricolor* (Vohník et al., 2015, 2019; Borovec & Vohník, 2018).

However, the function of *CCaMK/DMI3* in aquatic plants remains unclear. For example, in the

aquatic *Nelumbo nucifera*, which has also been reported to have *CCaMK/DMI3*, recent work

has documented a deletion in the kinase domain that likely renders it non-functional (Bravo *et*

al., 2016; Radhakrishnan *et al.*, 2020). These findings highlight the need for further research to

confirm whether *CCaMK/DMI3* and other symbiosis-related genes detected in aquatic

Alismatales remain functional and whether they have been co-opted for use in alternative

symbiotic relationships.

The loss of AMF associations in plants has been speculated to be due to adaptation to nutrient-

rich ecological niches, which are known to inhibit the formation of AMF (Read & Perez-Moreno,

2003). While aquatic environments are not wholly nutrient-rich, it is possible that the selective

pressures of freshwater and marine life, such as low oxygen or high salinity, make it harder to

maintain AMF (Nielsen *et al.*, 1999; Fusconi & Mucciarelli, 2018). Additionally, many aquatic

plants utilize their leaves, in addition to their roots, for uptake of nutrients which may make AMF

associations, which are traditionally thought to occur in roots, less necessary (Short & McRoy,

1984; Terrados & Williams, 1997). Overall, the Alismatales are a notable group for investigating

the evolution of plant-fungal symbiosis and the impact of aquatic transitions on host-microbe

interactions. This work serves to highlight the need for additional genome assembly and

annotation efforts for plants in this group in order to understand the molecular basis of these

interactions and the functional significance of retained symbiosis-related genes.

Acknowledgements

We would like to thank Jonathan A. Eisen for engaging in initial conversations around this work

and for feedback on this manuscript. We would like to thank the Darwin Tree of Life project

(https://www.darwintreeoflife.org) for making their genome assemblies publicly available without

restriction.

Competing interests

223 The authors declare no competing interests.

Author contributions

225 CLE conceived and designed the analysis, analyzed the data, prepared figures and/or tables,
226 wrote and reviewed drafts of the paper. JA prepared figures and/or tables, and wrote and
227 reviewed drafts of the paper. JES reviewed drafts of the paper.

Data availability

229 All code used in this work has been deposited on Github (casett/Seagrass_AMF) and archived
230 in Zenodo (DOI: 10.5281/zenodo.14218123). The genomic and transcriptomic data used in this
231 work are publicly available and were obtained from NCBI GenBank (Table S1).

Funding sources

CLE was supported by the National Science Foundation (NSF) under a NSF Ocean Sciences
Postdoctoral Fellowship (Award No. 2205744). JA was supported by the NSF REU The National
Summer Undergraduate Research Project, Award No. 2149582. JES is a CIFAR Fellow in the
program Fungal Kingdom: Threats and Opportunities and was partially supported by NSF
awards EF-2125066 and IOS-2134912. Computations were performed using the computer
clusters and data storage resources of the UC Riverside HPCC, which were funded by grants
from NSF (MRI-2215705, MRI-1429826) and NIH (1S10OD016290-01A1). The funders had no
role in study design, data collection and analysis, decision to publish, or preparation of the
manuscript.

References

Abramson BW, Novotny M, Hartwick NT, Colt K, Aevermann BD, Scheuermann RH,
Michael TP. **2021**. The genome and preliminary single-nuclei transcriptome of *Lemna minuta*
reveals mechanisms of invasiveness. *Plant physiology* **188**: 879–897.

An D, Zhou Y, Li C, Xiao Q, Wang T, Zhang Y, Wu Y, Li Y, Chao D-Y, Messing J, *et al.*
2019. Plant evolution and environmental adaptation unveiled by long-read whole-genome
sequencing of *Spirodela*. *Proceedings of the National Academy of Sciences* **116**: 18893–18899.

Bayer PE, Fraser MW, Martin BC, Petereit J, Severn-Ellis AA, Sinclair EA, Batley J,
Kendrick GA, Edwards D. **2022**. Not all pathways are the same – unique adaptations to
submerged environments emerge from comparative seagrass genomics. *bioRxiv*:
2022.11.22.517588.

Beck-Nielsen D, Madsen TV. **2001**. Occurrence of vesicular–arbuscular mycorrhiza in aquatic
macrophytes from lakes and streams. *Aquatic botany* **71**: 141–148.

Bellinger MR, Paudel R, Starnes S, Kambic L, Kantar MB, Wolfgruber T, Lamour K, Geib
S, Sim S, Miyasaka SC, *et al.* 2020. Taro Genome Assembly and Linkage Map Reveal QTLs
for Resistance to Taro Leaf Blight. *G3 Genes|Genomes|Genetics* **10**: 2763–2775.

Bennetzen JL, Schmutz J, Wang H, Percifield R, Hawkins J, Pontaroli AC, Estep M, Feng
L, Vaughn JN, Grimwood J, *et al.* 2012. Reference genome sequence of the model plant
Setaria. *Nature biotechnology* **30**: 555–561.

Bohrer KE, Friese CF, Amon JP. **2004**. Seasonal dynamics of arbuscular mycorrhizal fungi in
differing wetland habitats. *Mycorrhiza* **14**: 329–337.

Bonfante P, Anca IA. **2009**. Plants, mycorrhizal fungi, and bacteria: a network of interactions.
Annual review of microbiology **63**.

Borovec O, Vohník M. **2018**. Ontogenetic transition from specialized root hairs to specific root-
fungus symbiosis in the dominant Mediterranean seagrass *Posidonia oceanica*. *Scientific*
reports **8**: 10773.

Bravo A, York T, Pumplin N, Mueller LA, Harrison MJ. **2016**. Genes conserved for
arbuscular mycorrhizal symbiosis identified through phylogenomics. *Nature plants* **2**: 15208.

Brundrett MC. **2009**. Mycorrhizal associations and other means of nutrition of vascular plants:
understanding the global diversity of host plants by resolving conflicting information and
developing reliable means of diagnosis. *Plant and soil* **320**: 37–77.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. **2009**.
BLAST+: architecture and applications. *BMC bioinformatics* **10**.

Chen L-Y, Lu B, Morales-Briones DF, Moody ML, Liu F, Hu G-W, Huang C-H, Chen J-M,
Wang Q-F. **2022**. Phylogenomic Analyses of Alismatales Shed Light into Adaptations to Aquatic
Environments. *Molecular biology and evolution* **39**.

Clayton JS, Bagyaraj DJ. **1984**. Vesicular-arbuscular mycorrhizas in submerged aquatic plants
of New Zealand. *Aquatic botany* **19**: 251–262.

Darwin Tree of Life.

Delaux P-M, Séjalon-Delmas N, Bécard G, Ané J-M. **2013**. Evolution of the plant-microbe
symbiotic 'toolkit'. *Trends in plant science* **18**: 298–304.

Delaux P-M, Varala K, Edger PP, Coruzzi GM, Pires JC, Ané J-M. **2014**. Comparative
phylogenomics uncovers the impact of symbiotic associations on host genome evolution. *PLoS*
genetics **10**: e1004487.

D'Esposito D, Orrù L, Dattolo E, Bernardo L, Lamontanara A, Orsini L, Serra IA, Mazzuca

S, Procaccini G. **2016**. Transcriptome characterisation and simple sequence repeat marker
discovery in the seagrass *Posidonia oceanica*. *Scientific Data* **3**: 1–9.

Droc G, Larivière D, Guignon V, Yahiaoui N, This D, Garsmeur O, Dereeper A, Hamelin C,
Argout X, Dufayard J-F, *et al.* **2013**. The Banana Genome Hub. *Database: the journal of*
biological databases and curation **2013**: bat035.

Entrambasaguas L, Jahnke M, Biffali E, Borra M., Sanges R, Marín-Guirao L, Procaccini
G. **2017**. Tissue-specific transcriptomic profiling provides new insights into the reproductive
ecology and biology of the iconic seagrass species *Posidonia oceanica. Marine genomics* **35**:
51–61.

Frisse L, Martinez MA, Pirro S. **2022**. The Complete Genome Sequence of *Amorphophallus*
titanum, the Corpse Flower. *Biodiversity Genomes*.

Fusconi A, Mucciarelli M. **2018**. How important is arbuscular mycorrhizal colonization in
wetland and aquatic habitats? *Environmental and Experimental Botany* **155**: 128–141.

Gao Y, Zhang Y, Feng C, Chu H, Feng C, Wang H, Wu L, Yin S, Liu C, Chen H, *et al.* **2022**.
A chromosome-level genome assembly of *Amorphophallus konjac* provides insights into konjac
glucomannan biosynthesis. *Computational and structural biotechnology journal* **20**.

Heckman DS, Geiser DM, Eidell BR, Stauffer RL, Kardos NL, Hedges SB. **2001**. Molecular
evidence for the early colonization of land by fungi and plants. *Science (New York, N.Y.)* **293**.

Hirsch CN, Hirsch CD, Brohammer AB, Bowman MJ, Soifer I, Barad O, Shem-Tov D,
Baruch K, Lu F, Hernandez AG, *et al.* **2016**. Draft Assembly of Elite Inbred Line PH207
Provides Insights into Genomic and Transcriptome Diversity in Maize. *The Plant cell* **28**: 2700–
2714.

Hoang PTN, Fiebig A, Novák P, Macas J, Cao HX, Stepanenko A, Chen G, Borisjuk N,
Scholz U, Schubert I. **2020**. Chromosome-scale genome assembly for the duckweed *Spirodela*
intermedia, integrating cytogenetic maps, PacBio and Oxford Nanopore libraries. *Scientific*
reports **10**: 1–14.

Hoang PNT, Michael TP, Gilbert S, Chu P, Motley ST, Appenroth KJ, Schubert I, Lam E.
2018. Generating a high-confidence reference genome map of the Greater Duckweed by
integration of cytogenomic, optical mapping, and Oxford Nanopore technologies. *The Plant*
journal: for cell and molecular biology **96**.

Humphreys CP, Franks PJ, Rees M, Bidartondo MI, Leake JR, Beerling DJ. **2010**.
Mutualistic mycorrhiza-like symbiosis in the most ancient group of land plants. *Nature*
communications **1**: 103.

Khan AG, Belik M. **1995**. Occurence and Ecological Significance of Mycorrhizal Symbiosis in
Aquatic Plants. *Mycorrhiza*: 627–666.

Kohout P, Sýkorová Z, Ctvrtlíková M, Rydlová J, Suda J, Vohník M, Sudová R. **2012**.
Surprising spectra of root-associated fungi in submerged aquatic plants. *FEMS microbiology*
ecology **80**: 216–235.

Lamesch P, Berardini TZ, Li D, Swarbreck D, Wilks C, Sasidharan R, Muller R, Dreher K,

Alexander DL, Garcia-Hernandez M, *et al.* 2011. The *Arabidopsis* Information Resource
(TAIR): improved gene annotation and new tools. *Nucleic acids research* **40**: D1202–D1210.

Les DH, Cleland MA, Waycott M. **1997**. Phylogenetic Studies in Alismatidae, II: Evolution of
Marine Angiosperms (Seagrasses) and Hydrophily. *Systematic Botany* **22**: 443–463.

Li H, Ge Y, Zhang Z, Zhang H, Wang Y, Wang M, Zhao X, Yan J, Li Q, Qin L, *et al.* 2024.
Arbuscular mycorrhizal conserved genes are recruited for ectomycorrhizal symbiosis. *The New*
phytologist **242**: 1860–1864.

Li F, Yang J-J, Sun Z-Y, Wang L, Qi L-Y, Sina A, Liu Y-Q, Zhang H-M, Dang L-F, Wang S-J,
***et al.* 2023**. Plant-on-chip: Core morphogenesis processes in the tiny plant *Wolffia australiana*.
PNAS Nexus **2**: gad141.

Malandrakis E, Dadali O, Kavouras M, Danis T, Panagiotaki P, Miliou H, Tsioli S, Orfanidis
S, Küpper FC, Exadactylos A. **2017**. Identification of the abiotic stress-related transcription in
little Neptune grass *Cymodocea nodosa* with RNA-seq. *Marine genomics* **34**: 47–56.

Ma X, Olsen JL, Reusch TBH, Procaccini G, Kudrna D, Williams M, Grimwood J,
Rajasekar S, Jenkins J, Schmutz J, *et al.* 2021. Improved chromosome-level genome
assembly and annotation of the seagrass, *Zostera marina* (eelgrass). *F1000Research* **10**: 289.

Massa SI, Pearson GA, Aires T, Kube M, Olsen JL, Reinhardt R, Serrao EA, Arnaud-
Haond S. **2011**. Expressed sequence tags from heat-shocked seagrass *Zostera noltii*
(Hornemann) from its southern distribution range. *Marine genomics* **4**: 181–188.

Ma X, Vanneste S, Chang J, Ambrosino L, Barry K, Bayer T, Bobrov AA, Boston L,
Campbell JE, Chen H, *et al.* 2024. Seagrass genomes reveal ancient polyploidy and
adaptations to the marine environment. *Nature Plants* **10**: 240–255.

McCormick RF, Truong SK, Sreedasyam A, Jenkins J, Shu S, Sims D, Kennedy M,
Amirebrahimi M, Weers BD, McKinley B, *et al.* 2018. The *Sorghum bicolor* reference
genome: improved assembly, gene annotations, a transcriptome atlas, and signatures of
genome organization. *The Plant journal: for cell and molecular biology* **93**: 338–354.

Michael TP, Bryant D, Gutierrez R, Borisjuk N, Chu P, Zhang H, Xia J, Zhou J, Peng H, El
Baidouri M, *et al.* 2017. Comprehensive definition of genome features in *Spirodela polyrhiza* by
high-depth physical mapping and short-read DNA sequencing strategies. *The Plant journal: for*
cell and molecular biology **89**.

Michael TP, Ernst E, Hartwick N, Chu P, Bryant D, Gilbert S, Ortleb S, Baggs EL,
Sowjanya Sree K, Appenroth KJ, *et al.* 2021. Genome and time-of-day transcriptome of
Wolffia australiana link morphological minimization with gene loss and less growth control.
Genome research **31**: 225–238.

Ming R, VanBuren R, Liu Y, Yang M, Han Y, Li L-T, Zhang Q, Kim M-J, Schatz MC,
Campbell M, *et al.* 2013. Genome of the long-living sacred lotus (*Nelumbo nucifera* Gaertn.).
Genome biology **14**: 1–11.

Ming R, VanBuren R, Wai CM, Tang H, Schatz MC, Bowers JE, Lyons E, Wang M-L, Chen
J, Biggers E, *et al.* 2015. The pineapple genome and the evolution of CAM photosynthesis.
Nature genetics **47**: 1435–1442.

Motamayor JC, Mockaitis K, Schmutz J, Haiminen N, Iii DL, Cornejo O, Findley SD, Zheng
P, Utro F, Royaert S, *et al.* 2013. The genome sequence of the most widely cultivated cacao
type and its use to identify candidate genes regulating pod color. *Genome biology* **14**: 1–25.

Ndhlovu A, von der Heyden S. 2022. De novo assembly and annotation of the transcriptome
of the endangered seagrass *Zostera capensis*: Insights from differential gene expression under
thermal stress. *Marine genomics* **66**: 100984.

Nielsen SL, Thingstrup I, Wigand C. 1999. Apparent lack of vesicular–arbuscular mycorrhiza
(VAM) in the seagrasses *Zostera marina* L. and *Thalassia testudinum* Banks ex König. *Aquatic*
botany **63**: 261–266.

Ouyang S, Zhu W, Hamilton J, Lin H, Campbell M, Childs K, Thibaud-Nissen F, Malek RL,
Lee Y, Zheng L, *et al.* 2006. The TIGR Rice Genome Annotation Resource: improvements and
new features. *Nucleic acids research* **35**: D883–D887.

Park H, Park JH, Lee Y, Woo DU, Jeon HH, Sung YW, Shim S, Kim SH, Lee KO, Kim J-Y, *et*
***al.* 2021**. Genome of the world's smallest flowering plant, *Wolffia australiana*, helps explain its
specialized physiology and unique morphology. *Communications Biology* **4**: 1–12.

Parniske M. 2008. Arbuscular mycorrhiza: the mother of plant root endosymbioses. *Nature*
reviews. Microbiology **6**: 763–775.

Pasaribu B, Acosta K, Aylward A, Liang Y, Abramson BW, Colt K, Hartwick NT, Shanklin
J, Michael TP, Lam E. 2023. Genomics of turions from the Greater Duckweed reveal its
pathways for dormancy and re-emergence strategy. *The New phytologist* **239**: 116–131.

Pecrix Y, Staton SE, Sallet E, Lelandais-Brière C, Moreau S, Carrère S, Blein T, Jardinaud
M-F, Latrasse D, Zouine M, *et al.* 2018. Whole-genome landscape of *Medicago truncatula*
symbiotic genes. *Nature Plants* **4**: 1017–1025.

Qian Z-H, Li W, Wang Q-F, Liang S-C, Wu S, Li Z-Z, Chen J-M. 2024. The chromosome-level
genome of the submerged plant *Cryptocoryne crispatula* provides insights into the terrestrial–
freshwater transition in Araceae. *DNA research: an international journal for rapid publication of*
reports on genes and genomes **31**: dsae003.

Radhakrishnan GV, Keller J, Rich MK, Vernié T, Mbadinga Mbadinga DL, Vigneron N,
Cottret L, Clemente HS, Libourel C, Cheema J, *et al.* 2020. An ancestral signalling pathway
is conserved in intracellular symbioses-forming plant lineages. *Nature plants* **6**: 280–289.

Radhika KP, Rodrigues BF. 2007. Arbuscular Mycorrhizae in association with aquatic and
marshy plant species in Goa, India. *Aquatic botany* **86**: 291–294.

R Core Team. 2021. *R: A Language and Environment for Statistical Computing*.

Read DJ, Perez-Moreno J. 2003. Mycorrhizas and nutrient cycling in ecosystems - a journey
towards relevance? *The New phytologist* **157**: 475–492.

Redecker D, Kodner R, Graham LE. 2000. Glomalean fungi from the Ordovician. *Science*
(New York, N.Y.) **289**: 1920–1921.

Regier N, Baerlocher L, Münsterkötter M, Farinelli L, Cosio C. 2013. Analysis of the *Elodea*
nuttallii Transcriptome in Response to Mercury and Cadmium Pollution: Development of

Sensitive Tools for Rapid Ecotoxicological Testing.

Rowarth NM, Curtis BA, Einfeldt AL, Archibald JM, Lacroix CR, Gunawardena AHLAN.
2021. RNA-Seq analysis reveals potential regulators of programmed cell death and leaf
remodelling in lace plant (*Aponogeton madagascariensis*). *BMC plant biology* **21**: 1–20.

Ruocco M, Musacchia F, Olivé I, Costa MM, Barrote I, Santos R, Sanges R, Procaccini G,
Silva J. **2017**. Genomewide transcriptional reprogramming in the seagrass *Cymodocea nodosa*
under experimental ocean acidification. *Molecular ecology* **26**.

Schmutz J, McClean PE, Mamidi S, Wu GA, Cannon SB, Grimwood J, Jenkins J, Shu S,
Song Q, Chavarro C, *et al.* 2014. A reference genome for common bean and genome-wide
analysis of dual domestications. *Nature genetics* **46**: 707–713.

Short FT, McRoy CP. **1984**. Nitrogen Uptake by Leaves and Roots of the Seagrass *Zostera*
marina L. *Botanica marina* **27**: 547–556.

Soudzilovskaia NA, He J, Rahimlou S, Abarenkov K, Brundrett MC, Tedersoo L. **2022**.
FungalRoot v.2.0 - an empirical database of plant mycorrhizal traits: A response to Bueno et al.
(2021) 'Towards a consistent benchmark for plant mycorrhizal association databases'. *The New*
phytologist **235**: 1689–1691.

Šraj-Kržič N, Pongrac P, Klemenc M, Kladnik A, Regvar M, Gaberščik A. **2006**. Mycorrhizal
colonisation in plants from intermittent aquatic habitats. *Aquatic botany* **85**: 331–336.

Suzuki JY, Amore TD, Calla B, Palmer NA, Scully ED, Sattler SE, Sarath G, Lichty JS,
Myers RY, Keith LM, *et al.* 2017. Organ-specific transcriptome profiling of metabolic and
pigment biosynthesis pathways in the floral ornamental progenitor species *Anthurium amnicola*
Dressler. *Scientific reports* **7**: 1–15.

Tang H, Saina JK, Long Z-C, Chen J, Dai C. **2022**. De novo transcriptome assembly using
Illumina sequencing and development of EST-SSR markers in a monoecious herb *Sagittaria*
trifolia Linn. *PeerJ* **10**: e14268.

Terrados J, Williams SL. **1997**. Leaf versus root nitrogen uptake by the surfgrass *Phyllospadix*
torreyi. *Marine Ecology Progress Series* **149**: 267–277.

The International Brachypodium Initiative. **2010**. Genome sequencing and analysis of the
model grass *Brachypodium distachyon*. *Nature* **463**: 763–768.

Vohník M, Borovec O, Kolaříková Z, Sudová R, Réblová M. **2019**. Extensive sampling and
high-throughput sequencing reveal *Posidoniomyces atricolor* gen. et sp. nov. (Aigialaceae,
Pleosporales) as the dominant root mycobiont of the dominant Mediterranean seagrass
Posidonia oceanica. *MycoKeys* **55**: 59–86.

Vohník M, Borovec O, Župan I, Vondrášek D, Petrtýl M, Sudová R. **2015**. Anatomically and
morphologically unique dark septate endophytic association in the roots of the Mediterranean
endemic seagrass *Posidonia oceanica*. *Mycorrhiza* **25**: 663–672.

Vohník M, Josefiová J. **2024**. Novel epiphytic root-fungus symbiosis in the Indo-Pacific
seagrass *Thalassodendron ciliatum* from the Red Sea. *Mycorrhiza*.

Wang W, Haberer G, Gundlach H, Gläßer C, Nussbaumer T, Luo MC, Lomsadze A,

Borodovsky M, Kerstetter RA, Shanklin J, *et al.* 2014. The *Spirodela polyrhiza* genome
reveals insights into its neotenous reduction fast growth and aquatic lifestyle. *Nature*
communications **5**: 1–13.

Wang B, Qiu YL. **2006**. Phylogenetic distribution and evolution of mycorrhizas in land plants.
Mycorrhiza **16**.

Wang Z-F, Wu L-F, Chen L, Zhu W-G, Yu E-P, Xu F-X, Cao H-L. **2024**. Genome assembly of
Ottelia alismoides, a multiple-carbon utilisation aquatic plant. *BMC Genomic Data* **25**: 1–4.

Wang Y, Yang T, Wang D, Gou R, Jiang Y, Zhang G, Zheng Y, Gao D, Chen L, Zhang X, *et*
***al.* 2023**. Chromosome level genome assembly of colored calla lily (*Zantedeschia elliottiana*).
Scientific Data **10**: 1–7.

Wang B, Yeun LH, Xue J-Y, Liu Y, Ané J-M, Qiu Y-L. **2010**. Presence of three mycorrhizal
genes in the common ancestor of land plants suggests a key role of mycorrhizas in the
colonization of land by plants. *The New phytologist* **186**: 514–525.

Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, Grolemund G, Hayes
A, Henry L, Hester J, *et al.* 2019. Welcome to the Tidyverse. *Journal of Open Source Software*
4: 1686.

Wigand C, Stevenson JC. **1994**. The presence and possible ecological significance of
mycorrhizae of the submersed macrophyte, *Vallisneria americana*. *Estuaries* **17**: 206–215.

Yang Y, Chen X, Xu B, Li Y, Ma Y, Wang G. **2015**. Phenotype and transcriptome analysis
reveals chloroplast development and pigment biosynthesis together influenced the leaf color
formation in mutants of *Anthurium andraeanum* 'Sonate'. *Frontiers in plant science* **6**: 131822.

Yin J, Jiang L, Wang L, Han X, Guo W, Li C, Zhou Y, Denton M, Zhang P. **2021**. A high-
quality genome of taro (*Colocasia esculenta* (L.) Schott), one of the world's oldest crops.
Molecular ecology resources **21**: 68–77.

Zhang G-Q, Liu K-W, Li Z, Lohaus R, Hsiao Y-Y, Niu S-C, Wang J-Y, Lin Y-C, Xu Q, Chen
L-J, *et al.* 2017. The *Apostasia* genome and the evolution of orchids. *Nature* **549**: 379–383.

471
472

Figure 1. Loss of conserved symbiotic genes in aquatic plants. Heatmap showing initial blast and reciprocal blast results across marine, freshwater and terrestrial plants. Data type (genome [annotated], genome [unannotated], transcriptome) and FungalRoot v. 2 predicted AMF status (arbuscular mycorrhizal [AM], facultative arbuscular mycorrhizal [AM-NM], non-arbuscular mycorrhizal [NM]) are shown to aid interpretation.