The microbiome of *Calasterella californica* is shaped mostly by the genetics of

the host thalli

Ixchel González-Ramírez[1,2,a], Michael J. Song[3,a], Brent D. Mishler[1,2]

[1]University and Jepson Herbaria, University of California, Berkeley, CA

[2]Department of Integrative Biology, University of California, Berkeley, CA

[3]Skyline College, San Bruno, CA

[a]These authors contributed equally

Keywords: Liverwort, Bryophyte, Microbiome, California Conservation Genomics Project,

CCGP, *Methylobacterium*, Metagenomics

Short title: *Calasterella californica* microbiome composition

ABSTRACT

Premise of the study: Plant microbiomes have been shown to affect essential life processes like plant growth, defense mechanisms, and stress responses. Much of the work done on this topic has focused on vascular plants and model organisms. Liverwort biology is very different from that of vascular plants, making generalizations difficult. Here we characterize the microbiome of the liverwort *Calasterella californica* and explore whether climatic, geographic, and genetic factors are potential drivers of the microbiome composition.

Methods: We utilized the genomic data obtained through the California Conservation Genomics Project, which densely sampled individuals of *C. californica* across California. We selected 34 samples and utilized the MetaPhlAn pipeline to characterize the composition of the microbial communities for each sample.

Key results: We characterize the core microbiome of *Calasterella,* which is dominated by *Methylobacterium*. There is a weak association between the relative abundance of bacteria in the Hyphomicrobiales and liverwort size. Genetic distance was the most important factor in shaping beta diversity of the *Calasterella californica* microbiome. Environmental distance was only weakly predictive and geographic distance was not significant.

Conclusions: Leveraging genomic data obtained from field samples of a liverwort species allowed us to characterize the microbial diversity of *C. californica* and its variation across space. There is a group of taxa that are consistently associated with this plant. And the variation observed across samples seems to be linked to the genetic similarity of the liverworts.

INTRODUCTION

The community of bacteria, archaea, and small eukaryotes associated with an organism, known as the microbiome, has been shown to have a considerable role in shaping plant morphology, ecology, and evolution. Plant microbiomes play a central role in some of the most basic aspects of terrestrial plant life such as: mycorrhizae (Taylor et al., 1995), specialized nitrogen-fixation structures (Song et al., 2024), hormone signaling (reviewed in Li et al., 2023), and growth-promotion and disease protection (reviewed in Trivedi et al., 2020). Bryophytes have many well-characterized microbiome interactions, including the presence of mycorrhizae in liverworts and hornworts (reviewed in Pressel et al., 2010) and nitrogen-fixing cyanobionts in hornworts (Li et al., 2020). Thalloid liverworts in particular have been studied as proxies for early land-plant microbiomes (Alcaraz et al., 2018), which are hypothesized to have included nitrogen-fixing bacteria and methanotrophs (Knack et al. 2015).

Thalloid liverworts have been shown to associate with many members of *Methylobacterium* who live off of methanol excreted by the liverwort pores and in exchange promote host growth by secreting cytokinins (Alcaraz et al., 2018; Kutschera and Koopmann, 2005). Similarly, Alcaraz and colleagues (2018) found many other potentially symbiotic bacteria in the *Marchantia* microbiome that promote pathogen defense through the production of plant hormones in exchange for plant-produced polymers including growth-inducing genera like *Rhizobium* and organic degrading bacteria like *Lysobacter*. These bacteria may be actively recruited by the liverwort by inhibiting the growth of competing bacteria (Mewari and Kumar, 2008).

While the factors that shape the composition of the bryophyte microbiome are not well understood, some patterns have emerged. For example, it has been observed that bryophytes have a unique microbiome that is consistent over development (Bragina et al., 2012) and independent from geography (Opelt et al., 2007). Some studies suggest evidence for vertical

transmission of some beneficial *Burkholderia* taxa in moss (Bragina et al., 2013). Wicaksono et

al. (2023) found that *Riccia* liverworts recruit specific bacteria and that the surrounding soil is not

an important reservoir for them. However, they also found that enrichment of *Methylobacterium*

was dependent on field type (Wicaksono et al., 2023). Similarly, the bacteriome of *Marchantia*

inflexa liverworts were found to be highly shaped by habitat (Marks et al., 2017).

Figure 1. **a**. Collection locations of *Calasterella californica* for this study. **b**. Typical environment

and morphology of *C. californica*. **c**. Close up to the female thalli of *C. californica*.

Calasterella californica is a complex thalloid liverwort endemic to the west coast of North

America, extending from the southernmost part of Oregon to Baja California Norte in Mexico

(Fig. 1). This species is outstanding in the diversity of ecosystems in which it occurs, from humid

forests (*e.g.* in the Cascades region) to the desert (*e.g.* Anza Borrego). In all these ecosystems,

C. californica occurs in exposed areas, over bare soil or rock and plays an important role in soil

retention and secondary succession. In this paper, we characterize and compare the bacterial

metagenome of 34 samples of *C. californica* collected across the state of California. We explore

whether differences in microbiome composition across samples are associated with factors like

liverwort genetics, variation in environmental variables, or geographic distance between

samples. To achieve our goals, we utilize *C. californica* samples obtained as a part of the

California Conservation Genomics Project (CCGP), a state-funded conservation project to

provide high quality reference genomes for over 100 species and population-level resequencing

to study the effects of climate change and to inform conservation efforts (Shaffer et al., 2022;

Toffelmier et al., 2022; Fiedler et al. 2022; Beninde et al. 2022). While the stated goals of the

CCGP focus on landscape genomics of the host organisms," we propose that this population-

level sampling across the state also provides an opportunity to study metagenome variation and

bacterial ecology, and we provide a case study to demonstrate its potential.

METHODS

We selected thirty-four samples of *Calasterella californica* collected as part of the CCGP project

(Fig 1). The selection aimed to represent the geographic range, the environmental niche space,

and the genetic diversity of the 110 samples collected for the CCGP project. For each of the

samples, total genomic and symbiont DNA of one to four thalli was extracted using the Qiagen

DNeasy Plant Pro kit (California, USA). Library preparation and sequencing was carried out at

the UC Davis DNA Technologies & Expression Analysis Core Laboratory. Libraries were

sequenced using an Illumina NovaSeqX 300 (PE150) flow-cell.

Characterization of *C. californica* microbiome composition. For each sample, the paired-

end reads were mapped to the *Calasterella californica* isolate IGR150-G1 genome (González-

Ramírez et al., 2024) using bwa-mem (Li, 2013). We used samtools (Li et al., 2009) to filter

reads that mapped to the genome and reads that did not align to the genome were kept for

downstream analysis. We used MetaPhlAn on the Galaxy webserver (Galaxy Version

4.1.1+galaxy3; Giardine et al., 2005; Blanco-Míguez et al., 2023) to profile the composition of

microbial communities using these short-read data. MetaPhlAn can process raw reads without

the need for adapter trimming and quality filtering (Nicola Segata, MetaPhlAn developer; online

communication) and therefore the raw reads were used to maximize the amount of information.

Results were visualized using Krona (Galaxy Version 2.7.1+galaxy0; Ondov et al., 2011) and

with R (R Core Team 2021).

In order to compare the microbiome of field-collected samples vs tissue cultivated in-

vitro, we assessed the microbiome composition of the tissue used for the reference genome of

C. californica, which was grown from a spore in the lab. PacBio HiFi reads from González-

Ramírez et al. (2024) were run through the BugSeq pipeline (Fan et al., 2021). Metagenomic

classification was visualized using Krona (Ondov et al., 2011). This in-vitro sample required a

different pipeline, that can analyze long-read data, than the field-collected samples.

Exploring potential drivers of microbiome composition. To assess whether the microbiome

composition is correlated with (1) liverwort genetics, (2) environmental conditions, or (3) sample

geographic proximity, we performed mantel and partial-mantel tests using the ape package

(Paradis and Schliep, 2019) in R with 999 permutations. First, we calculated pairwise distance

matrices for each of the three variables. The liverwort genetic distance matrix was constructed

based on 23180 bi-allelic and uncorrelated SNPs using Manhattan distance.

To characterize the macro-environment of each sample, we obtained bioclimatic variables

values for 19 climatic variables from the WorldClim2 database (Fick et al., 2017) and used these

values to compute the pairwise euclidean environmental distance matrix. The geographical

distance matrix was constructed with the geographic coordinates of the collection points using

the `distm` function of the geosphere R package (Hijmans, 2024). Next, we constructed a

pairwise dissimilarity matrix based on the microbiome species relative abundance data from

MetaPhlAn using the Bray-Curtis dissimilarity index. Finally, we performed mantel tests between

the microbiome composition distance matrix and each of the three variables mentioned.

RESULTS

Composition of *Calasterella californica* microbiome. A total of 161 bacterial genera were

found in the in-vitro grown sample; this value was 177 across the 34 samples collected in the

field (Supp. Table S1). Fifty bacterial genera were shared between the in vivo and in vitro

samples: *Achromobacter, Acidovorax, Actinoplanes, Agreia, Agrobacterium, Alsobacter,*

Asticcacaulis, Aureimonas, Azospirillum, Belnapia, Brevundimonas, Burkholderia, Caulobacter,

Dankookia, Devosia, Duganella, Dyadobacter, Flavobacterium, Geodermatophilus,

Hansschlegelia, Herbaspirillum, Herbiconiux, Hymenobacter, Klebsiella, Massilia,

Mesorhizobium, Methylobacterium, Methylophilus, Microbacterium, Mucilaginibacter,

Mycolicibacterium, Neorhizobium, Nostoc, Novosphingobium, Paraburkholderia, Pedobacter,

Phenylobacterium, Polaromonas, Pseudomonas, Pseudorhodoferax, Rhizobium, Rhodoferax,

Sphingobium, Sphingomonas, Spirosoma, Tardiphaga, Variovorax, Williamsia, Xanthomonas,

and *Xylophilus*. Nonetheless, the proportions of these taxa vary substantially between the field-

collected and in-vitro samples (Supp. Fig. S1).

Microbial communities associated with individual *C. californica* samples were composed

on average of 27.85 genera (range: 15 – 50). Across the 177 genera represented in the 34 field-

collected samples, there are six genera of bacteria that were present in more than 70% of the

samples: *Methylobacterium* (97%), *Spirosoma* (85%), *Pseudomonas* (79%), *Inquilinus* (0.74%),

Dankookia (74%), and *Sphingomonas* (71%). All of these genera, with the exception of

Inquilinus, were also found in the liverwort grown from a spore in the lab. Furthermore, except

for *Dankookia*, all these genera have been reported to be endophytic in seed-plants (Afzal et al.

2019), often having beneficial effects on their plant hosts. The presence of these bacteria in the

microbiome across the *C. californica* samples suggests that these bacteria may represent the

core microbiome associated with *C. californica*, and that they could be vertically transmitted,

since the spore-grown plant had them too.

Drivers of variation in microbiome composition. When we compared the microbiome

composition across samples, we found that members of the Hyphomicrobiales were abundant in

the majority of samples, especially taxa in the Methylobacteriaceae (Fig. 2). *Methylobacterium*

was found in high abundance in many but not all of our samples. Notably, these taxa did not

constitute a large proportion in the *in vitro* sample (Supp. Table S1). Since it has been

hypothesized that members of the Hyphomicrobiales (syn. Rhizobiales), which contains many

nitrogen-fixing taxa and methanogens, promote growth, we tested whether there was an

association between thallus width (as a proxy for thallus size) and the proportion of the

microbiome constituted by members of the Hyphomicrobiales (Figure 4). We found a weak yet

significant association (F-statistic = 5.223 on 1 and 31 DF, p = 0.0293, adjusted r squared =

0.1166). Interestingly, Hyphomicrobiales were found to be relatively abundant across all

samples of C. californica (Supp. Fig. S1).

Figure 2. Relative abundance (in percentage of bacterial reads identified) of different bacteria families across *C. californica* samples.

We found no significant correlation of microbiome composition with geographic distance or environmental distance (Fig. 3). We did, however, find a significant correlation with genetic distance (z-stat = 1076314, p-value = 0.03). This correlation was still significant when accounting for geographic distance (r statistic = 0.205, p-value = 0.023; partial mantel test). When visualizing the similarity of samples following their microbiome community and genetics of the host liverworts (Supp. Fig. S2), while the clustering is not identical, large groups remain congruent. We also found a weakly significant association between microbiome community

181   distance and ecological distance when accounting for geographic distance (r statistic = 0.133, p-

182   value = 0.095; partial mantel test).

184   **Figure 3.** Correlation of pairwise distances between microbiome composition and three

185   potentially explanatory variables. **a.** Geographic distance, **b.** Environmental distance, and **c.**

186   Genetic distance.

 Figure 4. Mean thallus width by proportion of Hyphomicrobiales for *Calasterella* samples. A line

of best fit was added using a linear regression model (standard error in grey).

DISCUSSION

These findings corroborate previous studies that found that liverworts recruit specific bacteria,

especially *Methylobacterium* (Alcaraz et al., 2018; Kutschera and Koopmann, 2005), and that

there is an effect of the environment on microbiome composition (Wicaksono et al., 2023; Marks

et al., 2017). We also found evidence of a correlation between the relative abundance of

Hyphomicrobiales microbes in an individual's microbiome and its thallus size. Given that there

could be an association between genetic similarity and environment due to selection or other

factors, we also compared genetic distances among the host plants to test whether genetic

factors or environment were more strongly correlated with microbiome community composition.

We found that genetic distance was the most important factor in shaping beta diversity of the

Calasterella microbiome, but that the environment was also weakly predictive. The joint

importance of both environmental factors and host genetics in shaping the plant microbiome has

also been found in duckweeds (Wei and Tan, 2023), *Boechera* (Wagner et al., 2016), and maize

(He et al., 2024). Notably, these patterns are contrary to the major patterns observed in humans

in which the environment is more important than genetics in shaping the gut microbiome

(Rothschild et al., 2018).

There has been much recent work on vascular plants to investigate the role of host

genetics in shaping the root and leaf microbiome (reviewed in Zhang et al., 2023). However,

liverwort morphology is very different from that of vascular plants, making these assessments

hard to generalize. A recent study on the microbiome of the liverwort *Riccia* suggests that

bacterial communities associated with liverworts are highly stable, showing little variation across

environmental variables like soil type (Wicaksono et al. 2023), which is consistent with our

results. With the recent publication of high-quality liverwort genomes (González-Ramírez et al.,

2024; Diop et al., 2020; Linde et al., 2023), future work can now begin to assess how genetics

shapes the bryophyte metagenome and similarities and differences with vascular plants.

Additionally, our methodological approach does not allow us to make inferences about whether

the bacteria identified here are endophytic or epiphytic; further studies could categorize these

groups.

This demonstration of the power of the CCGP dataset to elucidate the ecology of the

liverwort microbiome using metagenomic techniques suggests that this approach should be

applied to all of the taxa of the CCGP that have a microbiome. Song et al. (2024) has previously

used similar approaches focusing on the *Azolla* endosymbionts, in a well characterized

system—the leaf pocket. However, many of the taxa of the CCGP may also have microbiomes

that could be studied, including taxa for which leaf or whole animal sampling was performed:

Quercus tomentella (Mead et al., 2024), *Linanthus parryae* (Anghel et al., 2022), *Hetaerina*

americana (Grether et al., 2023), *Branchinecta lynchi* (Kieran Blair et al., 2023), *Polyergus*

mexicanus (Cash et al., 2024), and many others (see https://www.ccgproject.org/reference-

genomes-tracking). We encourage future collaboration to address the multiple avenues of

research that these data from large scale genomic approaches can facilitate.

ACKNOWLEDGEMENTS

The authors thank Carl Rothfels for his constructive comments and edits to the manuscript.

Thank you to the California Conservation Genomics Project, with funding provided to the

University of California by the State of California, State Budget Act of 2019 [UC Award ID RSI-

19-690224]. IGR was supported by a UC Mexus-CONACyT fellowship (number 709967), a

Plant Science Fellowship by Oak Spring Garden Foundation, and the Philomathia Graduate

Fellowship in Environmental Sciences, at UC Berkeley.

AUTHOR CONTRIBUTIONS

IGR and MJS designed the study, performed the analyses and wrote the first draft of the

manuscript. BDM supervised the work and edited the manuscript.

DATA AVAILABILITY STATEMENT

All the data and code used for the analyses is hosted in the GitHub repository:

https://github.com/ixchelgzlzr/microbiome_Ccal.git

LITERATURE CITED

Afzal, I., Z. K. Shinwari, S. Sikandar, and S. Shahzad. 2019. Plant beneficial endophytic bacteria: Mechanisms, diversity, host range and genetic determinants. *Microbiological research*, *221*, 36-49.

Alcaraz, L.D., M. Peimbert, H.R. Barajas, A.E. Dorantes-Acosta, J.L. Bowman, and M.A. Arteaga-Vázquez. 2018. *Marchantia* liverworts as a proxy to plants' basal microbiomes. *Scientific Reports*, 8: 12712.

Anghel, I.G., S.J. Jacobs, M. Escalona, M.P. Marimuthu, C.W. Fairbairn, E. Beraut, O. Nguyen, E. Toffelmier, H.B. Shaffer, and F. Zapata. 2022. Reference genome of the color polymorphic desert annual plant sandblossoms, *Linanthus parryae*. *Journal of Heredity*, *113*(6), pp.712-721.

Beninde, J., E. Toffelmier, and H. B. Shaffer. 2022. A brief history of population genetic research in California and an evaluation of its utility for conservation decision-making. *Journal of Heredity,* 113: 604–614.

Blanco-Míguez, A., F. Beghini, F. Cumbo, L.J. McIver, K.N. Thompson, M. Zolfo, P. Manghi, L. Dubois, K.D. Huang, A.M. Thomas, and W.A. Nickols. 2023. Extending and improving metagenomic taxonomic profiling with uncharacterized species using MetaPhlAn 4. *Nature Biotechnology*, 41(11), 1633-1644.

Bragina A., M. Cardinale, C. Berg, and G. Berg. 2013. Vertical transmission explains the specific Burkholderia pattern in Sphagnum mosses at multi-geographic scale. *Frontiers in Microbiology*, 4: 394.

Bragina, A., C. Berg, M. Cardinale, A. Shcherbakov, V. Chebotar, and G. Berg. 2012. Sphagnum mosses harbour highly specific bacterial diversity during their whole lifecycle. *The ISME journal*, *6*(4): 802-813.

Cash, E.I., M. Escalona, P.S. Ward, R. Sahasrabudhe, C. Miller, E. Toffelmier, C. Fairbairn, W. Seligmann, H.B. Shaffer, and N.D. Tsutsui. 2024. Reference genome of the kidnapper ant, *Polyergus mexicanus*. *Journal of Heredity*, p.esae047.

Diop, S.I., O. Subotic, A. Giraldo-Fonseca, M. Waller, A. Kirbis, A. Neubauer, G. Potente, R. Murray-Watson, F. Boskovic, Z. Bont, and Z. Hock. 2020. A pseudomolecule-scale genome assembly of the liverwort Marchantia polymorpha. *The Plant Journal*, 101(6), 1378-1396.

Fan, J., S. Huang, and S. D. Chorlton. 2021. BugSeq: a highly accurate cloud platform for long-read metagenomic analyses. *BMC bioinformatics*, 22, 1-12.

287   Fick, S.E. and R.J. Hijmans. 2017. WorldClim 2: new 1km spatial resolution climate surfaces for
288   global land areas. *International Journal of Climatology,* 37 (12): 4302-4315.
289

290   Fiedler, P. L., B. Erickson, M. Esgro, M. Gold, J. M. Hull, J. M. Norris, B. Shapiro, et al. 2022.
291   Seizing the moment: The opportunity and relevance of the California Conservation
292   Genomics Project to state and federal conservation policy. *Journal of Heredity,* 113:
293   589–596.
294

295   Giardine, B., C. Riemer, R.C. Hardison, R. Burhans, L. Elnitski, P. Shah, Y. Zhang, D.
296   Blankenberg, I. Albert, J. Taylor, and W. Miller. 2005. Galaxy: a platform for interactive
297   large-scale genome analysis. *Genome research*, 15(10), 1451-1455.
298

299   González-Ramírez, I.S., M. Escalona, C. Miller, N. Chumchim, M.P. Marimuthu, O. Nguyen,
300   M.J. Song, and B.D. Mishler. 2024. A genome assembly for the California endemic
301   liverwort Calasterella californica. *Journal of Heredity,* p.esae065.
302

303   Grether, G.F., J. Beninde, E. Beraut, N. Chumchim, M. Escalona, Z.G. MacDonald, C. Miller, R.
304   Sahasrabudhe, A.M. Shedlock, E. Toffelmier, and H.B. Shaffer. 2023. Reference
305   genome for the American rubyspot damselfly, *Hetaerina americana*. *Journal of Heredity*,
306   *114*(4), pp.385-394.
307

308   He, X., D. Wang, Y. Jiang, M. Li, M. Delgado-Baquerizo, C. McLaughlin, C. Marcon, L. Guo, M.
309   Baer, Y.A. Moya, and N. Von Wirén. 2024. Heritable microbiome variation is correlated
310   with source environment in locally adapted maize varieties. *Nature Plants*, 10(4), 598-
311   617.
312

313   Hijmans R. 2024. geosphere: Spherical Trigonometry. R package version 1.5-20,
314   <https://CRAN.R-project.org/package=geosphere>.
315

316   Kieran Blair, S.R., A. Schreier, M. Escalona, A.J. Finger, S.E. Joslin, R. Sahasrabudhe, M.P.
317   Marimuthu, O. Nguyen, N. Chumchim, E.R. Morris, and H. Mangelson. 2023. A draft
318   reference genome of the Vernal Pool Fairy Shrimp, *Branchinecta lynchi*. *Journal of
319   Heredity*, *114*(1), pp.81-87.
320

321   Knack, J. J., L. W. Wilcox, P-M. Delaux, J-M. Ané, M. J. Piotrowski, M. E. Cook, J. M. Graham,
322   and L. E. Graham. 2015. Microbiomes of streptophyte algae and bryophytes suggest
323   that a functional suite of microbiota fostered plant colonization of land. *International
324   Journal of Plant Sciences,* 176, (5): 405-420.
325

326   Kutschera, U., and V. Koopmann. 2005. Growth in liverworts of the Marchantiales is promoted
327   by epiphytic methylobacteria. *Naturwissenschaften,* 92: 347-349.
328

Li, H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv preprint arXiv:1303.3997.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, R.
Durbin, and 1000 Genome Project Data Processing Subgroup, 2009. The sequence
alignment/map format and SAMtools. *Bioinformatics*, 25(16): 2078-2079.

Li, J. H., M. M. Aslam, Y. Y. Gao, L. Dai, G. F. Hao, Z. Wei, M. X. Chen, and F. Dini-Andreote.
2023. Microbiome-mediated signal transduction within the plant holobiont. *Trends in
Microbiology,* 31(6): 616-628.

Li, F. W., T. Nishiyama, M. Waller, E. Frangedakis, J. Keller, Z. Li, N. Fernandez-Pozo et al.
2020. *Anthoceros* genomes illuminate the origin of land plants and the unique biology of
hornworts. *Nature plants,* 6(3): 259-272.

Linde, A. M., S. Singh, J. L. Bowman, M. Eklund, N. Cronberg, and U. Lagercrantz. 2023.
Genome evolution in plants: complex thalloid liverworts (Marchantiopsida). *Genome
Biology and Evolution*, 15(3), evad014.

Marks, R.A., J.J. Smith, Q. Cronk, and D.N. McLetchie. 2018.Variation in the bacteriome of the
tropical liverwort, *Marchantia inflexa*, between the sexes and across habitats. *Symbiosis,*
75: 93–101.

Mead, A., S.T. Fitz-Gibbon, M. Escalona, E. Beraut, S. Sacco, M.P. Marimuthu, O. Nguyen,
and V.L. Sork. 2024. The genome assembly of Island Oak (*Quercus tomentella*), a
relictual island tree species. *Journal of Heredity*, *115*(2), pp.221-229.

Mewari, N., and P. Kumar. Antimicrobial Activity of Extracts of *Marchantia polymorpha*. 2008.
Pharmaceutical biology, *46*(10-11), pp.819-822.

Ondov, B. D., N. H. Bergman, and A. M. Phillippy. 2011. Interactive metagenomic visualization
in a Web browser. *BMC bioinformatics*, 12, 1-10.

Opelt K., C. Berg, S. Schönmann, L. Eberl, G. Berg. 2007. High specificity but contrasting
biodiversity of *Sphagnum*-associated bacterial and plant communities in bog ecosystems
independent of the geographical region. *ISME J.*, 1:502–16.

Paradis, E., and K. Schliep. 2019. ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics*, 35(3), 526-528.

Pressel, S., M. I. Bidartondo, R. Ligrone, and J. G. Duckett. 2010. Fungal symbioses in
bryophytes: new insights in the twenty first century. *Phytotaxa,* 9(1): 238-253.

R Core Team. 2021. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Rothschild, D., O. Weissbrod, E. Barkan, A. Kurilshikov, T. Korem, D. Zeevi, P.I. Costea, A.
Godneva, I.N. Kalka, N. Bar, and S. Shilo. 2018. Environment dominates over host
genetics in shaping human gut microbiota. *Nature*, 555(7695), 210-215.
378

379 Shaffer, H. B., E. Toffelmier, R. B. Corbett-Detig, M. Escalona, B. Erickson, P. Fiedler, M. Gold,
380   R.J. Harrigan, S. Hodges, T.K. Luckau, and C. Miller. 2022. Landscape Genomics to
381   Enable Conservation Actions: The California Conservation Genomics Project. *Journal of*
382   *Heredity,* 113: 577–588.
383

384 Song, M.J., F-W Li, F. Freund, C.M. Tribble, E. Toffelmier, C. Miller, H.B. Shaffer, and C.
385   Rothfels. 2024. The nitrogen-fixing fern *Azolla* has a complex microbiome characterized
386   by multiple modes of transmission. *bioRxiv* https://doi.org/10.1101/2024.05.20.592813
387

388 Taylor, T. N., W. Remy, H. Hass, and H. Kerp. 1995. Fossil arbuscular mycorrhizae from the
389   Early Devonian. *Mycologia,* 87(4): 560-573.
390

391 Toffelmier, E., J. Beninde, and H. B. Shaffer. 2022. The phylogeny of California, and how it
392   informs setting multispecies conservation priorities. *Journal of Heredity,* 113: 597–603.
393

394 Trivedi, P., J. E. Leach, S. G. Tringe, T. Sa, and B. K. Singh. 2020. Plant–microbiome
395   interactions: from community assembly to plant health. *Nature reviews microbiology,*
396   18(11): 607-621.
397

398 Wagner, M. R., D. S. Lundberg, T. G. Del Rio, S. G. Tringe, J. L. Dangl, and T. Mitchell-Olds.
399   2016. Host genotype and age shape the leaf and root microbiomes of a wild perennial
400   plant. *Nature communications*, 7(1): 12151.
401

402 Wei, N., and J. Tan. 2023. Environment and host genetics influence the biogeography of plant
403   microbiome structure. *Microbial Ecology*, 86(4), 2858-2868.
404

405 Wicaksono, W.A., B. Semler, M. Pöltl, C. Berg, G. Berg, and T. Cernava. 2023. The microbiome
406   of *Riccia* liverworts is an important reservoir for microbial diversity in temporary
407   agricultural crusts. *Environmental Microbiome,* 18: 46
408

409 Zhang, J., W. Liu, J. Bu, Y. Lin, and Y. Bai. 2023. Host genetics regulate the plant microbiome.
410   *Current Opinion in Microbiology*, 72, 102268.
411

Supplemental Information

Supplemental Table S1. Genera of bacteria in the microbial community associated with in-vitro and field-collected specimens of *C. californica*.

In vitro	In vivo	Unique in vitro	Unique in vivo	Shared
Aaosphaeria	Achromobacter	Aaosphaeria	Acinetobacter	Achromobacter
Acanthamoeba	Acidovorax	Acanthamoeba	Aeromicrobium	Acidovorax
Achromobacter	Acinetobacter	Acidobacterium	Altererythrobacter	Actinoplanes
Acidobacterium	Actinoplanes	Actinomadura	Amnibacterium	Agreia
Acidovorax	Aeromicrobium	Acytostelium	Anaerospora	Agrobacterium
Actinomadura	Agreia	Afipia	Aquabacterium	Alsobacter
Actinoplanes	Agrobacterium	Agromyces	Aquincola	Asticcacaulis
Acytostelium	Alsobacter	Alternaria	Arcicella	Aureimonas
Afipia	Altererythrobacter	Ancylobacter	Azohydromonas	Azospirillum
Agreia	Amnibacterium	Ankyra	Beijerinckiaceae_unclassified	Belnapia
Agrobacterium	Anaerospora	Aphanomyces	Betaproteobacteria_unclassified	Brevundimonas
Agromyces	Aquabacterium	Aquibium	Bifidobacterium	Burkholderia
Alsobacter	Aquincola	Arenibacter	Blastococcus	Caulobacter
Alternaria	Arcicella	Aspergillus	Bosea	Dankookia
Ancylobacter	Asticcacaulis	Aureococcus	Burkholderiales_unclassified	Devosia
Ankyra	Aureimonas	Bacidia	Caballeronia	Duganella
Aphanomyces	Azohydromonas	Bacillus	Calothrix	Dyadobacter
Aquibium	Azospirillum	Bartonella	Cellulomonas	Flavobacterium
Arenibacter	Beijerinckiaceae_unclassified	Beauveria	Chiayiivirga	Geodermatophilus
Aspergillus	Belnapia	Bipolaris	Chimaeribacter	Hansschlegelia
Asticcacaulis	Betaproteobacteria_unclassified	Blastochloris	Chlorogloea	Herbaspirillum
Aureimonas	Bifidobacterium	Bodomorpha	Chryseobacterium	Herbiconiux
Aureococcus	Blastococcus	Botrytis	Clostridium	Hymenobacter
Azospirillum	Bosea	Bradyrhizobium	Conexibacter	Klebsiella
Bacidia	Brevundimonas	Cadophora	Corallococcus	Massilia
Bacillus	Burkholderia	Caenimonas	Curtobacterium	Mesorhizobium
Bartonella	Burkholderiales_unclassified	Canariomyces	Cutibacterium	Methylobacterium
Beauveria	Caballeronia	Candida	Cylindrospermum	Methylophilus
Belnapia	Calothrix	Candidatus Methylopumilus	Desmonostoc	Microbacterium
Bipolaris	Caulobacter	Capnocytophaga	Emticicia	Mucilaginibacter
Blastochloris	Cellulomonas	Capronia	Enterobacter	Mycolicibacterium
Bodomorpha	Chiayiivirga	Ceraceosorus	Enterococcus	Neorhizobium
Botrytis	Chimaeribacter	Cercospora	Erwinia	Nostoc
Bradyrhizobium	Chlorogloea	Chaetomium	Friedmanniella	Novosphingobium
Brevundimonas	Chryseobacterium	Chlorella	Frigoribacterium	Paraburkholderia

Burkholderia	Clostridium	Chromohalobacter	Frondihabitans	Pedobacter
Cadophora	Conexibacter	Cladophialophora	Fulvivirgaceae_unclassified	Phenylobacterium
Caenimonas	Corallococcus	Cokeromyces	Gammaproteobacteria_unclassified	Polaromonas
Canariomyces	Curtobacterium	Colletotrichum	Gemmataceae_unclassified	Pseudomonas
Candida	Cutibacterium	Comamonas	Gemmatirosa	Pseudorhodoferax
Candidatus Methylopumilus	Cylindrospermum	Coniophora	Geodermatophilaceae_unclassified	Rhizobium
Capnocytophaga	Dankookia	Coniosporium	GGB35955	Rhodoferax
Capronia	Desmonostoc	Cordyceps	GGB44695	Sphingobium
Caulobacter	Devosia	Cryptococcus	GGB52803	Sphingomonas
Ceraceosorus	Duganella	Cryptosporidium	GGB53655	Spirosoma
Cercospora	Dyadobacter	Cupriavidus	GGB64492	Tardiphaga
Chaetomium	Emticicia	Daldinia	GGB64512	Variovorax
Chlorella	Enterobacter	Deinococcus	GGB65415	Williamsia
Chromohalobacter	Enterococcus	Delftia	GGB68992	Xanthomonas
Cladophialophora	Erwinia	Diaphorobacter	GGB69028	Xylophilus
Cokeromyces	Flavobacterium	Diaporthe	GGB71872	
Colletotrichum	Friedmanniella	Dichotomopilus	GGB72163	
Comamonas	Frigoribacterium	Dickeya	GGB77784	
Coniophora	Frondihabitans	Dictyostelium	GGB78124	
Coniosporium	Fulvivirgaceae_unclassified	Didymosphaeria	GGB78370	
Cordyceps	Gammaproteobacteria_unclassified	Dioszegia	GGB79878	
Cryptococcus	Gemmataceae_unclassified	Dissoconium	Granulicella	
Cryptosporidium	Gemmatirosa	Drechmeria	Hungatella	
Cupriavidus	Geodermatophilaceae_unclassified	Drepanopeziza	Hyalangium	
Daldinia	Geodermatophilus	Dyella	Hyphomicrobiales_unclassified	
Dankookia	GGB35955	Eimeria	Ideonella	
Deinococcus	GGB44695	Emericellopsis	Inquilinus	
Delftia	GGB52803	Endocarpon	Intrasporangiaceae_unclassified	
Devosia	GGB53655	Epidermidibacterium	Janibacter	
Diaphorobacter	GGB64492	Exophiala	Kamptonema	
Diaporthe	GGB64512	Exserohilum	Kibdelosporangium	
Dichotomopilus	GGB65415	Falsiroseomonas	Kineococcus	
Dickeya	GGB68992	Ferranicluibacter	Kineosporia	
Dictyostelium	GGB69028	Fomitopsis	Kocuria	
Didymosphaeria	GGB71872	Fusarium	Larkinella	
Dioszegia	GGB72163	Gaeumannomyces	Leifsonia	
Dissoconium	GGB77784	Geminella	Luteibacter	
Drechmeria	GGB78124	Gemmobacter	Marmoricola	
Drepanopeziza	GGB78370	Glaciihabitans	Methylopila	
Duganella	GGB79878	Gynurincola	Methylorubrum	
Dyadobacter	Granulicella	Hafniomonas	Microbacteriaceae_unclassified	

Dyella	Hansschlegelia	Halomonas	Microcoleus
Eimeria	Herbaspirillum	Hartmannibacter	Microlunatus
Emericellopsis	Herbiconiux	Herminiimonas	Microvirga
Endocarpon	Hungatella	Heterobasidion	Modestobacter
Epidermidibacterium	Hyalangium	Homo	Myxococcus
Exophiala	Hymenobacter	Hydrogenophaga	Naasia
Exserohilum	Hyphomicrobiales_unclassified	Hyphopichia	Nakamurella
Falsiroseomonas	Ideonella	Hypoxylon	Nevskia
Ferranicluibacter	Inquilinus	Ilyonectria	Nibricoccus
Flavobacterium	Intrasporangiaceae_unclassified	Janthinobacterium	Nitrososphaerales_unclassified
Fomitopsis	Janibacter	Kalmanozyma	Nitrosospira
Fusarium	Kamptonema	Kickxella	Nocardioides
Gaeumannomyces	Kibdelosporangium	Labilibacter	Noviherbaspirillum
Geminella	Kineococcus	Labilithrix	Ohtaekwangia
Gemmobacter	Kineosporia	Lasiodiplodia	Paludisphaera
Geodermatophilus	Klebsiella	Lasiosphaeria	Panacagrimonas
Glaciihabitans	Kocuria	Lentinula	Pantoea
Gynurincola	Larkinella	Leptographium	Patulibacter
Hafniomonas	Leifsonia	Letharia	Pedococcus
Halomonas	Luteibacter	Leucobacter	Perlucidibaca
Hansschlegelia	Marmoricola	Ligilactobacillus	Phormidesmis
Hartmannibacter	Massilia	Lipomyces	Pseudarthrobacter
Herbaspirillum	Mesorhizobium	Lithohypha	Pseudonocardia
Herbiconiux	Methylobacterium	Litorilituus	Pseudorhizobium
Herminiimonas	Methylophilus	Malassezia	Pseudosporangium
Heterobasidion	Methylopila	Martelella	Pseudoxanthomonas
Homo	Methylorubrum	Maudiozyma	Rhizobacter
Hydrogenophaga	Microbacteriaceae_unclassified	Meira	Rhizobiaceae_unclassified
Hymenobacter	Microbacterium	Melampsora	Rhodanobacter
Hyphopichia	Microcoleus	Mesoflavibacter	Rhodanobacteraceae_unclassified
Hypoxylon	Microlunatus	Metarhizium	Rhodococcus
Ilyonectria	Microvirga	Metarhizobium	Rhodocytophaga
Janthinobacterium	Modestobacter	Methylicorpusculum	Rhodoplanes
Kalmanozyma	Mucilaginibacter	Methylobacillus	Rhodospirillaceae_unclassified
Kickxella	Mycolicibacterium	Methylogaea	Rhodospirillales_unclassified
Klebsiella	Myxococcus	Methyloradius	Roseococcus
Labilibacter	Naasia	Methylotenera	Roseomonas
Labilithrix	Nakamurella	Methylovorus	Saccharibacillus
Lasiodiplodia	Neorhizobium	Microdochium	Sanguibacter
Lasiosphaeria	Nevskia	Micromonospora	Serinibacter
Lentinula	Nibricoccus	Mixia	Simulacricoccus
Leptographium	Nitrososphaerales_unclassified	Mollisia	Skermanella

Letharia	Nitrosospira	Mucor	Solirubrobacter
Leucobacter	Nocardioides	Mycoavidus	Sphingobacteriaceae_unclassified
Ligilactobacillus	Nostoc	Mycobacterium	Stenotrophomonas
Lipomyces	Noviherbaspirillum	Nannizzia	Synechocystis
Lithohypha	Novosphingobium	Nannocystis	Tahibacter
Litorilituus	Ohtaekwangia	Neocystis	Tolypothrix
Malassezia	Paludisphaera	Neurospora	Urbifossiella
Martelella	Panacagrimonas	Niabella	Verrucomicrobium
Massilia	Pantoea	Nocardia	Virgisporangium
Maudiozyma	Paraburkholderia	Nocardiopsis	
Meira	Patulibacter	Nonomuraea	
Melampsora	Pedobacter	Oscillibacter	
Mesoflavibacter	Pedococcus	Paenibacillus	
Mesorhizobium	Perlucidibaca	Paeniroseomonas	
Metarhizium	Phenylobacterium	Pandoraea	
Metarhizobium	Phormidesmis	Paracidovorax	
Methylicorpusculum	Polaromonas	Paracoccus	
Methylobacillus	Pseudarthrobacter	Paraphaeosphaeria	
Methylobacterium	Pseudomonas	Pararhizobium	
Methylogaea	Pseudonocardia	Parietochloris	
Methylophilus	Pseudorhizobium	Penicilliopsis	
Methyloradius	Pseudorhodoferax	Penicillium	
Methylotenera	Pseudosporangium	Pestalotiopsis	
Methylovorus	Pseudoxanthomonas	Photorhabdus	
Microbacterium	Rhizobacter	Phyllobacterium	
Microdochium	Rhizobiaceae_unclassified	Phytophthora	
Micromonospora	Rhizobium	Pleomorphomonas	
Mixia	Rhodanobacter	Podospora	
Mollisia	Rhodanobacteraceae_unclassified	Polychytrium	
Mucilaginibacter	Rhodococcus	Porogramme	
Mucor	Rhodocytophaga	Priceomyces	
Mycoavidus	Rhodoferax	Pseudophaeobacter	
Mycobacterium	Rhodoplanes	Pseudoroseomonas	
Mycolicibacterium	Rhodospirillaceae_unclassified	Pseudovirgaria	
Nannizzia	Rhodospirillales_unclassified	Pseudozyma	
Nannocystis	Roseococcus	Psychrobacter	
Neocystis	Roseomonas	Puccinia	
Neorhizobium	Saccharibacillus	Purpureocillium	
Neurospora	Sanguibacter	Pyrenophora	
Niabella	Serinibacter	Pyricularia	
Nocardia	Simulacricoccus	Rahnella	
Nocardiopsis	Skermanella	Ramlibacter	
Nonomuraea	Solirubrobacter	Recurvomyces	

Nostoc	Sphingobacteriaceae_unclassified	Rhizophagus
Novosphingobium	Sphingobium	Rhodoalgimonas
Oscillibacter	Sphingomonas	Rhodofomes
Paenibacillus	Spirosoma	Rhodophyticola
Paeniroseomonas	Stenotrophomonas	Rhodopseudomonas
Pandoraea	Synechocystis	Rhodotorula
Paraburkholderia	Tahibacter	Roseicella
Paracidovorax	Tardiphaga	Rosenbergiella
Paracoccus	Tolypothrix	Saxophila
Paraphaeosphaeria	Urbifossiella	Scheffersomyces
Pararhizobium	Variovorax	Schizophyllum
Parietochloris	Verrucomicrobium	Shewanella
Pedobacter	Virgisporangium	Shimia
Penicilliopsis	Williamsia	Shinella
Penicillium	Xanthomonas	Siccirubricoccus
Pestalotiopsis	Xylophilus	Sinorhizobium
Phenylobacterium		Sphaerulina
Photorhabdus		Sporisorium
Phyllobacterium		Sporothrix
Phytophthora		Stanieria
Pleomorphomonas		Streptomyces
Podospora		Sugiyamaella
Polaromonas		Suillus
Polychytrium		Taibaiella
Porogramme		Talaromyces
Priceomyces		Terriglobus
Pseudomonas		Tetrahymena
Pseudophaeobacter		Thalassotalea
Pseudorhodoferax		Thermomonospora
Pseudoroseomonas		Thermothelomyces
Pseudovirgaria		Thermothielavioides
Pseudozyma		Thiothrix
Psychrobacter		Tilletiopsis
Puccinia		Trametes
Purpureocillium		Trichoderma
Pyrenophora		Trichophyton
Pyricularia		Trichosporon
Rahnella		Truncatella
Ramlibacter		Tuber
Recurvomyces		Ustilaginoidea
Rhizobium		Ustilago
Rhizophagus		Vanrija
Rhodoalgimonas		Verticillium

Rhodoferax	Vibrio
Rhodofomes	Winogradskyella
Rhodophyticola	Xenophilus
Rhodopseudomonas	Zavarzinia
Rhodotorula	
Roseicella	
Rosenbergiella	
Saxophila	
Scheffersomyces	
Schizophyllum	
Shewanella	
Shimia	
Shinella	
Siccirubricoccus	
Sinorhizobium	
Sphaerulina	
Sphingobium	
Sphingomonas	
Spirosoma	
Sporisorium	
Sporothrix	
Stanieria	
Streptomyces	
Sugiyamaella	
Suillus	
Taibaiella	
Talaromyces	
Tardiphaga	
Terriglobus	
Tetrahymena	
Thalassotalea	
Thermomonospora	
Thermothelomyces	
Thermothielavioides	
Thiothrix	
Tilletiopsis	
Trametes	
Trichoderma	
Trichophyton	
Trichosporon	
Truncatella	
Tuber	
Ustilaginoidea	
Ustilago	

| Vanrija |
| Variovorax |
| Verticillium |
| Vibrio |
| Williamsia |
| Winogradskyella |
| Xanthomonas |
| Xenophilus |
| Xylophilus |
| Zavarzinia |

417 **Supplemental Figure S1.** Microbiome composition of samples of *C. californica* at the family
418 taxonomic level.

419
420
421
422 **Supplemental Figure S2.** Dendrograms reflecting the clustering of samples by the similarity of
423 their microbiome composition (left) and by the similarity of the liverwort genetics (right).
424
425

Microbiome taxa composition   **Liverwort genetic composition**

426
427

Figure legends

Figure 1. **a**. Collection locations of *Calasterella californica* for this study. **b**. Typical environment

and morphology of *C. californica*. **c**. Close up to the female thalli of *C. californica*.

Figure 2. Relative abundance (in percentage of bacterial reads identified) of different bacteria

families across *C. californica* samples.

Figure 3. Correlation of pairwise distances between microbiome composition and three

potentially explanatory variables. **a.** Geographic distance, **b.** Environmental distance, and **c.**

Genetic distance.

Figure 4. Mean thallus width by proportion of Hyphomicrobiales for *Calasterella* samples. A line

of best fit was added using a linear regression model (standard error in grey).