A non-invasive approach to awake mouse fMRI compatible with multi-modal techniques

Sam Laxer[1,2], Amr Eed[1], Miranda Bellyou[1], Peter Zeman[1], Kyle M Gilbert[1,2], Mohammad Naderi[1], Ravi S Menon[1,2]

[1]Centre for Functional and Metabolic Mapping, Western University, London, Ontario, Canada
[2]Department of Medical Biophysics, Western University, London, Ontario, Canada

Abstract

Mouse functional magnetic resonance imaging (fMRI) studies contribute significantly to neuroimaging research, playing an important role in advancing our understanding of the human brain. Efforts to scan mice in the awake state and avoid the use of anesthesia have demonstrated many benefits. However, the majority of current awake fMRI approaches involve head fixation through invasive surgical means. These approaches are limited for several reasons including increased subject morbidity and mortality, prolonged study timelines due to surgical recovery, a higher risk of susceptibility artifacts contaminating T2*-weighted images and obstructed access to the head for multi-modal recording techniques such as electrophysiology and fibre photometry. To address this need, we propose a non-invasive restraint to scan mice in the awake state. The restraint was designed to be compatible with brain stimulation and recording approaches. It was evaluated on the basis of motion, fMRI data quality, and animal stress levels, and subsequently compared to a conventional headpost restraint. We found the proposed approach was effective at restraining mice across a broad range of weights without the need for any anesthesia for setup. Conservatively, acquiring 25% more data with the non-invasive restraint compensates for censored frames in high motion runs and results in comparable network spatial specificity to less motion-vulnerable headpost approaches. This validation of network robustness paired with its non-invasive nature makes it a favourable approach for applications requiring high-throughput and/or multimodal imaging studies.

1 Introduction

Functional magnetic resonance imaging (fMRI) is an important tool in neuroscience for characterizing brain-wide networks in healthy animals and disease models. Mice have been particularly beneficial due to the wide range of available transgenic lines and the high translational value due to their similar neurophysiology compared to humans (Mandino et al., 2020, 2021; N. Xu et al., 2022) . However, motion needs to be minimized to acquire useful data (Power et al., 2012). To control motion, there are currently 3 main scanning approaches: 1) using anesthesia, 2) scanning awake using an invasive restraint, and 3) scanning awake using a non-invasive restraint.

Anesthesia in mice has been common practice in fMRI research due to its effectiveness, however an increasing number of studies have shown its confounding effects on brain hemodynamics and networks (Dinh et al., 2021; Gao et al., 2017; Gutierrez-Barragan et al., 2022). These include damped and lagged hemodynamic responses compared to the unanesthetized brain and altered functional connectivity. Therefore, when the research goal is understanding physiology or maintaining a high translational value, scanning mice in an awake state provides considerable advantages.

This motivated the development of awake invasive approaches, often utilizing surgically implanted headposts as a fixation point on the skull (Chen et al., 2020; Gutierrez-Barragan et al., 2022). This approach avoids the negative effects of anesthesia (after the initial surgery) and in some cases allows for comparable motion control to anesthesia. Despite these benefits, the use of a headpost poses challenges in many applications. These include: 1) the headpost obstructing access to the skull when simultaneous multimodal recording is desired (such as combining fMRI with fibre photometry, optogenetics, or electrophysiology), 2) vulnerable or highly valuable mouse lines being unable or posing too great of a risk to undergo headpost implantation surgery (Szilagyi et al., 2018; Wu et al., 2020), 3) the headposts, cement or associated surgical procedures causing MRI distortion artifacts, or 4) having high-throughput studies with a very large number of subjects makes surgical approaches impractical. One solution is to develop non-invasive awake imaging approaches that forego the use of a headpost while maintaining awake imaging.

Non-invasive restraints have been developed in the past (Fadel et al., 2022; Ferris, 2022; Harris et al., 2015; Madularu et al., 2017) and are quite common for rat studies. However, the current mouse systems suffer from tedious animal setup procedures often requiring the brief administration of anesthesia to position the mice and some holders are not compatible with the small inner diameter size of high-performance gradient inserts. The success of rat systems is due to the larger size of the animal and their better ability to be trained compared to mice (Ellenbroek & Youn, 2016; Parker et al., 2014). Thus, there remains a need for a simple, effective non-invasive restraint ideally requiring no anesthesia for animal setup.

Another barrier concerning awake imaging approaches is stress. Animal stress has been investigated and shown to impact hemodynamics and results in stress-related motion, while also imposing a negative impact on the animal's wellbeing (Low et al., 2016). Consequently, a large effort has been made to develop restraint features and associated procedures to minimize stress. To this end, various labs have developed acclimation procedures to gradually expose subjects to restraint and the scanning environment (i.e., dark and loud environment mimicking the auditory and visual experience of the scanner) (Chen et al., 2020; Desjardins et al., 2019; Dinh et al., 2021; Gutierrez-Barragan et al., 2022; Takata et al., 2018). Additionally, care has been taken to design restraints such that subject comfort is improved to further reduce stress (W. Xu et al., 2022).

Here, we present an approach to non-invasive awake-mouse MRI with a focus on
compatibility with multi-modal experiments – all while remaining small enough to fit inside the
60-mm-diameter gradient coil found in many small bore animal MRI scanners. The developed
restraint allows for effective motion control and no need for anesthesia during animal setup.
Motion and dual-regression-based resting state network robustness were evaluated in
comparison with a typical headpost restraint approach. Lastly, stress was analyzed by
measuring weight and corticosterone changes between three different acclimation procedures
in both sexes.

2 Methods

2.1 Ethics Statement

All experimental methods described were performed in accordance with the guidelines of the Canadian Council on Animal Care policy on the care and use of experimental animals and an ethics protocol approved by the Animal Care Committee of The University of Western Ontario.

2.2 Subjects and Groups

This study involved 40 wildtype C57BL/6 mice from Jackson Laboratories (n = 13) and Charles River Laboratories (n = 28). Mice were divided into 4 groups (Table 1) with each group having roughly equally proportions of males and females. Groups 1, 2 and 3 underwent the non-invasive imaging protocol (with differing acclimation lengths) and Group 4 underwent a conventional headpost imaging protocol with the same acclimation protocol as the non-invasive 9-day protocol as shown in Table 2. Animals in Groups 1, 2, and 3 were single-housed in a facility with a 12-hour light/dark cycle (light from 7:00 am – 7:00 pm) and fed food and water ad-libitum. Animals were single-housed to replicate the treatment for cannula-implanted mice who are single-housed to minimize the risk of mouse cage mates damaging/removing each other's cannulas. Mice in Group 4 (acquired previously) were group-housed.

Table 1. Subject information. The first 4 rows display the sample sizes in each group (m = males, f = females).

Parameter	Value
Group 1 (Non-invasive – 4-day protocol)	4 m + 5 f
Group 2 (Non-invasive – 9-day protocol)	4 m + 5 f
Group 3 (Non-invasive – 13-day protocol)	7 m + 8 f
Group 4 (Headpost – 9-day protocol)	4 m + 3 f
Strain	C57BL/6
Age range	2 – 35 weeks
Weight range	16 – 37 g

2.3 Restraint designs

The non-invasive restraint is almost entirely 3D printed of polycarbonate (Figure *1*). It consists of a cylindrical body tube, a neck plate to keep the subject in position, and the combination of a bite bar and two head arches (one anterior at the snout and one posterior at the caudal end of the skull) to stabilize the head in all three dimensions. An in-house built single-loop transmit/receive coil was designed to fit over top of the head after the arches were secured.

Figure 1. CAD models of the non-invasive restraint (left) and headpost restraint (right). The single loop coil used for single transmit/receive is not shown on the non-invasive restraint to allow clear visualization of the arches over the mouse head.

The headpost-style restraint is also 3D printed using polycarbonate. It was initially inspired by a headpost designed by (Gutierrez-Barragan et al., 2022), with further modifications. Rather than permanently implanting the large 'head-bar wing', we implanted a small threaded collar on the skull to allow for the attachment of the 'head-bar wing' when the mouse is to be restrained. The motivation was to provide less of an impediment to the natural movement of a mouse in their cage when unrestrained.

All designs have been made freely available (https://doi.org/10.17605/OSF.IO/TYHCE).

2.4 Surgery

Mice in group 4 underwent surgery to have headposts implanted on their skulls. Mice were initially anesthetized with 4% isoflurane in oxygen and then switched to a 1.7 – 2 % mixture for maintenance. Once deeply anesthetized (confirmed using toe pinch), mice were injected with 0.05 mL/kg of Meloxicam to reduce postoperative pain and 0.05 mL/kg of Lidocaine to numb the surgical area. The hair on the scalp was removed with an electronic razor, eye ointment was applied, and a rectal thermometer was used to record temperature throughout the surgery. The mouse was placed in a stereotactic frame (71000 Automated Stereotaxic Instrument, RWD Life Science) and positioned with the skull level. After sterilizing the scalp, the skin above the skull was removed, and the soft tissue was cleared away. The skull was cleaned with saline and dried. The skull was then lightly scraped to provide a better surface for the bonding agent to adhere to. The bonding agent (All-Bond Universal, Bisco Dental) was applied to the skull and cured using an ultraviolet light. Dental cement (CoreFlo DC Lite, Bisco Dental) was subsequently applied, and the headpost was pressed onto the skull maintaining a high, constant pressure. The dental cement was cured, then the remaining exposed skull was covered with dental cement and cured. The mouse was then released from the stereotactic device and began a 10-day period of recovery prior to restraint acclimation. Additional injections of Meloxicam were given 24 and 48 hours after surgery.

2.5 Acclimation protocols

Three different acclimation protocols were used in the present study. Short (4-day; Group 1), medium (9-day; Groups 2 and 4) and long (13-day; Group 3) protocols were devised. The details for each protocol are shown in Table 2. These were designed to represent protocols typically used in recent literature (refer to (Mandino et al., 2023)).

Table 2. Protocol used to acclimate each Group. Group 1 underwent the 4-day protocol (left), Group 2 and the headpost group underwent the 9-day protocol (middle) and Group 3 underwent the 13-day protocol (right).

Day	Duration	Procedure
1	15 min	Manual handling (0 dB)
2	30 min	Restrained (0 dB)
3	45 min	Restrained (75 dB)
4	1 hr	Restrained (100 dB)

Day	Duration	Procedure
1	15 min	Manual handling (0 dB)
2	10 min	Restrained (0 dB)
3	20 min	Restrained (0 dB)
4	30 min	Restrained (50 dB)
5	45 min	Restrained (75 dB)
6	1 hr	Restrained (100 dB)
7	1 hr 15 min	Restrained (100 dB)
8	1 hr 30 min	Restrained (100 dB)
9	1 hr 30 min	Restrained (100 dB)

Day	Duration	Procedure
1	15 min	Manual handling (0 dB)
2	10 min	Restrained (0 dB)
3	15 min	Restrained (0 dB)
4	20 min	Restrained (50 dB)
5	30 min	Restrained (75 dB)
6	45 min	Restrained (75 dB)
7	45 min	Restrained (75 dB)
8	1 hr	Restrained (75 dB)
9	1 hr	Restrained (75 dB)
10	1 hr 15 min	Restrained (100 dB)
11	1 hr 30 min	Restrained (100 dB)
12	1 hr 30 min	Restrained (100 dB)
13	1 hr 30 min	Restrained (100 dB)

All acclimation sessions were started at approximately the same time each day to account for differences in circadian rhythm. Day 1 consisted of the same procedure for Groups 1, 2, and 3. Specifically, mice were introduced to the restraint and allowed to freely explore it for the first few minutes. For the remaining duration of the session, mice would be encouraged to walk through the restraint. This involved the researcher picking up the mouse by the tail and guiding the subject into the apparatus. After successfully walking through the tube, a chocolate sprinkle would be provided as a reward. Subsequently, the subject would be repeatedly encouraged to walk through the restraint to establish that movement. Importantly, day 1 does not involve restraint. Each of the following sessions starts with this exercise for the first few minutes to encourage the intended behaviour.

Restraining the mouse using the proposed non-invasive restraint involves 3 steps: 1) broad restraint, 2) bite bar, and 3) fine restraint. Step 1 consists of letting the mouse walk through the body tube as performed previously but stopping the subject after their head passes through the neck plate and carefully lowering the neck plate into position. This effectively prevents the subject from moving further forward or backward in the restraint. To gradually expose mice to being fully restrained, Day 2 of each session consists of keeping the mouse in this coarsely restrained position only. Day 3 of training onwards involves all 3 restraint steps. Step 2 of restraint uses the bite bar and anterior arch. This involves using the anterior arch to control the subject's snout and guide the top front teeth (incisors) into the bite bar groove. Once in position, the bite bar and anterior arch are secured in place with fastening screws. Finally, step 3 is implemented consisting of the posterior arch being positioned over the posterior aspect

175   of the subject's head. Effectively, the neckplate and body tube prevent broad motion, and the
176   combination of the bite bar, anterior arch and posterior arch prevent fine motion.
177
2.6 Mock scanner
It is common practice to design a "mock scanner" to simulate the scanning experience
(Dinh et al., 2021; Lindhardt et al., 2022). This has the added benefit of allowing for a more
gradual acclimation as subjects can be exposed to restraint and MRI sounds independently.
The mock scanner apparatus consists of a large box (65 cm wide, 52 cm long and 44 cm high)
with PVC pipes placed inside allowing for multiple animals to be acclimated simultaneously. A
Bluetooth speaker is also placed in the box to play MRI sounds at designated sound levels
(Table 2). The mock sounds were selected as the same sequences used to scan the subjects
during acquisition. The audio was recorded using a fibre optic microphone (OptiSLM,
OptoAcoustics Ltd) placed next to the magnet. Both the gradient-echo echo planar imaging
(GE-EPI) sequence and anatomical Turbo RARE sequence were recorded. The audio clips
were edited to create customized lengths such that each acclimation session included audio
from both sequences of increasing lengths and sound levels (refer to Table 1). Additionally,
sound-suppression foam was adhered to the inner surfaces of the mock scanner box to reduce
the noise that escaped the acclimation room (i.e., to prevent other studies in nearby spaces
from being affected). The foam is a multi-layer acoustic suppression medium consisting of a
high-density rubber base (approximately 2 mm) with a foam backing (approximately 5 mm).
195

2.7 MRI Acquisition
All acquisitions were performed at the Centre for Functional and Metabolic Mapping at
Western University. MRI scans were acquired on a 9.4 T, 31-cm horizontal-bore magnet
(Varian/Agilent, Yarnton, UK) with a 6-cm Magnex HD gradient insert and Bruker BioSpec
Avance NEO console running ParaVision 360 v3.3 (Bruker BioSpin Corp, Billerica, MA).
Groups 1, 2, and 3 had four 10-minute functional runs acquired per session (TR=1.5 s,
TE=12 ms, resolution=300x300x400 mm^3, slice gap=100 mm, matrix=64x32x30 voxels, 400
volumes). Following the functional scans, a T2-weighted Turbo RARE anatomical scan was
acquired (TR=5.5 s, TE=30 ms, resolution=150x150x400 mm^3, slice gap=100 mm, 192x96x30
voxels, flip angle=160 degrees). Group 4 was acquired independently and had slightly different
acquisition parameters. Namely, 2 functional runs were acquired per session with 600 volumes
each (15-min runs). All other parameters were the same.
208

2.8 fMRI Analysis
All fMRI data were analyzed using the Rodent Automated Bold Improvement of EPI
Sequences (RABIES version 0.5.1) (Desrosiers-Grégoire et al., 2024). All parameters used for
RABIES processing stages are provided in Supplementary Table 1. Images were motion
corrected using a trimmed mean across the acquired EPI frames in each run. They were then
registered to a common space using an unbiased template as an intermediate step to improve
registration (Avants et al., 2011). Image distortions due to susceptibility artifacts were corrected
by nonlinearly registering EPI frames to the corresponding subject's T2w anatomical image
(Wang et al., 2017). This is performed after first applying an inhomogeneity correction to all EPI
frames. Importantly, all transformations calculated during preprocessing (from motion
correction to registration and susceptibility correction) are concatenated and applied in a single
operation to avoid multiple stages of resampling. Final common space images were resampled
to 0.2 mm x 0.2 mm x 0.2 mm prior to further processing.

Frame censoring was carried out using a framewise displacement threshold of 0.15 mm.
All frames exceeding this value, along with the preceding frame and the two following frames
were removed (Power et al., 2012). If fewer than 200 volumes remained after censoring, the
run was excluded from further analyses. Following this, data were detrended on a voxel-wise
basis to remove linear drifts and converted to units of percent BOLD change through grand
mean scaling (i.e., dividing by the grand mean signal and multiplying by 100). The 6 motion
parameters calculated during motion correction were used as regressors to further clean the
data. Lastly, spatial smoothing (Abraham et al., 2014) was implemented using a full-width half
maximum value of 0.3 mm.
231
2.9 Motion Evaluation
Motion was estimated by registering EPI volumes and determining the displacements
for each run. Although typical motion analyses involve measuring the 6 rigid body parameters,
the voxel displacements between subsequent volumes are more important for data quality and
better predict motion artifacts. This is because low frequency motion can be much more readily
corrected for than high frequency spikes (Power et al., 2018; Soares et al., 2016). Here, we use
the mean voxel-wise framewise displacement (FD) metric as performed in RABIES. It is
calculated by taking the average of brain voxel displacements between each subsequent
volume. A more detailed description of this metric can be found in the RABIES documentation
(Desrosiers-Grégoire et al., 2024). This metric was used to compare motion levels between the
3 acclimatation protocols using the non-invasive restraint and the headpost restraint. The
Kruskal-Wallis test was performed to determine significant differences followed by Bonferroni-
corrected Wilcoxon rank sum test to determine which groups were significantly different.
245
2.10 Resting-State Network Analysis
Following preprocessing, fMRI data quality was analyzed to evaluate the performance
of the non-invasive restraint. Due to the subjective nature of evaluating the robustness of
resting-state networks, we decided to compare our data using an externally derived component
representing the well-established somatomotor network (Pagani et al., 2023). This was done
by running dual-regression on our data using the prior component and then using Permutation
Analysis of Linear Models (PALM) (Winkler et al., 2014) to attain group statistical maps
representing how much each group contains the prior component. The external networks were
determined from two datasets comprising data from several institutions with a combined sample
size of 366 animals (Desrosiers-Grégoire et al., 2024).
First, we compared the network differences between the 3 non-invasive restraint
acclimation protocols and the headpost group using the prior component. To ensure the same
amount of data was used, the total number of frames acquired in the headpost group was
calculated and the same quantity of data was randomly selected from each non-invasive group
(note this is prior to frame censoring). Group maps were then generated using PALM and z-
scores were shown. To evaluate network quality, spatial specificity of the maps was measured
using the Dice Similarity Coefficient (DSC) between the prior component with a z-threshold of
3.1 and the group map with a z-threshold of 3.0. The prior component threshold was chosen to
represent a p value of 0.001 and the group map z threshold was selected to achieve the best
overlap between the headpost group and the prior network. Multiple z values were tested to
determine this optimal value. This same value was used for the non-invasive groups to fairly
compare between groups. Additionally, the voxel-wise correspondence of z-values between the
group and prior networks were obtained using the Pearson correlation coefficient (r) in the

region of the network (by masking both prior and group networks by the voxels in the prior
network that were greater or equal to 3.1).
To understand the relative effects of motion and sample size on network specificity and
amplitude correspondence (i.e. the z-score of a voxel in the prior map vs the z-score of the
same voxel in the restraint group maps), the non-invasive data from each acclimation protocol
were aggregated and separated into 5 groups according to the amount of motion present
measured by the average run FD (namely motion group 1-5, Supplementary Table 2). Within
each motion group, runs would be randomly sampled to form groups with 30, 60, 90, 120, 150,
180, and 210 minutes of acquisition. This was accomplished through bootstrapping (randomly
sampling the data without replacement). For each group, 10 random samples were selected,
and PALM was used to attain group maps. Network specificity was again evaluated using the
DSC of the group map and the prior component map. Similar to the acclimation group
comparison, a z-value threshold of 3.1 was used for the prior component (representing a p
value of 0.001) and the z-value threshold of the group networks was 3.0. For each cumulative
acquisition time, a one-way ANOVA was used to determine if groups significantly differed from
one another. Tukey's test for multiple comparisons was then used to determine the specific
groups that significantly differed.
The effects of cumulative group acquisition time and motion on network spatial specificity
were further investigated by determining linear trendlines that represented the data in
Supplementary Figure 1. All functional runs were aggregated and ranked by their average
framewise displacement (FD) values, in ascending order of motion. Runs were grouped
sequentially, starting with pairs, and the corresponding group map for each pair was generated.
This map was then used to calculate the DSC. The process was repeated with progressively
larger group sizes, including triplets, quartets, and so on, up to groups of ten (Supplementary
Figure 1).
294
2.11 Stress response
Animal weight and video monitoring were used to assess stress of all mice throughout
acclimation and scanning. A subset of mice (n = 21) was used to quantitatively assess
corticosterone changes throughout the short (4-day), medium (9-day) and long (13-day)
acclimation sessions. In contrast to the conventional extraction of CORT through blood serum
which may cause a confounding stress response (Eleftheriou et al., 2020; Rowland & Toth,
2019; Russo et al., 2021), we employed extraction from fecal pellets similar to (Russo et al.,
2021) This was performed by collecting the fecal pellets defecated during each session. CORT
was extracted and the concentrations were determined using an ELISA following the supplier's
manual (Arbor Assays DetectX Corticosterone Chemiluminescent Immunoassay Kit, catalog
number K014-C). All concentrations were measured in duplicate and are expressed as ng of
CORT per mg of fecal matter.
Corticosterone was analyzed on days 1, 2, and 4 for the 4-day protocol, days 1, 2, 4,
and 9 for the 9-day protocol, and days 1, 2, 4, 9, and 13 for the 13-day protocol. All samples
were read from a luminescence-capable plate reader (SpectraMax® M5 Multimode Plate
Reader, Molecular Devices, USA). One-way ANOVA was used to compare CORT
concentrations in each acclimation group. No corticosterone analysis was performed on the
headpost group as it was a dataset acquired previously for a separate study. However,
corticosterone data from head fixed animals using similar headposts has been obtained by
others (Lindhardt et al., 2022).

3 Results

3.1 Motion Evaluation

We first analysed scan motion levels to determine if motion depends on the acclimation protocol used (Figure 2). The framewise displacement for each voxel was measured and the average across each run was computed. Figure *2* (a) shows a representative motion plot with the 6 rigid body parameters (3 translations – top, and 3 rotations – middle). The corresponding mean voxel-wise displacements are shown below in Figure *2* (a). A histogram is plotted in Figure *2* (b) showing all FD measures for all acquired volumes. This metric was then used in Figure *2* (c) to compare motion between cohorts. Motion was significantly higher in the 4-day group than the 9-day group, and motion was significantly lower in the headpost group than all other groups. Note, after censoring, 12.5% of the total number of acquired volumes were censored and 19 of the 226 runs were excluded due to being left with fewer than 200 volumes after censoring. A comprehensive comparison of the censored volumes based on group can be found in Supplementary Table *2*.

Figure 2. Scan motion levels for the different restraints and acclimation protocols. (a) Representative plot displaying the motion parameters (3 translations – top, 3 rotations – middle, and the corresponding mean voxel-wise framewise displacements – bottom). The plot is expressed in FD magnitude (left y-axis) and as a fraction of the in-plane voxel size (300 μm – right y-axis). (b) A histogram displaying mean voxel-wise FD values for all volumes in a run (e.g., the mean value of FD values displayed in (b) bottom) for all subjects. The red vertical line indicates the in-plane voxel size (300 μm). (c) Boxplots comparing the motion between the four cohorts. Outliers are shown as red '+'. The mean run voxel-wise FD values are calculated by taking the average of brain voxel displacements between each subsequent volume (* = p < 0.05, ** = p < 0.01, *** = p < 0.001, Kruskal-Wallis rank sum test followed by Bonferroni-corrected Wilcoxon rank sum test).

3.2 Resting-State Networks

To avoid the bias of using components generated from this dataset and promote
standardization across centres, dual regression was performed using previously generated
components through independent component analysis that established well-known resting-
state networks (Desrosiers-Grégoire et al., 2024). To confirm the networks are recoverable in
our datasets, dual regression of the somatomotor network was performed for each scan and
data were grouped by acclimation group (Figure *3*). The DSC representing the amount of
overlap and correlation coefficient representing the amplitude correspondence between the
networks is overlaid above each network. Note, only a sample slice is shown however the DSC
and correlation coefficients were calculated from the entire volume.

349
350

351
Figure 3. Network quality comparison of the 3 non-invasive protocols and the headpost group using the Dice Similarity
Coefficient (DSC) and amplitude correspondence (using the Pearson correlation coefficient, r) with the externally derived
somatomotor component (Desrosiers-Grégoire et al., 2024). Note, DSC and correlation values were calculated using the
whole volume but only one sample slice is shown. The top left network map shows the externally derived component. The
top right shows the contribution in the headpost group overlayed on top of the prior component. The bottom row shows the
same comparison except for the 4-Day, 9-Day and 13-Day groups, respectively. Corresponding DSC and r values are
annotated over each network.

359
360

We further investigated the relationship between motion, sample size and network
quality by looking at how varying levels of motion and cumulative group acquisition time affect
quality (i.e., DSC and amplitude correlation) relative to the prior network component (Figure *4*).
Refer to Supplementary Figure *2* to see the resulting correlation results for a single sample
from each motion level and cumulative acquisition time. Past 120 min of acquired data, both
DSC and amplitude correlation analyses show clear statistical differences between headpost

and non-invasive groups. Taking this a step further, all functional runs were aggregated and
ranked by their average framewise displacement (FD) values, in ascending order of motion.
Runs were grouped sequentially, starting with pairs, and the corresponding group map for each
pair was generated. This map was then used to calculate the DSC. The process was repeated
with progressively larger group sizes, including triplets, quartets, and so on, up to groups of ten.

372
Figure 4. Group network quality assessment for the headpost and aggregated non-invasive data relative to the prior network
component. Groups have varying levels of motion and cumulative acquisition time. (a) Sample slices of the z-maps of the
prior component (z values between 3.1 and 20 illustrated in green/brown) and the group network (z values of 3.0 to 7.5
illustrated in red/blue). Motion Level 1 indicates the lowest amount of motion and Motion Level 5 indicates high motion. The
range of average FD values for each group are displayed below each Motion Level on the left. (b) Corresponding Dice
Similarity Coefficient (DSC) values to the network overlaps shown in (a). (c) Correspondence of z-values compared voxel-
wise between the prior map and the group map masked to include voxels in the prior network region. Error bars in (b) and (c)
represent the standard deviation of 10 random samples selected from the respective group. Groups that significantly differed
from the headpost group for each cumulative acquisition time are shown (* = p < 0.05, ** = p < 0.01, *** = p < 0.001, for
Tukey's test for multiple comparisons following one-way ANOVA).

3.3 Stress
Mouse stress was approximated using weight loss and fecal corticosterone
concentration over the course of the acclimation sessions. Weight was measured each day of

acclimation. Values were normalized to the weight on day 1 and plotted for each cohort
separately (Figure *5* (a) and (b)). Paired t-tests were conducted to test for statistical differences
between first and last days of acclimation for each group. Headpost, 9-day and 13-day groups
all had significant weight losses. However, the headpost group only showed a significant
difference when both sexes were aggregated. When males and females were analyzed
separately, no significant difference was found (Supplementary Figure 3). Fecal corticosterone
metabolite (FCM) was measured from a subset of mice from each non-invasive group. Results
are shown in Figure *5* (c) and Supplementary Figure 3 (c) and (d). Note, the headpost group
was acquired previously and fecal samples were not collected. One-way ANOVA was
conducted. Paired t-tests could not be performed since several groups had missing samples
(due to the subject not defecating during the respective acclimation session). There was no
significant difference in corticosterone levels between sampling days across the acclimation
protocols.
400

4-Day Protocol   **9-Day Protocol**   **13-Day Protocol**

Figure 5. Weight and fecal corticosterone changes across acclimation sessions. (a) Weight changes for all subjects (n = 7) in
the headpost group normalized to the weight on day 1. The weight on the last day is significantly less than the first day
(paired t-test, p < 0.05). (b) Weight changes of all subjects in the 4-day, 9-day and 13-day groups over the course of the
different acclimation lengths normalized to the weight on day 1. Note the headpost group underwent the same 9-day

acclimation session as the 9-day non-invasive group. Both the 9-day group and 13-day group showed significant weight
decreases (paired t-test, $p < 0.01$). (c) Corticosterone concentration changes over the course of the three acclimation
durations. Note, the headpost group was acquired in a separate study and corticosterone samples were not collected. There
were no significant CORT changes across acclimation sessions (one-way ANOVA, $p > 0.05$).

4 Discussion

All subjects used in the current study successfully used the restraint and accompanying acclimation protocols. The restraint contains several adjustable features to tailor the fit for each mouse regardless of size (i.e., 16 – 37 g as tested here). Similar to previous non-invasive restraints (Madularu et al., 2017), our restraint is 3D printed and modular in nature allowing for easy modification. Furthermore, no anesthesia is required for setup and experienced handlers can position acclimated mice in less than 5 min.

Three different acclimation procedures were developed representing a range of protocols typically found in recent awake mouse MRI literature (see (Mandino et al., 2023; Zeng et al., 2022) for comprehensive acclimation protocol comparisons). Based on our motion analysis, mice in the 4-day protocol did move significantly more than the 9-day group. Indeed, 20.6% of volumes were censored and 13.9% of runs were excluded due to being above the motion threshold (Supplementary Table 2). Even so, the network specificity between the 4-day and 9-day groups were almost identical at 0.64 and 0.65, respectively. Similarly, the 13-day group also had a spatial specificity of 0.65 and did not significantly differ in motion compared to the other two groups.

Non-invasive restraints are more vulnerable to residual motion compared to headpost restraints, since the latter directly fixates the skull. However, we have found the impact of the motion confound reduces tremendously with increased sample size as demonstrated in Figure *4*. Since motion tends to affect a relatively constant proportion of data, an increased sample size will allow for increased statistical power after censoring of motion-affected frames. Even at the extreme end, in untrained young human subjects, Smith and colleagues evaluated how harmful this motion is to data quality using similar metrics outlined here. In the end, they conclude a conservative design would be to acquire 2 to 3 times more data than would be expected for the same study without high-motion subjects (Smith et al., 2022). Our data suggest a much lower quantity of data is required. Despite allowing more than 3x the motion than the headpost cohort (Figure *2* (c)), acquiring 25% more data with the non-invasive restraint will compensate for censored volumes assuming most motion is at the high end allowed by the restraint.

The effects of motion are only seen past 50 min of cumulative group acquisition time. From Supplementary Figure 1, this is evident from the emergence of the negative slope of the linear trendlines. The slope only becomes noticeable in the 50 min group and above. The groups with less cumulative acquisition time contain other sources of variation such that the effects of motion are not apparent. Individual network variability can cause such variation (Bergmann et al., 2020; Fu et al., 2023). Similarly, in Figure 4 the effects of motion are only seen at 90 min as can be seen by the lower spatial specificity for motion groups 4 and 5 – though it is only apparent between the headpost group and the lower motion groups at higher amounts of data when the other sources of variation are presumably averaged out.

Sample size is a current topic of controversy among animal imaging researchers (Grandjean et al., 2024; Mandino et al., 2023). There is motivation from an ethics perspective to use smaller sample sizes, whereas from a statistical perspective, large sample sizes are preferred. Fortunately, here the slightly higher motion corrupted data benefits the same amount from increased group cumulative acquisition time compared to the headpost restraint (Figure *4*, Supplementary Figure 1). For instance, past 2 hours of cumulative acquisition time, all groups have a spatial specificity above 0.5 (Figure 4 (b)) and an amplitude correspondence greater than 0.7 (Figure 4 (c)). The results indicate the amount of data is much more important than the amount of motion during the acquisition. Indeed, with long cumulative acquisition times

(approaching 210 min), the spatial specificity difference between motion groups seems to be
solely based on the number of frames censored. It is possible higher amounts of motion (i.e.,
greater than the in-plane voxel size) would further degrade the networks, but that was not
investigated here as almost all (98.7%) displacements were under the in-plane voxel size
(Figure *2* (b)). Surprisingly, we do not see this same relationship when looking at the amplitude
correspondence. Approaching higher cumulative group acquisition times, the headpost group
emerges with a statistically significant increase over all non-invasive motion level groups,
however no clear relationship emerges among motion level groups like we see in the spatial
specificity analyses. This suggests that this amplitude correspondence metric is independent
of motion, at least at the levels observed here. However, increased cumulative acquisition time
still increases correlation values of all groups indicating differences between the headpost and
non-invasive restraint can be addressed by simply acquiring more data.
Stress has been shown to be a critical measure for awake fMRI as it has been shown to
impact functional connectivity (Grandjean et al., 2016; Maron-Katz et al., 2016). There is a
statistically significant weight loss between the first and last days of acclimation in the headpost
group, and the 9-day and 13-day non-invasive groups. Although no loss is observed in the 4-
day group, this is likely because the 4-day duration is too short to observe the change. Although
these weight losses are statistically significant, they remain below the critical threshold of 15%,
which is considered a concern for animal welfare by most animal care committees.
FCM was measured and compared for several selected days throughout each non-
invasive acclimation group. Specifically, fecal samples were used over serum samples to
minimize unrelated stress to the animals (Kroll et al., 2021). For the same reason, day 1 was
selected as our baseline as it is well known that CORT is only detectable in feces hours after
the stressful event (Rowland & Toth, 2019). It is also sensible to assume the FCM measurement
on Day 1 is in fact a baseline measurement because 1) any CORT would not be detectable
within 20 min (maximum duration between stress event onset and fecal excretion) of the
stressor, and 2) the procedure on day 1 is the least stressful and likely would not result in any
significant increase in CORT (e.g., Lindhardt et al. demonstrated that even a more intensive
protocol involving 20 minutes of restraint on Day 1 did not result in a significant increase in
corticosterone (Lindhardt et al., 2022)).
Surprisingly, no significant change in corticosterone concentration was observed in any
of the acclimation groups. Consequently, these results would indicate no relationship between
acclimation session and stress in mice and stress would not explain the observed difference in
motion between the 4-day and 9-day acclimation groups. This is an unexpected result
compared to the literature (Ferris et al., 2011; Gutierrez-Barragan et al., 2022; Lindhardt et al.,
2022; Low et al., 2016). This could be due to the measurement of FCM rather than more acutely
sensitive measures of CORT such as through plasma samples. FCM is known to integrate the
CORT levels over relatively long periods of time (hours to days) (Rowland & Toth, 2019). It is
possible our maximum 1.5-hour stressor did not produce a detectable CORT increase over the
48-hour integration time. (Lindhardt et al., 2022) used fecal corticosterone measurements and
did find a difference on day 11 of acclimation compared to day 1. However, they used a more
aggressive acclimation protocol than used here, therefore it is possible the animals showed a
larger stress increase on the first few days. Those animals also underwent surgery which
imposes an additional variable to consider.
Due to the results of this study, along with indications from related literature, the authors
recommend future corticosterone experiments be acquired over longer durations each session.
That is to say that single time-point measurements of CORT can be misleading and are difficult
to interpret. Further, it should be noted that it is still not fully understood what increases in

corticosterone represent and indeed, corticosterone levels naturally vary widely dependent on
many factors (Rowland & Toth, 2019). While it is generally thought that increases correspond
with increases in negative stress, it has also been postulated that elevations in HPA and basal
glucocorticoid levels may be linked with stress adaptation, a positive outcome rather than
excess stress (Rowland & Toth, 2019). More stress-targeted studies are required to both
characterize and better understand the relationship of corticosterone concentration on
perceived stress.
513

4.1 Limitations

514
Like all T2*-weighted sequences, care should be taken to minimize the effect of all major
sources of susceptibility artifacts. This includes avoiding the use of magnetic materials and
avoid pockets of air near tissue. One such source relevant to the proposed restraint was the
susceptibility artifact caused by the ears of the mouse. Since the posterior arch can fold the
ears onto the head, care should be taken to ensure the ears are folded towards the posterior
and lateral regions of the head rather than anterior and medial.
Another limitation is only one network (i.e., cortical somatomotor network) was used for
network spatial specificity and amplitude correspondence measures. It is possible other cortical
networks or even subcortical networks would have different results.
Although the restraint is designed to allow for tasks to be readily performed, task-based
imaging was not conducted in the current study. It can be postulated that tasks may cause task-
induced motion, the effects of which are amplified in restraints allowing for increased baseline
levels of motion. However, since resting-state networks were clearly identifiable, it is likely task-
based networks will only be clearer due to the higher activation and statistical power associated
with task-based protocols (Zhao et al., 2023). An additional limitation of this work is it does not
consider intra-volume motion. Consequently, acquisitions with large movements that occur and
return to the original position during the same TR may not be robustly corrected and can also
have degrading effects (Kim et al., 1999) which may explain some of the variance observed in
the network spatial specificity analyses. Lastly, this study focused on the network measures
using group ICA-based analyses. Although this is a very common approach used in resting-
state fMRI studies, investigation into the effects of motion and cumulative group data quantity
on other resting-state analysis approaches including seed-based and connectivity matrix-based
approaches were not conducted.
538

5 Conclusion

A non-invasive restraint was developed and evaluated along with three different
acclimation protocols. After evaluating motion and network quality with an externally generated
well-established network, the acclimation protocol was not found to change network spatial
specificity despite the short 4-day acclimation group having increased motion. However, longer
acclimation protocols (9-day and 13-day) did have improved amplitude correspondence with
the prior network. Separately analyzing the effect of motion and amount of data on network
quality showed a relatively small effect of motion compared to the large effect of amount of data
indicating increasing amount of scanning time (at the group and subject levels) is much more
important than the motion contamination allowed by the proposed approach. Invasive headpost
restraints have pushed the mouse fMRI field forward tremendously allowing for awake scanning
with very little motion. However, to increase translational utility, more focus should be put on
the development of non-invasive restraints to better align with scanning practices in humans.
By acquiring approximately 25% more data to conservatively compensate for censoring of high
motion frames, group resting state networks can be acquired at comparable quality to headpost
approaches. The proposed restraint is useful especially in situations where high-throughput
scanning is required without the use of any anesthesia for subject positioning, and if ample
unobstructed skull access is necessary, e.g., for multi-modal electrophysiology/optical/fMRI
studies. This awake MRI approach represents a step forward in increased translational utility
between mouse and human neuroimaging.
559

Data and Code Availability

560
561 All data and code have been made freely available (https://doi.org/10.17605/OSF.IO/TYHCE).
562

Author Contributions

563
564 S.L., and R.M. designed the study, S.L., A.E., M.B., and M.N. collected and analyzed the data,
565 S.L., K.M.G., P.Z., and R.M. designed the non-invasive restraint, S.L. wrote the initial draft of
566 the manuscript with feedback and review from all authors.
567

Funding

568
569 This work was supported by funding from the Canadian Institutes for Health Research (FDN-
570 148453), a Brain Canada Platform Support Grant and a Canada First Research Excellence
571 Fund award to BrainsCAN and a New Frontiers in Research Fund Transformation award
572 (NFRFT-2022-00051).
573

Declaration of Competing Interests

574
575 Nothing to declare.
576

Acknowledgements

577
578 We thank Alex Li for his help setting up and troubleshooting MRI scans; Gabriel Desrosiers-
579 Grégoire, Gabriel A. Devenyi, and Mallar Chakravarty from The Douglas Research Centre for
580 technical assistance with RABIES; and the Animal Care Committee from Western University
581 and veterinarians for feedback and suggestions while developing the non-invasive restraint.
582

References

Abraham, A., Pedregosa, F., Eickenberg, M., Gervais, P., Mueller, A., Kossaifi, J., Gramfort, A., Thirion, B. & Varoquaux, G. (2014). Machine learning for neuroimaging with scikit-learn. *Frontiers in Neuroinformatics*, *8*, 14. https://doi.org/10.3389/fninf.2014.00014

Avants, B. B., Tustison, N. J., Song, G., Cook, P. A., Klein, A. & Gee, J. C. (2011). A reproducible evaluation of ANTs similarity metric performance in brain image registration. *NeuroImage*, *54*(3), 2033–2044. https://doi.org/10.1016/j.neuroimage.2010.09.025

Bergmann, E., Gofman, X., Kavushansky, A. & Kahn, I. (2020). Individual variability in functional connectivity architecture of the mouse brain. *Communications Biology*, *3*(1), 738. https://doi.org/10.1038/s42003-020-01472-5

Chen, X., Tong, C., Han, Z., Zhang, K., Bo, B., Feng, Y. & Liang, Z. (2020). Sensory evoked fMRI paradigms in awake mice. *NeuroImage*, *204*, 116242. https://doi.org/10.1016/j.neuroimage.2019.116242

Desjardins, M., Kılıç, K., Thunemann, M., Mateo, C., Holland, D., Ferri, C. G. L., Cremonesi, J. A., Li, B., Cheng, Q., Weldy, K. L., Saisan, P. A., Kleinfeld, D., Komiyama, T., Liu, T. T., Bussell, R., Wong, E. C., Scadeng, M., Dunn, A. K., Boas, D. A., … Devor, A. (2019). Awake Mouse Imaging: From Two-Photon Microscopy to Blood Oxygen Level–Dependent Functional Magnetic Resonance Imaging. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, *4*(6), 533–542. https://doi.org/10.1016/j.bpsc.2018.12.002

Desrosiers-Grégoire, G., Devenyi, G. A., Grandjean, J. & Chakravarty, M. M. (2024). A standardized image processing and data quality platform for rodent fMRI. *Nature Communications*, *15*(1), 6708. https://doi.org/10.1038/s41467-024-50826-8

Dinh, T. N. A., Jung, W. B., Shim, H.-J. & Kim, S.-G. (2021). Characteristics of fMRI responses to visual stimulation in anesthetized vs. awake mice. *NeuroImage*, *226*, 117542. https://doi.org/10.1016/j.neuroimage.2020.117542

Eleftheriou, A., Palme, R. & Boonstra, R. (2020). Assessment of the Stress Response in North American Deermice: Laboratory and Field Validation of Two Enzyme Immunoassays for Fecal Corticosterone Metabolites. *Animals*, *10*(7), 1120. https://doi.org/10.3390/ani10071120

Ellenbroek, B. & Youn, J. (2016). Rodent models in neuroscience research: is it a rat race? *Disease Models & Mechanisms*, *9*(10), 1079–1087. https://doi.org/10.1242/dmm.026120

Fadel, L. C., Patel, I. V., Romero, J., Tan, I.-C., Kesler, S. R., Rao, V., Subasinghe, S. A. A. S., Ray, R. S., Yustein, J. T., Allen, M. J., Gibson, B. W., Verlinden, J. J., Fayn, S., Ruggiero, N., Ortiz, C., Hipskind, E., Feng, A., Iheanacho, C., Wang, A. & Pautler, R. G. (2022). A Mouse Holder for Awake Functional Imaging in Unanesthetized Mice:

Applications in 31P Spectroscopy, Manganese-Enhanced Magnetic Resonance Imaging
Studies, and Resting-State Functional Magnetic Resonance Imaging. *Biosensors*, *12*(8),
616. https://doi.org/10.3390/bios12080616

Ferris, C. F. (2022). Applications in Awake Animal Magnetic Resonance Imaging. *Frontiers in*
Neuroscience, *16*, 854377. https://doi.org/10.3389/fnins.2022.854377

Ferris, C. F., Smerkers, B., Kulkarni, P., Caffrey, M., Afacan, O., Toddes, S., Stolberg, T. &
Febo, M. (2011). Functional magnetic resonance imaging in awake animals. *Reviews in*
the Neurosciences, *22*(6), 665–674. https://doi.org/10.1515/rns.2011.050

Fu, Z., Liu, J., Salman, M. S., Sui, J. & Calhoun, V. D. (2023). Functional connectivity
uniqueness and variability? Linkages with cognitive and psychiatric problems in children.
Nature Mental Health, *1*(12), 956–970. https://doi.org/10.1038/s44220-023-00151-8

Gao, Y.-R., Ma, Y., Zhang, Q., Winder, A. T., Liang, Z., Antinori, L., Drew, P. J. & Zhang, N.
(2017). Time to wake up: Studying neurovascular coupling and brain-wide circuit function
in the un-anesthetized animal. *NeuroImage*, *153*, 382–398.
https://doi.org/10.1016/j.neuroimage.2016.11.069

Grandjean, J., Azzinnari, D., Seuwen, A., Sigrist, H., Seifritz, E., Pryce, C. R. & Rudin, M.
(2016). Chronic psychosocial stress in mice leads to changes in brain functional
connectivity and metabolite levels comparable to human depression. *NeuroImage*, *142*,
544–552. https://doi.org/10.1016/j.neuroimage.2016.08.013

Grandjean, J., Lake, E. M. R., Pagani, M. & Mandino, F. (2024). What N Is N-ough for MRI-
Based Animal Neuroimaging? *ENeuro*, *11*(3), ENEURO.0531-23.2024.
https://doi.org/10.1523/eneuro.0531-23.2024

Gutierrez-Barragan, D., Singh, N. A., Alvino, F. G., Coletta, L., Rocchi, F., Guzman, E. D.,
Galbusera, A., Uboldi, M., Panzeri, S. & Gozzi, A. (2022). Unique spatiotemporal fMRI
dynamics in the awake mouse brain. *Current Biology*.
https://doi.org/10.1016/j.cub.2021.12.015

Harris, A. P., Lennen, R. J., Marshall, I., Jansen, M. A., Pernet, C. R., Brydges, N. M., Duguid,
I. C. & Holmes, M. C. (2015). Imaging learned fear circuitry in awake mice using fMRI.
European Journal of Neuroscience, *42*(5), 2125–2134. https://doi.org/10.1111/ejn.12939

Kim, B., Boes, J. L., Bland, P. H., Chenevert, T. L. & Meyer, C. R. (1999). Motion correction in
fMRI via registration of individual slices into an anatomical volume. *Magnetic Resonance in*
Medicine, *41*(5), 964–972. https://doi.org/10.1002/(sici)1522-2594(199905)41:5<964::aid-
mrm16>3.0.co;2-d

Kroll, T., Kornadt-Beck, N., Oskamp, A., Elmenhorst, D., Touma, C., Palme, R. & Bauer, A.
(2021). Additional Assessment of Fecal Corticosterone Metabolites Improves Visual Rating
in the Evaluation of Stress Responses of Laboratory Rats. *Animals*, *11*(3), 710.
https://doi.org/10.3390/ani11030710

Lindhardt, T. B., Gutiérrez-Jiménez, E., Liang, Z. & Hansen, B. (2022). Male and Female
C57BL/6 Mice Respond Differently to Awake Magnetic Resonance Imaging Habituation.
Frontiers in Neuroscience, 16, 853527. https://doi.org/10.3389/fnins.2022.853527

Low, L. A., Bauer, L. C., Pitcher, M. H. & Bushnell, M. C. (2016). Restraint training for awake
functional brain scanning of rodents can cause long-lasting changes in pain and stress
responses. *PAIN, 157*(8), 1761–1772. https://doi.org/10.1097/j.pain.0000000000000579

Madularu, D., Mathieu, A. P., Kumaragamage, C., Reynolds, L. M., Near, J., Flores, C. &
Rajah, M. N. (2017). A non-invasive restraining system for awake mouse imaging. *Journal*
of Neuroscience Methods, 287, 53–57. https://doi.org/10.1016/j.jneumeth.2017.06.008

Mandino, F., Cerri, D. H., Garin, C. M., Straathof, M., Tilborg, G. A. F. van, Chakravarty, M.
M., Dhenain, M., Dijkhuizen, R. M., Gozzi, A., Hess, A., Keilholz, S. D., Lerch, J. P., Shih,
Y.-Y. I. & Grandjean, J. (2020). Animal Functional Magnetic Resonance Imaging: Trends
and Path Toward Standardization. *Frontiers in Neuroinformatics, 13*, 78.
https://doi.org/10.3389/fninf.2019.00078

Mandino, F., Vrooman, R. M., Foo, H. E., Yeow, L. Y., Bolton, T. A. W., Salvan, P., Teoh, C.
L., Lee, C. Y., Beauchamp, A., Luo, S., Bi, R., Zhang, J., Lim, G. H. T., Low, N., Sallet, J.,
Gigg, J., Lerch, J. P., Mars, R. B., Olivo, M., … Grandjean, J. (2021). A triple-network
organization for the mouse brain. *Molecular Psychiatry*, 1–8.
https://doi.org/10.1038/s41380-021-01298-5

Mandino, F., Vujic, S., Grandjean, J. & Lake, E. M. R. (2023). Where do we stand on fMRI in
awake mice? *Cerebral Cortex*, bhad478. https://doi.org/10.1093/cercor/bhad478

Maron-Katz, A., Vaisvaser, S., Lin, T., Hendler, T. & Shamir, R. (2016). A large-scale
perspective on stress-induced alterations in resting-state networks. *Scientific Reports, 6*(1),
21503. https://doi.org/10.1038/srep21503

Pagani, M., Gutierrez-Barragan, D., Guzman, A. E. de, Xu, T. & Gozzi, A. (2023). Mapping
and comparing fMRI connectivity networks across species. *Communications Biology, 6*(1),
1238. https://doi.org/10.1038/s42003-023-05629-w

Parker, C. C., Chen, H., Flagel, S. B., Geurts, A. M., Richards, J. B., Robinson, T. E., Woods,
L. C. S. & Palmer, A. A. (2014). Rats are the smart choice: Rationale for a renewed focus
on rats in behavioral genetics. *Neuropharmacology, 76*, 250–258.
https://doi.org/10.1016/j.neuropharm.2013.05.047

Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L. & Petersen, S. E. (2012).
Spurious but systematic correlations in functional connectivity MRI networks arise from
subject motion. *NeuroImage, 59*(3), 2142–2154.
https://doi.org/10.1016/j.neuroimage.2011.10.018

Power, J. D., Plitt, M., Gotts, S. J., Kundu, P., Voon, V., Bandettini, P. A. & Martin, A. (2018).
Ridding fMRI data of motion-related influences: Removal of signals with distinct spatial and

694 physical bases in multiecho data. *Proceedings of the National Academy of Sciences*,
695 *115*(9), E2105–E2114. https://doi.org/10.1073/pnas.1720985115

696 Rowland, N. E. & Toth, L. A. (2019). Analytic and Interpretational Pitfalls to Measuring Fecal
697 Corticosterone Metabolites in Laboratory Rats and Mice. *Comparative Medicine*, *69*(5),
698 337–349. https://doi.org/10.30802/aalas-cm-18-000119

699 Russo, G., Helluy, X., Behroozi, M. & Manahan-Vaughan, D. (2021). Gradual Restraint
700 Habituation for Awake Functional Magnetic Resonance Imaging Combined With a Sparse
701 Imaging Paradigm Reduces Motion Artifacts and Stress Levels in Rodents. *Frontiers in
702 Neuroscience*, *15*, 805679. https://doi.org/10.3389/fnins.2021.805679

703 Smith, J., Wilkey, E., Clarke, B., Shanley, L., Men, V., Fair, D. & Sabb, F. W. (2022). Can this
704 data be saved? Techniques for high motion in resting state scans of first grade children.
705 *Developmental Cognitive Neuroscience*, *58*, 101178.
706 https://doi.org/10.1016/j.dcn.2022.101178

707 Soares, J. M., Magalhães, R., Moreira, P. S., Sousa, A., Ganz, E., Sampaio, A., Alves, V.,
708 Marques, P. & Sousa, N. (2016). A Hitchhiker's Guide to Functional Magnetic Resonance
709 Imaging. *Frontiers in Neuroscience*, *10*, 515. https://doi.org/10.3389/fnins.2016.00515

710 Szilagyi, K., Zieger, M. A., Li, J. & Kacena, M. A. (2018). Improving Post-Operative Outcomes
711 in Aged and Diabetic Obese Mice. *Laboratory Animal Science Professional*, *6*(3), 65–67.

712 Takata, N., Sugiura, Y., Yoshida, K., Koizumi, M., Hiroshi, N., Honda, K., Yano, R., Komaki,
713 Y., Matsui, K., Suematsu, M., Mimura, M., Okano, H. & Tanaka, K. F. (2018). Optogenetic
714 astrocyte activation evokes BOLD fMRI response with oxygen consumption without
715 neuronal activity modulation. *Glia*, *66*(9), 2013–2023. https://doi.org/10.1002/glia.23454

716 Wang, S., Peterson, D. J., Gatenby, J. C., Li, W., Grabowski, T. J. & Madhyastha, T. M.
717 (2017). Evaluation of Field Map and Nonlinear Registration Methods for Correction of
718 Susceptibility Artifacts in Diffusion MRI. *Frontiers in Neuroinformatics*, *11*, 17.
719 https://doi.org/10.3389/fninf.2017.00017

720 Winkler, A. M., Ridgway, G. R., Webster, M. A., Smith, S. M. & Nichols, T. E. (2014).
721 Permutation inference for the general linear model. *NeuroImage*, *92*(100), 381–397.
722 https://doi.org/10.1016/j.neuroimage.2014.01.060

723 Wu, T., Wang, X., Zhang, R., Jiao, Y., Yu, W., Su, D., Zhao, Y. & Tian, J. (2020). Mice with
724 pre-existing tumors are vulnerable to postoperative cognitive dysfunction. *Brain Research*,
725 *1732*, 146650. https://doi.org/10.1016/j.brainres.2020.146650

726 Xu, N., LaGrow, T. J., Anumba, N., Lee, A., Zhang, X., Yousefi, B., Bassil, Y., Clavijo, G. P.,
727 Sharghi, V. K., Maltbie, E., Meyer-Baese, L., Nezafati, M., Pan, W.-J. & Keilholz, S. (2022).
728 Functional Connectivity of the Brain Across Rodents and Humans. *Frontiers in
729 Neuroscience*, *16*, 816331. https://doi.org/10.3389/fnins.2022.816331

Xu, W., Pei, M., Zhang, K., Tong, C., Bo, B., Feng, J., Zhang, X.-Y. & Liang, Z. (2022). A
systematically optimized awake mouse fMRI paradigm. *BioRxiv*, 2022.11.16.516376.
https://doi.org/10.1101/2022.11.16.516376

Zeng, H., Jiang, Y., Beer-Hammer, S. & Yu, X. (2022). Awake Mouse fMRI and Pupillary
Recordings in the Ultra-High Magnetic Field. *Frontiers in Neuroscience*, *16*, 886709.
https://doi.org/10.3389/fnins.2022.886709

Zhao, W., Makowski, C., Hagler, D. J., Garavan, H. P., Thompson, W. K., Greene, D. J.,
Jernigan, T. L. & Dale, A. M. (2023). Task fMRI paradigms may capture more behaviorally
relevant information than resting-state functional connectivity. *NeuroImage*, *270*, 119946.
https://doi.org/10.1016/j.neuroimage.2023.119946

740
741

Supplementary Figures and Tables

744 Supplementary Table 1. RABIES (version 0.5.1) processing stage commands.

Processing Stage	Parameters
Preprocessing	--anat_robust_inho_cor apply=true,masking=false,brain_extraction=false,template_registration=SyN --commonspace_resampling 0.2x0.2x0.2 --bold2anat_coreg masking=true,brain_extraction=false,registration=SyN
Confound Correction	--smoothing_filter 0.3 --conf_list mot_6 --frame_censoring FD_censoring=true,FD_threshold=0.15,DVARS_censoring=false,minimum_timepoint=200
Analysis	--data_diagnosis

747 Supplementary Table 2. Percentage of volumes censored for each group. The first 4 rows represent the data divided based on
748 acclimation group. The last 6 rows represent the non-invasive data grouped by average framewise displacement per run. The
749 range of average run framewise displacements is reported under each group. Note, motion level 1 has fewer runs than the
750 other motion groups to consist of runs that better match the average motion in the headpost group. *Motion Level 6 was not
751 included in analyses due to a lack of data from censoring.

Group	Total Runs	Censored Volumes	Runs Excluded
4-Day (0.039 – 0.13 mm)	36	20.6%	5 (13.9%)
9-Day (0.015 – 0.14 mm)	74	16.2%	4 (5.41%)
13-Day (0.014 – 0.16 mm)	102	22.1%	10 (9.80%)
Headpost (0.018 – 0.027 mm)	14	0.0%	0 (0.00%)
Motion Level 1 (0.014 – 0.030 mm)	24	0.3%	0 (0.00%)
Motion Level 2 (0.031 – 0.043 mm)	38	4.2%	0 (0.00%)
Motion Level 3 (0.044 – 0.059 mm)	38	8.1%	0 (0.00%)
Motion Level 4 (0.059 – 0.071 mm)	38	14.5%	0 (0.00%)
Motion Level 5 (0.071 – 0.091 mm)	38	26.0%	0 (0.00%)
Motion Level 6* (0.091 – 0.16 mm)	36	71.5%	19 (53.8%)

756
Supplementary Figure 1. Relationship between spatial specificity (Dice Similarity Coefficient, DSC) and average group
framewise displacement (FD) for different group cumulative acquisition times. All functional runs were aggregated and sorted
based on the average FD value. Group network spatial specificity was determined for all groups and plotted. Linear trendlines
were fitted to each cumulative acquisition time groupings.

Supplementary Figure 2. Correspondence between z-values of the prior network and corresponding group networks on a voxel-
wise basis for voxels in the prior network region. One sample from each group network is shown. The correlation coefficient is
annotated on each plot. The gap between -3.1 and 3.1 results from masking the prior network for voxels with values >= +/-3.1
and selecting those voxels for the correlation analysis. The corresponding Pearson Correlation coefficient (r value) is annotated
on each plot.

769
770
771 Supplementary Figure 3. Weight and fecal corticosterone (CORT) metabolite over the 3 acclimation days. (a) Weight split up
772 by sex for the 4-day, 9-day, 13-day and headpost groups. (b) Same as (a) except both sexes aggregated. (c) CORT split up
773 by sex for the 4-day, 9-day, and 13-day groups. (d) Same as (c) except both sexes aggregated. CORT concentrations are
774 displayed in ng of corticosterone per mg of fecal matter. Data are displayed as group means (data points) and standard
775 deviation (error bars).