Elucidation of the biosynthetic pathway of reserpine

Jiaqing Cao[1], Jingxiao Zhong[1,2,3], Feng Li[1,4], Yindi Jiang[1,*]

[1] Key Laboratory of Quantitative Synthetic Biology, Shenzhen Institute of Synthetic Biology,

Shenzhen Institutes of Advanced Technology, Chinese Academy of Sciences, Shenzhen 518055,

China.

[2] School of Life Sciences, Henan University, Kaifeng 475004, China.

[3] Shenzhen Research Institute of Henan University, Shenzhen 518000, China.

[4] University of Chinese Academy of Sciences, Beijing 100049, China.

[*]E-mail: yd.jiang@siat.ac.cn

Abstract

Reserpine is a landmark natural product that has profoundly influenced our understanding of

neurotransmitter biology and cardiovascular medicine. Its complex structure, featuring a unique

C3 β-configuration and five consecutive chiral centers, has inspired generations of synthetic

chemists to develop innovative strategies for its construction. However, the biosynthetic logic

underlying nature's assembly of this intricate molecule has remained elusive. Here, we decipher

the biosynthetic pathway of reserpine in *Rauvolfia verticillata*, revealing that α-configured

strictosidine serves as the biosynthetic precursor. The crucial C3 β-configuration is established

through a two-step enzymatic epimerization, orchestrated by a flavin-dependent oxidase and a

NADPH-dependent reductase. The consecutive chiral centers are constructed through coordinated

action of distinct enzyme families. Through the identification of eight biosynthetic enzymes,

including those catalyzing late-stage methoxylation, we successfully reconstituted the biosynthesis

of rauvomitorine G, a key intermediate in reserpine formation. This work unveils nature's elegant

approach to stereoselective synthesis of complex alkaloids and provides valuable biocatalytic tools

for molecular functionalization.

Introduction

Reserpine is a structurally complex monoterpene indole alkaloid (MIA) that has profoundly influenced our fundamental understanding of biomedicine and chemistry. Since its isolation from *Rauvolfia serpentina* in 1952, reserpine has been instrumental for deciphering the complex mechanisms of neurotransmitter storage and release, particularly regarding monoamine neurotransmitters such as serotonin and norepinephrine[1,2,3]. Its ability to deplete these neurotransmitters has provided groundbreaking insights into the pathophysiology of hypertension and various psychiatric disorders[4,5]. Furthermore, reserpine's clinical application as antihypertensive and antipsychotic agents paved the way for the development of more targeted therapeutics, revolutionizing the treatment paradigms in cardiovascular and mental health medicine[6].

Beyond its biomedical significance, the complex structure of reserpine has challenged and inspired generations of synthetic chemists, leading to numerous total syntheses and innovative synthetic methodologies[7]. The molecule's pentacyclic skeleton features a unique C3β stereochemistry and five consecutive chiral centers densely packed in the E ring (Fig. 1a). Most total syntheses of reserpine begin with the construction of a pentasubstituted E-ring, typically achieved through various synthetic strategies including Diels-Alder reaction, double Michael addition, and radical cyclization (Fig. 1b)[8,9,10,11,12]. This elaborated E-ring subsequently undergoes Pictet-Spengler condensation with 6-methoxytryptamine to generate a pentacyclic intermediate bearing an α-configuration at C3. Access to the desired β-configuration at C3 has largely relied on R. B. Woodward's landmark conformational control strategy, which involves locking the molecule in an unfavorable conformation through lactonization, followed by acid treatment to achieve late-stage C3 epimerization[8]. Despite these sophisticated strategies developed by synthetic chemists, the biosynthetic pathway by which nature constructs such complex natural products remains elusive.

Isotope feeding studies have established tryptamine as the biosynthetic precursor of reserpine[13]. Its condensation with monoterpene-derived secologanin yields two diastereomers, vincoside and strictosidine[14]. The shared C3 configuration between vincoside and reserpine has historically

suggested vincoside as the biosynthetic precursor of reserpine[7]. However, this proposal contradicts

established MIA biosynthetic pathways, where strictosidine, not vincoside, functions as the

common precursor[15]. This discrepancy raises an intriguing possibility that strictosidine might serve

as the reserpine precursor, necessitating subsequent stereochemical inversion to establish the

characteristic C3β configuration of reserpine.

Here we report the elucidation of the reserpine biosynthetic pathway in *R. verticillata* through an

integrated approach combining transcriptome mining, heterologous expression, and biochemical

characterization (Fig. 1c and 2). Our findings reveal that strictosidine, not vincoside, serves as the

authentic precursor in reserpine biosynthesis. We uncovered the enzymatic basis for C3

epimerization in *Rauvolfia*, which proceeds through a two-step process: an FAD-dependent

oxidase *Rv*YOO generates an imine intermediate, which is subsequently reduced by a medium-

chain dehydrogenase/reductase (MDR) *Rv*DYR1 to yield the 3-epi product. Furthermore, we

revealed nature's elegant strategy for constructing consecutive chiral centers in ring E, where two

stereocenters are established through stereoselective reduction by an MDR *Rv*YOS, and the final

stereocenter is installed via cytochrome P450 monooxygenase *Rv*CYP71D820. Through the

identification of eight biosynthetic enzymes, including P450 monooxygenase *Rv*CYP72A270 and

O-methyltransferase *Rv*11OMT for late-stage methoxylation, we successfully reconstituted

rauvomitorine G, a crucial intermediate in reserpine biosynthesis. This work advances our

understanding of complex alkaloid biosynthesis, revealing both the enzymatic logic underlying

reserpine biosynthesis and providing valuable biocatalytic tools for selective functionalization of

complex molecular scaffolds.

Results

To elucidate the biosynthesis of reserpine, we first examined the distribution of monoterpene

indole alkaloids (MIAs) in *Rauvolfia verticillata*, a medicinal plant used in China as a source of

reserpine. Liquid chromatography-mass spectrometry (LC-MS) analysis of tissue extracts revealed

several key MIAs, including reserpine (**11**), deserpidine (**13**), α-yohimbine (**2**), and 3-epi-α-

yohimbine (**4**) (Supplementary Fig. 1). Having confirmed the presence of these MIAs across

different tissues, we sought to identify the genes responsible for their biosynthesis. To facilitate

biosynthetic gene discovery, we generated tissue-specific transcriptome datasets in triplicate.

Formation of α-yohimbine through strictosidine pathway

A distinct structural feature of reserpine is its β-configuration at C3, which contrasts with the α-

configuration predominant in most MIAs. The isolation of vincoside and vincoside lactam, which

share this β-configuration at C3, from *Rauvolfia* species has long suggested vincoside as the

biosynthetic precursor of reserpine (Supplementary Fig. 2)[7,16]. However, the stepwise conversion

from vincoside to reserpine remains unsolved.

Strictosidine (**1**), a stereoisomer of vincoside and the common MIA precursor, is stereospecifically

synthesized from tryptamine and secologanin by strictosidine synthase (STR). The subsequent

deglycosylation by strictosidine glucosidase (SGD) generates a reactive iminium intermediate,

which various reductases can transform into diverse indole alkaloids[15]. While *Rauvolfia* SGD

showed specificity for **1** deglycosylation, it exhibits no activity towards vincoside[17]. Notably,

Catharanthus roseus SGD (*Cr*SGD) can deglycosylate both **1** and vincoside, suggesting the

possible existence of an unidentified vincoside-specific SGD in *Rauvolfia*[18].

The prevalence of C3 β-configured intermediates in *Rauvolfia* is evidenced by numerous isolated

compounds, including vincoside lactam, 3-epi-α-yohimbine (**4**), rauvomitorine G (**9**), deserpidic

acid methyl ester (**12**), and rescinnamine, with **4** representing the structurally simplest intermediate

(Supplementary Fig. 2)[19,20,21]. Recent studies have elucidated the biosynthesis of α-yohimbine (**2**),

the C3 epimer of **4**, in *R. tetraphylla*[22]. This pathway proceeds via strictosidine (**1**), involving SGD-

catalyzed formation of a reactive imine intermediate and subsequent reduction by yohimbane

synthase (YOS), a medium-chain dehydrogenase/reductase (MDR).

These findings led us to propose two potential biosynthetic pathways for 3-epi-α-yohimbine (**4**)

(Supplementary Fig. 3). Pathway A involves condensation of tryptamine and secologanin to form

vincoside (**1a**), followed by conversion to **4** through an unidentified SGD and an MDR.

Alternatively, pathway B begins with STR-mediated condensation of tryptamine and secologanin

to produce strictosidine (**1**)[23], which is converted to α-yohimbine (**2**) by characterized SGD and

YOS enzymes[22,24]. Subsequently, stereochemical inversion of C-3 of **2** could be catalyzed by an

unidentified epimerase to yield **4**.

To investigate pathway A, we chemically synthesized a mixture of strictosidine (**1**) and vincoside

(**1a**). Five putative candidate genes potentially capable of vincoside deglycosylation were

identified in the *R. verticillata* transcriptome using *Cr*SGD as a query. Each gene was cloned from

R. verticillata cDNA and expressed in *Nicotiana benthamiana*. LC-MS analysis revealed that

while *Rv*SGD1 effectively consumed strictosidine (**1**), vincoside remained unchanged in all SGD

candidate assays (Supplementary Fig. 4), suggesting that vincoside is not the source of 3-*epi*-α-

yohimbine (**4**).

We then explored pathway B through reconstituting α-yohimbine (**2**) in planta (Fig. 3a). Previous

work in *R. tetraphylla* showed that YOS reduces strictosidine aglycone to form a mixture of

yohimbine-type alkaloids, including α-yohimbine (**2**), yohimbine (**2a**), and corynanthine (**2c**)[22].

BLAST analysis of the *R. verticillata* transcriptome using *Rt*YOS as a query revealed a candidate

gene, *Rv*YOS, sharing 94.4% sequence identity. To assess functional conservation, we

coexpressed *Rv*YOS with *Catharanthus roseus* STR (*Cr*STR) and *Rv*SGD in *N. benthamiana*.

Following infiltration of tryptamine and secologanin, this system successfully produced both **2** and

2a (Supplementary Fig. 5). The formation of these products was further confirmed *in vitro* using

purified *Cr*STR, *Rv*SGD, and *Rv*YOS enzymes (Supplementary Fig. 5). Although *Rv*YOS

generated **2**, its lack of product specificity led us to screen all MDRs sharing over 30% sequence

identity for enzymes that could selectively produce **2** (Fig. 3b). As none exhibited the desired

specificity, we proceeded with a chemical standard of **2** for subsequent experiments.

Enzymatic epimerization of α-yohimbine (2)

We next investigated the enzymatic basis of α-yohimbine (**2**) epimerization to 3-epi-α-yohimbine

(**4**). Epimerization could be catalyzed by epimerases, which are crucial in the primary metabolism

of carbohydrates and amino acids, as well as in secondary metabolite biosynthesis[25,26]. During

morphine biosynthesis in opium poppy, a reticuline epimerase (REPI) catalyzed the

stereoconversion of *S*-reticuline to *R*-reticuline through sequential oxidation and reduction with its

two domains[27]. The N-terminal cytochrome P450 domain oxidizes *S*-reticuline to an iminium

intermediate, which was stereo-specifically reduced by the C-terminal aldo-keto reductase (AKR)

domain to *R*-reticuline in a NADPH-dependent manner (Supplementary Fig. 6). Thus, we

speculated that epimerization of α-yohimbine (**2**) happened in a similar manner that **2** was oxidized

to iminium intermediate 3-dehydro-α-yohimbine (**3**), followed by reduction from the convex face

to give 3-epi-α-yohimbine (**4**) (Fig. 3a).

Our search for REPI homologs in the *R. verticillata* transcriptome failed to identify candidates

containing both P450 and AKR domains. Although the CYP450 domain in REPI belongs to the

CYP82 family, expression of seven CYP82 gene candidates from *R. verticillata* in *N. benthamiana*

leaves co-infiltrated with **2** yielded no detectable oxidized products. We then considered

alternative oxidases other than cytochrome P450s. In vinblastine biosynthesis in *C. roseus*, the

flavin-dependent precondylocarpine acetate synthase (*Cr*PAS) oxidizes the tertiary amine in

stemmadenine acetate to an iminium intermediate[28]. This led us to investigate PAS homologs as

potential oxidants for **2**. Among eight *Cr*PAS homologs identified in the *R. verticillata*

transcriptome, one candidate sharing 58.0% identity with *Cr*PAS, which we named α-yohimbine

oxidase (*Rv*YOO), catalyzed the conversion of **2** to a product with an exact mass of $[M+H]^+$

353.1857, consistent with an oxidized product of **2** (calculated for $C_{21}H_{25}N_2O_3^+$: 353.1860) (Fig.

3c). The structure was confirmed using synthetic 3-dehydro-α-yohimbine (**3**), which showed

identical retention time, exact mass, and MS/MS fragmentation patterns (Supplementary Fig. 7).

Further investigation of *Rv*YOO's substrate scope through in planta expression and infiltration

with α-yohimbine stereoisomers revealed that α-yohimbine (**2**), yohimbine (**2a**), and *β*-yohimbine

(**2b**) were oxidized, while corynanthine (**2c**) and 3-epi-α-yohimbine (**4**) were not substrates

(Supplementary Fig. 7). Notably, the inability of *Rv*YOO to oxidize **4** suggests that once formed,

4 cannot be coverted back to **3** through the same enzymatic mechanism, indicating the irreversible

nature of this transformation.

For the subsequent reduction of 3-dehydro-α-yohimbine (**3**), we focused on MDR enzymes, which

are known to catalyze iminium reduction in MIA biosynthesis[29,30,31,32]. A well-characterized

example is tetrahydroalstonine synthase (THAS) in *C. roseus* that performs 1,2-reduction of

iminium strictosidine aglycone to yield tetrahydroalstonine[33]. Co-expression of either of the two

3-dehydro-α-yohimbine reductases (*Rv*DYR1 and *Rv*DYR2) with *Rv*YOO in *N. benthamiana*

infiltrated with α-yohimbine (**2**) resulted in the formation of a product that co-eluted with authentic

3-epi-α-yohimbine (**4**) standard (Fig. 3d). However, due to incomplete substrate conversion by

*Rv*YOO, we observed a mixture of **2** and **4** in *N. benthamiana* leaves expressing active MDRs,

which prevented definitive conclusions about the stereoselectivity of *Rv*DYR1 and *Rv*DYR2

toward **3**.

To unambiguously determine stereoselectivity, we expressed and purified N-terminal His6-tagged

*Rv*DYR1 and *Rv*DYR2 for *in vitro* characterization (Supplementary Fig. 8). Incubation of

synthetic **3** with purified enzymes and NADPH predominantly yielded 3-epi-α-yohimbine (**4**) over

α-yohimbine (**2**) (Fig. 3e). Notably, **2** could be generated by NADPH alone, while the formation

of **4** required both NADPH and active *Rv*DYR enzymes, indicating *Rv*DYR enzymes' preference

for C3β stereoselectivity. Interestingly, despite high sequence similarity among *Rv*DYR1,

*Rv*DYR2, and *Rv*DYR15260, *Rv*DYR15260 showed no activity toward the substrate

(Supplementary Fig. 9). This lack of activity might be attributed to certain non-conserved residues

among these three enzymes. While we were preparing this manuscript, two independent studies

reported similar epimerization mechanisms for MIAs in plants[34,35]. To further understand the

molecular basis of substrate recognition and and stereochemical control, structural studies

including X-ray crystallography and site-directed mutagenesis are currently underway.

Stereospecific C18 hydroxylation establishing the fifth chiral center in ring E

The next stage of reserpine biosynthesis involves two hydroxylation events at C11 and C18 as well

as two methylation events at O11 and O17. The isolation of 18β-hydroxy-3-epi-α-yohimbine (**5**)

and rauvomitorine G (**9**) from *Rauvolfia* suggested that stereospecific hydroxylation at C18 of **4**

occurs first, followed by methoxylation at C11 (Fig. 4a)[19,20]. Given that no C18 hydroxylase has

been previously reported, sequence homology-based approaches were not feasible for identifying

the responsible enzyme. Therefore, we employed a co-expression analysis strategy using *Rv*DYR1

as a bait, as the expression of upstream biosynthetic genes showed similar tissue-specific

expression patterns and high correlation (Fig. 4b and 4c). We selected 30 P450 enzymes (Pearson's

r>0.7) for functional characterization (Supplementary Fig 10). Transient expression of

*Rv*CYP71D820 in *N. benthamiana* leaves infiltrated with **4** resulted in the formation of a new

product with an exact mass (*m/z* [M+H]$^+$ = 371.1963) matching hydroxylated **4** (calculated for

$C_{21}H_{27}N_2O_4^+$: 371.1965) (Fig. 4d). We validated this finding *in vitro* using *Rv*CYP71D820 in yeast

microsomal fractions, which efficiently converted **4** to the same product observed in planta (Fig.

4d). MS/MS analysis of this product revealed an intact indole moiety (*m/z* $[M+H]^+$ = 144.0807)

(Supplementary Fig. 11), indicating hydroxylation occurred outside the indole ring. The product's

identity was confirmed using a chemically synthesized standard of **5**, prepared from deserpidine

(**13**) in two steps, which showed identical retention time, exact mass, and MS/MS fragmentation

patterns (Supplementary Fig. 11). These results establish *Rv*CYP71D820 as the enzyme

responsible for stereospecific C18 hydroxylation of **4**, completing the construction of five

consecutive chiral centers in the E ring of reserpine.

Enzymatic late-stage methoxylation

We next sought to identify the oxidase and methyltransferase responsible for C11 methoxylation

of **5**. During vindoline biosynthesis in *C. roseus*, tabersonine 16-hydroxylases (T16Hs) of the

CYP71D family catalyze C16 hydroxylation of tabersonine to form 16-hydroxytabersonine,

followed by *O*-methylation catalyzed by 16-hydroxytabersonine 16-*O*-methyltransferase (16OMT)

to complete methoxy group installation[36,37,38]. Given that the C16 position of tabersonine is

structurally equivalent to the C11 position in reserpine, we searched for T16H and 16OMT

homologs in the *R. verticillata* transcriptome. When T16H homologs proved inactive toward **5**, we

rescreened P450 candidate genes co-expressed with *Rv*DYR1. Among these candidates,

coexpression of *Rv*CYP72A270 with *Rv*CYP71D820 in *N. benthamiana* leaves infiltrated with **4**

resulted in a product that co-eluted with synthetic 11,18β-dihydroxy-3-epi-α-yohimbine (**6**) (Fig.

4e and Supplementary Fig. 12). Next, screening of 16OMT homologs in *N. benthamiana* leaves

infiltrated with **6** revealed that only *Rv*11OMT could catalyze the formation of rauvomitorine G

(**9**) (Fig. 4d and Supplementary Fig. 13). We further validated this activity using purified N-

terminal His6-tagged *Rv*11OMT *in vitro* (Fig. 4f). Finally, co-expression of *Rv*CYP71D820,

*Rv*CYP72A270, and *Rv*11OMT in *N. benthamiana* leaves treated with **4** successfully yielded **9**

(Fig. 4g), confirming the sequential activities of these enzymes in the pathway.

Branched pathway

During the co-expression of *Rv*CYP72A270 and *Rv*CYP71D820 in *N. benthamiana* infiltrated

with **4**, we observed a minor peak sharing the exact mass with **5** but exhibiting a different retention

time (Supplementary Fig. 14). MS/MS analysis of the peak indicated hydroxylation on the indole

ring (Supplementary Fig. 14). Given *Rv*CYP72A270's C11 hydroxylation activity on **5**, we

proposed that *Rv*CYP72A270 could also oxidize **4** to form the 11-hydroxy-3-epi-α-yohimbine (**7**).

This hypothesis was confirmed when we detected **7** in *N. benthamiana* expressing *Rv*CYP72A270

and treated with **4** (Supplementary Fig. 15). Based on *Rv*11OMT's known ability to methylate the

C11 hydroxyl group of **6**, we investigated its activity toward **7**. Co-expression of *Rv*CYP72A270

and *Rv*11OMT in *N. benthamiana* infiltrated with **4** yielded a product peak (**8**) corresponding to

the methoxylated derivative (Supplementary Fig. 15). The MS/MS fragmentation pattern of **8**

confirmed the presence of a methoxy group on the indole ring (Supplementary Fig. 15). However,

without authentic standards for **7** and **8**, their exact structures could not be unambiguously

confirmed. Furthermore, whether **8** could serve as a substrate for *Rv*CYP71D820 to generate **9**

remain to be determined. These findings suggest a potential branched pathway from **4** to **9**, adding

another layer of complexity to the biosynthetic route (Supplementary Fig. 16).

Remaining biosynthetic steps

According to our biosynthetic hypothesis, the next step involves the *O*-methylation of the C17

hydroxyl group of rauvomitorine G (**9**) to yield reserpic acid methyl ester (**10**) (Fig. 2). Guided by

the robust co-expression patterns observed among our newly identified biosynthetic genes (Fig.

4b and 4c), we initially employed co-expression analysis to identify the putative C17 *O*-

methyltransferase. We heterologously expressed 30 methyltransferases that showed strong co-

expression correlation (r>0.5) with *Rv*11OMT in *N. benthamiana*, using **9** as the substrate.

However, none of these candidates exhibited the desired methyltransferase activity. Further

exploration of the *R. verticillata* transcriptome for homologs of well-characterized

methyltransferases from the MIA biosynthetic pathway, including *Cr*16OMT and loganic acid *O*-

methyltransferase (LAMT), also proved unsuccessful (Supplementary Fig. 17)[38,39]. No activity

was detected toward either **9** or 18β-hydroxy-3-epi-α-yohimbine (**5**), the latter being a potential

substrate for C17 *O*-methylation in the biosynthesis of deserpidine (**13**), a demethoxylated

reserpine derivative with significant biological activities[40]. This challenging identification might

be attributed to the unique nature of the C17 hydroxyl group of **9**. Typical plant *O*-

methyltransferases target hydroxyl groups with relatively lower pKa values, such as phenolic

hydroxyl or carboxyl groups[41]. However, the C17 hydroxyl group in **9** has a significantly higher

pKa, making it less prone to form oxyanions for nucleophilic attack on *S*-adenosylmethionine.

This distinctive property suggest that an unusual methyltransferase might be required to facilitate

the deprotonation and subsequent methylation of this challenging hydroxyl group.

The final step of reserpine biosynthesis involves the transfer of a 3,4,5-trimethoxybenzoyl (TMB)

group to reserpic acid methyl ester (**10**) (Fig. 2). Such acylation steps are typically catalyzed by

two enzyme families: BAHD-acyltransferases (BAHD-ATs) and serine carboxypeptidase-like

acyltransferases (SCPL-ATs)[42]. Recent studies of leonurine biosynthesis in two *Leonurus* species

have demonstrated that 4-hydroxy-3,5-dimethoxybenzoic acid (syringic acid) is first glycosylated

by a UDP-glucosyltransferase (UGT) to form the glucose ester of syringic acid, which then

undergoes SCPL-catalyzed acylation with 4-guanidinobutanol to yield leonurine (Supplementary

Fig. 18)[43]. Given the structural similarity between syringic acid and TMB, we hypothesized that a

similar pathway operates in reserpine (**11**) biosynthesis, where a UGT glycosylates 3,4,5-

trimethoxybenzoic acid (TMBA) to form 1-*O*-TMB-β-D-glucopyranoside (TMB-Glc), followed

by SCPL-mediated acylation of **10** (Supplementary Fig. 18).

To identify the UGT responsible for TMBA glycosylation, we performed transcriptome analysis

using the *Leonurus* UGT sequence as a query and identified seven UGT candidate genes

(Supplementary Fig. 19). These candidates were cloned from *R. verticillata* cDNA and

heterologously expressed in *E. coli* BL21. In vitro assays with purified candidate enzymes, TMBA,

and uridine diphosphate glucose (UDP-Glc) revealed that *Rv*UGT1 and *Rv*UGT2 catalyzed TMBA

glycosylation, yielding a product identical to the synthetic TMB-Glc standard in retention time,

exact mass, and MS/MS fragmentation pattern (Supplementary Fig. 20).

For the final acylation step, we identified 15 SCPL candidate genes from the transcriptome using

the *Leonurus* SCPL sequence as a query (Supplementary Fig. 21). These candidates possessed the

characteristic N-terminal signal peptides typical of SCPLs, which require precise post-translational

processing for activation (Supplementary Fig. 22). Despite coexpressing full-length versions of

each SCPL candidate with *Rv*UGT1 in both *S. cerevisiae* and *N. benthamiana*, reserpine (**11**)

formation was not detected when supplementing with reserpic acid methyl ester (**10**) and TMBA.

This lack of activity likely stems from the complex post-translational modifications required for

SCPL maturation, including signal peptide cleavage, protein folding in the endoplasmic reticulum,

and potential glycosylation[42,44]. These plant-specific processing events may not be adequately

reproduced in heterologous hosts, suggesting the need for optimized expression conditions or

alternative expression systems that better mimic the native cellular environment.

Reconstitution of rauvomitorine G (9)

Having identified all biosynthetic enzymes responsible for the production of reserpine biosynthetic

intermediate **9**, we aimed to reconstitute **9** in tobacco. We overexpressed eight genes starting from

STR to *Rv*11OMT in tobacco and supplemented tryptamine and secologanin as substrates. LC-MS

analysis revealed only trace amounts of **3**, **4**, **5** and **9** (Fig. 5a). The production efficiency of **9** was

significantly compromised likely by the extensive length of the biosynthetic pathway and low

efficiency of certain enzymes, particularly *Rv*YOS, which exhibits low catalytic activity and

generates multiple yohimbine isomers that can serve as competing substrates for downstream

enzymes. To better validate the entire pathway, we divided it into two segments for separate

verification: the first segment from tryptamine and secologanin to **2**, and the second segment from

2 to **9**. When tested separately, both segments showed markedly improved performance. The first

segment, comprising *Cr*STR, *Rv*SGD, and *Rv*YOS produced **2** (Supplementary Fig. 5). Similarly,

the second segment, containing *Rv*YOO, *Rv*DYR1, *Rv*CYP72A270, *Rv*CYP71D820, and

*Rv*11OMT, efficiently converted externally supplied **2** to **9** (Fig. 5b and Supplementary Fig. 23).

These results confirmed the functionality of all biosynthetic steps and suggested that pathway

optimization strategies, such as protein engineering of *Rv*YOS and balanced expression of pathway

genes, could enhance the overall production of **9**.

Conclusion

In conclusion, we have deciphered the biosynthetic pathway of reserpine in *Rauvolfia verticillata*,

resolving a long-standing question in natural product chemistry. The molecule's complex

architecture, featuring a unique C3β stereochemistry and five consecutive chiral centers,

exemplifies sophisticated natural product biosynthesis. Our investigation revealed the enzymatic

basis for C3 epimerization, which proceeds through an oxidation-reduction sequence catalyzed by

an FAD-dependent oxidase (*Rv*YOO) and a medium-chain dehydrogenase/reductase (*Rv*DYR1)

to establish the crucial C3β stereochemistry. The pathway systematically constructs the five

consecutive chiral centers: the first two derive from secologanin, the third and fourth are installed
via stereoselective reduction by *Rv*YOS, and the final stereocenter is established through
cytochrome P450-catalyzed hydroxylation. Notably, while chemical synthesis typically introduces
the methoxy group at an early stage, nature employs a distinct late-stage strategy using a P450-
dependent methyltransferase, highlighting the contrasting approaches between biosynthetic and
synthetic chemistry. This work not only provides fundamental insights into the biosynthesis of
complex alkaloids but also establishes a foundation for metabolic engineering efforts aimed at
producing reserpine and related pharmaceutically important compounds. The enzymatic tools
uncovered here may enable the development of chemoenzymatic approaches for accessing
structurally diverse alkaloids with potential therapeutic applications.

Methods

Candidate gene cloning
The full-length sequences of candidate genes were amplified from cDNA using the primers listed in
Supplementary Table 2 with 2 × Phanta Max Master Mix according to the manufacturer's instructions. PCR
products were purified and ligated into different linearized expression vectors: 3Ω1 vector (digested with BsaI),
and pOPINF and pOPINM vectors (both digested with HindIII-HF and KpnI-HF) using the ClonExpress® II one
step cloning kit. The ligation reactions were transformed into chemically competent *E. coli* DH5α cells
(AlpaLifeBio, Shenzhen, China). Transformed cells were plated on LB agar plates containing either 50 μg/mL
spectinomycin for 3Ω1 constructs or 50 μg/mL ampicillin for pOPINF/pOPINM constructs. Single colonies were
picked and cultured overnight in 3 mL selective LB medium at 37 °C with shaking at 220 rpm. Plasmids were
isolated and verified by Sanger sequencing. Confirmed positive strains were mixed with an equal volume of 50%
glycerol and stored at −80 °C as frozen stocks.

364
Transformation of *Agrobacterium tumefaciens*
The 3Ω1 plasmids containing the full-length candidate gene were transformed into chemically competent *A.*
tumefaciens strain GV3101 (AngYuBio., Shanghai, China) following the manufacturer's instructions. The
transformed cells were plated onto LB agar plates containing 50 μg/mL rifampicin, 50 μg/mL gentamycin, and
25 μg/mL spectinomycin, and incubated at 28 °C for 2-3 days. Single colonies were picked and grown overnight
at 28 °C in 5 mL of LB medium containing 50 μg/mL rifampicin, 50 μg/mL gentamycin, and 25 μg/mL
spectinomycin. The overnight cultures were mixed with an equal volume of 50% glycerol and stored at −80 °C
as frozen stocks.

373
Transient expression of candidate genes in *N. benthamiana*
Agrobacterium GV3101 strains harboring the constructs of candidate genes were cultivated in 3 mL LB medium
supplemented with 25 μg/mL rifampicin, 25 μg/ml gentamycin, and 50 μg/ml spectinomycin overnight at 28 °C
with a shaking of 220 rpm. The overnight cultures were centrifuged at 4,000 g for 10 min at room temperature.
The cell pellet was resuspended in 3 mL infiltration buffer (50 mM MES, 22.2 mM glucose, 10 mM $MgCl_2$ and
100 μM acetosyringone). After centrifugation at 4,000 g for 10 min, the pellet was resuspended in 5 mL
infiltration buffer and adjusted to an OD_{600} of 0.6. The bacterial suspension was incubated at room temperature
for 2 h in the dark with gentle shaking before being infiltrated into the abaxial (bottom) side of 4-week-old *N.*
benthamiana leaves using a needleless 1 mL syringe. Four days post-infiltration, substrate solutions (200 μM in
water or in 1 % DMSO) were infiltrated into the same areas of Agrobacterium-infiltrated leaves using a
needleless 1 mL syringe. Substrate-infiltrated leaves were harvested 2 days after substrate introduction for
metabolite analysis. Each biological replicate consisted of leaves from independent *N. benthamiana* plants (n≥3).

386
Heterologous expression of candidate genes in *E. coli* BL21 (DE3)
Chemically competent *E. coli* BL21 (DE3) cells (AlpaLifeBio) were transformed with pOPINF or pOPINM
plasmids containing the coding sequences of interest. Transformants were selected on LB agar plates
supplemented with 100 μg/mL ampicillin and verified by colony PCR using vector-specific sequencing primers
(Supplementary Table 3). For each strain, a single colony was used to inoculate 10 mL of LB medium containing
100 μg/mL ampicillin, which was cultured overnight at 37 °C in a temperature-controlled shaker incubator. The
overnight culture was then used to inoculate 400 mL of TB media with 100 μg/mL ampicillin. Cultures were
grown at 37 °C with a shaking of 220 rpm until reaching an OD_{600} of 0.6-0.8. At this point, the temperature was
lowered to 18 °C, and protein expression was induced by adding isopropyl *β*-D-1-thiogalactopyranoside (IPTG)
to a final concentration of 0.5 mM. After approximately 16 hours of induction, the cultures were centrifuged at
8,000 rpm at 4 °C for 10 min to harvest the cells.
398
Protein purification
Bacterial cells were resuspended in lysis buffer (100 mM Tris-HCl, 100 mM NaCl, 5% glycerol (*v/v*), pH 8.0)
and disrupted using a high-pressure cell disruption equipment (ATS Engineering, Suzhou, China). The remaining
steps were performed at 4 °C or on ice. The cell lysates were clarified by centrifugation 12,000 rpm for 30 min,
and the supernatant (crude enzyme) was collected. The clarified lysate was loaded onto a column packed with
Ni-NTA Beads 6FF agarose resin (Smart-Lifesciences). The column was washed with Tris-HCl buffer before
stepwise elution using increasing concentrations of imidazole (30−300 mM) in Tris-HCl buffer. Elution fractions
were collected separately and analyzed by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-
PAGE) with Coomassie Blue staining. Fractions containing the purified protein were combined and concentrated
using Amicon Ultra-15 Centrifugal Filter Units (10 kDa molecular weight cutoff) via centrifugation at 3,500
rpm for 1 h at 4 °C. The concentrated protein solution (<0.5 mL) underwent two rounds of buffer exchange with
ice-cold HEPES buffer (50 mM HEPES, pH 7.5, 100 mM NaCl) to remove imidazole. The final protein
preparation was diluted to approximately 2 mL with ice-cold HEPES buffer. Protein concentrations were
determined using a NanoDrop spectrophotometer. Aliquots of the purified protein were flash-frozen in liquid
nitrogen and stored at −80 °C until further use.
414
Enzymatic in vitro assays
Enzymatic assays for reductase activity
The reductase activity of recombinant *Rv*YOS was assessed using tryptamine and secologanin as substrate.
Reaction mixtures (100 μL total volume) contained 1 μM *Cr*STR, 1 μM *Rv*SGD, 2 μM *Rv*YOS, 100 μM NADPH,
25 μM tryptamine, and 25 μM secologanin in 50 mM HEPES (pH 7.4). The reactions were incubated at 30 °C
with shaking at 500 rpm for 8 h. The reactions were quenched by adding an equal volume of methanol, followed
by vacuum concentration to dryness. The residues were reconstituted in 100 μL methanol and centrifugation at
15,000 × g for 10 min. The supernatants were filtered through 0.22 μm PTFE syringe filters for LC-MS analysis
(LC-MS Method 1).
424
The reductase activity of recombinant *Rv*DYR1 and *Rv*DYR2 was assessed using 3-dehydro-α-yohimbine (**3**)
as substrate. Reaction mixtures (50 μL total volume) contained 1 μM purified reductase, 100 μM NADPH, and
40 μM **3** in 50 mM HEPES (pH 7.4). Reactions were incubated at 30°C with shaking at 500 rpm for 12 h. Boiled
enzymes (90 °C, 10 min) served as negative controls. Reactions were quenched by adding an equal volume of
methanol, followed by centrifugation at 15,000 × g for 10 min. The supernatants were filtered through 0.22 μm
PTFE syringe filters for LC-MS analysis (LC-MS Method 2).
431
Enzymatic assays for *O*-methyltransferase activity
The *O*-methyltransferase activity of recombinant *Rv*11OMT candidate was assessed using 50 μM 11,18β-
hydroxy-3-epi-α-yohimbine (**6**) as substrate. Reaction mixtures (50 μL total volume) contained 1 μM *Rv*11OMT,
250 μM *S*-adenosyl methionine (SAM), and 200 μM ascorbate in 50 mM HEPES (pH 7.4). Reactions were
incubated at 30 °C with shaking at 500 rpm for 12 h and quenched by adding an equal volume of methanol,
followed by centrifugation at 15,000 × g for 10 min. Sample preparation for LC-MS analysis was performed as
described above for reductase activity assays.

<u>Enzymatic assays for UGTs activity</u>
The UGT activity of recombinant *Rv*UGT1 and *Rv*UGT2 was assessed using trimethylgallic acid and UDP-
glucose as substrate. Reaction mixtures (100 μL total volume) contained 2 μM purified enzymes, 50 μM
trimethylgallic acid, and 25 μM UDP-glucose in 50 mM HEPES (pH 7.4). The reactions were treated as for
reductase activity assays, and analysis was performed as described in the LC-MS analysis (LC-MS Method 3).

Heterologous expression of *Rv*CYP71D820 in *Saccharomyces cerevisiae*
The full-length coding sequence of *Rv*CYP71D820 and *Arabidopsis thaliana* NADPH-cytochrome P450
reductase (*At*CPR) were cloned into the yeast expression vector pESC-Ura under the control of GAL1 and
GAL10, respectively. The resulting construct was transformed into *S. cerevisiae* WAT11 chemically competent
cells (Coolaber, Beijing, China) using the lithium acetate/single-stranded carrier DNA/polyethylene glycol
(LiAc/SS carrier DNA/PEG) method as described previously[45]. Transformants were selected on synthetic
defined medium lacking uracil (SD-Ura). Positive clones were identified by colony PCR using vector-specific
primers listed in Supplementary Table 4.

Single colonies of recombinant *S. cerevisiae* strains were cultured in 10 ml of SD-Ura liquid medium containing
2% glucose and grown at 30 °C for 24 h with a shaking of 220 rpm. To induce protein expression, the cells were
harvested by centrifugation (1,000 rpm, 5 min) at room temperature and resuspended in 500 mL of SD-Ura
medium supplemented with 2% glucose and 1.8% galactose. The cultures were then incubated at 30 °C with a
shaking of 220 rpm for approximately 3-4 days. Yeast cells were harvested by centrifugation (1,000 g, 5 min)
for microsomal protein preparation.

Preparation of *Rv*CYP71D820-containing microsome
Isolation of yeast microsomal protein followed a previously described protocol[46]. The cell pellet was
resuspended in TEK buffer (50 mM Tris-HCl, 1 mM EDTA, 100 mM potassium chloride, pH 7.4) at a ratio of
1 mL buffer per 0.5 g wet cell pellet mass and incubated at room temperature for 5 min. Cells were harvested by
centrifugation (1,000 g, 5 min) and then resuspended in 50 mL ice-cold TES B buffer (50 mM Tris-HCl, 1 mM
EDTA, 600 mM sorbitol, pH 7.4). The cell suspensions were lysed using an ultra-high-pressure homogenizer
(1100 bar, 5 min, 4 °C). Cell debris was removed by centrifugation (10,000 g, 15 min, 4 °C). The supernatant
was further centrifuged at 87,000 × g for 60 min at 4 °C. The resulting microsomal pellet was resuspended in 2
mL of ice-cold TEG buffer (50 mM Tris-HCl, 1 mM EDTA, 20% [*v/v*] glycerol, pH 7.4). The protein
concentration in the microsomal fraction was measured using the Bio-Rad Protein Assay Kit. Aliquots (100 μL)
were snap-frozen in liquid nitrogen and stored at −80 °C until further use.

***In vitro* assays of *Rv*CYP71D820**
The enzymatic activity of *Rv*CYP71D820 was assessed in a reaction mixture (100 μL final volume) containing
40 μM substrate, 2 mM NADPH, 50 mM phosphate buffer (pH 8.0), and 30 μL microsomal fraction. The reaction
mixture was incubated at 30 °C for 16 h. Reactions were terminated by the addition of 200 μL methanol. The
reaction mixture was concentrated to dryness using a vacuum concentrator (Eppendorf, Hamburg, Germany) at
30 °C. The residue was redissolved in 100 μL methanol and subjected to analysis.

Metabolite extraction
For *R. serpentina* tissue analysis, freeze-dried tissue was ground to a fine powder, and 10 mg was accurately
weighed into a microcentrifuge tube. The tissue was extracted with 300 μL methanol in an ultrasonic bath (SB-
3200DT, Scientz, Ningbo, China) at room temperature for 30 min. The extracts were centrifuged at 15,000 rpm
for 5 min at room temperature, and the supernatant was filtered through a 0.22 μm PES syringe filter into HPLC
vials for analysis.

For *Nicotiana benthamiana* samples, leaves were collected with a 1.0-cm diameter cork borer. The leaf discs
were transferred to 2 mL microcentrifuge tubes and lyophilized overnight in a freeze dryer (SCIENTZ-10N,
Ningbo Scientz Biotechnology Co., Ltd). The dried tissue was extracted with 600 μL methanol and homogenized

using a homogenizer (SWE-D6, Servicebio, Hubei, China). The homogenate was centrifuged at 15,000 rpm for
10 min, and the supernatant was filtered through a 0.22 μm PES syringe filter into HPLC vials for analysis.
493
HPLC analysis and purification
Analytical high performance liquid chromatography (HPLC) was performed on an Agilent 1260 Series liquid
chromatography system equipped with a multiple wavelength detector and a COSMOSIL 5C18-MS-II column
(250 mm × 4.6 mm, 5 μm particle size). The injection volume was 10 μL and the flow rate was 1.0 mL/min
using 0.1% formic acid in water as mobile phase A and acetonitrile as mobile phase B. Samples were analyzed
with the following liquid chromatography gradient: 0-5 min, 10% B isocratic; 5-25 min, 10% to 50% B; 25-30
min, 50% to 95% B;30-40 min, 10% B isocratic.
501
Semi-preparative HPLC was carried out using the same Agilent 1260 Series system equipped with either a
COSMOSIL 5C18-MS-II column (250×10 mm, 5 μm particle size) or an XBridge BEH C18 OBD Prep column
(250 mm ×10 mm, 5 μm particle size, , 130Å pore size). The injection volume was 50 μL and the follow rate
was 3 mL/min using 0.1% formic acid in water as mobile phase A and acetonitrile as mobile phase B. For
compound isolation, the following liquid chromatography gradient was applied: 0-10 min, 10% B isocratic; 10-
40 min, 10% to 50% B; 40-50 min, 50% to 95% B; 50-60 min, 10% B isocratic. The fractions were collected
based on UV absorption at 254 nm.
509
LC-MS analysis
Method 1
Liquid chromatography-mass spectrometry (LCMS) analysis was performed using a SIL-30AC HPLC system
(SHIMADZU, Kyoto, Japan) coupled to a ZenoTOF 7600 System (Ultra-High-Resolution Quadruple-Time-
of-Flight) mass spectrometer (AB SCIEX, Framingham, MA, USA). Chromatographic separation was achieved
using a Phenomenex Kinetex XB-C18 column (100 mm × 2.1 mm, 1.7 μm particle size; 100 Å pore size)
maintained at 40 °C. The injection volume was 3 μL and the flow rate was 0.3 mL/min using 0.1 % formic acid
in water as mobile phase A and acetonitrile as mobile phase B. Authentic standards were prepared in methanol
at concentrations ranging from 20-100 μM. The liquid chromatography gradient elution program was as follows:
0-5 min, 5 % B isocratic; 5-15 min, 5 % B to 50 % B; 15-20 min, 50 % B to 95 % B; 20-22 min, 95 % B isocratic;
22-25 min, 5 % B isocratic. Mass spectrometry was performed in positive electrospray ionization mode using
data-independent acquisition (DIA). Operating parameters were as follows: capillary voltage, 3500 V; end plate
offset, 500 V; nebulizer pressure, 2.5 bar; drying gas (N_2) temperature, 250 °C; drying gas flow rate, 11 L/min.
Mass spectra were acquired over the range *m/z* 50 to 1000. For MS/MS analysis, fragmentation was triggered at
an absolute threshold of 100 with a total cycle time of 0.3-0.5 s and collision energy of 35 eV.
525
Method 2
Additional LC-MS analysis was performed using a Vanquish liquid chromatography system (Thermo Fischer
Scientific) connected to an IMPACT II System (High Resolution Quadrupole-Time-of-Flight) mass
spectrometer (Bruker). Separation was achieved using a Waters Acquity BEH C18 column (100 mm × 2.1 mm,
2.5 μm particle size) operated at 40 °C with a flow rate of 0.3 mL/min and injection volume of 3 μL. The mass
spectrometry was operated in positive electrospray ionization mode (ESI^+) with the following parameters:
capillary voltage, 3500 V; end plate offset, 500 V; nebulizer pressure, 2.5 bar; drying gas (N_2) temperature,
250 °C; drying gas flow rate, 8 L/min. Data acquisition was performed in autoMS/MS mode at 12 Hz over *m/z*
50-1000. Fragmentation was triggered on an absolute threshold of 500 with a cycle time of 0.5 s. Collision
energy was applied using a stepping gradient either from 11.0 to 44.0 eV or from 35.0 to 52.5 eV.
536
Method 3
Additional LC-MS analysis was performed using a Vanquish liquid chromatography system (Thermo Fischer
Scientific) connected to an IMPACT II System (High Resolution Quadrupole-Time-of-Flight) mass
spectrometer (Bruker). Separation was achieved using a Waters Acquity BEH C18 column (100 mm × 2.1 mm,
2.5 μm particle size) operated at 40 °C with a flow rate of 0.3 mL/min and injection volume of 3 μL. The mass
spectrometry was operated in positive electrospray ionization mode (ESI^-) with the following parameters:
capillary voltage, 4000 V; end plate offset, 500 V; nebulizer pressure, 2.5 bar; drying gas (N_2) temperature,

250 °C; drying gas flow rate, 8 L/min. Data acquisition was performed in autoMS/MS mode at 12 Hz over *m/z*
50-1000. Fragmentation was triggered on an absolute threshold of 500 with a cycle time of 0.5 s. Collision
energy was applied using a stepping gradient from 17.5 to 35 eV.
547

Data availability

549 The gene sequences of all characterized enzymes are provided in Supplementary Table 1. The

550 transcriptome data generated in this study have been deposited in the National Genomics Data

551 Center BioProject databased with the accession number PRJCA035431.

References

1. Müller, J. M., Schlittler, E. & Bein, H. J. Reserpin, der sedative Wirkstoff aus *Rauwolfia*
serpentina Benth. *Experientia* 8, **338** (1952).

2. Carlsson, A., Lindqvist, M. & Magnusson, T. 3,4-Dihydroxyphenylalanine and 5-
hydroxytryptophan as reserpine antagonists. *Nature* **180**, 1200 (1957).

3. Brodie, B. B., Olin, J. S., Kuntzman, R. G. & Shore, P. A. Possible interrelationship between
release of brain norepinephrine and serotonin by reserpine. *Science* **125**, 1293-1294 (1957).

4. Vakil, R. J. *Rauwolfia serpentina* in the treatment of high blood pressure: a review of the
literature. *Circulation* **12**, 220-229 (1955).

5. Braslow, J. T. & Marder, S. R. History of Psychopharmacology. *Annu. Rev. Clin. Psychol*. **15**,
25-50 (2019).

6. Carlsson, A. A paradigm shift in brain research. *Science* **294**, 1021-1024 (2001).

7. Chen, F. -E. & Huang, J. Reserpine: a challenge for total synthesis of natural products. *Chem.*
Rev. **105**, 4671-4706 (2005).

8. Woodward, R. B. et al. The total synthesis of reserpine. *J. Am. Chem. Soc*. **78**, 9, 2023-2025
(1956).

9. Chu, C. -S., Liao, C. -C. & Rao, P. D. A formal synthesis of (±)-reserpine from methyl vanillate.
Chem. Commun. 1537-1538 (1996).

10. Mehta, G. & Reddy, D. S. A formal synthesis of reserpine: hydrindane approach to the
Woodward's ring-E precursor. *J. Chem. Soc. Perkin Trans*. **1**, 1399-1404 (2000).

11. Stork, G. The stereospecific synthesis of reserpine. *Pure Appl. Chem*. **61**, 439-442 (1989).

12. Gomez, A. M., Lopez, J. C. & Fraser-Reid, B. Serial radical cyclization of pyranose-derived
dienes in the stereocontrolled synthesis of densely functionalized cyclohexanes. A route to
Woodward's reserpine precursor. *J. Org. Chem*. **59**, 4048-4050 (1994).

13. Leete, E. The Biogenesis of the Rauwolfia alkaloids. I. The incorporation of tryptophan into
ajmaline. *J. Am. Chem. Soc*. **82**, 6338-6342 (1960).

14. Battersby, A. R., Burnett, A. R. Parsons, P. G. Alkaloid biosynthesis. Part XV. Partial
synthesis and isolation of vincoside and isovincoside: biosynthesis of the three major classes of
indole alkaloids from vincoside. *J. Chem. Soc. C* 1193-1200 (1969).

15. O'Connor, S. E. & Maresh, J. J. Chemistry and biology of monoterpene indole alkaloid
biosynthesis. *Nat. Prod. Rep*. **23**, 532-547 (2006).

16. Gerasimenko, I., Sheludko, Y., Unger, M. & Stöckigt, J. Development of an efficient system
for the separation of indole alkaloids by high performance liquid chromatography and its
applications. *Phytochem. Anal*. **12**, 96-103 (2001).

17. Gerasimenko, I. et al. Heterologous expression of a *Rauvolfia* cDNA encoding strictosidine
glucosidase, a biosynthetic key to over 2000 monoterpene indole alkaloids. *Eur. J. Biochem*. **269**,
2204-2213 (2002).

18. Yerkes, N. et al. Substrate specificity and diastereoselectivity of strictosidine glucosidase, a
key enzyme in monoterpene indole alkaloid biosynthesis. *Bioorg. Med. Chem. Lett*. **18**, 3095-3098
(2008).

19. Zhan, G. et al. Monoterpene indole alkaloids with diverse skeletons from the stems of
Rauvolfia vomitoria and their acetylcholinesterase inhibitory activities. *Phytochemistry* **177**,
112450 (2020).

20. Falkenhagen, H. et al. Indole alkaloids from "hairy roots" of *Rauwolfia serpentina. Can. J.
Chem.* **71***, 2201-2203 (1993).

21. Kumar, S., Kumari, D. & Singh, B. Genus *Rauvolfia*: A review of its ethnopharmacology,
phytochemistry, quality control/quality assurance, pharmacological activities and clinical evidence.
J. Ethnopharmacol. **295**, 115327 (2022).

22. Stander, E. A. et al. The *Rauvolfia tetraphylla* genome suggests multiple distinct biosynthetic
routes for yohimbine monoterpene indole alkaloids. *Commun. Biol*. **6**, 1197 (2023).

23. Chen, R. et al. Molecular cloning and characterization of the strictosidine synthase gene
from *Rauwolfia verticillata. Russ. J. Plant Physiol*. **55**, 670–675 (2008).

24. Xu, H. et al. Cloning and characterization of the strictosidine-β-D-glucosidase (SGD) gene
from *Rauvolfia verticillata*. *Agric. Sci. Technol*. **13**, 1406-1409 (2012).

25. Allard, S. T., Giraud, M. F. & Naismith, J. H. Epimerases: structure, function and mechanism.
Cell Mol. Life Sci. **58**, 1650-1665 (2001).

26. Garg, A. et al. Elucidation of the cryptic epimerase activity of redox-inactive ketoreductase
domains from modular polyketide synthases by tandem equilibrium isotope exchange. *J. Am.
Chem. Soc*. **136**, 10190-10193 (2014).

27. Farrow, S. C., Hagel, J. M., Beaudoin, G. A. W., Burns, D. C. & Facchini, P. J. Stereochemical
inversion of (*S*)-reticuline by a cytochrome P450 fusion in opium poppy. *Nat. Chem. Biol*. **11**, 728-
732 (2015).

28. Caputi, L. et al. Missing enzymes in the biosynthesis of the anticancer drug vinblastine in
Madagascar periwinkle. *Science* **360**, 1235-1239 (2018).

29. Langley, C. et al. Expansion of the catalytic repertoire of alcohol dehydrogenases in plant
metabolism. *Angew. Chem. Int. Ed. Engl*. **61**, e202210934 (2022).

30. Trenti, F. et al. Early and late steps of quinine biosynthesis. *Org. Lett*. **23**, 1793-1797 (2021).

31. Schotte, C. et al. Directed biosynthesis of mitragynine stereoisomer. *J. Am. Chem. Soc.* **145**,
4957-4963 (2023).

32. Kim, K. et al. Biosynthesis of kratom opioids. *New Phytol*. **240**, 757-769 (2023).

33. Stavrinides, A. et al. Unlocking the diversity of alkaloids in *Catharanthus roseus*: nuclear
localization suggests metabolic channeling in secondary metabolism. *Chem. Biol*. **22**, 336-341
(2015).

34. McDonald, A. et al. Enzymatic epimerization of monoterpene indole alkaloids in Kratom.
bioRxiv 2024.12.13.628308; doi: https://doi.org/10.1101/2024.12.13.628308.

35. Hwang, J. et al. Ancient gene clusters initiate monoterpene indole alkaloid biosynthesis and C-
3 stereochemistry inversion. bioRxiv 2025.01.07.631695. doi: https://doi.org/10.1101/
2025.01.07.631695.

36. Besseau S. et al. A pair of tabersonine 16-hydroxylases initiates the synthesis of vindoline in
an organ-dependent manner in *Catharanthus roseus*. *Plant Physiol*. **163**, 1792-1803 (2013).

37. St-Pierre, B. & De Luca, V. A Cytochrome P450 monooxygenase catalyzes the first step in the
conversion of tabersonine to vindoline in *Catharanthus roseus*. *Plant Physiol*. **109**, 131-139 (1995).

38. Levac, D., Murata, J., Kim, W. S. & De Luca, V. Application of carborundum abrasion for
investigating the leaf epidermis: molecular cloning of *Catharanthus roseus* 16-
hydroxytabersonine-16-*O*-methyltransferase. *Plant J*. **53**, 225-236 (2008).

39. Murata, J., Roepke, J., Gordon, H. & De Luca, V. The leaf epidermome of *Catharanthus roseus*
reveals its biochemical specialization. *Plant Cell* **20**, 524-542 (2008).

40. Schneider, J. A. et al. Pharmacological studies with deserpidine, a new alkaloid from *Rauwolfia*
canescens. *J. Pharmacol. Exp. Ther*. **114**, 10-13 (1955).

41. Lam, K. C., Ibrahim, R. K., Behdad, B. & Dayanandan, S. Structure, function, and evolution
of plant *O*-methyltransferases. *Genome* **50**, 1001-1013 (2007).

42. Bontpart, T., Cheynier, V., Ageorges, A. & Terrier, N. BAHD or SCPL acyltransferase? What

a dilemma for acylation in the world of plant phenolic compounds. *New Phytol*. **208**, 695-707
(2015).

43. Li, P. et al. Multiomics analyses of two *Leonurus* species illuminate leonurine biosynthesis
and its evolution. *Mol. Plant* **17**, 158-177 (2024).

44. Zhao, Y. et al. Flavano-3-ol galloylation-related functional gene cluster and the functional
diversification of *SCPL* paralogs in *Camellia* sp. *J. Agric. Food Chem*. **71**, 488-498 (2023).

45. Gietz, R. D. &Woods, R. A. Transformation of yeast by lithium acetate/single-stranded carrier
DNA/polyethylene glycol method, *Method. Enzymol.* **350**, 87-96 (2002).

46. Pompson, D., Louerat, B., Bronine, A. & Urban, P. Yeast expression of animal and plant P450s
in optimized redox environments. *Methods Enzymol*. **272**, 51-64 (1996).
655

Acknowledgements

We thank Dr. David Nelson (University of Tennessee) for his assistance with naming

*Rv*CYP72A270 and *Rv*CYP71D820 identified in this study. We are grateful to the Shenzhen

Synthetic Biology Infrastructure for instrument support and technical assistance. This research was

supported by the National Key R&D Program of China (2023YFA0915500) and Shenzhen Science

and Technology Program (KJZD20230923115906013) to Y. J, National Natural Science

Foundation of China (82104023) to J. C.
663

664 **Author contributions**

665 J. C., J. Z, and F. L. performed the experiments and analyzed the data. J. C. and Y. J. designed the

666 experiments, wrote and revised the manuscript. Y. J. supervised the project. All authors discussed

667 the results and commented on the manuscript.
668

669 **Competing interests**

670 The authors declare no competing interests.
671

Fig. 1: Structural features, synthetic history, and biosynthetic pathway of reserpine.

a, Chemical structure of reserpine (**11**) highlighting its key structural features. **b**, Previous synthetic approaches exemplified by Woodward's synthesis. **c**, The biosynthetic pathway of reserpine identified in this work. TMB represents 3,4,5-trimethoxybenzoyl group.

Fig. 2: The biosynthetic pathway of reserpine in *R. verticillata*.

Solid arrows indicate characterized steps; dashed arrows indicate proposed steps. Names of characterized enzymes are shown in blue. AT represents acyltransferase.

Fig. 3: Identification and characterization of enzymes involved in α-yohimbine epimerization.

a, Proposed biosynthetic pathway from tryptamine and secologanin to 3-epi-α-yohimbine (**4**) via
α-yohimbine (**2**) and 3-dehydro-α-yohimbine (**3**) intermediates. **b**, Phylogenetic analysis of
medium-chain dehydrogenase/reductase (MDR) enzymes, with colored boxes indicating different
reaction types catalyzed by characterized enzymes. Bootstrap values are shown as circles of
varying sizes. **c**, LC-MS analysis showing oxidation of α-yohimbine (**2**) by *Rv*YOO expressed in
N. benthamiana leaves compared to empty vector (EV) control and standard (ST) of **3**. **d**, LC-MS
analysis demonstrating conversion of α-yohimbine (**2**) to 3-epi-α-yohimbine (**4**) by co-expression
of *Rv*YOO with *Rv*DYR1 or *Rv*DYR2 in *N. benthamiana*. **e**, In vitro characterization of purified
*Rv*DYR1 and *Rv*DYR2 showing stereoselective reduction of synthetic 3-dehydro-α-yohimbine (**3**)
to 3-epi-α-yohimbine (**4**) in the presence of NADPH.

Fig. 4: Discovery and characterization of enzymes responsible for C11 and C18 hydroxylation and *O*-methylation.

a, Proposed biosynthetic pathway from 3-epi-α-yohimbine (**4**) to rauvomitorine G (**9**) via sequential hydroxylation and *O*-methylation reactions. **b**, Tissue-specific expression patterns of biosynthetic genes in *R. verticillata* stem, leaf, and root tissues; **c**, Correlation matrix showing Pearson correlation coefficients between expression levels of identified biosynthetic genes. **d**, LC-MS analysis of *Rv*CYP71D820-catalyzed C18 hydroxylation of **4** in *N. benthamiana* (leaf icon) and *in vitro* using yeast microsomes (tube icon), compared with synthetic standard of **5**. **e**, LC-MS analysis showing formation of 11,18β-dihydroxy-3-epi-α-yohimbine (**6**) through co-expression of *Rv*CYP71D820 and *Rv*CYP72A270 *in N. benthamiana*, compared with synthetic standard of **6**. **f**, LC-MS analysis of *Rv*11OMT-catalyzed *O*-methylation of **6** in *N. benthamiana* and *in vitro* using purified enzyme, compared with synthetic standard of **9**. **g**, LC-MS analysis of the reconstitution of **9** from **4** through co-expression of *Rv*CYP71D820, *Rv*CYP72A270, and *Rv*11OMT in *N. benthamiana*. EV, empty vector control; ST, synthetic standard.

Fig. 5: Reconstitution of rauvomitorine G biosynthesis.

a, Reconstitution of the complete biosynthetic pathway in tobacco using eight enzymes with
tryptamine and secologanin as substrates. Extracted ion chromatograms (EIC) show the detection
of intermediates **3**, **4**, **5** and final product **9**. **b**, Stepwise reconstitution and validation of the
biosynthetic pathway from α-yohimbine (**2**) to rauvomitorine G (**9**), showing the sequential
enzymatic transformations and corresponding intermediates. Bar graphs indicate peak areas of
pathway intermediates (n = 3 biological replicates, error bars represent standard deviation).