Affinity-guided labeling reveals P2X7 nanoscale membrane
redistribution during microglial activation.

Benoit Arnould,[1,†] Adeline Martz,[1,‡] Pauline Belzanne, [2,‡] Francisco Andrés Peralta[1,3] Federico
Cevoli[1], Volodya Hovhannisyan[4], Yannick Goumon[4], Eric Hosy[2], Alexandre Specht[5], and Thomas
Grutter[1,3,*].

[1]Laboratoire de Chémo-Biologie Synthétique et Thérapeutique (CBST) UMR 7199, équipe
Ingénierie Canaux Ioniques, Centre National de la Recherche Scientifique, Université de
Strasbourg, Faculté de Pharmacie, F-67400 Illkirch, France. [2]Interdisciplinary Institute for
Neuroscience, CNRS, Université de Bordeaux, IINS, UMR 5297, Bordeaux, France. [3]University
of Strasbourg Institute for Advanced Studies (USIAS), 67000 Strasbourg, France. [4]Centre
National de la Recherche Scientifique-Unité Propre de Recherche (CNRS-UPR) 3212, Institut
des Neurosciences Cellulaires et Intégratives, Unistra, Strasbourg, France. [5]Laboratoire de
Chémo-Biologie Synthétique et Thérapeutique (CBST) UMR 7199, équipe NanoParticules
Intelligentes, Centre National de la Recherche Scientifique, Université de Strasbourg, Faculté de
Pharmacie, F-67400 Illkirch, France.

†Present address: Department of Chemistry, Washington University in St. Louis, St. Louis,
Missouri 63130, USA.

‡These authors contribute equally.

*To whom correspondence should be addressed.

E-mail: grutter@unistra.fr).

Author Contributions: Conceptualization: TG; Data curation: PB, EH; Formal analysis: BA, AM,
PB, FAP, EH, TG; Funding acquisition: TG; Investigation: BA, AM, PB, FAP, FC; Methodology:
TG, BA, AM, PB; Project administration: TG; Resources: VH, YG; Supervision: TG, AS, EH;
Visualization: TG; Writing—original draft: TG; Writing—review & editing: TG, BA, AS, EH.

Competing Interest Statement: Authors declare that they have no competing interests.

Keywords: purinergic receptors; inflammation; super-resolution microscopy; single-molecule.

This PDF file includes:

Main Text
Figures 1 to 6
32

34
35

Abstract

ATP-gated purinergic P2X7 receptors are crucial ion channels involved in inflammation. They sense abnormal ATP release during stress or injury and are considered promising clinical targets for therapeutic intervention. However, despite their predominant expression in immune cells such as microglia, there is limited information on P2X7 membrane expression and regulation during inflammation at the single-molecule level, necessitating new labeling approaches to visualize P2X7 in native cells. Here, we present **X7-uP**, an unbiased, affinity-guided P2X7 chemical labeling reagent that selectively biotinylates endogenous P2X7 in BV2 cells, a murine microglia model, allowing subsequent labeling with streptavidin-Alexa 647 tailored for super-resolution imaging. We uncovered a nanoscale microglial P2X7 redistribution mechanism where evenly spaced individual receptors in quiescent cells undergo upregulation and clustering in response to the pro-inflammatory agent lipopolysaccharide and ATP, leading to synergistic interleukin-1β release. Our method thus offers a new approach to revealing endogenous P2X7 expression at the single-molecule level.

Introduction

ATP serves as the primary energy carrier in cells and also functions as an extracellular signaling molecule. Normally, extracellular ATP levels are typically low in healthy interstitial fluid, but they rise at sites of stress or cellular injury, acting as a danger signal recognized as a damage-associated molecular pattern (DAMP). Extracellular ATP is detected by two families of purinergic receptors, metabotropic P2Y receptors and ionotropic P2X receptors (P2X1 to P2X7), with P2X7 specifically sensing abnormal ATP release in the mid-micromolar to low millimolar range (1-3). P2X7 is a transmembrane, non-selective cation channel (Na^+, K^+, Ca^{2+}) predominantly expressed in immune cells, such as macrophages and microglia (the resident macrophages of the central nervous system), where it plays a crucial role in inflammation and immunity (4, 5). Upon ATP binding, P2X7 activation triggers K^+ efflux, promoting the assembly of the NLRP3 inflammasome and leading to the maturation and secretion of pro-inflammatory mediators, including interleukin-1β (IL-1β) (6-11). P2X7 activation can cause cell death and is linked to the pathogenesis of several major disorders, including inflammatory pain (12), autoimmune diseases (13), tumor progression (14, 15), psychiatric conditions (16), anxiety-like behavior (17), and neurodegenerative diseases (18, 19), making P2X7 a clinically relevant target.

At the cellular level, P2X7 is expressed at the plasma membrane and associates with lipid rafts, which are membrane microdomains known to play a key role in the onset of inflammation (20-25). However, it is unknown how P2X7 is strictly distributed on the plasma membrane of cells and how its spatial and functional features intertwine in inflammatory conditions. Given the crucial role of P2X7 in inflammation and the growing interest in developing P2X7-targeted therapeutics, new approaches are needed to visualize P2X7 expression in immune cells at the single molecule level.

Numerous studies have investigated P2X7 expression in microglia using various experimental approaches, including functional assays (6, 26-28), radiolabeled ligand binding (29), immunofluorescence staining (30, 31), electron microscopy (32), and genetic manipulations (33). Although these studies confirmed the dominant expression of P2X7 in microglia, they provided only low-resolution cell imaging of P2X7, primarily due to the limitations of conventional light microscopy, which imposes a resolution restricted by the diffraction of light. Consequently, the detailed organization of P2X7 below ~250 nm remains unknown. This is a vital point, as recent data suggest that the spatial organization of cell membrane receptors may be a common regulatory mechanism for cellular signal transduction (34). However, it remains unclear whether such membrane organization similarly affects P2X7 signaling pathways.

Super-resolution microscopy techniques overcome this diffraction limit, enabling the visualization of fluorescently labeled membrane receptors at nanometric resolution (35, 36). To the best of our

knowledge, only two studies have reported super-resolved images of P2X7 (37, 38). The first study
found P2X7 in lipid microdomains of an osteoblastic cell line (38), but the use of polyclonal
antibodies to stain P2X7 raised questions about their selectivity (39). The second study involved
fusing P2X7 to the photo-convertible fluorescent Dendra2 protein to track its dynamic organization
in hippocampal neurons (37). While this study provided valuable insights into the nanometer-scale
distribution of P2X7, it had three experimental biases. First, heterologous overexpression of P2X7-
Dendra2 might not reflect native expression levels. Second, the C-terminal fusion of Dendra2 to
P2X7 after the "ballast", a region that is critically involved in inflammation (5), could affect P2X7's
pathogenicity. Third, there is growing evidence suggesting that neurons do not (or scarcely)
express P2X7 (33), questioning the physiological relevance of investigating P2X7 nanoscale
organization in hippocampal neurons. Therefore, these issues prompted us to develop a new,
genetic-free strategy able to decorate endogenous P2X7 in native cells with fluorophores suitable
for super-resolution imaging.
102
Among existing methods for labeling endogenous proteins in living cells, the affinity-driven reaction
strategy involving protein-ligand interaction is of particular interest (40). It relies on the formation of
a covalent bond on the protein that is guided by the affinity of a ligand bearing a reactive moiety.
The efficiency of the labeling reaction depends on the chemical function of the reactive species.
The recently developed *N*-acyl-*N*-alkyl sulfonamide (NASA) chemistry offers a suitable platform
that enables the covalent transfer, under native cell conditions, of functional ligands to nucleophilic
amino acid residues (typically lysine residues) located in close proximity to the ligand-binding site
(41) (Figure 1A). In addition, NASA electrophilicity can be finely tuned by introducing of an electron-
withdrawing moiety to the *N*-alky group, such as cyanomethyl, so that kinetic labeling can be
considerably enhanced and carried out within minutes under physiological conditions (41).
113
Here, we report **X7-uP**, a P2**X7-u**nbiased **P**urinergic labeling reagent that efficiently biotinylates
native P2X7 lysine residues via a *N*-cyanomethyl NASA-based chemical reaction driven by the
affinity of AZ10606120, a highly selective P2X7 allosteric ligand. Using streptavidin-Alexa Fluor®
647 (Strept-A 647), we reveal the nanoscale distribution of P2X7 in BV2 cells at the single molecule
level by direct stochastic optical-reconstruction microscopy (dSTORM). We show that P2X7 is
homogenously dispersed on the plasma membrane as single receptors in quiescent (non-
inflammatory condition) microglia. However, upon exposure to the pro-inflammatory agent
lipopolysaccharide (LPS), a pathogen-associated molecular pattern (PAMP) from Gram-negative
bacteria, combined with ATP or its potent synthetic analog BzATP, P2X7 undergoes significant
upregulation and clustering, resulting in the synergistic release of IL-1β. Our findings suggest that
the dynamic clustering of P2X7 on the plasma membrane is a crucial mechanism underlying IL-1β
secretion.
126
127
128 **Results**
129
Design and synthesis of X7-uP
X7-uP was rationally designed by leveraging available X-ray structures of the panda P2X7 receptor
(pdP2X7) bound to non-competitive P2X7 antagonists (42). These antagonists bind to an interfacial
allosteric binding pocket in pdP2X7, distinct from ATP-binding sites, and accessible from the upper
side of the trimeric receptor (42). Due to the trimeric assembly of P2X subunits, three allosteric
binding sites are present in P2X7 (Figure 1B). We focused on AZ10606120, a selective antagonist
with high-affinity for human (hP2X7) and rat P2X7 (rP2X7), with no effect on other P2X receptors
(43-46). The X-ray structure revealed several lysine residues localized in proximity to AZ10606120
(< 15 Å, measured from α-carbon of K72, K81, K82, K110 and K300, and the solvent-exposed
hydroxyl of AZ10606120) (Figure 1B inset). Since most of these lysine residues are unique to P2X7
(Figure 1 — figure supplement 1A), we hypothesized that NASA chemistry could selectively label
P2X7. We modified AZ10606120 by replacing its solvent-exposed hydroxyethyl group with the
reactive *N*-cyanomethyl NASA derivative, allowing a labeling reaction with the ε-amino group of

nearby P2X7 lysine residues, forming a stable amide bond (Figure 1A). We used molecular docking
to identify molecule **1** as a promising molecular scaffold built from AZ10606120 (Figure 1 — figure
supplement 1B). The distance separating the NASA electrophilic carbon of **1** and each α-carbon of
nearby P2X7 residues was compatible with a proximity-driven reaction with rP2X7 ($\leq$ 16.1 Å, Figure
1 — figure supplement 1C), the receptor species used to experimentally validate the approach (see
below). Thus, we designed and synthesized **X7-uP**, an extended version of **1**, containing an OEG
(olygoethylene glycol) biotin tag (Figure 1C). **X7-uP** is expected to covalently transfer the biotin tag
to P2X7, which, in turn, can serve as a versatile platform for subsequent cell surface labeling with
commercially available, cell-impermeable biotin-binding protein-conjugated molecular probes,
including Alexa 647, suitable for super-resolution microscopy (Figure 1A). **X7-uP** was synthesized
as described in Methods (Figure 1 — figure supplement 2).
154
X7-uP inhibits and biotinylates heterologously expressed rP2X7 in HEK293T cell
To address possible loss of affinity due to chemical modifications of the pharmacophore, we first
assessed the ability of **X7-uP** to bind to P2X7 by measuring agonist-evoked current inhibition using
whole-cell patch-clamp electrophysiology. In HEK293T cells transiently transfected with rP2X7, we
recorded inward currents induced by eight successive applications of 10 μM 2'(3')-*O*-(4-
benzoylbenzoyl)ATP (BzATP), a potent P2X7 agonist (Figure 2A). As expected, currents facilitated
upon successive BzATP applications, a hallmark feature of P2X7 (47). Application of 1 μM of **X7-
uP** for 10 s after washout of the fourth BzATP application did not induce inward currents but strongly
inhibited currents induced by co-application for 2 s with BzATP (79.3 $\pm$ 5.6%, mean $\pm$ standard
error of the mean (s.e.m.), n = 4; control without inhibitors: -15.2 $\pm$ 4.4%, n = 4), as expected for a
potent antagonist. Inhibition was reversible, as BzATP-evoked currents fully restored upon **X7-uP**
washout (Figure 2A). Compared to the parental AZ10606120 compound, for which full inhibition
was observed (99.7 $\pm$ 0.1%, n = 4) at 1 μM (Figure 2A and B), these data suggest that the chemical
modification of AZ10606120 has a minimal impact on the inhibition of P2X7 by **X7-uP**.
169
To test labeling efficiency, HEK293T cells were transiently transfected with c-myc tagged rP2X7
(P2X7c-myc) and incubated at 20°C with 1 μM **X7-uP** for 60 min. **X7-uP** labeling was performed
under physiological conditions (1 μM for 10 min at 37°C in FBS-free DMEM, see Methods), followed
by extensive washing to remove excess **X7-uP**. After cell lysis, biotinylated proteins were pulled
down with NeutrAvidin Agarose, separated on SDS-PAGE, and revealed with an anti-c-myc
antibody in Western blot analysis. A strong and unique band was detected in the Western blot at
the expected apparent molecular mass of monomeric P2X7c-myc (~75-80 kDa) (Figure 2C).
However, this band disappeared upon co-incubation with AZ10606120 (10 μM), or in the absence
of **X7-uP** (lane 1), while robust signals corresponding to P2X7c-myc and β-actin expression were
still detected in the corresponding input controls (Figure 2C). Similarly, no signal was detected in
non-transfected cells treated with **X7-uP** (Figure 2 — figure supplement 1A). To further
demonstrate the pharmacological specificity of P2X7 labeling by **X7-uP**, we used another non-
competitive P2X7 antagonist, A740003 (10 μM), which targets the same allosteric binding cavity
as AZ10606120 in pdP2X7 but is chemically different from AZ10606120 (48). As expected, no
labeling was detected when A740003 was co-incubated with **X7-uP** (Figure 2C). Taken together,
these data strongly suggest that **X7-uP** biotinylates P2X7 residues by targeting the unique P2X7
allosteric binding site.
187
We next conducted a kinetic analysis of P2X7 labeling by incubating cells with **X7-uP** at different
concentrations and monitoring labeling through quantitative Western blotting analysis. The time
course of P2X7 labeling at 1 μM of **X7-uP** is shown in Figure 2C, D, with labeled bands detected
as early as 5 min of incubation time. The data were fitted to Eq. (1), derived according to a kinetic
model in which an irreversible chemical reaction follows a reversible ligand binding reaction in a
large excess of ligand (see Methods) (41), providing the pseudo-first-order reaction rate (k_{app}) of
the labeling reaction. Kinetics of labeling carried out at different **X7-uP** concentrations allowed us
to plot k_{app} values against **X7-uP** concentrations (Figure 2 — figure supplement 1B-D). The data

were fitted to Eq. (2), providing the labeling rate constant k_L (0.011 ± 0.003 s^{-1}, mean of triplicate ±
standard deviation (s.d.)), the dissociation constant K_d (7.3 ± 2.7 μM, mean ± s.d.), and the second-
order rate constant (k_L/ K_d = 1.5 × 10^3 M^{-1}s^{-1}). The second-order rate constant value was in the
same order of magnitude as those of previously described labeling reagents based on NASA
chemistry (41).

Next, we determined **X7-uP** labeling yield by quantifying the amount of P2X7c-myc in the
supernatant after pulling down biotinylated P2X7c-myc (Figure 2 — figure supplement 2A). This
amount, corresponding to unbiotinylated P2X7c-myc, was then compared to the maximal amount
of unbiotinylated P2X7c-myc carried out in the absence of **X7-uP**, thus providing labeling yield.
Quantitative Western blotting analysis revealed that 65 ± 3% (mean of triplicate ± s.e.m.) of P2X7c-
myc were labeled by 1 μM of **X7-uP** for 60 min in HEK293T cells (Figure 2 — figure supplement
2B).

X7-uP labeling is selective for P2X7
Having established the efficacy of P2X7 labeling by Western blot, we next assessed the selectivity
of **X7-uP** towards the P2X family. We performed **X7-uP** labeling in HEK293T cells transiently
transfected with various fluorescently tagged rat P2X subunits (P2X1 to P2X6). The biotinylated
cells were then visualized using Strept-A 647 labeling through confocal microscopy. These P2X
subtypes are closely related to P2X7 but are not expected to exhibit the AZ10606120 allosteric site.

As expected, confocal microscopy revealed a strong Alexa 647 fluorescence signal in the periphery
of cells expressing only P2X7 tagged with the monomeric Scarlet (mScarlet) fluorescent protein
(Figure 3A). In contrast, no Alexa 647 signal was detected in cells transiently transfected with other
P2X subunits tagged with GFP, although a strong GFP signal was observed for all constructs,
confirming correct expression (Figure 3B and C). Furthermore, co-incubation with AZ10606120 (10
μM) or A740003 (10 μM) during **X7-uP** labeling abolished the Alexa 647 signal, but did not affect
mScarlet signal, further confirming the specificity of P2X7 labeling (Figure 3A and C). These data
demonstrate the high selectivity of **X7-uP** for P2X7 labeling.

To further validate our method for single-molecule localization microscopy, we employed dSTORM
to visualize the fluorescence emission of individual Alexa 647 fluorophores bound to P2X7
expressed on the plasma membrane of HEK293T cells. Our results confirmed the expression of
P2X7 at the cell surface, consistent with confocal microscopy data (Figure 3 — figure supplement
1A). Importantly, no emission was detected when labeling was conducted in the presence of
AZ10606120 (10 μM), or when Strept-A 647 was added to cells not treated with **X7-uP** (Figure 3
— figure supplement 1B and C), indicating that the observed emission in dSTORM specifically
originated from P2X7.

X7-uP labels K82 and K117 in rP2X7
We next identified the labeling site of **X7-uP**. Since it has been shown that the NASA reagent
preferentially reacts with the ε-amino group of lysine side chains (41), we individually mutated K82,
K110, K117, and K300 into alanine, an amino acid residue that cannot react with NASA, in the
P2X7-mScarlet background. These residues were selected based on the predicted proximity of
their equivalent positions in pdP2X7 to the electrophilic NASA group of **1** in pdP2X7 (Figure 4A and
Figure 1 — figure supplement 1A). We expressed these mutants individually in HEK293T cells and
monitored **X7-uP** labeling by quantifying Alexa 647 signals through confocal microscopy. While
K110A and K300A mutants did not affect **X7-uP** labeling, where Alexa 647 fluorescence
quantification showed comparable levels to that of P2X7-mScarlet (Figure 4C and Figure 4 — figure
supplement 1), a significant reduction in Alexa 647 fluorescence was observed for K82A (79.6 ±
1.0 %, n = 154 cells, P < 0.0001, t-test to the respective control P2X7-mScarlet) and K117A (61.5
± 2.1%, n = 155 cells, P < 0.0001). Importantly, a strong mScarlet signal was detected for all
mutants, indicating successful expression of mutants in cells (Figure 4B and Figure 4 — figure
supplement 1). Alexa 647 fluorescence reduction at K82A was significantly different from that at

K117A ($P < 0.0001$, t-test), suggesting a larger contribution of K82 to **X7-uP** labeling. The double
mutant K82A/K117A further reduced the signal by 87.3 ± 0.4% (n = 199 cells, $P < 0.0001$, t-test to
respective control P2X7-mScarlet, K82A or K117A), without affecting P2X7 expression (Figure 4B
and C). To exclude the possibility that introduced mutations affected **X7-uP** binding, we further
demonstrated that the double K82A/K117A mutant did not affect the ability of **X7-uP** to inhibit
BzATP-evoked currents (Figure 4D and E). These data support the hypothesis that both K82 and
K117 are the sites of **X7-uP** labeling in rP2X7.
257
X7-uP reveals uniform membrane distribution of P2X7 in quiescent BV2 cells
Our data demonstrate the high selectivity of **X7-uP** for P2X7 in HEK293T cells. We next
investigated whether its application extends to native P2X7-expressing microglia to reveal
endogenous P2X7 expression. We chose BV2 cells, a cultured murine microglia model where the
functional expression of P2X7 has been previously documented (26, 27). In addition, the activation
of microglia to mimic inflammation can be effectively stimulated *in vitro* using both LPS and ATP
(17). Although K117 is not conserved in mouse P2X7, the highly conserved K82 should ensure the
successful labeling of mouse P2X7 with **X7-uP** (Figure 1 — figure supplement 1A).
266
We first confirmed by ELISA assay that, under untreated conditions, BV2 cells are quiescent as
they do not release IL-1β (Figure 5A, B). We also confirmed P2X7 expression by Western blot and
confocal microscopy following 1 μM **X7-uP** labeling for 10 min in FBS-free DMEM (Figure 5 —
figure supplement 1). As observed for HEK293T cells, a strong and unique band was detected
using a P2X7 antibody only in the presence of **X7-uP** in the pulldown Western blot, at the excepted
apparent molecular mass of monomeric mouse P2X7 (~68 kDa, unglycosylated form; see lanes 3
and 4 in Figure 5 — figure supplement 1A). This band disappeared upon co-incubation with
AZ10606120 or A740003, while controls in the supernatant consistently showed endogenous P2X7
expression in all lanes, though non-specific bands were also observed, presumably due to the
limited specificity of commercially available P2X7 antibodies (Figure 5 — figure supplement 1B).
Quantitative Western blotting analysis revealed that 80 ± 1% (mean of triplicate ± s.e.m.) of total
P2X7 were labeled in BV2 cells (Figure 2 — figure supplement 2A and B). Confocal microscopy
revealed a strong Alexa 647 fluorescence signal in the periphery of BV2 cells that was reduced
upon co-incubation with AZ10606120 or A740003 (Figure 5 — figure supplement 1C and D).
Control with Strept-A 647 alone showed an extremely low fluorescence background level (Figure 5
— figure supplement 1C and D), demonstrating high **X7-uP** labeling specificity in BV2 cells.
283
To assess how BV2 activation affects P2X7 molecular organization, we employed dSTORM to
track single fluorophores tagged to endogenous receptors. In this super-resolution technique, each
fluorophore blinks several times before becoming silent due to bleaching or reaching a stable dark
state. As a result, clusters of detections could result either from multiple blinks of a single receptor
or represent the sum of multiple fluorophore emissions from clustered receptors. To quantify
nanoscale object properties, we used Metamorph to obtain pixelized global images for global
quantification (Figure 5 — figure supplement 2D), and SR-Tesseler, an open-source segmentation
framework based on Voronoï tessellation from localized molecule coordinates (49), to analyze
cluster properties (Figure 5H and I).
293
In control untreated condition (quiescent cells), dSTORM revealed punctated images, while no
detections occurred without **X7-uP**, further confirming high labeling specificity in BV2 cells (Figure
5D, and Figure 5 — figure supplement 2A and B). Detections are organized into clusters of 44 nm
in size (median, 25% percentile 18 nm, 75% percentile 80 nm), with on average 70 ± 15 (mean ±
s.e.m.) detections per cluster (n = 4 cells, Figure 5G). Distribution analysis suggests that detections
of fluorescent streptavidin conjugates are originating from one single receptor. On average, every
P2X7 receptor is separated from its neighbor by a mean inter-cluster distance of 292 ± 5 nm (Figure

5 — figure supplement 2G). These data suggest that P2X7 is homogenously dispersed on the cell
surface of quiescent BV2 cells.
303
Since P2X7 has been suggested to be associated with lipid rafts (20-25), we pre-incubated cells
with 15 mM methyl-β-cyclodextrin (MβCD), a lipid-raft disrupting agent, for 15 min, and labeled
P2X7 with **X7-uP** after extensive washout to remove remaining MβCD (Figure 5A). MβCD treatment
had no effect on IL-1β release, the number of detections per cluster, cluster size, or inter-cluster
distance (Figure 5B, F and G, Figure 5 — figure supplement 2C and G). We interpret these results
to suggest that disrupting lipid rafts does not alter P2X7 distribution in quiescent BV2 cells.
310
X7-uP reveals P2X7 clustering and upregulation in activated BV2 cells
We analyzed changes in cluster properties following BV2 activation by LPS and P2X agonists. To
reliably correlate P2X7 spatial distribution with the inflammatory response, we labeled the same
BV2 cells used for IL-1β release assays with **X7-uP** (Figure 5A). Previous studies have proposed
a two-signal, synergistic model for IL-1β release in microglia and other cell types (5, 6, 10, 50-52),
where the first signal involves LPS priming, leading to the intracellular accumulation of pro-IL-1β,
while the second involves ATP-dependent P2X7 stimulation, promoting caspase-1 activation and
the release of mature IL-1β. We confirmed this model by observing robust IL-1β release when BV2
cells were exposed to 1 mg/mL LPS for 24 hours, followed by 1 mM ATP (or 300 mM BzATP) for
30 minutes (LPS+ATP or LPS+BzATP) (Figure 5B). No IL-1β release was detected when ATP or
BzATP was applied alone, while LPS alone induced significant release. However, this release was
significantly less than that induced by co-treatments, representing only 35% of the release
observed with LPS+ATP and 21% with LPS+BzATP (Figure 5B), confirming the synergistic effect.
Co-incubation with AZ10606120 or A740003 during agonist exposure reduced IL-1β release to
levels observed with LPS alone, demonstrating the role of P2X7 in IL-1β release (Figure 5A and
C).
327
dSTORM images following treatments are shown in Figure 5D. Global single-molecule labeling on
the cell surface was significantly more pronounced in cells treated with LPS+ATP or LPS+BzATP,
and to a lesser extent, though not significantly, in those treated with LPS alone compared to
untreated control (Figure 5E). Similar results were obtained using confocal microscopy, although
the effect of LPS alone was significant (Figure 5 — figure supplement 3). Neither ATP nor BzATP
application alone affected the number of detections per cluster, cluster size, or inter-cluster
distance. At the cluster level, both LPS and LPS+ATP or LPS+BzATP treatments increased the
number of detections per cluster (Figure 5E and G and Figure 5 — figure supplement 2D and E),
with no change in inter-cluster distance or average intensity per sub-pixel (Figure 5 — figure
supplement 2F and G), indicating no increase in detection density per pixel. On pixelized images,
larger clusters (area greater than 0.025 mm^2, insert Figure 5F) emerged following LPS treatment,
with or without agonists. Tessellation analysis revealed that the number of fluorophores within
clusters significantly increased from ~1.5 in untreated cells to 4.1 ± 0.4 for LPS, 4.9 ± 0.5 for
LPS+ATP, and 5.2 ± 0.5 for LPS+BzATP (Figure 5H and I). These data demonstrate that pro-
inflammatory conditions induce P2X7 clustering by upregulating the average number of P2X7
receptors within clusters, increasing it from one to three, and favoring the formation of larger
clusters (Figure 6). The strong P2X7 clustering induced by the combination of LPS and P2X7
agonists mirrors IL-1β secretion.
346

Discussion

We present a rapid and efficient method for the selective labeling of P2X7 in native cells. This approach allowed us to uncover a nanoscale redistribution mechanism of individual endogenous P2X7 receptors on the plasma membrane of activated BV2 microglial cells, leading to the synergistic release of the pro-inflammatory cytokine IL-1β.

Our strategy employs an affinity-guided approach, through which we developed **X7-uP**, a selective P2X7 biotinylating reagent that covalently attaches a biotin tag to lysine residues on native P2X7. In rat P2X7 (rP2X7), we identified two lysine residues as labeling sites: K82, a highly conserved residue unique to P2X7, and K117, a residue present only in rP2X7. We found that K82 is the primary labeling site in rP2X7, and given that K117 is absent in mouse P2X7, it is likely that K82 is the only labeled site in BV2-expressing P2X7.

We demonstrate the effectiveness of **X7-uP** in both heterologous and native expression systems. Compared to existing methods, **X7-uP** offers several advantages. First, it combines *N*-cyanomethyl NASA chemistry with the high-affinity AZ10606120 ligand, enabling rapid labeling in microglia (within 10 min) at a low **X7-uP** concentration (1 μM) in physiological-like buffer (e.g., FBS-free DMEM). Second, because AZ10606120 binds to a unique P2X7 allosteric site (48), **X7-uP** is highly selective for P2X7, with no labeling observed in cells expressing other P2X or when specific P2X7 antagonists are present. This selectivity makes **X7-uP** particularly suitable for detecting P2X7 in native tissues, addressing the specificity issues seen with available antibodies (39). Third, **X7-uP** provides a versatile platform to deliver various probes to P2X7, including those for pulldown assays and super-resolution imaging. The strong biotin-streptavidin (or neutravidin) interaction ensures specificity and stability, two crucial criteria for biochemical assays and cell imaging.

Our results not only confirm P2X7 expression in microglia, as previously reported (6, 26-33), but also reveal its nanoscale localization at the cell surface using dSTORM. In quiescent cells, P2X7 is evenly distributed across the plasma membrane, likely as individual receptors with an average nearest-neighbor distance of ~300 nm, indicating the absence of clustering. To test whether P2X7 associates with cholesterol-enriched lipid rafts, we pre-treated cells with MβCD, a cholesterol-depleting agent that perturbs lipid rafts. This treatment did not affect P2X7 distribution, although we have previously shown in HEK293T cells that MβCD increases P2X7 single-channel mean open time without altering unitary conductance (53). These findings suggest that in quiescent BV2 microglial cells, perturbating lipid rafts does not alter the uniform distribution of individual P2X7 receptors, which likely reside within cholesterol-enriched nanodomains.

Under inflammatory conditions, we observed significant nanoscale reorganization of P2X7. Both LPS and ATP (orBzATP) trigger P2X7 upregulation and clustering, increasing the overall number of surface receptors and the number of receptors per cluster, from one to three (Figure 6). By labeling BV2 cells with **X7-uP** shortly after IL-1β release, we were able to correlate the nanoscale distribution of P2X7 with the functional state of BV2 cells, consistent with the two-signal, synergistic model for IL-1β secretion observed in microglia and other cell types (5, 6, 10, 50-52). In this model, LPS priming leads to intracellular accumulation of pro-IL-1β, while ATP stimulation activates P2X7, triggering NLRP3 inflammasome activation and the subsequent release of mature IL-1β. Although the mechanism behind the overall increase in surface receptors remains unclear, a recent study suggests that the a1 subunit of the Na$^+$/K$^+$-ATPase (NKAα1) forms a complex with P2X7 in microglia, including BV2 cells, and that LPS+ATP induces NKAα1 internalization (17). This internalization appears to release P2X7 from NKAα1, allowing P2X7 to exist in its free form. We speculate that the internalization of NKAα1 induced by both LPS and ATP exposes previously masked P2X7 sites, including the allosteric AZ10606120 sites, thus making them accessible for **X7-uP** labeling.

In sequence, the first priming step induced by LPS triggers P2X7 clustering and tends to increase
the number of cell-surface receptors. This modest increase in P2X7 density may explain the lack
of functional upregulation in the amplitude of the P2X7 current component observed in BV2 cells
following LPS treatment (26, 27). The second step, triggered by ATP exposure, further elevates
surface receptor levels and activates the P2X7-mediated NLRP3 inflammasome, which cleaves
pro-IL-1β into its mature form and facilitates its release. Previous evidence suggests that P2X7 and
NLRP3 co-localize in mouse microglia (54), indicating that NLRP3 may reside beneath P2X7
clusters. We propose that the spatial reorganization and clustering of P2X7 during the LPS-induced
priming phase at sites of active inflammation is an efficient mechanism to ensure synergistic IL-1β
secretion in response to ATP. This clustering mechanism represents a finely tuned regulatory
process that effectively links PAMP-induced signaling to DAMP-evoked responses.
410
411 In conclusion, **X7-uP** is a powerful tool for visualizing P2X7 localization at the nanoscale on the cell
412 surface of BV2 cells. We anticipate that **X7-uP** will become a valuable molecular probe for
413 investigating P2X7 organization in other native cell types, such as macrophages, and for resolving
414 the ongoing debate about P2X7 expression in neurons (55, 56).
415
416
417

Materials and Methods

Chemical synthesis

All chemicals were purchased from Sigma-Aldrich, Acros Organics or Alfa Aesar in analytical grade.
An agilent MM-ESI-ACI-SQ MSD 1200 SL spectrometer or an Agilant LC-MS Agilent RRLC
1200SL/ESI QTof 6520 was used for ESI analysis. ^{1}H NMR and ^{13}C NMR spectra were recorded
at 400 or 700, and 100 or 175 MHz, respectively, using the following NMR Bruker instruments: a
400 MHz spectrometer equipped with an Avance III console and a BBO H/F z-gradient probe or a
700 MHz spectrometer equipped with a TCI z-gradient cryoprobe and an Avance III-HD console.
Coupling constants (J) are quoted in hertz (Hz) and chemical shifts (δ) are given in parts per million
(ppm) using the residue solvent peaks (CDCl$_3$: 7.26 ppm, MeOD: 3.31 ppm, DMSO-d$_6$: 2.50 ppm
for ^{1}H NMR and CDCl$_3$: 77.16 ppm, MeOD: 49 ppm, DMSO-d$_6$: 39.52 ppm for ^{13}C NMR). The
attributions are given in the following manner: chemical shift followed by the multiplicity in
parenthesis (s, d, t, q, m, dd, dt, br corresponding respectively to singlet, doublet, triplet, quadruplet,
multiplet, doublet of a doublet, doublet of a triplet, broad; number of protons and coupling constant
in Hz).

HPLC analyses were performed on a Waters® high-performance chromatography system (1525
pump, 2996 detector) equipped with a Thermo Betabasics 5-micron analytical column (4.6, 250
nm). A gradient solution was applied, progressing from 100% mQ H$_2$O acidified with 0.01% TFA to
100% acetonitrile over 30 min, followed by 10 min at 100% acetonitrile. HPLC purifications were
carried out on a Waters® high-performance chromatography system (600 double body pump, 2996
detector) equipped with a Thermo Betabasics 5-micron semi-preparative column (10, 250 nm),
using the same gradient as described above.

5-aminoquinolin-2(1H)-one (3):

3

443
To a solution of 5-nitroquinolin-2(1H)-one (**2**) (1 g, 5.26 mmol, 1 eq.) dissolved in N,N-
dimethylformamide (DMF, 40 mL) froze at -196°C was added dry palladium on activated coal (10%)
(448 mg, 2.21 mmol, 0.8 eq.). Solution was let to warm up at room temperature and gaseous
dihydrogen was bubbled in the solution for 5 minutes. Reaction was then let under dihydrogen
atmosphere and stirred for 16 h. Crude product was then filtrated on a celite pad. The pad was
further rinsed with DMF (100 mL) and the filtration product was evaporated to yield a grey solid
(825 mg, 98 %).

^{1}H NMR (400 MHz, MeOD): δ 8.19 (dd, J = 9.7, 1.0 Hz, 1H), 7.26 (m, 1H), 6.63 (dt, J = 8.1, 1.0 Hz,
1H), 6.55 (dd, J = 8.1, 1.0 Hz, 1H), 6.47 (d, J = 9.7 Hz, 1H). See Figure 1 — figure supplement 3
for spectrum.

2-(adamantan-1-yl)acetyl chloride (4):

4

456
457 2-(1-adamantyl)acetic acid (1.1 g, 5.67 mmol, 1 eq.) was placed in an oven-dried flask equipped
458 with a reflux. Thionyl chloride (10 mL) was added and the solution was heat to reflux under argon
459 atmosphere for 3 h. Solvent was then evaporated and the resulting yellow oil was further dried
460 under vacuum. Obtained compound was immediately used in the next step.

462 **2-(adamantan-1-yl)-*N*-(2-oxo-1,2-dihydroquinolin-5-yl)acetamide (5):**

5

463
464 To a stirred solution of 5-aminoquinolin-2(1H)-one (**3**) (825 mg, 5.15 mmol, 1 eq.) in anhydrous
465 tetrahydrofuran (THF) was added freshly distilled triethylamine (1,11 mL, 8,24 mmol, 1,6 eq.). The
466 mixture was cooled to 0°C and a solution of 2-(-adamantan-1-yl)acetyl chloride (1.2 g, 5.67 mmol,
467 1.1 eq) (**4**) dissolved in dry THF (5 mL) was added dropwise over the course of 15 minutes to form
468 a grey solution. The solution was kept at 0°C for 1 h and brought to 20 °C for 2 more hours. The
469 solution was then evaporated and the obtained solid was rinsed 3 times with CH_2Cl_2 (50 mL). The
470 solid was further dried to yield the expected compound as a grey solid (1.4 g, 81 %). The product
471 was then used without further purification.
472 ^{1}H NMR (400 MHz, MeOD): δ 8.06 (dd, *J* = 9.8, 0.8 Hz, 1H), 7.58 – 7.54 (m, 1H), 7.33 (dd, *J* = 7.8,
473 1.1 Hz, 1H), 7.26 (m, 1H), 6.65 (d, *J* = 9.8 Hz, 1H), 2.26 (s, 2H), 2.03 (m, 3H), 1.86 – 1.70 (m, 12H).
474 See Figure 1 — figure supplement 4 for spectrum.
475
476 **2-(adamantan-1-yl)-*N*-(2-chloroquinolin-5-yl)acetamide (6)**
477

6

479 To a stirred solution of 2-(adamantan-1-yl)-*N*-(2-oxo-1,2-dihydroquinolin-5-yl)acetamide (**5**) (725
480 mg, 2.15 mmol, 1 eq.) dissolved in 1,2-dichloroethane (10 mL) was added freshly distilled POCl₃
481 (618 μL, 6.45 mmol, 3 eq.). The solution was brought to reflux for 16 h and the solvent was
482 evaporated. Crude product was dissolved in CH_2Cl_2, filtrated and evaporated to yield the expected
483 compound as an orange solid (760 mg, 99 %).
484 ^{1}H NMR (400 MHz, MeOD): δ 8.43 (dd, *J* = 8.8, 0.8 Hz, 1H), 7.87 – 7.79 (m, 2H), 7.72 (dd, *J* = 7.2,
485 1.5 Hz, 1H), 7.59 – 7.55 (m, 1H), 2.30 (s, 2H), 2.03 (m, 3H), 1.86 – 1.72 (m, 12H). See Figure 1 —
486 figure supplement 5 for spectrum.
487
488 **2-(adamantan-1-yl)-*N*-(2-((2-aminoethyl)amino)quinolin-5-yl)acetamide (7):**
489

7

491 To a stirred solution of 2-(adamantan-1-yl)-N-(2-chloroquinolin-5-yl)acetamide (**6**) (488 mg, 1.36
492 mmol, 1 eq.) in freshly distilled ethanol (10 mL) was added dry K₂CO₃ (376 mg, 2.72 mmol, 2 eq.)

and anhydrous ethylene diamine (4.5 mL, 68 mmol, 50 eq.). The mixture was stirred for 48 h at reflux. The solution was evaporated and subsequently extracted with 3 × 50 mL CH_2Cl_2. Organic layers were combined, washed with 25 mL of brine and dried over $MgSO_4$, filtered and evaporated. The crude extract was retaken in a minimum of CH_2Cl_2 and diethyl ether was added dropwise until apparition of opalescence. Heptane was then slowly added to precipitate the pure product (262 mg, 51 %) as a yellow solid.

[1]H NMR (400 MHz, MeOD): δ 7.96 (d, J = 9.2 Hz, 1H), 7.56 − 7.46 (m, 2H), 7.23 (dd, J = 7.2, 1.4 Hz, 1H), 6.80 (d, J = 9.2 Hz, 1H), 3.57 (t, J = 6.2 Hz, 2H), 2.91 (t, J = 6.2 Hz, 2H), 2.25 (s, 2H), 2.03 (m, 3H), 1.87 − 1.68 (m, 12H). See Figure 1 — figure supplement 6 for spectrum.

N-(2-((5-(2-(adamantan-1-yl)acetamido)quinolin-2-yl)amino)ethyl)-4-sulfamoylbenzamide (8):

8

To a stirred solution of 2-(adamantan-1-yl)-N-(2-((2-aminoethyl)amino)quinolin-5-yl)acetamide (**7**) (105 mg, 0.28 mmol, 1 eq.) in dry DMF (2.5 mL) is added 4-sulfamoylbenzoic acid (67 mg, 0.33 mmol, 1.2 eq.), 1-Hydroxybenzotriazole (45 mg, 0.33 mmol, 1.2 eq.) and DIEA (129 μL, 0.75 mmol, 2.7 eq.). The solution was cooled down to 0°C and EDC (64 mg, 0.33 mmol, 1.2 eq.) was added. The mixture was then slowly warmed up to 20°C, and stirred at 20°C for 18 h. Solvent was evaporated and the crude product was purified by invert phase flash chromatography (gradient acetonitrile: H_2O (0.1 % TFA), 0:1 to 1:0 over 30 min, retention time: 16.2 min) to yield pure compounds as an orange solid (100 mg, 64 %).

[1]H NMR (400 MHz, MeOD): δ 8.19 (d, J = 9.5 Hz, 1H), 8.02 − 7.91 (m, 4H), 7.70 (m, 2H), 7.45 (dd, J = 7.5, 1.3 Hz, 1H), 7.04 (d, J = 9.5 Hz, 1H), 3.83 (t, J = 6.2 Hz, 2H), 3.73 (t, J = 6.2 Hz, 2H), 2.28 (s, 2H), 2.02 (s, 3H), 1.88 − 1.68 (m, 12H).

[13]C NMR (100 MHz, MeOD): δ 173.92, 169.94, 165.28, 160.81, 156.37, 148.36, 138.62, 136.24, 133.03, 129.52, 127.77, 122.82, 119.63, 119.19, 114.59, 65.67, 55.16, 52.36, 44.52, 38.31, 34.83, 30.50. See Figure 1 — figure supplement 7 for spectra.

MS (ESI): m/z [M[+]] calculated for $C_{30}H_{36}N_5O_4S^+$: 562.2483, found 562.2516.

tert-butyl **(15-((4-((2-((5-(2-(adamantan-1-yl)acetamido)quinolin-2-yl)amino)ethyl)carbamoyl)phenyl)sulfonamido)-15-oxo-3,6,9,12-tetraoxapentadecyl)carbamate (10):**

10

To a stirred solution of N-(2-((5-(2-(adamantan-1-yl)acetamido)quinolin-2-yl)amino)ethyl)-4-sulfamoylbenzamide (**8**) (43 mg, 76 μmol, 1 eq.) in dry DMF (1 mL) was added COOH-OEG4-Boc (**9**) (33.6 mg, 92 μmol, 1.2 eq.), DIEA (33 μL, 191 μmol, 2.5 eq.) and DMAP (2 mg, 15 μmol, 0.2

eq.). Solution was cooled down to 0°C and EDC was added (17.6 mg, 92 µmol, 1.2 eq.). Solution
was let to slowly warm up to 20°C for 16 h. Every 24 h during 72 h was added 1 eq. of carboxylic
acid, 1 eq. of EDC and 2 eq. of DIEA at 0°C. Solvent was then evaporated and 50 mL of a 1:1
CH_2Cl_2:H_2O solution was added. Organic layer was extracted with 2 × 20 mL of CH_2Cl_2. Organics
layers were then combined, dry over $MgSO_4$, filtered and evaporated. Compound was purified by
HPLC purification (acetonitrile:H_2O (0.1% TFA), 0:1 to 1:0 over 30 min, retention time: 17.7 min).
Pure fractions were combined, extracted with 3 × 20 mL, dry over $MgSO_4$ and evaporated to yield
pure compound as a dark orange oil (57 mg, 82 %).
^{1}H NMR (400 MHz, $CDCl_3$): δ 8.20 (m, 1H), 8.01 (m, 4H), 7.42 (m, 3H), 6.85 (s, 1H), 3.79 (m, 2H),
3.60 (m, 18H), 3.26 (s, 2H), 2.48 (s, 2H), 2.22 (s, 2H), 1.98 (s, 3H), 1.77 − 1.59 (m, 12H), 1.41 (s,
9H).
^{13}C NMR (100 MHz, CDCl3): δ 171.08, 170.38, 168.12, 166.33, 156.37, 153.55, 139.80, 137.58,
134.55, 133.23, 128.28, 127.87, 121.02, 115.21, 113.45, 108.43, 79.43, 70.45, 70.38, 70.22, 70.13,
66.13, 53.42, 51.57, 42.68, 36.69, 33.47, 28.51. See Figure 1 — figure supplement 8 for spectra.
543
544   **_N_-(2-((5-(2-(adamantan-1-yl)acetamido)quinolin-2-yl)amino)ethyl)-4-(_N_-(17-oxo-21-**
545   **((3a_R_,4_R_,6a_S_)-2-oxohexahydro-1_H_-thieno[3,4-_d_]imidazol-4-yl)-4,7,10,13-tetraoxa-16-**
546   **azahenicosanoyl)sulfamoyl)benzamide (11):**

11

547
To a stirred solution tert-butyl (15-((4-((2-((5-(2-(adamantan-1-yl)acetamido)quinolin-2-
yl)amino)ethyl)carbamoyl)phenyl)sulfonamido)-15-oxo-3,6,9,12-tetraoxapentadecyl)carbamate
(**10**) (7.6 mg, 8 µmol, 1 eq.) in 700 µl of CH_2Cl_2 was added 100 µl of TFA and the solution was left
under argon for 1 h at 20°C. After completed deprotection of the amine (confirmed by HPLC),
solvent was evaporated and the residue was further dried by 3 coevaporation with anhydrous
toluene (5 mL). Anhydrous DMF was then added in the flask (1 mL), followed by N-
methylmorpholine (2 µL, 17 µmol, 2 eq.), biotin (3 mg, 13 µmol, 1.5 eq.) and 4-(4,6-dimethoxy-
1,3,5-triazin-2-yl)-4-methyl-morpholinium chloride (DMTMM, 3.4 mg, 13 µmol, 1.5 eq.). Solution
was left 2 h at 20°C. Solvent was then evaporated and the pure product was obtained by HPLC
purification (acetonitrile: H_2O (0.1% TFA), 0:1 to 1:0 over 30 min, retention time: 18.1 min) to yield
an orange solid (7 mg, 80 %).
^{1}H NMR (500 MHz, MeOD): δ 10.01 (s, 1H), 9.01 (br, 1H), 8.29 (s, 1H), 8.07 − 7.96 (m, 4H), 7.82
(t, J = 5.6 Hz, 1H), 7.76 − 7.69 (br, 1H), 7.60 (br, 1H), 7.09, (s, 1H), 6.41 (s, 1H), 6.36 (s, 1H), 4.31,
(dd, J = 7.6, 4.2 Hz, 1H), 4.13 (dd, J = 7.6, 4.4 Hz, 1H), 3.76 (br, 2H), 3.64 (br, 2H), 3.55 − 3.35 (m,
16H), 3.18 (q, J = 6.2 Hz, 2H), 3.12 − 3.07 (m, 1H), 2.82 (dd, J = 12.6, 5.3 Hz, 1H), 2.58 (J = 12.6,
2H), 2.21 (m, 2H), 2.06 (t, J = 7.3 Hz, 2H), 1.97 (s, 3H) 1.72 − 1.57 (m, 12H), 1.53 − 1.41 (m, 3H),
1.35 − 1.23 (m, 3H).
^{13}C NMR (100 MHz, MeOD): δ 174.74, 172.09, 170.58, 168.23, 164.72, 159.59, 153.15, 137.88,
136.88, 135.05, 132.40, 128.07, 127.70, 122.07, 116.79, 115.21, 113.51, 70.14, 70.12, 70.00,
69.17, 65.80, 61.98, 60.26, 55.58, 50.55, 42.46, 39.67, 38.97, 36.48, 35.33, 33.05, 28.79, 28.33,
28.09, 25.44. See Figure 1 — figure supplement 9 for spectra.
569

570 *N*-(2-((5-(2-(adamantan-1-yl)acetamido)quinolin-2-   yl)amino)ethyl)-4-(*N*-(cyanomethyl)-*N*-
571 **(17-oxo-21-((3a*R*,4*R*,6a*S*)-2-oxo-   hexahydro-1*H*-thieno[3,4-*d*]imidazol-4-yl)-4,7,10,13-**
572 **tetraoxa-16-azahenico- sanoyl)sulfamoyl)benzamide (X7-uP):**

573 **X7-uP**

574
575 To a stirred solution of N-(2-((5-(2-(adamantan-1-yl)acetamido)quinolin-2-yl)amino)ethyl)-4-(N-(17-
576 oxo-21-((3aR,4R,6aS)-2-oxohexahydro-1H-thieno[3,4-d]imidazol-4-yl)-4,7,10,13-tetraoxa-16-
577 azahenicosanoyl)sulfamoyl)benzamide (**11**) (4 mg, 4 µmol, 1 eq.), previously dried with 5
578 coevaporation with anhydrous toluene (5 mL), in dried DMF (0.5 mL) was added freshly distilled
579 DIEA (32 µl, 190 µmol, 50 eq.) and iodoacetonitrile (13 µl, 190 µmol, 50 eq.). The latter was dried
580 using alumina plug prior use. The solution was stirred for 16 h at 20°C in the dark and solvent was
581 then evaporated. Pure compound was obtained by HPLC purification (acetonitrile: H_2O (0.1% TFA),
582 0:1 to 1:0 over 30 min, retention time: 19.1 min) to yield a brown solid (3.3 mg, 80 %).
583 ^{1}H NMR (700 MHz, DMSO): δ 10.02 (s, 1H), 9.08 (br, 1H), 8.29 (s, 1H), 8.14 (d, *J* = 8.3 Hz, 2H),
584 8.08 (d, *J* = 8.3 Hz, 2H), 7.83 (t, *J* = 5.6 Hz, 1H), 7.76 – 7.72 (br, 1H), 7.60 (br, 1H), 7.09 (s, 1H),
585 6.42 (s, 1H), 6.37 (s, 1H), 5.02 (s, 2H), 4.31 (dd, *J* = 7.8, 5.0 Hz, 1H), 4.13 (dd, *J* = 7.7, 4.4 Hz, 1H),
586 3.77 (s, 2H), 3.66 (s, 2H), 3.60 (t, *J* = 6.1 Hz, 2H), 3.53 – 3.42 (m, 12H), 3.39 (t, *J* = 6.0 Hz, 2H),
587 3.18 (q, *J* = 6.0 Hz, 2H), 3.12 – 3.07 (m, 1H), 2.94 (t, *J* = 6.1 Hz, 2H), 2.82 (dd, *J* = 12.6, 5.1 Hz,
588 1H), 2.58 (d, *J* = 12.6 Hz, 2H), 2.22 (m, 2H), 2.07 (t, *J* = 7.4 Hz, 2H), 1.97 (s, 3H), 1.73 – 1.58 (m,
589 12H), 1.53 – 1.43 (m, 4H), 1.34 – 1.23 (m, 2H).
590 ^{13}C NMR (175 MHz, DMSO): δ 170.91, 169.04, 168.67, 164.13, 161.47, 161.06, 156.95, 156.77,
591 138.73, 137.84, 133.90, 129.02, 128.80, 127.17, 126.67, 116.81, 116.40, 114.96, 114.70, 68.52,
592 68.47, 68.44, 68.42, 68.31, 68.27, 63.96, 59.78, 57.94, 54.15, 53.63, 48.88, 40.83, 38.72, 37.15,
593 35.13, 34.63, 34.52, 33.82, 32.69, 31.52, 29.50, 29.42, 26.92, 26.78, 26.76, 23.99. See Figure 1
594 — figure supplement 10 for spectra.
595 MS (ESI): m/z [M+] calculated for $C_{53}H_{71}N_9O_{11}S_2^+$: 1074.4787, found 1074.4797.

596
597 **Molecular modelling**
598 The P2X7 structure resolved with the allosteric inhibitor AZ10606120 was retrieved form the Protein
599 Data Bank (PDB, code 5U1W) (48). The inhibitor was removed, and docking simulations were
600 performed using AutoDock Vina 1.5.6. A 30 Å box centered on the quinoline core of AZ10606120
601 was used for docking, set before ligand removal. The cartoon shown in Figure 6 was created using
602 tetrameric biotin-bound streptavidin (PDB code 1MK5) and rP2X7 (PDB code 6U9W). Tetrameric
603 biotin-bound streptavidin was positioned relative to K82, using the maximum 18.5 Å distance
604 between the α-carbon of K82 and the carboxylate carbon of the bound biotin.

605
606 **Cell culture and transfection**
607 HEK293T cells (ATCC) and mouse BV2 microglia cells (kind gift of Dr. R. Schlichter, Institute of
608 Cellular and Integrative Neuroscience, University of Strasbourg, France) were cultured in high-
609 glucose Dulbecco's modified Eagle's medium (DMEM ref. 31966-021) supplemented with 10%
610 (v/v) heat-inactivated fetal bovine serum (FBS), 100 units/mL penicillin and 100 µg/mL streptomycin
611 (Gibco Life Technologies, USA). Cells were grown at 37°C in a humidified cell incubator with 5%
612 CO_2.

613

For transfection, cells were grown to 70-80% confluence, and the calcium phosphate precipitation
method was employed. The cDNA encoding rat P2X7-mScarlet (57), P2X1-GFP, P2X2-GFP,
P2X3-GFP, P2X4-GFP, P2X5-GFP, or P2X6-GFP (all kind gifts from Dr. F. Rassendren, Institut de
Génomique Fonctionelle, University of Montpellier, France) were contained within pcDNA3.1(+)
plasmids (Invitrogen, USA). rP2X7-myc was obtained as previously described (53). 24 h after
transfection, the medium was replaced with fresh medium. For whole-cell patch-clamp
experiments, cells were co-transfected with the rP2X7 construct (0.8 μg) and an eGFP (0.3 μg)
which allowed to identify cells that had undergone efficient transfection. For biochemical
experiments, each 100 mm dish was transfected with P2X7-myc construct (10 μg). For confocal
imaging and dSTORM experiments, HEK293T cells were transfected with 1 μg of the indicated
construct.
625
Mutagenesis
627   Site-directed mutations were introduced into the P2X7-mScarlet cDNA in the pcDNA3.1(+) using
628   KAPA HiFi HotStart PCR kit (Cliniscience, France) as previously described (58). All mutations were
629   confirmed by DNA sequencing.
630
Electrophysiology patch-clamp
Electrophysiological recordings in whole-cell configuration were carried out as previously described
(53). Briefly, patch pipettes were pulled from borosilicate glass capillaries and microforged to yield
a resistance of 3-5 MΩ. Cells were voltage-clamped to -60 mV using EPC10 USB amplifier (HEKA,
Reutlingen, Germany), and data were recorded with PATCHMASTER software (version V2X90.5).
Ligands were applied via a perfusion system, using three capillary tubes placed directly over the
cell of interest.
638
2′(3′)-O-(4-benzoylbenzoyl) adenosine 5′-triphosphate (BzATP) (triethylammonium salt, Alomone
labs, Israel) was used as P2X7 agonist at a concentration of 10 μM. AZ10606120 and **X7-uP** were
used at the indicated concentrations. BzATP was dissolved in Normal Extracellular Solution (NES),
containing 140 mM NaCl, 2.8 mM KCl, 1 mM CaCl2, 0.1 mM MgCl2, 10 mM HEPES, and 10 mM
glucose, with a pH of 7.32–7.33. AZ10606120 and **X7-uP** were prepared as concentrated stocks
in DMSO and diluted in NES to the desired working concentration (< 0.1% DMSO). Only one cell
was patched per coverslip.
646
Labeling reaction with X7-uP
X7-uP was stored as a DMSO stock solution (24.9 mM) and diluted in FBS-free DMEM to achieve
the desired working concentration for labeling (< 0.1% DMSO). After three washing steps with
freshly prepared sterile-filtered PBS+ buffer (containing 137 mM NaCl, 2.68 mM KCl, 10 mM
Na$_2$HPO$_4$, 1.76 mM KH$_2$PO$_4$, 1 mM MgCl$_2$ and 0.4 mM CaCl$_2$), cells were incubated with **X7-uP**
diluted in FBS-free DMEM for the indicated time at 37°C. Following incubation, cells were washed
three times with PBS+ buffer and treated as indicated. For confocal and dSTORM experiments,
biotinylated cells were incubated with 1 μg/mL of Strept-A 647 (S21374, Invitrogen, USA) for 10
min in PBS+ buffer containing 1% bovine serum albumin (BSA) at 37°C. After three additional
washes with PBS+ buffer, cells were fixed as described below.
657
Pull Down, SDS-PAGE and Western blot
Cells were solubilized by vortexing in a lysis buffer (20 mM HEPES, 100 mM NaCl, 1 % Triton-X, 5
mM EDTA, 1 % protease inhibitor (Thermo Fischer, Waltham, MA, USA)) at 4°C for 90 min.
Following centrifugation (14,000 rpm, 10 min), supernatants were collected, mixed with the
NuPAGE LDS loading buffer (Thermo Fisher) containing 70 mM dithiothreitol (DTT), boiled for 10
min, and run on a NuPAGE Novex Bis-Tris 4%-12% (Thermo Fischer, USA) in MOPS buffer.
Proteins were transferred to a nitrocellulose membrane using semi-wet transfer (TransBlot Turbo
system, BioRad, CA, USA), and the membrane was blocked for 30 min in TPBS buffer (PBS buffer
supplemented with 1% milk powder, 0.5% BSA and 0.05 % Tween-20). The membrane was then
incubated overnight at 4 °C under gentle agitation with primary antibodies: 1:500 anti-c-Myc mouse

antibody (13-2500, Invitrogen, Thermo Fischer), 1:500 rabbit anti-P2X7 antibody (APR-008,
Alomone Labs, Israel), or mouse 1:5,000 anti-β-actin (A5441, Sigma Aldrich). Proteins were
visualized using 1:10,000 horseradish peroxidase-conjugated secondary antibodies directed
against mouse or rabbit (NA9310, GE Life Sciences and 31460, Invitrogen, respectively) and a
chemiluminescent substrate (Amersham ECL Select Western Blotting Detection, Ge Life Sciences,
MA, USA) on an Imager 600 (Amersham, IL, USA).
For pull-down experiments, solubilized cells (as described above) were incubated overnight at 4
°C under gentle agitation with Pierce NeutrAvidine agarose resin (29200, Thermo Fischer).
Samples preparation and Western blotting was carried out as described above.
677
Coverslip preparation
For confocal imaging and super-resolution experiments, coverslips (0117580, No. 1.5H,
Marenfield, Germany) were cleaned by incubation in freshly made piranha solution (H_2SO_4:H_2O_2
50% 3:1) at 60°C for 30 min. The cleaning solution was replaced with fresh piranha solution, and
the coverslips were sonicated for an additional 30 min. After discarding the piranha solution,
coverslips were washed 10 times with ultra-pure water and sonicated in ultrapure water (10×3
min each) to ensure complete removal of any cleaning residue. Coverslips were then dispatched
in 12-well plates and exposed to UV light for 30 min. For HEK293T cells, the coverslips were treated
with poly-L-lysine for 1 h at 37°C and subsequently washed with ultra-pure water before use. For
BV2 cells, no treatment with poly-L-lysine was performed.
688
BV-2 microglia stimulation
BV2 microglia cells were seeded on treated coverslips (as described above) in a 12-well plate
containing standard DMEM medium with 10% FBS. After 24h, cells were incubated in DMEM
medium with 1% FBS for 24 h, either left untreated or treated with *Escherichia coli* O111:B4 LPS
(1 ng/mL, ref. L2630, Sigma Aldrich, USA). After washing once with freshly prepared sterile-filtered
PBS+, cells were incubated in PBS+, either in the absence (vehicle) or supplemented with 1 mM
adenosine 5′-triphosphate (ATP disodium salt hydrate, ref. A7699, Sigma Aldrich) for 30 min, 300
μM BzATP for 30 min, or 15 mM MbCD (Sigma Aldrich, USA) for 15 min at 37°C in a humidified
cell incubator with 5% CO_2. After treatments, supernatants were collected for the IL-1β ELISA
assay, and cells were labeled with **X7-uP** as described above.
699
700 **ELISA assay**
701 IL-1β levels were measured using the IL-1β Mouse Uncoated ELISA Kit (ref. 88-7013, Invitrogen,
702 USA) following the manufacturer's protocol. Developed plates were read using a SAFAS Monaco
703 plate reader with SP2000 version 7 software.
704
Cell fixation
Cells were fixed by 4% paraformaldehyde (PFA) in PBS, pH 8.0 for 20 min at 20°C. Fixed cells
were then washed three times and incubated with fresh PBS buffer for 3×5 min, followed by
incubation in 100 mM glycine (in PBS buffer) for 15 min at 20°C. After three washes, cells were
stored at 4°C until use. All washing steps were performed with freshly made, sterile-filtered PBS
buffer.
711
Cytochemistry
For confocal imaging, coverslips containing fixed cells were treated with 4 μg/mL of Hoechst 33342
pentahydrate (bis-benzimide) (H21491, Molecular Probes, Life Technologies, USA) in PBS buffer
for 10 min. The staining solution was discarded, and coverslips were washed three times with PBS
buffer, rinsed with ultra-pure water, mounted on microscopy slides with the Prolong Gold Antifade
(P36930, Invitrogen, USA), and allowed to dry overnight in the dark before use.
718
719 For confocal imaging of BV2 cells shown in Figure 5 — figure supplement 3, coverslips were rinsed
720 with ultra-pure water, mounted on microscopy slides using the Prolong Diamond Antifade with DAPI
721 (P36962, Invitrogen, USA), and allowed to dry overnight in the dark before use.

722
For dSTORM experiments in HEK293T cells, coverslips were incubated at room temperature for 6
minutes in a PBS solution containing 1:5,000 fluorescent microspheres (Tetraspeck, Invitrogen,
USA) as fiducial markers for lateral drift correction during image reconstruction, rinsed with ultra-
pure water, mounted on microscopy slides on a Vectashield® H-1000 (Vector Laboratories, USA)
containing a 50 mM TRIS-glycerol (obtained by diluting 5% v/v TRIS pH 8.0 in glycerol and filtered
on 0.22 μm filter) in a 1:4 ratio, and sealed using dental cement (Picodent, Germany) (59).
729
For dSTORM experiments in BV2 cells, coverslips were first incubated at room temperature for 6
minutes in a PBS solution containing 1:5,000 fluorescent microspheres (Tetraspeck, Invitrogen,
USA). The coverslips were then mounted in a Ludin chamber (Life Imaging Services), where a
buffer composed of an oxygen scavenger (glucose oxidase) and a reducing agent (2-
mercaptoethylamine) was added (600 mL). The Ludin chamber was sealed with an 18-mm
coverslip to prevent oxygen exchange.
736
737 **Confocal imaging**
738 Confocal imaging was captured with Leica SPE (63x oil immersion objective, N.A. 1.4). Excitation
739 (λexc) and emission (λem) wavelengths were as follows: DAPI, Hoechst (λexc = 405 nm, λem =
740 430 – 480 nm), GFP (λexc = 488 nm, λem = 500 – 545 nm), mScarlet (λexc = 561 nm, λem = 570
741 – 610 nm) and Alexa Fluor-647 (λexc 635 nm, λem = 650 – 700 nm).
742
dSTORM imaging and reconstruction
dSTORM acquisitions in HEK293T cells were conducted on a homemade system built on a Nikon
Eclipse Ti microscope, equipped with a 100x oil immersion objective (N.A. 1.49) (60). A 642 nm
laser line was used to excite the Alexa Fluor-647 fluorophore, set to a power of 134 mW (resulting
in an intensity of 2.7 kW/cm^2). Samples were imaged with an EM-CCD camera (Hamamatsu,
ImagEM) maintained at -60°C. Z-stabilization was ensured by the Nikon Perfect Focus System
integrated into the microscope. Acquisition was performed in TIRF mode controlled using Micro-
Manager 1.4.23 (61). Single-molecule localization was achieved by analysis a stack of 15,000
images with the ThunderSTORM ImageJ plugin (62). The following parameters were used: image
filtering – Difference-of-Gaussians filter (sigma 1 = 1.2 px, sigma 2 = 1.9 px), approximate
localization of molecules: centroid of connected components (peak intensity threshold std
(Wave.F1), sub-pixel localization of molecules: PSF: Integrated Gaussian (fitting radius: 6 px, fitting
method: Weighted Least squares, initial sigma: 1.6 px)). Results were filtered by sigma, localization
precision, and intensities values: 110 nm < sigma > 220 nm, precision < 25 nm and intensity <
2000.
758
dSTORM acquisitions in BV2 cells were performed on a commercial LEICA DMi8 (Leica, Germany)
inverted microscope equipped with anti-vibrational table (TMC, USA) to minimize drift, along with
a Leica HC PL APO 100X oil immersion TIRF objective (NA 1.47). For sample excitation, the
microscope was equipped with a fiber-coupled laser launch composed of the following
wavelengths: 405, 488, 532, 561, and 642 nm (Roper Scientific, Evry, France). Samples were
imaged with an EMCCD camera (Teledyne Photometrics). Z stabilization was guaranteed by the
Leica auto focus system integrated into the microscope. The 642 nm laser was used at a constant
power to excite the Alexa Fluor-647 fluorophore, and the 405 nm laser was adjusted throughout
the acquisition to control the level of single molecules per frame. Image acquisition was performed
in TIRF mode controlled by MetaMorph software (Molecular Devices), with a 30 ms frame duration
and a stack of 20,000 frames per acquisition on a 512 x 512-pixel ROI (pixel size = 160 nm).
770
771 Super-resolved images were reconstructed using the PALMTracer plugin for MetaMorph (63). The
772 localizations were first extracted using a consistent intensity threshold across the entire dataset.
773 Subsequently, super-resolved images were generated from these localizations with a pixel size of
774 40 nm. The density of localization was then extracted on each image, based on ROIs determined
775 around the plasma membrane.

Cluster size measurement

Density-based clustering was performed using SR-Tesseler software (49) with the following input parameters: density factor 1 = 1; density factor 2 = 0.7, max length = 250 nm and min locs = 10; max locs = 200, max distance = 2 μm. The data obtained were further analyzed with GraphPad Prism (version 8.0.2).

Kinetic analysis of X7-uP labeling

The **X7-uP** labeling reaction follows a kinetic model in which a reversible ligand binding reaction (here **X7-uP**) precedes an irreversible chemical reaction, as described previously (41). In the presence of large excess of **X7-uP** relative to P2X7 sites, the pseudo-first-order reaction rate constant (k_{app}) is given by Eq. 1:

$$f(t) = \exp\left(-k_{app}t\right)$$

where t is the labeling reaction time at a given **X7-uP** concentration. The relationship between k_{app} and **X7-uP** concentration is described by Eq. 2:

$$k_{app} = \frac{k_L}{1 + K_d/[\text{X7-uP}]}$$

where K_d is the dissociation constant of **X7-uP** and k_L is the rate constant of the irreversible chemical reaction.

Data analysis

All experiments were conducted with at least three independent experiments. Statistical significance was assumed when the *P*-value was < 0.05. Graphs were generated using RStudio (version 1.4.1717) and ggplot2 package (version 3.3.5) or GraphPad prism.

For confocal data, membrane fluorescence intensities were manually selected using the ROI function of ImageJ (version 2.3.0/1.53f). Data were then analyzed using in-house written R scripts, employing the following packages: ggplot2, reshape2 (version 1.4.4), readxl (version 1.3.1), dplyr (version 1.0.7), Rmisc (version 1.5), and naniar (version .6.1).

For confocal data shown in Figure 5 — figure supplement 3, membrane intensities were automatically selected. A threshold was first applied to distinguish particles from the background, after which all particles >20 pixels in size were automatically selected and combined for each cell. After manually adjusting the selections, the integrated density (mean gray value x pixel number) was determined using ImageJ's Measure function. Data were then analyzed with GraphPad.

For cluster analysis, data extracted from SR-Tesseler (cluster diameter, number of detections per cluster, inter-cluster distance) were computed with GraphPad Prism for each cell.

Acknowledgments

We are grateful to Pr. P. Didier for training on dSTORM in HEK293T cells, Dr. F. Rassendren for providing P2X7-mScarlet and P2X-GFP constructs, and Pr. R. Schlichter for providing BV2 cells. This work was supported by the Agence Nationale de la Recherche (Grant ANR-20-CE14-0016-02), the Ministère de la Recherche (PhD grant to B.A., the International Center for Frontier Research in Chemistry (Labex CSC-TGR-18) (PhD grant to F.C.), The Région Grand Est (PhD grant to F.C.), the "École Universitaire de Recherche" Euridol (Programme d'investissement d'Avenir, ANR-17-EURE-0022) (PhD grant extension to B.A.) and was achieved within the NeuroStra Interdisciplinary Thematic Institute of the ITI 2021-2028 program of the University of

829 Strasbourg, CNRS and Inserm. This work has also benefitted from support provided by the
830 University of Strasbourg Institute for Advanced Study (USIAS) for a fellowship, within the French
831 national program "Investment for the future" (Idex-Unistra).

832
833 References

1. Burnstock G (2004) Introduction: P2 receptors. *Curr Top Med Chem* 4(8):793-803.
2. Jacobson KA*, et al.* (2020) Update of P2Y receptor pharmacology: IUPHAR Review 27. *Br
J Pharmacol* 177(11):2413-2433.
3. Illes P*, et al.* (2021) Update of P2X receptor properties and their pharmacology: IUPHAR
Review 30. *Br J Pharmacol* 178(3):489-514.
4. North RA (2002) Molecular physiology of P2X receptors. *Physiol Rev* 82(4):1013-1067.
5. Di Virgilio F, Dal Ben D, Sarti AC, Giuliani AL, Falzoni S (2017) The P2X7 Receptor in
Infection and Inflammation. *Immunity* 47(1):15-31.
6. Ferrari D*, et al.* (1996) Mouse microglial cells express a plasma membrane pore gated by
extracellular ATP. *J Immunol* 156(4):1531-1539.
7. Ferrari D*, et al.* (1997) Extracellular ATP triggers IL-1 beta release by activating the
purinergic P2Z receptor of human macrophages. *J Immunol* 159(3):1451-1458.
8. Monif M, Burnstock G, Williams DA (2010) Microglia: proliferation and activation driven
by the P2X7 receptor. *Int J Biochem Cell Biol* 42(11):1753-1756.
9. Bhattacharya A, Biber K (2016) The microglial ATP-gated ion channel P2X7 as a CNS drug
target. *Glia* 64(10):1772-1787.
10. He Y, Taylor N, Fourgeaud L, Bhattacharya A (2017) The role of microglial P2X7:
modulation of cell death and cytokine release. *J Neuroinflammation* 14(1):135.
11. Illes P (2020) P2X7 Receptors Amplify CNS Damage in Neurodegenerative Diseases. *Int J
Mol Sci* 21(17).
12. Inoue K, Tsuda M (2021) Nociceptive signaling mediated by P2X3, P2X4 and P2X7
receptors. *Biochem Pharmacol* 187:114309.
13. Cao F*, et al.* (2019) P2X7 receptor: A potential therapeutic target for autoimmune
diseases. *Autoimmun Rev* 18(8):767-777.
14. Kan LK, Williams D, Drummond K, O'Brien T, Monif M (2019) The role of microglia and
P2X7 receptors in gliomas. *J Neuroimmunol* 332:138-146.
15. Douguet L*, et al.* (2021) A small-molecule P2RX7 activator promotes anti-tumor immune
responses and sensitizes lung tumor to immunotherapy. *Nat Commun* 12(1):653.
16. Huang Z, Tan S (2021) P2X7 Receptor as a Potential Target for Major Depressive
Disorder. *Curr Drug Targets* 22(10):1108-1120.
17. Huang S*, et al.* (2024) Disruption of the Na(+)/K(+)-ATPase-purinergic P2X7 receptor
complex in microglia promotes stress-induced anxiety. *Immunity* 57(3):495-512 e411.
18. Martin E*, et al.* (2019) New role of P2X7 receptor in an Alzheimer's disease mouse
model. *Mol Psychiatry* 24(1):108-125.
19. Volonte C, Apolloni S, Skaper SD, Burnstock G (2012) P2X7 receptors: channels, pores
and more. *CNS Neurol Disord Drug Targets* 11(6):705-721.
20. Garcia-Marcos M*, et al.* (2006) Characterization and comparison of raft-like membranes
isolated by two different methods from rat submandibular gland cells. *Biochim Biophys
Acta* 1758(6):796-806.
21. Gonnord P*, et al.* (2009) Palmitoylation of the P2X7 receptor, an ATP-gated channel,
controls its expression and association with lipid rafts. *FASEB J* 23(3):795-805.

22.   Barth K*, et al.* (2008) Characterization of the molecular interaction between caveolin-1
876   and the P2X receptors 4 and 7 in E10 mouse lung alveolar epithelial cells. *Int J Biochem*
877   *Cell Biol* 40(10):2230-2239.
878   23.   Barth K*, et al.* (2007) Caveolin-1 influences P2X7 receptor expression and localization in
879   mouse lung alveolar epithelial cells. *FEBS J* 274(12):3021-3033.
880   24.   Garcia-Marcos M, Pochet S, Marino A, Dehaye JP (2006) P2X7 and phospholipid
881   signalling: the search of the "missing link" in epithelial cells. *Cell Signal* 18(12):2098-
882   2104.
883   25.   Weinhold K, Krause-Buchholz U, Rodel G, Kasper M, Barth K (2010) Interaction and
884   interrelation of P2X7 and P2X4 receptor complexes in mouse lung epithelial cells. *Cell*
885   *Mol Life Sci* 67(15):2631-2642.
886   26.   Raouf R, Chabot-Dore AJ, Ase AR, Blais D, Seguela P (2007) Differential regulation of
887   microglial P2X4 and P2X7 ATP receptors following LPS-induced activation.
888   *Neuropharmacology* 53(4):496-504.
889   27.   Trang M, Schmalzing G, Muller CE, Markwardt F (2020) Dissection of P2X4 and P2X7
890   Receptor Current Components in BV-2 Microglia. *Int J Mol Sci* 21(22).
891   28.   Janks L, Sharma CVR, Egan TM (2018) A central role for P2X7 receptors in human
892   microglia. *J Neuroinflammation* 15(1):325.
893   29.   Lord B*, et al.* (2015) A novel radioligand for the ATP-gated ion channel P2X7: [3H] JNJ-
894   54232334. *Eur J Pharmacol* 765:551-559.
895   30.   Choi HB, Ryu JK, Kim SU, McLarnon JG (2007) Modulation of the purinergic P2X7
896   receptor attenuates lipopolysaccharide-mediated microglial activation and neuronal
897   damage in inflamed brain. *J Neurosci* 27(18):4957-4968.
898   31.   Monif M, Reid CA, Powell KL, Smart ML, Williams DA (2009) The P2X7 receptor drives
899   microglial activation and proliferation: a trophic role for P2X7R pore. *J Neurosci*
900   29(12):3781-3791.
901   32.   Monif M*, et al.* (2016) Interleukin-1beta has trophic effects in microglia and its release is
902   mediated by P2X7R pore. *J Neuroinflammation* 13(1):173.
903   33.   Kaczmarek-Hajek K*, et al.* (2018) Re-evaluation of neuronal P2X7 expression using novel
904   mouse models and a P2X7-specific nanobody. *Elife* 7.
905   34.   Manz BN, Groves JT (2010) Spatial organization and signal transduction at intercellular
906   junctions. *Nat Rev Mol Cell Biol* 11(5):342-352.
907   35.   Choquet D, Sainlos M, Sibarita JB (2021) Advanced imaging and labelling methods to
908   decipher brain cell organization and function. *Nat Rev Neurosci* 22(4):237-255.
909   36.   Schermelleh L*, et al.* (2019) Super-resolution microscopy demystified. *Nat Cell Biol*
910   21(1):72-84.
911   37.   Shrivastava AN, Rodriguez PC, Triller A, Renner M (2013) Dynamic micro-organization of
912   P2X7 receptors revealed by PALM based single particle tracking. *Front Cell Neurosci*
913   7:232.
914   38.   Gangadharan V, Nohe A, Caplan J, Czymmek K, Duncan RL (2015) Caveolin-1 regulates
915   P2X7 receptor signaling in osteoblasts. *Am J Physiol Cell Physiol* 308(1):C41-50.
916   39.   Kaczmarek-Hajek K, Lorinczi E, Hausmann R, Nicke A (2012) Molecular and functional
917   properties of P2X receptors--recent progress and persisting challenges. *Purinergic Signal*
918   8(3):375-417.
919   40.   Shiraiwa K, Cheng R, Nonaka H, Tamura T, Hamachi I (2020) Chemical Tools for
920   Endogenous Protein Labeling and Profiling. *Cell Chem Biol* 27(8):970-985.

41. Tamura T, *et al.* (2018) Rapid labelling and covalent inhibition of intracellular native
proteins using ligand-directed N-acyl-N-alkyl sulfonamide. *Nat Commun* 9(1):1870.
42. Karasawa A, Kawate T (2016) Structural basis for subtype-specific inhibition of the P2X7
receptor. *Elife* 5.
43. Guile SD, *et al.* (2009) Antagonists of the P2X(7) receptor. From lead identification to
drug development. *J Med Chem* 52(10):3123-3141.
44. Michel AD, *et al.* (2007) Direct labelling of the human P2X7 receptor and identification
of positive and negative cooperativity of binding. *Br J Pharmacol* 151(1):103-114.
45. Bhattacharya A, *et al.* (2013) Pharmacological characterization of a novel centrally
permeable P2X7 receptor antagonist: JNJ-47965567. *Br J Pharmacol* 170(3):624-640.
46. Allsopp RC, Dayl S, Schmid R, Evans RJ (2017) Unique residues in the ATP gated human
P2X7 receptor define a novel allosteric binding pocket for the selective antagonist
AZ10606120. *Sci Rep* 7(1):725.
47. Surprenant A, Rassendren F, Kawashima E, North RA, Buell G (1996) The cytolytic P2Z
receptor for extracellular ATP identified as a P2X receptor (P2X7). *Science*
272(5262):735-738.
48. Karasawa A, Kawate T (2017) Expression and Purification of a Mammalian P2X7
Receptor from Sf9 Insect Cells. *Bio Protoc* 7(17).
49. Levet F, *et al.* (2015) SR-Tesseler: a method to segment and quantify localization-based
super-resolution microscopy data. *Nat Methods* 12(11):1065-1071.
50. Perregaux DG, McNiff P, Laliberte R, Conklyn M, Gabel CA (2000) ATP acts as an agonist
to promote stimulus-induced secretion of IL-1 beta and IL-18 in human blood. *J Immunol*
165(8):4615-4623.
51. Ferrari D, *et al.* (2006) The P2X7 receptor: a key player in IL-1 processing and release. *J*
Immunol 176(7):3877-3883.
52. Swanson KV, Deng M, Ting JP (2019) The NLRP3 inflammasome: molecular activation
and regulation to therapeutics. *Nat Rev Immunol* 19(8):477-489.
53. Dunning K, *et al.* (2021) P2X7 Receptors and TMEM16 Channels Are Functionally
Coupled with Implications for Macropore Formation and Current Facilitation. *Int J Mol*
Sci 22(12).
54. Franceschini A, *et al.* (2015) The P2X7 receptor directly interacts with the NLRP3
inflammasome scaffold protein. *FASEB J* 29(6):2450-2461.
55. Miras-Portugal MT, Sebastian-Serrano A, de Diego Garcia L, Diaz-Hernandez M (2017)
Neuronal P2X7 Receptor: Involvement in Neuronal Physiology and Pathology. *J Neurosci*
37(30):7063-7072.
56. Illes P, Khan TM, Rubini P (2017) Neuronal P2X7 Receptors Revisited: Do They Really
Exist? *J Neurosci* 37(30):7049-7062.
57. Bindels DS, *et al.* (2017) mScarlet: a bright monomeric red fluorescent protein for
cellular imaging. *Nat Methods* 14(1):53-56.
58. Habermacher C, *et al.* (2016) Photo-switchable tweezers illuminate pore-opening
motions of an ATP-gated P2X ion channel. *Elife* 5:e11050.
59. Olivier N, Keller D, Rajan VS, Gonczy P, Manley S (2013) Simple buffers for 3D STORM
microscopy. *Biomed Opt Express* 4(6):885-899.
60. Glushonkov O, Real E, Boutant E, Mely Y, Didier P (2018) Optimized protocol for
combined PALM-dSTORM imaging. *Sci Rep* 8(1):8749.

61.   Edelstein AD*, et al.* (2014) Advanced methods of microscope control using muManager
software. *J Biol Methods* 1(2).
62.   Ovesny M, Krizek P, Borkovec J, Svindrych Z, Hagen GM (2014) ThunderSTORM: a
comprehensive ImageJ plug-in for PALM and STORM data analysis and super-resolution
imaging. *Bioinformatics* 30(16):2389-2390.
63.   Butler C*, et al.* (2022) Multi-Dimensional Spectral Single Molecule Localization
Microscopy. *Front Bioinform* 2:813494.

973
974

975
976 **Figures**
977

978
979
Figure 1. Affinity-guided labeling strategy for P2X7. (**A**) Schematic representation of the P2X7
labeling strategy using ligand-directed *N*-cyanomethyl NASA chemistry. The biotin tag on the NASA
molecule enables super-resolution imaging of nanoscale P2X7 localization using Strept-A 647
probe through a highly specific biotinylation reaction involving endogenous lysine (K) residues. Lg:
P2X7 ligand. (**B**) Crystal structure of panda P2X7 (pdP2X7) shown in ribbon representation, bound
to AZ10606120 depicted as spheres (PDB:5U1W) (42, 48). One of the three ATP-binding sites and
the approximate location of the membrane are also indicated. Inset, enlarged view of the
AZ10606120-binding pocket, rotated 180°. Distances (in Å) between the α-carbon of selected
lysines and the hydroxyl group of AZ10606120 are displayed. Note that K300 is not visible in this
view. (**C**) Chemical structure of **X7-uP**.
990
991

993
Figure 2. X7-uP is a potent P2X7 inhibitor that rapidly labels ectopically expressed P2X7 in
HEK293T cells. (**A**) Whole-cell currents evoked by 10 μM BzATP are reversibly inhibited by co-
applying 1 μM **X7-uP** (upper trace) or 1 μM AZ10606120 (middle trace) in cells transiently
transfected with rP2X7. Inhibitors were pre-applied alone for 8 s before 2 seconds of co-
application. In the control (absence of inhibitors), BzATP-induced currents further increased,
demonstrating current facilitation, which is an expected feature of P2X7 activation (47). (**B**)
Summary of whole-cell inhibition (n = 4 cells for each condition). Bars represent mean ± s.e.m.
(**C**) Western blot analysis of P2X7 labeling by **X7-uP**. Cells transiently transfected with P2X7c-
myc were treated with 1 μM **X7-uP** for 0-60 minutes, in the absence or presence of 10 μM
AZ10606120 or 10 μM A740003 (as indicated), followed by extensive washing. After cell lysis,
biotinylated proteins were pulled down, separated on SDS-PAGE, and Western blotting was
revealed using an anti-c-myc antibody (@c-myc). Molecular mass markers are shown on the
right. Control of P2X7c-myc expression is presented in the corresponding input. β-Actin was used
as a loading control. (**D**) Time course plot of P2X7 labeling with 1 μM **X7-uP**. Data (mean ±
s.e.m., n = 3 independent transfections) were fitted with Eq. (1) to determine the pseudo-first-
order reaction rate k_{app} (mean ± s.e.m.).

1012

1013
Figure 3. X7-uP labeling is highly selective for P2X7. (**A-B**) Confocal images of HEK293T
cells transiently transfected with either P2X7-mScarlet (**A**) or various P2X subunits tagged with
GFP (P2X1-GFP, P2X2-GFP, P2X3-GFP, P2X4-GFP, P2X5-GFP, or P2X6-GFP) (**B**) were
labeled with **X7-uP** and revealed using Strept-A 647 (red) in FBS-free DMEM. Labeling was
performed in the presence of 10 μM AZ10606120 or 10 μM A740003 (**A**). Nuclei were stained
with Hoechst (blue). For clarity, mScarlet and GFP signals are displayed in green. Scale bars, 10
μm. (**C**) Quantification of Alexa 647 fluorescence. Bars represent mean ± standard deviation
(s.d.) (n = 75-129 cells, *t*-test comparisons to P2X7-mScarlet, ****P < 0.0001.

1022
1023

1024
Figure 4. X7-uP labels K82 and K117 in rat P2X7. (**A**) Molecular docking of **1** (same pose as
shown in Figure 1 — figure supplement 1C) in pdP2X7, showing distances (in Å) between the
reactive carbonyl of **1** (stick representation) and selected α-carbons of nearby residues (blue).
Residues shown in parentheses correspond to equivalent rP2X7 residues. (**B**) Confocal images
of HEK293T cells transiently transfected with different P2X7 constructs: P2X7-mScarlet, K82A,
K117A, and K82A/K117A. Scale bars, 10 μm. (**C**) Quantification of Alexa 647 fluorescence. Bars
represent mean ± s.d. (n = 90-190 cells, *t*-test comparisons to indicated conditions, ****$P <$
0.0001). (**D**) Whole-cell currents evoked by 10 μM BzATP are reversibly inhibited by co-
application of 0.5 μM **X7-uP** (upper trace) to BzATP in a cell transiently transfected with the
double mutant K82A/K117A. The control (absence of **X7-uP**) is shown in the bottom trace. (**E**)
Summary of whole-cell inhibition for K82A/K117A (n = 7 cells for **X7-uP** and 5 cells for control).
Bars represent mean ± s.e.m.; Mann-Whitney test (**$P <$ 0.005).
1037
1038

Figure 5. dSTORM data revealed nanoscale P2X7 plasma membrane localization in BV2 cells. (**A**) Cartoon and experimental timeline of BV2 cell treatments. IL-1β release was assessed in the supernatant (sup), and the same cells were labeled with 1 mM **X7-uP** after extensive washout. (**B**) Quantification of IL-1β release by ELISA following the indicated treatments: LPS (1 mg/mL for 24 h), ATP (1 mM for 30 min), BzATP (300 mM for 30 min), and MβCD (15 mM for 15

min). Bars represent mean ± s.e.m. (n = 12 samples from 3 independent experiments). Data were
compared using Kruskal-Wallis followed by Dunn's multiple comparisons (*P = 0.0208, #P =
0.0362, **P = 0.0014, ****P < 0.0001). (**C**) Normalized quantification of IL-1β release induced by
LPS+ATP or LPS+BzATP in the presence of P2X7 inhibitors AZ10606120 or A740003. Bars
represent mean ± s.e.m. (n = 6 samples from 6 independent experiments). One-way ANOVA with
Dunnett's multiple comparisons to control condition for ATP data (****P < 0.0001). Kruskal-Wallis
followed by Dunn's multiple comparisons to control condition for BzATP data (*P = 0.0414, *** P =
0.0006). (**D**) Bright-field and dSTORM images of **X7-uP**-labeled BV2 cells revealed with Strept-A
647 corresponding to experiments shown in panel b. Scale bars, 10 mm. Insets: Magnified
dSTORM images. Scale bars, 1 mm. (**E**) Quantification of single P2X7 localization density. Bars
represent mean ± s.e.m. (each data point represent a cell, n = 3 independent experiments). One-
way ANOVA with Tukey's multiple comparisons (*P < 0.019, #P < 0.0194, @P < 0.0477, ***P <
0.0002, ###P < 0.0008, ****P < 0.0001). (**F**) Relative frequency of cluster size. Inset: percentage
of clusters larger than 0.025 mm^2. (**G**) Number of detections per cluster. Bars represent mean ±
s.e.m. One-way ANOVA with Tukey's multiple comparisons (*P < 0.0197, #P < 0.0277, @P <
0.0389, ****P < 0.0001). (**H**) Images showing tessellation analysis of cells treated either with
LPS+ATP or left untreated. Inset: magnification. Scale bars, 200 nm. (**I**) Number of fluorophores
per cluster. Bars represent mean ± s.e.m. One-way ANOVA with Tukey's multiple comparisons
(****P < 0.0001).
1064

1065
Figure 6. Nanoscale redistribution of individual P2X7 receptors in microglia under pro-
inflammatory conditions at the plasma membrane. The cartoon illustrates two distinct clusters
of P2X7 receptors (blue), each adorned with one, two, or three fluorescently tagged tetrameric
biotin-bound streptavidin (red). In untreated cells, each cluster contains an average of 1.5
fluorophores per P2X7 receptor. Treatment with LPS and ATP promotes P2X7 clustering by
increasing the average number of fluorophores per cluster to between 4 and 5, resulting in an
increased number of P2X7 receptors per cluster, from one to three. This redistribution
synergistically triggers IL-1β release.

1074