**Caudata macrogenetics: Geographic attributes and lineage age predict global patterns of**

**mitochondrial genetic variation in salamanders**

**Short running title:** Macrogenetics in salamanders worldwide

**Abstract**

**Aim:** Genetic diversity contains valuable information about ecological and evolutionary aspects

of species. Intraspecific genetic variation is shaped by species' natural history traits and by

characteristics of geography and climate within their ranges. Amphibians are of ecological and

conservation interest because of their global distribution, deep history, trait diversity, and roles

within ecological communities. Here, we studied genetic variation within salamanders to

investigate predictors of nucleotide diversity and spatial patterns of genetic differentiation.

**Location:** Global.

**Time Period:** Present.

**Major Taxa Studied:** Salamanders.

**Methods:** We repurposed mitochondrial DNA sequences and ecological data from open-access

databases for 220 salamander species. We calculated nucleotide diversity ($\pi$) and tested for

isolation by distance (IBD) and isolation by environment (IBE). We analyzed these three

variables with random forest and phylogenetic comparative methods using 28 predictors

expected to be associated with genetic variation.

**Results:** We recovered 8,108 Cytb sequences with their associated geographic coordinates, of

which 7,007 sequences were manually curated by us. Range size, lineage age, and sample size

were important predictors of genetic variation. We found higher diversity in regions including

the Neotropics and central-eastern Europe. The absence of phylogenetic signal in $\pi$, IBD, and

IBE suggests that genetic variation is shaped by local ecological and geographical factors rather

than by shared ancestry.

**Main Conclusions:** Our finding of range size as an important predictor aligns with theoretical

expectations that species with larger ranges tend to harbor more genetic diversity. Furthermore,

lineage age being an important predictor is in line with the clade-age hypothesis, in which

species with longer divergence times have higher genetic diversity because they have had more

time to accumulate genetic variation. Our results underscore the importance of integrating spatial

data into macrogenetic studies, providing valuable information for future studies and

conservation strategies targeting regions with high or low genetic diversity.

**Keywords**

Amphibians, clade-age hypothesis, isolation-by-distance, isolation-by-environment, nucleotide

diversity, species range size.

## Introduction

Genetic variation plays a foundational role in maintaining biological diversity (Kardos et al., 2021). One significant evolutionary factor that predicts genetic variation within populations is the number of individuals that contribute DNA to the next generation (the effective population size; Wright, 1931). Genetic variation is proportional to population size, as explained by the neutral theory of molecular evolution (Kimura, 1968). Gene flow between populations is another key factor influencing variation because it can add and maintain novel alleles, counteract the effects of genetic drift and selection, and enable population and species persistence (Park et al., 2024). Natural selection also has an important role in affecting neutral diversity of populations through background or linked selection (Corbett-Detig et al., 2015). Understanding this variation and which factors influence it across different geographic and environmental contexts is essential for elucidating population dynamics and will help us identify populations with conservation needs (Pauls et al., 2013). For example, declining levels of genetic variation may be further exacerbated under global change scenarios due to the reduction of suitable habitat for species (Schierenbeck, 2017).

Climatic and geographic factors shape patterns of dispersal and population size through mechanisms such as habitat connectivity, environmental stability, and barriers to gene flow. These processes, in turn, determine the extent and distribution of genetic variation within species (e.g., Hanson et al., 2017). For example, geographic distance can limit gene flow among populations, resulting in patterns of isolation-by-distance (IBD), where genetic dissimilarity increases as populations become more spatially separated (Wright, 1943). Similarly, environmental differences can restrict gene flow by selecting against migrants non-adapted to the new environmental conditions, leading to isolation-by-environment (IBE), which reflects the influence of habitat heterogeneity and ecological gradients on genetic variation (Wang and Bradburd, 2014). Historical climatic fluctuations and complex topography often act together to promote cycles of population contraction and expansion, which generate and maintain genetic variation. In the alpine toad *Scutiger ningshanensis*, for instance, repeated glacial cycles during the Pleistocene, combined with uplifts of mountain ranges, likely fragmented populations and created refugia, increasing opportunities for genetic differentiation (Meng et al., 2014). Life

history traits, particularly those related to reproduction, also influence genetic diversity by

affecting effective population size and gene flow (Ellegren and Galtier, 2016). For example,

species with high fecundity, widespread dispersal, or external fertilization can maintain large and

connected populations, which reduces the effects of genetic drift and promotes diversity. Broad-

scale analyses across hundreds of plant and animal species further support the idea that

reproductive traits and climatic variables jointly shape genetic diversity by influencing

demographic stability and dispersal potential (De Kort et al., 2021), suggesting that these

variables act as consistent predictors of genetic variation across taxa.

Studies on focal taxa within a limited geographic range provide the resolution needed to

understand local genetic patterns and idiosyncrasies (e.g., Fonseca et al., 2021). It is equally

important to investigate genetic variation on broader geographic and taxonomic scales, to have a

global perspective of the role of ecological and geographic factors on evolutionary patterns.

Macrogenetics is an approach to analyze open-access DNA sequences across large scales and

investigate patterns of intraspecific genetic diversity across higher taxonomic groups (Blanchet

et al. 2017). Macrogenetic research has gained popularity over the past decade, driven by the

increasing availability of molecular resources (e.g., GenBank), computational tools (e.g.,

machine learning), and trait databases (Leigh et al., 2021). Despite limitations, particularly the

genetic markers represented by available data (see Millette et al., 2021; Paz-Vinas et al., 2021),

efforts to understand large-scale patterns and predictors of intraspecific genetic diversity have

been undertaken across various taxonomic groups. A study of global genetic diversity in insects

attributed high levels of mitochondrial DNA (mtDNA) genetic diversity in the subtropics to

temperature and climatic stability (French et al., 2023). Another study of nucleotide diversity

using mtDNA for animals and chloroplast DNA for plants in over 38,000 species found that

latitude significantly correlates with genetic diversity, with tropical species exhibiting higher

intraspecific diversity (Fonseca et al., 2023). These studies highlight the potential of

macrogenetics to uncover important predictors of intraspecific genetic diversity across broad sets

of taxa globally.

Amphibians contain remarkable diversity in life history strategies and the geographic and

climatic variability of their habitats. Recent studies of amphibian genetic diversity indicate that

biogeographic region can influence which factors predict intraspecific genetic variation. For

example, in Nearctic amphibians, taxonomic family, the number of sequences, and latitude were

key predictors of intraspecific variation (Barrow et al., 2021), whereas range size, elevation,

latitude, and precipitation predicted genetic variation in Neotropical amphibians (Amador et al.,

2024). In the Americas, Lawrence et al. (2023) found a strong relationship between

environmental variables and amphibian expected heterozygosity ($H_E$), and that amphibian $H_E$

decreased with sample size (number of individuals). Within amphibians, salamanders are

particularly valuable for macroevolutionary studies because of their broad distribution (excluding

Oceania, the Afrotropics, and Antarctica), manageable number of species, and unique traits.

They display remarkable ecological and life history diversity that can directly shape population

dynamics and, consequently, genetic structure. For instance, body size variation, from the tiny

*Thorius arboreus* (~20 mm) to the 1.8-meter-long *Andrias japonicus,* can influence dispersal

ability and home range size, thereby affecting gene flow and population structure. Likewise,

salamanders include generalist species like the tiger salamander, *Ambystoma tigrinum*, versus

habitat specialist species adapted to aquatic, terrestrial, or arboreal environments, which can

promote or constrain connectivity across landscapes.

One trait that may influence patterns of genetic diversity in salamanders is developmental mode.

Species with direct development (i.e., without a larval stage) often have limited dispersal, which

could lead to smaller, isolated populations favoring population differentiation (Paz et al. 2015;

Liedtke et al. 2022). In contrast, species with a larval stage and dispersing terrestrial adults may

maintain more interconnected populations, reducing genetic differentiation (e.g., Zamudio and

Wieczorek 2007). Some salamanders exhibit paedomorphosis, a reproductive strategy in which

adults retain larval characteristics, which reduces gene flow by allowing persistent aquatic life

cycles in species that could otherwise be terrestrial (Denoël et al., 2005). Recent studies have

begun to untangle the relationships between species-level traits and genetic variation. For

example, Segovia-Ramírez et al. (2023) used genomic data from 62 Neotropical salamander

species (one individual per species) to demonstrate that precipitation variability and snout-vent

length (body size) were significant predictors of genomic diversity, perhaps related to their roles

in shaping population persistence and abundance. Parsons et al. (2024) combined genetic,

geographic, climatic, and life history data from salamanders and used a machine learning model

to identify predictors of unrecognized genetic lineages. They found that Caudata hidden diversity

is the result of variation in climatic variables. A more comprehensive, georeferenced view of

intraspecific genetic variation in salamanders, encompassing more species and potential

predictors (e.g., climatic and natural history traits), would provide useful insights into the

evolutionary and demographic processes driving variation in this group.

Here, we combine molecular, phenotypic, and environmental open-access data to investigate the

determinants of intraspecific genetic variation in salamanders. We used machine learning

techniques and phylogenetic comparative methods to determine whether natural history traits or

climatic or environmental variables can predict genetic variation within species on a global scale.

We hypothesized that reproductive traits, such as the total number of eggs (a proxy for fitness)

and reproductive mode (e.g., larval vs. direct development), would predict intraspecific genetic

variation in salamanders. Species that produce more offspring may maintain larger, more stable

populations, which can preserve higher levels of genetic diversity over time (Lacy, 1987; Reed

and Frankham, 2003). Additionally, reproductive mode could influence dispersal, leading to

lower expected connectivity and more genetic differentiation in species with direct development

compared to those with aquatic larvae. We also predicted that traits associated with total species

population size (e.g., range size) would be positively correlated with genetic variation, as larger

ranges may support more numerous and genetically diverse populations (Frankham, 1996). We

hypothesized that species with older evolutionary age (lineage age) may have higher genetic

diversity because they have had more time to accumulate variation than younger species (the

clade-age hypothesis; see McPeek and Brown, 2007; Scholl and Wiens, 2016). Finally, we

expected high spatial autocorrelation in nucleotide diversity of salamanders overall based on

their limited dispersal capabilities, which can lead to genetic clustering at small scales (Fusco et

al., 2021).

**Methods**

*Data collection*

We followed the taxonomic classification proposed by AmphibiaWeb (amphibiaweb.org) and
Amphibian Species of the World (https://amphibiansoftheworld.amnh.org/) and prepared a
detailed count of all species included in this study (Supplementary Material 1, SM1). These
databases were also used to reconcile species names and verify taxonomic consistency across
datasets. When discrepancies or outdated names were found, we updated species names to match
current taxonomy, with a description of our decisions in SM1. We repurposed Caudata DNA
sequences, natural history traits, and geographic and environmental data from open-access
databases. Following the approach of recent research on the determinants of genetic diversity of
multiple amphibian species (e.g., Barrow et al., 2021; Amador et al., 2024), we assembled
sequences of the cytochrome-b (Cytb) mitochondrial gene. This gene is suitable for genetic
diversity studies because it contains both conserved and variable regions, is commonly used in
amphibian single-locus phylogenetic and population genetic studies, and is therefore the most
abundant gene in open-access databases for amphibians. Sequences were obtained from
GenBank (National Center for Biotechnology Information, NCBI), phylogatR (Pelletier et al.
2022), the *phruta* R package (Román□Palacios, 2023), and Amador et al. (2024). We included
species with at least 10 sequences, seeking a better representation of intraspecific genetic
variation. Our dataset contained species with high variation in sampling effort, such as those with
more than 200 sequences. To account for this sampling bias, we also conducted a set of analyses
on randomly subsampled datasets that included 10–20 sequences per species. Alignments were
generated, visualized, and edited with the MUSCLE aligner v.3.8.31 (Edgar, 2004) in AliView
v.1.28 (Larsson, 2014) and saved in FASTA format.

The geographic coordinates (latitude and longitude in decimal degrees format) previously
associated with each Cytb sequence were recovered from GenBank, phylogatR, and Amador et
al. (2024); corresponding to 31 species (14.1% of the total species) and 1,169 sequences (14.3%
of the total sequences). We then manually retrieved the remaining geographic coordinates
associated with the sequences (corresponding to 189 species and 7,007 sequences) following a

tutorial prepared for this purpose (see Supplementary Material 2). We georeferenced localities

using either the function geocode() of the *tidygeocoder* R package (Cambon et al., 2021) or the

GEOLocate web application (Rios and Bart, 2010). We only retained sequences that were

georeferenced with an error of less than 20 km, resulting in 12 species with fewer than 10

georeferenced sequences.

We calculated both the total range size and the sampled range size of DNA sequences for each

species as follows. For total range size, we obtained the geographic range map in shapefile

format for each species from the International Union for Conservation of Nature (IUCN) portal

using the getIUCN() function of the *rasterSp* R package (RS-eco, 2023). Species ranges not

available in the IUCN portal (24 species) were generated with minimum convex polygons using

the sequence coordinates and the function mcp() in the R package *adehabitatHR* (Calenge, 2024;

see Supplementary Material). This same approach was also used to obtain the geographic range

of the sampled sequences for all species (i.e., using only the sequence occurrences). Range size

was calculated from shapefiles using the areaPolygon() function of the *geosphere* R package

(Hijmans, 2022). Elevation and latitude (min, max, and mean) were obtained from the retrieved

geographic coordinates using the function get_elev_point() of the *elevatr* R package (Hollister et

al., 2025) for elevation, and custom R scripts for latitude (see Supplementary Material 3). We

also obtained precipitation and temperature data (mean and standard deviation) from WorldClim

v2.1 (Fick and Hijmans, 2017) using the function worldclim_global() of the *geodata* R package

v0.6-2 (Hijmans et al., 2024).

We used the same natural history and geographic traits as in Amador et al., (2024) with slight

additions and omissions, choosing predictors we expected to be related to genetic variation (e.g.,

body size, reproductive mode, elevation, latitude, species range size, precipitation and

temperature; Table S1). We omitted the variable activity because nearly all salamanders (except

three of the 220 species) included in our dataset were nocturnal. Eight new predictors were added

to the matrix (28 predictors in total), which included sampling effort/coordinates (i.e., number of

sequences with associated coordinates), sequence length/coordinates (i.e., number of base pairs

of sequences with associated coordinates), snout-vent length in millimeters (SVL),

paedomorphism (whether or not juvenile features are retained as an adult), litter size min

(minimum number of offspring or eggs per clutch), litter size max (maximum number of

offspring or eggs per clutch), latitudinal midpoint (mean of the minimum and maximum

latitudinal values), and lineage age (estimated species divergence times obtained from Stewart

and Wiens, 2025) (Table S1). We obtained trait information from the open-access database

AmphiBIO (Oliveira et al. 2017), species accounts in AmphibiaWeb (AmphibiaWeb, 2024), and

relevant literature.

*Measures of genetic variation*

We evaluated overall genetic diversity (range-wide nucleotide diversity) and spatial patterns of

genetic variation (IBD and IBE) within salamander species. We calculated nucleotide diversity

($\pi$) with the function nuc.div() of the *pegas* R package (Paradis, 2010) for all sequences ($\pi_A$) and

sequences with associated coordinates only ($\pi_S$). Genetic distances (*gendist*) for each species

were calculated based on the raw difference between sequences, using the dist.dna() function of

the *ape* R package (Paradis and Schliep, 2019). We tested IBD and IBE for each salamander

species as follows. Geographic distance (*geodist*) was calculated between each pair of sequence

coordinates using Euclidean distance with the dist() function of the *stats* package (R Core Team,

2023) in R. Before calculating environmental distances, we used the function scale() in R to

standardize variables measured in different units (e.g., temperature - °C or precipitation - mm).

Environmental distance (*envdist*) was calculated between each pair of sequence coordinates

based on the 19 variables of the WorldClim v2.1 database using the vegdist() function of the

*vegan* R package (Oksanen et al., 2024). This function calculates a distance matrix, selecting all

environmental variables using Euclidean distances. We performed Multiple Matrix Regression

with Randomization (MMRR) analyses for each species (Wang, 2013). MMRR quantifies the

relative effects of IBD and IBE, represented as distance matrices (*geodist* and *envdist*,

respectively), in explaining genetic variation, represented by genetic distances. We compared the

observed correlations between *geodist* vs. *gendist* (IBD) and *envdist* vs. *gendist* (IBE) to the

correlations from 1,000 permutations. Species were coded as 'Yes' if they did present IBD and

IBE or 'No' if they did not based on an assessment of the associated p-values from the

permutation tests. To account for multiple comparisons, we applied a strict Bonferroni correction

(Bonferroni, 1936) to reduce the p-value for assessing significance. We used the output (yes or

no IBD; yes or no IBE) as binary response variables for further analyses.

We mapped $\pi_S$ to visualize the distribution of genetic diversity across the globe and assess

differences among biogeographic realms (Nearctic, Neotropic, Oriental, and Palearctic). Briefly,

following Amador et al. (2024), we combined sequences within a species based on a maximum

distance of 100 km between coordinates, and calculated $\pi_S$ for each locality (see Supplementary

Material 4). We then mapped $\pi_S$ for four different grid sizes (1–4), which correspond with one to

four decimal degree grids, or approximately 110 km$^2$ to 440 km$^2$. We also visualized the number

of sequences per grid cell for the four resolutions. For these visualizations, we used the R

packages *ggplot2* (Wickham, 2016), *ggspatial* (Dunnington, 2023), and *sf* (Pebesma, 2018;

Pebesma and Bivand, 2023). Finally, we tested for spatial autocorrelation for $\pi$ and the number

of sequences separately with the Moran's I test using the moran.test() function in the *spdep* R

package (Bivand et al., 2013).

To further understand the potential link between historical demography and the patterns of

genetic variation we found, we used a common statistic for evaluating population size change,

Tajima's D (Tajima, 1989), using the tajima.test() function of the *pegas* package. This test is

compatible with different interpretations related to natural selection versus demographic change,

making it difficult to distinguish among hypotheses (Yang, 2014). Therefore, we evaluate these

results for comparison rather than include the statistic as a variable in our primary analyses. We

followed Fonseca et al. (2023) in assuming that the demographic change interpretation is likely

more relevant for global comparisons. Thus, in our study, we interpret Tajima's D values close to

zero as representative of neutrality (no population size change over time), positive values as

evidence of recent population contraction, and negative values as recent population expansion.

*Predicting genetic variation using random forests (RF)*

We used random forest (RF) regression to predict $\pi$ and RF classification to predict IBD and IBE

based on a set of geographic range characteristics and ecological variables. We built RF models

with the function randomForest() in the *randomForest* R package (Liaw & Wiener, 2002), with

10,000 trees and 100 permutations per model. We used the tuneRF() function to obtain the

optimal mtry parameter (number of variables that are randomly sampled as candidates at each

split) and then ran a new RF regression or classification analysis using the best mtry value. The

data was split into a training set (70% of the data) for building models and a testing set (30%) for

making predictions. For RF regression, we assessed models using the R-squared ($R^2$) statistic,

which represents the proportion of the variance explained by the model, with a higher $R^2$ value

indicating a better fit and more predictive power. The relative importance of the predictor

variables for $\pi_A$ and $\pi_S$ was evaluated based on the percentage increase in mean squared error

(%IncMSE).

In RF classification, in addition to the original models, we also tried upsampling and

downsampling to address the large class imbalance within our datasets (yes IBD = 56 species, no

IBD = 161 species; yes IBE = 39 species, no IBE = 178 species; based on MMRR results with

Bonferroni correction). For downsampling, the majority cases (no IBD/IBE) were randomly

subsampled to equal the number of minority cases (yes IBD/IBE). For upsampling, minority

cases were randomly duplicated to equal the number of majority cases. We chose the best model

based on the lowest out-of-bag (OOB) error rate. To assess overfitting and improve cross-

validation in the model, we split our data into a training set and a test set (70% train, 30% test).

To evaluate predictions of the RF classification models, we created a confusion matrix with the

confusionMatrix() function of the *caret* R package (Kuhn, 2008) and tested whether model

accuracy was significantly better than the no information rate (NIR). Finally, we assessed the

relative importance of the variables based on the Mean Decrease Accuracy (MDA) for the RF

classification models of IBD and IBE.

*Evaluating predictors of genetic variation with phylogenetic comparative methods*

For phylogenetic analyses, we downloaded a phylogeny subset from VertLife.org for the species

in our dataset, which represents phylogenetic relationships inferred by Jetz and Pyron (2018).

We used a subset of 180 species for these analyses because 40 species from our dataset are not

included in the available phylogeny. Using a custom R script (Supplementary Material 3), we

edited the name of 12 species in the tree to match the names in the dataset (e.g., some

*Speleomantes* and *Hypselotriton* species). To assess the potential effect of phylogenetic history

in our models, we tested for phylogenetic signal in $\pi$ using Pagel's $\lambda$ (lambda; Pagel, 1999) and

Blomberg's K (Blomberg et al., 2003) implemented in the *phytools* R package (Revell, 2012)

with the function phylosig(). For IBD and IBE, we used the *D* statistic implemented in the

package *caper* with the function phylo.d() to assess phylogenetic signal.

We used phylogenetic independent contrasts (PIC; Felsenstein, 1985) to investigate the

evolutionary correlation between $\pi$ and the most important predictors identified in the RF

analyses. We used the pic() function of the *ape* R package to compute PIC for both response and

explanatory variables. We then fit linear models in R for both the uncorrected and

phylogenetically corrected (PIC) response and predictor variables. We also used phylogenetic

generalized linear mixed models (PGLMMs) to analyze binary traits as described in Ives and

Helmus (2011). PGLMMs allow us to account for phylogenetic covariance while reducing Type

I error rates. This method was used to analyze IBD and IBE as binary dependent variables (0 for

no IBD/IBE, 1 for yes IBD/IBE) versus the most important predictors based on RF results.

Because of the nature of our data (binary outcomes), we chose binomial as the error family with

a logit link function. We compared IBD vs. sampling range size, number of sequences,

maximum latitude, and mean precipitation; and IBE vs. sampling range size, number of

sequences, mean elevation, and mean latitude. We implemented this analysis using the function

pglmm() in the *phyr* R package (Li et al., 2020), including a phylogenetic random effect to

account for the non-independence among species due to shared evolutionary history.

**Results**

*Data summary*

We obtained 12,961 Cytb sequences from 220 species distributed globally, of which 8,937 sequences were retrieved directly from GenBank, 2,349 sequences from phylogatR, 1,657 from the *phruta* package, and 296 from Amador et al. (2024). Of these, we obtained geographic coordinates for 8,108 sequences from 219 species. A single species, *Plethodon sherando*, did not have associated coordinates, therefore was not included in IBD and IBE analyses; and 11 species had between four and nine sequences with coordinates. The final dataset included 220 species in nine families (representing >26% of species and 90% of families globally), three response variables ($\pi$, IBD, and IBE), and 28 predictors (Table S2). All sequences associated with geographic coordinates and R scripts used to retrieve sequences and coordinates are available as Supplementary Material 4.

*Patterns of genetic variation in global salamanders*

Nucleotide diversity within species ranged from $\pi_A = 0$ to $\pi_A = 0.087$, with *Pseudoeurycea lineola* exhibiting the highest diversity when all sequences were analyzed. When considering only sequences with associated coordinates, values ranged from $\pi_S = 0$ to $\pi_S = 0.089$, with *Desmognathus amphileucus* showing the highest diversity. . We found that nucleotide diversity for all sequences ($\pi_A$) and for the sequences with associated coordinates ($\pi_S$) were highly correlated (Figure S1; Pearson's correlation $r = 0.95$, $t(218) = 46.18$, p-value < 0.001). Based on this validation, we conducted further analyses using sequences with associated coordinates ($\pi_S$). *Bolitoglossa yariguiensis* was the only species with a value of $\pi = 0$ (Table S2; Figure S2). We found that $\pi$ and the number of sequences were not randomly distributed across the world map (Figure 1, Figures S3 – S5). For example, regions with high genetic diversity included the Ecuadorian Amazon and central-southern Mexico in the Neotropics; and central-eastern Europe in the Palearctic. Areas with low genetic diversity were identified in northern North America (Nearctic), western Europe, and central-western China in the Oriental realm (Figure 1). Furthermore, we found statistically significant, but weak, positive spatial autocorrelation in both

$\pi_S$ (Moran's I = 0.1435, p-value < 0.0001) and the number of sequences (Moran's I = 0.4136, p-

value < 0.0001).

Most species did not present IBD and IBE patterns, with 161 species (74.2%) showing no IBD

and 178 species (82%) showing no IBE (Table S2; Figure S6). We found that only 21 species

had significant patterns for both IBD and IBE (Table S2). The Neotropical and Oriental realms

had the lowest percentages of species that showed IBD and IBE (IBD: 13.3% and 24.4%, IBE:

13.3% and 10.8% respectively; Figure S7). The Palearctic region had the highest percentage of

species that showed IBD (31%) and IBE (25.4%) patterns (Figure S7).

Genetic variation was randomly distributed across the phylogeny (Figure 2). For example, we

found species in the same genus with dissimilar $\pi$ values, such as *Plethodon* (range = 0.0008 –

0.0698) and *Eurycea* (range = 0.0015 – 0.0768) in Plethodontidae, *Paramesotriton* (range =

0.001 – 0.0767) in Salamandridae, and *Hynobius* (range = 0.0008 – 0.0853) in Hynobiidae. We

found no or low phylogenetic signal in $\pi$ based on the method used, for example, with Pagel's $\lambda$

= 0.0005, p-value (based on LR test) > 0.05; and Blomberg's K = 0.053, p-value (based on 1000

randomizations) > 0.05 (Figure S8). We found no significant phylogenetic signal in IBD

(Estimated $D_{IBD}$ = 0.945, p-value = 0.249), and significant but moderate phylogenetic signal in

IBE (Estimated $D_{IBE}$ = 0.782, p-value = 0.012) (Figure S9).

Our results from assessing Tajima's D indicate that many species in our dataset (n = 109) have

undergone recent population expansion (i.e., negative Tajima's D, more recent mutations than

expected under neutrality; Figure S10). Positive Tajima's D values indicate more alleles at

intermediate frequencies than expected under neutrality, possibly because of a recent population

contraction or bottleneck, as we observed in 51 salamander species (Figure S10). Fifty-nine

species do not show strong evidence for population expansion or contraction (Tajima's D ≈ 0)

suggesting no deviation from expectations under neutrality (Table S2; Figure S10).

*Predictors of genetic variation in global salamanders*

Our random forest analyses showed an influence of sampling effort on genetic variation

estimates, with the number of sequences (with associated geographic coordinates) as a top

predictor of IBD, and IBE (Figure 3). Additionally, we compared both species range size

variables in our dataset, total range size versus sampling range size, and we found a strong,

positive linear relationship (Figure S11; Pearson's correlation $r = 0.87$, $t(218) = 25.82$, p-value <

0.001). These results suggest the sampled range for most species adequately represented the total

range size. Nevertheless, we ran RF analyses using both total range size or sampling range size

and found similar results between the two variables (Figures 3, S12). Because the range size

variables were highly correlated with one another and captured similar aspects of the species'

biology, we subsequently describe the results using the sampling range size.

We found that sampling range size and lineage age were the best predictors of $\pi$ and were two of

the best predictors of IBD and IBE in random forest analyses (Figure 3). Random forest

regressions showed that taxonomic family was the fourth-best predictor of $\pi$ following mean

temperature (BIO1) considering the Increase in Mean Squared Error (%IncMSE) measure

(Figure 3a, S13). Validation of the RF regression results for $\pi$ based on the $R^2$ score indicated

good model performance ($R^2 = 0.95$). Mean precipitation (BIO12) and lineage age were other

important predictors of IBD (Figure 3b, S14). For IBE, RF showed that latitude was another

important predictor (Figure 3c, S15). In RF classification analyses, the best IBD and IBE models

were obtained when we downsampled our dataset. Our IBD RF model had high accuracy

(88.1%; 95% Confidence Interval CI: 77.82% – 94.7%), low sensitivity (56.25%), and very high

specificity (98.04%); our model was statistically better than predicting only the majority class, p-

value [Acc > NIR] = 0.0116. The IBE RF model had high accuracy (94.74%; CI: 85.38% –

98.9%), low sensitivity (57.14%), and very high specificity (100%); however, the model's

accuracy was not statistically better than the NIR, p-value [Acc > NIR] = 0.0688.

Sampling range size had a significant, positive relationship with $\pi$, showing that genetic diversity

within species increases with the geographic area sampled (Figure 4A; $R^2 = 0.09$; p-value <

0.001). We found a similar pattern using phylogenetic independent contrasts (PIC), where

sampling range size and $\pi$ had a strong, positive relationship (Figure 4B; $R^2 = 0.11$; p-value <

0.001). The relationship between lineage age and $\pi$ was positive and significant, in which older

species had higher genetic diversity in comparison with younger ones (Figure S16A; $R^2 = 0.11$,

p-value < 0.001), but this correlation was not significant taking the phylogeny into account

(Figure S16B; $R^2$ = 0.004, p-value: 0.405). We found a significant, positive relationship between
mean temperature (BIO1) and π, suggesting that species living in areas with higher temperatures
have more genetic diversity (Figure S17A). However, PIC did not show a significant correlation
between temperature (BIO1) and π after correcting for phylogenetic relatedness (Figures S17B).
430
The number of sequences associated with geographic coordinates, lineage age, and sampling
range size were top predictors for IBD and IBE models. There were clear differences between
the number of sequences sampled and IBD/IBE patterns (p-value < 0.01), in which species with
more sequences tended to show significant IBD/IBE patterns (Figure 5A, D). Species with
significant IBD/IBE had slightly larger sampling range sizes on average (Figure 5C, F) and also
tended to have older lineage ages (Figure 5E, S18). When temperate and tropical species were
considered separately, we found a similar trend in which species with larger ranges were more
likely to exhibit IBD and IBE, but results were not significant (Figure S19). Phylogenetic
generalized linear mixed models (PGLMM) found that the number of sequences had a
statistically significant, positive relationship with IBD (β = 0.01655, SE = 4.10 x $10^{-3}$, p <
0.001), and IBE (β = 0.0158626, SE = 4.27 × $10^{-3}$, p < 0.001). These results suggest that as
sampling effort increases, the likelihood of detecting an IBD or IBE pattern increases (Table S3).
Coefficients, standard errors, Z-scores, and p-values of the other predictors (i.e., sampling range
size and precipitation for IBD, and lineage age for IBE) are shown in Table S3.
445

**Discussion**

In this study, we analyzed 12,961 Cytb sequences from 220 global salamander species and identified key predictors of intraspecific diversity ($\pi$) and patterns of spatial genetic variation (IBD and IBE). Species range size and lineage age were important predictors of all three aspects of genetic variation. Specifically, species occupying larger geographic areas tended to have greater overall nucleotide diversity and to be more geographically and environmentally structured. The results for $\pi$ are consistent with population genetics theory (Wright, 1931), in that large geographic ranges can support larger populations that harbor more genetic diversity because of reduced genetic drift (Lacy, 1987). The significant relationship found between range size and spatial genetic differentiation (i.e., IBD and IBE) could be explained by the greater potential for geographic and environmental isolation between populations at distant locations in species with large ranges (Eckert et al., 2008). However, species range size alone does not explain much of the variation in IBD or IBE presence (small effect sizes recovered in logistic regression models), and the strong influence of the number of sequences sampled highlights one of the major challenges for macrogenetic studies—limited data availability for many species worldwide.

Our results provide only partial support for the clade-age hypothesis (e.g., random forest results). Although lineage age showed a positive and significant relationship with intraspecific genetic variation across global salamanders, there is no association when the phylogeny is included (see PIC and PGLMM results). For example, we tested whether older salamander lineages tend to accumulate more intraspecific genetic diversity than younger ones, the clade-age hypothesis (see McPeek and Brown, 2007), and we found that there is a significant non-phylogenetic correlation, but this relationship disappears once phylogenetic relatedness is taken into account. Similar patterns have been observed in other vertebrate groups, where evolutionary time is recovered as one of the best predictors of genetic diversity (e.g., Theodoridis et al., 2020 in mammals). Our results suggests that the observed effect of lineage age on genetic diversity could be explained by shared evolutionary history rather than independent accumulation of variation through time. The strong phylogenetic signal detected for lineage age (Pagel's $\lambda = 0.9999$, $p < 0.001$) indicates that this trait is almost entirely structured by shared ancestry. This high phylogenetic dependence

reinforces that comparisons involving lineage age must account for shared ancestry (PIC or

PGLMM models) to avoid inflated associations with other traits, such as $\pi$, IBD or IBE (e.g., Lai

et al., 2025). Thus, our results indicate that spatial (range size) and environmental factors

(precipitation, temperature) may play a stronger role than lineage age in shaping intraspecific

genetic variation across Caudata. This suggests that long-term ecological and demographic

stability, rather than lineage age, may better explain the maintenance of high intraspecific genetic

variation in some salamander lineages (e.g., Pan et al., 2019; Iannella et al., 2025). The

recovered relationship between lineage age and intraspecific genetic variation may be

confounded by lineage-level traits (e.g., life history traits, species range size) that are conserved

phylogenetically. In fact, we found a positive and significant relationship between lineage age

and sampling range size, as recently proposed by Alzate et al. (2025).

*Global patterns of nucleotide diversity ($\pi$)*

Our results provide evidence of the uneven distribution of nucleotide diversity ($\pi_S$) across the

globe. We observed that certain hotspots of genetic diversity, like the Ecuadorian Amazon and

central Europe, southwestern and central North America, and southeast Asia, align with regions

of high biodiversity (e.g., Mittermeier et al., 2011). Contrastingly, low genetic diversity in

regions such as northern North America or western Europe could reflect recent recolonizations

or historical bottlenecks (e.g., Riberon et al., 2002; Bonato et al., 2018; Auteri et al., 2022). Our

results only partially contrasted with those of Miraldo et al. (2016), who found that amphibian

genetic diversity decreased at higher latitudes. Although our map revealed a broadly similar

geographic pattern, latitude itself (mean, maximum, minimum, or midpoint) was not an

important predictor of genetic diversity in salamanders. Instead, the predictors that best

explained genetic diversity, such as species range size and environmental factors, tended to be

associated with regions closer to the equator. The high diversity we found in the Neotropics (16

species) could be explained by a combination of factors including tropical climate stability, high

topographic heterogeneity, or long evolutionary history in the Neotropics (Antonelli, 2022).

Another plausible explanation for the observed high levels of $\pi$ in some regions is that they

coincide with glacial refugia, areas where species persisted and maintained high levels of genetic

diversity through time (e.g., Ramírez⬜Barahona and Eguiarte, 2013). The Andean foothills were

a major glacial refugium, particularly the central Andes in the Neotropical region (e.g., Escobar

et al., 2021). In the Nearctic, the Southern Appalachians and coastal areas of the Pacific

Northwest are considered potential glacial refugia (e.g., Shafer et al., 2010). In the Palearctic, the

Caucasus Mountain range has been identified as a potential refugium during glacial periods (e.g.,

*Triturus* newts; Wielstra et al., 2013). The Korean Peninsula and the mountainous regions of

China are two examples of glacial refugia in the Oriental region (Fu and Wen, 2023). Low levels

of $\pi$ in northern North America are consistent with the impact of historical and demographic

processes in the region, such as Pleistocene glaciations and associated range contractions in

salamanders (Rovito and Schoville, 2017; López-Delgado and Meirmans, 2022). We showed that

species experiencing expansion, based on Tajima's D, presented low levels of nucleotide

diversity (Figure S20). This result is expected when recent expansion from a small population

occurred after a bottleneck, leading to a limited gene pool carried by a few founding individuals,

such as range expansions after the last glacial maximum (Mayr, 2001). Interestingly, the

occupation of refugia by salamander species may have contributed to the extinction of older

lineages and fostered more recent diversification events, consistent with patterns observed in the

regions mentioned above (see Tingley and Dubey, 2012). We observed that species experiencing

demographic expansion often have young evolutionary age (Figure S21), reflecting recent

diversification events associated with demographic growth.

*Global patterns of isolation-by-distance (IBD) and environment (IBE)*

The limited evidence for both IBD and IBE across species (only 21 species, or ~10%, showed

both patterns) suggests that gene flow in many salamander species may not be constrained by

geography or environment. Instead, the genetic variation in these species is shaped more by

demographic processes such as bottlenecks or population expansions than current spatial or

ecological gradients (e.g., Sexton et al., 2024). For instance, some salamander species may

maintain high gene flow that can homogenize genetic variation, reducing patterns of IBD or IBE.

In contrast, others with strong philopatry may exhibit spatial genetic structure influenced more

by historical processes that disrupt patterns of genetic isolation associated with IBD and IBE

(Wake, 2009; Sexton et al., 2014). For example, montane populations isolated in separate refugia

in response to climatic changes might differentiate and exhibit highly restricted gene flow even

at short distances or among similar environments, as reported in plethodontid salamanders

(Kozak and Wiens, 2006; Rovito, 2017).

These results underscore the importance of considering biogeographic and ecological context

when evaluating the drivers of intraspecific genetic variation on a global scale. The proportions

of species showing IBD and IBE differed slightly across biogeographic realms, with the

Palearctic including the most species with these patterns. One example is the fire salamander

(*Salamandra salamandra*), one of the most common and widespread salamander species in

Europe, which was found to be strongly structured in a landscape genetics analysis (Bani et al.,

2015). The Neotropical and Oriental regions had lower proportions of species with IBD or IBE

patterns compared to the other regions, but these differences were relatively minor. This contrast

between realms suggests that historical biogeography and environmental complexity are key to

understanding spatial genetic variation. For example, the incidence of IBD/IBE in the Nearctic

and Palearctic could reflect the consequence of glacial cycles (Schmitt, 2007), while the low

incidence in tropical regions may reflect more recent colonization dynamics that occurred in the

late Pleistocene (Carnaval et al., 2009).

*Predictors of genetic variation in global salamanders*

Two of the best predictors of nucleotide diversity in this study, the number of sequences and

range size, could both be considered proxies of demographic processes. Species with more

available sequences could truly be more abundant in nature, reflecting both a true demographic

property as well as sampling bias towards species that are more frequently encountered. Either

way, sampling more sequences could provide a greater likelihood of detecting genetic variation

because more samples analyzed will capture the full range of genetic diversity present within a

species. When we analyzed $\pi$, IBD, and IBE using a standardized sampling effort per species (10

to 20 sequences), we note that the number of sequences was not an important predictor in $\pi$ and

IBE based on the random forest models, but was the most important predictor of IBD. Similarly,

species sampling range size was not a key predictor for IBD and IBE but was an important

predictor for $\pi$ in this subsampled dataset (Figure S22). Lineage age was one of the best

predictors for $\pi$ and IBE in the subsampled dataset (Figure S22). These results suggest that

differences in sampling effort across species can provide challenges for macrogenetic studies and

should be considered carefully in future studies.

The positive correlations found between number of sequences, range size, and $\pi$ in our dataset

are consistent with the prediction that total population size (with number of sequences and range

size as potential proxies) corresponds with genetic diversity (Figure S23). Species with larger

geographic ranges can have larger and more balanced populations, reflecting the relationship

between effective population size and genetic variation (Waples, 2022). Geographic range size

was also one of the best predictors of IBD and IBE, like previous findings in a wide range of

taxonomic groups (Pelletier and Carstens, 2018). A large range can harbor more differentiation,

where population subdivision could be influenced by different ecological and evolutionary

pressures and undergo contrasting demographic histories (Lowe et al., 2017). The relationships

between range size and patterns of genetic variation we identified in global salamanders is

partially consistent with previous regional studies, such as Amador et al. (2024), which found a

similar pattern in Neotropical amphibians for $\pi$, but not for IBD or IBE. Similar to previous

meta-analyses of IBD and IBE (Crispo and Hendry, 2005; Jenkins et al. 2010), we found that

identifying global predictors of spatial genetic variation is challenging because of the many

potential factors that influence the presence and detection of these patterns.

Climatic variables, including different aspects of temperature and precipitation, were among the

top ten predictors of genetic variation in RF analyses, suggesting an important role of climate in

diversity and population differentiation within salamanders. Mean temperature (BIO1) was the

third best predictor for $\pi$. The association between higher genetic variation and warmer

temperatures is consistent with the evolutionary speed hypothesis (Wright et al., 2010; Gillman

and Wright, 2014), where genetic diversity is expected to increase with temperature. Variation in

precipitation (BIO12) was the second most important predictor for IBD after the number of

sequences. A possible explanation for this association is that current stable climates, such as

those in tropical environments, promote spatial genetic variation, thus supporting the climate

stability hypothesis (Janzen, 1967; Stevens, 1989). Although the relationships between these

climatic variables and the response variables we examined were relatively weak, previous work

has also highlighted the role of climate in explaining variation within salamanders. Specifically,

phylogeographic structure measured using species delimitation methods was best explained by

variation in climate (Parsons et al., 2024). Like other studies on birds (Smith et al., 2017) and

amphibians (Barrow et al., 2021), our analyses suggest that life history and ecological traits are

less important than geographic or climatic variation for predicting intraspecific diversity in

salamanders.

*Phylogenetic, taxonomic, and conservation implications*

We did not find phylogenetic signal in $\pi$, although taxonomic family was the fourth-best

predictor of $\pi$ in the final Percentage Increase in Mean Squared Error (%IncMSE) RF model,

suggesting that some variation can be explained by evolutionary history, but not all. Sample sizes

were limited within some families, but diversity estimates in those families were consistently

high (e.g., Sirenidae) or low (e.g., Cryptobranchidae), indicating family can be a useful predictor

of intraspecific variation. On the other hand, discrepancies in $\pi$ within genera of family

Plethodontidae, such as observed in *Plethodon* or *Eurycea*, emphasize the influence of local

ecological and recent demographic processes over shared ancestry for influencing variation

within species. Processes leading to homoplasy (e.g., convergence, parallel evolution) dilute

phylogenetic signal (Klingenberg and Gidaszewski, 2010). Homoplasy is a common

phenomenon in salamander evolution; for example, paedomorphosis has evolved independently

several times (Wake, 2009). Likewise, levels of genetic variation may be similar in salamander

species that do not share a recent common ancestor (Kozak and Wiens, 2006). Phylogenetic

signal in IBD was not statistically significant either, suggesting other factors independent of

phylogenetic history influence these patterns. This result was consistent with a previous study on

Neotropical amphibians (Amador et al. 2024), which found no phylogenetic signal in IBD or IBE.

Significant phylogenetic signal in nucleotide diversity, however, was previously found in

amphibian datasets including frogs (Barrow et al. 2021; Amador et al. 2024).

Species with higher $\pi$ values could contain cryptic diversity and would be useful target species

for exploring species boundaries in future studies (e.g., Parsons et al. 2022). Using a smaller

dataset than the present study (83 salamander species total, most from the COI mitochondrial

gene), Parsons et al. (2024) found that ~2/3 of species in the dataset showed strong

phylogeographic structure, determined by single locus molecular species delimitation methods.

Several species with high values of π in our dataset match species from their study, indicating

they have hidden genetic lineages that should be further explored (*Mertensiella caucasica,*

*Bolitoglossa rufescens, Plethodon cinereus, Eurycea cirrigera, Batrachoseps attenuatus,*

*Ommatotriton vittatus, Aneides flavipunctatus*), ideally incorporating nuclear genomic data and

thorough geographic sampling.

Salamanders are within one of the most threatened vertebrate groups (Luedtke et al., 2023) and

require innovative syntheses of available information to highlight priorities for conservation

strategies. Nucleotide diversity could be a useful proxy for conservation status (Petit-Marty,

2021). The relationship we identified between species range size and genetic variation highlights

the vulnerability of species with small ranges. Conservation status was an important predictor for

π, but sample sizes were limited within several categories. Seventy-seven (35%) of the 220

species in our dataset are threatened based on the IUCN Red List; 12 as Critical Endangered

(CR), 30 Endangered (EN), and 35 Vulnerable (VU). These salamanders should receive special

attention, especially if they occupy small ranges and have low genetic diversity.

*Limitations and challenges*

One major limitation for macrogenetic studies is the uneven spatial sampling of available

sequences, which can lead to an incomplete picture of the distribution of genetic diversity. There

are also sampling biases among taxa, such that some species are well characterized while others

are omitted from global comparisons. Another limitation is the use of a single mitochondrial

gene to calculate genetic diversity and genetic distances. Although mtDNA is widely used in

phylogenetic and population genetic studies because it evolves quickly (Galtier et al., 2009), it

provides an incomplete picture of intraspecific diversity because it is a single locus, haploid, and

solely maternally inherited (Moore, 1995). Moreover, although there is a potential association

between lineage age and genetic diversity, it is important to note that divergence times estimated

from different loci can have distinct coalescence histories and are often younger than species

ages. This suggests that a relationship between divergence times across loci and genetic variation

inferred from a single locus (mtDNA) is possible but not necessarily straightforward (Pie and

Caron, 2023). However, single locus macrogenetic studies are still valuable because the amount

and availability of data elucidate patterns associated with historical processes and variability

within species. Furthermore, the large genome size of salamanders hampers the use of nuclear

genomic data in studies on this scale. These limitations reinforce the need for additional

geographic sampling of high-quality tissues that include broader taxonomic coverage and

facilitate new genomic sequencing efforts.

The most important challenge for our study was the time required to retrieve the geographic

coordinates associated with each Cytb sequence. Most sequences in open access databases such

as GenBank do not have their latitude and longitude associated with them, which limits studies

of spatial genetic variation. Efforts such as phylogatR (Pelletier et al. 2022) aggregate genetic

sequences associated with georeferenced occurrences from global databases using automated

pipelines, but researchers providing the sequences must include the proper metadata when

uploading their data for this approach to work. We encourage scientists to share spatial data and

museum catalogue numbers when applicable along with sequences when they are published,

ensuring data reproducibility and extendibility.

*Conclusions*

Our study underscores the value of integrating spatial information into macrogenetic studies. Our

results support the hypotheses that variables associated with population size are positively

correlated with genetic variation and that salamanders tend to exhibit spatial patterns of genetic

variation that also differ across biogeographic realms. Based on our results, we reject the

hypothesis that traits associated with reproductive strategies predict mitochondrial genetic

variation in salamanders; and we partially support the hypothesis that older species tend to

accumulate higher generic variation, as explained above. Our results suggest the importance of

range-wide climatic variation, local environmental heterogeneity, and historical demographic

events for influencing genetic variation. Future amphibian macrogenetic and genomic studies can

incorporate more explicit tests of range expansion and contraction to better understand these

processes. Finally, our findings have broad implications for establishing conservation priorities

and highlighting regions where sampling has been historically scarce, particularly in biodiversity

hotspots that warrant further attention.

**Data and code availability statement**

The data and code (e.g., R scripts) that support the results of this research are openly available at

https://github.com/luchoamador/amphibian_macrogenetics. Large folders and other

supplementary materials are available in Dryad https://doi.org/10.5061/dryad.51c59zwjx.

**References**

Alzate, A., Rozzi, R., Velasco, J. A., Robertson, D. R., Zizka, A., Tobias, J. A., Hill, A., Bacon,

C. D., Janzen, T., Pellissier, L., Van Der Plas, F., Rosindell, J., & Onstein, R. E. (2025).

Evolutionary age correlates with range size across plants and animals. *Nature Communications*,

*16*(1), 7894. https://doi.org/10.1038/s41467-025-62124-y

Amador, L., Arroyo□Torres, I., & Barrow, L. N. (2024). Machine learning and phylogenetic

models identify predictors of genetic variation in Neotropical amphibians. *Journal of*

*Biogeography*, 51(5), 909–923. https://doi.org/10.1111/jbi.14795

AmphibiaWeb. (2024). <https://amphibiaweb.org> University of California, Berkeley, CA, USA.

Accessed 3 October 2024.

Antonelli, A. (2022). The rise and fall of Neotropical biodiversity. *Botanical Journal of the*

*Linnean Society*, 199(1), 8–24. https://doi.org/10.1093/botlinnean/boab061

Auteri, G. G., Marchán-Rivadeneira, M. R., Olson, D. H., & Knowles, L. L. (2022). Landscape

connectivity among coastal giant salamander (*Dicamptodon tenebrosus*) populations shows no

association with land use, fire frequency, or river drainage but exhibits genetic signatures of

potential conservation concern. *PLOS ONE*, *17*(6), e0268882.

https://doi.org/10.1371/journal.pone.0268882

Bani, L., Pisa, G., Luppi, M., Spilotros, G., Fabbri, E., Randi, E., & Orioli, V. (2015). Ecological

connectivity assessment in a strongly structured fire salamander (*Salamandra salamandra* )

population. *Ecology and Evolution*, *5*(16), 3472–3485. https://doi.org/10.1002/ece3.1617

Barrow, L. N., Masiero Da Fonseca, E., Thompson, C. E. P., & Carstens, B. C. (2021).

Predicting amphibian intraspecific diversity with machine learning: Challenges and prospects for

integrating traits, geography, and genetic data. *Molecular Ecology Resources*, 21(8), 2818–2831.

https://doi.org/10.1111/1755-0998.13303

Bivand R, Pebesma E, Gómez-Rubio V. (2013). *Applied spatial data analysis with R, Second*

*edition*. Springer, NY. https://asdar-book.org/.

Blanchet, S., Prunier, J. G., & De Kort, H. (2017). Time to Go Bigger: Emerging Patterns in

Macrogenetics. *Trends in Genetics*, *33*(9), 579–580. https://doi.org/10.1016/j.tig.2017.06.007

Blomberg, S. P., Garland, T., & Ives, A. R. (2003). Testing for phylogenetic signal in

comparative data: behavioral traits are more labile. *Evolution*, 57(4), 717–745.

Bonato, L., Corbetta, A., Giovine, G., Romanazzi, E., Šunje, E., Vernesi, C., & Crestanello, B.

(2018). Diversity among peripheral populations: Genetic and evolutionary differentiation of

*Salamandra atra* at the southern edge of the Alps. *Journal of Zoological Systematics and*

*Evolutionary Research*, *56*(4), 533–548. https://doi.org/10.1111/jzs.12224

Bonferroni, C. E. (1936) Teoria statistica delle classi e calcolo delle probabilita. Pubblicazioni

del R Istituto Superiore di Scienze Economiche e Commerciali di Firenze 8, 3-62, 1936.

Calenge C (2024). adehabitatHR: Home Range Estimation. R package version 0.4.22,

<https://CRAN.R-project.org/package=adehabitatHR>.

Cambon, J., Hernangómez, D., Belanger, C., & Possenriede, D. (2021). tidygeocoder: An R

package for geocoding. *Journal of Open Source Software*, *6*(65), 3544.

https://doi.org/10.21105/joss.03544

Carnaval, A. C., Hickerson, M. J., Haddad, C. F. B., Rodrigues, M. T., & Moritz, C. (2009).

Stability Predicts Genetic Diversity in the Brazilian Atlantic Forest Hotspot. *Science*, *323*(5915),

785–789. https://doi.org/10.1126/science.1166955

Corbett-Detig, R. B., Hartl, D. L., & Sackton, T. B. (2015). Natural Selection Constrains Neutral

Diversity across A Wide Range of Species. *PLOS Biology*, *13*(4), e1002112.

https://doi.org/10.1371/journal.pbio.1002112

Crispo, E., & Hendry, A. P. (2005). Does time since colonization influence isolation by distance?

A meta-analysis. *Conservation Genetics*, *6*(5), 665–682. https://doi.org/10.1007/s10592-005-

9026-4

De Kort, H., Prunier, J. G., Ducatez, S., Honnay, O., Baguette, M., Stevens, V. M., & Blanchet,

S. (2021). Life history, climate and biogeography interactively affect worldwide genetic diversity

of plant and animal populations. *Nature Communications*, *12*(1), 516.

https://doi.org/10.1038/s41467-021-20958-2

Denoël, M., Joly, P., & Whiteman, H. H. (2005). Evolutionary ecology of facultative

paedomorphosis in newts and salamanders. *Biological Reviews*, *80*(4), 663–671.

https://doi.org/10.1017/S1464793105006858

Dunnington, D. (2023). ggspatial: Spatial Data Framework for ggplot2. R package version 1.1.9,

<https://CRAN.R-project.org/package=ggspatial>.

Eckert, C. G., Samis, K. E., & Lougheed, S. C. (2008). Genetic variation across species'

geographical ranges: The central–marginal hypothesis and beyond. *Molecular Ecology*, *17*(5),

1170–1188. https://doi.org/10.1111/j.1365-294X.2007.03659.x

Edgar, R. C. (2004). MUSCLE: a multiple sequence alignment method with reduced time and

space complexity. *BMC Bioinformatics*, *5*(1), 113. https://doi.org/10.1186/1471-2105-5-113

Ellegren, H., & Galtier, N. (2016). Determinants of genetic diversity. *Nature Reviews Genetics*,

*17*(7), 422–433. https://doi.org/10.1038/nrg.2016.58

Escobar, S., Helmstetter, A. J., Jarvie, S., Montúfar, R., Balslev, H., & Couvreur, T. L. P. (2021).

Pleistocene climatic fluctuations promoted alternative evolutionary histories in *Phytelephas*

*aequatorialis* , an endemic palm from western Ecuador. *Journal of Biogeography*, *48*(5), 1023–

1037. https://doi.org/10.1111/jbi.14055

Felsenstein, J. (1985). Phylogenies and the Comparative Method. *The American Naturalist*,

*125*(1,), 1–15.

Fick, S.E. and R.J. Hijmans, (2017). WorldClim 2: new 1km spatial resolution climate surfaces

for global land areas. *International Journal of Climatology* 37 (12): 4302-4315.

Fonseca, E. M., Garda, A. A., Oliveira, E. F., Camurugi, F., Magalhães, F. D. M., Lanna, F. M.,

Zurano, J. P., Marques, R., Vences, M., & Gehara, M. (2021). The riverine thruway hypothesis:

Rivers as a key mediator of gene flow for the aquatic paradoxical frog *Pseudis tocantins* (Anura,

Hylidae). *Landscape Ecology*, *36*(10), 3049–3060. https://doi.org/10.1007/s10980-021-01257-z

Fonseca, E. M., Pelletier, T. A., Decker, S. K., Parsons, D. J., & Carstens, B. C. (2023).

Pleistocene glaciations caused the latitudinal gradient of within-species genetic diversity.

*Evolution Letters*, *7*(5), 331–338. https://doi.org/10.1093/evlett/qrad030

Frankham, R. (1996). Relationship of Genetic Variation to Population Size in Wildlife.

*Conservation Biology*, *10*(6), 1500–1508. https://doi.org/10.1046/j.1523-1739.1996.10061500.x

French, C. M., Bertola, L. D., Carnaval, A. C., Economo, E. P., Kass, J. M., Lohman, D. J.,

Marske, K. A., Meier, R., Overcast, I., Rominger, A. J., Staniczenko, P. P. A., & Hickerson, M. J.

(2023). Global determinants of insect mitochondrial genetic diversity. *Nature Communications*,

*14*(1), 5276. https://doi.org/10.1038/s41467-023-40936-0

Fu, J., & Wen, L. (2023). Impacts of Quaternary glaciation, geological history and geography on

animal species history in continental East Asia: A phylogeographic review. *Molecular Ecology*,

*32*(16), 4497–4514. https://doi.org/10.1111/mec.17053

Fusco, N. A., Pehek, E., & Munshi☐South, J. (2021). Urbanization reduces gene flow but not

genetic diversity of stream salamander populations in the New York City metropolitan area.

*Evolutionary Applications*, *14*(1), 99–116. https://doi.org/10.1111/eva.13025

Galtier, N., Nabholz, B., Glémin, S., & Hurst, G. D. D. (2009). Mitochondrial DNA as a marker

of molecular diversity: A reappraisal. *Molecular Ecology*, 18(22), 4541–4550.

https://doi.org/10.1111/j.1365-294X.2009.04380.x

Gillman, L. N., & Wright, S. D. (2014). Species richness and evolutionary speed: The influence

of temperature, water and area. *Journal of Biogeography*, *41*(1), 39–51.

https://doi.org/10.1111/jbi.12173

Hanson, J. O., Rhodes, J. R., Riginos, C., & Fuller, R. A. (2017). Environmental and geographic

variables are effective surrogates for genetic variation in conservation planning. *Proceedings of*

*the National Academy of Sciences*, *114*(48), 12755–12760.

https://doi.org/10.1073/pnas.1711009114

Hijmans R (2022). geosphere: Spherical Trigonometry. R package version 1.5-18,

<https://CRAN.R-project.org/package=geosphere>.

Hijmans RJ, Barbosa M, Ghosh A, Mandel A (2024). geodata: Download Geographic Data. R

package version 0.6-2, <https://CRAN.R-project.org/package=geodata>.

Hollister J, Shah T, Nowosad J, Robitaille A, Beck M, Johnson M (2025). *elevatr: Access*

*Elevation Data from Various APIs*. doi:10.5281/zenodo.8335450, R package version 0.99.1,

https://github.com/usepa/elevatr/.

Iannella, M., Cerasoli, F., Lunghi, E., Console, G., Biondi, M., & Sillero, N. (2025). Climate

Change Effects on the Only Western Palearctic Plethodontids: Range Changes and Possible

Depletion of Intraspecific Genetic Diversity. *Journal of Biogeography*, *52*(3), 686–698.

https://doi.org/10.1111/jbi.15064

Ives, A. R., & Helmus, M. R. (2011). Generalized linear mixed models for phylogenetic analyses

of community structure. *Ecological Monographs*, *81*(3), 511–525. https://doi.org/10.1890/10-

1264.1

Janzen, D. H. (1967). Why Mountain Passes are Higher in the Tropics. *The American Naturalist*,

*101*(919), 233–249. https://doi.org/10.1086/282487

Jenkins, D. G., Carey, M., Czerniewska, J., Fletcher, J., Hether, T., Jones, A., Knight, S., Knox,

J., Long, T., Mannino, M., McGuire, M., Riffle, A., Segelsky, S., Shappell, L., Sterner, A.,

822 Strickler, T., & Tursi, R. (2010). A meta□analysis of isolation by distance: Relic or reference

823 standard for landscape genetics? *Ecography*, *33*(2), 315–320. https://doi.org/10.1111/j.1600-

824 0587.2010.06285.x

825 Jetz, W., & Pyron, R. A. (2018). The interplay of past diversification and evolutionary isolation

826 with present imperilment across the amphibian tree of life. *Nature Ecology & Evolution*, *2*(5),

827 850–858. https://doi.org/10.1038/s41559-018-0515-5

828 Kardos, M., Armstrong, E. E., Fitzpatrick, S. W., Hauser, S., Hedrick, P. W., Miller, J. M.,

829 Tallmon, D. A., & Funk, W. C. (2021). The crucial role of genome-wide genetic variation in

830 conservation. *Proceedings of the National Academy of Sciences*, *118*(48), e2104642118.

831 https://doi.org/10.1073/pnas.2104642118

832 Kimura, M. (1968). Evolutionary rate at the molecular level. *Nature*, 217(5129), 624-626.

833 https://doi.org/10.1038/217624a0

834 Klingenberg, C. P., & Gidaszewski, N. A. (2010). Testing and Quantifying Phylogenetic Signals

835 and Homoplasy in Morphometric Data. *Systematic Biology*, 59(3), 245–261.

836 https://doi.org/10.1093/sysbio/syp106

837 Kozak, K. H., & Wiens, J. J. (2006). Does niche conservatism promote speciation? A case study

838 in North American salamanders. *Evolution*, *60*(12), 2604–2621. https://doi.org/10.1111/j.0014-

839 3820.2006.tb01893.x

840 Kuhn, M. (2008). "Building Predictive Models in R Using the caret Package." *Journal of*
841 *Statistical Software*, **28**(5), 1–26. doi:10.18637/jss.v028.i05,
842 https://www.jstatsoft.org/index.php/jss/article/view/v028i05.Lacy, R. C. (1987). Loss of Genetic
843 Diversity from Managed Populations: Interacting Effects of Drift, Mutation, Immigration,
844 Selection, and Population Subdivision. *Conservation Biology*, *1*(2), 143–158.

845 Lai, J., He, Y., Hou, M., Zhang, A., Wang, G., & Mao, L. (2025). Evaluating the relative

846 importance of phylogeny and predictors in phylogenetic generalized linear models using the

847 phylolm.hp R package. *Plant Diversity*, *47*(5), 709–717.

848 https://doi.org/10.1016/j.pld.2025.06.003Larsson, A. (2014). AliView: A fast and lightweight

849 alignment viewer and editor for large datasets. *Bioinformatics*, *30*(22), 3276–3278.

850 https://doi.org/10.1093/bioinformatics/btu531

Lawrence, E. R., Pedersen, E. J., & Fraser, D. J. (2023). Macrogenetics reveals multifaceted

influences of environmental variation on vertebrate population genetic diversity across the

Americas. *Molecular Ecology*, *32*(16), 4557–4569. https://doi.org/10.1111/mec.17059

Lee☐Yaw, J. A., & Irwin, D. E. (2012). Large geographic range size reflects a patchwork of

divergent lineages in the long☐toed salamander (*Ambystoma macrodactylum*). *Journal of*

*Evolutionary Biology*, *25*(11), 2276–2287. https://doi.org/10.1111/j.1420-9101.2012.02604.x

Leigh, D. M., Van Rees, C. B., Millette, K. L., Breed, M. F., Schmidt, C., Bertola, L. D., Hand,

B. K., Hunter, M. E., Jensen, E. L., Kershaw, F., Liggins, L., Luikart, G., Manel, S., Mergeay, J.,

Miller, J. M., Segelbacher, G., Hoban, S., & Paz-Vinas, I. (2021). Opportunities and challenges

of macrogenetic studies. *Nature Reviews Genetics*, *22*(12), 791–807.

https://doi.org/10.1038/s41576-021-00394-0

Li, D., Dinnage, R., Nell, L. A., Helmus, M. R., & Ives, A. R. (2020). phyr: An r package for

phylogenetic species☐distribution modelling in ecological communities. Methods in Ecology

and Evolution, 11(11), 1455–1463. https://doi.org/10.1111/2041-210X.13471

Liaw, A., Wiener, M. (2002). Classification and Regression by randomForest. *R News*, 2(3), 18-

22. <https://CRAN.R-project.org/doc/Rnews/>.

López☐Delgado, J., & Meirmans, P. G. (2022). History or demography? Determining the drivers

of genetic variation in North American plants. *Molecular Ecology*, 31(7), 1951–1962.

https://doi.org/10.1111/mec.16230

Lowe, W. H., Kovach, R. P., & Allendorf, F. W. (2017). Population Genetics and Demography

Unite Ecology and Evolution. *Trends in Ecology & Evolution*, *32*(2), 141–152.

https://doi.org/10.1016/j.tree.2016.12.002

Luedtke, J. A., Chanson, J., Neam, K., Hobin, L., Maciel, A. O., Catenazzi, A., Borzée, A.,

Hamidy, A., Aowphol, A., Jean, A., Sosa-Bartuano, Á., Fong G., A., De Silva, A., Fouquet, A.,

Angulo, A., Kidov, A. A., Muñoz Saravia, A., Diesmos, A. C., Tominaga, A., … Stuart, S. N.

(2023). Ongoing declines for the world's amphibians in the face of emerging threats. *Nature*,

*622*(7982), 308–314. https://doi.org/10.1038/s41586-023-06578-4

Malcher, G., Amorim, A.L., Ferreira, P. *et al.* (2023). First evaluation of the population genetics

and aspects of the evolutionary history of the Amazonian snook, *Centropomus irae*, and its

association with the Amazon plume. *Hydrobiologia* 850, 2115–2125.

https://doi.org/10.1007/s10750-023-05223-5

Mayr, E. (2001). *What Evolution Is*. New York, NY: Basic Books.

McPeek, M. A., & Brown, J. M. (2007). Clade Age and Not Diversification Rate Explains

Species Richness among Animal Taxa. *The American Naturalist*, *169*(4), E97–E106.

https://doi.org/10.1086/512135

Meng, H., Li, X., & Qiao, P. (2014). Population Structure, Historical Biogeography and

Demographic History of the Alpine Toad *Scutiger ningshanensis* in the Tsinling Mountains of

Central China. *PLoS ONE*, *9*(6), e100729. https://doi.org/10.1371/journal.pone.0100729

Millette, K. L., Fugère, V., Debyser, C., Greiner, A., Chain, F. J. J., & Gonzalez, A. (2021).

Refining analyses of existing data sets is valuable for macrogenetics: A response to Paz☐Vinas,

Jensen et al., (2021). *Ecology Letters*, *24*(6), 1285–1286. https://doi.org/10.1111/ele.13733

Liedtke, H.C., Wiens, J.J. & Gomez-Mestre, I. (2022). The evolution of reproductive modes and

life cycles in amphibians. Nat Commun 13, 7039. https://doi.org/10.1038/s41467-022-34474-4

Miraldo, A., Li, S., Borregaard, M. K., Flórez-Rodríguez, A., Gopalakrishnan, S., Rizvanovic,

M., Wang, Z., Rahbek, C., Marske, K. A., & Nogués-Bravo, D. (2016). An Anthropocene map of

genetic diversity. *Science*, 353(6307), 1532–1535. https://doi.org/10.1126/science.aaf4381

Mittermeier, R. A., Turner, W. R., Larsen, F. W., Brooks, T. M., & Gascon, C. (2011). Global

Biodiversity Conservation: The Critical Role of Hotspots. In F. E. Zachos & J. C. Habel (Eds.),

*Biodiversity Hotspots* (pp. 3–22). Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-642-

20992-5_1

Moore, W. S. (1995). Inferring phylogenies from mtDNA variation: mitochondrial-gene trees

versus nuclear-gene trees. Evolution, 49(4), 718-726.

Oksanen J, Simpson G, Blanchet F, Kindt R, Legendre P, Minchin P, O'Hara R, Solymos P,

Stevens M, Szoecs E, Wagner H, Barbour M, Bedward M, Bolker B, Borcard D, Carvalho G,

Chirico M, De Caceres M, Durand S, Evangelista H, FitzJohn R, Friendly M, Furneaux B,

Hannigan G, Hill M, Lahti L, McGlinn D, Ouellette M, Ribeiro Cunha E, Smith T, Stier A, Ter

Braak C, Weedon J (2024). vegan: Community Ecology Package. R package version 2.6-8,

<https://CRAN.R-project.org/package=vegan>

Oliveira, B. F., São-Pedro, V. A., Santos-Barrera, G., Penone, C., & Costa, G. C. (2017).

AmphiBIO, a global database for amphibian ecological traits. *Scientific Data*, 4(1), 170123.

https://doi.org/10.1038/sdata.2017.123

Pagel, M. (1999). Inferring the historical patterns of biological evolution. *Nature*, *401*(6756),

877–884. https://doi.org/10.1038/44766

Pan, T., Wang, H., Orozcoterwengel, P., Hu, C.-C., Wu, G.-Y., Qian, L.-F., Sun, Z.-L., Shi, W.-

B., Yan, P., Wu, X.-B., & Zhang, B.-W. (2019). Long-term sky islands generate highly divergent

lineages of a narrowly distributed stream salamander (Pachyhynobius shangchengensis) in mid-

latitude mountains of East Asia. *BMC Evolutionary Biology*, *19*(1), 1.

https://doi.org/10.1186/s12862-018-1333-8

Paradis, E. (2010). pegas: An R package for population genetics with an integrated–modular

approach. *Bioinformatics*, *26*(3), 419–420. https://doi.org/10.1093/bioinformatics/btp696

Paradis, E., & Schliep, K. (2019). ape 5.0: An environment for modern phylogenetics and

evolutionary analyses in R. *Bioinformatics*, 35(3), 526–528.

https://doi.org/10.1093/bioinformatics/bty633

Parsons, D. J., Pelletier, T. A., Wieringa, J. G., Duckett, D. J., & Carstens, B. C. (2022). Analysis

of biodiversity data suggests that mammal species are hidden in predictable places. *Proceedings*

*of the National Academy of Sciences*, *119*(14), e2103400119.

https://doi.org/10.1073/pnas.2103400119

Parsons, D. J., Green, A. E., Carstens, B. C., & Pelletier, T. A. (2024). Predicting genetic

biodiversity in salamanders using geographic, climatic, and life history traits. *PLOS ONE*, *19*(10),

e0310932. https://doi.org/10.1371/journal.pone.0310932

Park, K. Y., Lucas, M., Chaulk, A., Matter, S. F., Roland, J., & Keyghobadi, N. (2024).

Immigration allows population persistence and maintains genetic diversity despite an attempted

experimental extinction. *Royal Society Open Science*, *11*(7), 240557.

https://doi.org/10.1098/rsos.240557

Pauls, S. U., Nowak, C., Bálint, M., & Pfenninger, M. (2013). The impact of global climate

change on genetic diversity within populations and species. *Molecular Ecology*, *22*(4), 925–946.

https://doi.org/10.1111/mec.12152

Paz, A., Ibáñez, R., Lips, K.R. and Crawford, A.J. (2015), Testing the role of ecology and life

history in structuring genetic variation across a landscape: a trait-based phylogeographic

approach. *Molecular Ecology*, 24: 3723-3737. https://doi-

org.libproxy.unm.edu/10.1111/mec.13275

PazVinas, I., Jensen, E. L., Bertola, L. D., Breed, M. F., Hand, B. K., Hunter, M. E., Kershaw,

F., Leigh, D. M., Luikart, G., Mergeay, J., Miller, J. M., Van Rees, C. B., Segelbacher, G., &

Hoban, S. (2021). Macrogenetic studies must not ignore limitations of genetic markers and scale.

*Ecology Letters*, *24*(6), 1282–1284. https://doi.org/10.1111/ele.13732

Pebesma, E., 2018. Simple Features for R: Standardized Support for Spatial Vector Data. The R

Journal 10 (1), 439-446, https://doi.org/10.32614/RJ-2018-009

Pebesma, E., & Bivand, R. (2023). Spatial Data Science: With Applications in R. Chapman and

Hall/CRC. https://doi.org/10.1201/9780429459016

Pelletier, T. A., & Carstens, B. C. (2018). Geographical range size and latitude predict

population genetic structure in a global survey. *Biology Letters*, 14(1), 20170566.

https://doi.org/10.1098/rsbl.2017.0566

Pelletier, T. A., Parsons, D. J., Decker, S. K., Crouch, S., Franz, E., Ohrstrom, J., & Carstens, B.

C. (2022). phylogatR: Phylogeographic data aggregation and repurposing. *Molecular Ecology*

*Resources*, 22(8), 2830–2842. https://doi.org/10.1111/1755-0998.13673

PetitMarty, N., VázquezLuis, M., & Hendriks, I. E. (2021). Use of the nucleotide diversity in

COI mitochondrial gene as an early diagnostic of conservation status of animal species.

*Conservation Letters*, *14*(1), e12756. https://doi.org/10.1111/conl.12756

Pie, M. R., & Caron, F. S. (2023). Substantial variation in species ages among vertebrate clades.

*Evolutionary Journal of the Linnean Society*, *2*(1), kzad006.

https://doi.org/10.1093/evolinnean/kzad006

R Core Team (2023). R: A Language and Environment for Statistical Computing. R Foundation

for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.

RamírezBarahona, S., & Eguiarte, L. E. (2013). The role of glacial cycles in promoting genetic

diversity in the Neotropics: The case of cloud forests during the Last Glacial Maximum. *Ecology*

*and Evolution*, *3*(3), 725–738. https://doi.org/10.1002/ece3.483

Reed, D. H., & Frankham, R. (2003). Correlation between Fitness and Genetic Diversity.

*Conservation Biology*, *17*(1), 230–237. https://doi.org/10.1046/j.1523-1739.2003.01236.x

Revell, L. J. (2012). phytools: An R package for phylogenetic comparative biology (and other

things): *phytools: R package*. *Methods in Ecology and Evolution*, *3*(2), 217–223.

https://doi.org/10.1111/j.2041-210X.2011.00169.x

Riberon, A., Sotiriou, E., Miaud, C., Andreone, F., & Taberlet, P. (2002). Lack of genetic

diversity in *Salamandra lanzai* revealed by Cytochrome b gene sequences. *Copeia*, *2002*(1),

229–232.

Rios, N. E. & Bart, H. L. (2010). GEOLocate (Version 3.22). [Computer software]. Belle Chasse,

LA: Tulane University Museum of Natural History.

Román□Palacios, C. (2023). The phruta r package: Increasing access, reproducibility and

transparency in phylogenetic analyses. *Methods in Ecology and Evolution*, 14(9), 2284–2299.

https://doi.org/10.1111/2041-210X.14147

Rovito, S. M. (2017). The Geography of Speciation in Neotropical Salamanders. *Herpetologica*,

*73*(3), 229–241. https://doi.org/10.1655/HERPETOLOGICA-D-16-00077.1

Rovito, S. M., & Schoville, S. D. (2017). Testing models of refugial isolation, colonization and

population connectivity in two species of montane salamanders. *Heredity*, *119*(4), 265–274.

https://doi.org/10.1038/hdy.2017.31

RS-eco (2023). rasterSp: R Package to rasterize and summarise IUCN range maps. R package

version 0.0.1, <https://github.com/RS-eco/rasterSp>.

Schierenbeck, K. A. (2017). Population-level genetic variation and climate change in a

biodiversity hotspot. *Annals of Botany*, *119*(2), 215–228. https://doi.org/10.1093/aob/mcw214

Schmitt, T. (2007). Molecular biogeography of Europe: Pleistocene cycles and postglacial trends.

*Frontiers in Zoology*, *4*(1), 11. https://doi.org/10.1186/1742-9994-4-11

Scholl, J. P., & Wiens, J. J. (2016). Diversification rates and species richness across the Tree of

Life. *Proceedings of the Royal Society B: Biological Sciences*, *283*(1838), 20161334.

https://doi.org/10.1098/rspb.2016.1334

Segovia□Ramírez, M. G., Ramírez□Sánchez, O., Decena Segarra, L. P., Rios□Carlos, H., &

Rovito, S. M. (2023). Determinants of genetic diversity in Neotropical salamanders

(Plethodontidae: Bolitoglossini). *Ecology and Evolution*, *13*(11), e10707.

https://doi.org/10.1002/ece3.10707

Sexton, J. P., Hangartner, S. B., & Hoffmann, A. A. (2014). Genetic isolation by environment or

distance: which pattern of gene flow is most common? Special Section. *Evolution*, *68*(1), 1–15.

https://doi.org/10.1111/evo.12258

Sexton, J. P., Clemens, M., Bell, N., Hall, J., Fyfe, V., & Hoffmann, A. A. (2024). Patterns and

effects of gene flow on adaptation across spatial scales: Implications for management. *Journal of*

*Evolutionary Biology*, *37*(6), 732–745. https://doi.org/10.1093/jeb/voae064

Shafer, A. B. A., Cullingham, C. I., Côté, S. D., & Coltman, D. W. (2010). Of glaciers and

refugia: A decade of study sheds new light on the phylogeography of northwestern North

America. *Molecular Ecology*, *19*(21), 4589–4621. https://doi.org/10.1111/j.1365-

294X.2010.04828.x

Smith, B. T., Seeholzer, G. F., Harvey, M. G., Cuervo, A. M., & Brumfield, R. T. (2017). A

latitudinal phylogeographic diversity gradient in birds. *PLOS Biology*, *15*(4), e2001073.

https://doi.org/10.1371/journal.pbio.2001073

Stevens, G. C. (1989). The Latitudinal Gradient in Geographical Range: How so Many Species

Coexist in the Tropics. *The American Naturalist*, *133*(2), 240–256.

https://doi.org/10.1086/284913

Stewart, A. A., & Wiens, J. J. (2025). A time-calibrated salamander phylogeny including 765

species and 503 genes. *Molecular Phylogenetics and Evolution*, *204*, 108272.

https://doi.org/10.1016/j.ympev.2024.108272

Tajima, F. (1989). Statistical methods to test for nucleotide mutation hypothesis by DNA

polymorphism. *Genetics* 123:585-595.

Theodoridis, S., Fordham, D. A., Brown, S. C., Li, S., Rahbek, C., & Nogues-Bravo, D. (2020).

Evolutionary history and past climate change shape the distribution of genetic diversity in

terrestrial mammals. *Nature Communications*, *11*(1), 2557. https://doi.org/10.1038/s41467-020-

16449-5

Tingley, R., & Dubey, S. (2012). Disparity in the timing of vertebrate diversification events

between the northern and southern hemispheres. *BMC Evolutionary Biology*, *12*(1), 244.

https://doi.org/10.1186/1471-2148-12-244

Wake, D. B. (2009). What Salamanders Have Taught Us About Evolution. *Annual Review of*

*Ecology, Evolution, and Systematics*, *40*(1), 333–352.

https://doi.org/10.1146/annurev.ecolsys.39.110707.173552

Wang, I. J. (2013). Examining the full effects of landscape heterogeneity on spatial genetic

variation: a multiple matrix regression approach for quantifying geographic and ecological

isolation: special section. *Evolution*, *67*(12), 3403–3411. https://doi.org/10.1111/evo.12134

Wang, I. J., & Bradburd, G. S. (2014). Isolation by environment. *Molecular Ecology*, *23*(23),

5649–5662. https://doi.org/10.1111/mec.12938

Waples, R. S. (2022). What is $N_e$, anyway? *Journal of Heredity*, 113(4), 371–379.

https://doi.org/10.1093/jhered/esac023

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

Wielstra, B., Crnobrnja-Isailović, J., Litvinchuk, S. N., Reijnen, B. T., Skidmore, A. K.,

Sotiropoulos, K., Toxopeus, A. G., Tzankov, N., Vukov, T., & Arntzen, J. W. (2013). Tracing

glacial refugia of *Triturus* newts based on mitochondrial DNA phylogeography and species

distribution modeling. *Frontiers in Zoology*, *10*(1), 13. https://doi.org/10.1186/1742-9994-10-13

Wright, S. (1931). Evolution in Mendelian populations. *Genetics*, *16*(2), 97–159.

https://doi.org/10.1093/genetics/16.2.97

Wright, S. (1943). Isolation by distance. *Genetics*, *28*(2), 114–138.

https://doi.org/10.1093/genetics/28.2.114

Wright, S. D., Gillman, L. N., Ross, H. A., & Keeling, D. J. (2010). Energy and the tempo of

evolution in amphibians. *Global Ecology and Biogeography*, *19*(5), 733–740.

https://doi.org/10.1111/j.1466-8238.2010.00549.x

Yang, Z. (2014). Molecular evolution: a statistical approach. Oxford University Press, Oxford.

Zamudio, K. R. & Wieczorek, A. M. (2007). Fine-scale spatial genetic structure and dispersal

among spotted salamander (*Ambystoma maculatum*) breeding populations. *Molecular Ecology*,

16: 257-274. https://doi-org.libproxy.unm.edu/10.1111/j.1365-294X.2006.03139.x

 **Figures**

 **Figure 1.** Maps of A) mean nucleotide diversity ($\pi$) and B) mean number of sequences with

 coordinates (log scale) of global salamanders. Biogeographic realms are indicated with shading:

 Nearctic (brown), Neotropic (green), Oriental (blue), and Palearctic (purple). Boxplots of

 nucleotide diversity for each biogeographic realm are shown in map A. The number of species

 and mean number of sequences with coordinates for each biogeographic realm are shown in map

 B. For visualization purposes, we show grid cell = 4° (approximately 440 km$^2$) for mapping.

 **Figure 2.** Phylogeny of Caudata species included in this study, showing the values of nucleotide

 diversity ($\pi$) as gray barplots to the right and the presence (brown) or absence (teal) of isolation-

 by-distance (IBD) and isolation-by-environment (IBE) per species. Relationships are based on

 Jetz et al. (2018). *Notophthalmus viridescens* silhouette was taken from phylopic.org.

 **Figure 3.** Random Forest (RF) results for A) nucleotide diversity ($\pi$, using RF regression), B)

 isolation-by-distance (IBD, using RF classification), and C) isolation-by-environment (IBE,

 using RF classification). All predictors except the number of base pairs, and the total range size

 were used in these RF models, but only the top 10 best predictors for each model are shown here.

 Range size* = sampling range size based on localities with associated DNA sequences; # of

 sequences[+] = number of sequences associated with geographic coordinates; SVL = snout-vent

 length; mm = millimeters; BIO1 = Annual Mean Temperature; BIO4 = Temperature Seasonality

 (standard deviation ×100); BIO12 = Annual Precipitation; BIO15 = Precipitation Seasonality

 (Coefficient of variation).

 **Figure 4.** A) Comparison of sampling range size (in log scale) with nucleotide diversity ($\pi$) for

 each species. B) Comparison of phylogenetic independent contrast (PIC) results between the

 sampling range size (log scale) and nucleotide diversity ($\pi$). Linear regression results are

 indicated in each panel.

 **Figure 5.** Boxplots comparing the number of sequences (A, D) and the sampling range size (C,

 F) between species with (Yes, brown color) and without (No, teal color) isolation-by-distance

1084 (IBD) and isolation by environment (IBE). Comparisons of precipitation (BIO12) with IBD (B),

1085 and lineage age with IBE (E) are also shown. Each panel includes the results of a chi-squared test.

1086 Black dots represent outliers.

1087

1088

1089

**Figure 1.** Maps of A) mean nucleotide diversity (π) and B) mean number of sequences with

coordinates (log scale) of global salamanders. Biogeographic realms are indicated with shading:

Nearctic (brown), Neotropic (green), Oriental (blue), and Palearctic (purple). Boxplots of

nucleotide diversity for each biogeographic realm are shown in map A. The number of species

and mean number of sequences with coordinates for each biogeographic realm are shown in map

B. For visualization purposes, we show grid cell = 4° (approximately 440 km$^2$) for mapping.

1096

**Figure 2.** Phylogeny of Caudata species included in this study, showing the values of nucleotide

diversity (π) as gray barplots to the right and the presence (brown) or absence (teal) of isolation-

by-distance (IBD) and isolation-by-environment (IBE) per species. Relationships are based on

Jetz et al. (2018). *Notophthalmus viridescens* silhouette was taken from phylopic.org.

1101

**Figure 3.** Random Forest (RF) results for A) nucleotide diversity (π, using RF regression), B) isolation-by-distance (IBD, using RF classification), and C) isolation-by-environment (IBE, using RF classification). All predictors except the number of base pairs and the total range size were used in these RF models, but only the top 10 best predictors for each model are shown here. Range size* = sampling range size based on localities with associated DNA sequences; # of sequences[+] = number of sequences associated with geographic coordinates; SVL = snout-vent length; mm = millimeters; BIO1 = Annual Mean Temperature; BIO4 = Temperature Seasonality (standard deviation ×100); BIO12 = Annual Precipitation; BIO15 = Precipitation Seasonality (Coefficient of variation).

1113

**Figure 4.** A) Comparison of sampling range size (in log scale) with nucleotide diversity ($\pi$) for

each species. B) Comparison of phylogenetic independent contrast (PIC) results between the

sampling range size (log scale) and nucleotide diversity ($\pi$). Linear regression results are

indicated in each panel.

1119
1120 **Figure 5.** Boxplots comparing the number of sequences (A, D) and the sampling range size (C,

1121 F) between species with (Yes, brown color) and without (No, teal color) isolation-by-distance

1122 (IBD) and isolation by environment (IBE). Comparisons of precipitation with IBD (B), and

1123 lineage age with IBE (E) are also shown. Each panel includes the results of a chi-squared test.

1124 Black dots represent outliers.