Customized feminine hygiene wash containing postbiotics from *Lactobacillus* spp. to

treat Urinary Tract Infections (UTI).

Veena G Nair [a], David Raj Chellappan[b], Remya Devi Durai[c], Y.B.R.D Rajesh[d], Dhiviya

Narbhavi [e], Anupriya A[f], Prabhusaran N [g], Saisubramanian Nagarajan [a*]

* Corresponding Author: Dr. Sai Subramanian N, Antimicrobial Resistance Lab, Centre

for Research in Infectious diseases, School of Chemical and Biotechnology, SASTRA

Deemed to be University, Thanjavur 613401, Tamil Nadu, India. Phone: +91 4362

2641011

email: sai@scbt.sastra.edu

[a] Antimicrobial Resistance Lab, Centre for Research in Infectious diseases, School of

Chemical and Biotechnology, SASTRA Deemed to be University, Thanjavur 613401,

Tamil Nadu, India.

[b] Central Animal House Facility, SASTRA Deemed to be University, Thanjavur, 613401,

India

[c] Pharmaceutical Technology Laboratory, School of Chemical and Biotechnology,

SASTRA Deemed to be University, Thanjavur, 613401, India

[d] Department of Chemistry, School of Chemical and Biotechnology, SASTRA Deemed

University, Thanjavur – 613 401, Tamil Nadu, India.

[e] Assistant Professor, Department of Obstetrics and Gynaecology, TSRMMCH&RC,

Tiruchirappalli, Tamil Nadu, India

[f] Associate Professor, Department of Microbiology, TSRMMCH&RC, Tiruchirappalli,

Tamil Nadu, India

^g **Research Faculty, Institutional Research Board TSRMMCH&RC, Tiruchirappalli,**

Tamil Nadu, India

Abstract

The rising prevalence of antimicrobial resistance has intensified the search for innovative

therapeutic strategies, particularly in the prevention of urinary tract infections (UTIs). This

study presents the development and evaluation of a novel postbiotics vaginal wash

formulated to prevent UTIs by utilizing metabolites derived from indigenous vaginal

Lactobacillus spp. The primary objective was to create a cost-effective, stable, and non-

invasive solution targeting uropathogenic bacteria. Key metabolites, including tryptamine, (-

)-terpinen-4-ol, and itaconic anhydride, were identified from cell free supernatant of vaginal

Lactobacillus and incorporated into a poloxamer 407-based formulation. *In vitro* assays

demonstrated significant bioactivity against uropathogenic bacteria, effectively inhibiting

bacterial colonization and biofilm formation. Preclinical validation was conducted using

BALB/c mice models to assess both the safety and efficacy of the vaginal wash. Results

indicated a substantial reduction in infection rates among treated mice, with no observed

adverse effects, confirming the formulation's safety profile. In conclusion, this novel

postbiotics vaginal wash represents a promising non-invasive therapeutic approach for UTI

prevention. By harnessing bioactive metabolites from human vaginal Lactobacillus spp., the

formulation offers a potential solution to combat antimicrobial resistance while improving

women's health outcomes. Further clinical studies are warranted to validate these findings

and explore broader applications in clinical practice, paving the way for new strategies in

managing UTIs and enhancing overall female health.

Introduction

The growing issue of antibiotic resistance on a global scale is a pressing concern, significantly compromising the effectiveness of commonly used antibiotics against widespread bacterial infections. The 2022 Global Antimicrobial Resistance and Use Surveillance System (GLASS) report, also identifies extremely high levels of resistance in common pathogens. The global rapid situation assessment based on median and updated country-level reported rates shows that in 76 countries the projected resistance burden on third generation cephalosporins is as high as 42% rate of resistance in *E. coli*, and for methicillin-resistant *Staphylococcus aureus* it is up to 35%. Particularly for UTIs linked to *E. coli*, disturbingly one in five cases were less susceptible to conventional antibiotics such as ampicillin, co-trimoxazole and fluoroquinolones in 2020(1). This worrisome trend is making it increasingly challenging to effectively combat and treat common infections, emphasizing the urgent need for global strategies to address antibiotic resistance.

Over the recent years, there has been a noticeable surge in the pharmaceutical industry's interest in developing new formulations that incorporate beneficial microorganisms, often tailored to the specific needs of the host.(2) Nevertheless, the heightened focus on the clinical use of probiotics necessitates consideration when administering them to populations with health challenges. Recently, the Food and Drug Administration created a new category called "live biotherapeutic products" (LBP). As a result, the documentation and demonstration of quality, safety and efficacy for new products (including LBPs) are adapted to specific properties such as risk profiles stemming from both target population characteristics along with those related more directly to product strains or components. This approach enables a comprehensive assessment of the global benefit-risk ratio concerning their intended use.(3)

The pharmacological aspects of probiotics are more intricate compared to conventional drugs

due to their multifaceted impact on host cells, mucosal organs, and the entire organism.

Evaluating their effects is challenging as they operate through diverse mechanisms, either

individually or synergistically(4). The reports describe different ways vaginal lactobacilli can

exert their functions as follows: (a) producing antimicrobial substances such as organic acids,

hydrogen peroxide, bacteriocins and biosurfactants; or enzymes like arginine deaminase; (b)

adhesion to epithelial cells, mucus components or extracellular matrix for colonization and

domination of this ecological niche environment by autoaggregation co-aggregations,

biofilms in the vagina mucosa, competitive exclusion competition for nutrients and

immunomodulation.(5, 6)

Although probiotics are generally considered safe, concerns persist about the potential risks

associated with administering live bacteria in specific situations, such as during pregnancy or

while preparing for motherhood(7, 8). Additionally, ensuring the stability of probiotic

products—so that they maintain enough viable bacteria during handling, transportation, and

storage—presents practical challenges in industrial production and marketing(9). As a result

of these challenges, there has been considerable research focus on postbiotics, inactivated

probiotics, and the products formed during the breakdown of probiotic bacteria. This research

is increasingly demonstrating their potential to promote health and treat diseases(10).

Postbiotics were defined as "a preparation of inanimate microorganisms and/or their

components that confers a health benefit on the host" by The International Scientific

Association for Probiotics and Prebiotics (ISAPP) published its consensus statement

regarding Postscript in 2021 (9). In recent years, there has been increasing research focus on

the significant effects of *Lactobacillus* bioactive compounds (postbiotics) on vaginal health

(10–13). One of the most important biological activities for postbiotics is to reduce vaginal

pH to 3.8–4, which has antimicrobial effects resulting in significant inhibition of pathogenic

microorganisms(14–16). Research indicates that vaginal *Lactobacilli* and the bioactive

compounds they produce, including hydrogen peroxide, lactic acid, surfactants, and

ribosomally synthesized antimicrobial peptides (bacteriocins), are critical in suppressing the

growth and proliferation of vaginal pathogens(17, 18). Most likely due to bacteriocins, cell-

free supernatants exerted the highest scale of antibacterial activity (19). For instance, Cell

free supernatants from *Lactobacillus* and *Bifidobacterium* genera have demonstrated

antibacterial properties against enteroinvasive *E. coli*(20). Exopolysaccharides (EPS) from

Bifidobacterium bifidum are used to promote the growth of *Lactobacilli* and other anaerobes

& inhibit *Enterococci*, *Enterobacteriacea*, and *B. fragilis* (21). Furthermore, the antibacterial

metabolite reuterin, produced by *Lactobacillus reuteri*, is believed to act by oxidizing thiol

groups in pathogenic gut bacteria(22).

While most studies on postbiotics highlight common metabolites such as lactic acid, our

earlier works have identified unique bioactive compounds from human vaginal *Lactobacillus*

with distinct therapeutic properties. These metabolites include tryptamine, a biogenic

amine(23) ; (-)-terpinen-4-ol, a naturally occurring monoterpene (Manuscript under review);

and itaconic anhydride, the cyclic form of itaconic acid (Manuscript under Review). Unlike

typical postbiotics, these compounds exhibit specific bioactivities such as antibiofilm, efflux

inhibitory, and antipersister activities.

Tryptamine disrupts the exopolysaccharide matrix of bacterial biofilms, enhancing their

susceptibility to antimicrobial agents. (-)-Terpinen-4-ol demonstrates significant efflux

inhibitory activity, counteracting the mechanisms employed by uropathogenic bacteria to

expel antibiotics. Itaconic anhydride works synergistically with (-)-terpinen-4-ol to mitigate

the virulence of persister cells in uropathogenic *E. coli*, providing a robust defense against

persistent bacterial infections. Leveraging these findings, we have developed a customized

vaginal wash formulation incorporating these bioactive postbiotics with a poloxamer 407

base, a biocompatible polymer known for its stability and suitability for mucosal applications. Preclinical validation conducted on BALB/c mice models has demonstrated the efficacy and safety of this innovative formulation, underscoring its potential for clinical translation. This novel formulation represents a significant advancement in UTI prevention, addressing the urgent need for new therapies that can combat antimicrobial resistance and improve women's health.

Materials and Methods

In vitro **cell line infection and toxicity study**

T24 bladder epithelial cells and RAW macrophages (NCCS, Pune were utilized to assess the efficacy and cytotoxicity of various treatments targeting bacterial infections. T24 cells, derived from transitional bladder carcinoma, were chosen for their close resemblance to primary human bladder epithelial cells, making them an ideal model for studying urinary tract infections (UTIs). The cells were grown in McCoy's 5A medium, enhanced with 10% fetal bovine serum (FBS), and plated in a 24-well plate at a density of 1×10^5 cells/mL. To provide optimal conditions for growth, the T24 cells were incubated for 16 hours at 37°C in a 5% CO_2/95% air atmosphere.

Following a 2-hour incubation period at 37°C with 5% CO_2, the bacterial culture was prepared at a multiplicity of infection (MOI) of one and added to 24-well plates. Treatments were administered for 24 hours, after which cells were rinsed with sterile PBS to eliminate unbound bacteria. Cells were then trypsinized to detach and recover adhered bacteria, and bacterial counts were determined via serial dilution on Luria-Bertani agar plates. Each assay was performed in triplicate and repeated twice for validation.

For cytotoxicity assessments on RAW macrophages, the MTT assay was used. Cells grown

to 70–80% confluence were trypsinized, resuspended in fresh media, and seeded into 96-well

plates. Following 24 hours of incubation, fresh media containing treatment compounds and

MTT (0.5 mg/mL) was replaced. The formazan crystals were dissolved with DMSO after the

two-hour incubation period, and the cell viability was evaluated by measuring the absorbance

at 595 nm to assess cell viability(24).

Biofilm formation

A 96-well microtiter plate was used, with 200 μL of test medium and 20 μL of a 48-hour LB

broth culture as the inoculum. After incubating at 37°C for 48 hours under static conditions,

unbound cells were washed with sterile PBS. Biofilm-bound cells were stained with 1%

Crystal Violet (CV) for 15 minutes, washed with distilled water, and the bound dye was

extracted with 70% ethanol. The absorbance of the extracted CV was measured at 595 nm.

Viable cells were determined by scraping the biofilms from the wells, performing a 10-fold

dilution, and spread plating to enumerate colony-forming units (CFU). (25) Treatments were

administered with a triple combination of postbiotics at different pH levels.

Rheological analysis

The viscosity of the formulation was measured using a Brookfield Viscometer (LVDV-II +

Pro) with Spindle No. 64 at different temperatures. The formulation was equilibrated at the

initial temperature and its viscosity was measured. The sample was then subjected to

incremental temperature increases, with viscosity readings taken at each step after allowing

the formulation to stabilize. The results, indicating viscosity changes with temperature,

demonstrated the formulation's suitability for prolonged retention and enhanced therapeutic

efficacy, as a significant increase in viscosity with rising temperature suggests better

performance under therapeutic conditions.

 ***In vivo* infection study using BALB/c Mice.**

The reported animal experiments are confirmed to be compliant with ARRIVE guidelines by

the Institutional Animal Ethical Committee of SASTRA Deemed to be University (Approval

number: (740/SASTRA/IAEC/RPP)).

The *In vivo* study was organized into five groups (N=6 per group): disease control, placebo,

customized vaginal wash, probiotic wash, and commercial wash. A pilot study was conducted

before the main study to optimize the experimental parameters and determine the appropriate

study duration. The main study, spanning 14 days, involved vaginal induction of *E. coli*

UTI89 GFP in BALB/c mice by administering 20 µL of a bacterial suspension (10^6CFU/mL)

directly into the vaginal canal using a sterile pipette under aseptic conditions, ensuring

precise and uniform colonization. Following induction, the mice were monitored for signs of

infection, such as inflammation and vaginal discharge, with body weight recorded regularly.

Treatment with the vaginal wash formulations was applied daily. Urine samples were

collected on days 8, 11, and 14, and vaginal discharge samples were collected on days 3, 4, 5,

6, 8, 10, and 12. Blood samples were collected for serum creatinine assays (AGAPPE

Diagnostics Ltd, Mumbai, India) to assess renal function. Mice were sacrificed, and their

organs were homogenized to enumerate bacterial load.

For the preparation of the probiotic wash, 0.01 M lactic acid/sodium lactate buffer (pH 3.2)

was cooled to 4 °C. To this buffer, 18% Poloxamer and freeze-dried bacterial powder were

slowly added with continuous agitation. *Lactobacillus jensenii* was introduced at a

concentration 10^9 CFU/mL, and the mixture was left at 4 °C until a clear solution was

obtained, achieving a final pH of 3.8.

The Poloxamer-407 formulation containing postbiotics was prepared by combining 18%

Poloxamer with tryptamine (4 µg/mL), (-)-Terpinen-4-ol (5 µg/mL), and itaconic anhydride

(8 µg/mL). For the commercial vaginal wash, 1 mL of the product was diluted with 99 mL

of distilled water to create a solution suitable for administration in BALB/c mice. This

dilution was performed to ensure safety and minimize potential irritation, as well as to

adhere to the volume constraints appropriate for the species used in this study

Results

Triple combination of postbiotics reduces intracellular bioburden of _E. coli_ UTI89 in

T24 bladder cells, inhibits biofilm _invitro_ and exhibits non-toxicity in Raw macrophages.

The study evaluated the efficacy of postbiotic metabolites—tryptamine of 4 µg/ml, (-)-

terpinen-4-ol (5 µg/ml), and itaconic anhydride (8 µg/ml) —in reducing bacterial load using

an _In vitro_ infection in T24 bladder cells and preformed biofilm assays (Figure.1A). These

concentrations were selected based on our previous research, which indicated that these

specific levels of tryptamine, (-)-terpinen-4-ol, and itaconic anhydride exhibited optimal

antimicrobial activity against relevant bacterial strains. This prior work demonstrated that

these concentrations effectively reduced bacterial load while maintaining cell viability,

thereby supporting their use in the current study. Treatments included individual metabolites,

synergistic combinations, and a triple-compound mixture. Notably, the combination of all

three metabolites resulted in a significant 9-log reduction in bacterial load, underscoring their

potent synergistic effect in combating bacterial infections, particularly within T24 bladder

cells (Figure.1B). Additionally, in vitro toxicity studies using Raw macrophages were

conducted to assess the safety of these compounds. The MTT assay revealed no significant

cytotoxicity at concentrations of 4 µg/ml for tryptamine, 5 µg/ml for (-)-terpinen-4-ol, and 10

µg/ml for itaconic anhydride, indicating that these compounds are safe within the tested

concentration range (Supplementary Figure.1). These findings suggest that the triple

combination of postbiotics not only effectively reduces bacterial burden but also maintains a

safety profile, making it a promising candidate for therapeutic applications.

Figure 1. *In vitro* **cell line infection study with T24 Bladder cells of the metabolites and its synergistic effects.** A) Biofilm CFU Reduction in Preformed Biofilms Treated with Individual and Combined Metabolites B) Synergistic Effects of Metabolite Combinations on Bacterial Load in T24 Bladder Cells. One tail t test was performed to determine the significance, **$p<0.01$, ***$p<0.001$, n=3, EDTA (0.1%) used as a positive control for preformed biofilm.

Development of a customized vaginal wash for enhanced intravaginal delivery of vaginal *Lactobacillus* metabolites

Poloxamer 407 (procured from Sigma-Aldrich), a copolymer known for its unique liquid-to-gel transition upon temperature alteration, forms the base of the formulation (Figure. 3). Poloxamer 407, has a molecular weight of approximately 12.6 kDa and is hydrophilic in nature owing to its polyoxyethylene content (~70%) it serves as the key component of the

formulation. This copolymer's thermosensitive behavior, transitioning from a liquid to a gel

state in response to temperature changes, enables precise control over gelation kinetics. The

addition of Tryptamine (4 µg/ml), (-)-terpinen-4-ol (5 µg/ml), and itaconic anhydride

(8µg/ml) further enhances the formulation's therapeutic efficacy.

The optimal concentration of poloxamer for vaginal drug delivery systems is typically in the

range of 18-21% w/w. At an optimized concentration of 18%, Poloxamer 407 confers the

formulation with ideal rheological properties, ensuring optimal spreadability and retention

within the vaginal region. At room temperature (25 °C), the formulation exhibits low

viscosity, facilitating ease of application and mucosal coverage (Figure. 2B). Upon exposure

to body temperature (37 °C), the formulation undergoes in situ gelation, ensuring prolonged

retention within the vaginal environment (Figure. 2A). The rheological study using

Brookfield Viscometer (LVDV-II+Pro, Brookfield Engineering, USA with spindle no. 64)

demonstrated that the viscosity of the formulation increases significantly with the rise in

temperature, indicating its suitability for prolonged retention and enhanced therapeutic

efficacy (Figure 2C).

Figure 2 Sol-Gel Transformation of the customized vaginal wash formulation using

Poloxamer-407, where F1 is the *Lactobacillus* extracted metabolites incorporated in 18%

Poloxamer 407 and F2 is the procured metabolites incorporated in 18% Poloxamer 407 A)

Gel form at 37^0C B) Solution form at below 20 ^{0}C C) Rheological analysis of the formulation

to check the viscosity using Brookfield Viscometer (LVDV-II + Pro, Brookfield Engineering,

USA) with spindle no. 64. ELF means Extracted metabolites from *Lactobacillus* incorporated

in poloxamer 407 and PLF procured compounds incorporated in poloxamer 407.

Biofilm inhibition and bacterial growth suppression by the Customized Vaginal Wash

The findings from our study demonstrate the effectiveness of the customized vaginal wash

formulation in significantly reducing biofilm formation and bacterial viability across a range

of pH levels (Figure. 3A), highlighting its robust inhibitory capabilities. The observed

decrease in biofilm density and structural integrity, as evidenced by microscopy (Figure .3B)

and crystal violet assays (Supplementary Figure3C-D), suggests that the formulation

effectively disrupts the biofilm matrix, a key factor in preventing persistent bacterial

colonization and infection. The 9-log reduction in colony-forming units (CFUs) (Figure 3E)

further confirms the formulation's capacity to lower bacterial viability within the biofilm,

suggesting that it not only inhibits biofilm matrix formation but also actively reduces the

existing bacterial population (Figure 3E).

Figure. 3 Inhibition of Biofilm formation matrix on cover glass by the customized vaginal wash formulation A) Biofilm formation after treatment with different pH (2, 3, 4.5, 6, 7 and 8) of customized formulation B) Microscopic images of biofilm after treatment with different pH of customized formulation C) Plate showing Crystal violet assay for biofilm D) Quantification of crystal violet assay E) Colony forming units. One tail *t test* was performed to determine the significance., ***p<0.001, n=3

Complementing these findings, scanning electron microscopy (SEM) and fluorescence imaging were employed to assess bacterial growth inhibition following treatment with the customized formulation. SEM images revealed a significant reduction in bacterial density, coupled with noticeable morphological abnormalities in the treated groups compared to the untreated control (Figure. 4A). Fluorescence imaging further demonstrated decreased bacterial fluorescence intensity, indicating diminished bacterial viability post-treatment (Figure. 4B).

A. Untreated *E. coli* UTI89   **B. Treated *E. coli* UTI89 with formulation**

Figure. 4 Scanning Electron Microscope and Fluorescent images shows the inhibition of bacterial growth after treating with the formulation. A. Untreated *E. coli* UTI89, B. Treated with formulation. Scale bar =20µm

Customized vaginal wash formulation retains its stability and biological activity for 3 months at 4°C.

Results revealed that despite the prolonged storage period at 4°C for 3 months, the customized *Lactobacillus*-derived wash retained its ability to prevent biofilm formation of clinically relevant bacterial strains, including *E. coli* UTI89, *E. coli* CFT073, *E. coli* F11, *Klebsiella pneumoniae*, Methicillin-Resistant *Staphylococcus aureus* (MRSA), and *Pseudomonas aeruginosa*. (Supplementary Figure 2A) and there was ~ 9 log reduction in the growth of all tested bacterial strains (Supplementary Figure 2B). This broad-spectrum efficacy underscores the wash's potential utility in combating various infections encountered in clinical settings.

Customized Vaginal Wash prevents vaginal inflammation in BALB/c mice without affecting body weight.

In this study, the efficacy of various vaginal wash formulations was evaluated using a BALB/c mouse model of vaginal infection. The study included five groups (N=6 per group): disease control, placebo, customized vaginal wash, probiotic wash, and commercial wash.

Detailed schematics of the animal study are provided in Figure 5. On day 0, a 20 µL suspension containing 10^6 CFU/mL of *E. coli* UTI89 GFP was instilled into the vaginal cavity of BALB/c mice to induce infection. The inoculation was performed under aseptic conditions to ensure precise bacterial colonization within the vaginal canal. Treatment with the respective vaginal wash formulations began on day 3 post-infection, continuing through day 14 (Supplementary Figure. 3).

Figure 5: Schematics of Animal Study

By day 8, observations revealed significant differences among the treatment groups. The disease control, probiotic wash, and commercial wash groups exhibited pronounced

inflammation, including redness, swelling, and the formation of papules. In contrast, the customized vaginal wash-treated group showed significantly reduced inflammation, with vaginal health closely resembling that of the normal control group (Figure.6A).

Throughout the study, the body weight of the mice was monitored. While the baseline weight ranged between 35-40 grams, by day 8, the disease control, probiotic wash, and commercial wash groups exhibited a substantial weight loss of approximately 10 grams. Notably, mice treated with the customized vaginal wash maintained their baseline weight, indicating better overall health and the efficacy of the customized formulation in managing the infection (Figure.6B).

Figure 6. Effect of vaginal wash formulations on inflammation and body weight in BALB/c Mice infected with *E. coli* UTI89 GFP. A) Inflammation of vagina of BALB/c Mice, Blue arrow mark indicated the inflammation B) Body weight measurements of

BALB/c mice. One tail t test was performed to determine the significance. *p<0.05,

**p<0.01. Data represented as a mean ± standard error of the mean (SEM) (N= 6).

Customized vaginal wash reduced bacterial load in vaginal discharge.

Following formulation treatment, bacterial load in the vaginal discharge of BALB/c mice was

quantified by plate counts on Day 14 post infection (Figure 7A), Since *E. coli* UTI89 GFP

strain was used for infection, it was directly employed for fluorescent imaging (Figure 7B).

Plate counts (Figure 7A) revealed that the metabolite-based formulation achieved a

significant 8-10 log reduction in bacterial count compared to the untreated control, probiotic

formulation, and commercial wash, demonstrating its efficacy in inhibiting bacterial growth

and colonization. Both the probiotic wash and commercial wash displayed ~ 2 log reduction

in cell counts relative to untreated infected control. Similarly, fluorescent images on Day 6,

Day 8, and Day 12 post-infection revealed that, relative to commercial wash, which failed to

curtail the growth of *E. coli* UTI89 strain, metabolite wash completely mitigated *E. coli*

UTI89 growth (Figure 7B).

Figure. 7 Bacterial loads in vaginal discharge of BALB/c Mice post infection. A) Quantifications were performed in triplicate and represented as a mean ± standard error of the mean (SEM) (N= 6) B) Fluorescent images showing the bacterial load in the vaginal discharge of BALB/c Mice at Day 4, Day 8, and Day 12 post infection. (N= 6)

Absence of bacteria in urine/ organs after treatment with customized vaginal wash.

We assessed the efficacy of a metabolite-based wash in preventing urinary tract infections

(UTIs) and related complications in BALB/c mice. Infection was induced using *E. coli*

UTI89 GFP, and bacterial load in urine was quantified at multiple time points post-infection

(Figure. 82A). Fluorescent imaging of urine samples collected on Day 8, Day 11, and Day 14

revealed bacterial colonization dynamics within the urinary tract (Figure. 8B). Remarkably,

no *E. coli* UTI89 GFP was detected in the urine of mice treated with the metabolite wash,

indicating successful prevention of UTI development.

A)

B)

Figure. 8 Bacterial loads in Urine of BALB/c Mice post infection. "0" on the y-axis denotes

unsuccessful urine collection. Log 1 on the y-axis represents the baseline (No bacterial count)

B) Fluorescent images showing the bacterial load in the urine of BALB/c Mice at Day 8, Day

11, and Day 14 post infection.

Additionally, we evaluated the bacterial burden in vital organs implicated in UTIs, including

the kidney, urinary bladder, and vagina. Mice were sacrificed, and the organs were

homogenized to enumerate bacterial load. Mice treated with the metabolite wash exhibited no

bacterial load relative to either untreated control or treatments with probiotic wash/

commercial wash, highlighting its protective effects against cystitis and vaginal infections

(Figure. 9). The normal creatinine levels were maintained in these mice during metabolite

wash, further demonstrating the safety and efficacy of the metabolite wash in preventing

infection and preserving renal function (Supplementary Figure. 4).Thus the formulated

metabolite wash was superior in successfully treating UTI caused by UPEC in mice models

relative to either a commercial wash/probiotic wash. In conjunction with non toxic nature, the

customized postbiotic formulation carries a strong translational potential and can be

successfully employed as a stand-alone/ supplemental therapy to treat UTI infections.

Figure. 9 Bacterial bioburdens in organs (Kidney, Vagina and Urinary bladder)

Quantifications were performed in triplicate and represented as a mean ± standard error of the

mean (SEM) (N= 6)

Discussion

Although postbiotic supplements are not as widely available as probiotics yet, postbiotics are

still superior due to their purity, ease of preparation, long shelf life, capacity for mass

production, precise action, and ability to elicit more targeted responses through ligand-

receptor interactions.(26). Because postbiotics don't include any bacteria—dead, live, or

fragmented—they have a higher safety profile than probiotics. The increasing interest in

postbiotics as a safer and more effective alternative to probiotics in feminine hygiene

products is noteworthy. Postbiotics, which are non-viable microbial metabolites, offer several

advantages over traditional probiotics, particularly in formulations aimed at preventing

urinary tract infections Postbiotics are both natural and offer a unique and promising method

for reestablishing host-microbe balance with low toxicity and consistent functional features

since their lack of bacterial components reduces the possibility of potentially hazardous

bacterial dissemination (Figure. 1 &Supplementary figure 1). The advantages of using

postbiotics in vaginal washes are they do not contain live bacteria, which significantly

reduces the risk of infections, especially in vulnerable populations. This absence of viable

microorganisms means there's no potential for bacterial translocation, making them safer than

probiotic formulations that may include live or dead bacteria. Postbiotics can elicit specific

responses through ligand-receptor interactions, allowing for more precise therapeutic effects.

This targeted action can enhance their effectiveness in managing bacterial infections and

maintaining vaginal health. The formulation of postbiotics can be more economical than

probiotics, making them accessible to a broader range of consumers across different

socioeconomic groups (27, 28) Due to their non-viable nature, postbiotics are easier to

formulate and have a longer shelf life compared to probiotics(29). While postbiotic

formulations are emerging in the market, many existing feminine hygiene products still rely

on probiotics. Most probiotic-based hygiene washes incorporate live bacteria such as

Lactobacillus species, which are believed to help restore the natural flora of the vagina and

prevent infections. However, these products may carry risks associated with bacterial

viability and potential contamination(30). The present invention relates to a novel feminine

hygiene wash formulation designed to improve women's health and well-being. The

formulation comprises a combination of postbiotics and poloxamer 407, offering a fragrance-

free, pH balanced solution that effectively prevents urinary tract infections (UTIs). This

invention provides a cost-effective alternative to antibiotics and aims to reach all socioeconomic groups, thereby enhancing the quality of life for women worldwide.

The in *vivo* findings suggest that the customized vaginal wash offers a promising therapeutic strategy for managing bacterial infections and preventing UTIs, outperforming both probiotic and commercial formulations. Further investigation in clinical settings is warranted to validate these results and explore the potential for broader applications in human health.

Acknowledgement

The authors acknowledge SASTRA Deemed University for the infrastructural support. We would like to express our gratitude to the Centre for Advanced Research in Indian Systems of Medicine (CARISM), supported by the FIST Biochemical Engineering-II grant (SR/FST/ET-1/2020/614), for their generous support. Additionally, we extend our sincere thanks to Ms. Sudha V for her indispensable assistance, notably in conducting GC-MS analysis. The authors would like to thank Trichy SRM Medical college and Research Centre for the collection and isolation vaginal probiotics.

.

Funding

This work was supported by the Women Scientist-B, Department of Science and Technology, Government of India under grant number DST/WOS-B/HN-32/2021and NSS thankfully acknowledge Indian Council of Medical research funded project (ICMR) (Grant number: EMDR/SG/14/2024/01-04170) to conduct this study.

Conflict of Interest

The authors have filed a patent for a formulation containing postbiotics derived from
Lactobacilli to treat induced urinary tract infections in a mouse model (Patent Application
Number: 202441057663; Reference Number: TEMP/E-1/67129/2024-CHE).

References

1. Kiiru S, Maina J, Katana J, Mwaniki J, Asiimwe BB, Mshana SE, Keenan K, Gillespie
SH, Stelling J, Holden MTG, Kiiru J. 2023. Bacterial etiology of urinary tract
infections in patients treated at Kenyan health facilities and their resistance towards
commonly used antibiotics. PLoS One 18:e0277279.

2. LeBegue CE, Love BL, Wyatt MD. 2020. Microbes as Drugs: The Potential of
Pharmabiotics. Pharmacotherapy: The Journal of Human Pharmacology and Drug
Therapy 40:102–106.

3. Cordaillat-Simmons M, Rouanet A, Pot B. 2020. Live biotherapeutic products: the
importance of a defined regulatory framework. Exp Mol Med 52:1397–1406.

4. Nader-Macías MEF, Juárez Tomás MS. 2015. Profiles and technological requirements
of urogenital probiotics. Adv Drug Deliv Rev 92:84–104.

5. Reid G. 2001. Probiotic agents to protect the urogenital tract against infection.
American Journal of Clinical Nutrition 73.

6.   De Gregorio PR, Tomás MSJ, Terraf MCL, Nader-Macías MEF. 2014. In vitro and in
vivo effects of beneficial vaginal lactobacilli on pathogens responsible for urogenital
tract infections. J Med Microbiol 63:685–696.

7.   Suez J, Zmora N, Segal E, Elinav E. 2019. The pros, cons, and many unknowns of
probiotics. Nat Med 25:716–729.

8.   Zhang T, Zhang W, Feng C, Kwok L-Y, He Q, Sun Z. 2022. Stronger gut microbiome
modulatory effects by postbiotics than probiotics in a mouse colitis model. NPJ Sci
Food 6:53.

9.   Salminen S, Collado MC, Endo A, Hill C, Lebeer S, Quigley EMM, Sanders ME,
Shamir R, Swann JR, Szajewska H, Vinderola G. 2021. The International Scientific
Association of Probiotics and Prebiotics (ISAPP) consensus statement on the definition
and scope of postbiotics. Nat Rev Gastroenterol Hepatol 18:649–667.

10.   Nataraj BH, Ali SA, Behare P V., Yadav H. 2020. Postbiotics-parabiotics: the new
horizons in microbial biotherapy and functional foods. Microb Cell Fact 19:168.

11.   Croatti V, Parolin C, Giordani B, Foschi C, Fedi S, Vitali B. 2022. Lactobacilli
extracellular vesicles: potential postbiotics to support the vaginal microbiota
homeostasis. Microb Cell Fact 21:237.

12.   Abramov V, Khlebnikov V, Kosarev I, Bairamova G, Vasilenko R, Suzina N, Machulin
A, Sakulin V, Kulikova N, Vasilenko N, Karlyshev A, Uversky V, Chikindas ML,
Melnikov V. 2014. Probiotic Properties of Lactobacillus crispatus 2,029: Homeostatic
Interaction with Cervicovaginal Epithelial Cells and Antagonistic Activity to
Genitourinary Pathogens. Probiotics Antimicrob Proteins 6:165–176.

13.   Tachedjian G, Aldunate M, Bradshaw CS, Cone RA. 2017. The role of lactic acid
production by probiotic Lactobacillus species in vaginal health. Res Microbiol
168:782–792.

14.   Ventolini G. 2016. Progresses in Vaginal Microflora Physiology and Implications for
Bacterial Vaginosis and Candidiasis. Women's Health 12:283–291.

15.   Pericolini E, Gabrielli E, Ballet N, Sabbatini S, Roselletti E, Cayzeele Decherf A,
Pélerin F, Luciano E, Perito S, Jüsten P, Vecchiarelli A. 2017. Therapeutic activity of a
Saccharomyces cerevisiae -based probiotic and inactivated whole yeast on vaginal
candidiasis. Virulence 8:74–90.

16.   Abbasi A, Rad AH, Ghasempour Z, Sabahi S, Kafil HS, Hasannezhad P, Rahbar Saadat

Y, Shahbazi N. 2022. The biological activities of postbiotics in gastrointestinal

disorders. Crit Rev Food Sci Nutr 62:5983–6004.

17.   Ventolini G. 2013. New Insides on Vaginal Immunity and Recurrent Infections. J Genit

Syst Disord 02.

18.   Fettweis JM, Serrano MG, Girerd PH, Jefferson KK, Buck GA. 2012. A New Era of

the Vaginal Microbiome: Advances Using Next-Generation Sequencing. Chem

Biodivers 9:965–976.

19.   Sarkar A, Mandal S. 2016. Bifidobacteria—Insight into clinical outcomes and

mechanisms of its probiotic action. Microbiol Res 192:159–171.

20.   Scarpellini E, Rinninella E, Basilico M, Colomier E, Rasetti C, Larussa T, Santori P,

Abenavoli L. 2021. From Pre- and Probiotics to Post-Biotics: A Narrative Review. Int J

Environ Res Public Health 19:37.

21.   Wu M-H, Pan T-M, Wu Y-J, Chang S-J, Chang M-S, Hu C-Y. 2010.

Exopolysaccharide activities from probiotic bifidobacterium: Immunomodulatory

effects (on J774A.1 macrophages) and antimicrobial properties. Int J Food Microbiol

144:104–110.

22. Schaefer L, Auchtung TA, Hermans KE, Whitehead D, Borhan B, Britton RA. 2010.

The antimicrobial compound reuterin (3-hydroxypropionaldehyde) induces oxidative

stress via interaction with thiol groups. Microbiology (N Y) 156:1589–1599.

23. Nair VG, Srinandan CS, Rajesh YBRD, Narbhavi D, Anupriya A, Prabhusaran N,

Nagarajan S. 2024. Biogenic amine tryptamine in human vaginal probiotic isolates

mediates matrix inhibition and thwarts uropathogenic E. coli biofilm. Sci Rep

14:15387.

24. Peng C-C, Chen K-C, Peng RY, Su C-H, Hsieh-Li HM. 2006. Human urinary bladder

cancer T24 cells are susceptible to the Antrodia camphorata extracts. Cancer Lett

243:109–119.

25. Miryala S, Nair VG, Chandramohan S, Srinandan CS. 2020. Matrix inhibition by

Salmonella excludes uropathogenic E. coli from biofilm. FEMS Microbiol Ecol 97.

26. Nataraj BH, Ali SA, Behare P V., Yadav H. 2020. Postbiotics-parabiotics: the new

horizons in microbial biotherapy and functional foods. Microb Cell Fact 19:168.

27.   Vinderola G, Druart C, Gosálbez L, Salminen S, Vinot N, Lebeer S. 2023. Postbiotics

in the medical field under the perspective of the ISAPP definition: scientific,

regulatory, and marketing considerations. Front Pharmacol 14.

28.   Żółkiewicz J, Marzec A, Ruszczyński M, Feleszko W. 2020. Postbiotics—A Step

Beyond Pre- and Probiotics. Nutrients 12:2189.

29.   Babu AS, Guruprasath N, Adeyeye SAO, Sankarganesh P, Kumar AG, Sivapriya T.

2023. A Critical Analysis of Postbiotics: Exploring their Potential Impact on the Health

and Food Industries. J Pure Appl Microbiol 17:2041–2059.

30.   Ji J, Jin W, Liu S, Jiao Z, Li X. 2023. Probiotics, prebiotics, and postbiotics in health

and disease. MedComm (Beijing) 4.

Figures

Figure 1. *In vitro* **cell line infection study with T24 Bladder cells of the metabolites and its synergistic effects.** A) Biofilm CFU Reduction in Preformed Biofilms Treated with Individual and Combined Metabolites B) Synergistic Effects of Metabolite Combinations on Bacterial Load in T24 Bladder Cells. One tail t test was performed to determine the significance, **$p<0.01$, ***$p<0.001$, n=3, EDTA (0.1%) used as a positive control for preformed biofilm

Figure 2 Sol-Gel Transformation of the customized vaginal wash formulation using Poloxamer-407, where F1 is the *Lactobacillus* extracted metabolites incorporated in 18% Poloxamer 407 and F2 is the procured metabolites incorporated in 18% Poloxamer 407 A) Gel form at 37^0C B) Solution form at below $20\ ^0$C C) Rheological analysis of the formulation to check the viscosity using Brookfield Viscometer (LVDV-II + Pro, Brookfield Engineering, USA) with spindle no. 64. ELF means Extracted metabolites from *Lactobacillus* incorporated in poloxamer 407 and PLF procured compounds incorporated in poloxamer 407.

Figure. 3 Inhibition of Biofilm formation matrix on cover glass by the customized vaginal wash formulation A) Biofilm formation after treatment with different pH (2, 3, 4.5, 6, 7 and 8) of customized formulation B) Microscopic images of biofilm after treatment with different pH of customized formulation C) Plate showing Crystal violet assay for biofilm D) Quantification of crystal violet assay E) Colony forming units. One tail *t test* was performed to determine the significance., ***p<0.001, n=3

Figure. 4 Scanning Electron Microscope and Fluorescent images shows the inhibition of bacterial growth after treating with the formulation. A. Untreated *E. coli* UTI89, B. Treated with formulation. Scale bar =20µm

Figure 5: Schematics of Animal Study

Figure 6. Effect of vaginal wash formulations on inflammation and body weight in BALB/c Mice infected with *E. coli* UTI89 GFP. A) Inflammation of vagina of BALB/c Mice, Blue arrow mark indicated the inflammation B) Body weight measurements of BALB/c mice. One tail t test was performed to determine the significance. *p<0.05, **p<0.01. Data represented as a mean ± standard error of the mean (SEM) (N= 6).

A)

B)

Figure. 7 Bacterial loads in vaginal discharge of BALB/c Mice post infection. A)
Quantifications were performed in triplicate and represented as a mean ± standard
error of the mean (SEM) (N= 6) B) Fluorescent images showing the bacterial load in

the vaginal discharge of BALB/c Mice at Day 4, Day 8, and Day 12 post infection. (N= 6)

A)

B)

Figure. 8 Bacterial loads in Urine of BALB/c Mice post infection. "0" on the y-axis denotes unsuccessful urine collection. Log 1 on the y-axis represents the baseline (No bacterial count) B) Fluorescent images showing the bacterial load in the urine of BALB/c Mice at Day 8, Day 11, and Day 14 post infection.

Figure. 9 Bacterial bioburdens in organs (Kidney, Vagina and Urinary bladder)
Quantifications were performed in triplicate and represented as a mean ± standard error
of the mean (SEM) (N= 6)