1

Heterozygosity at a conserved candidate sex determination locus is associated

with female development in the clonal raider ant (*Ooceraea biroi*)

Kip D. Lacy[1*], Jina Lee[1], Kathryn Rozen-Gagnon[2,3], Wei Wang[3], Thomas S. Carroll[3], Daniel

J.C. Kronauer[1,4*]

[1]Laboratory of Social Evolution and Behavior, The Rockefeller University, New York, NY,

USA.

[2]Department of Molecular Genetics, University of Toronto, Toronto, Canada.

[3]Bioinformatics Resource Center, The Rockefeller University, New York, NY, USA.

[4]Howard Hughes Medical Institute, New York, NY, USA.

[*]Email: klacy@rockefeller.edu (K.D.L.); dkronauer@rockefeller.edu (D.J.C.K)

Abstract

Sex determination is a developmental switch that triggers sex-specific developmental programs. This switch is "flipped" by the expression of genes that promote male- or female-specific development. Many lineages have evolved sex chromosomes that act as primary signals for sex determination. However, haplodiploidy (males are haploid and females are diploid), which occurs in ca. 12% of animal species, is incompatible with sex chromosomes. Haplodiploid taxa must, therefore, rely on other strategies for sex determination. One mechanism, "complementary sex determination" (CSD), uses heterozygosity as a proxy for diploidy. In CSD, heterozygosity at a sex determination locus triggers female development, while hemizygosity or homozygosity permits male development. CSD loci have been mapped in honeybees and two ant species, but we know little about their evolutionary history. Here, we investigate sex determination in the clonal raider ant, *Ooceraea biroi*. We identified a 46kb candidate CSD locus at which all females are heterozygous, but most diploid males are homozygous for either allele. As expected for CSD loci, the candidate locus has more alleles than most other loci, resulting in a peak of nucleotide diversity. This peak negligibly affects the amino acid sequences of protein-coding genes, suggesting that heterozygosity of a non-coding genomic sequence triggers female development. This locus is distinct from the CSD locus in honeybees but homologous to a CSD locus mapped in two distantly related ant species, implying that this molecular mechanism has been conserved since a common ancestor that lived approximately 112 million years ago.

Introduction

Whether an animal develops as a male or a female is typically determined by a switch early in development. This switch, known as sex determination, is a multistep process beginning with primary sex determination signals triggering sex-specific splicing of downstream transcription factors that encode sexual identity (Williams and Carroll 2009). Although the downstream transcription factors are evolutionarily conserved, primary sex determination signals are evolutionarily labile and comprise a diverse range of environmental and genetic mechanisms (Bachtrog et al. 2014). Genetically encoded primary signals (sex-determining genes or chromosomes) are usually found in only one sex. Such systems do not work in haplodiploid taxa (an estimated 12% of animal species, including ants (Normark 2003)), where any allele on any homologous chromosome can be transmitted from diploid females to their haploid male sons, and all alleles experience selection to be viably transmitted from both sexes (Whiting 1935). Haplodiploids, therefore, require alternative sex determination mechanisms, but we know little about the molecular details.

Some parasitoid wasps use a mechanism involving maternal imprinting (Verhulst et al. 2010; Zou et al. 2020). However, in many Hymenoptera, inbred crosses produce diploid males, which led to the hypothesis that female development is triggered by heterozygosity as a proxy for diploidy (Whiting 1933; Whiting 1943). Under this "complementary sex determination" (CSD) hypothesis, different alleles at a given sex determination locus "complement" one another, meaning that heterozygosity triggers female development (**Fig. 1A**). By contrast, the presence of a single allele (either because the individual is haploid or because a diploid individual is homozygous at the sex determination locus) permits male development. Under CSD, diploid males produce diploid sperm and, therefore, are functionally sterile (van

Wilgenburg et al. 2006). This has several evolutionary consequences (Cook and Crozier 1995);

at the population level, there is selection against homozygosity, often meaning obligatory

avoidance of inbreeding (Page 1980). At the gene level, there is negative frequency-dependent

(balancing) selection on CSD loci, resulting in the co-occurrence of many alleles at similar

frequencies within populations (Laidlaw et al. 1956).

Having been a favored hypothesis for more than half a century, CSD was demonstrated in

the honeybee *Apis mellifera* in 2003 (Beye et al. 2003). In *A. mellifera*, biallelic heteromers of

the protein encoded by the *complementary sex determiner* (*csd*) gene trigger female

development, whereas monoallelic homomers permit male development (Beye et al. 2003; Beye

et al. 2013; Otte et al. 2023; Seiler and Beye 2024). Although this primary sex determination

signal differs from primary signals in other insects, it nonetheless converges upon the pathway

that transduces sex determination signals in holometabolous insects (Bopp et al. 2014; Sawanth

et al. 2016; Wexler et al. 2019). The Csd protein sex-specifically splices the closely linked

splicing factor *feminizer* (*fem*) (the honeybee version of *transformer* (*tra*) in *Drosophila* and

other insects), which in turn leads to the production of sex-specific isoforms of *doublesex* (*dsx*), a

transcription factor that regulates sex-specific development (Hasselmann et al. 2008; Gempe et

al. 2009; Verhulst et al. 2010; Zou et al. 2020). Intriguingly, *csd* is a paralog of *fem*. Following

the prediction that CSD loci should evolve under balancing selection, many *csd* alleles are held

at similar frequencies, and the locus bears signatures of adaptive evolution (Hasselmann and

Beye 2004), with amino acid-level heterozygosity in certain protein domains required for female

development (Otte et al. 2023).

CSD is assumed to occur in many Hymenoptera that occasionally produce diploid males

(van Wilgenburg et al. 2006). Until recently, however, heterozygosity-dependent female

development had not been demonstrated outside of *A. mellifera*, which remained the only species

for which a CSD locus had been genetically mapped. A recent genetic mapping study in the

wasp *Lysiphlebus fabarum* identified at least one and as many as four candidate sex

determination loci (Matthey-Doret et al. 2019) (the multi-locus extension of the CSD hypothesis

proposes that heterozygosity at any of multiple sex determination loci is sufficient to trigger

female development (Crozier 1971) (**Fig. 1B**)). However, a requirement of heterozygosity at

these loci for female development was not demonstrated. In the ant *Vollenhovia emeryi*, a

mapping study found two sex determination QTL (Miyakawa and Mikheyev 2015), with follow-

up work suggesting that these function as a multi-locus CSD system (Miyakawa and Miyakawa

2023). One of these QTL (*V.emeryiCsdQTL1*) contains two closely linked *tra* homologs,

reminiscent of the locus containing *csd* and *fem* in *A. mellifera*. By contrast, the other QTL

(*V.emeryiCsdQTL2*) only contains genes without annotated functions in sexual development.

Although *V. emeryi* sex determination is likely integrated through *tra* and *dsx*, the molecular

basis of their primary sex determination signals remains unclear (Miyakawa et al. 2018;

Miyakawa and Miyakawa 2023).

Recently, a sex determination locus was identified in the Argentine ant, *Linepithema*

humile (Pan et al. 2024). Consistent with CSD, all pairwise heterozygous combinations of the

seven identified alleles were found in females, whereas diploid males were invariably

homozygous for one of the seven alleles. Moreover, these alleles occurred at roughly equal

frequencies within populations, suggesting evolution under balancing selection. The mapped

locus is located within a noncoding genomic region tightly linked to *ant noncoding transformer*

splicing regulator (*ANTSR*), a long noncoding RNA (lncRNA) with previously uncharacterized

function. *ANTSR* knockdown in female-destined embryos leads to male-specific splicing of *tra*,

suggesting that this lncRNA transduces the primary sex determination signal encoded by the

heterozygosity (or hemi- or homozygosity) of the linked noncoding region (Pan et al. 2024).

Because the synteny of this region (including the presence of a lncRNA putatively homologous

to *ANTSR*) is conserved across the ants, bees, and vespoid wasps, it was suggested that this sex

determination locus may be deeply conserved among the Aculeata (Pan et al. 2024). However,

synteny is insufficient to demonstrate functional conservation, and genetic mapping of sex

determination has yet to be performed in most of these taxa.

To explore the evolution of sex determination across ants, we investigated the topic in the

clonal raider ant, *Ooceraea biroi.* Diploid males occur sporadically in this species, suggesting

that, like other ants, it might employ CSD (Kronauer et al. 2012). *O. biroi* reproduces via a mode

of parthenogenesis in which two haploid nuclei from a single meiosis fuse to produce diploid

offspring (central fusion automixis) (Oxley et al. 2014). Although heterozygosity is maintained

with high fidelity despite meiotic crossover recombination (Lacy et al. 2024), losses of

heterozygosity occur occasionally, probably due to the inheritance of one recombined and one

non-recombined version of a homologous chromosome (Kronauer et al. 2012; Oxley et al. 2014;

Trible et al. 2023; Lacy et al. 2024). Therefore, if *O. biroi* uses CSD, diploid males might result

from losses of heterozygosity at sex determination loci (**Fig. 1C**), similar to what is thought to

occur in other asexual Hymenoptera that produce diploid males (Rabeling and Kronauer 2012;

Matthey-Doret et al. 2019). Here, we use whole genome sequencing to map a sex determination

locus, demonstrate that most diploid males carry a loss of heterozygosity at this locus, assess the

homology of this locus with other sex determination loci, and explore whether balancing

selection has shaped evolution at this locus.

Results

<u>Identification of a candidate sex determination locus on chromosome 4</u>

To map sex determination loci in *O. biroi*, we first identified males based on sexual dimorphism and distinguished diploid from haploid males based on heterozygosity at eight genetic markers (**Supplementary Table S1, File S1, and Fig. S1**). We then sequenced the genomes of 16 diploid males and compared them with 19 previously sequenced genomes of diploid females (Trible et al. 2023; Lacy et al. 2024) (descriptions and metadata for all genomes sequenced in this study are in **Supplementary Table S2**). CSD loci must be heterozygous to trigger female development but can be homozygous in diploid males. To identify such sites, we calculated a "CSD index" for each single nucleotide polymorphism (SNP), which equals zero if any female is homozygous and equals the proportion of diploid males that are homozygous if all females are heterozygous. We found a peak of the CSD index on chromosome 4, which deviated significantly from the null distribution of CSD index values (**Fig. 2A**). At this peak, homozygosity levels differed significantly between males and females (**Supplementary Fig. S2**).

To corroborate our results, we looked for evidence of segmental losses of heterozygosity along chromosome 4. Eleven out of 16 sequenced diploid males bore segmental losses of heterozygosity on chromosome 4 (**Fig. 2B**). The intersection of these LOH segments comprised a 46kb peak at which all females were heterozygous, and most (11 out of 16) diploid males were homozygous. Under the CSD hypothesis, homozygosity for any allele at CSD loci permits male development. Consistently, *O. biroi* diploid males were homozygous for either allele at the 46kb region on chromosome 4 (**Table 1, Supplementary Fig. S3**).

The stretches of homozygosity found in diploid males could have resulted from

segmental deletions resulting from improperly repaired DNA damage or rare copy-neutral losses

of heterozygosity resulting from thelytokous parthenogenesis (Kronauer et al. 2012; Oxley et al.

2014; Trible et al. 2023; Lacy et al. 2024). To distinguish between these two scenarios, we

analyzed read depth and found that runs of homozygosity were not accompanied by changes in

copy number (**Supplementary Fig. S3**). This implies that diploid males arise when rare losses of

heterozygosity, which result from crossover recombination during meiosis, span the CSD locus.

Surprisingly, five of the 16 diploid males were not homozygous in this 46kb region. One

possible explanation is that these individuals had mutations in other genes involved in the sex-

determination pathway. We therefore inspected all unique mutations and losses of heterozygosity

in these five diploid males, but did not find any in genes with annotated sex determining

functions (**Supplementary File S2**). These individuals may carry mutations or losses of

heterozygosity in genes or regulatory elements with unannotated sex determination function.

Alternatively, male development may have been triggered by rare stochastic perturbations to

gene expression or splicing. We address these possibilities further in the Discussion.

O. biroi's candidate CSD locus is homologous to another ant sex determination locus but not to

honeybee *csd*

To assess homology to CSD loci mapped in other species, we identified *O. biroi*

orthologs of the genes found in other mapped sex determination loci. We found no homology

between the genes within the *O. biroi* CSD index peak and any of the genes within the putative

L. fabarum CSD loci (**Supplementary Table S3**). By identifying orthologs of genes within the

two sex determination QTL identified in *V. emeryi* (Miyakawa and Mikheyev 2015)

(**Supplementary Table S4**), we also identified homology to the *L. humile* CSD locus, because a

subset of *V.emeryiCsdQTL2* is homologous to the CSD identified in *L. humile* (Pan et al. 2024).

Homology to *V.emeryiCsdQTL2* was scattered across one arm of *O. biroi* chromosome 4,

including the 46kb CSD index peak, and homology to the *L. humile* CSD locus mapped within

the *O. biroi* CSD index peak (**Fig. 3A**). The *O. biroi* CSD index peak contains four protein-

coding genes that are also present in *V.emeryiCsdQTL2*, including *HCF*, *COPA*, an

uncharacterized gene, and *CRELD2*. Notably, the candidate CSD region in *O. biroi* also includes

a large stretch of sequence with no annotated genes downstream of an uncharacterized lncRNA.

LncRNAs often retain their functions and synteny with surrounding genes despite divergence in

sequence homology (Chodroff et al. 2010; Ulitsky et al. 2011; Quinn et al. 2016), raising the

possibility that, although this lncRNA has limited sequence homology with *L. humile ANTSR*, it

may have a homologous function (Pan et al. 2024). Regardless of the underlying molecular

mechanism, our data suggest that this CSD locus is conserved between *O. biroi*, *L. humile*, and

V. emeryi (**Fig. 3B**). This would mean that this locus is ancient, as these three species belong to

different ant subfamilies that diverged roughly 112 million years ago (Borowiec et al. 2025).

We performed our genetic mapping in clonal line A because this was the only clonal line

in which we found diploid males (see below). However, different clonal lines of *O. biroi* vary in

which portions of the genome are ancestrally homozygous (Oxley et al. 2014). Therefore, it was

unclear whether heterozygosity at the mapped CSD locus is required for female development in

all clonal lines of *O. biroi*. To investigate this, we inspected heterozygosity in diploid females

from six different clonal lines by retrieving previously published genome sequences from clonal

line B (Lacy et al. 2024) and sequencing genomes of diploid females from four additional clonal

lines (from both the native and invasive ranges of *O. biroi* (Kronauer et al. 2012; Trible et al.

2020)). All females from all six clonal lines (including 26 diploid females from clonal line B)

were heterozygous at the CSD index peak, consistent with its putative role as a CSD locus in all

O. biroi (**Fig. 3C**).

<u>The CSD index peak has high diversity in a non-coding region.</u>

Because CSD loci are expected to evolve under balancing selection, we expect many

different alleles to exist at those loci. For example, as many as 19 *csd* alleles were found to

segregate in honeybee populations (Hasselmann et al. 2008), and seven alleles were found to

segregate in a population of *L. humile*, despite a recent genetic bottleneck (Pan et al. 2024).

Looking for elevated genetic diversity is impossible by only studying genomes from a single

clonal line (i.e., asexual descendants of a single diploid female), so we looked across diploid

females from six different clonal lines. We calculated nucleotide diversity (π) in 5kb sliding

windows (step size = 1kb) across the genome and found a nucleotide diversity peak that fell

within the CSD index peak (**Fig. 4A, Supplementary Fig. S4**), between positions 1,717,000 and

1,730,000 on chromosome 4 (**Fig. 4B**).

We hypothesized that this nucleotide diversity peak resulted from many alleles found

among clonal lines of *O. biroi* at this locus. Identifying these alleles requires DNA sequences of

individual haplotypes, whereas short-read genome sequences of diploid females only provide

diploid genotypes. To investigate allelic diversity, we retrieved previously published haploid

male genome sequences from clonal lines A and B (Lacy et al. 2024) and sequenced whole

genomes of haploid males from additional clonal lines. We assembled their genomes *de novo* to

identify the different haplotypes found in this region. In addition to the haplotype found in the

reference genome, we identified six additional haplotypes (or "alleles") in the five clonal lines

for which we sequenced at least one haploid male genome (**Fig. 4C**, **Table 2**). These alleles

differ substantially in their DNA sequences in the region with high nucleotide diversity (**Fig.**

4C).

To investigate which genetic elements might act to determine sex in a heterozygosity-

dependent manner, we inspected all annotated genes in the CSD index peak (**Fig. 4D**). Within

this region, none of the protein-coding genes contained substantial heterozygosity that affected

amino acid identity (**Supplementary File S3**). To determine whether any genes or exons were

missing from our annotation, we performed RNA sequencing of embryos and several other life

stages (including female and male adults) using several different technologies (see Methods).

Briefly, we used long-read RNA sequencing (PacBio IsoSeq) to extend gene models and used

several different library preparation methods followed by standard short-read (Illumina)

sequencing: standard poly-A tail library prep to capture mRNAs, ribosomal RNA depletion to

capture lncRNAs that are not poly-adenylated, and small RNA sequencing to annotate miRNAs,

endo-siRNAs, and putative piRNAs. Although this improved the *O. biroi* genome annotation by

adding new genes and improving existing gene models (**Supplementary Table S5, File S4**), no

new genes or exons were identified near the CSD index peak.

To look for sex-biased gene expression and splicing, we performed differential gene

expression and differential transcript usage analysis between the transcriptomes of three male

and three female early-stage (white) pupae. We used pupae for this experiment because this is

the earliest life stage at which males and females can be readily distinguished. Primary sex

determination signals act early during embryonic development (Sawanth et al. 2016) and thus

need not be differentially expressed later in development. However, these genes can still be

expressed as late as the pupal stage (Beye et al. 2003). Many genes were differentially expressed

between males and females (24.6% of expressed genes) or had one or more differentially used

exons, including four genes near the CSD index peak (LOC105285605, LOC105283850,

LOC105283849, and LOC105283844) (**Supplementary Tables S5 and S6; Fig. S5**). However,

none of these overlapped with the region of elevated genetic diversity. Very few reads from the

male and female pupae samples aligned to the lncRNA, and although more reads aligned in

RNA-seq libraries from eggs prepared using ribosomal RNA depletion, these data are

insufficient for a rigorous assessment of sex-specific expression at this locus (**Supplementary**

Fig. S6). Therefore, it remains unclear whether this lncRNA plays a role in sex determination in

O. biroi.

The peak of nucleotide diversity minimally affects amino acid sequences and is primarily

located in a non-coding genomic region (**Fig. 4B-D**). Thus, a non-coding genetic element within

this region may instruct sex determination in *O. biroi* in a heterozygosity-dependent manner.

This is similar to *L. humile*, where the mapped CSD locus had high variability in non-coding and

non-exonic regions (Pan et al. 2024). RNA interference of the nearby lncRNA, *ANTSR*, led to

male-specific splicing of *tra*, raising the possibility that heterozygosity or homozygosity in the

non-exonic and non-coding *L. humile* CSD locus affects the expression level of *ANTSR*, which

instructs sex-specific splicing of *tra* (Pan et al. 2024). Because the peak of nucleotide diversity

within our CSD index peak minimally affects protein-coding sequences and the region is closely

linked to a lncRNA that is putatively an ortholog of *L. humile ANTSR*, our data suggest that the

mechanism of CSD in *O. biroi* may be conserved with *L. humile*.

Diploid male production differs across *O. biroi* clonal lines.

268  Over the course of our study, we sampled many males from clonal lines A and B, and a

269 few males from four additional clonal lines. Despite the fact that we maintain more colonies of

270 clonal line B than of clonal line A in the lab, all the diploid males we detected came from clonal

271 line A. We, therefore, looked back at data from previously genotyped males and found that this

272 pattern held across a previous study (Kronauer et al. 2012). In total, from clonal line A, 22 out of

273 114 males were diploid, whereas all 78 males genotyped from clonal line B were haploid (**Table**

274 **3**). This disparity is statistically significant (χ^2=15.4, df=1, p<0.0001), suggesting that the two

275 clonal lines differ in their propensity to produce diploid males.

276  One possible explanation for this phenomenon could be that *O. biroi* has multi-locus

277 CSD (**Fig. 1B**), but that clonal line A is ancestrally homozygous for all but one of the loci,

278 whereas clonal line B retains heterozygosity at two or more loci. Because multi-locus CSD

279 requires homozygosity at all sex determination loci for diploids to develop as males, in this

280 scenario, two or more losses of heterozygosity would be needed for diploid males to develop in

281 clonal line B. In contrast, only one loss of heterozygosity would be required in clonal line A,

282 which we used for our genetic mapping. However, we identified no additional candidate loci in

283 our mapping study, and other possibilities are equally likely, including that clonal lines A is

284 more prone to loss of heterozygosity, or is more permissive to diploid male development than

285 other clonal lines.

287 *O. biroi* lacks a *tra*-containing CSD locus.

288  A second QTL region identified in *V. emeryi* (*V.emeryiCsdQTL1*) contains two closely

289 linked *tra* homologs, similar to the closely linked honeybee *tra* homologs, *csd* and *fem*

290 (Miyakawa and Mikheyev 2015). This, along with the discovery of duplicated *tra* homologs that

undergo concerted evolution in bumblebees and ants (Schmieder et al. 2012; Privman et al.

2013) has led to the hypothesis that the function of *tra* homologs as CSD loci is conserved with

the *csd*-containing region of honeybees (Schmieder et al. 2012; Miyakawa and Mikheyev 2015).

However, other work has suggested that *tra* duplications occurred independently in honeybees,

bumblebees, and ants (Hasselmann et al. 2008; Koch et al. 2014), and it remains to be

demonstrated that either of these *tra* homologs acts as a primary CSD signal in *V. emeryi*. To test

whether the *tra* homolog-containing QTL (*V.emeryiCsdQTL1*) might be involved in CSD in *O.*

biroi, we searched for homology to this QTL in the *O. biroi* genome and identified several

regions scattered across *O. biroi* chromosome 2 (**Fig. 5A**). However, we did not find a CSD

index peak on chromosome 2, and no sex determination locus with homology to the *V. emeryi*

QTL was found in *L. humile* (Pan et al. 2024). Thus, the function of *V.emeryiCsdQTL1* as a

primary sex determination signal might have evolved after this lineage separated from other ants

(**Fig. 5B**). Alternatively, the sex determination activity of *V.emeryiCsdQTL1* could be ancestral

to these three ant species and independently lost in *O. biroi* (at least in clonal line A) and *L.*

humile.

Intriguingly, while clonal lines B, D, I, and M retain heterozygosity at all sites with

homology to *V.emeryiQTL1* (**Fig. 5C**), clonal lines A and C are homozygous in a region

homologous to a portion of *V.emeryiQTL1*. We note, however, that heterozygosity in this region

is limited, without elevated nucleotide diversity (**Supplementary Fig. S4**), and there are only

two nonsynonymous heterozygous variants in this region, with clonal line B conspicuously

lacking any nonsynonymous heterozygosity. Therefore, it seems unlikely that this region bears a

CSD locus in *O. biroi*. We also note that all clonal lines are heterozygous in the region

containing *O. biroi*'s *tra* ortholog (**Fig. 5C**), and that this gene is sex-specifically spliced

(**Supplementary Fig. S7**), similar to *tra* homologs in other Holometabola (Hasselmann et al.

2008; Miyakawa and Miyakawa 2023). This contrasts with the *tra-like* gene in *V. emeryi*, which,

like *A. mellifera csd*, is not sex-specifically spliced (Miyakawa and Miyakawa 2023).

<u>Discussion</u>

Genetically mapping sex determination loci in diverse taxa is key to understanding how

sex determination evolves. Here, we investigated the mode of sex determination in the clonal

raider ant, *O. biroi*. First, we mapped a candidate sex determination locus that was heterozygous

in all females, but homozygous for either allele in most diploid males (**Fig. 2**). Next, we showed

that most diploid males bear rare losses of heterozygosity that arise during thelytokous

parthenogenesis (**Fig. 2**). We then showed that this locus is homologous to a sex determination

locus found in two other ant species—*V. emeryi* and *L. humile* (**Fig. 3**), suggesting that this

mechanism of sex determination may be conserved among formicoid ants. By sequencing whole

genomes of haploid males and diploid females from other clonal lines of *O. biroi,* we showed

that the mapped locus corresponds to a region with high genetic diversity, following the

expectation that CSD loci should evolve under balancing selection (**Fig. 4**). This diversity peaks

in a noncoding genomic region, and we found limited evidence for functional heterozygosity in

nearby protein-coding sequences, suggesting that heterozygosity in this noncoding sequence may

trigger female development. Finally, we demonstrated that *O. biroi* CSD in clonal line A does

not map to the *tra*-containing sex determination locus previously identified in *V. emeryi*, but it

remains unclear whether *O. biroi* ancestrally has single-locus or multi-locus CSD (**Fig. 5**).

The presence of diploid males in many ant taxa previously led to the conclusion that most

ants employ CSD (van Wilgenburg et al. 2006). Along with *L. humile* (Pan et al. 2024) and *V.*

emeryi (Miyakawa and Mikheyev 2015), *O. biroi* is now one of three ant species for which

heterozygosity-dependent female development has been demonstrated at mapped CSD loci. Our

mapped locus appears to be homologous to *V.emeryiCsdQTL2* and to the locus identified in *L.*

humile (**Fig. 3**). As was found in *L. humile*, this locus colocalizes with a peak of nucleotide

diversity (**Fig. 4**), with seven alleles found among five clonal lines (**Fig. 4C, Table 2**), consistent

with the expectation that CSD loci should harbor many alleles due to evolution under balancing

selection. This implies that the role of this locus in CSD evolved before the divergence of the

Dorylinae from the other formicoid ants roughly 112 million years ago (Borowiec et al. 2025).

Although synteny analyses raised the possibility that this locus arose early within the evolution

of the Aculeata (Pan et al. 2024), genetic mapping studies of additional representative species of

different ant subfamilies (especially in the poneroid clade and the Leptanillinae) and across other

Aculeata will be required to determine when the function of this locus in CSD originated and

how it is distributed across taxa.

In *L. humile*, a lncRNA (*ANTSR*) closely linked to this locus plays a role in sex

determination (Pan et al. 2024). Although we observed some expression of the putatively

homologous lncRNA in *O. biroi* eggs of unknown sex, we were unable to assess sex-specific

expression. The reason is that males in *O. biroi* are exceedingly rare and morphologically only

identifiable at the pupal stage, well after primary sex determination signals are transduced to

self-sustaining sex-specific splicing of downstream transcription factors during the early stages

of embryonic development (Gempe and Beye 2011). Therefore, we likely missed differential

gene expression relevant to sex determination that occurs before the pupal stage.

We did not find a peak of the CSD index at the *O. biroi tra* homolog (**Fig. 5**), indicating

that, as in *L. humile*, the mode of CSD is not homologous to the system in *A. mellifera* mediated

by the *transformer* homologs *csd* and *fem*. This was previously suspected for *O. biroi* due to the

presence of only a single *tra* homolog (Oxley et al. 2014) rather than the two closely linked *tra*

homologs found in other ants (Privman et al. 2013). In *V. emeryi*, however, one of the two Csd

QTLs (*V.emeryiCsdQTL1*) spans the *tra* locus, raising the possibility that the molecular

mechanism of CSD might, in part, be conserved between *A. mellifera* and *V. emeryi* (Miyakawa

and Mikheyev 2015). It has not been demonstrated that either of *V. emeryi*'s *tra* homologs act as

a primary sex determination signal. Our results raise the possibility that the role of

V.emeryiCsdQTL1 as a primary sex determination signal evolved independently of *A. mellifera*

csd and after *V. emeryi* diverged from *L. humile*. However, further investigations into the

mechanisms of sex determination in *V. emeryi* and other ants are required to reconstruct the

evolutionary history of *V.emeryiCsdQTL1*'s function in hymenopteran sex determination.

Curiously, diploid males were only produced in clonal line A, even though many males

were collected from other clonal lines (**Table 3**). One possible explanation is that *O. biroi* has

multi-locus CSD, and clonal line A is ancestrally homozygous for all but one of the loci. Multi-

locus CSD has been suggested to limit the extent of diploid male production in asexual species

under some circumstances (Vorburger 2013; Matthey-Doret et al. 2019). Clonal line A is

ancestrally homozygous for a portion of the region homologous to the second *V. emeryi* QTL,

although it retains heterozygosity in the region overlapping *tra* (**Fig. 5C**). However, this region

lacks elevated genetic diversity and functional heterozygosity, making it a poor candidate for a

second, undetected, CSD locus. Further study would be required to determine whether *O. biroi*

has a second CSD locus. However, there may be other explanations for diploid males being

found exclusively in clonal line A. For example, clonal line A could be more likely to lose

heterozygosity than other clonal lines, or diploid males in clonal line A could be more likely to

develop and survive. Currently, we have no data to distinguish between these hypotheses.

384   Another peculiarity of our study is that five of the 16 sequenced diploid males retained

heterozygosity at the mapped CSD locus. One possible explanation for this would be that our

mapped locus is, in fact, a false positive. However, this seems unlikely given that 1) those losses

of heterozygosity occurred independently in multiple genetic backgrounds within clonal line A

(**Supplementary Fig. S3**), 2) a peak of nucleotide diversity co-occurs with the mapped locus

(**Fig. 4**), and 3) this locus is homologous to CSD loci mapped in two other ant species (**Fig. 3**).

Stochastic gene expression or splicing fluctuations may have caused these individuals to develop

as males despite having female genotypes. Certain allelic combinations of *A. mellifera csd*

trigger female development with incomplete penetrance (Beye et al. 2013), and the allelic

combination found in clonal line A might similarly be permissive to occasional male

development. Alternatively, these diploid males could result from mutations in or losses of

heterozygosity at different genetic loci. We note that it might be easier to find rare stochastically

produced diploid males in laboratory colonies of *O. biroi* than in sexual species. This is because

genetically encoded diploid males only result from loss of heterozygosity, which is rare in *O.*

biroi (Oxley et al. 2014; Lacy et al. 2024). By contrast, in sexually reproducing populations,

diploid males homozygous for sex determination loci will arise much more frequently due to

matings between individuals that share an allele at the sex determination locus (this effect is

exacerbated in invasive species like *L. humile* that have undergone recent genetic bottlenecks).

Thus, if stochastically produced diploid males occur at very low baseline levels in ants, such

males will rarely be detected in sexual species. In *O. biroi*, however, they could make up a

substantial proportion of diploid males, simply because diploid males caused by homozygosity at

the CSD locus are likewise extremely rare.

Finally, we note that heterozygosity-dependent female development, a fundamental tenet

of CSD, has major implications for the evolution of breeding systems and reproductive modes.

Due to the high fitness cost of producing sterile diploid males, species with CSD should

obligatorily outbreed or have other mechanisms of maintaining heterozygosity (van Wilgenburg

et al. 2006). In this study, most diploid males resulted from rare crossover-associated losses of

heterozygosity that spanned the putative CSD locus (the shortest loss of heterozygosity at the

mapped locus was 46kb and, therefore, likely resulted from crossover recombination rather than

gene conversion). Intriguingly, in many taxa, asexual lineages reproduce via mechanisms that

lead to complete homozygosity in a single generation (Suomalainen et al. 1987; Ma and

Schwander 2017). However, among the social Hymenoptera (and other Vespoids), the known

modes of asexual reproduction all maintain at least some heterozygosity (Rabeling and Kronauer

2012; Ma and Schwander 2017). For formicoid ants, bees, and other taxa with CSD, asexual

reproduction with high rates of heterozygosity loss would incur a steep fitness penalty due to the

production of sterile diploid males. This may help explain the unusual inheritance system

recently described in *O. biroi*, where following crossover recombination, reciprocally

recombined chromatids are faithfully co-inherited so that heterozygosity is rarely lost (Lacy et al.

2024).

How this ancient putative CSD locus triggers female development in a heterozygosity-

dependent manner remains unknown. In *L. humile*, RNAi experiments suggested that the

expression level of the lncRNA *ANTSR* affects sex-specific splicing of *tra*, even though

heterozygosity in the exons of *ANTSR* is not required for this effect (Pan et al. 2024). Thus, the

heterozygosity-dependent function of the candidate CSD locus may depend on local regulatory

interactions between homologous chromosomes, similar to a phenomenon known as transvection

(Duncan 2002). To better understand how heterozygosity induces female development in ants,

future studies will need to fine-map and functionally characterize this causal locus.

<u>Materials and Methods</u>

<u>Software Versions</u>

The versions of all software used for analyses are provided in **Supplementary Table S8**.

<u>Identifying Male Ploidy</u>

Clonal raider ant colonies are composed of female ants that reproduce asexually via a

type of thelytokous parthenogenesis termed central fusion automixis (Oxley et al. 2014). Males

are produced only sporadically (Oxley et al. 2014). Females of this species are diploid, whereas

males can either be haploid (presumably arising via a failure of central fusion following female

meiosis) or diploid (presumably arising due to rare losses of heterozygosity at CSD loci). Males

are easily recognizable due to substantial sexual dimorphism (males have eyes and wings,

whereas females are blind and flightless, and males also have different body shape and more

darkly pigmented cuticles than females). We sampled them opportunistically whenever they

were observed in colonies. We distinguished diploid males from haploid males based on

heterozygosity at several unlinked genetic markers (**Supplementary Table S1**). For this, we

disrupted one leg from each male using a Qiagen TissueLyser II and extracted DNA using

Qiagen's QIAmp DNA Micro Kit. We then PCR-amplified each marker and genotyped using

Sanger sequencing (for most markers) or via gel electrophoresis either after restriction digestion

(for two markers) or directly after amplification (for one marker at which the different alleles are

different sizes). Details of the genotyping methods can be found in **Supplementary Table S1**.

<u>Genome Sequencing and Preliminary Analysis</u>

For each library, we disrupted an individual ant with a metal bead and Qiagen's

TissueLyser II and extracted DNA with Qiagen's QIAmp DNA Micro Kit. We prepared Nextera

Flex Illumina DNA sequencing libraries for 16 diploid males from clonal line A, six diploid

females from different clonal lines, and seven haploid males from different clonal lines. We

sequenced these libraries using an Illumina NovaSeq 6000. We also accessed libraries for

individual genomes from previous studies. The information for all DNA sequencing libraries

used in this study can be found in **Supplementary Table S2**.

We trimmed reads using Trimmomatic 0.36 (Bolger et al. 2014), aligned them to the *O.*

biroi reference genome (McKenzie and Kronauer 2018) (Obir_v5.4, GenBank assembly

accession: GCA_003672135.1) using bwa mem (Li 2013), and then sorted, deduplicated, and

indexed using picard (http://broadinstitute.github.io/picard/). We called variants using GATK

HaplotypeCaller (version 4.2) (Poplin et al. 2018). To exclude falsely collapsed regions in the

reference genome, we filtered against sites found to be heterozygous in haploid males. We also

screened out any variants for which putatively heterozygous samples had a proportionate minor

allelic depth less than 0.25 and/or for which putatively homozygous samples had a minor allelic

depth greater than zero. We performed custom filtering of variants and calculated statistics using

pyvcf 0.6.8 (https://github.com/jamescasbon/PyVCF), and performed sliding window analyses

using pybedtools 0.9.0 (Dale et al. 2011).

<u>CSD Index Mapping</u>

CSD loci are heterozygous in females but can be homozygous in diploid males. Thus,

traditional association mapping cannot identify CSD loci because particular alleles do not trigger

female development. Therefore, to map candidate CSD loci, we used a "CSD Index," which

equals zero if any female is homozygous, but if all females are heterozygous, it equals the

proportion of diploid males that are homozygous. Before calculating this index, we had to

identify ancestrally heterozygous sites. All individuals within an *O. biroi* clonal line are

descended asexually from a single diploid female ancestor, and CSD loci would have been

heterozygous in that common ancestor. Putatively ancestrally heterozygous variants are those for

which at least one diploid individual is heterozygous, and all other diploid individuals are either

heterozygous for the same two alleles, or homozygous for one of the two alleles for which other

diploid(s) are heterozygous (Oxley et al. 2014; Lacy et al. 2024). After identifying putatively

ancestrally heterozygous SNPs, we calculated the CSD index for each SNP.

To assess the likelihood that our observed CSD index values were due to chance (false

positives), we randomly shuffled the identities of males and females to generate a null

distribution of CSD index values for each SNP. In doing so, we maintained the underlying

genetic structure by only shuffling sex labels within colonies. We computed p-values by

calculating the proportion of permutations with values greater than or equal to the observed CSD

index. To correct for multiple testing and control the false discovery rate across the genome-wide

analysis, we applied the Benjamini-Hochberg False Discovery Rate correction and then recorded

the mean CSD index value in sliding windows.

To directly evaluate the statistical association between homozygosity and sex, we applied

Fisher's Exact Test to assess whether SNP homozygosity frequencies differed significantly

between males and females and used the Benjamini-Hochberg False Discovery Rate correction

to adjust p-values.

<u>Losses of Heterozygosity</u>

500   We identified losses of heterozygosity as ancestrally heterozygous SNPs that had become

homozygous in a given sample. To assess whether homozygosity for either allele at the mapped

region permitted male development, we randomly chose a reference haploid male and assigned

allelic identity based on whether a focal sample was homozygous for the same allele or the

alternate allele to that possessed by the reference haploid male. To clearly illustrate losses of

heterozygosity, we used the ordinal position of SNPs rather than the position of SNPs in the

reference genome assembly.

507   Homozygosity in whole genome sequencing data can result from deletions, for which the

affected region would have half the read depth of the genome-wide average, or copy-neutral

losses of heterozygosity (which result rarely from meiotic recombination and central fusion

parthenogenesis (Kronauer et al. 2012; Rabeling and Kronauer 2012; Oxley et al. 2014; Lacy et

al. 2024)). To determine which of these processes caused homozygosity in diploid males, we

identified runs of homozygosity using "bcftools roh" (Narasimhan et al. 2016), obtained read

depths at all SNPs that passed filtering within these regions using "samtools depth –aa"

(Danecek et al. 2021) and then normalized by dividing by the genome-wide median read depth.

For samples without losses of heterozygosity on chromosome 4, we obtained normalized read

depth at all SNPs that passed filtering on chromosome 4.

<u>Homology Between CSD Loci Across Species</u>

To determine whether our mapped CSD locus bore homology to those identified in other

ant species, we used OrthoFinder (Emms and Kelly 2019) to identify orthologs of the genes

present in the two CSD QTL in *V. emeryi* (Miyakawa and Mikheyev 2015). The locus identified

in *L. humile* (Pan et al. 2024) is homologous to a subset of one of the *V. emeryi* QTL. In this

process, we noticed that one of the genes found within our mapped CSD locus (*cysteine-rich*

with EGF-like domain protein 2, or CRELD2) was also present in a second copy in a contig that

was erroneously duplicated during the genome assembly process (Chr4:21571-39095). We

masked this contig and realigned sequencing data for all subsequent analyses.

<u>Balancing Selection</u>

Because CSD loci are expected to evolve under negative frequency-dependent

(balancing) selection, many different alleles should segregate in populations. To assess this, we

used scikit-allel (Miles et al. 2019) to calculate nucleotide diversity across six different clonal

lines and reported nucleotide diversity in sliding windows.

From inspecting DNA sequencing alignments to the reference genome, it became clear

that the high genetic variation within the mapped CSD locus likely resulted from several highly

differentiated haplotypes in this region. To investigate this further, we sequenced whole genomes

of haploid males from clonal lines other than A and B. We assembled the genomes of each

haploid male de novo using SPAdes (Bankevich et al. 2012; Prjibelski et al. 2020), found the

contig bearing homology to the mapped CSD locus in the reference genome assembly using

blastn (Camacho et al. 2009), manually scaffolded contigs as needed, and pairwise aligned each

alternate allele to the reference genome using nucmer and mummer (Kurtz et al. 2004) to identify

differences between alleles.

RNAseq for Annotation Improvements

To avoid missing any previously unannotated genes in the mapped CSD locus region, we

performed RNA sequencing following various approaches. To start, we extracted total RNA

from multiple samples, including separate pooled tissue from several different life stages (eggs,

young larvae, fourth instar larvae, prepupae, pupae, and adults), different adult tissues (antennae,

heads, thoraces, legs, abdomens), and individual male and female pupae using a TriZol RNA

extraction method followed by precipitation in isopropanol at -20C overnight. We used the

different life stage extractions for long read RNA-seq, rRNA depletion RNA-seq, and small

RNA-seq. We used the different adult tissues for long read RNA-seq, and the individual male

and female pupae for mRNA-seq. Long read RNA-seq was performed on a PacBio Sequel II,

and all other sequencing was performed on an Illumina NovaSeq 6000. For rRNA depletion, we

used Illumina's TruSeq Stranded Total RNA RiboZero kit. For small RNA-seq, we prepared

libraries using Perkin Elmer's NEXTFLEX Small RNA-Seq Kit v3 for Illumina Platforms.

Annotation Improvements – Long Read Transcriptome Assembly

The GenBank reference genome (GCA_003672135) for *O. biroi* (Obir_v5.4) was

retrieved from NCBI in FASTA format alongside the GenBank (GCA_003672135) and RefSeq

(GCF_003672135.1) transcriptomes in GTF format. Additionally, a curated RefSeq annotation

was retrieved from Zenodo (https://doi.org/10.5281/zenodo.10079884) in GTF format. RefSeq

assemblies were chromosome name-mapped to GenBank chromosome names using the

Bioconductor GenomeInfoDb package (Arora et al. 2024) and the chromosome name maps

available in NCBI.

The PacBio toolset's pbmm2 (https://github.com/PacificBiosciences/pbmm2), a PacBio

wrapper for Minimap2 (Li 2018), was used for indexing and alignment of pooled full-length,

non-concatemer long reads to the GenBank reference genome using the parameters "--preset

ISOSEQ -j 8". Following the generation of aligned long reads in BAM format, Stringtie2

(Kovaka et al. 2019) was used with parameters "-L" for long read transcriptome assembly, and

the curated RefSeq annotation was supplied as a guide for assembly using the "-G" parameter.

The resulting gene models were exported for further analysis and integration as a gzipped GTF

file.

Annotation Improvements – Short Read Transcriptome Assembly

For Illumina RNA-seq data, FastQ quality control was performed using Rfastp (Wang

and Carroll 2024). Indexing and alignment of short reads to the GenBank genome was performed

using Hisat2 (Kim et al. 2019) with default parameters, and samtools was used to generate sorted

and indexed BAM files for further analysis. Stringtie2 was then used to assemble transcriptomes

using the default parameters. Gene models were then combined across replicates using

Stringtie2's merge function with parameters "-F 0.5 -T 2 -f 0.1" and were exported for further

analysis and integration as a gzipped GTF file.

Annotation Improvements – Integrating Short and Long Read Transcriptomes with the RefSeq

Assembly.

Integration of long read and short read generated gene models with the established and

curated RefSeq transcriptome was performed using custom scripts in R available from GitHub

(https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025) and described here.

First, conflicting gene models where single long read genes map to multiple RefSeq

genes and vice versa were identified. As generated long read gene models are supported by

multiple full-length, non-concatemer long read alignments, attention was focused on the 720

instances of multiple long read generated genes overlapping single RefSeq genes.

Single exon long read genes were removed as these are often found to be artifacts from

PacBio sequencing. Following this, long read gene models that split RefSeq gene models but

matched with GenBank gene models were accepted with the corresponding RefSeq models

rejected.

Next, RefSeq gene models that conflicted within themselves were removed. This includes

genes that were overlapping in their exons but marked as separate genes.

After tidying RefSeq gene models, long read gene models that split RefSeq gene models

but matched with short read gene models were accepted, with the corresponding RefSeq models

rejected. After this gene model polishing, only 132 gene models were left where multiple long

read generated genes overlapped single RefSeq genes.

To potentially rescue internally primed transcripts caused by long stretches of genomic

polyAs, long read transcripts were assessed for polyA nucleotide content downstream from their

endpoints, and these distributions were visualized to identify cut-offs for putative internally

primed transcripts. Transcripts with at least 70 percent as 10bp downstream of the last exon or

with a polyA stretch > 4 in 10bp of the last exons were labeled as putative internally primed

transcripts. For putative internally primed transcripts, which also split RefSeq genes, short read

gene models were used to guide the rescued models when both short read and RefSeq agreed on

a single gene model.

Finally, for the remaining long read gene models that split RefSeq gene models, gaps

between long read gene models were assessed in the short read and RefSeq gene models. If

agreement in exon and junction structure was found between short read and RefSeq gene models,

the long read gene models were rejected. This left a final 27 genes where long read gene models

split RefSeq gene models, and these were manually assessed in IGV alongside short read and

long read alignments to reject or accept.

With a final set of long read gene models, all RefSeq and short read transcripts that fell

within single genes were then added to the long read transcriptome to create an integrated

transcriptome. Duplicate transcripts were then removed from the integrated transcriptome, and

transcripts and genes were annotated for their sources (long read, short read, RefSeq) and any

overlap with the established RefSeq, GenBank, and Ensembl gene models included. Following

the creation of the integrated gene models, SQANTI3 (Pardo-Palacios et al. 2024) was used to

provide a final corrected set of gene models as well as additional QC, as described below.

Assessment of Annotation Quality and Identifying Novel Genes, Transcripts and Exons

SQANTI3 was run with default parameters on the integrated transcriptome against the

curated RefSeq and GenBank gene models to generate QC and descriptive statistics on novel and

non-canonical splicing events. Gffcompare software (Pertea and Pertea 2020) was run against the

curated RefSeq and GenBank gene models to provide statistics on novel exons, transcripts, and

genes.

BUSCO (Simão et al. 2015) was run on the integrated, curated RefSeq and GenBank

gene models with the parameters "-l insecta_odb10 -m tran" to provide measures of

transcriptome completeness based on a set of core gene models. All these metrics were

summarized into tables using custom R scripts available on GitHub

(https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025).

<u>Annotation of CDS and Functional Annotation of Genes</u>

Annotation of coding regions was performed using TransDecoder (Haas 2023) to add

CDS information to our integrated gene models GTF. TransDecoder was run with default

parameters except for providing the "--single_best_only" to the TransDecoder.Predict function to

get only one prediction per transcript. CDS information was merged with SQANTI3's GeneMark

CDS annotation for a final set of CDSs. With this coding information, gene models were read

into R, translated into protein sequences using the BioStrings package, and protein per transcript

sequences exported as FASTA format for use in downstream tools. For functional annotation,

both eggNOG (Huerta-Cepas et al. 2019) and InterProScan (Jones et al. 2014) were used using

default parameters with our protein sequences FASTA to provide predictions for functional

annotations of genes including their putative GO, KEGG, PFAM and Panther categories.

Annotations were summarized to gene levels from transcript level annotation using R.

<u>Annotation of lncRNAs</u>

FEELnc (Wucher et al. 2017) was used to identify putative lncRNAs within our final

integrated gene models. First, the FEELnc_filter.pl script was run with default parameters, and

then the FEELnc_codpot.pl script was run with parameters "--mode=shuffle" to generate our

putative set of lncRNAs for annotation of our integrated gene model GTF.

<u>Annotation of miRNAs</u>

Small RNA-seq reads were adaptor clipped for a "TGGAATTCTCGGGTGCCAAGG" adapter

using CLIPflexR (Rozen-Gagnon et al. 2021) and aligned to the genome using miRDeep2's

mapper.pl script with parameters "-c -j -m -l 18" (Friedländer et al. 2012). Aligned data were

then processed using miRDeep2's miRDeep2.pl script with a FASTA of known *Drosophila*

miRNAs retrieved from miRBase (Kozomara et al. 2019). The resulting annotated miRNA

genomic locations were merged across samples and added into our integrated gene model GTF.

Annotation of piRNAs

For the identification of piRNA clusters, small read RNA-seq was processed using the pipeline

workflow described within the Pilfer (Ray and Pandey 2018) toolkit to generate piRNA cluster

BED files per sample. The sample piRNA clusters were then merged across samples to produce a

final piRNA cluster set, which was then included in our integrated gene model GTF.

Annotation of tRNAs, snRNAs, snoRNAs and rRNAs

Additional annotations for tRNAs, snRNAs, snoRNAs and rRNAs were mapped from RefSeq

annotation onto our integrated gene model GTF.

Sex-specific Gene Expression

To identify genes with sex-specific expression, we aligned Illumina mRNA-seq reads

from three female and three male pupae using STAR (v2.5.0a) (Dobin et al. 2013). We counted

and assigned reads to genes and transcripts using RSEM (v1.2.28) (Li and Dewey 2011). We

analyzed differential gene expression using DESeq2 (Love et al. 2014), filtering out genes

shorter than ten bases or with less than ten counts across all six samples. We hierarchically

clustered the differentially expressed genes by Euclidian distance using the R package pheatmap

(v1.0.12) (Kolde 2024). We used DEXSeq (v1.50.0) (Anders et al. 2012) to identify differential

exon usage.

Data and Software Availability

All DNA and RNA sequencing data are publicly available at the National Center for

Biotechnology Information Sequence Read Archive under accession number PRJNA1075055.

All other data are available in the article and Supporting Information. All code is available on

GitHub (genome annotation:

https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025; differential gene

expression and exon usage: https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq;

all other code and analyses: https://github.com/kipdlacy/SexDetermination_Obiroi).

Contributions

K.D.L., J.L., and D.J.C.K. designed research; J.L. analyzed data and visualized results for gene-

expression and exon usage analyses; K.R.G., W.W., and T.C. improved the genome annotation

with input from K.D.L.; K.D.L. performed research, analyzed data, and visualized results for all

other parts of the manuscript; K.D.L. and D.J.C.K. wrote the manuscript; D.J.C.K. supervised the

project.

Competing interests

The authors declare no competing interests.

Acknowledgements

We thank Stephany Valdés-Rodríguez, Leonora Olivos-Cisneros, Alejandra Hurtado-Giraldo,

Sean McKenzie, and other members of the Kronauer laboratory for helping us obtain male ants

and for valuable discussions; Jen Balacco and the Rockefeller University Reference Genome

Resource Center for long read RNA sequencing; the Weill Cornell Genomics Resources Core

Facility for Small RNA sequencing; Connie Zhao and the Rockefeller University Genomics

Resource Center for rRNA-depletion sequencing and all other Illumina sequencing; Tom Kay,

Matteo Rossi, and Yukina Chiba for valuable comments on an earlier version of the manuscript.

This work was supported by a Gabrielle H. Reem and Herbert J. Kayden Early-Career

Innovation Award and the National Institute of General Medical Sciences of the National

Institutes of Health under award no. R35GM127007, both to D.J.C.K. The content is solely the

responsibility of the authors and does not necessarily represent the official views of the National

Institutes of Health. This work was also supported by the Howard Hughes Medical Institute,

where D.J.C.K. is an Investigator. This is Clonal Raider Ant Project paper number 36.

716
717 **References**
718

Anders S, Reyes A, Huber W. 2012. Detecting differential usage of exons from RNA-seq data.
Genome Research 22:2008–2017.

Arora S, Morgan M, Carlson M, Pagès H. 2024. GenomeInfoDb: Utilities for manipulating
chromosome names, including modifying them to follow a particular naming style.

Bachtrog D, Mank JE, Peichel CL, Kirkpatrick M, Otto SP, Ashman TL, Hahn MW, Kitano J,
Mayrose I, Ming R, et al. 2014. Sex determination: why so many ways of doing it? *PLoS*
Biology 12:e1001899.

Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko
SI, Pham S, Prjibelski AD, et al. 2012. SPAdes: a new genome assembly algorithm and
its applications to single-cell sequencing. *Journal of Computational Biology* 19:455–477.

Beye M, Hasselmann M, Fondrk MK, Page RE, Omholt SW. 2003. The gene *csd* is the primary
signal for sexual development in the honeybee and encodes an SR-type protein. *Cell*
114:419–429.

Beye M, Seelmann C, Gempe T, Hasselmann M, Vekemans X, Fondrk MK, Page RE. 2013.
Gradual molecular evolution of a sex determination switch through incomplete
penetrance of femaleness. *Current Biology* 23:2559–2564.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics* 30:2114–2120.

Bopp D;, Saccone G;, Beye M. 2014. Sex determination in insects: variations on a common
theme. *Sexual Development* 8:20–28.

Borowiec ML, Rabeling C, Brady SG, Fisher BL, Schultz TR, Ward PS. 2019. Compositional
heterogeneity and outgroup choice influence the internal phylogeny of the ants.
Molecular Phylogenetics and Evolution 134:111–121.

Borowiec ML, Zhang YM, Neves K, Ramalho MO, Fisher BL, Lucky A, Moreau CS. 2025.
Evaluating UCE data adequacy and integrating uncertainty in a comprehensive
phylogeny of ants. *Systematic Biology*:syaf001.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009.
BLAST+: architecture and applications. *BMC Bioinformatics* 10:1–9.

Chodroff RA, Goodstadt L, Sirey TM, Oliver PL, Davies KE, Green ED, Molnár Z, Ponting CP.
2010. Long noncoding RNA genes: conservation of sequence and brain expression
among diverse amniotes. *Genome Biology* 11:R72.

Cook JM, Crozier RH. 1995. Sex determination and population biology in the Hymenoptera.
Trends in Ecology & Evolution 10:281–286.

Crozier RH. 1971. Heterozygosity and sex determination in haplo-diploidy. *The American*
Naturalist 105:399–412.

Dale RK, Pedersen BS, Quinlan AR. 2011. Pybedtools: a flexible Python library for
manipulating genomic datasets and annotations. *Bioinformatics* 27:3423–3424.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T,
McCarthy SA, Davies RM, et al. 2021. Twelve years of SAMtools and BCFtools.
GigaScience 10:giab008.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras
TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29:15–21.

Duncan IW. 2002. Transvection effects in *Drosophila*. *Annual Review of Genetics* 36:521–556.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biology* 20:238.

Friedländer MR, Mackowiak SD, Li N, Chen W, Rajewsky N. 2012. miRDeep2 accurately
identifies known and hundreds of novel microRNA genes in seven animal clades. *Nucleic*
Acids Research 40:37–52.

Gempe T, Beye M. 2011. Function and evolution of sex determination mechanisms, genes and
pathways in insects. *BioEssays* 33:52–60.

Gempe T, Hasselmann M, Schiøtt M, Hause G, Otte M, Beye M. 2009. Sex determination in
honeybees: Two separate mechanisms induce and maintain the female pathway. *PLoS*
Biology 7:e1000222.

Haas B. 2023. TransDecoder (find coding regions within transcripts). Available from:
https://github.com/TransDecoder/TransDecoder

Hasselmann M, Beye M. 2004. Signatures of selection among sex-determining alleles of the
honey bee. *Proceedings of the National Academy of Sciences* 101:4888–4893.

Hasselmann M, Gempe T, Schiøtt M, Nunes-Silva CG, Otte M, Beye M. 2008. Evidence for the
evolutionary nascence of a novel sex determination pathway in honeybees. *Nature*
454:519–522.

Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H, Mende DR,
Letunic I, Rattei T, Jensen LJ, et al. 2019. eggNOG 5.0: a hierarchical, functionally and
phylogenetically annotated orthology resource based on 5090 organisms and 2502
viruses. *Nucleic Acids Research* 47:D309–D314.

Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell
A, Nuka G, et al. 2014. InterProScan 5: genome-scale protein function classification.
Bioinformatics 30:1236–1240.

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology* 37:907–915.

Koch V, Nissen I, Schmitt BD, Beye M. 2014. Independent Evolutionary Origin of fem
Paralogous Genes and Complementary Sex Determination in Hymenopteran Insects.
PLOS ONE 9:e91883.

Kolde R. 2024. pheatmap: Pretty Heatmaps. Available from:
https://github.com/raivokolde/pheatmap

Kovaka S, Zimin AV, Pertea GM, Razaghi R, Salzberg SL, Pertea M. 2019. Transcriptome
assembly from long-read RNA-seq alignments with StringTie2. *Genome Biology* 20:278.

Kozomara A, Birgaoanu M, Griffiths-Jones S. 2019. miRBase: from microRNA sequences to
function. *Nucleic Acids Research* 47:D155–D162.

Kronauer DJC, Pierce NE, Keller L. 2012. Asexual reproduction in introduced and native
populations of the ant *Cerapachys biroi*. *Molecular Ecology* 21:5221–5235.

Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL. 2004.
Versatile and open software for comparing large genomes. *Genome Biology* 5:R12.

Lacy KD, Hart T, Kronauer DJC. 2024. Co-inheritance of recombined chromatids maintains
heterozygosity in a parthenogenetic ant. *Nature Ecology & Evolution* 8:1522–1533.

Laidlaw HH Jr, Gomes FP, Kerr WE. 1956. Estimation of the number of lethal alleles in a
panmictic population of *Apis mellifera L. Genetics* 41:179–188.

Li B, Dewey CN. 2011. RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. *BMC Bioinformatics* 12:323.

Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
Available from: http://arxiv.org/abs/1303.3997

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34:3094–
3100.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biology* 15:550.

Ma W-J, Schwander T. 2017. Patterns and mechanisms in instances of endosymbiont☐induced
parthenogenesis. *Journal of Evolutionary Biology* 30:868–888.

Matthey-Doret C, van der Kooi CJ, Jeffries DL, Bast J, Dennis AB, Vorburger C, Schwander T.
2019. Mapping of multiple complementary sex determination loci in a parasitoid wasp.
Genome Biology and Evolution 11:2954–2962.

McKenzie SK, Kronauer DJC. 2018. The genomic architecture and molecular evolution of ant
odorant receptors. *Genome Research* 28:1757–1765.

Miles A, Ralph P, Rae S, Pisupati R. 2019. cggh/scikit-allel: v1.2.1. Available from:
https://zenodo.org/records/3238280

Miyakawa MO, Mikheyev AS. 2015. QTL mapping of sex determination loci supports an
ancient pathway in ants and honey bees. *PLoS Genetics* 11:e1005656.

Miyakawa MO, Miyakawa H. 2023. *Transformer* gene regulates feminization under two
complementary sex determination loci in the ant, *Vollenhovia emeryi. Insect*
Biochemistry and Molecular Biology 156:103938.

Miyakawa MO, Tsuchida K, Miyakawa H. 2018. The doublesex gene integrates multi-locus
complementary sex determination signals in the Japanese ant, *Vollenhovia emeryi. Insect*
Biochemistry and Molecular Biology 94:42–49.

830 Narasimhan V, Danecek P, Scally A, Xue Y, Tyler-Smith C, Durbin R. 2016. BCFtools/RoH: a
831   hidden Markov model approach for detecting autozygosity from next-generation
832   sequencing data. *Bioinformatics* 32:1749–1751.

833 Normark BB. 2003. The evolution of alternative genetic systems in insects. *Annual Review of*
834   *Entomology* 48:397–423.

835 Otte M, Netschitailo O, Weidtkamp-Peters S, Seidel CAM, Beye M. 2023. Recognition of
836   polymorphic Csd proteins determines sex in the honeybee. *Science Advances* 9:eadg4239.

837 Oxley PR, Ji L, Fetter-Pruneda I, McKenzie SK, Li C, Hu H, Zhang G, Kronauer DJC. 2014.
838   The genome of the clonal raider ant *Cerapachys biroi*. *Current Biology* 24:451–458.

839 Page RE Jr. 1980. The evolution of multiple mating behavior by honey bee queens (*Apis*
840   *mellifera L.*). *Genetics* 96:263–273.

841 Pan Q, Darras H, Keller L. 2024. LncRNA gene ANTSR coordinates complementary sex
842   determination in the Argentine ant. *Science Advances* 10:eadp1532.

843 Pardo-Palacios FJ, Arzalluz-Luque A, Kondratova L, Salguero P, Mestre-Tomás J, Amorín R,
844   Estevan-Morió E, Liu T, Nanni A, McIntyre L, et al. 2024. SQANTI3: curation of long-
845   read transcriptomes for accurate identification of known and novel isoforms. *Nature*
846   *Methods* 21:793–797.

847 Pertea G, Pertea M. 2020. GFF Utilities: GffRead and GffCompare. Available from:
848   https://f1000research.com/articles/9-304

849 Poplin R, Ruano-Rubio V, DePristo MA, Fennell TJ, Carneiro MO, Auwera GAV der, Kling
850   DE, Gauthier LD, Levy-Moonshine A, Roazen D, et al. 2018. Scaling accurate genetic
851   variant discovery to tens of thousands of samples. Available from:
852   https://www.biorxiv.org/content/10.1101/201178v3

853 Privman E, Wurm Y, Keller L. 2013. Duplication and concerted evolution in a master sex
854   determiner under balancing selection. *Proceedings of the Royal Society B: Biological*
855   *Sciences* 280:20122968.

856 Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. 2020. Using SPAdes De
857   Novo Assembler. *Current Protocols in Bioinformatics* 70:e102.

858 Quinn JJ, Zhang QC, Georgiev P, Ilik IA, Akhtar A, Chang HY. 2016. Rapid evolutionary
859   turnover underlies conserved lncRNA–genome interactions. *Genes & Development*
860   30:191–207.

861 Rabeling C, Kronauer DJC. 2012. Thelytokous parthenogenesis in eusocial Hymenoptera.
862   *Annual Review of Entomology* 58:273–292.

863 Ray R, Pandey P. 2018. piRNA analysis framework from small RNA-Seq data by a novel cluster
864   prediction tool - PILFER. *Genomics* 110:355–365.

Rozen-Gagnon K, Gu M, Luna JM, Luo J-D, Yi S, Novack S, Jacobson E, Wang W, Paul MR,
Scheel TKH, et al. 2021. Argonaute-CLIP delineates versatile, functional RNAi networks
in *Aedes aegypti*, a major vector of human viruses. *Cell Host & Microbe* 29:834–848.

Sawanth SK, Gopinath G, Sambrani N, Arunkumar KP. 2016. The autoregulatory loop: A
common mechanism of regulation of key sex determining genes in insects. *J Biosci*
41:283–294.

Schmieder S, Colinet D, Poirié M. 2012. Tracing back the nascence of a new sex-determination
pathway to the ancestor of bees and ants. *Nature Communications* 3:1–7.

Seiler J, Beye M. 2024. Honeybees' novel complementary sex-determining system: function and
origin. *Trends in Genetics* 40:969–981.

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO:
assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics 31:3210–3212.

Suomalainen E, Saura A, Lokki J. 1987. Cytology and Evolution in Parthenogenesis. CRC Press

Trible W, Chandra V, Lacy KD, Limón G, McKenzie SK, Olivos-Cisneros L, Arsenault SV,
Kronauer DJC. 2023. A caste differentiation mutant elucidates the evolution of socially
parasitic ants. *Current Biology* 33:1047–1058.

Trible W, Mckenzie SK, Kronauer DJC. 2020. Globally invasive populations of the clonal raider
ant are derived from Bangladesh. *Biology Letters* 16:20200105.

Ulitsky I, Shkumatava A, Jan CH, Sive H, Bartel DP. 2011. Conserved function of lincRNAs in
vertebrate embryonic development despite rapid sequence evolution. *Cell* 147:1537–
1550.

Verhulst EC, Beukeboom LW, van de Zande L. 2010. Maternal control of haplodiploid sex
determination in the wasp *Nasonia*. *Science* 328:620–623.

Vorburger C. 2013. Thelytoky and Sex Determination in the Hymenoptera: Mutual Constraints.
Sexual Development 8:50–58.

Wang W, Carroll T. 2024. Rfastp: An ultra-fast and all-in-one fastq preprocessor (quality
control, adapter, low quality and polyX trimming) and UMI sequence parsing). Available
from: 10.18129/B9.bioc.Rfastp

Wexler J, Delaney EK, Belles X, Schal C, Wada-Katsumata A, Amicucci MJ, Kopp A. 2019.
Hemimetabolous insects elucidate the origin of sexual development via alternative
splicing. *eLife* 8:e47490.

Whiting PW. 1933. Selective fertilization and sex-determination in Hymenoptera. *Science*
78:537–538.

Whiting PW. 1935. Genic balance, sex-determination and selective fertilization in Hymenoptera.
Proceedings of the American Philosophical Society 75:517–520.

Whiting PW. 1943. Multiple alleles in complementary sex determination of *Habrobracon*.
Genetics 28:365–382.

van Wilgenburg E, Driessen G, Beukeboom LW. 2006. Single locus complementary sex
determination in Hymenoptera: an "unintelligent" design? *Frontiers in Zoology* 3:1.

Williams TM, Carroll SB. 2009. Genetic and molecular insights into the development and
evolution of sexual dimorphism. *Nature Reviews Genetics* 10:797–804.

Wucher V, Legeai F, Hédan B, Rizk G, Lagoutte L, Leeb T, Jagannathan V, Cadieu E, David A,
Lohi H, et al. 2017. FEELnc: a tool for long non-coding RNA annotation and its
application to the dog transcriptome. *Nucleic Acids Research* 45:e57.

Zou Y, Geuverink E, Beukeboom LW, Verhulst EC, van de Zande L. 2020. A chimeric gene
paternally instructs female sex determination in the haplodiploid wasp *Nasonia*. *Science*
370:1115–1118.

<u>**Fig. 1.**</u> **Theoretical expectations for complementary sex determination (CSD). (A)** Cartoon depicting

a mating between a diploid female with two different alleles at a sex determination locus and a haploid

male bearing one of these two alleles. Half of the sexually produced diploid offspring are expected to

develop as males. **(B)** Table illustrating multi-locus CSD in diploids, in which heterozygosity of at least

one sex determination locus is required for female development. By contrast, only homozygosity at all

loci results in diploid male development. **(C)** Cartoon depicting how CSD might work in asexual species

such as the clonal raider ant. If diploid males arise from losses of heterozygosity at sex loci, then

homozygosity for either of the two alleles should trigger male development. Offspring proportions reflect

empirical observations in the clonal raider ant (Kronauer et al. 2012; Oxley et al. 2014).

924
925

Fig. 2. Whole genome sequencing reveals a candidate sex determination locus on chromosome 4. (A)

Karyoplot depicting the mean CSD index p-value in 50kb windows with a 15kb sliding interval. The CSD

index peak is shown as a green dotted line. The significance threshold (p=0.05) and FDR-corrected

significance threshold (p=0.002) are indicated by gold and brown lines, respectively. The grey histogram

shows the number of ancestrally heterozygous SNPs in 300kb windows. **(B)** Stacked bar plots for all

diploids used for mapping, with one horizontal line for each ordered putatively ancestrally heterozygous

SNP on chromosome 4. For each sample, SNPs that retain ancestral heterozygosity are drawn as grey

lines, whereas SNPs that have lost ancestral heterozygosity are drawn as black lines.

	Heterozygous	Homozygous Allele 1	Homozygous Allele 2
Diploid Females	19	0	0
Diploid Males	5	2	9

934 **Table 1.** Number of samples heterozygous and homozygous for each allele in the 46kb region on

935 chromosome 4.

Fig. 3. The putative sex determination locus identified in *O. biroi* is conserved across formicoid ants.

(**A**) Karyoplot depicting the 14 chromosomes in the *O. biroi* genome, chromosome 4 from the *L. humile*

genome (Pan et al. 2024), and one contig from linkage group 14 of the *V. emeryi* reference genome

(Miyakawa and Mikheyev 2015). Note that the plots from the different species are not drawn to scale.

Homology to protein-coding genes identified within *V.emeryiCsdQTL2* is drawn in green. The location of

the *O. biroi* CSD index peak is indicated. (**B**) A phylogeny of the ant subfamilies (adapted from

(Borowiec et al. 2019)), with the presence of a sex determination locus homologous to the peak of our

CSD index, or the absence of data shown. The yellow shaded background denotes the formicoid clade.

(**C**) Karyoplot for *O. biroi* chromosome 4, depicting homology to *V.emeryiCsdQTL2* and ancestral

heterozygosity for six different clonal lines (A, B, C, D, I, and M). Grey histograms depict the number of

ancestrally heterozygous SNPs in 100kb windows.

Fig. 4. The CSD index peak is characterized by high genetic diversity in a non-coding region. (A &

B) Nucleotide diversity across the length of *O. biroi* chromosome 4 in 5kb windows (step size=1kb) **(A)**

and across the vicinity of the CSD index peak in 100bp windows (step size=20bp) **(B)**. **(C)** The number of

differences per 100bp window between each alternate *de novo* assembled allele and the reference genome

allele. **(D)** Annotated genes in the vicinity of the CSD index peak. Black boxes depict exons and thin lines

depict introns. The lncRNA *ANTSR* is indicated in bold. Arrows indicate the names of genes in close

proximity. The CSD index peak is shown as a green dotted line in A, and with green shading in B-D.

Clonal Line	CSD Index Peak Haplotypes
A	A1 / A2
B	B1 / A1
C	? / ?
D	D1 / D2
I	I1 / ?
M	M1 / A1

957 **Table 2.** The haplotypes present at the CSD index peak for each studied *O. biroi* clonal line. Because no

958 haploid males were sequenced from clonal line C, and only one haploid male was sequenced from clonal

959 line I, the haplotypes for these two lines remain incompletely known. We denote the unidentified

960 haplotypes with question marks.

Clonal Line	Haploid Males	Diploid Males
A	92	22
B	78	0
C	6	0
D	4	0
I	1	0
M	2	0

961 **Table 3.** Number of haploid and diploid males sampled from different *O. biroi* clonal lines.

Fig. 5. **Whether a second, *tra*-containing CSD QTL from *V. emeryi* is conserved across ants remains**

ambiguous. (**A**) Karyoplot depicting the 14 chromosomes in the *O. biroi* genome and one contig from

linkage group 13 of the *V. emeryi* reference genome. Homology to protein-coding genes identified within

V.emeryiCsdQTL1 is drawn in purple. The locations of the *O. biroi* CSD index peak and the homolog of

transformer are indicated. (**B**) A phylogeny of the ant subfamilies (adapted from (Borowiec et al. 2019)),

with the presence or absence of this second putative sex determination locus, or the absence of data

shown. The yellow shaded background denotes the formicoid clade. (**C**) Karyoplot for *O. biroi*

chromosome 2, depicting homology to *V.emeryiCsdQTL1* and ancestral heterozygosity for each clonal

line. Grey histograms depict the number of ancestrally heterozygous SNPs in 100kb windows. The region

of homology that is ancestrally homozygous in clonal lines A and C is labeled.

Supplementary Figures

<u>**Supplementary Fig. S1.**</u> **Genome-wide heterozygosity levels.** Scatterplot depicting the

proportion of base pairs in the ~220Mb *O. biroi* genome that are heterozygous in each individual

for which we conducted whole genome sequencing. Haploid males have no legitimate

heterozygosity—the small elevation above zero is due to heterozygous genotype calls at

erroneously assembled regions of the reference genome.

Supplementary Fig. S2. **Homozygosity levels differ between males and females at the**

putative CSD locus. Manhattan plot depicting FDR-corrected p-values from Fisher's Exact

Tests on whether homozygosity levels differ between males and females at each SNP. The red

line indicates the significance threshold (p=0.05), and the vertical green line marks the peak of

the CSD index.

<u>Supplementary Fig. S3.</u> Diploid males result from copy-neutral losses of heterozygosity for

either allele. (A) Karyoplot depicting losses (green) or retention (gray) of heterozygosity across

the vicinity of the peak of the CSD index in diploid males and females. Codes at the left (C16,

C17, etc.) indicate the stock colonies from which each ant was sampled. The magenta box

surrounds the CSD index peak. Dark green indicates that the sample has the same allele /

haplotype as the reference haploid male whereas light green indicates that the sample has the

alternative allele. Switches between alleles in homozygous regions within samples (i.e., changes

from light green to dark green or vice versa) represent recombination events that occurred in

generations prior to the meiosis that produced each diploid male. Because the reference haploid

male was from colony C16, fewer generations have elapsed since the common ancestor of the

reference haploid male and the diploid males from that colony, and thus fewer recombination

events have accumulated. By contrast, more generations have elapsed since the common ancestor

of the reference haploid male and the diploid males from colonies C17, C18, and C1, which is

reflected in the higher number of recombination events that accumulated across generations. **(B)**

Normalized read depth across either all SNPs that passed filtering on chromosome 4 (gray, for

samples without losses of heterozygosity on chromosome 4) or all SNPs that passed filtering

within that sample's loss of heterozygosity region (green). Box plots show median (center line),

interquartile range (IQR) (box limits) and 1.5 x IQR (whiskers). Data points that fall outside 1.5

x IQR are shown individually. Violin plots show the kernel probability density, meaning that the

proportion of the data located at an x-axis value is represented by the width of the outlined area.

1008

1009 <u>**Supplementary Fig. S4. Genome-wide nucleotide diversity.**</u> Nucleotide diversity across the

1010 genome in 5kb windows with a 1kb sliding interval. In the genome-wide plot (top), the red line

1011 depicts the peak of the CSD index from genetic mapping. In the zoomed-in plots for

1012 chromosomes 4 (middle) and 2 (bottom), the green and purple vertical shading indicates

1013 homology to *V. emeryi* CSD QTL 2 and 1, respectively.

1014

Supplementary Fig. S5. Sex-specific gene expression. (**A**) Heatmap of genes that were

significantly differentially expressed between sexes (adjusted p-value < 0.05). Each row

represents a gene, and each column represents a sample. The trees at the left and the top of the

heat map depict hierarchical clustering of genes and samples, respectively. Z-scores of log-

transformed expression levels are shown as a color gradient. (**B**) Volcano plot depicting

differential gene expression between males and females. Each gene is represented by a dot,

which is colored gray if not differentially expressed, green if the expression difference is at least

2-fold ($\log \geq 1$), or blue if the expression difference is statistically significant (adjusted p-value

< 0.05). Red indicates genes that are significantly differentially expressed and have at least a 2-

fold difference in gene expression. (**C**) Expression levels of genes in the vicinity of the CSD

index peak. Normalized counts are shown for each male and female sample, with error bars

depicting the mean and SEM for each sex. The genomic coordinates, name, and ID are given for

each gene. Genes with significantly different expression between sexes (adjusted p-value < 0.05)

are bolded in the table and marked with an asterisk on the plot. The CSD index peak is shown in

green, and the dashed box denotes the peak of nuclear diversity.

1030

Supplementary Fig. S6. RNAseq reads aligned to the unannotated lncRNA flanking the

putative CSD locus. RNAseq read depth in the vicinity of the unannotated lncRNA in nine

different libraries. The peak of the CSD index is indicated in light green shading, but note that

1034   this region extends further downstream. The nucleotide diversity peak is indicated by a

1035   horizontal green bar.

1036

1037

Supplementary Fig. S7. The *O. biroi* transformer homolog is sex-specifically spliced. The c-

terminal region of *tra* exon four is spliced differently in females (pink) and males (blue),

producing a truncated protein in males and a functional, full-length TRA protein in females. (**A**)

Diagram of transformer splicing depicting full-length (functional) and truncated (non-functional)

transcripts. The black arrow indicates the transcription start site, the pink line indicates the

female-specific splice junction that allows the production of functional protein, and the black line

indicates the splice junction that leads to a truncated protein. (**B**) The number of reads spanning

the female splice junction required to produce functional protein from three female and three

male short-read RNAseq libraries. (**C**) The number of full-length and truncated *tra* transcripts in

six long read RNA sequencing (IsoSeq) libraries from different tissues and sexes.

1048   **Supplementary Tables**

Primer Name	5.4 Scaffold	5.4 Position	Forward 5'-3'	Reverse 5'-3'	Genotyping Method
Obir Ploidy 12SV	Chr12	12801234	TAGATCGAGT GAGGTATCTT	AGCGTTAA GAGTAGAT TAAG	Visualize differently sized alleles (i.e., different versions of a structural variant [SV]) using gel electrophoresis
Obir Ploidy 4RE	Chr4	5840887	CCACCCTCTC TCTCGTCTC	TGCCGATT CATATTTC GAGATAAA C	Separately digest products with the restriction enzymes (REs) EcoRI and BsmI and visualize genotypes via gel electrophoresis
Obir Ploidy 5RE	Chr5	1963279	GCGATAGAA GAGTGCTCGT ATT	GCCAGTCA GTCTGTCA TTCA	Separately digest products with the restriction enzymes (REs) EcoRI and AvaII and visualize genotypes via gel electrophoresis
Obir Ploidy 2SNP	Chr2	18082503	CACGTACAAT GAGGGAACC TAA	TCGCGATC ATGAAATC TAACCT	Sanger sequence and call SNPs
Obir Ploidy 12SNP	Chr12	8100152	GTCAGTGCGC TGTTCATTTC	AGATTATA GGGCGTTC CAATTCT	Sanger sequence and call SNPs
Obir Ploidy 14SNP	Chr14	6377499	CACGTGAACG CTACGATGAA	GGGTTGAG GATGTCTA AGGTAATG	Sanger sequence and call SNPs
Obir Ploidy 13SNPa	Chr13	6262370	CGTTACACTC CTGTGTCGAA TC	GCACGTAG CATTCTCTC TTTCT	Sanger sequence and call SNPs
Obir Ploidy 13SNPb	Chr13	7173724	GCCATACATC CTGCACTCTT AC	CGACGAGC ATGCGTCT AAA	Sanger sequence and call SNPs

1049   **Supplementary Table S1. Primers for ploidy assessment via heterozygosity.** 5.4 refers to the

1050   reference genome version (GCA_003672135.1).

Sample name	Library prep method	Clonal line	Stock colony	Sample type	Mean read depth	BioProject
LineA_DiploidFemale_1	Nextera Flex	A	C16	Diploid Female	30.1	PRJNA923657
LineA_DiploidFemale_2	Nextera Flex	A	C16	Diploid Female	27.4	PRJNA923657
LineB_DiploidFemale_1	Nextera Flex	B	STC6	Diploid Female	28.3	PRJNA923657
LineA_DiploidFemale_6	Nextera Flex	A	C18	Diploid Female	30	PRJNA923657
LineA_DiploidFemale_5	Nextera Flex	A	C1	Diploid Female	41.5	PRJNA923657
LineA_DiploidFemale_3	Nextera Flex	A	C16	Diploid Female	26.6	PRJNA947942
LineA_DiploidFemale_4	Nextera Flex	A	C16	Diploid Female	24.1	PRJNA947942
LineB_DiploidFemale_2	Nextera Flex	B	STC6	Diploid Female	27	PRJNA947942
LineB_DiploidFemale_3	Nextera Flex	B	STC6	Diploid Female	47.1	PRJNA947942
LineA_HaploidMale_1	Nextera Flex	A	C16	Haploid Male	26.1	PRJNA947942
LineA_HaploidMale_2	Nextera Flex	A	C16	Haploid Male	26.1	PRJNA947942
LineA_HaploidMale_3	Nextera Flex	A	C16	Haploid Male	27.6	PRJNA947942
LineA_HaploidMale_4	Nextera Flex	A	C16	Haploid Male	25.8	PRJNA947942
LineB_HaploidMale_1	Nextera Flex	B	STC6	Haploid Male	26.3	PRJNA947942
LineB_HaploidMale_2	Nextera Flex	B	STC6	Haploid Male	27.6	PRJNA947942
LineB_HaploidMale_3	Nextera Flex	B	STC6	Haploid Male	31.7	PRJNA947942
LineB_DiploidFemale_5	Nextera Flex	B	STC12	Diploid Female	42.4	PRJNA947942
LineB_DiploidFemale_4	Nextera Flex	B	STC1	Diploid Female	31.7	PRJNA947942
LineA_HaploidMale_5	Nextera Flex	A	C1	Haploid Male	30.3	PRJNA947942
LineA_HaploidMale_6	Nextera Flex	A	C18	Haploid Male	25.3	PRJNA947942
LineB_HaploidMale_4	Nextera Flex	B	STC1	Haploid Male	31	PRJNA947942
LineB_HaploidMale_5	Nextera	B	STC12	Haploid	32.8	PRJNA947

	Flex			Male		942
DiploidMale_1	Nextera Flex	A	C16	Diploid Male	57.9	PRJNA107 5055
DiploidMale_2	Nextera Flex	A	C16	Diploid Male	32.7	PRJNA107 5055
DiploidMale_3	Nextera Flex	A	C16	Diploid Male	31.5	PRJNA107 5055
DiploidMale_4	Nextera Flex	A	C18	Diploid Male	38.3	PRJNA107 5055
DiploidMale_5	Nextera Flex	A	C17	Diploid Male	47.7	PRJNA107 5055
DiploidMale_6	Nextera Flex	A	C16	Diploid Male	28.4	PRJNA107 5055
DiploidMale_7	Nextera Flex	A	C16	Diploid Male	32.0	PRJNA107 5055
DiploidMale_8	Nextera Flex	A	C2	Diploid Male	30.2	PRJNA107 5055
DiploidMale_9	Nextera Flex	A	C16	Diploid Male	41.6	PRJNA107 5055
DiploidMale_10	Nextera Flex	A	C16	Diploid Male	28.1	PRJNA107 5055
DiploidMale_11	Nextera Flex	A	C18	Diploid Male	40.2	PRJNA107 5055
DiploidMale_12	Nextera Flex	A	C16	Diploid Male	38.5	PRJNA107 5055
DiploidMale_13	Nextera Flex	A	C16	Diploid Male	35.4	PRJNA107 5055
DiploidMale_14	Nextera Flex	A	C16	Diploid Male	34.4	PRJNA107 5055
DiploidMale_15	Nextera Flex	A	C1	Diploid Male	31.5	PRJNA107 5055
DiploidMale_16	Nextera Flex	A	C1	Diploid Male	32.2	PRJNA107 5055
LineC_DiploidFemale	Nextera Flex	C	?	Diploid Female	39.4	PRJNA107 5055
LineD_DiploidFemale_2	Nextera Flex	D	BG9	Diploid Female	35.2	PRJNA107 5055
LineD_DiploidFemale_1	Nextera Flex	D	?	Diploid Female	36.9	PRJNA107 5055
LineI_DiploidFemale	Nextera Flex	I	?	Diploid Female	36.1	PRJNA107 5055
LineM_DiploidFemale	Nextera Flex	M	?	Diploid Female	32.3	PRJNA107 5055
LineB_HaploidMale	Nextera Flex	B	?	Haploid Male	28.3	PRJNA107 5055
LineC_HaploidMale	Nextera	C	C9	Haploid	27.6	PRJNA107

	Flex			Male		5055
LineD_HaploidMale_1	Nextera Flex	D	BG7	Haploid Male	25.2	PRJNA107 5055
LineD_HaploidMale_2	Nextera Flex	D	BG9	Haploid Male	34.9	PRJNA107 5055
LineI_HaploidMale	Nextera Flex	I	BG2	Haploid Male	21.2	PRJNA107 5055
LineM_HaploidMale_1	Nextera Flex	M	BG14	Haploid Male	70.7	PRJNA107 5055
LineM_HaploidMale_2	Nextera Flex	M	BG14	Haploid Male	34.0	PRJNA107 5055
LineB_DiploidFemale	Nextera Flex	B	STC6	Diploid Female	42.8	PRJNA107 5055
ShortReadPedigree1_Dau ghter	Nextera Flex	B	STC6 – dsRed	Diploid Female	39.1	PRJNA947 942
ShortReadPedigree1_Mot her	Nextera Flex	B	STC6 – dsRed	Diploid Female	41.4	PRJNA947 942
ShortReadPedigree2_Dau ghter	Nextera Flex	B	STC6 – dsRed	Diploid Female	42	PRJNA947 942
ShortReadPedigree2_Mot her	Nextera Flex	B	STC6 – dsRed	Diploid Female	42.7	PRJNA947 942
LinkedReadPedigree2_Sis ter_1	TELL-Seq	B	STC6 – dsRed	Diploid Female	74.7	PRJNA947 942
LinkedReadPedigree2_Sis ter_2	TELL-Seq	B	STC6 – dsRed	Diploid Female	196.7	PRJNA947 942
LinkedReadPedigree1_M other	TELL-Seq	B	STC6 – dsRed	Diploid Female	221.5	PRJNA947 942
LinkedReadPedigree1_Da ughter	TELL-Seq	B	STC6 – dsRed	Diploid Female	90	PRJNA947 942
LinkedReadPedigree3_Da ughter_1	TELL-Seq	B	STC6 – dsRed	Diploid Female	73.6	PRJNA947 942
LinkedReadPedigree3_M other	TELL-Seq	B	STC6 – dsRed	Diploid Female	50.1	PRJNA947 942
LinkedReadPedigree3_Da ughter_2	TELL-Seq	B	STC6 – dsRed	Diploid Female	100.8	PRJNA947 942
LinkedReadPedigree4_M other	TELL-Seq	B	STC6 – dsRed	Diploid Female	92.4	PRJNA947 942
LinkedReadPedigree4_Da ughter_1	TELL-Seq	B	STC6 – dsRed	Diploid Female	91.1	PRJNA947 942
LinkedReadPedigree4_Da ughter_2	TELL-Seq	B	STC6 – dsRed	Diploid Female	81	PRJNA947 942
LinkedReadPedigree5_Sis ter_1	TELL-Seq	B	STC6 – dsRed	Diploid Female	75.5	PRJNA947 942
LinkedReadPedigree5_Sis ter_2	TELL-Seq	B	STC6 – dsRed	Diploid Female	70	PRJNA947 942
LinkedReadPedigree6_Sis	TELL-	B	STC6 –	Diploid	77.6	PRJNA947

ter_1	Seq		dsRed	Female		942
LinkedReadPedigree6_Sister_2	TELL-Seq	B	STC6 – dsRed	Diploid Female	69.3	PRJNA947942
LinkedReadPedigree7_Sister_1	TELL-Seq	B	STC6 - dsRed	Diploid Female	91.9	PRJNA947942
LinkedReadPedigree7_Sister_2	TELL-Seq	B	STC6 - dsRed	Diploid Female	82.3	PRJNA947942

1051 **<u>Supplementary Table S2.</u> Metadata for all DNA whole-genome shotgun sequencing**

1052 **libraries included in this study.** Question marks indicate male samples for which the clonal line

1053 is known, but the stock colony of origin was not recorded.

1054

L. fabarum CSD "region"	L. fabarum Gene ID	O. biroi homolog	O. biroi genomic locus
2	LF001188	XP_026823582.1	Chr2:8902488-8903163
2	LF001187	XP_011341081.1	Chr11:370661-370885
2	LF001186	XP_011337689.1	Chr9:12701112-12701246
2	LF001185	XP_011337690.1	Chr9:12701868-12702623
2	LF001185	XP_011337690.1	Chr9:12701868-12702623
2	LF001184	XP_019887284.1	Chr9:12784516-12784735
2	LF001183	XP_011337709.1	Chr9:12786107-12786502
2	LF001182	XP_011337688.2	Chr9:12691705-12691868
2	LF001182	XP_011337688.2	Chr9:12691705-12691868
2	LF001178	XP_011342643.1	Chr4:17418475-17418687
2	LF001175	XP_026825191.1	Chr4:17449644-17450001
2	LF001174	XP_011345175.1	Chr4:2098225-2098482
2	LF001173	XP_011351264.2	Chr4:17451378-17451779
2	LF001172	XP_011348371.2	Chr3:4120987-4121232
2	LF001171	XP_019886864.2	Chr4:13108487-13108728
2	LF001170	XP_011335781.1	Chr4:13104605-13104701
2	LF001169	XP_026824725.1	Chr4:13096987-13097135
2	LF001169	XP_026824725.1	Chr4:13096987-13097135
2	LF001168	XP_011334862.1	Chr1:5429156-5429315
2	LF001167	XP_011346080.1	Chr4:15317225-15317286
2	LF001166	XP_011346280.1	Chr4:15323952-15324443
2	LF001165	XP_011346265.1	Chr4:15325281-15325355
2	LF001164	XP_011333902.1	Chr9:13671279-13671529
2	LF001163	XP_011342394.2	Chr12:8945912-8946647
2	LF001162	XP_011347542.2	Chr5:13878906-13879057
2	LF001160	XP_011342397.1	Chr12:8898038-8898047
2	LF001159	XP_011342399.1	Chr12:8838643-8839361
2	LF001158	XP_026826742.1	Chr7:9069620-9070081
2	LF001157	XP_011336745.2	Chr7:9059981-9060079
2	LF001156	XP_011342557.1	Chr7:8937548-8938389
2	LF001155	XP_011333056.1	Chr2:8038775-8039241
2	LF001154	XP_011333053.1	Chr2:8074012-8074175
2	LF001153	XP_011333054.1	Chr2:8074146-8074195
2	LF001152	XP_011333051.1	Chr2:8149348-8149419
2	LF001151	XP_026827816.1	Chr8:9271044-9271606
2	LF001149	XP_026825880.1	Chr6:3921592-3922644
2	LF001148	XP_011338311.1	Chr7:9788316-9790754
2	LF001147	XP_011347933.1	Chr12:10190445-10190537
2	LF001146	XP_011335214.3	Chr11:13374668-13374686
2	LF001146	XP_011335214.3	Chr11:13374668-13374686
2	LF001146	XP_011335214.3	Chr11:13374668-13374686
2	LF001146	XP_011335214.3	Chr11:13374668-13374686
2	LF001146	XP_011335214.3	Chr11:13374668-13374686
2	LF001145	XP_026827302.1	Chr7:9709496-9709894
3	LF000382	XP_011346847.2	Chr1:10948255-10948488
3	LF000382	XP_011346847.2	Chr1:10948255-10948488
3	LF000383	XP_011346847.2	Chr1:10948255-10948488
3	LF000384	XP_011333915.1	Chr9:13583746-13584022
3	LF000386	XP_011338617.1	Chr2:15483180-15483410
3	LF000387	XP_011338616.2	Chr2:15484445-15484662
3	LF000390	XP_011338613.1	Chr2:15505270-15505403
3	LF000391	XP_026829310.1	Chr10:7183955-7184266
3	LF000392	XP_026826758.1	Chr7:5505525-5506433
3	LF000392	XP_026826758.1	Chr7:5505525-5506433
3	LF000393	XP_011341974.1	Chr7:5453558-5453828
3	LF000394	XP_019886900.1	Chr10:2038764-2039594
3	LF000395	XP_011349203.1	Chr4:4954193-4954244
3	LF000396	XP_011345154.1	Chr4:2157855-2158361

3	LF000397	XP_011352274.1	Chr7:203734-203802
3	LF000398	XP_026830084.1	Chr12:9934340-9935728
3	LF000399	XP_026830285.1	Chr13:8584249-8584824
3	LF000400	XP_011335143.1	Chr11:13391322-13391400
3	LF000401	XP_011336214.1	Chr11:11716747-11716790
3	LF000402	XP_011345301.1	Chr13:2249289-2249472
3	LF000403	XP_026830488.1	Chr13:2248176-2248407
3	LF000403	XP_026830488.1	Chr13:2248176-2248407
3	LF000403	XP_026830488.1	Chr13:2248176-2248407
3	LF000404	XP_011345919.1	Chr12:8096652-8097413
3	LF000405	XP_019889529.1	Chr13:9111829-9111940
3	LF000407	XP_011337549.1	Chr9:15451205-15451282
3	LF000408	XP_011351557.1	Chr5:4614020-4614156
3	LF000410	XP_011334973.2	Chr11:13498556-13498818
3	LF000411	XP_011337970.1	Chr7:12764736-12766373
3	LF002285	XP_011338781.1	Chr2:16117746-16118104
3	LF002286	XP_019887425.1	Chr2:16125070-16125404
3	LF002286	XP_019887425.1	Chr2:16125070-16125404
3	LF002287	XP_011338074.1	Chr7:12214725-12214961
3	LF002288	XP_011335962.1	Chr1:20844717-20844860
3	LF002289	XP_026823646.1	Chr2:16637198-16637380
3	LF002289	XP_026823646.1	Chr2:16637198-16637380
3	LF002289	XP_026823646.1	Chr2:16637198-16637380
3	LF002289	XP_026823646.1	Chr2:16637198-16637380
3	LF002290	XP_011336005.1	Chr1:20544946-20545045
3	LF002291	XP_026831057.1	Chr2:16812379-16813501
3	LF002292	XP_026831104.1	Chr2:16819356-16820705
3	LF002293	XP_026831104.1	Chr2:16819356-16820705
3	LF002296	XP_026828394.1	Chr9:14899993-14900040
3	LF002297	XP_019886768.2	Chr1:19222308-19222451
3	LF002297	XP_019886768.2	Chr1:19222308-19222451
3	LF002297	XP_019886768.2	Chr1:19222308-19222451
3	LF002298	XP_019889777.2	Chr7:5141809-5142057
3	LF002299	XP_019886808.1	Chr2:16884037-16884431
3	LF002300	XP_026825006.1	Chr4:2377212-2377305
3	LF002301	XP_011348694.1	Chr4:2367942-2368174
3	LF002302	XP_026825006.1	Chr4:2377212-2377305
3	LF002303	XP_011348699.1	Chr4:2386146-2386411
3	LF002304	XP_011350345.1	Chr4:8051873-8052036
3	LF002305	XP_011350347.1	Chr4:8052766-8052969
3	LF002306	XP_011337620.1	Chr9:15576650-15576680
3	LF002307	XP_011337625.1	Chr9:15576536-15576978
3	LF002308	XP_026824247.1	Chr3:10590193-10590478
3	LF002309	XP_019888231.1	Chr7:8924371-8925765
3	LF002310	XP_026824253.1	Chr3:10586163-10586300
3	LF002312	XP_026826755.1	Chr7:5509815-5510612
3	LF002313	XP_011329194.1	Chr11:847027-847118
3	LF002314	XP_011349559.2	Chr13:9386369-9386463
3	LF002315	XP_011333093.2	Chr3:9794768-9794937
3	LF002315	XP_011333093.2	Chr3:9794768-9794937
3	LF002318	XP_011333739.1	Chr3:15560244-15560328
3	LF002319	XP_011333739.1	Chr3:15560244-15560328
3	LF002321	XP_011344935.1	Chr10:9297465-9297664
3	LF002322	XP_011342759.1	Chr4:17388837-17389067
3	LF002323	XP_011333096.1	Chr3:9801594-9801632
3	LF002324	XP_026830338.1	Chr13:2378794-2378931
3	LF002325	XP_011333675.1	Chr11:9497118-9497256
3	LF002326	XP_011347567.1	Chr5:13957042-13957184
3	LF002327	XP_026830797.1	Chr14:7238938-7239145

3	LF002328	XP_011343665.1	Chr14:7321276-7321392
3	LF002328	XP_011343665.1	Chr14:7321276-7321392
3	LF002329	XP_011341939.1	Chr7:5246471-5246686
3	LF002330	XP_011347566.1	Chr5:13962033-13962259
3	LF002331	XP_026825510.1	Chr5:13964307-13964333
3	LF002332	XP_011344606.1	Chr1:22794512-22794678
3	LF002333	XP_011347655.1	Chr5:15247775-15247953
3	LF002334	XP_011341957.2	Chr7:5429102-5429536
3	LF002335	XP_011338783.2	Chr2:16112384-16112752
3	LF002336	XP_011335942.1	Chr1:21613665-21614096
3	LF002337	XP_026823940.1	Chr3:14533163-14533248
3	LF002338	XP_011333620.2	Chr3:14494369-14496705
3	LF002338	XP_011333620.2	Chr3:14494369-14496705
3	LF002340	XP_011333615.1	Chr3:14487295-14487394
3	LF002341	XP_011333610.1	Chr3:14469678-14470225
3	LF002342	XP_026831025.1	Chr1:1900527-1900742
3	LF002343	XP_011342722.1	Chr2:9443066-9443173
3	LF002344	XP_011336746.2	Chr7:9059981-9060079
3	LF002345	XP_026829935.1	Chr12:8103395-8103580
3	LF002345	XP_026829935.1	Chr12:8103395-8103580
3	LF002346	XP_011345921.1	Chr12:8107693-8108806
3	LF002347	XP_019888787.2	Chr12:8119273-8119816
3	LF002348	XP_011345923.2	Chr12:8124085-8124275
3	LF002348	XP_011345923.2	Chr12:8124085-8124275
3	LF002349	XP_026829855.1	Chr12:8157914-8158229
3	LF002349	XP_026829855.1	Chr12:8157914-8158229
3	LF002349	XP_026829855.1	Chr12:8157914-8158229
3	LF002350	XP_011345927.1	Chr12:8171391-8171651
3	LF002351	XP_011345928.2	Chr12:8172293-8172529
3	LF002353	XP_011347547.1	Chr5:13890987-13891126
3	LF002354	XP_026831106.1	Chr2:16898162-16900440
3	LF002355	XP_019890050.1	Chr10:1331247-1331374
3	LF002356	XP_011337159.1	Chr5:1700584-1700746
3	LF002357	XP_011352656.1	Chr11:2527061-2527567
3	LF002358	XP_011337758.2	Chr13:2376845-2376933
3	LF002359	XP_011342746.2	Chr4:17388313-17388616
3	LF002359	XP_011342746.2	Chr4:17388313-17388616
3	LF002360	XP_011343447.1	Chr4:17131792-17131946
3	LF002361	XP_011331147.2	Chr4:11068734-11068785
3	LF002362	XP_011345798.1	Chr4:15213481-15213677
3	LF002363	XP_011345817.1	Chr4:15215508-15215954
3	LF002364	XP_019889473.2	Chr4:15220312-15220911
3	LF002365	XP_011340809.1	Chr3:2701684-2702137
3	LF002366	XP_026824052.1	Chr3:2697959-2698230
3	LF002367	XP_026824725.1	Chr4:13096987-13097135
3	LF008641	XP_026824039.1	Chr3:9118311-9118866
4	LF005364	XP_011334504.1	Chr7:2098841-2099004
4	LF005365	XP_011342313.1	Chr12:9424093-9424190
4	LF005366	XP_011334041.1	Chr3:1349035-1349223
4	LF011889	XP_019886420.2	Chr3:10475465-10475573
5	LF003030	XP_026831077.1	Chr1:13554921-13555017
6	LF010888	XP_011344990.1	Chr10:8509474-8509767
6	LF009377	XP_011345605.2	Chr6:3471683-3472093
6	LF009377	XP_011345605.2	Chr6:3471683-3472093
6	LF009377	XP_011345605.2	Chr6:3471683-3472093
6	LF009379	XP_026828517.1	Chr9:8831304-8831370
6	LF009379	XP_026828517.1	Chr9:8831304-8831370
6	LF009379	XP_026828517.1	Chr9:8831304-8831370
6	LF009380	XP_011336797.1	Chr9:8837429-8837507

6	LF009381	XP_019887101.1	Chr9:8727410-8727519
6	LF009382	XP_026831406.1	Chr1:24511192-24511317
6	LF009382	XP_026831406.1	Chr1:24511192-24511317
6	LF009383	XP_026824186.1	Chr1:20167532-20167679
6	LF009384	XP_011342660.2	Chr7:8337474-8337620
6	LF009385	XP_011342660.2	Chr7:8337474-8337620
6	LF009386	XP_026828439.1	Chr9:8725506-8725902
6	LF009387	XP_026826104.1	Chr6:3608454-3608638
6	LF009389	XP_011345586.1	Chr6:3545693-3545764
6	LF009390	XP_011345587.1	Chr6:3542266-3542719
6	LF009391	XP_011336812.1	Chr9:9037569-9038229
6	LF009392	XP_026828409.1	Chr9:8861220-8865200
6	LF009393	XP_026829596.1	Chr11:13057235-13057309
6	LF009394	XP_011329564.1	Chr8:3674260-3674331
6	LF007431	XP_026828915.1	Chr1:10434755-10434933
6	LF007431	XP_026828915.1	Chr1:10434755-10434933
6	LF007432	XP_011331671.1	Chr9:14807644-14807839
6	LF007433	XP_011342213.1	Chr10:12611527-12611623
6	LF007434	XP_011342212.1	Chr10:12609888-12610342
6	LF007434	XP_011342212.1	Chr10:12609888-12610342
6	LF007436	XP_011337291.1	Chr2:9115215-9115236
6	LF007437	XP_026828146.1	Chr9:4587802-4588041
6	LF007438	XP_026829425.1	Chr11:6693825-6694146
6	LF007439	XP_011331274.1	Chr11:6694246-6694911
6	LF007440	XP_019886109.1	Chr11:6695786-6696124
6	LF007442	XP_011336194.1	Chr4:12380182-12380380
6	LF007443	XP_026830331.1	Chr13:9384275-9384535
6	LF007444	XP_011345852.1	Chr12:7367565-7367651
6	LF007445	XP_011335257.2	Chr3:7019336-7019555
6	LF007446	XP_026824186.1	Chr1:20167532-20167679
6	LF007446	XP_026824186.1	Chr1:20167532-20167679
6	LF007446	XP_026824186.1	Chr1:20167532-20167679
6	LF007446	XP_026824186.1	Chr1:20167532-20167679
6	LF007447	XP_026826373.1	Chr6:5911003-5911053
6	LF007448	XP_026824186.1	Chr1:20167532-20167679
6	LF007449	XP_011334216.1	Chr7:4769776-4770045
6	LF007450	XP_011349065.2	Chr12:27005-29771
6	LF007451	XP_011335419.1	Chr10:4847255-4847344
6	LF007452	XP_011335417.1	Chr10:4846815-4846859
6	LF007453	XP_026827807.1	Chr1:9089478-9089776
6	LF007455	XP_011335835.1	Chr10:2185332-2185676
6	LF007456	XP_011335834.1	Chr10:2178787-2178863
6	LF007457	XP_011329223.1	Chr11:762131-762397
6	LF007458	XP_011329224.1	Chr11:760857-760889
6	LF007459	XP_019886855.1	Chr4:13082208-13083182
6	LF007460	XP_026824731.1	Chr4:13078696-13078998
6	LF006413	XP_011335818.1	Chr10:2240022-2241763
6	LF006414	XP_011335862.1	Chr10:2135525-2135797
6	LF006415	XP_011335859.1	Chr10:2134826-2134936
6	LF006416	XP_011343295.1	Chr1:9058776-9058815
6	LF006417	XP_011343292.1	Chr1:9057219-9057283
6	LF006418	XP_011343292.1	Chr1:9057219-9057283
6	LF006419	XP_026829398.1	Chr11:2305329-2305581
6	LF006419	XP_026829398.1	Chr11:2305329-2305581
6	LF006420	XP_026829401.1	Chr11:2275861-2276530
6	LF006421	XP_011329296.1	Chr11:2271877-2272179
6	LF006422	XP_026829393.1	Chr11:7634951-7634983
6	LF006423	XP_026829706.1	Chr11:12285934-12286060
6	LF006424	XP_011344595.1	Chr1:22825831-22826081

6	LF006424	XP_011344595.1	Chr1:22825831-22826081
6	LF006425	XP_011352013.1	Chr8:6310826-6311037
6	LF006426	XP_011331632.1	Chr9:14579839-14579928
6	LF006427	XP_011352461.1	Chr7:15410331-15410681
6	LF006429	XP_011343549.1	Chr2:751822-752041
6	LF006430	XP_011349203.1	Chr4:4954193-4954244
6	LF006431	XP_011343549.1	Chr2:751822-752041
6	LF006432	XP_026827963.1	Chr9:14575261-14575420
6	LF006433	XP_011352574.1	Chr11:3070833-3070971
6	LF006433	XP_011352574.1	Chr11:3070833-3070971
6	LF006434	XP_011337828.2	Chr9:5812211-5812382
6	LF006435	XP_011349204.1	Chr4:4957779-4958419
6	LF006436	XP_011337810.1	Chr9:5882093-5882191
6	LF006437	XP_026823896.1	Chr3:805152-805214
6	LF006438	XP_011340786.1	Chr3:2598144-2598261
6	LF006439	XP_026823896.1	Chr3:805152-805214
6	LF006440	XP_011337822.1	Chr9:5860948-5861173
6	LF006440	XP_011337822.1	Chr9:5860948-5861173
6	LF006441	XP_019887296.1	Chr9:5860223-5860442
6	LF006442	XP_011331327.2	Chr11:5860356-5861681
6	LF006443	XP_026829555.1	Chr11:5855375-5855507
6	LF006444	XP_026824731.1	Chr4:13078696-13078998
6	LF004259	XP_011345653.1	Chr6:3026242-3028545
6	LF004259	XP_011345653.1	Chr6:3026242-3028545
6	LF004260	XP_011345657.1	Chr6:3023351-3023556
6	LF004261	XP_011333413.1	Chr3:11511683-11511793
6	LF004262	XP_011351420.1	Chr5:6432092-6432208
6	LF004264	XP_026826171.1	Chr6:5325828-5326190
6	LF004265	XP_011346014.1	Chr6:5325587-5325696
6	LF004266	XP_026826168.1	Chr6:5323418-5323504
6	LF004267	XP_011345652.1	Chr6:3045096-3045236
6	LF004269	XP_011345654.1	Chr6:3040496-3040563
6	LF004270	XP_011339354.1	Chr1:3930870-3931115
6	LF004271	XP_026826524.1	Chr6:3008743-3016188
6	LF004271	XP_026826524.1	Chr6:3008743-3016188
6	LF004271	XP_026826524.1	Chr6:3008743-3016188
6	LF004271	XP_026826524.1	Chr6:3008743-3016188
6	LF004272	XP_026828184.1	Chr9:13111407-13111541
6	LF004273	XP_026828185.1	Chr9:13086623-13086800
6	LF004274	XP_011333961.2	Chr9:13105804-13106034
6	LF004275	XP_011348142.2	Chr4:1434786-1435024
6	LF004276	XP_011340055.1	Chr13:7326557-7326826
6	LF004277	XP_026830287.1	Chr13:5035511-5035847
6	LF004279	XP_019887713.2	Chr13:4920080-4920324
6	LF004280	XP_011340067.2	Chr13:4897303-4901754
6	LF004280	XP_011340067.2	Chr13:4897303-4901754
6	LF004280	XP_011340067.2	Chr13:4897303-4901754
6	LF004281	XP_011340070.1	Chr13:4895933-4896176
6	LF004282	XP_011336869.1	Chr9:7349357-7349991
6	LF004283	XP_011339358.1	Chr1:3933442-3933591
6	LF004285	XP_011339356.1	Chr1:3937027-3937044
6	LF004286	XP_011339359.1	Chr1:3941264-3941427
6	LF004287	XP_011339359.1	Chr1:3941264-3941427
6	LF004288	XP_011344952.1	Chr10:9007970-9008172
6	LF004289	XP_011344954.1	Chr10:9003166-9003200
6	LF004290	XP_011344958.1	Chr10:8994570-8994686
6	LF004291	XP_011329385.1	Chr10:1633362-1633470
6	LF004292	XP_011329393.1	Chr10:1634963-1635204
6	LF004293	XP_011329393.1	Chr10:1634963-1635204

6	LF004294	XP_011329389.1	Chr10:1645838-1646248
6	LF004295	XP_026829098.1	Chr10:1647341-1647849
6	LF004296	XP_011329398.1	Chr10:1650088-1651341
6	LF004296	XP_011329398.1	Chr10:1650088-1651341
6	LF004297	XP_011337200.1	Chr7:7915393-7915548
6	LF004298	XP_026827348.1	Chr7:7916367-7916552
6	LF004299	XP_026827076.1	Chr1:7279248-7279325
6	LF004300	XP_011345863.1	Chr12:7388227-7388336
6	LF004301	XP_011345866.1	Chr12:7391418-7391629
6	LF004302	XP_011345466.1	Chr2:14533815-14534021
6	LF004303	XP_011343443.1	Chr7:10202220-10202623
6	LF004305	XP_011329969.1	Chr8:13384806-13384905
6	LF004305	XP_011329969.1	Chr8:13384806-13384905
6	LF004306	XP_026825681.1	Chr5:10051561-10051588
6	LF004307	XP_011346998.2	Chr8:14292863-14293716
6	LF004308	XP_011337232.1	Chr7:7682544-7682629
6	LF004309	XP_011331820.1	Chr9:450924-451223
6	LF004310	XP_011330274.1	Chr4:10793001-10793258
6	LF004311	XP_011345060.1	Chr10:7937737-7937849
6	LF004312	XP_026827788.1	Chr1:9083319-9083473
6	LF004312	XP_026827788.1	Chr1:9083319-9083473
6	LF004313	XP_011343304.1	Chr1:9082377-9082587
6	LF004314	XP_011330198.1	Chr4:10755815-10755993
6	LF004315	XP_011334459.1	Chr7:2283750-2284486
6	LF004316	XP_011339244.1	Chr1:3588029-3588192
6	LF004317	XP_026828329.1	Chr9:14054627-14054801
6	LF004317	XP_026828329.1	Chr9:14054627-14054801
6	LF004318	XP_011331508.2	Chr9:14061763-14061837
6	LF004319	XP_011331505.1	Chr9:14062698-14062811
6	LF004320	XP_011334857.1	Chr1:5454703-5454882
6	LF005974	XP_026828018.1	Chr1:5439229-5439530
6	LF005975	XP_011334860.2	Chr1:5434057-5434711
6	LF005976	XP_026827305.1	Chr7:8473920-8474183
6	LF005977	XP_011330739.1	Chr8:10462228-10462411
6	LF005978	XP_011330737.1	Chr8:10468312-10468396
6	LF005979	XP_011349122.1	Chr12:277094-277179
6	LF005980	XP_026824516.1	Chr3:12947714-12949000
6	LF005981	XP_011332350.1	Chr6:1628389-1628441
6	LF005982	XP_011351118.1	Chr14:2865850-2866011
6	LF005983	XP_011345061.1	Chr10:7928695-7928868
6	LF005984	XP_011345062.1	Chr10:7924312-7924663
6	LF005985	XP_011338878.1	Chr1:1486009-1486266
6	LF005986	XP_011345091.1	Chr10:7358649-7358818
6	LF005987	XP_011345085.1	Chr10:7382381-7382723
6	LF005989	XP_011345087.1	Chr10:7383338-7383519
6	LF005991	XP_019888652.1	Chr10:7645163-7645175
6	LF005992	XP_011345080.1	Chr10:7647698-7647862
6	LF005993	XP_011345081.1	Chr10:7649238-7649498
6	LF005994	XP_011335587.1	Chr10:3611073-3611430
6	LF005995	XP_011335585.1	Chr10:3617454-3617666
6	LF005996	XP_011335586.1	Chr10:3618392-3618466
6	LF005998	XP_011335579.1	Chr10:3670372-3670597
6	LF005999	XP_011335574.1	Chr10:3798053-3798466
6	LF006000	XP_011335569.1	Chr10:3800730-3800831
6	LF006002	XP_019888003.1	Chr6:2510924-2511178
6	LF006003	XP_011341632.2	Chr6:2491411-2491420
6	LF006004	XP_011341633.1	Chr6:2485102-2485164
6	LF006005	XP_026826688.1	Chr6:2483634-2483969
6	LF006006	XP_011334809.1	Chr1:5682988-5683224

6	LF006007	XP_011333725.1	Chr3:15156949-15157037
6	LF006008	XP_026827630.1	Chr1:5696215-5696343
6	LF006009	XP_011334805.1	Chr1:5696804-5696984
6	LF006011	XP_026827900.1	Chr8:3473774-3473863
6	LF006012	XP_011346986.1	Chr8:14327158-14327217
6	LF006013	XP_011346817.1	Chr1:11047741-11047827
6	LF006014	XP_011346814.2	Chr1:11042895-11042925
6	LF006015	XP_011343298.1	Chr1:9066719-9067238
6	LF006016	XP_011343299.1	Chr1:9074220-9074383
6	LF006018	XP_011330337.1	Chr11:12610357-12610821
6	LF006018	XP_011330337.1	Chr11:12610357-12610821
6	LF006019	XP_011330537.1	Chr4:10831178-10831370
6	LF006020	XP_011330547.1	Chr4:10830112-10830472
6	LF006599	XP_026824000.1	Chr3:15073019-15073218
6	LF006601	XP_011333707.2	Chr3:15069612-15069744
6	LF006602	XP_026824000.1	Chr3:15073019-15073218
6	LF006602	XP_026824000.1	Chr3:15073019-15073218
6	LF006602	XP_026824000.1	Chr3:15073019-15073218
6	LF006602	XP_026824000.1	Chr3:15073019-15073218
6	LF006602	XP_026824000.1	Chr3:15073019-15073218
6	LF006603	XP_011333707.2	Chr3:15069612-15069744
6	LF006603	XP_011333707.2	Chr3:15069612-15069744
6	LF006604	XP_011333707.2	Chr3:15069612-15069744
6	LF006604	XP_011333707.2	Chr3:15069612-15069744
6	LF006605	XP_026824000.1	Chr3:15073019-15073218
6	LF006606	XP_011331848.1	Chr9:587811-587881
6	LF006607	XP_011333716.1	Chr3:15073981-15074049
6	LF006608	XP_011329905.2	Chr8:14088136-14088231
6	LF006609	XP_011329905.2	Chr8:14088136-14088231
6	LF006611	XP_011334791.1	Chr1:5731015-5731140
6	LF006613	XP_011334796.1	Chr1:5726449-5726931
6	LF006614	XP_011332839.1	Chr6:14149691-14150234
6	LF006615	XP_011332839.1	Chr6:14149691-14150234
6	LF006616	XP_011332821.2	Chr6:14152496-14152628
6	LF006616	XP_011332821.2	Chr6:14152496-14152628
6	LF006616	XP_011332821.2	Chr6:14152496-14152628
6	LF006617	XP_011332825.1	Chr6:14184424-14184557
6	LF006618	XP_011332827.2	Chr6:14191487-14191902
6	LF006619	XP_011332863.2	Chr6:14063075-14064223
6	LF006620	XP_011332866.3	Chr6:14083471-14083668
6	LF006621	XP_011332857.1	Chr6:14084068-14084244

1055   **Supplementary Table S3.** *O. biroi* **orthologs of genes located in the putative** *L. fabarum*

1056   **CSD "regions".**

V. emeryi CSD QTL	V. emeryi GeneID	OrthoFinder O. biroi "Ortholog"	O. biroi genomic locus
2	LOC105559778	LOC105285603	Chr4:1637384-1672949
2	LOC105559779	LOC105285604	Chr4:1673043-1674418
2	LOC105559781	LOC105285607	Chr4:1674523-1676002
2	LOC105559780	LOC105285605	Chr4:1675932-1678296
2	LOC105559782	LOC105283856, LOC113561733	Chr4:1727256-1729679; Chr4:22762-24749
2	LOC105559784	LOC105283852	Chr4:1729888-1737844
2	LOC105559783	LOC105283851	Chr4:1730326-1732211
2	LOC105559785	LOC105283850	Chr4:1738076-1749057
2	LOC105559786	LOC105282554	Chr4:7069030-7075460
2	LOC105559787	LOC105282557	Chr4:7075106-7080018
2	LOC105559788	LOC105286165	Chr4:4951574-4955391
2	LOC105559790	LOC105282568	Chr4:7141680-7152824
2	LOC105559789	LOC105282568	Chr4:7141680-7152824
2	LOC105559791	LOC105283915	Chr4:5377245-5381427
2	LOC105559793	LOC105283797	Chr4:2173362-2237652
1	LOC105558826	LOC105287916	Chr2:985469-993787
1	LOC105558815	LOC105283998	Chr2:14515051-14517463
1	LOC105558816	LOC105283997, LOC113563596	Chr2:14517342-14519998; Chr2:19219509-19221771
1	LOC105558817	LOC105283996, LOC113563597	Chr2:14519947-14522109; Chr2:19217723-19219230
1	LOC105558818	LOC105283999	Chr2:14522233-14522882
1	LOC105558819	LOC105278142	Chr2:16951611-16953740
1	LOC105558820	LOC105278143	Chr2:16947572-16951531
1	LOC105558821	LOC105278141	Chr2:16939029-16945082
1	LOC105558822	LOC105287131	Chr2:12765464-12768244
1	LOC105558823	LOC105278140	Chr2:16929798-16936497

1057 **<u>Supplementary Table S4.</u>** *O. biroi* **orthologs of genes located in the two** *V. emeryi* **CSD**

1058 **QTLs.**

	Gene_type	RU	RefSeq	GenBank
Genes	guide_RNA	0	1	0
	lncRNA	1767	1430	0
	miRNA	239	0	0
	misc_RNA	0	44	0
	piRNA	23596	0	0
	protein_coding	12839	11927	13022
	pseudogene	0	373	77
	rRNA	108	109	0
	snoRNA	15	16	0
	snRNA	22	26	0
	tRNA	194	202	0
Total Genes		**38780**	**14128**	**13099**
Transcripts	lncRNA	3795	1751	0
	miRNA	239	0	0
	mRNA	63620	24215	13099
	piRNA	23596	0	0
	rRNA	108	109	0
	snoRNA	15	16	0
	snRNA	22	26	0
	tRNA	194	202	0
	guide_RNA	0	1	0
	transcript	0	818	0
Total transcripts		**91589**	**27135**	**13099**
Total multi-exon transcripts		**65882**	**25945**	**11923**

1059 **Supplementary Table S5. Improvements over previous genome annotations, with the new**

1060 **(RU) annotation featuring genes and transcripts not found in previous annotation versions.**

Gene Name	Gene ID	Log 2-Fold Change	p-value	p-adjusted
unchar. protein	LOC105285599	0.11	0.84	0.92
unchar. protein	LOC105285600	0.11	0.73	0.85
paraneoplastic antigen Ma6E	LOC105285601	0.21	0.67	0.82
ataxin-7	LOC105285602	-0.33	0.063	0.17
Kank	LOC105285603	0.05	0.82	0.91
Bet1	LOC105285604	0.08	0.54	0.72
prob. ribosome prod. factor	LOC105285607	0.34	0.12	0.27
THUMP	LOC105285605	0.19	0.13	0.28
unchar. lncRNA	LOC105285608	0.12	0.87	0.93
Creld	LOC113561733	0.56	0.77	0.88
alphaCOP	LOC105283852	-0.17	0.21	0.40
unchar. protein	LOC105283851	9.95	NA	NA
Host cell factor	LOC105283850	-0.52	2.1E-04	1.8E-03
l(2)37Cg	LOC105283849	0.86	5.7E-04	4.2E-03
ribosomal protein S2	LOC105283847	-0.06	0.50	0.69
guanine deaminase	LOC105283846	0.03	0.88	0.94
unchar. protein	LOC105283844	1.77	6.1E-08	1.0E-07

1061 **<u>Supplementary Table S6.</u> Differential gene expression near the CSD index peak.**

1062 The log 2-fold change in expression levels between male and female samples, p-values (Wald

1063 test), and adjusted p-values (Benjamini-Hochberg adjustment) for 17 genes in the vicinity of the

1064 CSD index peak. The p-values are "NA" for LOC105283850 because one of the samples is an

1065 extreme outlier detected by Cook's distance.

Gene Name	Gene ID	Exon	Log 2-Fold Change	p-value	p-adjusted
unchar. protein	LOC105285599	E001	-0.24	0.37	0.84
		E002	0.09	0.6	0.91
		E003	0.21	0.63	0.92
unchar. protein	LOC105285600	E001	-0.59	0.32	0.82
		E002	-0.23	0.78	0.96
		E003	0.51	0.15	0.72
		E004	0.03	0.87	0.98
		E005	0.06	0.73	0.95
		E006	-0.54	0.31	0.82
		E007	-0.48	0.35	0.83
		E008	0.11	0.91	0.99
		E009	-0.66	0.65	0.92
		E010	0.08	0.92	0.99
		E011	0.11	0.91	0.99
paraneoplastic antigen Ma6E	LOC105285601	E001	-0.28	0.06	0.51
		E002	0.18	0.06	0.5
ataxin-7	LOC105285602	E001	2.03	0.1	0.63
		E002	0.12	0.98	1
		E003	0.06	0.75	0.95
		E004	0.05	0.69	0.94
		E005	-0.04	0.48	0.87
		E006	-0.48	0.01	0.17
		E007	0.12	0.55	0.89
Kank	LOC105285603	E001	-0.93	0.24	0.79
		E002	-0.65	0.05	0.47
		E003	-0.14	0.72	0.94
		E004	-1.04	0.44	0.86
		E005	-1.6	0.29	0.81
		E006	-0.55	0.08	0.57
		E007	-0.67	0.04	0.4
		E008	-0.63	0.08	0.58
		E009	0.01	0.88	0.98
		E010	-0.15	0.34	0.83
		E011	-0.28	0.2	0.76
		E012	-0.27	0.2	0.76
		E013	-0.27	0.18	0.74
		E014	0.07	0.3	0.81

		E015	0.11	0.07	0.54
		E016	0.18	0.26	0.8
		E017	NA	NA	NA
Bet1	LOC105285604	E001	2.7	0.05	0.47
		E002	0	0.99	1
		E003	-0.04	0.67	0.93
		E004	-0.01	0.82	0.97
		E005	0.05	0.94	1
		E006	9.44	0.23	0.78
prob. ribosome prod. factor	LOC105285607	E001	0.24	0.77	0.96
		E002	-0.12	0.77	0.96
		E003	0.05	0.7	0.94
		E004	0	1	1
		E005	0.3	0.55	0.89
THUMP	LOC105285605	E001	1.26	0.04	0.42
		E002	-0.12	0.49	0.87
		E003	-0.09	0.55	0.89
		E004	0.03	0.79	0.96
		E005	0.13	0.19	0.75
		E006	-0.03	0.96	1
		E007	-0.09	0.76	0.95
		E008	-0.09	0.68	0.93
		E009	3.9	0.000257	0.00951
unchar. lncRNA	LOC105285608	E001	-1.46	0.08	0.56
		E002	0.28	0.06	0.5
Creld	LOC113561733	E001	-0.06	0.62	0.92
		E002	-0.01	0.86	0.98
		E003	0.09	0.32	0.82
		E004	-0.15	0.6	0.91
alphaCOP	LOC105283852	E001	-0.29	0.3	0.81
		E002	-0.23	0.33	0.83
		E003	-0.21	0.25	0.79
		E004	-0.15	0.16	0.73
		E005	0.14	0.19	0.75
		E006	0.32	0.3	0.81
		E007	0.74	0.14	0.69
unchar. protein	LOC105283851	E001	-0.07	0.76	0.96
		E002	0.14	0.76	0.96
Host cell factor	LOC105283850	E001	-0.34	0.28	0.8

		E002	-0.33	0.24	0.79
		E003	-0.25	0.38	0.84
		E004	-0.34	0.36	0.84
		E005	-0.19	0.5	0.88
		E006	-0.19	0.39	0.85
		E007	0.75	0.12	0.67
		E008	-0.23	0.33	0.82
		E009	0.3	0.39	0.85
		E010	-0.05	0.93	1
		E011	0.44	0.34	0.83
		E012	0	0.59	0.91
		E013	0	0.35	0.83
		E014	0.45	0.12	0.67
		E015	0.04	0.2	0.77
		E016	-0.35	0.56	0.9
		E017	0.14	0.07	0.55
		E018	0.11	0.48	0.87
		E019	0.35	0.38	0.84
l(2)37Cg	LOC105283849	E001	0.04	0.54	0.89
		E002	-0.03	0.6	0.91
		E003	0	0.94	1
ribosomal protein S2	LOC105283847	E001	0.1	0.54	0.89
		E002	-0.03	0.45	0.86
		E003	-0.06	0.63	0.92
		E004	3.68	0.00233	0.06
guanine deaminase	LOC105283846	E001	0.29	0.31	0.82
		E002	0.15	0.52	0.89
		E003	0.15	0.37	0.84
		E004	0.12	0.55	0.89
		E005	0.1	0.58	0.9
		E006	-0.06	0.62	0.92
		E007	-0.09	0.41	0.85
		E008	-0.17	0.18	0.75
		E009	-0.26	0.07	0.54
unchar. protein	LOC105283844	E001	-0.52	0.000763	0.02
		E002	-0.54	0.00000519	0.000311
		E003	-0.56	0.000378	0.01
		E004	-0.91	0.00693	0.13
		E005	-0.85	0.00588	0.11

		E006	-0.92	0.000107	0.00449
		E007	-0.91	0.00434	0.09
		E008	-0.8	0.00161	0.04
		E009	-0.53	0.00458	0.1
		E010	-0.54	0.0000288	0.00143
		E011	-0.46	0.000237	0.00887
		E012	0.57	7.31E-07	0.0000543
		E013	0.79	0.00000443	0.00027
		E014	1.11	1.32E-07	0.0000115
		E015	1.07	3.96E-07	0.000031
		E016	0.86	0.000705	0.0219
		E017	0.8	0.000814	0.0245
		E018	1.29	0.000153	0.00613
		E019	1.29	0.000454	0.0152
		E020	1.13	0.000177	0.00694
		E021	0.99	0.00261	0.06
		E022	-0.89	0.55	0.89
		E023	1.32	0.000135	0.00547
		E024	1.11	0.07	0.55
		E025	-0.57	0.44	0.86
		E026	-0.24	0.32	0.82
		E027	0.09	0.00973	0.165
		E028	0.21	0.0003	0.0107
		E029	-0.59	2.27E-08	0.00000229
		E030	-0.23	0.00000842	0.000479
		E031	0.51	0.00158	0.0418

1066

1067 **<u>Supplementary Table S7.</u> Differential gene expression near the CSD index peak.**

1068 The log 2-fold change in expression levels between male and female samples, p-values

1069 (likelihood ratio test), and adjusted p-values (Benjamini-Hochberg adjustment) for each exon of

1070 17 genes in the vicinity of the CSD index peak. The p-values are "NA" for one of the exons of

1071 LOC105285603 because there is negligible expression of that exon (basemean=0).

Analysis	Program	Version	Analysis Available At:
Genetic Mapping	Trimmomatic	v0.36	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	bwa	v0.7.12-r1039	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	GATK	v4.2	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	vcftools	v0.1.17	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	pyvcf	v0.6.8	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	pybedtools	v0.9.0	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	pandas	v1.1.5	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	numpy	v.1.19.5	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	bcftools	v0.1.19-96b5f2294a	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	samtools	v1.8	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	OrthoFinder	v2.5.5	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	scikit-allel	v1.3.5	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	blast	v2.5.0	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	mummer	v4.0.0rc1	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	python	v3.6.15	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	R	v4.3.2	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	karyoploteR	1.28.0	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	ggplot2	v3.5.0	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	ggbeeswarm	v0.7.2	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	matplotlib	v3.3.4	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genome Annotation	pbmm2	v1.12.0	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome	stringtie2	v2.2.1	https://github.com/RockefellerUniversity/Obiroi_Ge

Annotation			neModels_2025
Genome Annotation	samtools	v1.6	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome Annotation	hisat2	v2.2.1	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome Annotation	transdecoder	v5.7.1	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome Annotation	squanti3	v3.5.1	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome Annotation	Repeatmasker	v4.1.7-p1	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome Annotation	FEELnc	v0.2.1	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome Annotation	BUSCO	v5.4.7	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome Annotation	CLIPflexR	v0.1.20	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome Annotation	miRDeep2	v0.1.3	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome Annotation	Pilfer	v1.0	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome Annotation	R	v4.0	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome Annotation	Bioconductor	v3.19	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Differential Gene Expression	multiqc	v1.9	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	RSEM	v1.2.28	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	STAR	v2.5.0a	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	R	v4.3.3	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	GenomicFeatures	v1.52.2	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	tximport	v1.28.0	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	DESeq2	v1.40.2	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq

Differential Gene Expression	EnhancedVolcano	v1.18.0	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	pheatmap	v1.0.12	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	dplyr	v1.1.4	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	reshape2	v1.4.4	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	tidyr	v1.3.1	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	tibble	v3.2.1	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	ggbreak	v.0.1.2	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Exon Usage	HTSeq	v2.0.5	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Exon Usage	DEXSeq	v.1.46.0	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Exon Usage	stageR	v.1.22.0	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq

1072   **<u>Supplementary Table S8.</u> Versions of software used for genomics analyses.**

1073