1

Heterozygosity at a conserved candidate sex determination locus is associated

with female development in the clonal raider ant

Kip D. Lacy[1*], Jina Lee[1], Kathryn Rozen-Gagnon[2,3], Wei Wang[3], Thomas S. Carroll[3], Daniel J.C. Kronauer[1,4*]

[1]Laboratory of Social Evolution and Behavior, The Rockefeller University, New York, NY, USA.

[2]Department of Molecular Genetics, University of Toronto, Toronto, Canada.

[3]Bioinformatics Resource Center, The Rockefeller University, New York, NY, USA.

[4]Howard Hughes Medical Institute, New York, NY, USA.

[*]Email: klacy@rockefeller.edu (K.D.L.); dkronauer@rockefeller.edu (D.J.C.K)

Abstract

Sex determination is a developmental switch that triggers sex-specific developmental programs. This switch is "flipped" by the expression of genes that promote male- or female-specific development. Many lineages have evolved sex chromosomes that act as primary signals for sex determination. However, haplodiploidy (males are haploid and females are diploid), which occurs in ca. 12% of animal species, is incompatible with sex chromosomes. Haplodiploid taxa must, therefore, rely on other strategies for sex determination. One mechanism, "complementary sex determination" (CSD), uses heterozygosity as a proxy for diploidy. In CSD, heterozygosity at a sex determination locus triggers female development, while hemizygosity or homozygosity permits male development. CSD loci have been mapped in honeybees and two ant species, but we know little about their evolutionary history. Here, we investigate sex determination in the clonal raider ant, *Ooceraea biroi*. We identified a 46kb candidate CSD locus at which all females are heterozygous, but diploid males are homozygous for either allele. As expected for CSD loci, the candidate locus has more alleles than most other loci, resulting in a peak of nucleotide diversity. This peak negligibly affects the amino acid sequences of protein-coding genes, suggesting that heterozygosity of a non-coding genomic sequence triggers female development. This locus is distinct from the CSD locus in honeybees but homologous to a CSD locus mapped in two distantly related ant species, implying that this molecular mechanism has been conserved since a common ancestor that lived approximately 112 million years ago.

Introduction

Whether an animal develops as a male or a female is typically determined by a switch early in development. This switch, known as sex determination, is a multistep process beginning with primary sex determination signals triggering sex-specific splicing of downstream transcription factors that encode sexual identity (Williams and Carroll 2009). Although the downstream transcription factors are evolutionarily conserved, primary sex determination signals are evolutionarily labile and comprise a diverse range of environmental and genetic mechanisms (Bachtrog et al. 2014). Genetically encoded primary signals (sex-determining genes or chromosomes) are usually found in only one sex. Such systems do not work in haplodiploid taxa (an estimated 12% of animal species, including ants (Normark 2003)), where any allele on any homologous chromosome can be transmitted from diploid females to their haploid male sons, and all alleles experience selection to be viably transmitted from both sexes (Whiting 1935). Haplodiploids, therefore, require alternative sex determination mechanisms, but we know little about the molecular details.

Some parasitoid wasps use a mechanism involving maternal imprinting (Verhulst et al. 2010; Zou et al. 2020). However, in many Hymenoptera, inbred crosses produce diploid males, which led to the hypothesis that female development is triggered by heterozygosity as a proxy for diploidy (Whiting 1933; Whiting 1943). Under this "complementary sex determination" (CSD) hypothesis, different alleles at a given sex determination locus "complement" one another, meaning that heterozygosity triggers female development (**Fig. 1A**). By contrast, the presence of a single allele (either because the individual is haploid or because a diploid individual is homozygous at the sex determination locus) permits male development. Under CSD, diploid males produce diploid sperm and, therefore, are functionally sterile (van

Wilgenburg et al. 2006). This imposes selection for homozygosity avoidance, often meaning

obligatory outbreeding. At the gene level, there is negative frequency-dependent (balancing)

selection on CSD loci, resulting in the co-occurrence of many alleles at similar frequencies

within populations.

Having been a favored hypothesis for more than half a century, CSD was demonstrated in

the honeybee *Apis mellifera* in 2003 (Beye et al. 2003). In *A. mellifera*, biallelic heteromers of

the protein encoded by the *complementary sex determiner* (*csd*) gene trigger female

development, whereas monoallelic homomers permit male development (Beye et al. 2003; Beye

et al. 2013; Otte et al. 2023; Seiler and Beye 2024). Although this primary sex determination

signal differs from primary signals in other insects, it nonetheless converges upon the pathway

that transduces sex determination signals in most insects (Bopp et al. 2014; Wexler et al. 2019).

The Csd protein sex-specifically splices the closely linked splicing factor *feminizer* (*fem*) (the

honeybee version of *transformer* (*tra*) in *Drosophila* and other insects), which in turn leads to the

production of sex-specific isoforms of *doublesex* (*dsx*), a transcription factor that regulates sex-

specific development (Hasselmann et al. 2008; Gempe et al. 2009; Verhulst et al. 2010; Zou et

al. 2020). Intriguingly, *csd* is a paralog of *fem*. Following the prediction that CSD loci should

evolve under balancing selection, many *csd* alleles are held at similar frequencies, and the locus

bears signatures of adaptive evolution (Hasselmann and Beye 2004), with amino acid-level

heterozygosity in certain protein domains required for female development (Otte et al. 2023).

CSD is assumed to occur in many Hymenoptera that occasionally produce diploid males

(van Wilgenburg et al. 2006). Until recently, however, heterozygosity-dependent female

development had not been demonstrated outside of *A. mellifera*, which remained the only species

for which a CSD locus had been genetically mapped. A recent genetic mapping study in the

wasp *Lysiphlebus fabarum* identified at least one and as many as four candidate sex

determination loci (Matthey-Doret et al. 2019) (the multi-locus extension of the CSD hypothesis

proposes that heterozygosity at any of multiple sex determination loci is sufficient to trigger

female development (Crozier 1971) (**Fig. 1B**)). However, a requirement of heterozygosity at

these loci for female development was not demonstrated. In the ant *Vollenhovia emeryi*, a

mapping study found two sex determination QTL (Miyakawa and Mikheyev 2015), with follow-

up work suggesting that these function as a multi-locus CSD system (Miyakawa and Miyakawa

2023). One of these QTL (*V.emeryiCsdQTL1*) contains two closely linked *tra* homologs,

reminiscent of the locus containing *csd* and *fem* in *A. mellifera*. By contrast, the other QTL

(*V.emeryiCsdQTL2*) only contains genes without annotated functions in sexual development.

Although *V. emeryi* sex determination is likely integrated through *tra* and *dsx*, the molecular

basis of their primary sex determination signals remains unclear (Miyakawa et al. 2018;

Miyakawa and Miyakawa 2023).

Recently, a sex determination locus was identified in the Argentine ant, *Linepithema*

humile (Pan et al. 2024). Consistent with CSD, all pairwise heterozygous combinations of the

seven identified alleles were found in females, whereas diploid males were invariably

homozygous for one of the seven alleles. Moreover, these alleles occurred at roughly equal

frequencies within populations, suggesting evolution under balancing selection. The mapped

locus is located within a noncoding genomic region tightly linked to *ant noncoding transformer*

splicing regulator (*ANTSR*), a long noncoding RNA (lncRNA) with previously uncharacterized

function. *ANTSR* knockdown in female-destined embryos leads to male-specific splicing of *tra*,

suggesting that this lncRNA transduces the primary sex determination signal encoded by the

heterozygosity (or hemi- or homozygosity) of the linked noncoding region (Pan et al. 2024).

Because the synteny of this region (including the presence of a lncRNA putatively homologous

to *ANTSR*) is conserved across the ants, bees, and vespoid wasps, it was suggested that this

molecular mechanism may be deeply conserved among the Aculeata (Pan et al. 2024). However,

genetic mapping of sex determination has yet to be performed in most of these taxa.

To explore the evolution of sex determination across ants, we investigated the topic in the

clonal raider ant, *Ooceraea biroi*. Diploid males occur sporadically in this species, suggesting

that, like other ants, it might employ CSD (Kronauer et al. 2012). *O. biroi* reproduces via a mode

of parthenogenesis in which two haploid nuclei from a single meiosis fuse to produce diploid

offspring (central fusion automixis) (Oxley et al. 2014). Although heterozygosity is maintained

with high fidelity despite meiotic crossover recombination (Lacy et al. 2024), losses of

heterozygosity occur occasionally, probably due to the inheritance of one recombined and one

non-recombined version of a homologous chromosome (Kronauer et al. 2012; Oxley et al. 2014;

Trible et al. 2023; Lacy et al. 2024). Therefore, if *O. biroi* uses CSD, diploid males might result

from losses of heterozygosity at sex determination loci (**Fig. 1C**). Here, we use whole genome

sequencing to map a sex determination locus, demonstrate that most diploid males carry a loss of

heterozygosity at this locus, assess the homology of this locus with other sex determination loci,

and explore whether balancing selection has shaped evolution at this locus.

Results

<u>Identification of a candidate sex determination locus on chromosome 4</u>

To map sex determination loci in *O. biroi*, we first identified males based on sexual

dimorphism and distinguished diploid from haploid males based on heterozygosity at seven

genetic markers (**supplementary table S1, file S1, and fig. S1**). We then sequenced the

genomes of 16 diploid males and compared them with 19 previously sequenced genomes of

diploid females (Trible et al. 2023; Lacy et al. 2024) (descriptions and metadata for all genomes

sequenced in this study are in **supplementary table S2**). CSD loci must be heterozygous to

trigger female development but can be homozygous in diploid males. To identify such sites, we

calculated a "CSD index" for each single nucleotide polymorphism (SNP), which equals zero if

any female is homozygous and equals the proportion of diploid males that are homozygous if all

females are heterozygous. We found a peak of the CSD index on chromosome 4 (**Fig. 2A**).

To corroborate our results, we looked for evidence of segmental losses of heterozygosity

along chromosome 4. Eleven out of 16 sequenced diploid males bore segmental losses of

heterozygosity on chromosome 4 (**Fig. 2B**). The intersection of these LOH segments comprised

a 46kb peak at which all females were heterozygous, and most (11 out of 16) diploid males were

homozygous. Under the CSD hypothesis, homozygosity for any allele at CSD loci permits male

development. Consistently, *O. biroi* diploid males were homozygous for either allele at the 46kb

region on chromosome 4 (**Table 1, supplementary fig. S2**).

The stretches of homozygosity found in diploid males could have resulted from

segmental deletions resulting from improperly repaired DNA damage, or rare copy-neutral losses

of heterozygosity resulting from thelytokous parthenogenesis (Kronauer et al. 2012; Oxley et al.

2014; Trible et al. 2023; Lacy et al. 2024). To distinguish between these two scenarios, we

analyzed read depth and found that runs of homozygosity were not accompanied by changes in

copy number (**supplementary fig. S2**). This implies that diploid males arise when rare losses of

heterozygosity, which result from crossover recombination during meiosis, span the CSD locus.

Surprisingly, five of the 16 diploid males were not homozygous in this 46kb region. One

possible explanation is that these individuals had mutations in other genes involved in the sex-

determination pathway. We therefore inspected all unique mutations and losses of heterozygosity

in these five diploid males, but did not find any in genes with annotated sex determining

functions (**supplementary file S2**). Male development may have been triggered by rare

stochastic perturbations to gene expression or splicing. Alternatively, these individuals may carry

mutations or losses of heterozygosity in genes or regulatory elements with unannotated sex

determination function.

<u>The *O. biroi* CSD locus is homologous to another ant sex determination locus but not to</u>

<u>honeybee *csd*</u>

To assess homology to CSD loci mapped in other species, we identified *O. biroi*

orthologs of the genes found in the sex determination QTL identified in *V. emeryi* (Miyakawa

and Mikheyev 2015) (**supplementary table S3**). This approach also identified homology to the

L. humile CSD locus, which was expected because a subset of *V.emeryiCsdQTL2* is homologous

to the CSD identified in *L. humile* (Pan et al. 2024). Homology to *V.emeryiCsdQTL2* was

scattered across one arm of *O. biroi* chromosome 4, including the 46kb CSD index peak, and

homology to the *L. humile* CSD locus mapped within the *O. biroi* CSD index peak (**Fig. 3A**).

The *O. biroi* CSD index peak contains four protein-coding genes that are also present in

V.emeryiCsdQTL2, including *HCF*, *COPA*, an uncharacterized gene, and *CRELD2*. Notably, the

candidate CSD region in *O. biroi* also includes a large stretch of sequence with no annotated

genes downstream of an uncharacterized lncRNA. LncRNAs often retain their functions and

synteny with surrounding genes despite divergence in sequence homology (Chodroff et al. 2010;

Ulitsky et al. 2011; Quinn et al. 2016), meaning that although this lncRNA has limited sequence

homology with *L. humile ANTSR*, it may have a homologous function (Pan et al. 2024). Because

this region is implicated in CSD in *L. humile*, our data suggest that the mechanistic basis of CSD

at this locus is conserved between *O. biroi*, *L. humile*, and *V. emeryi* (**Fig. 3B**). This would mean

that this sex determination locus is ancient, as these three species belong to different ant

subfamilies that diverged roughly 112 million years ago (Borowiec et al. 2025).

We performed our genetic mapping in clonal line A, but different clonal lines of *O. biroi*

vary in which portions of the genome are ancestrally homozygous (Oxley et al. 2014). Therefore,

it was unclear whether heterozygosity at the mapped CSD locus is required for female

development in all clonal lines of *O. biroi*. To investigate this, we inspected heterozygosity in

diploid females from six different clonal lines by retrieving previously published genome

sequences from clonal line B (Lacy et al. 2024) and sequencing genomes of diploid females from

four additional clonal lines (from both the native and invasive ranges of *O. biroi* (Kronauer et al.

2012; Trible et al. 2020)). Females from all six clonal lines were heterozygous at the CSD index

peak, consistent with its putative role as a CSD locus in all *O. biroi* (**Fig. 3C**).

<u>The CSD index peak has high diversity in a non-coding region.</u>

Because CSD loci are expected to evolve under balancing selection, we expect many

different alleles to exist at those loci. For example, as many as 19 *csd* alleles were found to

segregate in honeybee populations (Hasselmann et al. 2008), and seven alleles were found to

segregate in a population of *L. humile*, despite a recent genetic bottleneck (Pan et al. 2024).

Looking for elevated genetic diversity is impossible by only studying genomes from a single

clonal line (i.e., asexual descendants of a single diploid female), so we looked across diploid

females from six different clonal lines. We calculated nucleotide diversity (π) in 5kb sliding

windows (step size = 1kb) across the genome and found a nucleotide diversity peak that fell

within the CSD index peak (**Fig. 4A, supplementary fig. S3**), between positions 1,717,00 and

1,730,000 on chromosome 4 (**Fig. 4B**).

We hypothesized that this nucleotide diversity peak resulted from many alleles found

among clonal lines of *O. biroi* at this locus. Identifying these alleles requires DNA sequences of

individual haplotypes, whereas short-read genome sequences of diploid females only provide

diploid genotypes. To investigate allelic diversity, we retrieved previously published haploid

male genome sequences from clonal lines A and B (Lacy et al. 2024) and sequenced whole

genomes of haploid males from additional clonal lines. We assembled their genomes *de novo* to

identify the different haplotypes found in this region. In addition to the haplotype found in the

reference genome, we identified six additional haplotypes (or "alleles") in the five clonal lines

for which we sequenced at least one haploid male genome (**Fig. 4C**, **Table 2**). These alleles

differ substantially in their DNA sequences in the region with high nucleotide diversity (**Fig.**

4C).

To investigate which genetic elements might act to determine sex in a heterozygosity-

dependent manner, we inspected all annotated genes in the CSD index peak (**Fig. 4D**). Within

this region, none of the protein-coding genes contained substantial heterozygosity that affected

amino acid identity (**supplementary file S3**). To determine whether any genes or exons were

missing from our annotation, we performed RNA sequencing of embryos and several other life

stages (including female and male adults) using several different technologies (see Methods).

Briefly, we used long-read RNA sequencing (PacBio IsoSeq) to extend gene models and used

several different library preparation methods followed by standard short-read (Illumina)

sequencing: standard poly-A tail library prep to capture mRNAs, ribosomal RNA depletion to

capture lncRNAs that are not poly-adenylated, and small RNA sequencing to annotate miRNAs,

endo-siRNAs, and putative piRNAs. Although this improved the *O. biroi* genome annotation by

adding new genes and improving existing gene models (**supplementary table S4, file S4**), no

new genes or exons were identified near the CSD index peak.

To look for sex-biased gene expression, we performed differential gene expression and

differential transcript usage analysis between the transcriptomes of three male and three female

white pupae. We used pupae for this experiment because this is the earliest life stage at which

males and females can be readily distinguished. Many genes were differentially expressed

between males and females (24.6% of expressed genes) or had one or more differentially used

exons, including four genes near the CSD index peak (LOC105285605, LOC105283850,

LOC105283849, and LOC105283844) (**supplementary tables S5 and S6; fig. S4**). However,

none of these overlapped with the region of elevated genetic diversity. Very few reads aligned to

the lncRNA putatively homologous to *ANTSR*, which precluded a rigorous assessment of its

differential expression in *O. biroi*.

The peak of nucleotide diversity minimally affects amino acid sequences and is primarily

located in a non-coding genomic region (**Fig. 4B-D**). Thus, a non-coding genetic element within

this region may instruct sex determination in *O. biroi* in a heterozygosity-dependent manner.

This is similar to *L. humile*, where the mapped CSD locus had high variability in non-coding and

non-exonic regions (Pan et al. 2024). RNA interference of the nearby lncRNA, *ANTSR*, led to

male-specific splicing of *tra*, raising the possibility that heterozygosity or homozygosity in the

non-exonic and non-coding *L. humile* CSD locus affects the expression level of *ANTSR*, which

instructs sex-specific splicing of *tra* (Pan et al. 2024). Because the peak of nucleotide diversity

within our CSD index peak minimally affects protein-coding sequences and the region is closely

linked to a lncRNA that is putatively an ortholog of *L. humile ANTSR*, our data suggest that the

mechanism of CSD in *O. biroi* may be conserved with *L. humile*.

O. biroi may have one or multiple sex determination loci.

250   Our CSD index mapping results suggest that *O. biroi* has a single CSD locus rather than

multiple loci. However, a curious observation raises the possibility that there may be other

"ghost" CSD loci that went undetected in our mapping study. We sampled many males from

clonal lines A and B, and a few males from four additional clonal lines. Despite this, all the

diploid males we detected came from clonal line A. We, therefore, looked back at data from

previously genotyped males and found that this pattern held across a previous study (Kronauer et

al. 2012). In total, from clonal line A, 22 out of 114 males were diploid, whereas all 78 males

genotyped from clonal line B were haploid (**Table 3**). This disparity is statistically significant

(χ^2=15.4, df=1, p<0.0001), suggesting that the two clonal lines differ in their propensity to

produce diploid males.

260   One possible explanation for this phenomenon could be that *O. biroi* has two-locus CSD

(**Fig. 1B**), but that clonal line A is ancestrally homozygous for one of the loci, whereas clonal

line B retains heterozygosity at both loci. Because multi-locus CSD requires homozygosity at all

sex determination loci for diploids to develop as males, in this scenario, two losses of

heterozygosity would be needed for diploid males to develop in clonal line B. In contrast, only

one loss of heterozygosity would be required in clonal line A, which we used for our genetic

mapping.

267   A second QTL region identified in *V. emeryi* (*V.emeryiCsdQTL1*) contains two closely

linked *tra* homologs, similar to the closely linked honeybee *tra* homologs, *csd* and *fem*. This has

led to the hypothesis that its function as a CSD locus is conserved with the *csd*-containing region

of honeybees (Miyakawa and Mikheyev 2015). However, it remains to be demonstrated that

either of these *tra* homologs acts as a primary CSD signal in *V. emeryi*. To test whether this locus

might be involved in CSD in *O. biroi*, we searched for homology to *V.emeryiCsdQTL1* in the *O.*

biroi genome and identified several regions scattered across *O. biroi* chromosome 2 (**Fig. 5A**).

However, we did not find a CSD index peak on chromosome 2, and no sex determination locus

with homology to the *V. emeryi* QTL was found in *L. humile* (Pan et al. 2024). Thus, the function

of *V.emeryiCsdQTL1* as a primary sex determination signal might have evolved after this lineage

separated from other ants (**Fig. 5B**). Alternatively, the sex determination activity of

V.emeryiCsdQTL1 could be ancestral to these three ant species and independently lost in *O. biroi*

(at least in clonal line A) and *L. humile*.

Intriguingly, while clonal lines B, D, I, and M retain heterozygosity at all sites with

homology to *V.emeryiQTL1* (**Fig. 5C**), clonal lines A and C are homozygous in a region

homologous to a portion of *V.emeryiQTL1*. We note, however, that heterozygosity in this region

is limited—it has no elevated nucleotide diversity (**supplementary fig. S3**), and there are only

two nonsynonymous heterozygous variants in this region, with clonal line B conspicuously

lacking any nonsynonymous heterozygosity. Therefore, it seems unlikely that this region bears a

CSD locus in *O. biroi*.

Discussion

Genetically mapping sex determination loci in diverse taxa is key to understanding how

sex determination evolves. Here, we investigated the mode of sex determination in the clonal

raider ant, *O. biroi*. First, we mapped a sex determination locus that was heterozygous in all

females, but homozygous for either allele in diploid males (**Fig. 2**). Next, we showed that most

diploid males bear rare losses of heterozygosity that arise during thelytokous parthenogenesis

(**Fig. 2**). We then showed that this locus is homologous to a sex determination locus found in two

other ant species—*V. emeryi* and *L. humile* (**Fig. 3**), suggesting that this mechanism of sex

determination may be conserved among formicoid ants. By sequencing whole genomes of

haploid males and diploid females from other clonal lines of *O. biroi,* we showed that the

mapped locus corresponds to a region with high genetic diversity, following the expectation that

CSD loci should evolve under balancing selection (**Fig. 4**). This diversity peaks in a noncoding

genomic region, and we found limited evidence for functional heterozygosity in nearby protein-

coding sequences, suggesting that heterozygosity in this noncoding sequence may trigger female

development. Finally, we demonstrated that *O. biroi* CSD in clonal line A does not map to the

tra-containing sex determination locus previously identified in *V. emeryi*, but it remains unclear

whether *O. biroi* ancestrally has single-locus or multi-locus CSD (**Fig. 5**).

The presence of diploid males in many ant taxa previously led to the conclusion that most

ants employ CSD (van Wilgenburg et al. 2006). Along with *L. humile* (Pan et al. 2024) and *V.*

emeryi (Miyakawa and Mikheyev 2015), *O. biroi* is now one of three ant species for which

heterozygosity-dependent female development has been demonstrated at mapped CSD loci. Our

mapped locus appears to be homologous to *V.emeryiCsdQTL2* and to the locus identified in *L.*

humile (**Fig. 3**). As was found in *L. humile*, this locus colocalizes with a peak of nucleotide

diversity (**Fig. 4**), with seven alleles found among five clonal lines (**Fig. 4C, Table 2**), consistent

with the expectation that CSD loci should harbor many alleles due to evolution under balancing

selection. This implies that the role of this locus in CSD evolved before the divergence of the

Dorylinae from the other formicoid ants roughly 112 million years ago (Borowiec et al. 2025).

Although synteny analyses raised the possibility that this locus arose early within the evolution

of the Aculeata (Pan et al. 2024), genetic mapping studies of additional representative species of

different ant subfamilies (especially in the poneroid clade and the Leptanillinae) and across other

Aculeata will be required to determine when the function of this locus in CSD originated and

how it is distributed across taxa.

In *L. humile*, a lncRNA (*ANTSR*) closely linked to this locus plays a role in sex

determination (Pan et al. 2024). We did not observe substantial expression of the putative *ANTSR*

homolog in *O. biroi*. This is likely because the rarity of males in *O. biroi* required us to sample

males at the pupal stage, well after primary sex determination signals are transduced to self-

sustaining sex-specific splicing of downstream transcription factors during the early stages of

embryonic development (Gempe and Beye 2011). Therefore, we likely missed differential gene

expression relevant to sex determination that occurs before the pupal stage.

We did not find a peak of the CSD index at the *O. biroi tra* homolog (**Fig. 5**), indicating

that, as in *L. humile*, the mode of CSD is not homologous to the system in *A. mellifera* mediated

by the *transformer* homologs *csd* and *fem*. This was previously suspected for *O. biroi* due to the

presence of only a single *tra* homolog (Oxley et al. 2014) rather than the two closely linked *tra*

homologs found in other ants (Privman et al. 2013). In *V. emeryi*, however, one of the two Csd

QTLs (*V.emeryiCsdQTL1*) spans the *tra* locus, raising the possibility that the molecular

mechanism of CSD might, in part, be conserved between *A. mellifera* and *V. emeryi* (Miyakawa

and Mikheyev 2015). It has not been demonstrated that either of *V. emeryi*'s *tra* homologs act as

a primary sex determination signal. Our results raise the possibility that the role of

V.emeryiCsdQTL1 as a primary sex determination signal evolved independently of *A. mellifera*

csd and after *V. emeryi* diverged from *L. humile*. However, further investigations into the

mechanisms of sex determination in *V. emeryi* and other ants are required to reconstruct the

evolutionary history of *V.emeryiCsdQTL1*'s function in hymenopteran sex determination.

Curiously, diploid males were only produced in clonal line A, even though many males

were collected from other clonal lines (**Table 3**). One possible explanation is that *O. biroi* has

multi-locus CSD, and clonal line A is ancestrally homozygous for all but one of the loci. Clonal

line A is ancestrally homozygous for a portion of the region homologous to the second *V. emeryi*

QTL, although it retains heterozygosity in the region overlapping *tra* (**Fig. 5C**). However, this

region lacks elevated genetic diversity and functional heterozygosity, making it a poor candidate

for a second CSD locus. Further study would be required to determine whether *O. biroi* has a

second CSD locus. However, there may be other explanations for diploid males being found

exclusively in clonal line A. For example, clonal line A could be more likely to lose

heterozygosity than other clonal lines, or diploid males in clonal line A could be more likely to

develop and survive. Currently, we have no data to distinguish between these hypotheses.

Another peculiarity of our study is that five of the 16 sequenced diploid males retained

heterozygosity at the mapped CSD locus. Stochastic gene expression or splicing fluctuations

may have caused these individuals to develop as males despite having female genotypes. Certain

allelic combinations of *A. mellifera csd* trigger female development with incomplete penetrance

(Beye et al. 2013), and the allelic combination found in clonal line A might similarly be

permissive to occasional male development. Alternatively, these diploid males could result from

mutations in or losses of heterozygosity at different genetic loci. We note that it might be easier

to find rare stochastically produced diploid males in laboratory colonies of *O. biroi* than in

sexual species. This is because genetically encoded diploid males only result from loss of

heterozygosity, which is rare in *O. biroi* (Oxley et al. 2014; Lacy et al. 2024). By contrast, in

sexually reproducing populations, diploid males homozygous for sex determination loci will

arise much more frequently due to matings between individuals that share an allele at the sex

determination locus (this effect is exacerbated in invasive species like *L. humile* that have

undergone recent genetic bottlenecks). Thus, if stochastically produced diploid males occur at

very low baseline levels in ants, such males will rarely be detected in sexual species. In *O. biroi*,

however, they could make up a substantial proportion of diploid males, simply because diploid

males caused by homozygosity at the CSD locus are likewise extremely rare.

Finally, we note that heterozygosity-dependent female development, a fundamental tenet

of CSD, has major implications for breeding system evolution. Due to the high fitness cost of

producing sterile diploid males, species with CSD should obligatorily outbreed or have other

mechanisms of maintaining heterozygosity (van Wilgenburg et al. 2006). Intriguingly, in many

taxa, asexual lineages reproduce via mechanisms that lead to complete homozygosity in a single

generation (Suomalainen et al. 1987; Ma and Schwander 2017). However, among the social

Hymenoptera (and other Vespoids), the known modes of asexual reproduction all maintain at

least some heterozygosity (Rabeling and Kronauer 2012; Ma and Schwander 2017). For

formicoid ants, bees, and other taxa with CSD, asexual reproduction with high rates of

heterozygosity loss would incur a steep fitness penalty due to the production of sterile diploid

males. This may help explain the unusual inheritance system recently described in *O. biroi*,

where following crossover recombination, reciprocally recombined chromatids are faithfully co-

inherited so that no heterozygosity is lost (Lacy et al. 2024).

How this ancient putative CSD locus triggers female development in a heterozygosity-

dependent manner remains unknown. In *L. humile*, RNAi experiments suggested that the

expression level of the lncRNA *ANTSR* affects sex-specific splicing of *tra*, even though

heterozygosity in the exons of *ANTSR* is not required for this effect (Pan et al. 2024). Thus, the

heterozygosity-dependent function of the putative CSD locus may depend on local regulatory

interactions between homologous chromosomes, similar to a phenomenon known as transvection

(Duncan 2002). To better understand how heterozygosity induces female development in ants,

future studies will need to fine-map and functionally characterize this causal locus.

Materials and Methods

Software Versions

The versions of all software used for analyses are provided in **supplementary table S7**.

Identifying Male Ploidy

Clonal raider ant colonies are composed of female ants that reproduce asexually via a

type of thelytokous parthenogenesis termed central fusion automixis (Oxley et al. 2014). Males

are produced only sporadically (Oxley et al. 2014). Females of this species are diploid, whereas

males can either be haploid (presumably arising via a failure of central fusion following female

meiosis) or diploid (presumably arising due to rare losses of heterozygosity at CSD loci). Males

are easily recognizable due to substantial sexual dimorphism (males have eyes and wings,

whereas females are blind and flightless, and males also have different body shape and more

darkly pigmented cuticles than females). We sampled them opportunistically whenever they

were observed in colonies. We distinguished diploid males from haploid males based on

heterozygosity at several unlinked genetic markers (**supplementary table S1**). For this, we

disrupted one leg from each male using a Qiagen TissueLyser II and extracted DNA using

Qiagen's QIAmp DNA Micro Kit. We then PCR-amplified each marker and genotyped using

Sanger sequencing (for most markers) or via gel electrophoresis either after restriction digestion

(for two markers) or directly after amplification (for one marker at which the different alleles are

different sizes). Details of the genotyping methods can be found in **supplementary table S1**.

<u>Genome Sequencing and Preliminary Analysis</u>

For each library, we disrupted an individual ant with a metal bead and Qiagen's

TissueLyser II and extracted DNA with Qiagen's QIAmp DNA Micro Kit. We prepared Nextera

Flex Illumina DNA sequencing libraries for 16 diploid males from clonal line A, six diploid

females from different clonal lines, and seven haploid males from different clonal lines. We

sequenced these libraries using an Illumina NovaSeq 6000. We also accessed libraries for

individual genomes from previous studies. The information for all DNA sequencing libraries

used in this study can be found in **supplementary table S2**.

We trimmed reads using Trimmomatic 0.36 (Bolger et al. 2014), aligned them to the *O.*

biroi reference genome (McKenzie and Kronauer 2018) (Obir_v5.4, GenBank assembly

accession: GCA_003672135.1) using bwa mem (Li 2013), and then sorted, deduplicated, and

indexed using picard (http://broadinstitute.github.io/picard/). We called variants using GATK

HaplotypeCaller (version 4.2) (Poplin et al. 2018). To exclude falsely collapsed regions in the

reference genome, we filtered against sites found to be heterozygous in haploid males. We also

screened out any variants for which putatively heterozygous samples had a proportionate minor

allelic depth less than 0.25 and/or for which putatively homozygous samples had a minor allelic

depth greater than zero. We performed custom filtering of variants and calculated statistics using

pyvcf 0.6.8 (https://github.com/jamescasbon/PyVCF), and performed sliding window analyses

using pybedtools 0.9.0 (Dale et al. 2011).

<u>CSD Index Mapping</u>

By definition, CSD loci are heterozygous in females, but can be homozygous in diploid

males. Thus, traditional association mapping cannot identify CSD loci because particular alleles

do not trigger female development. Therefore, to map candidate CSD loci, we used a "CSD

Index," which equals zero if any female is homozygous, but if all females are heterozygous, it

equals the proportion of diploid males that are homozygous. Before calculating this index, we

had to identify ancestrally heterozygous sites. All individuals within an *O. biroi* clonal line are

descended asexually from a single diploid female ancestor, and CSD loci would have been

heterozygous in that common ancestor. Putatively ancestrally heterozygous variants are those for

which at least one diploid individual is heterozygous, and all other diploid individuals are either

heterozygous for the same two alleles, or homozygous for one of the two alleles for which other

diploid(s) are heterozygous (Oxley et al. 2014; Lacy et al. 2024). After identifying putatively

ancestrally heterozygous SNPs, we calculated the CSD index for each SNP and then recorded the

mean CSD index value in sliding windows.

<u>Losses of Heterozygosity</u>

We identified losses of heterozygosity as ancestrally heterozygous SNPs that had become

homozygous in a given sample. To assess whether homozygosity for either allele at the mapped

region permitted male development, we randomly chose a reference haploid male and assigned

allelic identity based on whether a focal sample was homozygous for the same allele or the

alternate allele to that possessed by the reference haploid male. To clearly illustrate losses of

heterozygosity, we used the ordinal position of SNPs rather than the position of SNPs in the

reference genome assembly.

Homozygosity in whole genome sequencing data can result from deletions, for which the

affected region would have half the read depth of the genome-wide average, or copy-neutral

losses of heterozygosity (which result rarely from meiotic recombination and central fusion

parthenogenesis (Kronauer et al. 2012; Rabeling and Kronauer 2012; Oxley et al. 2014; Lacy et

al. 2024)). To determine which of these processes caused homozygosity in diploid males, we

identified runs of homozygosity using "bcftools roh" (Narasimhan et al. 2016), obtained read

depths at all SNPs that passed filtering within these regions using "samtools depth –aa"

(Danecek et al. 2021) and then normalized by dividing by the genome-wide median read depth.

For samples without losses of heterozygosity on chromosome 4, we obtained normalized read

depth at all SNPs that passed filtering on chromosome 4.

<u>Homology Between CSD Loci Across Species</u>

To determine whether our mapped CSD locus bore homology to those identified in other

ant species, we used OrthoFinder (Emms and Kelly 2019) to identify orthologs of the genes

present in the two CSD QTL in *V. emeryi* (Miyakawa and Mikheyev 2015). The locus identified

in *L. humile* (Pan et al. 2024) is homologous to a subset of one of the *V. emeryi* QTL. In this

process, we noticed that one of the genes found within our mapped CSD locus (*cysteine-rich*

with EGF-like domain protein 2, or CRELD2) was also present in a second copy in a contig that

was erroneously duplicated during the genome assembly process (Chr4:21571-39095). We

masked this contig and realigned sequencing data for all subsequent analyses.

 Balancing Selection

Because CSD loci are expected to evolve under negative frequency-dependent

(balancing) selection, many different alleles should segregate in populations. To assess this, we

used scikit-allel (Miles et al. 2019) to calculate nucleotide diversity across six different clonal

lines and reported nucleotide diversity in sliding windows.

From inspecting DNA sequencing alignments to the reference genome, it became clear

that the high genetic variation within the mapped CSD locus likely resulted from several highly

differentiated haplotypes in this region. To investigate this further, we sequenced whole genomes

of haploid males from clonal lines other than A and B. We assembled the genomes of each

haploid male de novo using SPAdes (Bankevich et al. 2012; Prjibelski et al. 2020), found the

contig bearing homology to the mapped CSD locus in the reference genome assembly using

blastn (Camacho et al. 2009), manually scaffolded contigs as needed, and pairwise aligned each

alternate allele to the reference genome using nucmer and mummer (Kurtz et al. 2004) to identify

differences between alleles.

RNAseq for Annotation Improvements

To avoid missing any previously unannotated genes in the mapped CSD locus region, we

performed RNA sequencing following various approaches. To start, we extracted total RNA

from multiple samples, including separate pooled tissue from several different life stages (eggs,

young larvae, fourth instar larvae, prepupae, pupae, and adults), different adult tissues (antennae,

heads, thoraces, legs, abdomens), and individual male and female pupae using a TriZol RNA

extraction method followed by precipitation in isopropanol at -20C overnight. We used the

different life stage extractions for long read RNA-seq, rRNA depletion RNA-seq, and small

RNA-seq. We used the different adult tissues for long read RNA-seq, and the individual male

and female pupae for mRNA-seq. Long read RNA-seq was performed on a PacBio Sequel II,

and all other sequencing was performed on an Illumina NovaSeq 6000. For rRNA depletion, we

used Illumina's TruSeq Stranded Total RNA RiboZero kit. For small RNA-seq, we prepared

libraries using Perkin Elmer's NEXTFLEX Small RNA-Seq Kit v3 for Illumina Platforms.

<u>Annotation Improvements – Long Read Transcriptome Assembly</u>

The GenBank reference genome (GCA_003672135) for *O. biroi* (Obir_v5.4) was

retrieved from NCBI in FASTA format alongside the GenBank (GCA_003672135) and RefSeq

(GCF_003672135.1) transcriptomes in GTF format. Additionally, a curated RefSeq annotation

was retrieved from Zenodo (https://doi.org/10.5281/zenodo.10079884) in GTF format. RefSeq

assemblies were chromosome name-mapped to GenBank chromosome names using the

Bioconductor GenomeInfoDb package (Arora et al. 2024) and the chromosome name maps

available in NCBI.

The PacBio toolset's pbmm2 (https://github.com/PacificBiosciences/pbmm2), a PacBio

wrapper for Minimap2 (Li 2018), was used for indexing and alignment of pooled full-length,

non-concatemer long reads to the GenBank reference genome using the parameters "--preset

ISOSEQ -j 8". Following the generation of aligned long reads in BAM format, Stringtie2

(Kovaka et al. 2019) was used with parameters "-L" for long read transcriptome assembly, and

the curated RefSeq annotation was supplied as a guide for assembly using the "-G" parameter.

The resulting gene models were exported for further analysis and integration as a gzipped GTF

file.

Annotation Improvements – Short Read Transcriptome Assembly

For Illumina RNA-seq data, FastQ quality control was performed using Rfastp (Wang

and Carroll 2024). Indexing and alignment of short reads to the GenBank genome was performed

using Hisat2 (Kim et al. 2019) with default parameters, and samtools was used to generate sorted

and indexed BAM files for further analysis. Stringtie2 was then used to assemble transcriptomes

using the default parameters. Gene models were then combined across replicates using

Stringtie2's merge function with parameters "-F 0.5 -T 2 -f 0.1" and were exported for further

analysis and integration as a gzipped GTF file.

Annotation Improvements – Integrating Short and Long Read Transcriptomes with the RefSeq

Assembly.

Integration of long read and short read generated gene models with the established and

curated RefSeq transcriptome was performed using custom scripts in R available from GitHub

(https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025) and described here.

First, conflicting gene models where single long read genes map to multiple RefSeq

genes and vice versa were identified. As generated long read gene models are supported by

multiple full-length, non-concatemer long read alignments, attention was focused on the 720

instances of multiple long read generated genes overlapping single RefSeq genes.

Single exon long read genes were removed as these are often found to be artifacts from

PacBio sequencing. Following this, long read gene models that split RefSeq gene models but

matched with GenBank gene models were accepted with the corresponding RefSeq models

rejected.

Next, RefSeq gene models that conflicted within themselves were removed. This includes

genes that were overlapping in their exons but marked as separate genes.

After tidying RefSeq gene models, long read gene models that split RefSeq gene models

but matched with short read gene models were accepted, with the corresponding RefSeq models

rejected. After this gene model polishing, only 132 gene models were left where multiple long

read generated genes overlapped single RefSeq genes.

To potentially rescue internally primed transcripts caused by long stretches of genomic

polyAs, long read transcripts were assessed for polyA nucleotide content downstream from their

endpoints, and these distributions were visualized to identify cut-offs for putative internally

primed transcripts. Transcripts with at least 70 percent as 10bp downstream of the last exon or

with a polyA stretch > 4 in 10bp of the last exons were labeled as putative internally primed

transcripts. For putative internally primed transcripts, which also split RefSeq genes, short read

gene models were used to guide the rescued models when both short read and RefSeq agreed on

a single gene model.

Finally, for the remaining long read gene models that split RefSeq gene models, gaps

between long read gene models were assessed in the short read and RefSeq gene models. If

agreement in exon and junction structure was found between short read and RefSeq gene models,

the long read gene models were rejected. This left a final 27 genes where long read gene models

split RefSeq gene models, and these were manually assessed in IGV alongside short read and

long read alignments to reject or accept.

With a final set of long read gene models, all RefSeq and short read transcripts that fell

within single genes were then added to the long read transcriptome to create an integrated

transcriptome. Duplicate transcripts were then removed from the integrated transcriptome, and

transcripts and genes were annotated for their sources (long read, short read, RefSeq) and any

overlap with the established RefSeq, GenBank, and Ensembl gene models included. Following

the creation of the integrated gene models, SQANTI3 (Pardo-Palacios et al. 2024) was used to

provide a final corrected set of gene models as well as additional QC, as described below.

<u>Assessment of Annotation Quality and Identifying Novel Genes, Transcripts and Exons</u>

SQANTI3 was run with default parameters on the integrated transcriptome against the

curated RefSeq and GenBank gene models to generate QC and descriptive statistics on novel and

non-canonical splicing events. Gffcompare software (Pertea and Pertea 2020) was run against the

curated RefSeq and GenBank gene models to provide statistics on novel exons, transcripts, and

genes.

BUSCO (Simão et al. 2015) was run on the integrated, curated RefSeq and GenBank

gene models with the parameters "-l insecta_odb10 -m tran" to provide measures of

transcriptome completeness based on a set of core gene models. All these metrics were

summarized into tables using custom R scripts available on GitHub

(https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025).

<u>Annotation of CDS and Functional Annotation of Genes</u>

Annotation of coding regions was performed using TransDecoder (Haas 2023) to add

CDS information to our integrated gene models GTF. TransDecoder was run with default

parameters except for providing the "--single_best_only" to the TransDecoder.Predict function to

get only one prediction per transcript. CDS information was merged with SQANTI3's GeneMark

CDS annotation for a final set of CDSs. With this coding information, gene models were read

into R, translated into protein sequences using the BioStrings package, and protein per transcript

sequences exported as FASTA format for use in downstream tools. For functional annotation,

both eggNOG (Huerta-Cepas et al. 2019) and InterProScan (Jones et al. 2014) were used using

default parameters with our protein sequences FASTA to provide predictions for functional

annotations of genes including their putative GO, KEGG, PFAM and Panther categories.

Annotations were summarized to gene levels from transcript level annotation using R.

Annotation of lncRNAs

FEELnc (Wucher et al. 2017) was used to identify putative lncRNAs within our final

integrated gene models. First, the FEELnc_filter.pl script was run with default parameters, and

then the FEELnc_codpot.pl script was run with parameters "--mode=shuffle" to generate our

putative set of lncRNAs for annotation of our integrated gene model GTF.

Annotation of miRNAs

Small RNA-seq reads were adaptor clipped for a "TGGAATTCTCGGGTGCCAAGG" adapter

using CLIPflexR (Rozen-Gagnon et al. 2021) and aligned to the genome using miRDeep2's

mapper.pl script with parameters "-c -j -m -l 18" (Friedländer et al. 2012). Aligned data were

then processed using miRDeep2's miRDeep2.pl script with a FASTA of known *Drosophila*

miRNAs retrieved from miRBase (Kozomara et al. 2019). The resulting annotated miRNA

genomic locations were merged across samples and added into our integrated gene model GTF.

Annotation of piRNAs

For the identification of piRNA clusters, small read RNA-seq was processed using the pipeline

workflow described within the Pilfer (Ray and Pandey 2018) toolkit to generate piRNA cluster

BED files per sample. The sample piRNA clusters were then merged across samples to produce a

final piRNA cluster set, which was then included in our integrated gene model GTF.

Annotation of tRNAs, snRNAs, snoRNAs and rRNAs

Additional annotations for tRNAs, snRNAs, snoRNAs and rRNAs were mapped from RefSeq

annotation onto our integrated gene model GTF.

Sex-specific Gene Expression

To identify genes with sex-specific expression, we aligned Illumina mRNA-seq reads

from three female and three male pupae using STAR (v2.5.0a) (Dobin et al. 2013). We counted

and assigned reads to genes and transcripts using RSEM (v1.2.28) (Li and Dewey 2011). We

analyzed differential gene expression using DESeq2 (Love et al. 2014), filtering out genes

shorter than ten bases or with less than ten counts across all six samples. We hierarchically

clustered the differentially expressed genes by Euclidian distance using the R package pheatmap

(v1.0.12) (Kolde 2024). We used DEXSeq (v1.50.0) (Anders et al. 2012) to identify differential

exon usage.

Data and Software Availability

All DNA and RNA sequencing data are publicly available at the National Center for

Biotechnology Information Sequence Read Archive under accession number PRJNA1075055.

All other data are available in the article and Supporting Information. All code is available on

GitHub (genome annotation:

https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025; differential gene

expression and exon usage: https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq;

all other code and analyses: https://github.com/kipdlacy/SexDetermination_Obiroi).

Contributions

640 K.D.L., J.L., and D.J.C.K. designed research; J.L. analyzed data and visualized results for gene-

641 expression and exon usage analyses; K.R.G., W.W., and T.C. improved the genome annotation

642 with input from K.D.L.; K.D.L. performed research, analyzed data, and visualized results for all

643 other parts of the manuscript; K.D.L. and D.J.C.K. wrote the manuscript; D.J.C.K. supervised the

644 project.

Competing interests

647 The authors declare no competing interests.

Acknowledgements

We thank Stephany Valdés-Rodríguez, Leonora Olivos-Cisneros, Alejandra Hurtado-Giraldo,

Sean McKenzie, and other members of the Kronauer laboratory for helping us obtain male ants

and for valuable discussions; Jenn Balacco and the Rockefeller University Reference Genome

Resource Center for long read RNA sequencing; the Weill Cornell Genomics Resources Core

Facility for Small RNA sequencing; Connie Zhao and the Rockefeller University Genomics

Resource Center for rRNA-depletion sequencing and all other Illumina sequencing; Tom Kay,

Matteo Rossi, and Yukina Chiba for valuable comments on an earlier version of the manuscript.

This work was supported by a Gabrielle H. Reem and Herbert J. Kayden Early-Career

Innovation Award and the National Institute of General Medical Sciences of the National

Institutes of Health under award no. R35GM127007, both to D.J.C.K. The content is solely the

responsibility of the authors and does not necessarily represent the official views of the National

Institutes of Health. This work was also supported by the Howard Hughes Medical Institute,

where D.J.C.K. is an Investigator. This is Clonal Raider Ant Project paper number 36.

663
664   **<u>References</u>**
665

666   Anders S, Reyes A, Huber W. 2012. Detecting differential usage of exons from RNA-seq data.
667   *Genome Research* 22:2008–2017.

668   Arora S, Morgan M, Carlson M, Pagès H. 2024. GenomeInfoDb: Utilities for manipulating
669   chromosome names, including modifying them to follow a particular naming style.

670   Bachtrog D, Mank JE, Peichel CL, Kirkpatrick M, Otto SP, Ashman TL, Hahn MW, Kitano J,
671   Mayrose I, Ming R, et al. 2014. Sex determination: why so many ways of doing it? *PLoS*
672   *Biology* 12:e1001899.

673   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko
674   SI, Pham S, Prjibelski AD, et al. 2012. SPAdes: a new genome assembly algorithm and
675   its applications to single-cell sequencing. *Journal of Computational Biology* 19:455–477.

676   Beye M, Hasselmann M, Fondrk MK, Page RE, Omholt SW. 2003. The gene *csd* is the primary
677   signal for sexual development in the honeybee and encodes an SR-type protein. *Cell*
678   114:419–429.

679   Beye M, Seelmann C, Gempe T, Hasselmann M, Vekemans X, Fondrk MK, Page RE. 2013.
680   Gradual molecular evolution of a sex determination switch through incomplete
681   penetrance of femaleness. *Current Biology* 23:2559–2564.

682   Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence
683   data. *Bioinformatics* 30:2114–2120.

684   Bopp D;, Saccone G;, Beye M. 2014. Sex determination in insects: variations on a common
685   theme. *Sexual Development* 8:20–28.

686   Borowiec ML, Rabeling C, Brady SG, Fisher BL, Schultz TR, Ward PS. 2019. Compositional
687   heterogeneity and outgroup choice influence the internal phylogeny of the ants.
688   *Molecular Phylogenetics and Evolution* 134:111–121.

Borowiec ML, Zhang YM, Neves K, Ramalho MO, Fisher BL, Lucky A, Moreau CS. 2025.
Evaluating UCE data adequacy and integrating uncertainty in a comprehensive
phylogeny of ants. *Systematic Biology*:syaf001.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009.
BLAST+: architecture and applications. *BMC Bioinformatics* 10:1–9.

Chodroff RA, Goodstadt L, Sirey TM, Oliver PL, Davies KE, Green ED, Molnár Z, Ponting CP.
2010. Long noncoding RNA genes: conservation of sequence and brain expression
among diverse amniotes. *Genome Biology* 11:R72.

Crozier RH. 1971. Heterozygosity and sex determination in haplo-diploidy. *The American*
Naturalist 105:399–412.

Dale RK, Pedersen BS, Quinlan AR. 2011. Pybedtools: a flexible Python library for
manipulating genomic datasets and annotations. *Bioinformatics* 27:3423–3424.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T,
McCarthy SA, Davies RM, et al. 2021. Twelve years of SAMtools and BCFtools.
GigaScience 10:giab008.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras
TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29:15–21.

Duncan IW. 2002. Transvection effects in *Drosophila*. *Annual Review of Genetics* 36:521–556.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biology* 20:238.

Friedländer MR, Mackowiak SD, Li N, Chen W, Rajewsky N. 2012. miRDeep2 accurately
identifies known and hundreds of novel microRNA genes in seven animal clades. *Nucleic*
Acids Research 40:37–52.

Gempe T, Beye M. 2011. Function and evolution of sex determination mechanisms, genes and
pathways in insects. *BioEssays* 33:52–60.

Gempe T, Hasselmann M, Schiøtt M, Hause G, Otte M, Beye M. 2009. Sex determination in
honeybees: Two separate mechanisms induce and maintain the female pathway. *PLoS*
Biology 7:e1000222.

Haas B. 2023. TransDecoder (find coding regions within transcripts). Available from:
https://github.com/TransDecoder/TransDecoder

Hasselmann M, Beye M. 2004. Signatures of selection among sex-determining alleles of the
honey bee. *Proceedings of the National Academy of Sciences* 101:4888–4893.

Hasselmann M, Gempe T, Schiøtt M, Nunes-Silva CG, Otte M, Beye M. 2008. Evidence for the
evolutionary nascence of a novel sex determination pathway in honeybees. *Nature*
454:519–522.

Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H, Mende DR,
Letunic I, Rattei T, Jensen LJ, et al. 2019. eggNOG 5.0: a hierarchical, functionally and
phylogenetically annotated orthology resource based on 5090 organisms and 2502
viruses. *Nucleic Acids Research* 47:D309–D314.

Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell
A, Nuka G, et al. 2014. InterProScan 5: genome-scale protein function classification.
Bioinformatics 30:1236–1240.

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology* 37:907–915.

Kolde R. 2024. pheatmap: Pretty Heatmaps. Available from:
https://github.com/raivokolde/pheatmap

Kovaka S, Zimin AV, Pertea GM, Razaghi R, Salzberg SL, Pertea M. 2019. Transcriptome
assembly from long-read RNA-seq alignments with StringTie2. *Genome Biology* 20:278.

Kozomara A, Birgaoanu M, Griffiths-Jones S. 2019. miRBase: from microRNA sequences to
function. *Nucleic Acids Research* 47:D155–D162.

Kronauer DJC, Pierce NE, Keller L. 2012. Asexual reproduction in introduced and native
populations of the ant *Cerapachys biroi*. *Molecular Ecology* 21:5221–5235.

Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL. 2004.
Versatile and open software for comparing large genomes. *Genome Biology* 5:R12.

Lacy KD, Hart T, Kronauer DJC. 2024. Co-inheritance of recombined chromatids maintains
heterozygosity in a parthenogenetic ant. *Nature Ecology & Evolution* 8:1522–1533.

Li B, Dewey CN. 2011. RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. *BMC Bioinformatics* 12:323.

Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
Available from: http://arxiv.org/abs/1303.3997

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34:3094–
3100.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biology* 15:550.

Ma W-J, Schwander T. 2017. Patterns and mechanisms in instances of endosymbiont☐induced
parthenogenesis. *Journal of Evolutionary Biology* 30:868–888.

Matthey-Doret C, van der Kooi CJ, Jeffries DL, Bast J, Dennis AB, Vorburger C, Schwander T.
2019. Mapping of multiple complementary sex determination loci in a parasitoid wasp.
Genome Biology and Evolution 11:2954–2962.

McKenzie SK, Kronauer DJC. 2018. The genomic architecture and molecular evolution of ant
odorant receptors. *Genome Research* 28:1757–1765.

Miles A, Ralph P, Rae S, Pisupati R. 2019. cggh/scikit-allel: v1.2.1. Available from:
https://zenodo.org/records/3238280

Miyakawa MO, Mikheyev AS. 2015. QTL mapping of sex determination loci supports an
ancient pathway in ants and honey bees. *PLoS Genetics* 11:e1005656.

Miyakawa MO, Miyakawa H. 2023. *Transformer* gene regulates feminization under two
complementary sex determination loci in the ant, *Vollenhovia emeryi. Insect*
Biochemistry and Molecular Biology 156:103938.

Miyakawa MO, Tsuchida K, Miyakawa H. 2018. The doublesex gene integrates multi-locus
complementary sex determination signals in the Japanese ant, *Vollenhovia emeryi. Insect*
Biochemistry and Molecular Biology 94:42–49.

Narasimhan V, Danecek P, Scally A, Xue Y, Tyler-Smith C, Durbin R. 2016. BCFtools/RoH: a
hidden Markov model approach for detecting autozygosity from next-generation
sequencing data. *Bioinformatics* 32:1749–1751.

Normark BB. 2003. The evolution of alternative genetic systems in insects. *Annual Review of*
Entomology 48:397–423.

Otte M, Netschitailo O, Weidtkamp-Peters S, Seidel CAM, Beye M. 2023. Recognition of
polymorphic Csd proteins determines sex in the honeybee. *Science Advances* 9:eadg4239.

Oxley PR, Ji L, Fetter-Pruneda I, McKenzie SK, Li C, Hu H, Zhang G, Kronauer DJC. 2014.
The genome of the clonal raider ant *Cerapachys biroi. Current Biology* 24:451–458.

Pan Q, Darras H, Keller L. 2024. LncRNA gene ANTSR coordinates complementary sex
determination in the Argentine ant. *Science Advances* 10:eadp1532.

Pardo-Palacios FJ, Arzalluz-Luque A, Kondratova L, Salguero P, Mestre-Tomás J, Amorín R,
Estevan-Morió E, Liu T, Nanni A, McIntyre L, et al. 2024. SQANTI3: curation of long-
read transcriptomes for accurate identification of known and novel isoforms. *Nature*
Methods 21:793–797.

Pertea G, Pertea M. 2020. GFF Utilities: GffRead and GffCompare. Available from:
https://f1000research.com/articles/9-304

Poplin R, Ruano-Rubio V, DePristo MA, Fennell TJ, Carneiro MO, Auwera GAV der, Kling
DE, Gauthier LD, Levy-Moonshine A, Roazen D, et al. 2018. Scaling accurate genetic

789   variant discovery to tens of thousands of samples. Available from:
790   https://www.biorxiv.org/content/10.1101/201178v3

791   Privman E, Wurm Y, Keller L. 2013. Duplication and concerted evolution in a master sex
792   determiner under balancing selection. *Proceedings of the Royal Society B: Biological*
793   *Sciences* 280:20122968.

794   Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. 2020. Using SPAdes De
795   Novo Assembler. *Current Protocols in Bioinformatics* 70:e102.

796   Quinn JJ, Zhang QC, Georgiev P, Ilik IA, Akhtar A, Chang HY. 2016. Rapid evolutionary
797   turnover underlies conserved lncRNA–genome interactions. *Genes & Development*
798   30:191–207.

799   Rabeling C, Kronauer DJC. 2012. Thelytokous parthenogenesis in eusocial Hymenoptera.
800   *Annual Review of Entomology* 58:273–292.

801   Ray R, Pandey P. 2018. piRNA analysis framework from small RNA-Seq data by a novel cluster
802   prediction tool - PILFER. *Genomics* 110:355–365.

803   Rozen-Gagnon K, Gu M, Luna JM, Luo J-D, Yi S, Novack S, Jacobson E, Wang W, Paul MR,
804   Scheel TKH, et al. 2021. Argonaute-CLIP delineates versatile, functional RNAi networks
805   in *Aedes aegypti*, a major vector of human viruses. *Cell Host & Microbe* 29:834–848.

806   Seiler J, Beye M. 2024. Honeybees' novel complementary sex-determining system: function and
807   origin. *Trends in Genetics* 40:969–981.

808   Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO:
809   assessing genome assembly and annotation completeness with single-copy orthologs.
810   *Bioinformatics* 31:3210–3212.

811   Suomalainen E, Saura A, Lokki J. 1987. Cytology and Evolution in Parthenogenesis. CRC Press

812   Trible W, Chandra V, Lacy KD, Limón G, McKenzie SK, Olivos-Cisneros L, Arsenault SV,
813   Kronauer DJC. 2023. A caste differentiation mutant elucidates the evolution of socially
814   parasitic ants. *Current Biology* 33:1047–1058.

815   Trible W, Mckenzie SK, Kronauer DJC. 2020. Globally invasive populations of the clonal raider
816   ant are derived from Bangladesh. *Biology Letters* 16:20200105.

817   Ulitsky I, Shkumatava A, Jan CH, Sive H, Bartel DP. 2011. Conserved function of lincRNAs in
818   vertebrate embryonic development despite rapid sequence evolution. *Cell* 147:1537–
819   1550.

820   Verhulst EC, Beukeboom LW, van de Zande L. 2010. Maternal control of haplodiploid sex
821   determination in the wasp *Nasonia*. *Science* 328:620–623.

Wang W, Carroll T. 2024. Rfastp: An ultra-fast and all-in-one fastq preprocessor (quality
control, adapter, low quality and polyX trimming) and UMI sequence parsing). Available
from: 10.18129/B9.bioc.Rfastp

Wexler J, Delaney EK, Belles X, Schal C, Wada-Katsumata A, Amicucci MJ, Kopp A. 2019.
Hemimetabolous insects elucidate the origin of sexual development via alternative
splicing. *eLife* 8:e47490.

Whiting PW. 1933. Selective fertilization and sex-determination in Hymenoptera. *Science*
78:537–538.

Whiting PW. 1935. Genic balance, sex-determination and selective fertilization in Hymenoptera.
Proceedings of the American Philosophical Society 75:517–520.

Whiting PW. 1943. Multiple alleles in complementary sex determination of *Habrobracon*.
Genetics 28:365–382.

van Wilgenburg E, Driessen G, Beukeboom LW. 2006. Single locus complementary sex
determination in Hymenoptera: an "unintelligent" design? *Frontiers in Zoology* 3:1.

Williams TM, Carroll SB. 2009. Genetic and molecular insights into the development and
evolution of sexual dimorphism. *Nature Reviews Genetics* 10:797–804.

Wucher V, Legeai F, Hédan B, Rizk G, Lagoutte L, Leeb T, Jagannathan V, Cadieu E, David A,
Lohi H, et al. 2017. FEELnc: a tool for long non-coding RNA annotation and its
application to the dog transcriptome. *Nucleic Acids Research* 45:e57.

Zou Y, Geuverink E, Beukeboom LW, Verhulst EC, van de Zande L. 2020. A chimeric gene
paternally instructs female sex determination in the haplodiploid wasp *Nasonia*. *Science*
370:1115–1118.

Fig. 1. Theoretical expectations for complementary sex determination (CSD). (A) Cartoon depicting

a mating between a diploid female with two different alleles at a sex determination locus and a haploid

male bearing one of these two alleles. Half of the sexually produced diploid offspring are expected to

develop as males. **(B)** Table illustrating multi-locus CSD in diploids, in which heterozygosity of at least

one sex determination locus is required for female development. By contrast, only homozygosity at all

loci results in diploid male development. **(C)** Cartoon depicting how CSD might work in asexual species

such as the clonal raider ant. If diploid males arise from losses of heterozygosity at sex loci, then

homozygosity for either of the two alleles should trigger male development. Offspring proportions reflect

empirical observations in the clonal raider ant (Kronauer et al. 2012; Oxley et al. 2014).

855
856
<u>**Fig. 2.**</u> **Whole genome sequencing reveals a candidate sex determination locus on chromosome 4. (A)**

Karyoplot depicting the mean CSD index in 50kb windows with a 15kb sliding interval. For each SNP,

the CSD index equals zero if any female is homozygous. If all females are heterozygous, the CSD index

equals the proportion of homozygous diploid males. The CSD index peak is shown as a green dotted line.

The grey histogram shows the number of ancestrally heterozygous SNPs in 200kb windows. **(B)** Stacked

bar plots for all diploids used for mapping, with one horizontal line for each ordered putatively ancestrally

heterozygous SNP on chromosome 4. For each sample, SNPs that retain ancestral heterozygosity are

drawn as grey lines, whereas SNPs that have lost ancestral heterozygosity are drawn as black lines.

	Heterozygous	Homozygous Allele 1	Homozygous Allele 2
Diploid Females	19	0	0
Diploid Males	5	2	9

Table 1. Number of samples heterozygous and homozygous for each allele in the 46kb region on chromosome 4.

Fig. 3. The putative sex determination locus identified in *O. biroi* is conserved across formicoid ants.

(**A**) Karyoplot depicting the 14 chromosomes in the *O. biroi* genome, chromosome 4 from the *L. humile*

genome (Pan et al. 2024), and one contig from linkage group 14 of the *V. emeryi* reference genome

(Miyakawa and Mikheyev 2015). Note that the plots from the different species are not drawn to scale.

Homology to protein-coding genes identified within *V.emeryiCsdQTL2* is drawn in green. The location of

the *O. biroi* CSD index peak is indicated. (**B**) A phylogeny of the ant subfamilies (adapted from

(Borowiec et al. 2019)), with the presence of a sex determination locus homologous to the peak of our

CSD index, or the absence of data shown. The yellow shaded background denotes the formicoid clade.

(**C**) Karyoplot for *O. biroi* chromosome 4, depicting homology to *V.emeryiCsdQTL2* and ancestral

heterozygosity for six different clonal lines (A, B, C, D, I, and M). Grey histograms depict the number of

ancestrally heterozygous SNPs in 100kb windows.

Fig. 4. The CSD index peak is characterized by high genetic diversity in a non-coding region. (A &

B) Nucleotide diversity across the length of *O. biroi* chromosome 4 in 5kb windows (step size=1kb) **(A)**

and across the vicinity of the CSD index peak in 100bp windows (step size=20bp) **(B)**. **(C)** The number of

differences per 100bp window between each alternate *de novo* assembled allele and the reference genome

allele. **(D)** Annotated genes in the vicinity of the CSD index peak. Black boxes depict exons and thin lines

depict introns. The lncRNA *ANTSR* is indicated in bold. Arrows indicate the names of genes in close

proximity. The CSD index peak is shown as a green dotted line in A, and with green shading in B-D.

Clonal Line	CSD Index Peak Haplotypes
A	A1 / A2
B	B1 / A1
C	? / ?
D	D1 / D2
I	I1 / ?
M	M1 / A1

888   **Table 2.** The haplotypes present at the CSD index peak for each studied *O. biroi* clonal line.

Clonal Line	Haploid Males	Diploid Males
A	92	22
B	78	0
C	6	0
D	4	0
I	1	0
M	2	0

Table 3. Number of haploid and diploid males sampled from different *O. biroi* clonal lines.

Fig. 5. <u>**Fig. 5.**</u> **Whether a second, *tra*-containing CSD QTL from *V. emeryi* is conserved across ants remains**

ambiguous. (A) Karyoplot depicting the 14 chromosomes in the *O. biroi* genome and one contig from

linkage group 13 of the *V. emeryi* reference genome. Homology to protein-coding genes identified within

V.emeryiCsdQTL1 is drawn in purple. The locations of the *O. biroi* CSD index peak and the homolog of

transformer are indicated. **(B)** A phylogeny of the ant subfamilies (adapted from (Borowiec et al. 2019)),

with the presence or absence of this second putative sex determination locus, or the absence of data

shown. The yellow shaded background denotes the formicoid clade. **(C)** Karyoplot for *O. biroi*

chromosome 2, depicting homology to *V.emeryiCsdQTL1* and ancestral heterozygosity for each clonal

line. Grey histograms depict the number of ancestrally heterozygous SNPs in 100kb windows. The region

of homology that is ancestrally homozygous in clonal lines A and C is labeled.

Supplementary Figures

<u>**Supplementary Fig. S1.** Genome-wide heterozygosity levels.</u> Scatterplot depicting the

proportion of base pairs in the ~220Mb *O. biroi* genome that are heterozygous in each individual

for which we conducted whole genome sequencing. Haploid males have no legitimate

heterozygosity—the small elevation above zero is due to heterozygous genotype calls at

erroneously assembled regions of the reference genome.

Supplementary Fig. S2. Diploid males result from copy-neutral losses of heterozygosity for

either allele. (A) Karyoplot depicting losses (green) or retention (gray) of heterozygosity across

the vicinity of the peak of the CSD index in diploid males and females. Codes at the left (C16,

C17, etc.) indicate the stock colonies from which each ant was sampled. The magenta box

surrounds the CSD index peak. Dark green indicates that the sample has the same allele /

haplotype as the reference haploid male whereas light green indicates that the sample has the

alternative allele. Switches between alleles in homozygous regions within samples (i.e., changes

from light green to dark green or vice versa) represent recombination events that occurred in

generations prior to the meiosis that produced each diploid male. Because the reference haploid

male was from colony C16, fewer generations have elapsed since the common ancestor of the

reference haploid male and the diploid males from that colony, and thus fewer recombination

events have accumulated. By contrast, more generations have elapsed since the common ancestor

of the reference haploid male and the diploid males from colonies C17, C18, and C1, which is

reflected in the higher number of recombination events that accumulated across generations. **(B)**

Normalized read depth across either all SNPs that passed filtering on chromosome 4 (gray, for

samples without losses of heterozygosity on chromosome 4) or all SNPs that passed filtering

within that sample's loss of heterozygosity region (green). Box plots show median (center line),

interquartile range (IQR) (box limits) and 1.5 x IQR (whiskers). Data points that fall outside 1.5

x IQR are shown individually. Violin plots show the kernel probability density, meaning that the

proportion of the data located at an x-axis value is represented by the width of the outlined area.

Supplementary Fig. S3. Genome-wide nucleotide diversity. Nucleotide diversity across the

genome in 5kb windows with a 1kb sliding interval. In the genome-wide plot (top), the red line

depicts the peak of the CSD index from genetic mapping. In the zoomed-in plots for

chromosomes 4 (middle) and 2 (bottom), the green and purple vertical shading indicates

homology to *V. emeryi* CSD QTL 2 and 1, respectively.

Supplementary Fig. S4. Sex-specific gene expression. (**A**) Heatmap of genes that were

significantly differentially expressed between sexes (adjusted p-value < 0.05). Each row

represents a gene, and each column represents a sample. The trees at the left and the top of the

heat map depict hierarchical clustering of genes and samples, respectively. Z-scores of log-

transformed expression levels are shown as a color gradient. (**B**) Volcano plot depicting

differential gene expression between males and females. Each gene is represented by a dot,

which is colored gray if not differentially expressed, green if the expression difference is at least

2-fold ($\log \geq 1$), or blue if the expression difference is statistically significant (adjusted p-value

< 0.05). Red indicates genes that are significantly differentially expressed and have at least a 2-

fold difference in gene expression. (**C**) Expression levels of genes in the vicinity of the CSD

index peak. Normalized counts are shown for each male and female sample, with error bars

depicting the mean and SEM for each sex. The genomic coordinates, name, and ID are given for

each gene. Genes with significantly different expression between sexes (adjusted p-value < 0.05)

are bolded in the table and marked with an asterisk on the plot. The CSD index peak is shown in

green, and the dashed box denotes the peak of nuclear diversity.

Supplementary Tables

Primer Name	5.4 Scaffold	5.4 Position	Forward 5'-3'	Reverse 5'-3'	Genotyping Method
Obir Ploidy 12SV	Chr12	12801234	TAGATCGAGT GAGGTATCTT	AGCGTTAA GAGTAGAT TAAG	Visualize differently sized alleles (i.e., different versions of a structural variant [SV]) using gel electrophoresis
Obir Ploidy 4RE	Chr4	5840887	CCACCCTCTC TCTCGTCTC	TGCCGATT CATATTTC GAGATAAA C	Separately digest products with the restriction enzymes (REs) EcoRI and BsmI and visualize genotypes via gel electrophoresis
Obir Ploidy 5RE	Chr5	1963279	GCGATAGAA GAGTGCTCGT ATT	GCCAGTCA GTCTGTCA TTCA	Separately digest products with the restriction enzymes (REs) EcoRI and AvaII and visualize genotypes via gel electrophoresis
Obir Ploidy 2SNP	Chr2	18082503	CACGTACAAT GAGGGAACC TAA	TCGCGATC ATGAAATC TAACCT	Sanger sequence and call SNPs
Obir Ploidy 12SNP	Chr12	8100152	GTCAGTGCGC TGTTCATTTC	AGATTATA GGGCGTTC CAATTCT	Sanger sequence and call SNPs
Obir Ploidy 14SNP	Chr14	6377499	CACGTGAACG CTACGATGAA	GGGTTGAG GATGTCTA AGGTAATG	Sanger sequence and call SNPs
Obir Ploidy 13SNPa	Chr13	6262370	CGTTACACTC CTGTGTCGAA TC	GCACGTAG CATTCTCTC TTTCT	Sanger sequence and call SNPs
Obir Ploidy 13SNPb	Chr13	7173724	GCCATACATC CTGCACTCTT AC	CGACGAGC ATGCGTCT AAA	Sanger sequence and call SNPs

953 **Supplementary Table S1. Primers for ploidy assessment via heterozygosity.** 5.4 refers to the

954 reference genome version (GCA_003672135.1).

Sample name	Library prep method	Clonal line	Stock colony	Sample type	Mean read depth	BioProject
LineA_DiploidFemale_1	Nextera Flex	A	C16	Diploid Female	30.1	PRJNA923657
LineA_DiploidFemale_2	Nextera Flex	A	C16	Diploid Female	27.4	PRJNA923657
LineB_DiploidFemale_1	Nextera Flex	B	STC6	Diploid Female	28.3	PRJNA923657
LineA_DiploidFemale_6	Nextera Flex	A	C18	Diploid Female	30	PRJNA923657
LineA_DiploidFemale_5	Nextera Flex	A	C1	Diploid Female	41.5	PRJNA923657
LineA_DiploidFemale_3	Nextera Flex	A	C16	Diploid Female	26.6	PRJNA947942
LineA_DiploidFemale_4	Nextera Flex	A	C16	Diploid Female	24.1	PRJNA947942
LineB_DiploidFemale_2	Nextera Flex	B	STC6	Diploid Female	27	PRJNA947942
LineB_DiploidFemale_3	Nextera Flex	B	STC6	Diploid Female	47.1	PRJNA947942
LineA_HaploidMale_1	Nextera Flex	A	C16	Haploid Male	26.1	PRJNA947942
LineA_HaploidMale_2	Nextera Flex	A	C16	Haploid Male	26.1	PRJNA947942
LineA_HaploidMale_3	Nextera Flex	A	C16	Haploid Male	27.6	PRJNA947942
LineA_HaploidMale_4	Nextera Flex	A	C16	Haploid Male	25.8	PRJNA947942
LineB_HaploidMale_1	Nextera Flex	B	STC6	Haploid Male	26.3	PRJNA947942
LineB_HaploidMale_2	Nextera Flex	B	STC6	Haploid Male	27.6	PRJNA947942
LineB_HaploidMale_3	Nextera Flex	B	STC6	Haploid Male	31.7	PRJNA947942
LineB_DiploidFemale_5	Nextera Flex	B	STC12	Diploid Female	42.4	PRJNA947942
LineB_DiploidFemale_4	Nextera Flex	B	STC1	Diploid Female	31.7	PRJNA947942
LineA_HaploidMale_5	Nextera Flex	A	C1	Haploid Male	30.3	PRJNA947942
LineA_HaploidMale_6	Nextera Flex	A	C18	Haploid Male	25.3	PRJNA947942
LineB_HaploidMale_4	Nextera Flex	B	STC1	Haploid Male	31	PRJNA947942

LineB_HaploidMale_5	Nextera Flex	B	STC12	Haploid Male	32.8	PRJNA947942
DiploidMale_1	Nextera Flex	A	C16	Diploid Male	57.9	PRJNA1075055
DiploidMale_2	Nextera Flex	A	C16	Diploid Male	32.7	PRJNA1075055
DiploidMale_3	Nextera Flex	A	C16	Diploid Male	31.5	PRJNA1075055
DiploidMale_4	Nextera Flex	A	C18	Diploid Male	38.3	PRJNA1075055
DiploidMale_5	Nextera Flex	A	C17	Diploid Male	47.7	PRJNA1075055
DiploidMale_6	Nextera Flex	A	C16	Diploid Male	28.4	PRJNA1075055
DiploidMale_7	Nextera Flex	A	C16	Diploid Male	32.0	PRJNA1075055
DiploidMale_8	Nextera Flex	A	C2	Diploid Male	30.2	PRJNA1075055
DiploidMale_9	Nextera Flex	A	C16	Diploid Male	41.6	PRJNA1075055
DiploidMale_10	Nextera Flex	A	C16	Diploid Male	28.1	PRJNA1075055
DiploidMale_11	Nextera Flex	A	C18	Diploid Male	40.2	PRJNA1075055
DiploidMale_12	Nextera Flex	A	C16	Diploid Male	38.5	PRJNA1075055
DiploidMale_13	Nextera Flex	A	C16	Diploid Male	35.4	PRJNA1075055
DiploidMale_14	Nextera Flex	A	C16	Diploid Male	34.4	PRJNA1075055
DiploidMale_15	Nextera Flex	A	C1	Diploid Male	31.5	PRJNA1075055
DiploidMale_16	Nextera Flex	A	C1	Diploid Male	32.2	PRJNA1075055
LineC_DiploidFemale	Nextera Flex	C	?	Diploid Female	39.4	PRJNA1075055
LineD_DiploidFemale_2	Nextera Flex	D	BG9	Diploid Female	35.2	PRJNA1075055
LineD_DiploidFemale_1	Nextera Flex	D	?	Diploid Female	36.9	PRJNA1075055
LineI_DiploidFemale	Nextera Flex	I	?	Diploid Female	36.1	PRJNA1075055
LineM_DiploidFemale	Nextera Flex	M	?	Diploid Female	32.3	PRJNA1075055

LineB_HaploidMale	Nextera Flex	B	?	Haploid Male	28.3	PRJNA1075055
LineC_HaploidMale	Nextera Flex	C	C9	Haploid Male	27.6	PRJNA1075055
LineD_HaploidMale_1	Nextera Flex	D	BG7	Haploid Male	25.2	PRJNA1075055
LineD_HaploidMale_2	Nextera Flex	D	BG9	Haploid Male	34.9	PRJNA1075055
LineI_HaploidMale	Nextera Flex	I	BG2	Haploid Male	21.2	PRJNA1075055
LineM_HaploidMale_1	Nextera Flex	M	BG14	Haploid Male	70.7	PRJNA1075055
LineM_HaploidMale_2	Nextera Flex	M	BG14	Haploid Male	34.0	PRJNA1075055
LineB_DiploidFemale	Nextera Flex	B	STC6	Diploid Female	42.8	PRJNA1075055

955 **Supplementary Table S2. Metadata for all DNA whole-genome shotgun sequencing**

956 **libraries included in this study.** Question marks indicate male samples for which the clonal line

957 is known, but the stock colony of origin was not recorded.

V. emeryi CSD QTL	V. emeryi GeneID	OrthoFinder O. biroi "Ortholog"	O. biroi genomic locus
2	LOC105559778	LOC105285603	Chr4:1637384-1672949
2	LOC105559779	LOC105285604	Chr4:1673043-1674418
2	LOC105559781	LOC105285607	Chr4:1674523-1676002
2	LOC105559780	LOC105285605	Chr4:1675932-1678296
2	LOC105559782	LOC105283856, LOC113561733	Chr4:1727256-1729679; Chr4:22762-24749
2	LOC105559784	LOC105283852	Chr4:1729888-1737844
2	LOC105559783	LOC105283851	Chr4:1730326-1732211
2	LOC105559785	LOC105283850	Chr4:1738076-1749057
2	LOC105559786	LOC105282554	Chr4:7069030-7075460
2	LOC105559787	LOC105282557	Chr4:7075106-7080018
2	LOC105559788	LOC105286165	Chr4:4951574-4955391
2	LOC105559790	LOC105282568	Chr4:7141680-7152824
2	LOC105559789	LOC105282568	Chr4:7141680-7152824
2	LOC105559791	LOC105283915	Chr4:5377245-5381427
2	LOC105559793	LOC105283797	Chr4:2173362-2237652
1	LOC105558826	LOC105287916	Chr2:985469-993787
1	LOC105558815	LOC105283998	Chr2:14515051-14517463
1	LOC105558816	LOC105283997, LOC113563596	Chr2:14517342-14519998; Chr2:19219509-19221771
1	LOC105558817	LOC105283996, LOC113563597	Chr2:14519947-14522109; Chr2:19217723-19219230
1	LOC105558818	LOC105283999	Chr2:14522233-14522882
1	LOC105558819	LOC105278142	Chr2:16951611-16953740
1	LOC105558820	LOC105278143	Chr2:16947572-16951531
1	LOC105558821	LOC105278141	Chr2:16939029-16945082
1	LOC105558822	LOC105287131	Chr2:12765464-12768244
1	LOC105558823	LOC105278140	Chr2:16929798-16936497

958 **<u>Supplementary Table S3.</u>** *O. biroi* **orthologs of genes located in the two *V. emeryi* CSD**

959 **QTLs.**

	Gene_type	RU	RefSeq	GenBank
Genes	guide_RNA	0	1	0
	lncRNA	1767	1430	0
	miRNA	239	0	0
	misc_RNA	0	44	0
	piRNA	23596	0	0
	protein_coding	12839	11927	13022
	pseudogene	0	373	77
	rRNA	108	109	0
	snoRNA	15	16	0
	snRNA	22	26	0
	tRNA	194	202	0
Total Genes		**38780**	**14128**	**13099**
Transcripts	lncRNA	3795	1751	0
	miRNA	239	0	0
	mRNA	63620	24215	13099
	piRNA	23596	0	0
	rRNA	108	109	0
	snoRNA	15	16	0
	snRNA	22	26	0
	tRNA	194	202	0
	guide_RNA	0	1	0
	transcript	0	818	0
Total transcripts		**91589**	**27135**	**13099**
Total multi-exon transcripts		**65882**	**25945**	**11923**

960   **Supplementary Table S4. Improvements over previous genome annotations, with the new**

961   **(RU) annotation featuring genes and transcripts not found in previous annotation versions.**

Gene Name	Gene ID	Log 2-Fold Change	p-value	p-adjusted
unchar. protein	LOC105285599	0.11	0.84	0.92
unchar. protein	LOC105285600	0.11	0.73	0.85
paraneoplastic antigen Ma6E	LOC105285601	0.21	0.67	0.82
ataxin-7	LOC105285602	-0.33	0.063	0.17
Kank	LOC105285603	0.05	0.82	0.91
Bet1	LOC105285604	0.08	0.54	0.72
prob. ribosome prod. factor	LOC105285607	0.34	0.12	0.27
THUMP	LOC105285605	0.19	0.13	0.28
unchar. lncRNA	LOC105285608	0.12	0.87	0.93
Creld	LOC113561733	0.56	0.77	0.88
alphaCOP	LOC105283852	-0.17	0.21	0.40
unchar. protein	LOC105283851	9.95	NA	NA
Host cell factor	LOC105283850	-0.52	2.1E-04	1.8E-03
l(2)37Cg	LOC105283849	0.86	5.7E-04	4.2E-03
ribosomal protein S2	LOC105283847	-0.06	0.50	0.69
guanine deaminase	LOC105283846	0.03	0.88	0.94
unchar. protein	LOC105283844	1.77	6.1E-08	1.0E-07

962 **Supplementary Table S5. Differential gene expression near the CSD index peak.**

963 The log 2-fold change in expression levels between male and female samples, p-values (Wald

964 test), and adjusted p-values (Benjamini-Hochberg adjustment) for 17 genes in the vicinity of the

965 CSD index peak. The p-values are "NA" for LOC105283850 because one of the samples is an

966 extreme outlier detected by Cook's distance.

Gene Name	Gene ID	Exon	Log 2-Fold Change	p-value	p-adjusted
unchar. protein	LOC105285599	E001	-0.24	0.37	0.84
		E002	0.09	0.6	0.91
		E003	0.21	0.63	0.92
unchar. protein	LOC105285600	E001	-0.59	0.32	0.82
		E002	-0.23	0.78	0.96
		E003	0.51	0.15	0.72
		E004	0.03	0.87	0.98
		E005	0.06	0.73	0.95
		E006	-0.54	0.31	0.82
		E007	-0.48	0.35	0.83
		E008	0.11	0.91	0.99
		E009	-0.66	0.65	0.92
		E010	0.08	0.92	0.99
		E011	0.11	0.91	0.99
paraneoplastic antigen Ma6E	LOC105285601	E001	-0.28	0.06	0.51
		E002	0.18	0.06	0.5
ataxin-7	LOC105285602	E001	2.03	0.1	0.63
		E002	0.12	0.98	1
		E003	0.06	0.75	0.95
		E004	0.05	0.69	0.94
		E005	-0.04	0.48	0.87
		E006	-0.48	0.01	0.17
		E007	0.12	0.55	0.89
Kank	LOC105285603	E001	-0.93	0.24	0.79
		E002	-0.65	0.05	0.47
		E003	-0.14	0.72	0.94
		E004	-1.04	0.44	0.86
		E005	-1.6	0.29	0.81
		E006	-0.55	0.08	0.57
		E007	-0.67	0.04	0.4
		E008	-0.63	0.08	0.58
		E009	0.01	0.88	0.98
		E010	-0.15	0.34	0.83
		E011	-0.28	0.2	0.76
		E012	-0.27	0.2	0.76
		E013	-0.27	0.18	0.74
		E014	0.07	0.3	0.81

		E015	0.11	0.07	0.54
		E016	0.18	0.26	0.8
		E017	NA	NA	NA
Bet1	LOC105285604	E001	2.7	0.05	0.47
		E002	0	0.99	1
		E003	-0.04	0.67	0.93
		E004	-0.01	0.82	0.97
		E005	0.05	0.94	1
		E006	9.44	0.23	0.78
prob. ribosome prod. factor	LOC105285607	E001	0.24	0.77	0.96
		E002	-0.12	0.77	0.96
		E003	0.05	0.7	0.94
		E004	0	1	1
		E005	0.3	0.55	0.89
THUMP	LOC105285605	E001	1.26	0.04	0.42
		E002	-0.12	0.49	0.87
		E003	-0.09	0.55	0.89
		E004	0.03	0.79	0.96
		E005	0.13	0.19	0.75
		E006	-0.03	0.96	1
		E007	-0.09	0.76	0.95
		E008	-0.09	0.68	0.93
		E009	3.9	0.000257	0.00951
unchar. lncRNA	LOC105285608	E001	-1.46	0.08	0.56
		E002	0.28	0.06	0.5
Creld	LOC113561733	E001	-0.06	0.62	0.92
		E002	-0.01	0.86	0.98
		E003	0.09	0.32	0.82
		E004	-0.15	0.6	0.91
alphaCOP	LOC105283852	E001	-0.29	0.3	0.81
		E002	-0.23	0.33	0.83
		E003	-0.21	0.25	0.79
		E004	-0.15	0.16	0.73
		E005	0.14	0.19	0.75
		E006	0.32	0.3	0.81
		E007	0.74	0.14	0.69
unchar. protein	LOC105283851	E001	-0.07	0.76	0.96
		E002	0.14	0.76	0.96
Host cell factor	LOC105283850	E001	-0.34	0.28	0.8

		E002	-0.33	0.24	0.79
		E003	-0.25	0.38	0.84
		E004	-0.34	0.36	0.84
		E005	-0.19	0.5	0.88
		E006	-0.19	0.39	0.85
		E007	0.75	0.12	0.67
		E008	-0.23	0.33	0.82
		E009	0.3	0.39	0.85
		E010	-0.05	0.93	1
		E011	0.44	0.34	0.83
		E012	0	0.59	0.91
		E013	0	0.35	0.83
		E014	0.45	0.12	0.67
		E015	0.04	0.2	0.77
		E016	-0.35	0.56	0.9
		E017	0.14	0.07	0.55
		E018	0.11	0.48	0.87
		E019	0.35	0.38	0.84
l(2)37Cg	LOC105283849	E001	0.04	0.54	0.89
		E002	-0.03	0.6	0.91
		E003	0	0.94	1
ribosomal protein S2	LOC105283847	E001	0.1	0.54	0.89
		E002	-0.03	0.45	0.86
		E003	-0.06	0.63	0.92
		E004	3.68	0.00233	0.06
guanine deaminase	LOC105283846	E001	0.29	0.31	0.82
		E002	0.15	0.52	0.89
		E003	0.15	0.37	0.84
		E004	0.12	0.55	0.89
		E005	0.1	0.58	0.9
		E006	-0.06	0.62	0.92
		E007	-0.09	0.41	0.85
		E008	-0.17	0.18	0.75
		E009	-0.26	0.07	0.54
unchar. protein	LOC105283844	E001	-0.52	0.000763	0.02
		E002	-0.54	0.00000519	0.000311
		E003	-0.56	0.000378	0.01
		E004	-0.91	0.00693	0.13
		E005	-0.85	0.00588	0.11

	E006	-0.92	0.000107	0.00449
	E007	-0.91	0.00434	0.09
	E008	-0.8	0.00161	0.04
	E009	-0.53	0.00458	0.1
	E010	-0.54	0.0000288	0.00143
	E011	-0.46	0.000237	0.00887
	E012	0.57	7.31E-07	0.0000543
	E013	0.79	0.00000443	0.00027
	E014	1.11	1.32E-07	0.0000115
	E015	1.07	3.96E-07	0.000031
	E016	0.86	0.000705	0.0219
	E017	0.8	0.000814	0.0245
	E018	1.29	0.000153	0.00613
	E019	1.29	0.000454	0.0152
	E020	1.13	0.000177	0.00694
	E021	0.99	0.00261	0.06
	E022	-0.89	0.55	0.89
	E023	1.32	0.000135	0.00547
	E024	1.11	0.07	0.55
	E025	-0.57	0.44	0.86
	E026	-0.24	0.32	0.82
	E027	0.09	0.00973	0.165
	E028	0.21	0.0003	0.0107
	E029	-0.59	2.27E-08	0.00000229
	E030	-0.23	0.00000842	0.000479
	E031	0.51	0.00158	0.0418

968 **<u>Supplementary Table S6.</u> Differential gene expression near the CSD index peak.**

969 The log 2-fold change in expression levels between male and female samples, p-values

970 (likelihood ratio test), and adjusted p-values (Benjamini-Hochberg adjustment) for each exon of

971 17 genes in the vicinity of the CSD index peak. The p-values are "NA" for one of the exons of

972 LOC105285603 because there is negligible expression of that exon (basemean=0).

Analysis	Program	Version	Analysis Available At:
Genetic Mapping	Trimmomatic	v0.36	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	bwa	v0.7.12-r1039	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	GATK	v4.2	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	vcftools	v0.1.17	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	pyvcf	v0.6.8	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	pybedtools	v0.9.0	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	pandas	v1.1.5	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	numpy	v.1.19.5	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	bcftools	v0.1.19-96b5f2294a	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	samtools	v1.8	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	OrthoFinder	v2.5.5	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	scikit-allel	v1.3.5	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	blast	v2.5.0	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	mummer	v4.0.0rc1	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	python	v3.6.15	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	R	v4.3.2	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	karyoploteR	1.28.0	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	ggplot2	v3.5.0	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	ggbeeswarm	v0.7.2	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genetic Mapping	matplotlib	v3.3.4	https://github.com/kipdlacy/SexDeterminationMapping_Obiroi
Genome Annotation	pbmm2	v1.12.0	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome	stringtie2	v2.2.1	https://github.com/RockefellerUniversity/Obiroi_Ge

Annotation			neModels_2025
Genome Annotation	samtools	v1.6	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome Annotation	hisat2	v2.2.1	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome Annotation	transdecoder	v5.7.1	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome Annotation	squanti3	v3.5.1	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome Annotation	Repeatmasker	v4.1.7-p1	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome Annotation	FEELnc	v0.2.1	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome Annotation	BUSCO	v5.4.7	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome Annotation	CLIPflexR	v0.1.20	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome Annotation	miRDeep2	v0.1.3	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome Annotation	Pilfer	v1.0	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome Annotation	R	v4.0	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Genome Annotation	Bioconductor	v3.19	https://github.com/RockefellerUniversity/Obiroi_GeneModels_2025
Differential Gene Expression	multiqc	v1.9	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	RSEM	v1.2.28	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	STAR	v2.5.0a	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	R	v4.3.3	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	GenomicFeatures	v1.52.2	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	tximport	v1.28.0	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	DESeq2	v1.40.2	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq

Differential Gene Expression	EnhancedVolcano	v1.18.0	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	pheatmap	v1.0.12	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	dplyr	v1.1.4	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	reshape2	v1.4.4	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	tidyr	v1.3.1	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	tibble	v3.2.1	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Gene Expression	ggbreak	v.0.1.2	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Exon Usage	HTSeq	v2.0.5	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Exon Usage	DEXSeq	v.1.46.0	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq
Differential Exon Usage	stageR	v.1.22.0	https://github.com/jina-leemon/CSD_biroi_male-female-RNA-seq

973 **Supplementary Table S7. Versions of software used for genomics analyses.**