1

# BubR1 and Mad2 regulate adult midgut remodeling in *Drosophila* diapause

**Yuya Adachi[1], Hiroki Nagai[1,2], Takako Fujichika[3], Aya Takahashi[3], Masayuki Miura[1], and Yu-ichiro Nakajima[1]***

1. Graduate School of Pharmaceutical Sciences, The University of Tokyo, Japan
2. Institute of Science and Technology Austria, Austria
3. Department of Biological Sciences, Tokyo Metropolitan University, Japan

*Author for correspondence e-mail: nakaji97@g.ecc.u-tokyo.ac.jp

Tel: +81-3-5841-4863

**Text**:  28,227 characters

**Figures**: 5

**Supporting information**: Supplemental figures S1-S4, Supplemental table S1-S5

## Summary

Diapause is a survival strategy in which growth and aging are temporarily suspended, enabling animals to withstand unfavorable environments. Various insects, including the fruit fly *Drosophila*, enter reproductive diapause, or dormancy, in response to colder temperatures and/or shorter day lengths. During reproductive diapause, ovarian development halts, and non-reproductive organs also undergo remodeling at both morphological and metabolic levels: however, the mechanisms underlying this remodeling and its physiological impact remain largely unclear. Here, we show that the *Drosophila* adult midgut undergoes extensive remodeling in diapause, marked by a sustained suspension of growth due to the cell cycle arrest of intestinal stem cells (ISCs) that revert to normal upon returning to recovery conditions. During dormancy, BubR1 and Mad2, key regulators of mitosis, are highly expressed and localized in the cytoplasm of ISCs rather than at the kinetochore, and both BubR1 and Mad2 are essential for diapause-specific midgut remodeling. Furthermore, disruption of midgut growth arrest during diapause reduces the resistance to starvation in adult flies. Together, our findings identify a novel role for BubR1 and Mad2 in ISCs–promoting proper midgut remodeling during dormancy–and highlight the importance of this process for survival under adverse environments.

## Introduction

Wild animals have developed a wide array of survival strategies to cope with harsh environmental conditions such as temperature fluctuations, drought, and food shortages. Among these strategies, dormancy programs (e.g. hibernation, torpor, and diapause) are critical adaptations that involve a reduction in metabolic rate and body temperature, which conserves energy, extends lifespan, and increases the probability of survival for the species (Easwaran & Montell, 2023; Staples, 2016; Wilsterman et al., 2021). By temporarily halting growth and delaying organ development under stressful conditions such as starvation and low temperatures, dormancy helps reduce the energy expenditure required for growth and reproduction, and this synchronization of physiological processes with environmental conditions enhances the organism's ability to survive (Denlinger, 2022; Schiesari & O'Connor, 2013). Understanding the underlying mechanisms of dormancy may thus provide valuable insights into the regulation of stress resistance, aging, and reproduction.

Dormancy programs typically comprise three distinct phases: a preparatory phase in which environmental cues initiate dormancy; a dormancy phase characterized by reduced metabolism and activity; and a recovery phase wherein normal growth and activity resume (Gray et al., 1991; Harsimran Kaur Gill et al., 2017). To enter and exit dormancy at the appropriate time, animals must be able to successfully detect environmental conditions and coordinate internal physiological processes. As such, dormancy regulation generally requires central control mechanisms involving the brain and endocrine systems (Helfrich-Förster, 2024; Karp, 2021). While hormonal and neural regulation play pivotal roles in directing these processes, peripheral organs and tissues must also undergo significant remodeling during dormancy and subsequent reversion during recovery. For instance, during hibernation in certain mammals (e.g. thirteen-lined ground squirrels, Syrian hamsters) and diapause in insects, structural changes are observed in respiratory and digestive organs as well as in muscles and adipose tissues (Kim et al., 2006; Kim & Denlinger, 2009; Kurtz et al., 2021). Despite these substantial transformations and their vital importance, the mechanisms by which cellular compositions within individual organs and

tissues are precisely coordinated to adjust dormancy programs remain largely
unknown.

Animal species enter dormancy at different life stages, and in adult
insects, dormancy is often referred to as reproductive diapause, a state that
suppresses reproductive organ development and is associated with reduced
locomotor activity, lower metabolism, increased stress resistance, and extended
lifespan(Denlinger, 2022; Tatar & Yin, 2001). In species like *Drosophila*,
reproductive diapause is triggered by low temperatures and/or short-day
conditions (Hara & Yamamoto, 2022; Lirakis et al., 2018; Ojima et al., 2018;
Saunders et al., 1989). As with other diapausing insects, the reduction of juvenile
hormone and insulin signaling are critical for maintaining reproductive diapause
in *Drosophila* (Easwaran et al., 2022; Saunders et al., 1990; Schiesari et al.,
2016). Recent studies have used *D. melanogaster* to explore the genetic basis of
dormancy's complex biology and have resulted in the identification of circadian
neurons that regulates the endocrine organs, the mechanism of germline
maintenance, and sleeping behaviors (Easwaran et al., 2022; Kurogi et al., 2023;
Meiselman et al., 2022; Meyerhof et al., 2024; Schiesari et al., 2016). Interestingly,
similar to dormancy programs in other animals, somatic non-reproductive organs
such as the midgut undergo structural changes during reproductive diapause in
*Drosophila* (Kubrak et al., 2014); however, the mechanism by which the dormant
state regulates organ remodeling and its physiological impacts remain unclear.

The *Drosophila* adult midgut, composed of a monolayered epithelium
surrounded by visceral muscle, is home to resident intestinal stem cells (ISCs)
that self-renew and give rise to committed progenitors, enteroblasts (EBs) and
enteroendocrine progenitor cells (EEPs); these progenitors differentiate into
absorptive enterocytes (ECs) and secretory enteroendocrine cells (EEs),
respectively (Hung et al., 2020; Miguel-Aliaga et al., 2018; see also Fig. 2A, 2B).
To achieve adaptive responses to environmental changes at the organ level, the
ISC lineage exhibits characteristic behaviors that enable proper tissue
homeostasis and plasticity (Nagai et al., 2022; O'Brien et al., 2011). Furthermore,
the midgut plays a central role in regulating organism-level physiology,
influencing appetite, fecundity, immunity, and lifespan (Malita et al., 2022; White
et al., 2021; Yoshinari et al., 2024; Zhou et al., 2020). With its relatively simple

structure and conserved cell lineages, the *Drosophila* adult midgut serves an
ideal system for dissecting the mechanisms of organ remodeling associated with
environmental challenges like dormancy.

Here, we investigate the mechanistic underpinnings of organ remodeling
during diapause, using the *Drosophila* adult midgut as a model. Our results show
that intestinal growth halts during dormancy and resumes during recovery,
accompanied by changes in structural and metabolic gene expression. We
identify mitotic regulators BubR1 and Mad2, which are highly expressed in the
cytoplasm of ISCs, as key players in midgut remodeling during diapause, and
show that disruption of these genes leads to abnormal gut size and a reduced
capacity for starvation resistance. These findings suggest novel roles of BubR1
and Mad2 in controlling ISCs dynamics and contributing to midgut remodeling in
*Drosophila* diapause.

**Results**
**The adult midgut of *D. melanogaster* undergoes remodeling in response to**
**diapause and recovery**
In adult females of *D. melanogaster*, reproductive diapause can be induced by
exposing newly eclosed flies to cold temperatures (below 12°C), halting ovarian
development at an early stage (Anduaga et al., 2018). Following previous studies
(Easwaran et al., 2022; Kubrak et al., 2014; Saunders et al., 1989), we kept newly
eclosed flies at short-day photoperiods (10L:14D) and 10°C while non-diapause
control flies were kept at 12L:12D and 18°C (Fig. 1A). Under this suboptimal
condition, females were confirmed to have entered reproductive dormancy as no
yolk accumulation was observed in the ovaries (Fig. S1A). To assess changes in
midgut size during diapause, we measured the length and area of the midgut in
the wild-type fly (Canton-S) under diapause and non-diapause conditions for 1 to
5 weeks. Consistent with previous findings (Kubrak et al., 2014), we confirmed
that midgut length remained shorter during diapause compared to non-diapause
conditions (Fig. 1B, 1C), while becoming similar to that immediately after eclosion
(Fig. 1C). After one week of recovery from diapause, the midgut significantly
increased in size (Fig. 1B, 1D, Fig. S1B, S1C) and fully recovered by day 4 (Fig.

S1D). Of note, after 5 weeks of diapause, no significant midgut growth occurred

upon recovery (Fig. 1D), suggesting the existence of a time limit on the midgut's

ability to regrow. These findings together indicate that the midgut halts growth

during diapause but remains capable of rapid regrowth upon recovery.

The *Drosophila* adult midgut consists of anterior (R1-R2), middle (R3),

and posterior regions (R4-R5) (Buchon et al., 2013; Marianes & Spradling, 2013).

Since these regions respond differently to environmental changes due to the

divergent nature of ISCs and their progenies (Li & Jasper, 2016; Mattila et al.,

2024; Nagai et al., 2023; O'Brien et al., 2011), we analyzed midgut size changes

in the anterior and posterior regions under the 3-week diapause condition where

the difference between diapause and non-diapause midgut size is greatest (Fig.

1C). We found that the posterior midgut differed more significantly across

conditions (Fig. 1E) than the anterior midgut. Similar region-specific gut resizing

during diapause and recovery was also observed in the $w^{1118}$ strain (Fig. 1F),

suggesting that the posterior region is preferentially affected during dormancy. It

has been reported that reproductive development is also suppressed in

*Drosophila* adult males under the same conditions females enter diapause

(Kubrak et al., 2016). To assess diapause's effect on the midgut across sexes,

we measured midgut size in males under diapause and non-diapause conditions

but ultimately found no significant differences between the two (Fig. S1G). These

observations suggest that midgut remodeling in diapause is a sexually dimorphic

trait, appearing only in females.

To determine whether the observed midgut remodeling is specific to

diapause or is rather a general response to cold-induced stress, we exposed

female flies to the cold temperature conditions outlined above after the midgut

had matured and ensuring no diapause was induced (Ojima et al., 2018).

Comparing the midgut size after 3 weeks under cold and diapause conditions,

we found that midgut size was larger under cold conditions (Fig. S1E, S1F).

Furthermore, midgut growth was not observed when flies were transferred from

the cold condition to recovery, contrary to the transition observed during diapause

recovery (Fig. S1F). Altogether, these results suggest that midgut remodeling is

a diapause-specific response in adult females in which midgut growth is halted in

a regrowth-ready state. We term this midgut remodeling during diapause:

"dormancy-dependent midgut remodeling" (DDMR).

**Midgut remodeling as a common response in *Drosophila* species**

While the model organism *D. melanogaster* has recently been used as a genetic

model to study numerous aspects of diapause (Easwaran et al., 2022; Easwaran

& Montell, 2024; Meiselman et al., 2022; Ojima et al., 2018), it is still unclear

whether this species naturally enters diapause in the wild (Lirakis et al., 2018).

However, it has been established that some species of the genus *Drosophila*

undergo a programmed state of diapause in response to environmental changes

like short-day lengths that enables them to survive through the winter (Kimura,

1990; Lakovaara et al., 1972).

To explore whether DDMR is a general organ-level response during

reproductive dormancy, we examined two other *Drosophila* species: *D. bifasciata*

and *D. triauraria* (Fig. 1G), which inhabit northern regions and have been shown

to halt ovarian development in short-day seasons (Watabe, 1979; Yamada &

Yamamoto, 2011). We confirmed that these species exhibit reproductive

diapause under short-day conditions (SD, 10L: 14D), while such diapause was

not induced in long-day conditions (LD, 16L: 8D) at the same moderate

temperature (15°C) (Fig. 1H, S1H). In SD, midgut size was significantly smaller

compared to LD, and midgut regrowth was observed during recovery (Fig. 1I-1K).

Notably, this midgut growth arrest was not observed in non-diapause strains of

*D. triauraria* that originate from southern regions (Yamada & Yamamoto, 2011)

(Fig. S1I, S1J). These results support the notion that DDMR in *D. melanogaster*

represents a general organ-level response triggered during diapause in the

genus *Drosophila*.

**ISCs maintain cell cycle arrest and inhibition of mitosis during diapause**

In the adult midgut epithelium of *Drosophila*, ISCs serve as resident stem cells,

maintaining the self-renewing population and producing progenitors (EBs and

EEPs) that further differentiate into absorptive ECs and secretory EEs,

respectively (Fig. 2A, 2B; Hung et al., 2020; Miguel-Aliaga et al., 2018).

Previous studies have shown that alterations in cell number via ISC expansion

and reduction contribute to the substantial midgut size changes that correspond

with environmental fluctuations (Christensen et al., 2024; Nagai et al., 2022;

Stojanović et al., 2022). For example, after mating, juvenile hormones stimulate

ISC proliferation, leading to an increase in midgut size (Reiff et al., 2015).

Similarly, upon refeeding following starvation, ISC symmetric division is

promoted in the posterior midgut (O'Brien et al., 2011), resulting in ISC

expansion and a subsequent increase in total cell number as well as midgut

size.

To investigate whether changes in cell number contribute to midgut

remodeling in diapause, we first quantified the midgut cell abundance following

3 weeks of diapause. We observed a significant reduction in total cell number

during diapause compared to non-diapause conditions (Fig. 2C). We then

examined the number of ISC/EB lineage using the protein trap line *esg-GFP.*

The number of *esg+* cells, along with their relative proportion, was markedly

lower during diapause, especially in the posterior midgut (Fig. 2D, 2E, Fig.

S2A). Consistent with this observation, the ISC marker *Delta* (*Dl*) and the EB

marker *Su(H)* exhibited reduced expression in the diapausing midgut (Fig. S2B-

S2F). Moreover, *esg+* cell number dramatically increased during recovery from

diapause, coinciding with midgut regrowth (Fig. 2E). These findings suggest the

possibility that ISCs remain in a non-proliferative state during diapause and

resume proliferation upon recovery.

To test this possibility, we investigated ISC mitotic activity using the

anti-phospho-Histone H3 (PH3) immunostaining, a marker for mitotic cells.

Unexpectedly, we found that the number of ISCs in the M phase significantly

increased in both the anterior and posterior midgut during diapause (Fig. 2F,

2G). Despite the reduced number of ISCs (Fig. 2E, S2B, S2C), the increase in

mitotic cells suggests that ISCs may arrest the cell cycle in the M phase or,

alternatively, experience prolonged M phase during diapause.

In order to directly test ISC proliferative capacity, we conducted clonal

expansion assays using the mosaic analysis with a repressible cell marker

(MARCM) system (Fig. 2H) and found that clonal size was significantly reduced

during diapause, with fewer ISCs present in the clones (Fig. 2I, 2J). This mitotic

arrest was not observed during recovery from diapause, with the number of
mitotic cells returning to non-diapause levels (Fig. S2G).

To further examine the ISC lineage cell cycle state, we employed the
Fly-FUCCI system (Zielke et al., 2014) in combination with the *esg-QF* driver
during diapause. Using this system makes it possible to monitor the cell cycle at
cold temperatures, and we found that a significantly higher proportion of *esg*+
cells were in the G2/M phase during diapause compared to non-diapause
controls (Fig. 2K, 2L). Additionally, using the *CycB-GFP* line, we confirmed
increased *CycB* expression in ISCs during diapause (Fig. S2H), reinforcing the
idea that ISCs arrest in the G2/M phase. These results suggest that ISCs
undergo cell cycle arrest in the G2/M phase during dormancy.

Combined, our findings reveal that ISCs undergo drastic behavioral
changes during DDMR–that they effectively halt the cell cycle at a recoverable
state, reducing the number of cells produced post-eclosion and allowing for
proper size change during diapause.

**Molecular signatures of DDMR**
To explore the molecular signature of DDMR, we conducted a transcriptome
analysis of the midgut. Specifically, bulk RNA sequencing (RNA-seq) was
performed on samples of the entire adult midgut from two conditions: 3 weeks
diapause (3wD) and 3 weeks non-diapause control (3wC). We identified 3,391
genes that were differentially expressed during dormancy (Fig. 3A). Gene
Ontology (GO) enrichment analysis and Reactome pathway analyses revealed
that metabolic processes, including small molecule, lipid, and carboxylic acid
metabolism were significantly enriched in the 3wD midgut (Fig. S3A, S3B),
suggesting that large-scale metabolic remodeling occurs alongside structural
changes in the midgut during dormancy. Lipid metabolism-related genes such as
*bmm* and *pepck*, along with carbohydrate metabolism-related genes like *tobi*,
were differently expressed (Fig. S3C). Additionally, pathways related to
metabolism (e.g., Metabolism of Lipids), translation regulation (e.g., Translation),
and circadian gene degradation (e.g., Degradation of TIM) were among the top
processes identified (Fig. S3B). Notably, circadian genes *per* and *tim*, which are
upregulated in the entire body during diapause (Kučerová et al., 2016), also

showed significantly increased expression in the midgut during dormancy (Fig.
S3D). Furthermore, the expression of the antimicrobial peptide *Drs*, downstream
of the Toll pathway, dramatically increased during dormancy (Fig. S3E), reflecting
enhanced innate immunity, a hallmark of diapause at the organismal level
(Kubrak et al., 2014).

A detailed enrichment analysis was performed on the differentially
expressed genes, by categorizing the genes into two groups: those upregulated
during dormancy and those downregulated. In the upregulated gene group, GO
terms associated with cell adhesion and RNA processing (e.g., regulation of RNA
splicing) were identified as top-ranked categories (Fig. 3B). Genes associated
with RNA processing and RNA metabolism are also upregulated in the ovaries
during diapause, suggesting the existence of dormancy-specific translational
regulation conserved across organs (DiVito Evans et al., 2023). In the
downregulated gene group, by contrast, GO terms related to metabolic processes
(e.g., cellular lipid metabolic process) were prominently enriched (Fig 3C). During
diapause, extensive changes in the expression of genes involved in metabolic
pathways have been observed throughout the body (Kubrak et al., 2014;
Kučerová et al., 2016), and these findings suggest that gene expression changes
linked to DDMR globally follow a pattern similar to that observed in the entire
organism and other organs.

Next, to explore the molecular signature of DDMR recovery, RNA-seq
was performed using samples of the entire adult midgut from two conditions: 3
weeks diapause followed by recovery (3wD+R) and 3 weeks control followed by
recovery (3wC+R). We identified 526 genes with significant expression changes
between the 3wD+R and 3wC+R midguts (Fig. S3G). GO enrichment and
Reactome pathway analyses identified significant enrichment of genes
associated with muscle differentiation and smooth muscle contraction (Fig. S3H,
S3I). Given the extensive structural remodeling of muscles during insect
dormancy (Kim et al., 2006; Kim & Denlinger, 2009), these gene expression
changes related to muscle structure during recovery may contribute to the
resizing of the midgut. Consistent with this finding, *dilp3*, which codes the product
secreted by muscles during feeding and promote ISC proliferation (O'Brien et al.,
2011), was significantly upregulated in the 3wD+R midgut (Fig. S3J), suggesting

growth-promoting interactions between muscle and ISCs during recovery. In
addition, neuropeptide genes such as *NPF, Dh31,* and *CCHa2* showed significant
increases in expression (Fig. S3J), indicating that hormone secretion from EEs
may change during recovery, potentially influencing gut growth and inter-organ
communications. Overall, our findings suggest that the midgut undergoes large-
scale structural and metabolic remodeling associated with extensive gene
expression changes.
320
**BubR1 expression increases in the cytoplasm of ISCs during diapause**
Because ISC cell cycle are arrested during diapause (Fig. 2), we focused on
differentially expressed genes categorized under the GO term "cell cycle
checkpoint signaling" (Fig. 3D). G2/M phase checkpoint regulators such as
*mre11*, *mei-41* showed significant changes while *CG15237* and *BubR1* were
differentially expressed as M-phase checkpoint regulators. These gene
expression changes align with ISC behaviors during dormancy such as
prolonged mitotic phases and an increase in G2/M phase cells. Among these
genes, *BubR1* is particularly noteworthy due to its established roles as a mitotic
checkpoint regulator and mitosis timer (Karess et al., 2013) and its expression
in ISCs (Shinoda et al., 2023). Importantly, *BubR1* expression increased nearly
threefold during diapause (Fig. 3E), suggesting a major potential impact on ISC
behaviors and midgut remodeling.
To investigate *BubR1* expression at the cellular level, we used fly lines,
which allow for the detection of the endogenous expression pattern of *BubR1*
(*EGFP::BubR1^{WT}* and *GFP-BubR1*) to visualize its expression in midgut cells.
During reproductive dormancy, *BubR1* was strongly expressed in the cytoplasm
of ISCs, primarily in R4 regions (Fig. 3F, 3G, S3F). This cytoplasmic localization
during diapause contrasts with the localization of BubR1 associated with typical
spindle checkpoint during mitosis, where it accumulates around kinetochores
(Resende et al., 2018). Indeed, colchicine treatment, which induces mitotic
arrest through the spindle assembly checkpoint (SAC), caused strong BubR1
localization around kinetochores in ISCs under non-diapause conditions (Fig.
3H, 3I). In contrast, during diapause, despite ISCs remaining arrested in
mitosis, BubR1 remained cytoplasmic, with minimal signal around the

kinetochores (Fig. 3H, 3I). These findings suggest that the role of BubR1 during

DDMR is likely spindle checkpoint-independent and constitutes a unique

function in the midgut during the dormancy state.

**BubR1 expression in ISCs is essential for DDMR**

To explore the function of BubR1 in reproductive dormancy, we examined its

effects on midgut remodeling using two *BubR1* mutant alleles *BubR1$^{k06109}$* and

*BubR1$^{k03113}$*. qPCR analysis revealed that midgut *BubR1* expression in

diapause-induced flies was significantly reduced in both heterozygous mutants,

with *BubR1$^{k03113}$* showing 48% and *BubR1$^{k06109}$* showing 57% expression

compared to controls (Fig. 4A). We then measured midgut size during diapause

in these mutants and found a significant increase in midgut size compared to

control flies (Fig. 4B, 4C). This midgut enlargement was accompanied by an

increase in the total cell number, the number of mitotic ISCs, as well as the

number of *esg+* cells (Fig. 4D-4F). Importantly, when BubR1 function was

rescued in the *BubR1* trans-heterozygous mutant (*BubR1$^{k06109}$/BubR1$^{k03113}$*)

with an *EGFP::BubR1$^{WT}$* transgene (Shinoda et al., 2023), the total cell number

in the posterior region and mitotic cell number in the midgut returned to levels

comparable to controls (Fig. 4C-4E). These results strongly suggest that BubR1

is required for proper midgut remodeling during diapause. Notably, no ovarian

development was observed in any of the dormant mutants, confirming the

successful induction of ovarian dormancy despite impaired midgut remodeling

(data not shown). Furthermore, no significant differences in midgut size were

detected between these mutant backgrounds under non-dormant conditions

(Fig. S4A), consistent with the result that the number of mitotic cells did not

increase—or even decreased— in *BubR1* mutants (Fig. S4B). Collectively,

these results highlight a critical role for BubR1 in DDMR.

The results obtained from the mutants could potentially reflect non-

autonomous effects of *BubR1* expressed in other tissues. We thus investigated

whether DDMR is influenced by BubR1 activity specifically in adult ISCs. To this

end, we employed somatic tissue-specific CRISPR-mediated depletion (Bier et

al., 2018; Meltzer et al., 2019), combined with the temperature-sensitive Gal80$^{ts}$

system (McGuire et al., 2003) to induce *BubR1* mutations in ISC lineages

before eclosion (Fig. 4G, S4C). When adult flies were shifted to dormancy
conditions, we observed a significant increase in midgut size in ISC/EB-specific
*BubR1* knockout flies (Fig. 4H). Furthermore, the number of total and mitotic
cells also increased in these somatic *BubR1* knockout lines (Fig. 4I, 4J).
Altogether, our results demonstrate that *BubR1* function in ISCs is essential for
midgut remodeling during reproductive diapause.
Intriguingly, in *D. bifasciata* and *D. triauraria,* which undergo a
programmed state of diapause in response to short-day length, *BubR1*
expression in the midgut is higher in diapausing flies under SD compared to
non-diapausing flies under LD (Fig. S4D, S4E). These results suggest a
potential role of BubR1 in DDMR for the genus *Drosophila* more broadly.
390

**The MCC component Mad2 is involved in DDMR**

During mitosis, BubR1 forms a complex with Mad2, Bub3, and Cdc20, which
constitute key components of the mitotic checkpoint complex (MCC) and
function cooperatively at the kinetochores (Musacchio, 2015). Our RNA-seq
analysis indicated that, in addition to *BubR1,* the expression levels of *mad2*
increased significantly during dormancy (Fig. 5A), though it was not identified as
a differentially expressed gene. Intriguingly, Mad2 can interact with BubR1 in
the cytoplasm of mammalian cells to regulate insulin signaling and metabolism
(Choi et al., 2016). This raises the possibility that Mad2 may play a critical role
in DDMR, independently of its role in the MCC.
To assess this potential role, we utilized the *GFP-Mad2* line to examine
*mad2* expression in the adult midgut and found it to be highly expressed in the
cytoplasm of ISCs during diapause (Fig. 5B, 5C), reminiscent of the cytosolic
localization of BubR1 (Fig. 3F, S3F). To further investigate the function of Mad2
during DDMR, we analyzed midgut size using two *mad2* mutant alleles: *mad2$^{\Delta}$*
and *mad2$^{EY21687}$*. Our analysis revealed that heterozygous *mad2* mutants had
significantly larger midguts during dormancy compared to controls (Fig. 5D, 5E).
We also found that the number of *esg*+ cells increased in the anterior and
posterior region of the midgut (Fig. 5F). These findings suggest that during
reproductive diapause, Mad2 is localized in the cytoplasm of ISCs similarly to
BubR1 and that the two proteins together regulate DDMR.

**DDMR enhances starvation resistance**

Halting organ growth during dormancy is thought to conserve energy, thereby enhancing survival during periods of food scarcity, such as winter (Denlinger, 2022). In *Drosophila*, maintaining midgut size during reproductive diapause may contribute to increased resistance to starvation. To test this hypothesis, we compared starvation resistance between dormant *BubR1* mutants and controls. Although dormancy generally enhances starvation resistance compared to non-dormant states, *BubR1* $^{k03113}$ mutant flies exhibited significantly reduced starvation resistance during dormancy (Fig. 5G). These results suggest that BubR1/Mad2-mediated DDMR plays a critical role in enhancing starvation resistance at the organismal level.

## Discussion

In this study, we demonstrate that BubR1 and Mad2 play crucial roles in DDMR in *Drosophila* by modulating ISC dynamics to achieve diapause-specific states. While BubR1 and Mad2 are well-known as components of the MCC that localizes to the kinetochore, we found that in the diapausing midgut, they are predominantly localized in the cytoplasm of ISCs. Depletion of these proteins causes misregulated midgut growth and leads to reduced starvation resistance during diapause. Based on these findings, we propose that BubR1 and Mad2 act as key regulators of the ISC cell cycle during diapause, enabling a reversible state that facilitates recovery (Fig. 5H).

The *Drosophila* midgut undergoes extensive remodeling in response to various internal and external stimuli that directly or indirectly influence ISC activity (Christensen et al., 2024; Nagai et al., 2022; Stojanović et al., 2022). During starvation, for instance, the midgut size decreases as ECs undergo apoptosis, while upon refeeding, ISCs are stimulated by Dilp3 secreted from visceral muscles, promoting symmetrical ISC division and restoring midgut size (O'Brien et al., 2011). Similarly, during reproductive diapause—wherein food intake is reduced by less than half (Kubrak et al., 2014)—mild starvation responses may also contribute to midgut size reduction. Indeed, reproductive

arrest can be more readily induced by adding starvation conditions (Ojima et al.,

2018). Our RNA-seq data revealed increased *dilp3* expression and decreased

expression of the FoxO target gene *Thor* during recovery (Fig. S3J), suggesting

the possibility that Dilp3 secreted from muscles contributes to ISC activation as

it does in post-starvation refeeding. However, in contrast to the simple

starvation response, midgut cell death is minimal during diapause (Fig.S2I), and

ISC cell cycles are arrested at G2/M in a recoverable state (Fig. 2).

Furthermore, during reproductive arrest, germline stem cells in ovaries exhibit

behaviors markedly different from those observed during amino acid deficiency

responses (Easwaran et al., 2022). Collectively, these findings suggest that

organ-level responses during diapause are driven by a complex interplay of

environmental factors rather than by a simple starvation-induced mechanism.

Across species, developmental diapause typically involves a temporary

halt in the cell cycle that delays organ growth. For example, during pupal

diapause in the *Sarcophaga crassipalpis* brain and embryonic diapause in the

turquoise killifish, the cell cycle primarily arrests at the G0/G1 phase (Meller et

al., 2012; Tammariello & Denlinger, 1998). By contrast, most cells in silkworm

embryos are arrested in the G2 phase during diapause (Nakagaki et al., 1991).

In this study, we observed a significant increase in ISCs arrested at the G2/M

phase during DDMR. *Drosophila* oogenesis arrests during diapause due to

activation of the Chk2 DNA damage checkpoint (Easwaran et al., 2022), and

our findings suggest that similar mechanisms may govern growth arrest in the

midgut. Indeed, genes like *mre11* that are differentially expressed in the

diapausing midgut (Fig. 3D) are required for G2/M checkpoint activation (Song,

2005). Notably, ISCs exhibit active cell cycle arrest during diapause via

checkpoint mechanisms, rather than undergoing a passive delay induced by

low temperatures (Fig. S1E-S1F). This controlled cell cycle arrest in resident

stem cells may contribute to the pausing of organ growth in other diapausing or

hibernating animals and to facilitating rapid recovery once stem cell division

resumes.

BubR1 is a highly conserved eukaryotic protein that functions as a key

component of the SAC, forming the MCC with Mad2, Bub3, and Cdc20 to

prevent premature progression of cell division and ensure accurate

chromosomal segregation, and its disruption increases the risk of aneuploidy

(Baker et al., 2013; Karess et al., 2013; Resende et al., 2018). Beyond these

canonical functions, studies across cellular and organismal contexts have

uncovered additional roles for BubR1. For example, in mouse embryonic

fibroblasts, *BubR1* heterozygosity reduces *p53* and *p21* expression after DNA

damage, impairing proper G2/M phase arrest (Fang et al., 2006). Additionally,

BubR1, together with Mad2, recruits adaptor protein 2 to the insulin receptor

(IR) to promote IR endocytosis and reduce insulin sensitivity (Choi et al., 2016;

Park et al., 2023; Tang et al., 2020). Consistent with these findings, *BubR1*

deficiency in mice leads to premature aging and reduced lifespan, highlighting

its role in organism-wide responses (Baker et al., 2004). In *Drosophila*, BubR1

in the fat body negatively regulates *bmm* expression via IMD signaling during

starvation, suppressing lipid breakdown and enhancing lifespan (Liu et al.,

2024). These findings delineate a complex regulatory network involving BubR1

and associated proteins across different biological contexts. Although the

regulation and interacting partners of BubR1 and Mad2 during diapause have

yet to be identified, the observed changes in their expression in ISCs likely

contribute to organism-level responses, including enhanced starvation

resistance mediated by structural and metabolic changes in the midgut. Future

investigations into these emerging additional roles of BubR1/Mad2 will enhance

our understanding of organ remodeling in dormancy programs and provide new

insights into the control of resident stem cells.

## Acknowledgements

503    We thank R. Giet, B. Ohlstein, C. Sunkel, N. Shinoda, BDSC, Kyoto Stock Center,

504    NIG-FLY, KYORIN-Fly, and Developmental Studies Hybridoma Bank (DSHB) for

505    fly stocks and reagents; Y. Hara and S. Kondo for discussion.

## Funding

This work was supported by JSPS/MEXT KAKENHI (grant numbers JP22J01430

to H.N., JP21H04774, JP23H04766, JP24H00567 to M.M., and JP17H06332,

JP22H02762,     JP23K18134,     JP23H04696     to     Y.N.),     AMED-PRIME

(JP22gm6110025 to Y.N.), The Mochida Memorial Foundation for Medical and

Pharmaceutical Research (Y.N.), The Cell Science Research Foundation (Y.N.),

and Sadako O. Hirai Ban Award for Young Researchers (H.N.)

## Author contributions

Conceptualization: Y.A., Y.N.

Investigation: Y.A., H.N., Y.N.

Methodology: Y.A., H.N., T. F., A.T., Y.N.

Validation: Y.A., H.N., M.M., T. F., A.T., Y.N.

Data curation: Y.A., Y.N.

Writing – original draft: Y.A., H.N., Y.N.

Writing – review & editing: Y.A., H.N., A.T., M.M., Y.N.

Supervision: M.M., Y.N.

Funding acquisition: H.N., M.M., Y.N.

## Declaration of interests

527     The authors declare no competing interests.

## Materials and methods

### *Drosophila* stock and maintenance

All stocks were maintained on a standard diet containing 4% cornmeal, 6%

baker's yeast (Saf Yeast), 6% glucose (Wako, 049-31177), and 0.8% agar

(Kishida chemical, 260-01705) with 0.3% propionic acid (Tokyo Chemical

Industry, P0500) and 0.05% nipagin (Wako, 132-02635).

Canton-S was utilized as the wild-type strain. Transgenic fly lines were

obtained from the Bloomington Drosophila Stock Center (BDSC), NIG-Fly and

the Kyoto Stock Center. Unless otherwise indicated, strain descriptions can be

found at Flybase (http://flybase.bio.indiana.edu): *Dl-lacZ* (Röttgen et al.,

1998) (BDSC 11651), *esg-QF2* (Nagai et al., 2023)*, QUAS-Fly-FUCCI* (Zielke et

al., 2014) (BDSC 55108), *BubR1$^{k03113}$* (Kyoto 102258), *BubR1$^{k06109}$* (Kyoto

102444), *mad2$^{EY}$* (Bellen et al., 2011) (BDSC 22495), *tub-Gal80$^{ts}$* (BDSC 7017,

7019, described in Flybase and BDSC), *esg-Gal4* (Goto & Hayashi, 1999)

(Kyoto 109126), *UAS-GFP* (BDSC 1521), *UAS-Cas9.T2A.eGFP* (Trivedi et al.,

2020) (BDSC 94298), *FRT40A* (originated from BDSC 5074), *attP40{U6-gRNA}*

(NIG 2RG-1578). The following lines were gifts from fly community: *esg-

GFP[P01986]* (Le Bras & Van Doren, 2006), *Su(H)GBE-lacZ* (Furriols & Bray,

2001), *yw hs-flp; tub-Gal80 FRT40A; tub-Gal4 UAS-dsRed /TM6B* (Guo et al.,

2013)*, EGFP::BubR1$^{WT}$* (Shinoda et al., 2023)*, GFP-BubR1, GFP-Mad2* (Buffin

et al., 2005)*, mad2$^{Δ}$* (Buffin et al., 2007)*, CycB-GFP* (Huang & Raff, 1999). The

following other *Drosophila* species were obtained from KYORIN-Fly: *D.

bifasciata* (JZK3), *D. triauraria* (TM15-92), *D. triauraria* (KMJ1), *D. triauraria*

(OEB12)*. See Additional file: Table S1 for the genotypes in each figure. We

used female flies unless otherwise noted in the figures.

**Diapause and recovery induction**

To assess the effects of diapause on midgut growth using *D. melanogaster*, we

subjected  old virgin females less than 6 h to 10°C and 8 L:16D photoperiods

(Diapause condition), and 18°C at 12L:12D photoperiods (Control condition) for

1-5 weeks. We then allowed the flies to recover by transferring them into vials

with fresh food with dry yeast for 7 days at 25°C and 12L:12D photoperiods

(Recovery condition).

To assess the effects of the cold on midgut size, we allowed the flies to

grow in 18°C for 4 days after eclosion, and then subjected them to 10°C and

8 L:16D photoperiods for 3 weeks (Cold condition).

To assess the effects of diapause state on midgut growth in other

*Drosophila* species, we subjected virgin females less than 6 h old to 10 L:14D

photoperiods (SD, diapause) and 16L:8D photoperiods (LD, control) for 16 days.

The rearing temperature was kept consistent at 15°C. We then allowed the flies

to recover by transferring them into 16L:8D photoperiod condition.

**MARCM clone induction scheme**

For the mosaic analysis with a repressible cell marker (MARCM) analysis, MARCM-ready flies (*hsFLP; tub-Gal80,FRT40A/FRT40A; tub-Gal4,UAS-dsRed*) were cultured at 25°C throughout pupal development, except during heat shock induction. When female larvae began metamorphosis, white pupae were placed into an empty vial at 25°C and designated as being at 0 h after puparium formation (APF). MARCM labels one of the two daughter cells after mitotic division. To label dividing ISCs, MARCM-ready flies were heat shocked at 48 h APF for 1 h (the time during which ISCs undergo asymmetric divisions (Guo & Ohlstein, 2015)). After eclosion, the size of the growing ISC clones was observed with anti-Pros immunofluorescence staining under 1 week diapause or 1 week control conditions. EE-only clones, which can be induced by labeling daughter EE during asymmetric division were excluded by immunostaining against the EE marker Pros, and the size of clones containing two or more ISCs was quantified.

**Tissue-specific somatic mutation of BubR1**

Transgenic flies expressing BubR1-sgRNA (GGTGACGCCAGTTACTATTG) obtained from NIG gRNA stock collections were crossed to flies expressing *esg^{ts}>Cas9* (*esg-Gal4 tubGal80^{ts} / CyO; UAS-Cas9 / TM3, Act-GFP*). Experimental crosses were maintained at 18°C to utilize Gal80^{ts} for suppressing Gal4 activity. When female larvae began metamorphosis, white pupae were placed into an empty vial at 18°C. After 6 days, when ISC division was completed, pupae were transferred to 29°C and reared for 1 day to introduce somatic mutations. These pupae were then transferred back to 18°C and reared under diapause or control conditions for 3 weeks after eclosion.

**Immunofluorescence**

Adult midgut samples were dissected in 1 × PBS and fixed in 4% PFA for 30–45 min at room temperature (RT). The following primary antibodies were used with indicated dilution into 1 × PBS containing 0.5% BSA and 0.1% Triton X-100: rabbit anti-PH3 (Millipore 06–570, 1:1000), mouse anti-Prospero (DSHB MR1A, 1:100), rat anti-GFP (Nacalai tesque 04404–26, 1:400), rabbit anti-cDcp1 (Cell Signaling Technology 9578, 1:200), chicken anti-β-galactosidase (Abcam ab9361, 1:500), and mouse anti-Delta (DSHB C594.9B, 1:100). After overnight incubation

606 with primary antibodies at 4°C, samples were incubated with fluorescent
607 secondary antibodies (Jackson ImmunoResearch and Invitrogen, 1:500) for 1 h
608 at RT. Hoechst 33,342 (Invitrogen, final concentration: 10 μg/ml) was used to
609 visualize DNA. Samples were mounted in Slowfade Diamond (ThermoFisher,
610 S36963) and imaged with a Zeiss LSM880 confocal microscope.
611

**RT-qPCR**
Total RNA was purified from 7-10 midguts using the ReliaPrep RNA Tissue
Miniprep System (Promega). cDNA was made from 100 or 200 ng of RNA using
PrimeScript RT Reagent Kit (TaKaRa). Quantitative PCR was performed using
TB Green Premix Ex Taq II (TaKaRa) and the QuantStudio 6 Flex Real-Time PCR
System (ThermoFisher). *RpL32* was used as an internal control. Primers for *D.*
*bifasciata* and *D. triauraria* were designed based on the sequence homology to
*D. melanogaster* transcripts. Primer sequences are listed in Table S2.
620

**Bulk RNA-seq**
Total RNA was purified from 10 to 15 midguts using the ReliaPrep RNA Tissue
Miniprep System (Promega). A NanoDrop (ThermoFisher) was used to measure
the concentration of each RNA sample, and the integrity of RNA samples was
checked using a Bioanalyzer (Agilent). A total of 300 ng RNA for each sample
was used for cDNA library preparation and the subsequent sequencing was
performed on an DNB sequencing system with single reads of 50 base pairs. The
sequencing reads were aligned against the *Drosophila melanogaster* genome
(ENSEMBL, BDGP6.22) using Salmon (Patro et al., 2017) for transcript
quantification. The quantification data of each transcript was converted into gene-
level counts using Tximport (Soneson et al., 2015). Differential expression
analysis of two conditions/group (four biological replicates per condition) was
performed using the DESeq2 R package (ver 3.20). Genes with an adjusted *P*-
value < 0.1 were assigned as differentially expressed. For functional cluster
analysis of DEGs, GO BP enrichment analysis and Reactome pathway analysis
were performed with the ClusterProfiler R package.
637

**Colchicine treatment**

0.2 mM Colchicine was dissolved in 5% (w/v) sucrose solution. Filter paper

(Whatman 3MM) was soaked with 300 μL of reagents and placed into empty vials.

Newly-eclosed flies were transferred into these vials and fed with 5% sucrose

only (control) or colchicine solutions for 1 day.

**Starvation resistance**

To assay the starvation resistance, flies were fed with agar only food. For the

survival assay, dead flies (determined by immobility and a lack of response to

tapping) were counted daily.

**Quantification of midgut size**

The central line length and area of the midgut were quantified using the

Fiji/ImageJ free hand tool on images obtained with a Zeiss AXIO Zoom.V16

stereomicroscope.

**Quantification of cell number**

Cell number was quantified using Fiji/ImageJ. To count the total cell and *esg-*

*GFP*$^+$ cell number, Hoechst and GFP signal was processed as follows: (1)

despeckle, (2) binarization, (3) watershed, (4) analyze particle. Cell counts for

PH3$^+$ cells, *Su(H)GBE-lacZ*$^+$ cells, and Fly-Fucci$^+$ cells were manually counted

using the cell counter function.

For MARCM clone size quantification, clones containing two or more

ISCs were selected, and the number of cells in the clone was counted manually.

**Quantification of signal intensity**

The signal intensity of reporter lines (*EGFP::BubR1*$^{WT}$*, GFP-Mad2*) was

quantified using Fiji/ImageJ. The signal areas were selected as ROIs using the

polygon selection tool. The reporter intensity in each ROI was quantified using

the Measure command.

**Statistics**

Statistical analyses were performed using Excel and RStudio. Two-tailed *t* tests

were used for comparisons between two groups. One-way ANOVAs with post hoc

Tukey tests were performed when comparing three or more groups. Log-rank

tests were used for survival curve comparison. Significance is indicated in the

figures as follows: *$P \le 0.05$, **$P \le 0.01$, ***$P \le 0.001$, not significant

(N.S.): $P > 0.05$. Bar graphs show mean ± standard error. Boxplots show median

(thick line in the box), first and third quartiles (bottom and top of the box),

minimum value (lower whisker), and maximum value (upper whisker). Dots in bar

graphs and boxplots indicate individual values. Violin plots portray the distribution

of individual values.

**Figure legends**

**Figure 1. The adult midgut of *Drosophila* undergoes remodeling in**

**response to diapause and recovery**

(A) Schematic of diapause and recovery in *D. melanogaster*. LD = Light : Dark.

(B) Representative images of the midgut under the 3 weeks diapause (3wD) and

recovery (3wD+R) condition. The two orange lines indicate anterior and posterior

end of the midgut. Scale bars: 1 mm.

(C, D) Quantification of midgut length of wild-type flies (Canton-S) under

diapause and control conditions (C) and their recovery conditions (D). $n$=10-15

midguts for each condition. The same diapause samples are quantified in (C) and

(D).

(E) Quantification of midgut length and area in the anterior and posterior region.

Canton-S was analyzed. $n$=6 (3wC), 9 (3wD), and 21 (3wD+R) midguts for length

measurements and $n$=15 (3wC), 20 (3wD), 17(3wD+R) midguts for area

measurements.

(F) Quantification of midgut length and area in the anterior and posterior region.

$w^{1118}$ was analyzed. $n$=10 (3wC), 10 (3wD), 15 (3wD+R) midguts for length

measurements and $n$=10 (3wC), 11 (3wD), 15 (3wD+R) midguts for area

measurements.

(G) Female adult of *D. bifasciata* (JZK3) and *D. triauraria* (TM15-92)*.* Scale bars:

1 mm.

(H) Schematic of diapause and recovery conditions in *D. bifasciata* and *D.*

*triauraria.* Light: Dark= 16 : 8 (hours) for long day (LD) condition, 10 : 14 (hours)

for short day (SD) condition.

(I) Representative images of the midgut in *D. bifasciata* (JZK3) under control (LD),

diapause (SD), and recovery conditions. Scale bars: 1 mm.

(J, K) Quantification of midgut length in *D. bifasciata* (J) and *D. triauraria* (K). *n*=7-

12 midguts for each condition.

**Figure 2. ISCs maintain cell cycle arrest and mitotic inhibition throughout**

**diapause**

(A, B) Schematics of the structure (A) and cell types (B) of the adult midgut of

*D. melanogaster*. An intestinal stem cell (ISC) generates an enteroblast (EB) or

an enteroendocrine progenitor (EEP), which differentiates into an enterocyte

(EC) or an enteroendocrine cell (EE), respectively.

(C) Total cell number in the adult midgut of wild-type flies under 3 weeks control

(3wC), 3 weeks diapause (3wD), and recovery after 3 weeks diapause (3wD+R).

*n*=7 (3wC), 8 (3wD), 8 (3wD+R) midguts.

(D) Representative images of *esg-GFP*$^+$ cells in the posterior midgut under 3wC

and 3wD conditions. Scale bars: 100 μm.

(E) Quantification of *esg-GFP*$^+$ cell number. *n*=11 (3wC), 12 (3wD), 10 (3wD+R)

midguts.

(F) Representative images of anti-PH3 staining in 3wC and 3wD wild-type flies.

Scale bars: 100 μm.

(G) PH3+ cell number in the anterior and posterior regions of 3wC and 3wD flies.

*n*=6 (3wC), 8 (3wD) midguts.

(H) Schematics of MARCM clone induction. ISC was labeled before eclosion, and

clonal expansion was examined at 1 week after eclosion.

(I) Representative images of MARCM clones in 1wC and 1wD conditions. A single

clone derived from an ISC is enclosed within the orange lines. Scale bars: 50 μm.

(J) Quantification of clone size. *n*=21 (1wC),10 (1wD) clones.

(K, L) Representative images of the Fly-Fucci signals in the progenitor (esg+)

cells (K). Scale bars: 100 μm. The proportion of G2/M (GFP+ and RFP+ cells) is

higher under the 3wD condition (L). *n*=7 (3wC), 5 (3wD) midguts.

**Figure 3. BubR1 expression increases in the cytoplasm of ISCs during**

**diapause**

(A) MA plot of differentially expressed genes in a pair-wise comparison of 3wC

and 3wD conditions.

(B, C) Gene ontology (GO) enrichment for upregulated genes (B) and

downregulated genes (C) in 3wD.

(D) Cell-cycle-related GO terms (biological process, BP) enriched in 3wD (left)

and heat map of "cell cycle checkpoint signaling" genes (right). These GO terms

were identified through enrichment analysis of all differentially expressed genes.

(E) Expression level of *BubR1* in 3wC and 3wD midguts from RNA-seq data.

Count data was normalized with RLE (Relative Log Expression). *n*=4 (3wC), 4

(3wD) RNA samples.

(F) Representative images of EGFP::BubR1 in the midgut under 3wC and 3wD

conditions. Scale bars: 50 μm.

(G) Quantification of the EGFP::BubR1 signal intensity of (F). 30 cells from 4

midguts (3wC), 41 cells from 4 midguts (3wD).

(H) Representative images of BubR1 localization in ISCs. BubR1 is localized in

the cytosol under the 3wD condition while BubR1 is accumulated at the

kinetochore (KT) after colchicine treatment. Scale bars: 10 μm.

(I) The proportion of BubR1 localization in mitotic (PH3+) cells. 215 cells from 6

midguts (3wD), 31 cells from 4 midguts (Colchicine fed fly).

**Figure 4. BubR1 expression in ISCs is essential for DDMR**

(A) RT-qPCR of *BubR1* in control and *BubR1* heterozygous mutant flies under

the 3wC and 3wD condition. *n*=4 (3wC, *w$^{1118}$*), 3 (3wC, *BubR1$^{k06109}$/+*), 4 (3wC,

*BubR1$^{k03113}$/+*), 4 (3wD, *w$^{1118}$*), 4 (3wD, *BubR1$^{k06109}$/+*), 4 (3wD, *BubR1$^{k03113}$/+*)

RNA samples.

(B) Representative images of the midgut in control and *BubR1* heterozygous

mutant flies under the 3wD conditions. Scale bars: 1 mm.

(C) Quantification of the midgut length in control, *BubR1* heterozygous mutant,

and *EGFP::BubR1$^{WT}$* flies under 3wD. In *EGFP::BubR1$^{WT}$* flies, wild-type form of

*BubR1* is BAC rescued in the *BubR1* trans-heterozygous mutant background.

*n*=27 (*w$^{1118}$*), 22 (*BubR1$^{k06109}$/+*), 18 (*BubR1$^{k03113}$/+*), 23 (*EGFP::BubR1$^{WT}$*)

midguts.

(D) Total cell number in control, *BubR1* heterozygous mutants, and *EGFP::BubR1$^{WT}$* rescue flies under 3wD. *n*=10 (*w$^{1118}$*), 10 (*BubR1$^{k06109}$*/+), 15 (*BubR1$^{k03113}$*/+), 23 (*EGFP::BubR1$^{WT}$*) midguts.

(E) PH3+ cell number in control, *BubR1* heterozygous mutants, and *EGFP::BubR1$^{WT}$* rescue flies under 3wD. *n*=16 (*w$^{1118}$*), 10 (*BubR1$^{k06109}$*/+), 15 (*BubR1$^{k03113}$*/+), 22 (*EGFP::BubR1$^{WT}$*) midguts.

(F) Quantification of *esg*+ cell number in control, *BubR1* heterozygous mutants under 3wD. *n*=23 (*w$^{1118}$*), 11 (*BubR1$^{k06109}$*/+), 16 (*BubR1$^{k03113}$*/+) midguts.

(G) Schematic of CRISPR-mediated *BubR1* knockout in *esg*+ cells.

(H) Quantification of midgut length in control and *esg*+-specific *BubR1* knockout flies. *n*=10 (*esg$^{Cas9}$*>+),10 (*esg$^{Cas9}$*>*BubR1$^{CRISPR}$*) midguts.

(I) Total cell number in control and *esg*+-specific *BubR1* knockout flies. *n*=9 (*esg$^{Cas9}$*>+),9 (*esg$^{Cas9}$*>*BubR1$^{CRISPR}$*) midguts.

(J) PH3+ cell number in control and *esg*+-specific *BubR1* knockout flies. *n*=10 (*esg$^{Cas9}$*>+),10 (*esg$^{Cas9}$*>*BubR1$^{CRISPR}$*) midguts.

**Figure 5. The MCC component Mad2 is involved in DDMR**

(A) Expression level of *mad2* in 3wC and 3wD midguts from RNA-seq data. Count data was normalized with RLE.

(B) Representative images of *GFP-Mad2* midguts under 3wC and 3wD conditions.

(C) Quantification of GFP-Mad2 signal intensity of (B). 40 cells from 4 midguts (3wC), 52 cells from 4 midguts (3wD).

(D) Representative images of the midgut in control and *mad2* heterozygous mutant flies under 3wD. Scale bars: 1 mm.

(E) Quantification of midgut length in control and *mad2* heterozygous mutant flies under 3wD. *n*=12 (*w$^{1118}$*), 9 (*mad2$^{\Delta}$*/+), 7 (*mad2$^{EY}$*/+) midguts. The same *w$^{1118}$* control samples were used in Figure 5E and Figure 4C.

(F) Quantification of *esg*+ cell number in control and *mad2* heterozygous mutant flies under 3wD. *n*=23 (*w$^{1118}$*), 10 (*mad2$^{\Delta}$*/+), 6 (*mad2$^{EY}$*/+) midguts. The *w$^{1118}$* control are same sample as in Figure 4E.

(G) Starvation resistance in control and *BubR1* heterozygous mutant flies. *n*=128 (*w$^{1118}$*), 114 (*BubR1$^{k06109}$*/+), 103 (*BubR1$^{k03113}$*/+) flies.

(H) Graphical summary of midgut remodeling in diapause and recovery

conditions.

**Figure S1. Midgut remodeling in coordination with *Drosophila* reproductive**

**diapause**

(A) Representative image of ovary growth arrest in *D. melanogaster* diapause.

Scale bar: 1 mm.

(B) Schematic of control and recovery in *D. melanogaster*. LD = Light : Dark.

(C) Quantification of midgut length of wild-type flies (Canton-S) under control and

recovery conditions. *n*=6-13 midguts for each condition. The control samples are

same as in Figure 1C.

(D) Quantification of midgut length in the recovery condition for 1-7 days. *n*=8-21

midguts for each condition.

(E) Schematic of the diapause and cold conditions in *D. melanogaster*. LD =

Light : Dark. To set up the cold condition, we allowed the flies to grow in 18°C for

4 days after eclosion, and then held them at 10 °C with 8 L:16D photoperiods for

3 weeks.

(F) Quantification of midgut length under diapause and cold conditions after 3

weeks. *n*=9-21 midguts for each condition. The 3 weeks diapause samples are

the same shown in Figure 1C.

(G) Quantification of midgut length and area in wild-type male flies (Canton-S)

under 3wC and 3wD condition. *n*=7 (3wC), 7 (3wD) midguts.

(H) Representative images of ovaries in *D. bifasciata* (JZK3) and *D. triauraria*

(TM15-92) under long day (LD) and short day (SD) conditions. In SD, ovarian

growth arrest was induced. Scale bars: 1 mm.

(I, J) Quantification of midgut length of non-diapause strains of *D. triauraria* (KMJ1

and OEB12) under LD and SD conditions. *n*=11 (KMJ1, LD), 11 (KMJ1, SD), 8

(OEB12, LD), 13 (OEB12, SD) midguts. KMJ1 and OEB12 female flies do not

enter diapause state in the SD condition (Yamada & Yamamoto, 2011).

**Figure S2. ISCs remain in a non-proliferative state during diapause**

(A) The proportion of esg+ cells in total cells. *n*=11 (3wC), 12 (3wD), 10 (3wD+R)

midguts.

(B) Representative images of the posterior midgut under 3wC and 3wD

conditions in *Dl-lacZ* flies. Scale bars: 100 µm.

(C) Representative images of anti-Dl staining in 3wC and 3wD wild-type flies.

Scale bars: 100 µm.

(D) Representative images of the posterior midgut under 3wC and 3wD

conditions in *Su(H)GBE-lacZ* flies. Scale bars: 100 µm.

(E) Quantification of the Su(H)+ cell number. *n*=8 (3wC), 6 (3wD) midguts.

(F) The proportion of Su(H)+ cells in total cells. *n*=8 (3wC), 6 (3wD) midguts.

(G) PH3+ cell number in the midgut of 3wC, 3wD, and 3wD+R flies. n=6 (3wC),

8 (3wD), 6 (3wD+R) midguts. The 3wC and 3wD samples are the same shown in

Figure 2F.

(H) Representative images of the posterior midgut under 3wC and 3wD

conditions in *CycB-GFP* flies. Scale bars: 100 µm.

(I) Representative images of anti-cDcp1 staining in 3wC and 3wD wild-type flies.

Scale bars: 50 µm.

**Figure S3. Molecular signatures identified by RNA-seq analysis in DDMR**

(A) Gene ontology (GO) enrichment of 3wD transcriptome compared to 3wC. All

differentially expressed genes in 3wD are used for enrichment analysis.

(B) Reactome Pathway Enrichment in 3wD compared to 3wC. All differentially

expressed genes in 3wD are used for enrichment analysis.

(C) Expression levels of *bmm, pepck, tobi* from RNA-seq data. *n*=4 (3wC), 4

(3wD) RNA samples. Count data was normalized with RLE.

(D) Expression levels of *per, tim* from RNA-seq data. *n*=4 (3wC), 4 (3wD) RNA

samples. Count data was normalized with RLE.

(E) Expression levels of *Drs* from RNA-seq data. *n*=4 (3wC), 4 (3wD) RNA

samples. Count data was normalized with RLE.

(F) Representative images of GFP-BubR1 from 3wC and 3wD posterior midguts.

Scale bar: 100 µm.

(G) MA plot of differentially expressed genes in a pair-wise comparison of 3wC+R

and 3wD+R conditions.

(H) GO enrichment in 3wD+R compared to 3wC+R. All differentially expressed

genes in 3wD+R are used for enrichment analysis.

(I) Reactome Pathway Enrichment in 3wD+R compared to 3wC+R. All

differentially expressed genes in 3wD+R are used for enrichment analysis.

(J) Expression levels of *CCHa2, Dh31, dilp3, NPF, Thor* from RNA-seq data. *n*=4

(3wC+R), 4 (3wD+R) RNA samples. Count data was normalized with RLE.

**Figure S4. BubR1 functions in the midgut in coordination with *Drosophila***

**reproductive diapause**

(A) Quantification of midgut length in control and *BubR1* heterozygous mutants

under 3wC. *n*=10 (*w^{1118}*), 8 (*BubR1^{k06109}/*+), 8 (*BubR1^{k03113}/*+) midguts.

(B) PH3+ cell number in control, *BubR1* heterozygous mutants, and

*EGFP::BubR1^{WT}* flies under 3wC. *n*=10 (*w^{1118}*), 8 (*BubR1^{k06109}/*+), 8

(*BubR1^{k03113}/*+) midguts.

(C) RT-qPCR of *BubR1* in control and CRISPR-mediated *BubR1* knockout flies

under the 3wD condition. *n*=9 (*esg^{Cas9}>*+), 9 (*esg^{Cas9}>BubR1^{CRISPR}*) RNA

samples.

(D) RT-qPCR of *BubR1* in *D. bifasciata* (JZK3) midgut under the control (LD) and

diapause (SD) conditions. *n*=8 (LD), 9 (SD) RNA samples.

(E) RT-qPCR of *BubR1* in *D. triauraria* (TM15-92) midgut under the control (LD)

and diapause (SD) conditions. *n*=6 (LD), 8 (SD) RNA samples.

**References**

Anduaga, A. M., Nagy, D., Costa, R., & Kyriacou, C. P. (2018). Diapause in

Drosophila melanogaster – Photoperiodicity, cold tolerance and

metabolites. *Journal of Insect Physiology*, *105*, 46–53.

https://doi.org/10.1016/j.jinsphys.2018.01.003

Baker, D. J., Dawlaty, M. M., Wijshake, T., Jeganathan, K. B., Malureanu, L., Van

Ree, J. H., Crespo-Diaz, R., Reyes, S., Seaburg, L., Shapiro, V., Behfar, A.,

Terzic, A., Van De Sluis, B., & Van Deursen, J. M. (2013). Increased

expression of BubR1 protects against aneuploidy and cancer and extends

healthy     lifespan.     *Nature     Cell     Biology*,     *15*(1),     96–102.

https://doi.org/10.1038/ncb2643

Baker, D. J., Jeganathan, K. B., Cameron, J. D., Thompson, M., Juneja, S.,

Kopecka, A., Kumar, R., Jenkins, R. B., De Groen, P. C., Roche, P., & Van
Deursen, J. M. (2004). BubR1 insufficiency causes early onset of aging-
associated phenotypes and infertility in mice. *Nature Genetics*, *36*(7), 744–
749. https://doi.org/10.1038/ng1382

Bellen, H. J., Levis, R. W., He, Y., Carlson, J. W., Evans-Holm, M., Bae, E., Kim,
J., Metaxakis, A., Savakis, C., Schulze, K. L., Hoskins, R. A., & Spradling, A.
C. (2011). The Drosophila gene disruption project: Progress using
transposons with distinctive site specificities. *Genetics*, *188*(3), 731–743.
https://doi.org/10.1534/genetics.111.126995

Bier, E., Harrison, M. M., O'connor-Giles, K. M., & Wildonger, J. (2018). Advances
in engineering the fly genome with the CRISPR-Cas system. *Genetics*,
*208*(1), 1–18. https://doi.org/10.1534/genetics.117.1113

Buchon, N., Osman, D., David, F. P. A., Yu Fang, H., Boquete, J. P., Deplancke,
B., & Lemaitre, B. (2013). Morphological and Molecular Characterization of
Adult Midgut Compartmentalization in Drosophila. *Cell Reports*, *3*(5), 1725–
1738. https://doi.org/10.1016/j.celrep.2013.04.001

Buffin, E., Emre, D., & Karess, R. E. (2007). Flies without a spindle checkpoint.
*Nature Cell Biology*, *9*(5), 565–572. https://doi.org/10.1038/ncb1570

Buffin, E., Lefebvre, C., Huang, J., Gagou, M. E., & Karess, R. E. (2005).
Recruitment of Mad2 to the Kinetochore Requires the Rod/Zw10 Complex.
*Current Biology*, *15*(9), 856–861. https://doi.org/10.1016/j.cub.2005.03.052

Choi, E., Zhang, X., Xing, C., & Yu, H. (2016). Mitotic Checkpoint Regulators
Control Insulin Signaling and Metabolic Homeostasis. *Cell*, *166*(3), 567–581.
https://doi.org/10.1016/j.cell.2016.05.074

Christensen, C. F., Laurichesse, Q., Loudhaief, R., Colombani, J., & Andersen,
D. S. (2024). Drosophila activins adapt gut size to food intake and promote
regenerative growth. *Nature Communications*, *15*(1), 273.
https://doi.org/10.1038/s41467-023-44553-9

Denlinger, D. L. (2022). *Insect Diapause* (Vol. 464). Cambridge University Press.
https://doi.org/10.1017/9781108609364

DiVito Evans, A., Fairbanks, R. A., Schmidt, P., & Levine, M. T. (2023). Histone
methylation regulates reproductive diapause in Drosophila melanogaster.
*PLOS Genetics*, *19*(9), e1010906.

https://doi.org/10.1371/journal.pgen.1010906

Easwaran, S., & Montell, D. J. (2023). The molecular mechanisms of diapause
and diapause-like reversible arrest. In *Biochemical Society Transactions* (Vol.
51,    Issue    5,    pp.    1847–1856).    Portland    Press    Ltd.
https://doi.org/10.1042/BST20221431

Easwaran, S., & Montell, D. J. (2024). *A genome-wide association study*
*implicates the olfactory system in Drosophila melanogaster diapause-*
*associated    lifespan    extension    and    fecundity.*
https://doi.org/10.7554/eLife.98142.2

Easwaran, S., Van Ligten, M., Kui, M., & Montell, D. J. (2022). Enhanced germline
stem cell longevity in Drosophila diapause. *Nature Communications*, *13*(1).
https://doi.org/10.1038/s41467-022-28347-z

Fang, Y., Liu, T., Wang, X., Yang, Y. M., Deng, H., Kunicki, J., Traganos, F.,
Darzynkiewicz, Z., Lu, L., & Dai, W. (2006). BubR1 is involved in regulation
of    DNA    damage    responses.    *Oncogene*,    *25*(25),    3598–3605.
https://doi.org/10.1038/sj.onc.1209392

Goto, S., & Hayashi, S. (1999). Proximal to distal cell communication in the
*Drosophila* leg provides a basis for an intercalary mechanism of limb
patterning.    *Development*,    *126*(15),    3407–3413.
https://doi.org/10.1242/dev.126.15.3407

Gray, D. R., Logan, J. A., Ravlin, F. W., & Carlson, J. A. (1991). Toward a Model
of Gypsy Moth Egg Phenology: Using Respiration Rates of Individual Eggs
to    Determine    Temperature–Time    Requirements    of    Prediapause
Development. *Environmental    Entomology*,    *20*(6),    1645–1652.
https://doi.org/10.1093/ee/20.6.1645

Guo, Z., Driver, I., & Ohlstein, B. (2013). Injury-induced BMP signaling negatively
regulates *Drosophila* midgut homeostasis. *Journal of Cell Biology*, *201*(6),
945–961. https://doi.org/10.1083/jcb.201302049

Guo, Z., & Ohlstein, B. (2015). Bidirectional Notch signaling regulates Drosophila
intestinal    stem    cell    multipotency.    *Science*,    *350*(6263),    927.
https://doi.org/10.1126/science.aab0988

Hara, Y., & Yamamoto, D. (2022). Effects of Food and Temperature on Drosophila
melanogaster Reproductive Dormancy as Revealed by Quantification of a

GFP-Tagged Yolk Protein in the Ovary. *Frontiers in Physiology, 12.*
https://doi.org/10.3389/fphys.2021.803144

Harsimran Kaur Gill, Gaurav Goyal, & Gurminder Chahil. (2017). Insect
Diapause: A Review. *Journal of Agricultural Science and Technology A*, 7(7).
https://doi.org/10.17265/2161-6256/2017.07.002

Helfrich-Förster, C. (2024). Neuropeptidergic regulation of insect diapause by the
circadian clock. In *Current Opinion in Insect Science* (Vol. 63). Elsevier Inc.
https://doi.org/10.1016/j.cois.2024.101198

Huang, J.-Y., & Raff, J. W. (1999). The disappearance of cyclin B at the end of
mitosis is regulated spatially in Drosophila cells. In *The EMBO Journal* (Vol.
18, Issue 8). https://www.embopress.org

Hung, R.-J., Hu, Y., Kirchner, R., Liu, Y., Xu, C., Comjean, A., Tattikota, S. G., Li,
F., Song, W., Ho Sui, S., & Perrimon, N. (2020). A cell atlas of the adult
*Drosophila* midgut. *Proceedings of the National Academy of Sciences*,
*117*(3), 1514–1523. https://doi.org/10.1073/pnas.1916820117

Karess, R. E., Wassmann, K., & Rahmani, Z. (2013). New insights into the role
of BubR1 in mitosis and beyond. In *International Review of Cell and*
*Molecular Biology* (Vol. 306, pp. 223–273). Elsevier Inc.
https://doi.org/10.1016/B978-0-12-407694-5.00006-7

Karp, X. (2021). Hormonal Regulation of Diapause and Development in
Nematodes, Insects, and Fishes. In *Frontiers in Ecology and Evolution* (Vol.
9). Frontiers Media S.A. https://doi.org/10.3389/fevo.2021.735924

Kim, M., & Denlinger, D. L. (2009). Decrease in expression of beta-tubulin and
microtubule abundance in flight muscles during diapause in adults of Culex
pipiens. *Insect Molecular Biology*, *18*(3), 295–302.
https://doi.org/10.1111/j.1365-2583.2009.00870.x

Kim, M., Robich, R. M., Rinehart, J. P., & Denlinger, D. L. (2006). Upregulation of
two actin genes and redistribution of actin during diapause and cold stress
in the northern house mosquito, Culex pipiens. *Journal of Insect Physiology*,
*52*(11–12), 1226–1233. https://doi.org/10.1016/j.jinsphys.2006.09.007

Kimura, M. T. (1990). Quantitative response to photoperiod during reproductive
diapause in the Drosophila auraria species-complex. *Journal of Insect*
*Physiology*, *36*(3), 147–152. https://doi.org/10.1016/0022-1910(90)90115-V

Kubrak, O. I., Kučerová, L., Theopold, U., & Nässel, D. R. (2014). The sleeping beauty: How reproductive diapause affects hormone signaling, metabolism, immune response and somatic maintenance in Drosophila melanogaster. *PLoS ONE*, *9*(11). https://doi.org/10.1371/journal.pone.0113051

Kubrak, O. I., Kucerová, L., Theopold, U., Nylin, S., & Nässel, D. R. (2016). Characterization of reproductive dormancy in male Drosophila melanogaster. *Frontiers in Physiology*, *7*(NOV). https://doi.org/10.3389/fphys.2016.00572

Kučerová, L., Kubrak, O. I., Bengtsson, J. M., Strnad, H., Nylin, S., Theopold, U., & Nässel, D. R. (2016). Slowed aging during reproductive dormancy is reflected in genome-wide transcriptome changes in Drosophila melanogaster. *BMC Genomics*, *17*(1). https://doi.org/10.1186/s12864-016-2383-1

Kurogi, Y., Imura, E., Mizuno, Y., Hoshino, R., Nouzova, M., Matsuyama, S., Mizoguchi, A., Kondo, S., Tanimoto, H., Noriega, F. G., & Niwa, R. (2023). Female reproductive dormancy in Drosophila is regulated by DH31-producing neurons projecting into the corpus allatum. *Development (Cambridge, England)*, *150*(10). https://doi.org/10.1242/dev.201186

Kurtz, C. C., Otis, J. P., Regan, M. D., & Carey, H. V. (2021). How the gut and liver hibernate. In *Comparative Biochemistry and Physiology -Part A: Molecular and Integrative Physiology* (Vol. 253). Elsevier Inc. https://doi.org/10.1016/j.cbpa.2020.110875

LAKOVAARA, S., SAURA, A., KOREF-SANTIBAÑEZ, S., & EHRMAN, L. E. E. (1972). Aspects of diapause and its genetics in northern drosophilids. *Hereditas*, *70*(1), 89–96. https://doi.org/https://doi.org/10.1111/j.1601-5223.1972.tb00996.x

Le Bras, S., & Van Doren, M. (2006). Development of the male germline stem cell niche in Drosophila. *Developmental Biology*, *294*(1), 92–103. https://doi.org/10.1016/j.ydbio.2006.02.030

Li, H., & Jasper, H. (2016). Gastrointestinal stem cells in health and disease: from flies to humans. *Disease Models & Mechanisms*, *9*(5), 487–499. https://doi.org/10.1242/dmm.024232

Lirakis, M., Dolezal, M., & Schlötterer, C. (2018). Redefining reproductive dormancy in Drosophila as a general stress response to cold temperatures.

*Journal of Insect Physiology*, *107*, 175–185. https://doi.org/10.1016/j.jinsphys.2018.04.006

Liu, M., Yang, S., Yang, J., Feng, P., Luo, F., Zhang, Q., Yang, L., & Jiang, H. (2024). BubR1 controls starvation-induced lipolysis via IMD signaling pathway in Drosophila. *Aging*, *16*(4), 3257–3279. https://doi.org/10.18632/aging.205533

Malita, A., Kubrak, O., Koyama, T., Ahrentløv, N., Texada, M. J., Nagy, S., Halberg, K. V., & Rewitz, K. (2022). A gut-derived hormone suppresses sugar appetite and regulates food choice in Drosophila. *Nature Metabolism*, *4*(11), 1532–1550. https://doi.org/10.1038/s42255-022-00672-z

Marianes, A., & Spradling, A. C. (2013). Physiological and stem cell compartmentalization within the Drosophila midgut. *ELife*, *2*. https://doi.org/10.7554/eLife.00886

Mattila, J., Viitanen, A., Fabris, G., Strutynska, T., Korzelius, J., & Hietakangas, V. (2024). Stem cell mTOR signaling directs region-specific cell fate decisions during intestinal nutrient adaptation. *Science Advances*, *10*(6). https://doi.org/10.1126/sciadv.adi2671

McGuire, S. E., Le, P. T., Osborn, A. J., Matsumoto, K., & Davis, R. L. (2003). Spatiotemporal Rescue of Memory Dysfunction in *Drosophila*. *Science*, *302*(5651), 1765–1768. https://doi.org/10.1126/science.1089035

Meiselman, M. R., Alpert, M. H., Cui, X., Shea, J., Gregg, I., Gallio, M., & Yapici, N. (2022). Recovery from cold-induced reproductive dormancy is regulated by temperature-dependent AstC signaling. *Current Biology*, *32*(6), 1362-1375.e8. https://doi.org/10.1016/j.cub.2022.01.061

Meller, C. L., Meller, R., Simon, R. P., Culpepper, K. M., Podrabsky, J. E., Meller, C., Lynn, ;, Meller, R. ;, Simon, R. P. ;, & Culpepper, K. M. ; (2012). Cell Cycle Arrest Associated with Anoxia-Induced Quiescence, Anoxic Preconditioning, and Embryonic Diapause in Embryos of the Annual Killifish Austrofundulus Limnaeus Citation Details. In *Biology Faculty Publications and Presentations Biology*. https://pdxscholar.library.pdx.edu/bio_fac/47

Meltzer, H., Marom, E., Alyagor, I., Mayseless, O., Berkun, V., Segal-Gilboa, N., Unger, T., Luginbuhl, D., & Schuldiner, O. (2019). Tissue-specific (ts)CRISPR as an efficient strategy for in vivo screening in Drosophila. *Nature*

*Communications*, *10*(1), 2113. https://doi.org/10.1038/s41467-019-10140-0

Meyerhof, G. T., Easwaran, S., Bontempo, A. E., Montell, C., & Montell, D. J.
(2024). Altered circadian rhythm, sleep, and *rhodopsin 7* –dependent shade
preference during diapause in *Drosophila melanogaster*. *Proceedings of the*
*National*     *Academy*     *of*     *Sciences*,     *121*(27).
https://doi.org/10.1073/pnas.2400964121

Miguel-Aliaga, I., Jasper, H., & Lemaitre, B. (2018). Anatomy and Physiology of
the Digestive Tract of *Drosophila melanogaster*. *Genetics*, *210*(2), 357–396.
https://doi.org/10.1534/genetics.118.300224

Musacchio, A. (2015). The Molecular Biology of Spindle Assembly Checkpoint
Signaling Dynamics. In *Current Biology* (Vol. 25, Issue 20, pp. R1002–
R1018). Cell Press. https://doi.org/10.1016/j.cub.2015.08.051

Nagai, H., Miura, M., & Nakajima, Y. ichiro. (2022). Cellular mechanisms
underlying adult tissue plasticity in Drosophila. In *Fly* (Vol. 16, Issue 1, pp.
190–206).     Taylor     and     Francis     Ltd.
https://doi.org/10.1080/19336934.2022.2066952

Nagai, H., Nagai, L. A. E., Tasaki, S., Nakato, R., Umetsu, D., Kuranaga, E., Miura,
M., & Nakajima, Y. (2023). Nutrient-driven dedifferentiation of
enteroendocrine cells promotes adaptive intestinal growth in Drosophila.
*Developmental*     *Cell*,     *58*(18),     1764-1781.e10.
https://doi.org/10.1016/j.devcel.2023.08.022

Nakagaki, M., Takei, R., Nagashima, E., & Yaginuma, T. (1991). Cell cycles in
embryos of the silkworm, Bombyx mori: G2-arrest at diapause .stage. In *Arch*
*Dev Biol* (Vol. 200).

O'Brien, L. E., Soliman, S. S., Li, X., & Bilder, D. (2011). Altered modes of stem
cell division drive adaptive intestinal growth. *Cell*, *147*(3), 603–614.
https://doi.org/10.1016/j.cell.2011.08.048

Ojima, N., Hara, Y., Ito, H., & Yamamoto, D. (2018). Genetic dissection of stress-
induced reproductive arrest in Drosophila melanogaster females. *PLoS*
*Genetics*, *14*(6). https://doi.org/10.1371/journal.pgen.1007434

Park, J., Hall, C., Hubbard, B., LaMoia, T., Gaspar, R., Nasiri, A., Li, F., Zhang,
H., Kim, J., Haeusler, R. A., Accili, D., Shulman, G. I., Yu, H., & Choi, E.
(2023). MAD2-Dependent Insulin Receptor Endocytosis Regulates

Metabolic Homeostasis. *Diabetes*, *72*(12), 1781–1794.
https://doi.org/10.2337/db23-0314

Patro, R., Duggal, G., Love, M. I., Irizarry, R. A., & Kingsford, C. (2017). Salmon
provides fast and bias-aware quantification of transcript expression. *Nature*
*Methods*, *14*(4), 417–419. https://doi.org/10.1038/nmeth.4197

Reiff, T., Jacobson, J., Cognigni, P., Antonello, Z., Ballesta, E., Tan, K. J., Yew, J.
Y., Dominguez, M., & Miguel-Aliaga, I. (2015). Endocrine remodelling of the
adult intestine sustains reproduction in Drosophila. *ELife*, *4*.
https://doi.org/10.7554/eLife.06930

Resende, L. P., Monteiro, A., Brás, R., Lopes, T., & Sunkel, C. E. (2018).
Aneuploidy in intestinal stem cells promotes gut dysplasia in Drosophila.
*Journal of Cell Biology*, *217*(11), 3930–3946.
https://doi.org/10.1083/jcb.201804205

Röttgen, G., Wagner, T., & Hinz, U. (1998). A genetic screen for elements of the
network that regulates neurogenesis in Drosophila. *Molecular and General*
*Genetics MGG*, *257*(4), 442–451. https://doi.org/10.1007/s004380050668

Saunders, D. S., Henricht, V. C., & Gilbertt, L. I. (1989). Induction of diapause in
Drosophila melanogaster: Photoperiodic regulation and the impact of
arrhythmic clock mutations on time measurement (ovarian diapause). In
*Genetics* (Vol. 86).

Saunders, D. S., Richard, D. S., Applebaum, S. W., Ma, M., & Gilbert~, L. I. (1990).
Photoperiodic Diapause in Drosophila melanogaster Involves a Block to the
Juvenile Hormone Regulation of Ovarian Maturation. In *GENERAL AND*
*COMPARATIVE ENDOCRINOLOGY* (Vol. 79).

Schiesari, L., Andreatta, G., Kyriacou, C. P., O'Connor, M. B., & Costa, R. (2016).
The insulin-like proteins dILPs-2/5 determine diapause inducibility in
Drosophila. *PLoS ONE*, *11*(9). https://doi.org/10.1371/journal.pone.0163680

Schiesari, L., & O'Connor, M. B. (2013). *Diapause* (pp. 213–246).
https://doi.org/10.1016/B978-0-12-396968-2.00008-7

Shinoda, N., Horikoshi, M., Taira, Y., Muramoto, M., Hirayama, S., Murata, S., &
Miura, M. (2023). Caspase cleaves Drosophila BubR1 to modulate spindle
assembly checkpoint function and lifespan of the organism. *FEBS Journal*,
*290*(17), 4200–4223. https://doi.org/10.1111/febs.16811

Soneson, C., Love, M. I., & Robinson, M. D. (2015). Differential analyses for RNA-
seq: transcript-level estimates improve gene-level inferences.
*F1000Research*, *4*, 1521. https://doi.org/10.12688/f1000research.7563.2

Song, Y.-H. (2005). Drosophila melanogaster: a model for the study of DNA
damage checkpoint response. *Molecules and Cells*, *19*(2), 167–179.

Staples, J. F. (2016). Metabolic Flexibility: Hibernation, Torpor, and Estivation. In
*Comprehensive Physiology* (pp. 737–771). Wiley.
https://doi.org/10.1002/cphy.c140064

Stojanović, O., Miguel-Aliaga, I., & Trajkovski, M. (2022). Intestinal plasticity and
metabolism as regulators of organismal energy homeostasis. *Nature
Metabolism*, *4*(11), 1444–1458. https://doi.org/10.1038/s42255-022-00679-6

Tammariello, S. P., & Denlinger, D. L. (1998). G0/G1 Cell Cycle Arrest in the Brain
of Sarcophaga crassipalpis During Pupal Diapause and the Expression
Pattern of the Cell Cycle Regulator, Proliferating Cell Nuclear Antigen. In
*Insect Biochem. Molec. Biol* (Vol. 28, Issue 2).

Tang, R., Jiang, Z., Chen, F., Yu, W., Fan, K., Tan, J., Zhang, Z., Liu, X., Li, P., &
Yuan, K. (2020). The Kinase Activity of Drosophila BubR1 Is Required for
Insulin Signaling-Dependent Stem Cell Maintenance. *Cell Reports*, *31*(12),
107794. https://doi.org/10.1016/j.celrep.2020.107794

Tatar, M., & Yin, C.-M. (2001). Slow aging during insect reproductive diapause:
why butterflies, grasshoppers and flies are like worms. *Experimental
Gerontology*, *36*(4–6), 723–738. https://doi.org/10.1016/S0531-
5565(00)00238-2

Trivedi, D., Vinitha, C. M., Bisht, K., Janardan, V., Pandit, A., Basak, B., Shwetha,
H., Ramesh, N., & Raghu, P. (2020). A genome engineering resource to
uncover principles of cellular organization and tissue architecture by lipid
signaling. *ELife*, *9*, 1–23. https://doi.org/10.7554/ELIFE.55793

Watabe, H. (1979). *Drosophila Survey of Hokkaido, XXXVI. Seasonal Changes
in the Reproductive Condition of Wild and Domestic Species of Drosophila*.

White, M. A., Bonfini, A., Wolfner, M. F., & Buchon, N. (2021). *Drosophila
melanogaster* sex peptide regulates mated female midgut morphology and
physiology. *Proceedings of the National Academy of Sciences*, *118*(1).
https://doi.org/10.1073/pnas.2018112118

Wilsterman, K., Ballinger, M. A., & Williams, C. M. (2021). A unifying, eco-
physiological framework for animal dormancy. *Functional Ecology*, *35*(1),
11–31. https://doi.org/10.1111/1365-2435.13718

Yamada, H., & Yamamoto, M. T. (2011). Association between circadian clock
genes and diapause incidence in Drosophila triauraria. *PLoS ONE*, *6*(12).
https://doi.org/10.1371/journal.pone.0027493

Yoshinari, Y., Nishimura, T., Yoshii, T., Kondo, S., Tanimoto, H., Kobayashi, T.,
Matsuyama, M., & Niwa, R. (2024). A high-protein diet-responsive gut
hormone regulates behavioral and metabolic optimization in Drosophila
melanogaster. *Nature Communications*, *15*(1), 10819.
https://doi.org/10.1038/s41467-024-55050-y

Zhou, X., Ding, G., Li, J., Xiang, X., Rushworth, E., & Song, W. (2020).
Physiological and Pathological Regulation of Peripheral Metabolism by Gut-
Peptide Hormones in Drosophila. In *Frontiers in Physiology* (Vol. 11).
Frontiers Media S.A. https://doi.org/10.3389/fphys.2020.577717

Zielke, N., Korzelius, J., vanStraaten, M., Bender, K., Schuhknecht, G. F. P., Dutta,
D., Xiang, J., & Edgar, B. A. (2014). Fly-FUCCI: A Versatile Tool for Studying
Cell Proliferation in Complex Tissues. *Cell Reports*, *7*(2), 588–598.
https://doi.org/10.1016/j.celrep.2014.03.020

1185

# Figure 1

# Figure 2

**A**

Lumen

Epithelia

Visceral muscle

**B** Midgut epithelial cells

ISC → EEP → EE

ISC → EB → EC

**C**

Total cell number

condition
- 3wC
- 3wD
- 3wD+R

Anterior    Posterior

**D** *esg-GFP*

3wC

3wD

esg    Hoechst esg

**E**

esg+ number

condition
- 3wC
- 3wD
- 3wD+R

NS.  NS.

Anterior    Posterior

**F** *CantonS*

PH3    PH3 Hoechst

3wC

3wD

**G**

PH3+ cell number

condition
- 3wC
- 3wD

Anterior    Posterior

**H** MARCM clone induction scheme

pupation    48h APF    96h APF    1w old

Heat shock    eclosion    Dissect

Labeled ISC    Clonal expansion

**I**

1wC

1wD

MARCM Clone    Hoechst Clone Pros

**J**

Cell number/Clone

1wC    1wD

**K**

3wC

3wD

E2F.GFP CycB.RFP

**L**

Proportion

Cell cycle
- G0/G1
- S
- G2/M

3wC    3wD

# Figure 3

# Figure 4

# Figure 5