1 **Centriole Loss in Embryonic Development Disrupts Axonal Pathfinding and**

2 **Muscle Integrity**

Beatriz González[1,2], Júlia Sellés-Altés[1], Jèrica Pla-Parron[1], Judith Castro-Ribera[1], and Sofia

J. Araújo*[1,2]

[1]Department of Genetics, Microbiology and Statistics, Facultat de Biologia, Universitat de Barcelona

(UB), 08028 Barcelona, Spain

[2]Institut de Biomedicina Universitat de Barcelona (IBUB), Barcelona, Spain

*Corresponding author. email: S. J. Araújo sofiajaraujo@ub.edu,

Keywords: centriole; centrosome; axon; nervous system; muscle; Drosophila; Sas-4; Sas-6;

wiggliness, tortuosity

Running title: centriole loss disrupts axon and muscle development

Abstract

Centrosomes, the primary microtubule-organizing centers (MTOCs), are crucial for early

neuronal development, particularly in establishing polarity and promoting axon formation.

Traditionally considered non-essential in terminally differentiated cells, recent evidence

suggests that centrosomes play vital roles in specialized cellular contexts. In mammalian

neurons, centrosome-mediated microtubule remodeling is essential for axon elongation, with

centrosome dysfunction leading to axonal misrouting and growth defects. Although active

centrosomes have been detected in the tracheal terminal cells of *Drosophila melanogaster*,

their activity in neurons *in vivo* has not been observed. Their gradual loss during

embryogenesis has been reported as non-essential for organogenesis, as adult flies can

eclose without centrioles.

To investigate centrosome activity in neurons, we revisited *Drosophila Sas-4* mutants, which

exhibit centrosome loss (CL), and observed that 50% of homozygous mutant individuals fail

to hatch as larvae. We analysed their development, focusing on the expressivity and

penetrance of nervous system phenotypes, and examined centriole localization in neurons.

Our findings confirm the presence of centrosomes in motor and sensory neurons in *Drosophila*

and their localization near the nascent axon. CL conditions resulted in axonal misrouting and

muscle developmental abnormalities. Targeted downregulation of Sas-4 in the pioneer motor

neurons aCC and RP2 induced axon guidance errors, indicating an autonomous role for

centrosomes in axonal navigation. Colocalization of acetylated- and γ-tubulin with centrioles

in motor neurons further confirmed the presence of functional centrosomes in these cells.

Analysis of motor axons revealed that CL leads to axonal tortuosity, a characteristic associated

with neurodegeneration. This is the first direct association of CL with axonal morphological

phenotypes, highlighting the role of centrosomes in neuronal development and their broader

influence on nervous system structure and function.

Introduction

Centrosomes are small membrane-less organelles that serve as primary microtubule-

organizing centres (MTOCs). Composed of a mother-daughter centriole pair surrounded by

pericentriolar material (PCM), they anchor and nucleate microtubules (MTs) through γ-tubulin

ring complexes (γ-TuRCs) (Bettencourt-Dias & Glover, 2007). During mitosis, centrosomes

organize the spindle apparatus, while their role in interphase is more variable (Conduit *et al*,

2015). Abnormalities in centrosomes are linked to disorders like microcephaly and

neurodevelopmental disorders, highlighting the importance of centrosome function in early

neurodevelopment (Goundiam & Basto, 2020; Marthiens *et al*, 2013). However, how

centrosome numbers affect differentiated cells and their involvement in post-cell-division

organogenesis is not yet fully understood. Although traditionally studied in dividing cells, recent

insights suggest centrosomes also regulate cellular behaviours in differentiated cells,

contradicting the assumption that centrosome-based MTOCs are largely inactive in such

contexts.

In vertebrates, centrosomes are critical for embryonic development. For example, null

mutations in mouse *Cenpj* (also called *Sas4*) lead to centriole loss and embryonic arrest at

embryonic day 9, and *Sass6*-mutant mouse embryos arrest at mid-gestation (Bazzi &

Anderson, 2014; Grzonka & Bazzi, 2024). In contrast, *Drosophila melanogaster* embryos have

been reported to develop to adulthood using maternally supplied Sas-4 and Sas-6, despite

gradual centrosome loss (CL) (Basto *et al*, 2006; Fatalska *et al*, 2021; Gogendeau & Basto,

2010). However, these adults are sterile, uncoordinated, unable to form cilia, and die shortly

after eclosion (Basto *et al.*, 2006). Without maternal Sas-4, Sas-4 and Asterless (Asl)

Drosophila embryonic development arrests early (Stevens *et al*, 2007; Varmark *et al*, 2007),

as observed in *C. elegans Sas-4* mutants (Leidel & Gönczy, 2003). In *Drosophila* neural

progenitor and epithelial cells, centrosome absence impairs spindle assembly and orientation,

leading to defects like neural apoptosis and altered brain size, mimicking either human

microcephaly or tumourigenesis (Basto *et al.*, 2006; Castellanos *et al*, 2008; Poulton *et al*,

2014; Poulton *et al*, 2017).

Centrosomes are also important for differentiated cellular functions. In mouse fibroblasts,

myoblasts and endothelial cells, centrosome orientation is important for nuclear positioning

and cell migration(Chang *et al*, 2015; Gomes *et al*, 2005; Kushner *et al*, 2014). During kidney

development, CL leads to decreased branching morphogenesis and small dysplastic kidneys

at birth (Cheng *et al*, 2023). In *Drosophila* tracheal terminal cells (TCs), centrosome numbers

influence the formation of subcellular lumina during tubulogenesis (Ricolo *et al*, 2016).

Increasing centrosome numbers or depleting centrosomal proteins alters this process,

stressing their importance in cells with specialized architectures (Ricolo *et al*, 2021).

In neurons, centrosomes play critical roles as MTOCs during early development, particularly

in axon formation, a process reliant on microtubule remodelling to establish neuronal polarity

(Meka *et al*, 2022). In mammalian neurons, centrosomes nucleate MTs in the soma, which are

subsequently severed and transported into neurites via motor-driven sliding, enabling axon

elongation and polarity establishment (Ahmad & Baas, 1995; del Castillo *et al*, 2015; Stiess *et*

al, 2010; Yu *et al*, 1993). In hippocampal neurons *in vitro,* suppression of centrosome-

mediated functions prevents axonal growth and neurons with more than one centrosome

sprout more than one axon (Anda *et al*, 2005). In addition, aberrant centrosome localization,

leads to misplaced axonal outgrowth(Anda *et al*, 2010). In human iPSC-derived neurons,

depleting centrioles causes significant axonal defects, including mislocalization of axonal MT-

associated proteins and affected growth cone morphologies (Lindhout *et al*, 2021). However,

as neurons mature, γ-tubulin (γ-tub) remains essential, but centrosomes lose MTOC activity

and become dispensable in mammals and flies (Leask *et al*, 1997; Nguyen *et al*, 2011;

Sanchez-Huertas *et al*, 2016; Stiess *et al.*, 2010).

These findings collectively suggest that while centrosomes are essential in specific cells,

stages and contexts, many differentiated cells can adapt to their loss through alternative

MTOCs. However, in many cases, CL induces phenotypes which indicate its important function

in cytoskeletal remodelling and the acquisition of specialised cell shapes (Araújo, 2019).

To investigate the role of centrosomes in neuronal development, we revisited *Drosophila Sas-*

4 mutants with reduced centriole numbers and analysed their embryonic development. We

quantified the expressivity and penetrance of nervous system phenotypes and examined

centriole localization within neurons. Our findings indicate that loss of centrosomal protein

function results in variable expressivity and penetrance of embryonic phenotypes, which likely

accounts for inconsistencies in previously reported results. We observed that centrosomes

are present and active in *Drosophila* motor and peripheral nervous system (PNS) neurons.

Importantly, CL during embryogenesis leads to axonal misrouting and abnormalities in muscle

development, emphasising a critical role of centrosomes in proper nervous system and muscle

structure formation. Furthermore, we were able to specifically observe axonal phenotypes

associated with CL.

Results

Sas-4 mutant conditions induce axonal misrouting phenotypes

Spindle assembly abnormal 4 (Sas-4) protein is essential for centriole replication in *Drosophila* and early embryonic development (Basto *et al.*, 2006; Stevens *et al.*, 2007). However, in contrast to *Sas-4* maternal mutants, *Sas-4* zygotic mutant embryos initially progress through early development due to the maternal contribution but gradually lose centrioles during embryonic and larval stages. This progressive centriole loss makes these mutants an excellent model for studying the role of centrosomes in differentiated cells. By embryonic stages 15–16, centrioles have reported to be undetectable in 50–80% of cells (Basto *et al.*, 2006). By these stages, *Sas-4* mutant embryos exhibit defects in tracheal subcellular lumen formation, showing the importance of centrioles in some differentiated cells (Ricolo *et al.*, 2016). Nevertheless, *Sas-4* homozygous mutants have been reported to survive to adulthood, exhibiting early lethality shortly after eclosion(Basto *et al.*, 2006). This early adult lethality suggests a critical role for Sas-4 in maintaining cellular and developmental processes essential for organismal viability.

To investigate the role of centrosomes in embryonic development, we first analysed the viability of *Sas-4* loss-of-function (LOF) mutants during embryonic, larval and adult stages. As previously reported, we were able to raise *Sas-4* mutant adults; however, only a proportion of Sas-4 individuals were found to eclose. These mutants were uncoordinated and became trapped in the food upon eclosion. When isolated at the pupal stage and placed in Petri dishes, they were able to eclose and survive for longer, but with available food, they died within 24-36 hours, which is significantly less than the reported survival rates of unfed flies (2-3 days) (Kopeć, 1928; Pearl & Parker, 1921). These adult individuals exhibited severe motor impairments, barely moving and failing to extend their wings (Fig. 1A). To obtain survival rates, we quantified the percentage of *Sas-4* mutants that reached adulthood, compared to the total number expected, based on Mendelian proportions from a cross between heterozygous *Sas-*

4/TM3ftzlacZ individuals. Since the homozygous balancer chromosome embryos are early

embryonic lethal, we could detect a proportion of 33.3% mutant embryos reaching embryonic

stage 16. However, we only detected 16.7% of L1 larvae 36h after egg laying (AEL) (Fig. 1B,

n=407). This proportion was maintained to wandering L3, pupal and adult stages, suggesting

that once the barrier of embryonic stages is passed, centrosomes are no longer necessary for

viability, and individuals can reach adult stages as previously reported (Basto *et al.*, 2006).

However, our results show that only 50% of the homozygous embryos successfully reached

the larval and adult stages, indicating that CL induces embryonic lethality in the remaining

50% of cases.

We then asked if the *Sas-4* embryos exhibited mutant nervous system phenotypes. To do so,

we first analysed FasII positive neurons focusing on the trajectories of their motor axons,

particularly on the Intersegmental Nerve (ISN) (Fig. 1C). The penetrance of the ISN phenotype

was 52.8% (n=53) quantified as at least 2 defects per embryo, despite variability in the

expressivity of the phenotype (Fig. 1E and H and Sup. Fig. 1B). The aberrant pathways

included the following: synaptic phenotypes, which were evident when the nerve fibres

innervated the wrong target or failed to innervate (Fig. 1E and Sup. Fig. 1B, arrow); misrouting

phenotypes, which involved defects in nerves that do not follow the wild-type (wt) stereotypical

routes (Fig. 1E and Sup. Fig. 1B, asterisk); and crossing phenotypes, which included aberrant

crossing between segments (Fig. 1E arrowhead). The same type of anomalous phenotypes

was detected in *Sas-4/Df* mutant embryos (Fig. 1F and H), penetrance 56% (n=25), as well

as RNAi Sas-4 downregulation (Fig. 1G and H), penetrance 33.3% (n=15). We downregulated

Sas-4 using a *UAS-Sas-4RNAi* expressed in all post-mitotic neurons, by an elavGal4 driver

(Sup. Fig. 1 D and K) and specifically in a subset of neurons, such as the anterior corner cell

(aCC), posterior corner cell (pCC) and RP2 pioneers, using RN2-Gal4 (Fujioka *et al*, 2003).

Interestingly, when Sas-4 was specifically downregulated in neurons, we observed an

additional phenotype: fasciculation/defasciculation defects (Fig. 1G arrow) ISN misrouting and

defasciculation phenotypes (Fig. 1G and Sup. Fig. 1D, asterisk and arrowhead) as well as

shorter nerve extensions (Fig. 1 G, dot), also observed in some *Sas-4* and *Sas-4/Df* mutant

embryos (Fig. 1F' and Sup. Fig. 1 C). This suggested these motor axon phenotypes were

induced cell-autonomously. Furthermore, ubiquitous expression of *Sas-4* using a *pUbqSas-4*

insertion (Peel *et al*, 2007), could partially rescue the mutant phenotypes, particularly by

reducing the number of defects (Fig. 1 H and Sup. Fig. 1 E and K).

We followed by analysing the peripheral nervous system (PNS) phenotypes of *Sas-4* mutant

conditions specifically focusing on their axonal misrouting phenotypes. We could detect axonal

misrouting phenotypes in all three mutant conditions: *Sas-4* (penetrance 60% n=20), *Sas-4/Df*

(penetrance 55% n=20) and *UAS-Sas-4RNAi* (penetrance 60% n=20) (Fig. 1J-L and M)

expressed in PNS neurons using *P0163Gal4* (Hummel *et al*, 2000). We could partially rescue

these phenotypes by expressing *Sas-4* ubiquitously (Fig.1M and Sup. Fig. 1J and L). Taken

together, these results indicate that *Sas-4* conditions lead to mutant phenotypes in the nervous

system, characterized by incomplete penetrance and variable expressivity. These phenotypes

appear to be at least partially cell-autonomous, as Sas-4 downregulation in motor and sensory

neurons results in axonal misrouting errors.

***Sas-4* mutant conditions disrupt muscle development**

Since the phenotypes induced by RNAi expression exclusively in neurons did not fully replicate

the strength of the motor axon defects observed in *Sas-4* embryos, we next decided to

investigate the muscle phenotypes in *Sas-4* mutants. Muscles play a critical role in guiding

motor neurons by providing chemical and physical cues essential for accurate axonal

pathfinding and target selection (Jeong, 2021). These signals ensure proper neuromuscular

connections, essential for coordinated movement. In *Drosophila* mutants lacking muscular

structures, such as *twist* (*twi*) embryos, the absence of these cues disrupts motor axon

trajectories (Younossi-Hartenstein & Hartenstein, 1993).

In stage 16 *Drosophila* embryos, each abdominal hemisegment (A2–A7) displays a consistent

and repeating pattern of 30 somatic muscle fibres, which can be visualized using a myosin-

specific antibody (Fig. 2A and C). These 30 syncytial muscle fibres are innervated by two

motor nerves: the anterior ISN, which targets the dorsal muscles, and the posterior SN, which

innervates the ventral muscles (Fig. 2C, wt). We first analysed the muscle pattern of *Sas-4*

mutant embryos and could detect defects that ranged from misplaced to missing muscle fibres

(Fig. 2 B, B´). We then analysed these muscle fibre phenotypes in parallel to the axonal defects

and were able to correlate many of the axonal defects in *Sas-4* embryos to the location of the

muscle pattern disruptions (Fig. 2E" and Sup. Fig. 2B', asterisk). In fact, 100% of *Sas-4*

embryos with detectable muscle mutant phenotypes also exhibited ISN axonal trajectory

disruptions, while 11% of embryos displayed ISN axonal misrouting without any observable

muscle phenotypes (n = 27). Accordingly, Sas-4 downregulation in muscle cells, using a *Mef2*

driver, induced muscle mutant phenotypes in 80% of embryos, with ISN defects spatially

correlating with muscle defects in 63% of the cases (n = 19, Fig. 2G" asterisk). Sas-4

downregulation in muscle cells, using a *Mef2* driver together with a *Twist* (*Twi*) driver, induced

stronger, more penetrant ISN phenotypes that were dose dependent (Sup. Fig. 2E, F)

Consistent with these findings, downregulation of Sas-4 in pioneer neurons using the RN2

driver resulted in axonal misrouting phenotypes in 33% of the embryos (n = 15) and very mild

muscle mutant phenotypes in only 20% of the embryos (n = 10). Collectively, these findings

suggest that axonal misrouting phenotypes arise from two distinct contributions: an

autonomous component originating from their own CL and a non-autonomous one driven by

CL-affected muscle cells (Fig. 2C, *Sas-4*).

Sas-6 mutants have axonal guidance and muscle developmental defects

Centriole assembly depends on a small set of conserved proteins, the absence of any of which

results in centriole loss; *Sas-4* and *Sas-6* are key components of this process (Gartenmann *et*

al, 2020). To rule out the possibility that the observed axonal and muscle defects were

specifically due to mutations in *Sas-4*, we began by analysing the viability of *Sas-6* LOF

mutants during the embryonic, larval, and adult stages. We used a transallelic combination of

two CRISPR-Cas9 generated *Sas-6* induced deletions mutants, *Sas-6$^{\Delta a}$/Sas-6$^{\Delta b}$* (Z. Novak

and J. Raff, unpublished). We successfully raised *Sas-6* adults (Fig. 3A), which, like *Sas-4*

adults, died within 24-36 hours. To corroborate that this combination of *Sas-6* alleles induced

CL in embryos, we counted the number of centrioles in *Sas-6^{Δa}/Sas-6^{Δb}* embryos and observed

a reduction at stage 16 (Fig. 3B). Like *Sas-4* mutants, 33.3% of *Sas-6* mutant embryos

reached embryonic stage 16. However, compared to *Sas-4* mutants, in *Sas-6* we observed a

higher survival rate of 27.5% at the L1 larval stage (Fig. 3C, n=407). This proportion was

maintained through the wandering L3, pupal, and adult stages, suggesting that, like in *Sas-4*

mutants, centrosomes seem not to be essential for larval and pupal development once the

embryonic stages are completed.

We then investigated whether *Sas-6* embryos exhibited mutant axonal guidance and muscle

phenotypes. We then investigated whether *Sas-6* embryos exhibited mutant axonal guidance

and muscle phenotypes. We observed ISN axonal misrouting in *Sas-6* embryos (Fig. 3E, F),

with a penetrance of 47.6% (n=20), similar to that seen in *Sas-4* embryos. However, the

severity of the defects was milder than in *Sas-4*, as we did not observe any strong axonal

misrouting or crossing phenotypes. In most cases, the ISN axonal misrouting also correlated

with muscle phenotypes (Fig. 3E'' and F'', asterisk). Altogether, these findings support the role

of centrosomes in regulating axon guidance of motor neurons.

Pioneer neurons have centrioles localizing to the sites of axon outgrowth

In *Drosophila*, the aCC and RP2 motor neurons are pioneer neurons which influence growth

of the ISN (Sánchez-Soriano & Prokop, 2005) (Fig. 4 A). Pioneer neurons extend growth cones

that navigate a future fibre tract, and must interpret the full complexity of environmental cues

to define the path of the nerve (Harrison, 1910; Jacobs & Goodman, 1989). Therefore, we

investigated whether centrioles could be detected in the soma of these neurons in stage 12-

13 when they are starting to extend their axons. We found that centrioles were present in the

soma of both aCC (Fig. 4 B and C) and RP2 motor neurons (Fig. 4 D). Specifically, in the case

of the aCC, a centriole was observed in the quadrant where the axon exit point is first detected

in 72% of the cases (Fig. 4B n=75). Additionally, we did not detect centrioles in the quadrant

opposite the axon exit point (Fig. 2B n=75). In *Sas-4* mutants, centrioles were largely

undetected in both aCC/pCC neurons and RP2 neurons. (Fig. 4 E, F). At these stages, the

aCC and pCC neurons are very close together, making it difficult to differentiate between the

somas of the two neurons. Therefore, we quantified the number of centrioles in aCC+pCC and

RP2 neurons in both control and *Sas-4* mutant embryos. In control embryos, we observed

around 2.53 centrioles per pair of aCC and pCC neurons, while in *Sas-4* mutant, this number

was significantly reduced to 0.81 (Fig. 4G n=76). Similarly, for RP2 neurons, control embryos

exhibited 0.81 centriole per RP2 neuron, whereas *Sas-4* embryos showed a reduction to 0.26

(Fig. 4G n=76). This significant decrease in the humber of centrioles further supports the

critical role of Sas-4 in pioneer neurons of the ventral nerve cord (VNC) during *Drosophila*

embryogenesis.

Non-ciliated sensory neurons have centrioles

In *Drosophila* embryos, the dorsal cluster neurons project their axons in a highly organized

manner to form the sensory axonal tracts (Ghysen *et al*, 1986). During development, the axons

of these neurons extend dorsally and medially, connecting with their target regions and

fasciculating with the intersegmental nerve (ISN) to form a bundled tract for efficient sensory

signal routing (Hartenstein, 1988). Sensory neurons in *Drosophila* embryos are known to have

centrioles. These centrioles are crucial for forming basal bodies in ciliated cells, including type-

I sensory neurons, which are essential for the development of cilia (Jana *et al*, 2016). Since

we detected axonal guidance phenotypes in the axons extending from the dorsal cluster, we

tested for the presence of centrioles in dorsal cluster sensory neurons (Fig. 5A). We detected

centrioles in the soma of most of these neurons and quantified the number of centriole-positive

ddaC neurons (Fig. 5A'', B, and C', arrowhead), finding centrioles in 81.4% of the total (Fig.

5B, n=43). Importantly, we also detected centrioles in dorsal bipolar neuron (dbp, also known

as dbd), the proposed pioneer neuron of the dorsal cluster (Fig. 5A'' and C', arrow) (Martin *et*

al, 2008; Merritt & Whitington, 1995). In *Sas-4* embryos, the presence of centrioles in the

dorsal cluster neurons was significantly lower (Fig. 5D). In many embryos, we could not detect

any centrioles in the soma of the dorsal cluster neurons (Fig. 5E).

Based on our findings in the CNS and PNS, we decided to quantify the total number of

centrioles in *Sas-4* embryos and compare them to controls. To do so, we selected equal

segmental areas spanning the dorsal cluster and the lateral chordotonal organs (lch) and

counted the total number of centrioles in stage 16 control embryos, *Sas-4* homozygous

embryos displaying the mutant phenotype, *Sas-4* homozygous embryos with no mutant

phenotype and *Sas-4* mutants rescued by ubiquitous Sas-4 expression (Fig. 5F). We observed

that both mutants had approximately one-third of the centrioles found in the control group.

Embryos without defects had slightly higher centriole numbers compared to those with mutant

phenotypes. Additionally, ubiquitous expression of Sas-4 could increase centriole numbers.

These results confirm that *Sas-4* embryos gradually lose their centrioles. Significantly, we also

found a correlation between axonal guidance phenotypes and the numbers of centrioles per

embryo.

Pioneer motor neurons have active centrosomes

After revealing that ISN pioneers have centrioles, we were interested in knowing if these could

be active MTOCs. To determine if the centriole could organize a functional centrosome and

serve as a major site of neuronal microtubule nucleation in aCC and RP2 neurons, we

analysed the localization of acetylated-tubulin (ace-tub) and γ-tubulin (γ-tub). MTs are

nucleated and anchored by the microtubule nucleator γ-tub, which is embedded within the

pericentriolar material (PCM) of the centrosome (Lin *et al*, 2015). In both neurons we could

see stable acetylated MTs extending from the centrosomes (Fig. 6 A, B, arrowheads) and

these colocalized with γ-tub (Fig. 6C, D, arrowheads). We quantified the number of centrioles

colocalizing with γ-tub (Fig. 6E) and could detect co-localization in 50% of the cases, meaning

that there are centrioles in these neurons which are not active as MTOCs at these stages (Fig.

6C'' asterisk). In conclusion, these results suggest that at least some microtubule organization

in pioneer neurons of the VNC is dependent on the centrosome.

Centrosome loss induces motor axon fibre tortuosity

Axonal tortuosity (wiggliness) is a measure of the irregularities or abnormalities in the

morphological structure of axons which can be associated with neurodegeneration (Bando *et*

al, 2015). When axons are "wiggly," it suggests that they are not following their usual straight

or organized paths, possibly due to changes in axonal cytoskeletal structure or cell-adhesion,

but not much is known about what triggers axonal wiggliness. Qualitative examination of

confocal projections showed that aCC and RP2 axons are straighter in controls than in *Sas-4*

mutants, revealing the tortuosity of these axons (Fig. 7 A-E, asterisks). Apart from the

previously characterised phenotypes such as axonal crossings (Fig. 7 B, arrowhead) and

misrouting, we could observe axonal wiggliness at the CNS exit points (Fig. 7 B', D and E,

asterisks). Since we were interested in determining the proportion of wiggly axons in the

mutant, we quantified this for the aCC and RP2 neurons (Fig. 7 F). In *Sas-4* mutants we could

detect an average of 23% aCC and 30% RP2 wiggly axons per total number of aCC and RP2

axons, which was significantly different from the proportion of wiggly axons in the controls (6%

wiggly aCC and 8% wiggly RP2 axons). Therefore, CL induces axonal morphological changes

(Fig. 7G). Most importantly, we could observe specific axonal phenotypes in embryos with

reduced centriole numbers, confirming the importance of the centrosome for axonal extension.

Discussion

Here we provide evidence that the centrosome plays a role in axonal extension, navigation,

and muscle development *in vivo*. Previous studies have shown that *Sas-4* and *Sas-6* mutant

flies can reach adulthood (Basto *et al.*, 2006; Fatalska *et al.*, 2021; Peel *et al.*, 2007), which

initially seems incompatible with an important role for the centrosome during embryonic

development. However, we found that *Sas-4* homozygous mutations result in 50% lethality,

and a *Sas-6* transallelic combination causes 17% embryonic lethality.

Sas-4 and *Sas-6* LOF embryos display motor axon mutant phenotypes (Fig. 1 and 3),

revealing that centrosomes are important for axonal guidance. Additionally, *Sas-4* and *Sas-6*

LOF embryos exhibit muscle mutant phenotypes (Fig. 2 and 3), indicating that centrosomes

are also important for muscle development. Hence, the embryonic axon guidance phenotypes

have a dual contribution: an autonomous one from axonal CL and a non-autonomous one from

muscle CL. Since innervation does not influence the patterning, morphogenesis, maintenance,

or physiological development of somatic muscles in *Drosophila* embryos, the muscle defects

observed are most likely due to CL in muscle cells (Broadie & Bate, 1993). CL in muscle cells

may impair cilia formation and nuclear positioning, two processes important for muscle fibre

development (Loh *et al*, 2023; Metzger *et al*, 2012; Tillery *et al*, 2018). Expressing *Sas-4 RNAi*

under the control of the muscle-specific *Mef2-GAL4* driver induced penetrant muscle

phenotypes, consistent with previous results reporting the lethality of these embryos

(Schnorrer *et al*, 2010). *Sas-6* embryos exhibited milder muscle phenotypes than *Sas-4*,

aligning with previous findings that driving *Sas-6 RNAi* with a Mef2-GAL4 driver induced adult

pharate lethality (Schnorrer *et al.*, 2010).

What could explain the variability in the consequences of CL during embryonic development?

Many disease-causing mutations exhibit mild or no effects in some individuals, a phenomenon

known as incomplete phenotype penetrance. This phenomenon is still poorly understood, but

recent studies in *Drosophila* suggest it is stochastic, akin to flipping a coin (Marmion *et al*,

2023). Although the reported study focuses on fluctuations in signalling, a similar scenario

could be imagined for centrosomes. In cases of centrosome duplication impairment which

induce CL, distribution would vary among dividing cells as development proceeds. Early

embryonic mitotic cycles can partly explain the differences in phenotype expressivity. The first

13 mitotic cycles of *Drosophila* embryos are rapid, synchronous, and occur without cell

division. Subsequently, three rounds of patterned mitosis occur (mitoses 14, 15, and 16).

Following mitosis 16, many cells enter their first G1 phase (Edgar & O'Farrell, 1990). However,

the nervous system lineage does not enter a terminal interphase after mitosis 16 and

continues to divide according to an independent program (Bodmer & Jan, 1987; Hartenstein

et al, 1987). Similarly, muscle cells, continue dividing until embryonic stage 10 (Nabel-Rosen

et al, 2005). These extra cell divisions can account for the appearance, variability and severity

of the phenotypes observed, as centrosome loss is exacerbated with each cell division, which

depletes the maternal contribution. In agreement with this, when we quantified the number of

centrioles present in *Sas-4* embryos displaying axonal phenotypes, we could detect a

decrease of about 1/3 (Fig. 5), with consistent results when we quantified the numbers of

centrioles in *Sas-4* aCC, pCC and RP2 neurons (Fig. 4)

Centrosome localization in neurons and implications for axonal morphology

Here we show that centrioles are present in motor and sensory neurons and that in motor

neurons these centrioles colocalize with acetylated- and γ-tubulin suggesting they are active

MTOCs in these cells (Fig. 4, 5 and 6). Previous studies failed to detect colocalization of

centrosomes with γ-tub in neurons (Nguyen *et al.*, 2011). However, this study used Fizzy-

related-GFP (Fzr-GFP) to visualise centrioles and while it is clear Fzr is localizes to centrioles

during the cell cycle (Meghini *et al*, 2016), it is not clear it maintains this localization in all fully

differentiated cells (Djabrayan *et al*, 2014; Huynh *et al*, 2009). This, or differences in

microscopy techniques, could account for the inconsistency between our results and those of

others. Furthermore, centrosome localization and its ability to organise MTs is more likely to

be important at initial axon outgrowth stages, and not once the axon has already grown.

Regarding centrosome localization and activity in neurons, we show that axonal morphology

in embryos with fewer centrosomes is altered. This is the first example of a morphological

defect in axons associated with CL in *Drosophila*, and one we consider highly significant.

Under conditions of CL, we observe tortuous (wiggly) axons, suggesting that while CL does

not prevent neurons from extending axons, it does affect axonal outgrowth. CL could induce

this phenotype by disrupting the neuronal cytoskeleton and preventing MTs to form, be

severed and transported into the axons. It is known that alterations in the neuronal cytoskeleton, caused by signalling deregulation such as increased JNK activity, can result in axonal wiggliness (Karkali *et al*, 2023). Also, alterations in MT dynamics in mammalian neurons have been shown to decrease axonal elongation, but not its formation, giving rise to wiggly axons (Rochlin *et al*, 1996). In addition, axonal tortuosity is one of the hallmarks of neurodegeneration (Salvadores *et al*, 2020), which makes this phenotype a good entrance point for the analysis of the progression of various neurodegenerative diseases.

The loss of centrosome function has been documented to disrupt various developmental stages in vertebrates. In humans, CL is associated with hereditary spastic paraplegia, Bardet-Biedl syndrome, the formation of cystic kidneys, disrupted left-right asymmetry, and microcephaly (Badano *et al*, 2005; Jaiswal & Singh, 2021). Demonstrating that CL in *Drosophila* affects the development of the nervous and muscular systems paves the way for using this model organism to further explore these human diseases.

Acknowledgments

We thank M. Llimargas and J. Casanova for generously allowing us to use their laboratory for specific parts of this work and for their support throughout this study. We thank B. Estrada, C. González, K. Karkali, C. Klämbt, E. Martin-Blanco, J. Raff, S. Rumpf, Y. Zheng, the Bloomington Drosophila Stock Center (BDSC) and the Vienna Drosophila Stock Center (VDRC) for fly stocks and reagents. We extend our gratitude to Z. Novak and J. Raff for sharing two unpublished CRISPR-Cas9 made *Sas-6* fly stocks with us. Thanks also go to R. Mendez for technical assistance in all the experiments and M. Bosch for assistance and advice with confocal microscopy and software. B.G. is the recipient of a Beatriu de Pinós post-doctoral fellowship from the AGAUR (2022BP00125). This work was supported by a grant from the Spanish *Ministerio de Ciencia y Innovación* (PID2021-125860NB-I00), the Universitat de Barcelona, and a grant from the Generalitat de Catalunya (2021 SGR 1455).

Author contributions

Conceptualization: S.J.A. Methodology and Experiments: B.G., J.P., J.S.-A., J.C.-R. and

S.J.A. Analysis: B.G., J.P., J.S.-A. and S.J.A. Writing: S.J.A. Reviewing and editing: S.J.A. and

B.G. Supervision and Funding: S.J.A.

References

Ahmad FJ, Baas PW (1995) Microtubules released from the neuronal centrosome are transported into the axon. *Journal of Cell Science* 108 (Pt 8): 2761-2769

Anda FCd, Meletis K, Ge X, Rei D, Tsai L-H (2010) Centrosome Motility Is Essential for Initial Axon Formation in the Neocortex. *The Journal of Neuroscience* 30: 10391-10406

Anda FCd, Pollarolo G, Silva JSD, Camoletto PG, Feiguin F, Dotti CG (2005) Centrosome localization determines neuronal polarity. *Nature Cell Biology* 436: 704 - 708

Araújo SJ (2019) The Golgi Apparatus and Centriole, Functions, Interactions and Role in Disease - Centrosomes in Branching Morphogenesis. *Results Problems Cell Differ* 67: 323-336

Badano JL, Teslovich TM, Katsanis N (2005) The centrosome in human genetic disease. *Nature Reviews Genetics* 6: 194-205

Bando Y, Nomura T, Bochimoto H, Murakami K, Tanaka T, Watanabe T, Yoshida S (2015) Abnormal morphology of myelin and axon pathology in murine models of multiple sclerosis. *Neurochemistry International* 81: 16-27

Basto R, Lau J, Vinogradova T, Gardiol A, Woods CG, Khodjakov A, Raff JW (2006) Flies without centrioles. *Cell* 125: 1375-1386

Bazzi H, Anderson KV (2014) Acentriolar mitosis activates a p53-dependent apoptosis pathway in the mouse embryo. *Proc Natl Acad Sci U S A* 111: E1491-1500

Bettencourt-Dias M, Glover DM (2007) Centrosome biogenesis and function: centrosomics brings new understanding. *Nat Rev Mol Cell Biol* 8: 451-463

Bodmer R, Jan YN (1987) Morphological differentiation of the embryonic peripheral neurons in Drosophila. *Roux's Arch Dev Biol*: 69-77

Broadie K, Bate M (1993) Muscle development is independent of innervation during Drosophila embryogenesis. *Development* 119: 533-543

Castellanos E, Dominguez P, Gonzalez C (2008) Centrosome Dysfunction in Drosophila Neural Stem Cells Causes Tumors that Are Not Due to Genome Instability. *Current Biology* 18: 1209-1214

Chang W, Antoku S, Östlund C, Worman HJ, Gundersen GG (2015) Linker of nucleoskeleton and cytoskeleton (LINC) complex-mediated actin-dependent nuclear positioning orients centrosomes in migrating myoblasts. *Nucleus* 6: 77-88

Cheng T, Agwu C, Shim K, Wang B, Jain S, Mahjoub MR (2023) Aberrant centrosome biogenesis disrupts nephron and collecting duct progenitor growth and fate resulting in fibrocystic kidney disease. *Development* 150

Conduit PT, Wainman A, Raff JW (2015) Centrosome function and assembly in animal cells. *Nature Reviews Molecular Cell Biology* 16: 611-624

del Castillo U, Lu W, Winding M, Lakonishok M, Gelfand VI (2015) Pavarotti/MKLP1 regulates microtubule sliding and neurite outgrowth in Drosophila neurons. *Curr Biol* 25: 200-205

Djabrayan Nareg JV, Cruz J, de Miguel C, Franch-Marro X, Casanova J (2014) Specification of Differentiated Adult Progenitors via Inhibition of Endocycle Entry in the Drosophila Trachea. *Cell Reports* 9: 859-865

Edgar BA, O'Farrell PH (1990) The three postblastoderm cell cycles of Drosophila
embryogenesis are regulated in G2 by string. *Cell* 62: 469-480
Fatalska A, Stepinac E, Richter M, Kovacs L, Pietras Z, Puchinger M, Dong G, Dadlez M,
Glover DM (2021) The dimeric Golgi protein Gorab binds to Sas6 as a monomer to mediate
centriole duplication. *Elife* 10
Fujioka M, Lear BC, Landgraf M, Yusibova GL, Zhou J, Riley KM, Patel NH, Jaynes JB
(2003) Even-skipped, acting as a repressor, regulates axonal projections in Drosophila.
Development 130: 5385-5400
Gartenmann L, Vicente CC, Wainman A, Novak ZA, Sieber B, Richens JH, Raff JW (2020)
Drosophila Sas-6, Ana2 and Sas-4 self-organise into macromolecular structures that can be
used to probe centriole and centrosome assembly. *Journal of Cell Science* 133: jcs244574
Ghysen A, Dambly-Chaudière C, Aceves E, Jan L, Jan Y (1986) Sensory neurons and
peripheral pathways in Drosophila embryos. *Rouxs Arch Dev Biol* 195: 281-289
Gogendeau D, Basto R (2010) Centrioles in flies: The exception to the rule? *Semin Cell Dev*
Biol 21: 163-173
Gomes ER, Jani S, Gundersen GG (2005) Nuclear Movement Regulated by Cdc42, MRCK,
Myosin, and Actin Flow Establishes MTOC Polarization in Migrating Cells. *Cell* 121: 451-463
Goundiam O, Basto R (2020) Centrosomes in disease: how the same music can sound so
different? *Current opinion in structural biology* 66: 74-82
Grzonka M, Bazzi H (2024) Mouse SAS-6 is required for centriole formation in embryos and
integrity in embryonic stem cells. *eLife* 13: e94694
Harrison RG (1910) The outgrowth of the nerve fiber as a mode of protoplasmic movement.
Journal of Experimental Zoology 9: 787-846
Hartenstein V (1988) Development of Drosophila larval sensory organs: spatiotemporal
pattern of sensory neurones, peripheral axonal pathways and sensilla differentiation.
Development 102: 869-886
Hartenstein V, Rudloff E, Campos-Ortega JA (1987) The pattern of proliferation of the
neuroblasts in the wild-type embryo of Drosophila melanogaster. *Rouxs Arch Dev Biol* 196:
473-485
Hummel T, Krukkert K, Roos J, Davis G, Klämbt C (2000) Drosophila Futsch/22C10 Is a
MAP1B-like Protein Required for Dendritic and Axonal Development. *Neuron* 26: 357-370
Huynh MA, Stegmüller J, Litterman N, Bonni A (2009) Regulation of Cdh1-APC function in
axon growth by Cdh1 phosphorylation. *J Neurosci* 29: 4322-4327
Jacobs J, Goodman C (1989) Embryonic development of axon pathways in the Drosophila
CNS. II. Behavior of pioneer growth cones. *The Journal of Neuroscience* 9: 2412-2422
Jaiswal S, Singh P (2021) Centrosome dysfunction in human diseases. *Semin Cell Dev Biol*
110: 113-122
Jana SC, Bettencourt-Dias M, Durand B, Megraw TL (2016) Drosophila melanogaster as a
model for basal body research. *Cilia*: 1-7
Jeong S (2021) Molecular Mechanisms Underlying Motor Axon Guidance in Drosophila.
Molecules and Cells 44: 549-556
Karkali K, Vernon SW, Baines RA, Panayotou G, Martín-Blanco E (2023) Puckered and JNK
signaling in pioneer neurons coordinates the motor activity of the Drosophila embryo. *Nature*
Communications 14: 8186
Kopeć S (1928) On the Influence of Intermittent Starvation on the Longevity of the Imaginal
Stage of Drosophila Melanogaster. *Journal of Experimental Biology* 5: 204-211
Kushner EJ, Ferro LS, Liu J-Y, Durrant JR, Rogers SL, Dudley AC, Bautch VL (2014) Excess
centrosomes disrupt endothelial cell migration via centrosome scattering. *J Cell Biol* 206:
257-272
Leask A, Obrietan K, Stearns T (1997) Synaptically coupled central nervous system neurons
lack centrosomal γ-tubulin. *Neuroscience Letters* 229: 17-20
Leidel S, Gönczy P (2003) SAS-4 is essential for centrosome duplication in C elegans and is
recruited to daughter centrioles once per cell cycle. *Dev Cell* 4: 431-439
Lin T-c, Neuner A, Schiebel E (2015) Targeting of γ-tubulin complexes to microtubule
organizing centers: conservation and divergence. *Trends in Cell Biology* 25: 296-307

Lindhout FW, Portegies S, Kooistra R, Herstel LJ, Stucchi R, Hummel JJA, Scheefhals N, Katrukha EA, Altelaar M, MacGillavry HD *et al* (2021) Centrosome-mediated microtubule remodeling during axon formation in human iPSC-derived neurons. *The EMBO Journal* 40: e106798

Loh M, Bernard F, Guichet A (2023) Kinesin-1 promotes centrosome clustering and nuclear migration in the Drosophila oocyte. *Development* 150

Marmion RA, Simpkins AG, Barrett LA, Denberg DW, Zusman S, Schottenfeld-Roames J, Schüpbach T, Shvartsman SY (2023) Stochastic phenotypes in RAS-dependent developmental diseases. *Current Biology* 33: 807-816.e804

Marthiens V, Rujano MA, Pennetier C, Tessier S, Paul-Gilloteaux P, Basto R (2013) Centrosome amplification causes microcephaly. *Nat Cell Biol* 15: 731-740

Martin V, Mrkusich E, Steinel MC, Rice J, Merritt DJ, Whitington PM (2008) The L1-type cell adhesion molecule Neuroglian is necessary for maintenance of sensory axon advance in the Drosophila embryo. *Neural Dev* 3: 10

Meghini F, Martins T, Tait X, Fujimitsu K, Yamano H, Glover DM, Kimata Y (2016) Targeting of Fzr/Cdh1 for timely activation of the APC/C at the centrosome during mitotic exit. *Nat Commun* 7: 12607

Meka DP, Kobler O, Hong S, Friedrich CM, Wuesthoff S, Henis M, Schwanke B, Krisp C, Schmuelling N, Rueter R *et al* (2022) Centrosome-dependent microtubule modifications set the conditions for axon formation. *Cell Reports* 39: 110686

Merritt DJ, Whitington PM (1995) Central projections of sensory neurons in the Drosophila embryo correlate with sensory modality, soma position, and proneural gene function. *J Neurosci* 15: 1755-1767

Metzger T, Gache V, Xu M, Cadot B, Folker ES, Richardson BE, Gomes ER, Baylies MK (2012) MAP and kinesin-dependent nuclear positioning is required for skeletal muscle function. *Nature* 484: 120-124

Nabel-Rosen H, Toledano-Katchalski H, Volohonsky G, Volk T (2005) Cell Divisions in the Drosophila Embryonic Mesoderm Are Repressed via Posttranscriptional Regulation of string/cdc25 by HOW. *Current Biology* 15: 295-302

Nguyen MM, Stone MC, Rolls MM (2011) Microtubules are organized independently of the centrosome in Drosophila neurons. *Neural Development* 6: 38

Pearl R, Parker SL (1921) Experimental Studies on the Duration of Life. I. Introductory Discussion of the Duration of Life in Drosophila. *Am Nat* 55: 481-509

Peel N, Stevens NR, Basto R, Raff JW (2007) Overexpressing Centriole-Replication Proteins In Vivo Induces Centriole Overduplication and De Novo Formation. *Current Biology* 17: 834-843

Poulton John S, Cuningham John C, Peifer M (2014) Acentrosomal Drosophila Epithelial Cells Exhibit Abnormal Cell Division, Leading to Cell Death and Compensatory Proliferation. *Developmental Cell* 30: 731-745

Poulton JS, Cuningham JC, Peifer M (2017) Centrosome and spindle assembly checkpoint loss leads to neural apoptosis and reduced brain size. *Journal of Cell Biology* 216: 1255-1265

Ricolo D, Castro-Ribera J, Araújo SJ (2021) Cytoskeletal players in single-cell branching morphogenesis. *Developmental Biology* 477: 22-34

Ricolo D, Deligiannaki M, Casanova J, Araújo SJ (2016) Centrosome Amplification Increases Single-Cell Branching in Post-mitotic Cells. *Curr Biol* 26: 2805-2813

Rochlin MW, Wickline KM, Bridgman PC (1996) Microtubule Stability Decreases Axon Elongation but Not Axoplasm Production. *The Journal of Neuroscience* 16: 3236-3246

Salvadores N, Gerónimo-Olvera C, Court FA (2020) Axonal Degeneration in AD: The Contribution of Aβ and Tau. *Front Aging Neurosci* 12: 581767

Sanchez-Huertas C, Freixo F, Viais R, Lacasa C, Soriano E, Luders J (2016) Non-centrosomal nucleation mediated by augmin organizes microtubules in post-mitotic neurons and controls axonal microtubule polarity. *Nat Commun* 7: 12187

Sánchez-Soriano N, Prokop A (2005) The influence of pioneer neurons on a growing motor
nerve in Drosophila requires the neural cell adhesion molecule homolog FasciclinII. *J*
Neurosci 25: 78-87
Schnorrer F, Schönbauer C, Langer CCH, Dietzl G, Novatchkova M, Schernhuber K, Fellner
M, Azaryan A, Radolf M, Stark A *et al* (2010) Systematic genetic analysis of muscle
morphogenesis and function in Drosophila. *Nature* 464: 287-291
Stevens NR, Raposo AASF, Basto R, Johnston DS, Raff JW (2007) From Stem Cell to
Embryo without Centrioles. 17: 1498-1503
Stiess M, Maghelli N, Kapitein LC, Gomis-Rüth S, Wilsch-Bräuninger M, Hoogenraad CC,
Tolić-Nørrelykke IM, Bradke F (2010) Axon extension occurs independently of centrosomal
microtubule nucleation. *Science* 327: 704-707
Tillery MML, Blake-Hedges C, Zheng Y, Buchwalter RA, Megraw TL (2018) Centrosomal and
Non-Centrosomal Microtubule-Organizing Centers (MTOCs) in Drosophila melanogaster.
Cells 7
Varmark H, Llamazares S, Rebollo E, Lange B, Reina J, Schwarz H, Gonzalez C (2007)
Asterless is a centriolar protein required for centrosome function and embryo development in
Drosophila. *Curr Biol* 17: 1735-1745
Younossi-Hartenstein A, Hartenstein V (1993) The role of the tracheae and musculature
during pathfinding of Drosophila embryonic sensory axons. *Dev Biol* 158: 430-447
Yu W, Centonze VE, Ahmad FJ, Baas PW (1993) Microtubule nucleation and release from
the neuronal centrosome. *The Journal of cell biology* 122: 349-359
585

Figure legends

Fig. 1 – *Sas-4* loss of function induces axonal misrouting defects.

(A) *Sas-4* adult fly at 6h after eclosion. (B) Graphical representation of the percentage of *Sas-*

4 surviving individuals at different life cycle stages. (C) Highlight of the three main FasII

positive nerves over a microscope image of two embryonic lateral segments. (D-G) Details of

three segments from stage 16 embryos stained with anti-FasII antibody; scale bar 10 μm. (D'-

G') Confocal lateral views of stage 16 embryos stained with anti-FasII antibody; scale bar 50

μm. (H) Penetrance and expressivity of the ISN mutant phenotypes in different *Sas-4* mutant

conditions; control is *w* (n=35); *Sas-4^{s2214}* (n=53); *Sas-4^{s2214}/Df* (n=25); *RN2Gal4xUASSas-*

4RNAi (n=15); *UbqSas-4GFP;Sas-4/Df* (n=28). (I-L) Details of three segments from stage 16

embryos stained with anti-Futsch antibody; scale bar 5 μm (I'-L') Confocal lateral views of

stage 16 embryos stained with anti-Futsch antibody; scale bar 50 μm. (H) Penetrance and

expressivity of the sensory axon mutant phenotypes in different *Sas-4* mutant conditions;

control is *Sas-4/+* (n=20); *Sas-4^{s2214}* (n=20); *Sas-4^{s2214}/Df* (n=20); *P0163Gal4xUASSas-4RNAi*

(n=20); *UbqSas-4GFP;Sas-4/Df* (n=20).

Fig. 2 - *Sas-4* loss of function induces muscle defects.

(A,B) Stage 16 embryos stained with an antibody against myosin to mark the embryonic muscle structure in green. (A-A') Control embryo and (B-B') *Sas-4* mutant embryo. (C) Schematic representation of the motor axon muscle innervation pattern of a wt (wt) and *Sas-4* mutant embryo of *Drosophila*. TN, transverse nerve; ISN, intersegmental nerve, and SN, segmental nerve. (D-H) Confocal projections of four segments from stage 16 embryos of different mutant conditions stained with anti-myosin (green and D') and anti-FasII (magenta and D'') antibodies; scale bar 10 µm. (D''-H'') composite images with the two channels (anti-myosin in green and anti-FasII in magenta). (D'''-H''') Myosin-stained muscle single-stacks. Scale bar 10 µm. (I) Penetrance and expressivity of the ISN mutant phenotypes in different *Sas-4* mutant conditions; control is *Sas-4/+* (n=27); *Sas-4^{s2214}* (n=27); *Sas-4^{s2214}/Df* (n=25); *Mef2Gal4xUASSas-4RNAi* (n=19); *RN2Gal4xUASSas-4RNAi* (n=15). (J) Penetrance and expressivity of the muscle mutant phenotypes in different *Sas-4* mutant conditions; control is *w* (n=16); *Sas-4^{s2214}* (n=16); *Sas-4^{s2214}/Df* (n=10); *Mef2Gal4xUASSas-4RNAi* (n=10); *RN2Gal4xUASSas-4RNAi* (n=10).

Fig. 3 - *Sas-6* loss of function induces axonal misrouting and muscle defects.

(A) *Sas-6* adult fly at 6h after eclosion. (B) Quantification of the number of centrioles per stage 16 embryonic segmental area in Control, *Sas-6/+* embryos, (mean 165.9 ± 26.4, n=10) and *Sas-6* embryos displaying PNS mutant phenotypes (mean 53.2 ± 21.9, n=15); unpaired t-test with Welch's correction p<0.0001. (C) Graphical representation of the percentage of *Sas-6* surviving individuals at different life-cycle stages. (D) Penetrance and expressivity of the ISN mutant phenotypes in *Sas-6* embryos; controls are *Sas-6$^{\Delta a}$* (n=19) and *Sas-6$^{\Delta b}$* (n=8); mutants are *Sas-6$^{\Delta a/\Delta b}$* (n=42). (E, F) Confocal projections of four segments from stage 16 embryos of control and *Sas-6$^{\Delta a/\Delta b}$* mutants stained with anti-myosin (magenta and E', F') and anti-FasII (green and E, F) antibodies. (E''-F'') Composite projections with the two channels; scale bar

is 10 μm. (E'''-F''') Composite projections of the two channels showing the whole embryo;

scale bar is 50μm.

Fig. 4 - Pioneer neurons have centrioles localizing to the sites of axon outgrowth

(A) Confocal image showing 3 segments of the embryonic Ventral Nerve Cord (VNC) midline

marked with an antibody against FasII and GFP driven by the RN2 driver marking the aCC,

pCC and RP2 in green. (B) Confocal section showing an aCC neuron and its centriole at the

axonal exit point, marked with an antibody against CP309; quantification of centrioles present

in different aCC quadrants and the percentage of aCC with centrioles present at the axonal

exit point (72%, n=75). (C, D) CP309 marking all centrioles and the neuronal body drawn from

the channel marking the neuron (C', D'). (C'') aCC+pCC and RP2 (D'') marked in green and

CP309 in magenta. Scale bar is 5μm. (G) Quantification of centriole numbers in control and

Sas-4 aCC+pCC and RP2 pioneer neurons; Control RP2 average 0.81 (n=22), control

aCC+pCC average 2.53 (n=41), *Sas-4* RP2 average 0.26 (n=34) and *Sas-4* aCC+pCC

average 0.87 (n=38); unpaired t-test with Welch's correction control RP2 vs *Sas-4* RP2

p=0.0005, control aCC+pCC vs *Sas-4* aCC+pCC p<0.0001.

Fig. 5 – Non-ciliated sensory neurons have centrioles

(A) Dorsal cluster neurons in a stage 15 embryo marked with anti-Futsch (22C10) in magenta

to mark the sensory neurons and Asterless (Asl) in green to mark the centrioles. (A') Inverted

channels with neurons identified and (A'') cell bodies drawn to show the centrioles within each

neuron. Scale bar is 8 μm (B) Stage 16 dorsal cluster with the neurons identified and ddaC

marked in green with ppkGAL4 driving UASCD8GFP marking class IV dendritic arborisation

neurons, with centriole in cyan and Futsch in green. Scale bar is 5 μm. Quantification of the

number of ddaC neurons with centrioles (81.4% n=43). (C) Control and (D,E) dorsal clusters

marked with Futsch in magenta CP309 in green. (C', D' and E') CP309 marking all

centrosomes and cell bodies drawn from the other channel (C'', D'' and E''). (F) Quantification

of the number of centrioles per stage 16 embryonic segmental area in Control, *Sas-4/+*

embryos, (mean 179.4 ± 30.1, n=25), *Sas-4* embryos displaying PNS mutant phenotypes

(mean 68.2 ± 21.6, n=23), *Sas-4* embryos without mutant phenotypes (mean 86.3 ± 23.8,

n=24) and *Sas-4* embryos rescued by ubiquitous expression of Sas-4 (mean 157.3 ± 51.45,

n=23); unpaired t-test with Welch's correction control vs *Sas-4* p<0.0001, control vs *Sas-4* no

defects p<0.0001, control vs *Sas-4* rescue p=0.08 and *Sas-4* vs *Sas-4* no defects p=0.009.

Fig. 6 – Pioneer motor neurons have active centrosomes

(A) aCC+pCC and RP2 (B) marked in green, CP309 in magenta and ace-tub in blue; scale

bar 5 µm. (A', B') CP309 marking all centrioles and the neuronal body drawn from the channel

marking the neuron (A, B). (A'', B'') CP309 marking all centrioles and ace-tub antibody marking

stable MTs with the neuronal body drawn from the channel marking the neuron (A, B). (C)

aCC+pCC and RP2 (D) marked in green, CP309 in magenta and γ-tub in blue. (C', D') CP309

marking all centrioles and the neuronal body drawn from the channel marking the neuron (C,

D). (C'', D'') CP309 marking all centrioles and ace-tub antibody marking stable MTs with the

neuronal body drawn from the channel marking the neuron (C, D). (E) Quantification of the

total number of centrioles in aCC+pCC and RP2 neurons and quantification of the numbers of

these which colocalize with γ-tub (n=18).

Fig. 7 - *Sas-4* loss of function induces wiggly axon phenotypes

(A-E) Confocal images of stage 16 RN2GAL4, UASCD8GFP embryonic lateral segments

(A,B) or the ventral midline (C-F) in wt (A, C) or *Sas-4* mutant background (B, D, E). (A',B')

axons marked with anti-FasII and (A'', B'', D, E) axons marked with anti-GFP, detecting the

axonal membranes. Scale bar is 10 µm in A-B and x µm in C-E. (F) Quantification of the

proportion (wiggly/normal axons) of wiggly axons in control and *Sas-4* mutant aCC and RP2

neurons; Mann-Whitney test p=0.025 in control vs *Sas-4* aCC (n=8 embryos) and p=0.0065

in control vs *Sas-4* RP2 (n=8 embryos). (G) Schematic representation of the wiggly axon

phenotypes; in Sas-4 aCC and RP2 neurons, the absence of active centrosomes gives rise to

axonal tortuosity.

Fig. 1

Fig 2

RN2>GFP			
CTRL		***Sas-4^{s2214}***	
aCC + pCC	RP2	aCC + pCC	RP2

CP309

C | D | E | F

V / A (green arrows)

GFP

C' | D' | E' | F'

GFP CP309

C'' | D'' | E'' | F''

G

Sas-4 RP2

Control RP2

Sas-4 aCC + pCC

Control aCC + pCC

***

****

centriole number
0   1   2   3   4   5