The evolution of Hutchinsonian climatic niche hypervolumes in gymnosperms

Authors: Fernanda S. Caron[1], David F. R. P. Burslem[2], Juliano Morimoto[1],[3]*

Authors' affiliations:

[1] Programa de Pós-graduação em Ecologia e Conservação, Universidade Federal do Paraná,

Curitiba, 82590-300, Brazil

[2] School of Biological Sciences, University of Aberdeen, Cruickshank Building, Aberdeen

AB24 3UU, UK

[3] Institute of Mathematics, University of Aberdeen, King's College, Aberdeen AB24 3FX,

UK

Corresponding author:

Juliano Morimoto

Institute of Mathematics, University of Aberdeen, AB24 3UE, Aberdeen

+44 (0)1224 273218

juliano.morimoto@abdn.ac.uk

Short running title

The evolution of the multidimensional niche in non-flowering woody plants

ABSTRACT

The niche is a fundamental concept in ecology. One way to represent the niche is to use multidimensional geometry known as the Hutchinsonian niche hypervolume. However, hypervolume data are complex and the biological significance of niche hypervolume properties needs to be better understood. Here, we conducted for the first time a systematic and comprehensive test of the evolution of Hutchinsonian niche hypervolume properties (volume and holes) across 65 genera and 12 families of gymnosperms, which includes many species that are endangered or threatened. Using cutting-edge computational algorithms, we measured the evolution of geometric (i.e. volume) and topological (i.e. holes) properties of gymnosperm hypervolumes across a comprehensive calibrated phylogeny. Our comparative analysis revealed weak evidence of the non-independent evolution of niche hypervolume volume and no evidence of the non-independent evolution of hypervolume holes. We also found that genera and families with low hypervolume volume such as monotypic groups like *Gingko*, likely experienced shifts in hypervolume evolutionary rates. Our results show that geometric and topological properties of gymnosperm climatic niche hypervolumes evolve independently. This agrees with competitive exclusion hypothesis where extant groups are likely to be the ones which minimise niche overlap and competition.

Keywords: climatic niche; persistence homology; phylogenetic comparative data; phylogenetic signal

INTRODUCTION

The 'niche' is arguably one of the most important concepts in ecology (Chase, 2011; McCann & Gellner, 2020). Although definitions are debated (Hutchinson, 1957; Soberón, 2007; Wennekes et al., 2012), the concept of the niche has enabled unprecedented theoretical and empirical advances in our understanding of how species interact with their environment (e.g., Blonder, 2016; Soberón & Nakamura, 2009; Soberón, 2010; Winemiller et al., 2015; and references therein). In particular, the definition of niche formulated by Hutchinson – the Hutchinsonian niche hypervolume – is widespread in the literature and is simple to apply to the large quantities of ecological data that are increasingly available. In this context, the niche hypervolume can be defined as "an abstract mapping of population dynamics onto an environmental space, the axes of which are abiotic and biotic factors that influence birth and death rates" (verbatim from Holt, 2009; see also Hutchinson, 1957). Niche hypervolumes have had a major influence on the advancement of ecological theory (Soberón & Arroyo-Peña, 2017; Blonder, 2018; Jiménez et al., 2019; Mammola & Cardoso, 2020; Soberón & Peterson, 2020; Carrasco et al., 2022; Conceição & Morimoto, 2022), as well as to uncover insights into niche evolution (Blonder, 2018; Pili et al., 2020; Bates & Bertelsmeier, 2021), niche differentiation (Carvalho & Cardoso, 2020; Huang et al., 2024), biological invasion (Tingley et al., 2014; Helsen et al., 2020; Zhang et al., 2020), community diversity (Chrisholm & Pacala, 2010; Loke & Chrisholm, 2023), ecological specialisation (Bebber & Chaloner, 2022), and global biodiversity patterns (Beaugrand et al, 2020; but see Justus, 2019).

The niche hypervolume concept has evolved with advances in theoretical and empirical ecology (Chase & Leibhold, 2003; Holt, 2009; Wiens, 2011; Kearney et al., 2017; Letten et al., 2017). One aspect of the hypervolume concept that has recently emerged, but remains untested, is the idea that hypervolumes can have holes (Blonder, 2016). Specifically, Blonder (2016) proposed the existence of holes in niche hypervolumes and conjectured that these holes could indicate environmental conditions in which species are vulnerable to be outcompeted in interspecific competition or unaccounted eco-evolutionary processes (e.g. range shifts) that lead to vacant niches. One cannot rule out the possibility of missing data generating artefacts interpreted as holes, but it is plausible – although largely untested in large scales – that holes have ecological meaning (Blonder, 2016; Conceição and Morimoto, 2022). Note that, in this conjecture, holes are representative of a fundamental ecological feature that could evolve if, for example, sister lineages share the physiological or morphological vulnerabilities that facilitate invasion portions of environmental space by other lineages. Blonder (2016) tested the

holey niche concept in the morphological niche hypervolume of a community of co-existing

finches in the Galapagos, and concluded that "well-known datasets may be described in terms

of hypervolumes with holey geometries that are potentially consistent with unexplored

mechanisms" (Blonder, 2016). Thus, Blonder's conjecture could link niche hypervolume

properties that exist in high-dimensional space to real-world ecological processes, making the

study of hypervolume properties a paramount surrogate and key target of investigation to

understanding species' ecology. However, to date, we still lack large scale tests to directly

ascertain the evolution of Hutchinsonian climatic niche hypervolume geometric and

topological properties. Yet, the concept of climatic niche hypervolumes was originally created

to represent species' climatic occupancy and it what hypervolumes are widely used in many

applications today (Blonder, 2018; Vilas et al., 2022).

Studying niche hypervolume properties is not trivial. The incomplete information available for

defining the dimensions of the hypervolume topology poses a challenge. Patchy and

heterogeneous observations prevent reconstruction hypervolume topology from bioclimatic

variables with accuracy (e.g., Peterson et al., 2018; Jiménez et al., 2019), and statistical models

to "fill" hypervolume topologies are criticized (e.g., Qiao et al., 2017; Mammola, 2019;

Guillerme et al., 2020). These limitations can be particularly insidious when estimating

hypervolume holes as high dimensionality can pose a challenge (Blonder, 2016; Conceição and

Morimoto, 2022). We recently proposed the use of algebraic topology, specifically the concept

of persistence homology, to identify holes in hypervolumes and overcome the dimensionality

challenges (Conceição & Morimoto, 2022). However, neither we nor others have yet used

approaches to identify and derive biological meaning from holes and other properties from

ecological niche hypervolumes across species within an evolutionary framework. Such

comparative studies are needed to better understand if and how ecological niche hypervolume

properties, including holes, evolved.

In this paper we present the first comparative study of the evolution of niche hypervolume

properties (volume and holes). Plants are excellent models for studying hypervolume properties

because they minimise potential confounding effects of range shifts and migration observed in

animals (e.g., Valladares et al., 2014; Menchetti et al., 2019; Häfker et al., 2022; and references

therein). Many plants are threatened by climate change and understanding the evolution of

niche hypervolumes may uncover new insights into how to protect them (Antão et al., 2020;

Singh et al., 2023). Moreover, climate niche evolution in plants is representative of similar

processes in animals (Liu et al., 2020). In this study, we focused our attention on studying ecological niche hypervolume properties of gymnosperms. Gymnosperms non-flowering woody plants with major importance for many ecosystems worldwide. Gymnosperm trees are abundant in many subtropical, temperate, and boreal forests that collectively represent over one-third of the forested regions on the planet (Lesiv et al., 2022). Moreover, gymnosperms form the group of plants has assessed as having high extinction risks (~40% of its species classified as CR, EN, VU) despite their importance for cultural and provisioning ecosystem services (Forest et al., 2018; Tagliari et al., 2023), whereas other gymnosperms are q primary source of timber (e.g., *Pinus sp*). Gymnosperms are also relatively less diverse than flowering plants, likely due to direct competition (Bond, 1989; Coiffard et al., 2012; De Boer et al., 2012), which makes them a tractable but representative group for investigating the evolution of ecological niche hypervolume properties. Given gymnosperms' broader evolutionary and socio-ecological significance, we analysed the environmental niches of 65 gymnosperm genera distributed across 12 families (Figure 1B). We used genera and families, rather than species, as our taxonomic levels of comparison because the immense computational power to estimate hypervolume geometric and topological properties prevents a species-level analysis at this stage, even for high-performance computers. For example, calculating hypervolume holes in a single hypervolume with more than a few thousand of data points in dimensions >2 is virtually impossible with current algorithms, and we still lack a properly validated approach to rarefy the point cloud (i.e. reduce number of points) without losing topological information (Lazoviskis, Levi, Morimoto, *in* prep). We constructed niche hypervolumes with 31 environmental variables and applied our recently developed approach of topological data analysis to estimate hypervolume holes (Conceição & Morimoto, 2022). We hypothesised that if niche hypervolume properties captured conserved ecological processes, then hypervolume properties would display strong phylogenetic signals. For example, if closely related genera and families are more likely to have more similar climatic niches (e.g. over- or under-dispersed point clouds in niche space), then this would be captured by our comparative analysis of climatic niche properties such as volume. Our findings advance our conceptual and biological understanding of the multidimensional niche and the evolution of its properties.

MATERIALS AND METHODS

Data collection

We obtained occurrence data from the GBIF database (https://www.gbif.org/) for all available gymnosperms. The search was conducted for Cycadopsida (40,554 entries), Ginkgoopsida

(57,488 entries), Pinopsida (5,723,563 entries), and Gnetopsida (52,648 entries) (Table S1).

We processed the GBIF data to remove duplicated points, points with locations in impossible

places (e.g., the sea), and points with large uncertainty. We assumed that observations outside

native ranges were valid in the construction of the realised niche hypervolume and were

therefore maintained. In this respect, we interpreted the realised climatic hypervolume niche as

the multidimensional representation of all measured climatic conditions where the genus or

families have survived (either within its native range or naturalised in an alien area). Recent

studies have used hypervolumes to test the differences in climatic niche in invasive vs native

ranges (e.g. Tingley et al. 2014; Zhang et al., 2020) but this is not the scope of this paper. We

only kept points derived from human observation, as opposed to fossil records or preserved

specimens. The final dataset contained 2,483,316 entries, corresponding to 764 species. The

taxonomic classification for the gymnosperms was followed according to the World of Flora

Plant List (http://www.worldfloraonline.org/, last accessed Oct 13, 2023).

Apart from the gymnosperm data, we also gathered occurrence data for four outgroup families

(see Data analyses section). These families were Psilotaceae (23,865 entries), Calycanthaceae

(10,013 entries), Amborellaceae (327 entries), and Matoniaceae (836 entries) (Table S1). For

these families, we had a total of 16,361 entries after cleaning and processing the data,

corresponding to 23 species. Climatic data were obtained from the WorldClim database using

the getData function in raster v3.6-26 (Hijmans et al., 2023). This dataset was downloaded at

a resolution of 2.5 minutes of a degree, using the argument "var" as "bio" and "tmin".

Phylogenetic relationships for the genera and families for which we had ecological data were

retrieved from Stull et al. (2021). In this paper we only considered the climatic component of

the niche, as climatic variables represent the primary determinant of variation in plant

distributions globally (e.g. Walter, 1979). We acknowledge that numerous additional abiotic

and biotic factors contribute to niche diversification in plants, including soil properties,

elevation and topography, but incorporating these niche dimensions is beyond the scope of the

current paper.

Data analyses

We performed analyses of the evolution of the Hutchinsonian niche across gymnosperms at

the level of genera and families. We discuss the limitations of this approach in the discussion

section. We used 31 environmental variables to construct the climatic niche of each taxon.

First, we extracted the climatic variables for the occurrence points of each taxon using the

extract function in *raster* v3.6-26 (Hijmans et al., 2023). Then, we conducted a principal

component analysis (PCA) to reduce the dimensionality of these environmental variables to

three axes (i.e., PC1-3). A PCA was conducted for each gymnosperm genus and family using

the function prcomp in R (R Core Team, 2024). For each taxon, we also calculated their

centroid in the first three PC axes. For this, we calculated the mean of each PC axis for the taxa

individually and for the gymnosperms. Then, we estimated the Euclidean distance between

each taxon centroid to the gymnosperm centroid. While the hypervolume volume is based on

the extreme values of the PCA, the distance between centroids can be regarded as a measure

of the uniqueness of the taxon's climatic niche. Hypervolumes were estimated using the

hypervolume_gaussian function in *hypervolume* v.3.1-4 (Blonder, 2024) with default

parameters. This algorithm provides random points that "fill" the hypervolume space (see

Blonder et al., 2014) and from which properties of the hypervolumes can be estimated. As the

hypervolume analyses are computationally intensive, we had to reduce the dataset of the genera

and families that had more than 50,000 locality entries. The genera that had their occurrences

reduced were *Abies*, *Juniperus*, *Larix*, *Picea*, *Pinus*, and *Taxus*, and the families were

Cupressaceae and Pinaceae. Inspection of the occurrence points derived from the entire and

reduced dataset showed that this reduction in the locality entries did not affect the coverage of

the taxon occurrence (Figure S1). Finally, we estimated the phylogenetic signal and adaptive

shifts for the hypervolume properties and the distance of centroids. The phylogenetic signal

was calculated using the phylosig function in *phytools* v2.3-0 (Revell, 2024). The statistics

used were Pagel's λ (Pagel, 1999), with 1,000 iterations, and Blomberg's K (Blomberg et al.,

2003), with 1,000 simulations. We also mapped the hypervolume properties in the phylogeny

using the contMap function in *phytools* v2.3-0 (Revell, 2024). As a final analysis, we estimated

adaptive shifts in the evolution of the hypervolume volume using the bayou package v2.3-0

(Uyeda & Harmon, 2014). This package fits Bayesian reversible-jump multi-optima Ornstein–

Uhlenbeck (OU) models to phylogenetic comparative data, identifying the location and

magnitude of adaptive shifts (Uyeda & Harmon, 2014). We ran the analyses at the genus and

family levels using the default prior configurations. We ran the models three times at each level

for 10 million generations, considering 30% as burn-in. All variables were log-transformed

prior to the calculations. The phylogeny used was manipulated to be used at the genus and

family level, keeping either one random representative per genus or family for this. In addition,

we also estimated the dimension 2 holes in the 3D ecological hypervolumes using our

topological data analysis approach (Conceição & Morimoto, 2022) and the *TDAstats* package

(Wadwa et al., 2018). We focused on dimension 2 holes because they represent the boundaries

of 3D sub-volumes within the 3D hypervolume. We estimated maximum niche hypervolume
hole size (i.e. largest hole size) as the maximum "survival" of dimension 2 holes upon Vietoris-
Rips filtration and the average and standard error as the mean and standard error of the survival
of all dimension 2 holes identified for a given hypervolume (Conceição & Morimoto, 2022).
Our rationale in estimating dimension 2 holes was that, if holes exist in ecological niche
hypervolumes and evolve non-independently, we could infer their biological significance from
the shared ecological properties among species. It is important to note that we cannot rule out
that hypervolumes constructed using principal components could distort the holes that exist
thereby adding noise to our phylogenetic analysis (false negative). We currently lack a
systematic study to ascertain whether our approach, which is common practice in ecology,
conserves the topological structures of the niche hypervolume. Such knowledge will have far-
reaching implications for ecological studies as many studies rely on the ability of computational
algorithms to estimate the topology of niche hypervolumes accurately. All analyses were
performed in R v4.4-0 (R Core Team, 2024).
221

RESULTS
We first tested whether ecological niche hypervolume volume evolved non-independently
across genera of gymnosperms. We found weak evidence to support the non-independent
evolution of ecological niche hypervolume volume as only Blomberg's K at the genus level
was statistically significant (Table 1). Estimates of Pagel's λ at the genus and family levels and
Blomberg's K at the family level were not statistically significant (Table 1 and Table S2).

Next, to gain further insight into the weak support for the non-independent evolution of
hypervolume volume at the genus level, we measured adaptive shifts in evolutionary rates for
the hypervolume volume. Comparing the value of the adaptive regime (θ) in the root of the
phylogeny with the adaptive regimes leading to other lineages, it is possible to assess if these
new adaptive regimes suffer an increase in value or decrease. Thus, this can indicate if a smaller
or larger hypervolume volume has been favoured in these lineages regarding their ancestor. In
gymnosperms, the root value of the adaptive regime (θ) of the hypervolume volume was
estimated as 21.548. Our model showed five major shifts in the adaptive regimes at the genus
level leading to the evolution of smaller hypervolume volumes (Figure 1C). The clades which
went through these decreasing shifts encompassed (1) *Welwitschia* (θ = 18.614, PP [posterior
probability] = 0.438), (2) *Stangeria* (θ = 18.752, PP = 0.355), (3) *Gingko* (θ = 17.273, PP =
0.839), and the outgroup (4) *Calycanthus* (θ = 18.282, PP = 0.310), and (5) *Calycanthus* and

Amborella (θ = 19.169, PP = 0.361). These results were virtually identical across all three runs
and thus, are robust.

We also analysed the evolution of hypervolume holes (Blonder, 2016). However, none of our
analyses of the average size of dimension 2 holes (Figures S2A, S3B), standard error of
estimates of dimension 2 holes (Figures S2B, S3C), and maximum size of dimension 2 holes
(Figures S2C, S3D) showed evidence of non-independent evolution of hypervolume holes.
And while the overall shape and associated persistence diagrams differed amongst themselves
(Figure 1C), pairwise distance matrices of the dimension 2 holes did not reveal any clusters of
pairwise distances which appeared similar (Figures S2E, S3F). In fact, there was no difference
between the observed and randomised pairwise distance matrices (Figure S4).

Lastly, we tested if the relative distinctiveness of the niche of each genus and family of
gymnosperms evolved non-independently, but the results were not statistically significant
(Text S1). These results suggest that ecological niche hypervolume holes, like hypervolume
volume, likely capture more dynamic ecological processes that do not necessarily evolve non-
independently among lineages.

DISCUSSION
We studied the evolution of the Hutchinsonian climatic niche hypervolume properties.
Specifically, we focused on mapping how climatic niche hypervolume geometrical and
topological properties, such as volume and holes, evolved (or not) evolved in gymnosperms.
This gave us new evolutionary insights into the climatic niche hypervolumes, which remains
one of the most significant concepts in ecology today (Chase, 2011; McCann & Gellner, 2020).
Our data supported the idea that climatic niche hypervolume properties evolved independently
across lineages at the genus and family levels. Overall, there was no phylogenetic signal on
hypervolume volume except for a statistically significant estimate of Blomberg's *K* at the genus
but not family levels. Blomberg's *K* captures the degree of variation within and among lineages
and a statistically significant value of *K* suggests that within lineage variation supersedes
variation among lineages (Meireles et al., 2020). Therefore, the statistically significant results
at the genus but not family level were driven by the relatively homogenous hypervolume
volumes of genera within the Pinaceae, Araucariaceae, and Podocarpaceae families (Figure
1B). These are genera and families with widespread climatic tolerances which are represented
by hypervolumes with comparatively higher volumes in our analysis. Contrastingly, Pagel's λ
is a scalar that correlates a trait matrix with a phylogenetic matrix and was not statistically

significant in any of our estimates. This is not surprising given the variability of hypervolume

volume across gymnosperms found in our data both at the genus and family levels (Meireles

et al., 2020).

We found a nearly 2-fold difference between the smallest and largest hypervolume by volume

in our dataset. *Gingko, Stangeria*, and *Welwitschia* genera had the lowest hypervolume volume

estimates (Figure 1C). *Ginkgo* is a living fossil that has been deemed one of the most critically

endangered gymnosperms in the world and, therefore, has a relatively narrow ecological

distribution reflected in its hypervolume volume (Forest et al., 2018). Although not classified

as (critically) endangered by the IUCN Red List, *Welwitschia* has a unique ecology and

occupies arid and semi-arid environments in sub-Saharan Africa with relatively narrow

ecological distribution, also being considered a living fossil. Likewise, *Stangeria* is not

classified as (critically) endangered by the Red List but has a unique ecology and is endemic

to southern Africa (Schönland, 1918). These findings confirm that our hypervolume approach

captures ecological characteristics of the genera and families and, thus, that the lack of evidence

for the non-independent evolution of ecological niche hypervolume properties is meaningful.

The causes of such shifts are unknown and multifactorial, but the direct competition and

diversification of angiosperms likely contributed, although cannot fully explain this pattern.

Adaptive shift results at the family level are available in Text S1.

Multidimensional hypervolumes are complex high-dimensional geometries which poses

challenges for visualisation and computations of its properties. To study climatic niche

hypervolume properties across gymnosperms, we needed to make simplifying assumptions to

guarantee the feasibility of the analysis. One such assumption was that PCA transformation

and dimensionality reduction of our point cloud preserved ecological information. This

assumption is shared with most studies using the concept of niche hypervolumes (e.g., Pianka

et al., 2017; Zhang et al. 2020; Ellis, 2022) but it is worth highlighting that no systematic study

has yet tested the implications of PCA on point cloud structure and hypervolume properties

(but see also Mahoney et al., 2017; Lu et al. 2021). PCA acts as a projection of the point cloud

into lower dimensions and can introduce biases such as for example "filling in" holes which

are present in higher dimensions. Similar biases can occur from our decision to measure

hypervolume properties at the genus and family levels, since holes which are present in the

climatic niche hypervolume of individual species might be filled in when analysing the climatic

niche hypervolume of the genus or family. However, it is important to highlight that the

opposite is not true: we cannot have artificial holes or topological structure emerging after PCA
or at the genus or family levels if these structures are absent in the raw, species-level data.
Thus, our study provides a conservative estimate of the evolution of climatic niche
hypervolume properties. We are currently working to develop algorithms which directly test
the influence of PCA on climatic niche hypervolume properties, and which would allow the
computation of hypervolume properties (particularly holes) more efficiently. This will be the
scope of future work.
313
In conclusion, the niche is a cornerstone concept in ecology which has stimulated
unprecedented advances in theoretical and empirical studies. Hutchinsonian climatic niche
hypervolume, defined as multidimensional geometries, is widespread in the literature and likely
to become increasingly common for modelling ecological niches from open data. We studied
the properties of the climatic niche hypervolumes of gymnosperms and found evidence that at
the genus and family levels, the climatic niche hypervolume geometric and topological
properties evolve independently. Our results, which are the first to systematically investigate
how Hutchinsonian hypervolumes evolved, show that ecological niche hypervolume properties
likely capture dynamic ecological characteristics of the niche. This limits the range of
inferences which can be made from Hutchinsonian climatic niche hypervolumes in
evolutionary contexts.
325

Conflict of interest

The authors have no conflict of interest to declare.

Funding

FSC was funded through a graduate scholarship from CAPES (Grant 88887.923452/2023-00).
DB is supported by the Natural Environment Research Council [NERC: NE/X000281/1 and
NE/V000071/1]. JM is supported by the Biotechnology and Biological Sciences Research
Council [BBSRC: BB/V015249/1].

Data availability

Raw data is available as supplementary material.

REFERENCES

Antão, L.H., Weigel, B., Strona, G., Hällfors, M., Kaarlejärvi, E., Dallas, T., Opedal, Ø.H., Heliölä, J., Henttonen, H., Huitu, O. and Korpimäki, E., (2022). Climate change reshuffles northern species within their niches. *Nature Climate Change*, *12*(6), pp.587-592.

Bates, O. K., & Bertelsmeier, C. (2021). Climatic niche shifts in introduced species. Current Biology, 31(19), R1252-R1266.

Beaugrand, G., Kirby, R. & Goberville, E. (2020). The mathematical influence on global patterns of biodiversity. Ecology and Evolution, 10, 6494–6511.

Bebber, D. P., & Chaloner, T. M. (2022). Specialists, generalists and the shape of the ecological niche in fungi. The New Phytologist, 234(2), 345.

Blomberg, S. P., Garland Jr, T., & Ives, A. R. (2003). Testing for phylogenetic signal in comparative data: behavioral traits are more labile. Evolution, 57(4), 717-745.

Blonder, B. (2016). Do Hypervolumes Have Holes? The American Naturalist 187. PMID: 27028084, E93–E105.

Blonder, B. (2018). Hypervolume concepts in niche-and trait-based ecology. Ecography, 41, 1441–1455.

Blonder, B. (2024). hypervolume: High Dimensional Geometry, Set Operations, Projection, and Inference Using Kernel Density Estimation, Support Vector Machines, and Convex Hulls. R Package Version 3.1-4. https://CRAN.R-project.org/package=hypervolume

Blonder, B., Lamanna, C., Violle, C. & Enquist, B. J. (2014). The n-dimensional hypervolume. Global Ecology and Biogeography, 23, 595–609.

Bond, W. J. (1989). The tortoise and the hare: ecology of angiosperm dominance and gymnosperm persistence. Biological Journal of the Linnean Society, 36(3), 227-249.

Carrasco, J., Lisón, F., Jiménez, L., & Weintraub, A. (2022). A new method to estimate the ecological niche through n-dimensional hypervolumes that combines convex hulls and elliptical envelopes. bioRxiv, 2022-03.

Carvalho, J. C. & Cardoso, P. (2020). Decomposing the causes for niche differentiation between species using hypervolumes. Frontiers in Ecology and Evolution, 8, 243.

Chase, J. M. (2011). Ecological niche theory. The theory of ecology, 93–107.

Chase, J. M., & Leibold, M. A. (2003). Ecological niches: linking classical and contemporary
approaches. University of Chicago Press.

Chisholm, R. A., & Pacala, S. W. (2010). Niche and neutral models predict asymptotically
equivalent species abundance distributions in high-diversity ecological communities.
Proceedings of the National Academy of Sciences, 107(36), 15821-15825.

Coiffard, C., Gomez, B., Daviero-Gomez, V. & Dilcher, D. L. (2012). Rise to dominance of
angiosperm pioneers in European Cretaceous environments. Proceedings of the
National Academy of Sciences of the United States of America 109, 20955–20959.

Conceição, P. & Morimoto, J. (2022). 'Holey'niche! finding holes in niche hypervolumes
using persistence homology. Journal of Mathematical Biology, 84, 58.

De Boer, H. J., Eppinga, M. B., Wassen, M. J., & Dekker, S. C. (2012). A critical transition
in leaf evolution facilitated the Cretaceous angiosperm revolution. Nature
communications, 3(1), 1221.

Ellis, C. J. (2022). A hypervolume approach to niche specialism, tested for the old-growth
indicator status of calicioids. *The Lichenologist*, *54*(6), 379-387.

Forest, F., Moat, J., Baloch, E., Brummitt, N.A., Bachman, S.P., Ickert-Bond, S.,
Hollingsworth, P.M., Liston, A., Little, D.P., Mathews, S. and Rai, H., (2018).
Gymnosperms on the EDGE. *Scientific reports*, *8*(1), p.6053.

Guillerme, T., Puttick, M. N., Marcy, A. E., & Weisbecker, V. (2020). Shifting spaces: Which
disparity or dissimilarity measurement best summarize occupancy in
multidimensional spaces?. Ecology and evolution, 10(14), 7261-7275.

Häfker, N. S. et al. (2022). Animal behavior is central in shaping the realized diel light niche.
Communications Biology 5, 562.

Helsen, K. et al. (2020). Inter-and intraspecific trait variation shape multidimensional trait
overlap between two plant invaders and the invaded communities. Oikos, 129, 677–
688.

Hijmans, R. J., van Etten, J., Sumner, M., Cheng, J. Baston, D., Bevan, A., … & Wueest, R.
(2023). raster: Geographic Data Analysis and Modeling. R Package Version 3.6-26.
https://CRAN.R-project.org/package=raster

Holt, R. D. (2009). Bringing the Hutchinsonian niche into the 21st century: ecological and
evolutionary perspectives. Proceedings of the National Academy of Sciences 106,
19659– 19665.

Huang, J., Yu, R., Ding, Y., Xu, Y., Yao, J., & Zang, R. (2024). The Relationship between Trait-Based Functional Niche Hypervolume and Community Phylogenetic Structures of Typical Forests across Different Climatic Zones in China. Forests, 15(6), 954.

Hutchinson, G. E. (1957). Population studies-animal ecology and demography-concluding remarks in Cold Spring Harbor symposia on quantitative biology, 22, 415–427.

Jiménez, L., Soberón, J., Christen, J. A., & Soto, D. (2019). On the problem of modeling a fundamental niche from occurrence data. Ecological Modelling, 397, 74-83.

Justus, J. (2019). Ecological theory and the superfluous niche. philosophical topics, 47, 105–124.

Kearney, M., Simpson, S. J., Raubenheimer, D., & Helmuth, B. (2010). Modelling the ecological niche from functional traits. Philosophical Transactions of the Royal Society B: Biological Sciences, 365(1557), 3469-3483.

Lesiv, M., Schepaschenko, D., Buchhorn, M., See, L., Dürauer, M., Georgieva, I., ... & Fritz, S. (2022). Global forest management data for 2015 at a 100 m resolution. Scientific Data, 9(1), 199.

Letten, A. D., Ke, P. J., & Fukami, T. (2017). Linking modern coexistence theory and contemporary niche theory. Ecological Monographs, 87(2), 161-177.

Liu, H., Ye, Q., & Wiens, J. J. (2020). Climatic-niche evolution follows similar rules in plants and animals. Nature Ecology & Evolution, 4(5), 753-763.

Lu, M., Winner, K., & Jetz, W. (2021). A unifying framework for quantifying and comparing n-dimensional hypervolumes. *Methods in Ecology and Evolution*, *12*(10), 1953-1968.

Loke, L. H., & Chisholm, R. A. (2023). Unveiling the transition from niche to dispersal assembly in ecology. Nature, 618(7965), 537-542.

Mahony, C. R., Cannon, A. J., Wang, T., & Aitken, S. N. (2017). A closer look at novel climates: New methods and insights at continental to landscape scales. *Global Change Biology*, **23**(9), 3934–3955. https://doi.org/10.1111/gcb.13645

Mammola, S. (2019). Assessing similarity of n-dimensional hypervolumes: Which metric to use?. Journal of Biogeography, 46(9), 2012-2023.

Mammola, S. & Cardoso, P. (2020). Functional diversity metrics using kernel density n-dimensional hypervolumes. Methods in Ecology and Evolution, 11, 986–995.

McCann, K. S. & Gellner, G. (2020). Theoretical ecology: concepts and applications.

Menchetti, M., Guéguen, M. & Talavera, G. (2019). Spatio-temporal ecological niche modelling of multigenerational insect migrations. Proceedings of the Royal Society B, 286, 20191583.

Meireles, J. E., O'Meara, B., & Cavender-Bares, J. (2020). Linking leaf spectra to the plant
tree of life. Remote sensing of plant biodiversity, 155-172.

Qiao, H., Escobar, L. E., Saupe, E. E., Ji, L., & Soberón, J. (2017). A cautionary note on the
use of hypervolume kernel density estimators in ecological niche modelling. Global
Ecology and Biogeography, 26(9), 1066-1070.

Pagel, M. (1999). Inferring the historical patterns of biological evolution. Nature, 401(6756),
877-884.

Peterson, A. T., Cobos, M. E., & Jiménez-García, D. (2018). Major challenges for
correlational ecological niche model projections to future climate conditions. Annals
of the New York Academy of Sciences, 1429(1), 66-77.

Pianka, E. R., Vitt, L. J., Pelegrin, N., Fitzgerald, D. B., & Winemiller, K. O. (2017). Toward
a periodic table of niches, or exploring the lizard niche hypervolume. *The American*
Naturalist, *190*(5), 601-616.

Pili, A. N., Tingley, R., Sy, E. Y., Diesmos, M. L. L. & Diesmos, A. C. (2020). Niche shifts
and environmental non-equilibrium undermine the usefulness of ecological niche
models for invasion risk assessments. Scientific Reports, 10, 1–18.

R Core Team. (2024). R: A language and environment for statistical computing. Version
4.4.0. https://www.R-project.org/.

Revell, L. J. (2024). phytools 2.0: an updated R ecosystem for phylogenetic comparative
methods (and other things). PeerJ, 12, e16505.

Schönland, S. (1918). A summary of the distribution of the genera of south african flowering
plants: (With Special Reference to those found in the Divisions of Uitenhage And Port
Elizabeth.). Transactions of the Royal Society of South Africa, 7(1), 19-58.

Singh, B. K., Delgado-Baquerizo, M., Egidi, E., Guirado, E., Leach, J. E., Liu, H., & Trivedi,
P. (2023). Climate change impacts on plant pathogens, food security and paths
forward. Nature Reviews Microbiology, 21(10), 640-656.

Soberón, J. (2007). Grinnellian and Eltonian niches and geographic distributions of species.
Ecology letters 10, 1115–1123.

Soberón, J. M. (2010). Niche and area of distribution modeling: a population ecology
perspective. Ecography 33, 159–167.

Soberón, J., & Arroyo-Peña, B. (2017). Are fundamental niches larger than the realized?
Testing a 50-year-old prediction by Hutchinson. Plos one, 12(4), e0175138.

Soberón, J., & Peterson, A. T. (2020). What is the shape of the fundamental Grinnellian
niche?. Theoretical Ecology, 13(1), 105-115.

Soberón, J. & Nakamura, M. (2009). Niches and distributional areas: concepts, methods, and
assumptions. Proceedings of the National Academy of Sciences 106, 19644–19650.

Stull, G. W., Qu, X. J., Parins-Fukuchi, C., Yang, Y. Y., Yang, J. B., Yang, Z. Y., ... & Yi, T.
S. (2021). Gene duplications and phylogenomic conflict underlie major pulses of
phenotypic evolution in gymnosperms. Nature Plants, 7(8), 1015-1025.

Tagliari, M. M., Bogoni, J. A., Blanco, G. D., Cruz, A. P., & Peroni, N. (2023). Disrupting a
socio-ecological system: could traditional ecological knowledge be the key to
preserving the Araucaria Forest in Brazil under climate change? Climatic Change,
176(2), 2.

Tingley, R., Vallinoto, M., Sequeira, F., & Kearney, M. R. (2014). Realized niche shift during
a global biological invasion. Proceedings of the National Academy of Sciences,
111(28), 10233-10238.

Uyeda, J. C., & Harmon, L. J. (2014). A novel Bayesian method for inferring and interpreting
the dynamics of adaptive landscapes from phylogenetic comparative data. Systematic
biology, 63(6), 902-918.

Valladares, F. et al. (2014). The effects of phenotypic plasticity and local adaptation on
forecasts of species range shifts under climate change. Ecology letters 17, 1351–1364.

Vilas, D., Fletcher Jr, R. J., Siders, Z. A., & Chagaris, D. (2022). Understanding the temporal
dynamics of estimated environmental niche hypervolumes for marine fishes. Ecology
and Evolution, *12*(12), e9604.

Wadhwa, R.R., Williamson, D.F., Dhawan, A. and Scott, J.G. (2018). TDAstats: R pipeline
for computing persistent homology in topological data analysis. *Journal of open*
source software, *3*(28), p.860.

Walter, H. (1979) Vegetation of the Earth and ecological Systems of the geo-biosphere.
Second edition. Springer-Verlag New York Inc. 1979.

Wennekes, P. L., Rosindell, J. & Etienne, R. S. (2012). The neutral—niche debate: a philo-
sophical perspective. Acta biotheoretica 60, 257–271.

Wiens, J. J. (2011). The niche, biogeography and species interactions. Philosophical
Transactions of the Royal Society B: Biological Sciences, 366(1576), 2336-2350.

Winemiller, K. O., Fitzgerald, D. B., Bower, L. M. & Pianka, E. R. (2015). Functional traits,
convergent evolution, and periodic tables of niches. Ecology letters, 18, 737–751.

Zhang, Z., Mammola, S., McLay, C. L., Capinha, C. & Yokota, M. (2020). To invade or not
to invade? Exploring the niche-based processes underlying the failure of a biological

invasion using the invasive Chinese mitten crab. Science of the Total Environment,
728, 138815.

TABLES

Table 1. Phylogenetic signal using Blomberg's K (Blomberg et al., 2003) and Pagel's λ

(Pagel, 1999) of hypervolume volume and holes at the genus level. LogL (logL0) refers to the

log-likelihood estimates of the (null) models.

Trait	λ	logL	logL0	p	K	p
Hypervolume	0.817	-128.686	-129.517	0.197	0.606	**0.009**
Mean distance	1.000	5.322	5.322	1.000	0.425	0.358
Maximum distance	0.000	-24.736	-24.736	1.000	0.397	0.503
SD distance	0.000	-15.040	-15.039	1.000	0.378	0.653

FIGURES

Figure 1. Overview of the analyses performed in the present study. **A.** Steps to analyse the
data: collection of the occurrence data in GBIF database; extraction of the bioclimatic variables
for each occurrence; PCA analysis with the bioclimatic data; estimation of the climatic niche
hypervolume volume and properties; calculation of the phylogenetic signal of the hypervolume
properties. **B.** Occurrence data for each family studied. Legend indicates the number of genera
in each family. **C.** Hypervolume properties, phylogenetic signal, and adaptive shifts estimated.
Phylogeny is mapped with hypervolume volume, in which blue corresponds to small
hypervolume volume and red to large volumes. Time scale is represented at the base of the
phylogeny. Families represented by more than one genus are highlighted in the internal
branches. Circles represent the adaptive shifts with a posterior probability higher than 0.3.
Graphs at the right represent examples of hypervolumes estimated and the persistence
homology diagrams. Green: dimension 0, Blue = dimension 1, and Pink: dimension 2
homology.
527