Evolution and development of segmented body plan revealed by *engrailed* and *wnt1* gene expression in the annelid *Alitta virens*

Arsenii I. Kairov[1,2], Vitaly V. Kozin[1*]

[1]Department of Embryology, St. Petersburg State University, Universitetskaya nab. 7-9, 199034 St. Petersburg, Russia
[2]Laboratory of Morphogenesis Evolution, Koltzov Institute of Developmental Biology RAS, Vavilov st. 26, 119334, Moscow, Russia
[*]e-mail: v.kozin@spbu.ru

Abstract: Segmentation is one of the most striking features of bilaterians, and understanding its mechanisms provides insights into the evolution of body plans. In annelids, segmentation occurs at different developmental stages through a variety of processes, yet the molecular pathways remain underexplored. Aiming to compare segmentation patterns in ontogeny and phylogeny, we analysed the expression of *Avi-en* (homologous to *engrailed*) and *Avi-wnt1* in the nereidid polychaete *Alitta virens*. Using in situ hybridization, immunofluorescence, and cell proliferation assays, we mapped the spatiotemporal expression of these genes across embryonic, larval, and postlarval stages. We found that *Avi-en* was expressed in solid lateral domains early in the unsegmented protrochophore stage and progressed through a metameric pattern, while *Avi-wnt1* expression appeared later, also aligning with segmental boundaries. At the nectochaete stage, the posterior domain of *Avi-en* expression in the growth zone expanded and splitted into two due to increased cell proliferation. The postlarval segment primordium then developed progressively, culminating in the activation of *Avi-wnt1* at the posterior border. According to available published data, the revealed pattern of gradual segment formation is unique to nereidids. The observed divergence in gene expression and cell proliferation across annelids suggests that segmentation in bilaterians did not arise from a common ancestral mechanism. Our study enhances future progress in understanding the evolution of body patterning by providing a foundation for future comparisons.

Keywords: metamery, segment addition, larval development, metamorphosis, axial elongation, growth zone, segment polarity gene, Wnt signaling, Annelida, Nereididae

Introduction

The segmentation of the annelid body has been a central focus of evolutionary developmental biology for many years (Balavoine, 2014; Irvine and Martindale, 1996; Shimizu and Nakamoto, 2001; Zattara and Weisblat, 2020). Annelids exhibit a wide variety of anatomical patterns, including homonomously segmented errant polychaetes (which have more or less identical metameres), heteronomously segmented sedentary worms (whose metameres are functionally specialized and form different tagmata), and unsegmented sipunculids and echiurids. Segment formation occurs at different stages of ontogenesis, depending on the species: throughout life, as in oligochaetes; only in postembryonic

development, as in most polychaetes; exclusively during the larval period, as in Pectinariidae;
or only during embryogenesis, as in leeches. This results in varying numbers of segments,
from oligomeric to polymeric species. The mechanisms responsible for dividing tissue layers
into metameric units are equally diverse (Hannibal and Patel, 2013; Minelli and Fusco, 2004;
Simsek and Özbudak, 2023). In polychaetes, the larval ectoderm of a solid trunk region (the
somatic plate) is segmenting through the formation of metameric ciliary rings and
intersegmental furrows, which also subdivide mesodermal tissue (Anderson, 1973). Although
this boundary-driven segmentation is considered characteristic of polychaetes, some species
(*Platynereis, Scoloplo*s) show mesodermal bands with a teloblastic origin, suggesting clonal
segmental domains (Fischer and Arendt, 2013; Özpolat et al., 2017).

In contrast, the entire body of clitellates forms through the activity of a teloblastic
growth zone, where ectodermal and mesodermal daughter cells divide in a stereotypical
manner, intermix, and generate anatomical segments, often with a shift in register relative to
the clonal domains (Weisblat and Winchell, 2020). This embryonic pattern of clitellates and
some crustaceans is referred to as lineage-driven segmentation. However, teloblasts do not
persist beyond embryogenesis, and the growth zone organization in adult oligochaetes
remains unclear. Non-clitellate annelids (i.e., "polychaetes") also possess a growth zone, but
it lacks identifiable teloblasts and begins producing segments only later in larval life. The
precise nature of the growth zone in polychaetes, whether teloblastic or diffuse, remains
uncertain (Kairov and Kozin, 2023). Discrepancies in the formation of larval segments from
the trochophore's hyposphere and postlarval segments from the growth zone suggest different
developmental mechanisms and evolutionary implications. This is reflected in the ongoing
debate surrounding the theory of primary heteronomy of segments (Giangrande and Gambi,
1998; Iwanoff, 1928). Thus, the diversity of developmental and anatomical segmentation
patterns in annelids makes them an invaluable model for comparative evolutionary analysis.

The molecular and cellular basis of segmentation in annelids is still not fully
understood. The one clear consensus in the study of annelid segment formation is the critical
role of cell proliferation, which drives axial elongation (de Rosa et al., 2005; Paulus and
Müller, 2006; Ribeiro et al., 2021; Seaver et al., 2005; Shalaeva and Kozin, 2023; Zattara and
Bely, 2013). Several genes are known to exhibit metameric expression patterns, including
Wnt ligands, *engrailed*, *hes/hey*, *twist*, NK-genes, and members of the Hedgehog signaling
pathway (Bastin et al., 2015; Dray et al., 2010; Gazave et al., 2014; Kairov and Kozin, 2023;
Kozin et al., 2019, 2016; Prud'homme et al., 2003; Saudemont et al., 2008; Seaver et al.,
2001; Seaver and Kaneshige, 2006; Steinmetz et al., 2011; Thamm and Seaver, 2008).
Functional data are available for Wnt and Hedgehog signaling, which are essential for proper
segmentation (Dray et al., 2010; Niwa et al., 2013). Less is known about the other genes,
whether their metameric expression patterns are a cause of segmentation or merely a
consequence of other patterning systems.

A particularly important segment patterning mechanism likely involves the Wnt
signaling pathway. Wnt signaling is considered a conserved feature of bilaterians, playing a
central role in anterior-posterior patterning, with activity concentrated at the posterior end
(Martin and Kimelman, 2009; Petersen and Reddien, 2009). This is supported for annelids
both by Wnt expression patterns and functional data (Bastin et al., 2015; Dray et al., 2010;
Kozin et al., 2019; Pruitt et al., 2014; Ribeiro and Aguado, 2021). Furthermore, Wnt

signaling regulates cell proliferation in the posterior growth zones of various bilaterians (Bénazéraf and Pourquié, 2013; Constantinou et al., 2020; Martin and Kimelman, 2009; Mundaca-Escobar et al., 2022; Williams and Nagy, 2017), which is especially relevant in the context of segmentation.

Homologs of the segment polarity genes *engrailed* and *wingless* have an evolutionary conserved role in morphogenetic field delimitation. While their expression has been partially documented in nereidids (Kairov and Kozin, 2023; Prud'homme et al., 2003; Steinmetz et al., 2011) and other polychaetes (Seaver et al., 2001; Seaver and Kaneshige, 2006), no comprehensive analysis has been conducted to date. Specifically, previous studies have not explored continuous gene expression during larval and postlarval development, nor have they systematically compared the expression patterns of the orthologous genes. Here we examined the coordinated expression of segmental boundary markers *Avi-en* and *Avi-wnt1* throughout the developmental stages of *Alitta virens*, from embryo to juvenile, with the goal of identifying shared mechanistic principles underlying segment formation in annelids.

A. virens is a nereidid polychaete whose embryonic and larval development (Dondua, 1975; Kairov and Kozin, 2023; Sveshnikov, 1978) closely resembles that of *Platynereis dumerilii* (Fischer et al., 2010). The cleavage pattern is spiral and unequal (heteroquadrant), and gastrulation occurs through epiboly. The early non-segmented larva, known as a trochophore, is planktonic and lecithotrophic, relying on maternally provided yolk for nutrition. The trochophore possesses an equatorial ciliated band called the prototroch, which divides the larval body into two regions: the episphere (anterior half) and the hyposphere (posterior half). The episphere develops into head structures, including the prostomium, while the hyposphere forms the trunk. Additionally, a discontinuous ciliary band, the telotroch, is located at the posterior end of the hyposphere and delineates the area that will develop into the pygidium.

During gradual metamorphosis of nereidids (Fischer et al., 2010; Sveshnikov, 1978; Wilson, 1892), the trochophore transforms into a metameric larva. The hyposphere of the metatrochophore elongates via convergent extension, and the pygidial lobes separate from the trunk. The first morphological signs of metameric organization include additional ciliary bands called paratrochs, parapodia, and segmental grooves. The metatrochophore of nereidids develops four segments: an anterior underdeveloped cryptic (zero) segment and three chaetigerous segments. Following the emergence of head appendages and initiation of parapodia movements, the larva transforms into a nectochaete. The nectochaete remains planktonic before eventually settling on the substrate. This process aligns with the onset of feeding, which occurs asynchronously among individuals. Subsequent postlarval segments are formed through anamorphic growth, involving intercalation in front of the pygidium.

Materials and Methods

Animals and fixation

Adult *A. virens* worms were collected at the White Sea near the Marine Biological Station of St. Petersburg State University. The artificial fertilization and embryo culturing were performed as described earlier (Dondua, 1975). Embryos and early larval stages were

cultivated at +14°C, then temperature was increased to +20°C from the early nectochaete
stage. Larvae and juveniles were fixed in 4% paraformaldehyde on 1xPBS/0.1%Tween20
overnight at +4 °C, followed by washes in 1xPBS/0.1%Tween20 and dehydrated in 100%
methanole. Starting from mid metatrochophore stage (135 h.p.f.), the larvae were
anaesthetised in MgCl2 before fixation.

EdU labelling

To reveal cell proliferation, larvae and juveniles were incubated in 50 μM EdU (5-
ethynyl-2'-deoxyuridine) for 15 minutes before anesthetization and fixation. Click-reaction
was performed as described earlier (Shalaeva and Kozin, 2023).

Whole mount *in situ* hybridization (WMISH) and immunohistochemistry

Plasmids with cloned fragments of *Avi-en* and *Avi-wnt1* (~1,6 kb) were provided by
R. P. Kostyuchenko. Linearized plasmids were used for synthesis of DIG-labelled RNA
probes. From 50 to 100 specimens of each developmental stage were analysed using whole
mount in situ hybridization according to the protocol described earlier (Shalaeva et al., 2021).
Incubation with RNA probes lasted 48 h; staining with BCIP/NBT lasted from 20 to 40 h.
To combine WMISH and EdU detection we performed standard WMISH protocol
followed by standard click-reaction.
For combined WMISH and antibody labeling we used the same WMISH protocol
with following modifications: samples were incubated simultaneously with anti-DIG
antibodies and primary mouse anti-acetylated tubulin antibody (Sigma T6793, dilution
1:400). Antibody washing was followed by standard NBT/BCIP staining. Stained samples
were washed in 1xPBS/0.1%Tween20 and incubated in 5% sheep serum for 1 hour followed
by incubation with secondary antibodies (Sigma T5168 anti-mouse antibodies conjugated
with Alexa-488 fluorescent dye, dilution 1:500) overnight at +4 °C. The washed samples
were mounted in 90% glycerol.

Data Visualization

Confocal images were obtained using Leica TCS SPE confocal microscope. WMISH
results were visualized using DIC optics with Axio Imager D1 microscope (Carl Zeiss).
Confocal detection of the WMISH signal was performed as described earlier (Jekely and
Arendt, 2007). All schemes and figures were made with Adobe Illustrator, Adobe Photoshop
and ImageJ.

Results

Avi-en expression

Avi-en mRNA was first detected at 32 hours post-fertilization (h.p.f.) during the
protrochophore stage. Transcripts were localized in five groups of cells arranged in a
bilaterally symmetrical pattern (Fig. 1A, 1A1). A medial, unpaired domain was observed near

the vegetal pole (Fig. 1A, A1, black arrow). Additionally, two pairs of bilaterally symmetrical domains were located on the lateral sides of the hyposphere, with the anterior domains being larger than the others.

Fig. 1. *Avi-en* expression during early larval stages in *A. virens*. Red arrows — metameric domains, black arrow — unpaired domain, green arrow — expression in the presumptive growth zone, orange arrowheads — expression in the neuroectodermal cells, white asterisk — stomodeum, yellow dotted line — prototroch, green dotted line — medial line of the body, red dotted line — area of connection between metameric domains. Scale bar — 50 μm. (A) Protrochophore, 32 hours post fertilization (h.p.f.), lateral view, animal pole is to the top, ventral side is to the left. (A1) Protrochophore, 32 h.p.f., dorsal view, deep focus. (B) Protrochophore, 38 h.p.f., lateral view. (B1) Protrochophore, 38 h.p.f., ventro-caudal view. (C) Protrochophore, 45 h.p.f., ventral view, anterior is to the top. (C1) Protrochophore, 45 h.p.f., dorso-caudal view, anterior is to the top. (C2) Protrochophore , 45 h.p.f., caudal-lateral view, ventral side is to the top. (D) Early trochophore, 55 h.p.f., ventral view, anterior is to the top, deep focus. (D1) Early trochophore, 55 h.p.f., lateral view, ventral is to the left top corner. (E) Mid-trochophore, 66 h.p.f., ventral view, anterior is to the top. (E') The same object as in (E), left side, deep focus. White dotted line — border between ectoderm and mesoderm. Blue arrows — mesodermal domains. (E1) Mid-trochophore, 66 h.p.f., ventro-caudal view, anterior is to the top. (E2) Mid-trochophore, 66 h.p.f., lateral view, anterior is to the top right corner.

By 38 h.p.f., *Avi-en* mRNA expression was evident in rows of cells along the lateral sides, stretching in the anterior-posterior direction (Fig. 1B). These rows were organized into five groups of cells, which later align with the segment borders. The unpaired domain expanded in size and connected posteriorly with the lateral domains on the dorsal side (Fig. 1B1, black arrow).

At 45 h.p.f., the expression pattern consisted of four pairs of bands arranged
perpendicular to the anterior-posterior axis, located on the lateral sides of the body (Fig. 1C).
These bands corresponded to the distinct cell groups observed in the previous stage. The most
anterior band was the shortest, while the other three bands were longer and interconnected at
their ventrolateral ends (Fig. 1C, 1C2). The last pair of bands was the widest and
corresponded to the two undiverged domains observed in the 38 h.p.f. embryo (Fig. 1C, C1,
C2, red and green arrows). The posterior unpaired domain connected these wide posterior
bands on the dorsal side (Fig. 1C1, black arrow).
At the early trochophore stage (55 h.p.f.), the *Avi-en* expression pattern consists of
four pairs of bands (Fig. 1D, D1). Unlike the previous stage, these bands no longer connect at
the ventrolateral sides (Fig. 1D). The posterior unpaired domain persists on the dorsal side,
and there is an emerging tendency for the last pair of bands to separate into distinct domains
(Fig. 1D1, black arrow).
By the middle trochophore stage (66 h.p.f.), *Avi-en* mRNA was detected in five pairs
of stripes of ectodermal cells arranged in a bilaterally symmetrical pattern (Fig. 1E, 1E2). A
corresponding metameric signal was observed in internal (mesodermal) cells, aligning with
the superficial ectodermal domains (Fig. 1E'). The most anterior pair of stripes was the
shortest and restricted to the ventrolateral regions of the larva. The remaining stripes were
curved in an arc, with the left and right halves joining at the ventral side but not extending to
the dorsal midline. Cells located at the ventral side, which are neuroectodermal in nature,
exhibited a posterior phase shift relative to the lateral regions of expression, indicating a
mosaic pattern. The most posterior domain formed an almost circular shape, corresponding to
the prospective growth zone (Fig. 1E1).
At the early metatrochophore stage (116 h.p.f.), the *Avi-en* expression pattern
remained largely unchanged. There were five metameric transverse bands, which were
circular except for the most anterior one, which was interrupted on the ventral and dorsal
sides (Fig. 2A, A1). The expression pattern became more mosaic, as the lateral domain parts
were now distinct from the ventral (neuroectodermal) regions. Metatrochophores exhibited
external segmentation features, including segmental grooves and paratrochs (metameric
circular ciliary bands), enabling us to correlate metameric domains with segment borders.
The first and second stripes were located between the prototroch and the first intersegmental
groove. The third and fourth expression domains aligned with segmental grooves at the
boundaries between segments 1/2 and 2/3, respectively. The posterior-most expression band
corresponded to the border between the last segment and the area of the future pygidium (Fig.
2A1, green arrow).
In nereidid polychaetes, paratrochs are situated at the posterior part of each segment
(Bastin et al., 2023; Fischer et al., 2010), and the growth zone lies between the third paratroch
and the telotroch (Fischer et al., 2010; Nielsen, 2004; Starunov et al., 2015). The relative
positions of *engrailed* expression stripes and ciliary bands have been previously documented
only in *P. dumerilii* (Steinmetz et al., 2011). Using confocal microscopy, we observed that
the third and fourth metameric *Avi-en* domains were positioned posterior to the paratrochs
(Fig. 3A), corresponding to the anterior parts of chaetigerous segments 2 and 3. The
posterior-most circular domain was located between the last paratroch and the telotroch,
indicating that it lies at the anterior border of the pygidium and coincides with the prospective

growth zone. The two anterior expression stripes were not confined to morphological boundaries. The second expression domain was located posterior to the metatroch, while the first domain was anterior to it (Fig. 3). Consequently, the first and second domains correspond to the anterior regions of segment 0 (a cryptic segment) and segment 1, respectively.

Fig. 2. *Avi-en* expression during segmented larvae and juvenile stages in *A. virens*. Scale bars — 50 μm. Red arrows — metameric domains, green arrow — expression in the presumptive growth zone, orange arrowhead — expression in the neuroectodermal cells, yellow dotted line — prototroch, white asteriks — stomodeum or pharynx. White dotted lines emphasize metameric domains extended on two adjacent segments. (A), (B), (C), (C') — ventral view, anterior is to the top; (A1) — caudal view, ventral is to the right bottom corner; (D), (D1), (E), (E1) — ventral views, anterior is to the top left corner; (B1) — anterior view; (F), (H), (H1) — ventral views, anterior is to the left; (G), (G1) — dorsal views, anterior is to the left. (A) Early metatrochophore, 116 h.p.f. Expression is similar to that of the mid trochophore. (A1) Early metatrochophore, 116 h.p.f. *Avi-en*+ cell bands are circumferential. (B) Mid metatrochophore, 135 h.p.f. (B') *Avi-en*+ mesodermal cells (red dotted line) in the area of the future growth zone. (B1) Mid-metatrochophore, 135 h.p.f., there are 4 *Avi-en*+ groups of cells closer to the dorsal side. (B2) Mid-metatrochophore, dorso-lateral view, anterior is to the top. Metameric domains expand to the ventral and dorsal sides. (C) Late metatrochophore, 160 h.p.f. (C') Magnified image of the pharynx. Some specimens have *Avi-en*+ cells (yellow dotted line) in the anterior pharynx. (D) Early nectochaete, 180 h.p.f. (D1) Early nectochaete, the same object as in (D), magnified image of caudal end, deep focus. There are two adjacent rows of *Avi-en*+ cells in the growth zone (green arrowheads). (E) Mid-nectochaete, 200 h.p.f. (E1) Mid-nectochaete, the same object as in (E), magnified image of the caudal end, deep focus. There are two bands of *Avi-en*+ cells in the pygidium. (F) Juvenile with the forming 4[th] chaetigerous (1[st] postlarval) segment, 300 h.p.f. The expression signal is absent in neuroectodermal cells. (G) Juvenile with the 4[th] segment. Two rings of *Avi-en*+ cells in the pygidium. (G1) The same object as in (G), deep focus. Two rows of *Avi-en*+ cells indicate the formation of a new segment. (H), (H1) 5-segment juvenile.

At the mid-metatrochophore stage (135 h.p.f.), the expression pattern remains largely unchanged from the previous stage and consists of five metameric domains. The last two domains are circular, while the others are interrupted on the dorsal side of the body (Fig. 2B, 2B1, 2B2). Ventromedial neuroectodermal *Avi-en*-positive cells increase in number and shift further posteriorly relative to the flanking lateral fragments. This shift enhances the mosaic pattern, as neuroectodermal cells no longer form distinct bands with the ventrolateral

domains (Fig. 2B). Within the last expression domain, which corresponds to the prospective growth zone, *Avi-en* mRNA was detected in ectodermal and also in deep mesodermal cells (Fig. 2B, red dotted line). Additionally, *Avi-en* expression was observed in four groups of cells located in the episphere, closer to the dorsal side (Fig. 2B1).

At the late metatrochophore stage (160 h.p.f.), the *Avi-en* expression pattern becomes increasingly mosaic (Fig. 2C, 3B). Unlike the stable circular posterior domain, ventral neuroectodermal cells no longer form distinguishable bands with the ventrolateral ectodermal cells. However, the ventrolateral domains can still be grouped into four pairs. *Avi-en* mRNA disappeared from the lateral and dorsal regions of the larval body. The third and fourth metameric domains span both sides of the furrows, extending across the boundaries of two adjacent segments (Fig. 2C, white dotted line). Additionally, some specimens demonstrated *Avi-en*-positive cells in the pharynx (Fig. 2C').

Fig. 3. *Avi-en* and *Avi-wnt1* expression (red channel) relative to ciliary bands (green channel). Maximum projections of confocal Z-stacks of ventral views, anterior is to the top. Scale bar — 50 μm. White asterisk — stomodeum/pharynx. Par 1…3 — paratrochs of the 1st - 3rd segments, pt — prototroch, tt — telotroch, met — metatroch, red arrows — metameric expression domains, white arrowhead — expression domain in the growth zone, yellow arrow — terminal expression domain. (A) *Avi-en* expression, early metatrochophore. 3rd and 4th rows of expression are posterior to the 1st and 2nd paratrochs, 5th row is located between the 3rd paratroch and telotroch. (B) *Avi-en* expression, late metatrochophore. (C) *Avi-wnt1* expression, early metatrochophore. Metameric domains are located posterior to metatroch/paratrochs. The most posterior band of expression is located between the 3rd paratroch and telotroch, which exactly corresponds to the growth zone position.

Previously, we described the *Avi-en* expression pattern during the final stages of metamorphosis in *A. virens* (Kairov and Kozin, 2023). In this study, we confirm our earlier conclusions through additional analysis using morphological and proliferative markers. At the early nectochaete stage (180 h.p.f.), the most significant changes were observed in the posterior ring-shaped domain at the boundary between the third segment and the pygidium (Fig. 2D). This domain expanded along the anterior-posterior axis, forming two rows of cells instead of one, as seen in earlier stages. The expansion occurred progressively from the ventral to the dorsal side (Fig. 2D1). Anteriorly, the signal disappeared from the episphere.

At 200 h.p.f., nectochaetes exhibited two circular rows of *Avi-en*-positive cells in the anterior part of the pygidium (Fig. 2E, 2E1). These rows likely represent two parts of *Avi-en* expression domain that were adjacent at the early nectochaete stage but diverged during cell proliferation and pygidium growth.

To identify exact areas of cell proliferation, we performed an EdU assay. At 200 h.p.f., EdU incorporation was detected in the anterior end of the nectochaete (episphere), in

the trunk neuroectoderm, and prominently in the pygidium (Fig. 4A). In the pygidium, EdU labeling was especially intensive, with a lower density of labelled nuclei in its posterior region. During the formation of the fourth segment (Fig. 4B, 4C), the distribution of EdU-labelled nuclei resembled that in nectochaetes. However, after the separation of the new segment, it showed the highest labelling density, whereas the pygidium contained fewer EdU-labeled nuclei. This confirms that the pygidium and the emerging segment are regions of active cell proliferation.

Fig. 4. Cell proliferation reviled by EdU incorporation (red channel) during the first postlarval segment development in *A. virens* nectochaetes and juveniles. Maximum projections of confocal Z-stack of ventral views, anterior is to the top. Scale bar — 50 μm. White asterisk — pharynx, white dotted line — intersegmental borders, l.s. — larval segment, p.l.s. — postlarval segment, pyg — pygidium. (A) Mid-nectochaete, 200 h.p.f. (B) Juvenile with the forming 4th chaetigerous (1st postlarval) segment, 300 h.p.f. (C-D) 4-segment juvenile. (D) EdU assay combined with *Avi-en in situ* hybridization. *Avi-en*+ cells (cyan channel) colocalize with the EdU+ nuclei in the growth zone (white arrowhead).

Following the appearance of a segmental groove within the pygidium, specifically during the growth and morphogenesis of the fourth segment, the signal was no longer detected in the larval segments (Fig. 2F). Towards the posterior part of the body, two *Avi-en* expression domains were observed: one located at the anterior border of the developing fourth segment (the first postlarval segment), and the other along the anterior border of the pygidium, corresponding to the growth zone of juvenile worms. Once the first postlarval segment is fully formed, expression in neuroectodermal cells resumes anterior to the fourth segment (Fig. 2G'). *Avi-en*-positive cells were also detected at the anterior border of the fourth segment and the anterior border of the pygidium (Fig. 2G). In more mature individuals with elongated pygidium, two bands of cells were observed within the pygidium, similar to those seen in the middle nectochaete stage (Fig. 2G1).

The expression pattern of *Avi-en* in five-segment juveniles was found to be comparable to that observed in four-segment juveniles (Fig. 2H). The signal was detected in neural cells and at the boundaries between posterior segments. Within the pygidium, two circular rows of cells exhibiting the signal were retained: one in a more anterior position (at the anterior morphological boundary of the pygidium) and another located posterior to the

first one (Fig. 2H1). The distance between these two rings varied among individuals,
reflecting the ongoing growth process of the segment primordium. In fully differentiated
postlarval segments (the fourth and fifth chaetigers), metameric *Avi-en* expression domains
were positioned on both sides of the groove, spanning the territory of two adjacent segments
(Fig. 2H1, white dotted line).

Avi-wnt1 expression

Previously, *Avi-wnt1* expression was partially described in *A. virens* trochophores
(Kozin et al., 2019). In the present study, we detected the signal at an earlier stage than
previously reported and conducted a detailed analysis of segmented larvae and juveniles. At
38 h.p.f., *Avi-wnt1* transcripts were localized in superficial cells at the posterior pole of the
embryo (Fig. 5A). In 45 h.p.f. protrochophores, the posterior expression domain slightly
expanded, but no new domains appeared (Fig. 5B).

345
Fig. 5. *Avi-wnt1* expression in *A. virens* development. Black arrow — caudal domain of expression, red arrows -
metameric domains of expression. (A), (G), (H), (H1), (I), (J), (K) — ventral view, anterior is to the top left
corner; (B), (C), (E), (F), (F1) — ventral view, anterior is to the top; (C1), (E1), (E2) — lateral view, anterior is
to the top, ventral is to the left; (D) — caudal view, ventral is to the top. (E3) — dorso-caudal view, anterior is
to the top; (E2) — ventro-caudal view, anterior is to the top; (L), (L1) — dorsal view, anterior is to the top left
corner. (A) Protrochophore, 38h.p.f. Only the terminal posterior domain of expression is detected. (B)
Protrochophore, 45h.p.f. (C), (C1) Mid-trochophore, 66 h.p.f. Metameric bands arise at ventro-lateral sides. Red
dotted line — area of connection between metameric domains (D) Early metatrochophore, 106 h.p.f. (E)-(E3),
Early metatrochophore, 116 h.p.f. Metameric bands of expression are located along segmental borders. (E2),

(E3) Terminal posterior domain of expression consists of two parts. (F)-(F2) Mid-metatrochophore, 135 h.p.f. (G), (G') Late metatrochophore, 160 h.p.f. (G') Lateral view, deep focal plane, ventral is to the right. The terminal expression domain marks the hindgut. (H), (H1) Early nectochaete, 180 h.p.f. (I) Mid-nectochaete, 200 h.p.f. Metameric domains are not detectable. (J) Juvenile with the forming 4th chaetigerous (1st postlarval) segment, 300 h.p.f. (K) 4-segment juvenile. (L), (L1) 5-segment juvenile. Metameric expression domains mark the borders of postlarval segments.

By the mid-trochophore stage (66 h.p.f.), the *Avi-wnt1* expression pattern became metameric. Four transverse bands of ectodermal cells expressing *Avi-wnt1* emerged on both lateral sides of the hyposphere (Fig. 5C). The two posterior metameric bands formed an arc and were continuous on the ventral side of the body (Fig. 5C1, red dotted line). The posterior medial expression domain expanded anteriorly on the ventral side of the larval body and intersected with the last two metameric bands.

In metatrochophores, the *Avi-wnt1* expression pattern did not change significantly (Fig. 5D-E). The posterior expression domain in the proctodaeum region included two parts: a more ventral and a dorsal one (Fig. 5D, 5E2, 5E3). These domains incorporated both superficial and internal cells. The metameric expression bands extended along the segment grooves (Fig. 5E, 5E1) and posterior to the paratroch (Fig. 3C). Notably, these bands were positioned closer to the paratroch than the *Avi-en* bands. The most posterior metameric band was located in front of the telotroch, corresponding to the *Avi-en* expression ring in the future growth zone (Fig. 3C, green arrow). The anterior-most *Avi-wnt1* expressing band was situated posterior to the metatroch (Fig. 3C).

In middle and late metatrochophores, *Avi-wnt1* expression in metameric domains gradually faded and completely disappeared by the 200 h.p.f. nectochaete stage (Fig. 5F-5I). However, in segmented larvae, the signal was detected in a circular band at the posterior border of the third segment (Fig. 5H, 5H1). Terminal *Avi-wnt1* expression continued in the hindgut wall (on the ventral and dorsal sides) (Fig. 5G, 5G', 5H1, 5I).

At the onset of anamorphic growth and postlarval segment formation, the posterior medial *Avi-wnt1* expression domain decreased in length (Fig. 5J, 5K). The signal was detected in the most terminal cells of the hindgut. By the time the sixth segment formed, the *Avi-wnt1* expression pattern became metameric again. In addition to the signal in the hindgut (Fig. 5L1), mRNA was detected in cells at the posterior border of postlarval segments on the ventral and lateral sides of the body (Fig. 5L). We were unable to detect *Avi-wnt1* expression in the growth zone (anterior pygidium) at this stage.

Discussion

Analysis of *Avi-en* and *Avi-wg* expression throughout the developmental stages of *A. virens*

Avi-en expression was first detected at the protrochophore stage. Initially, the expression was bilaterally symmetrical, with a single unpaired domain at the posterior end and two pairs of lateral domains (Fig. 6). These lateral domains gradually expanded along the anterior-posterior (AP) axis, forming two longitudinal bands subdivided into five distinct units. This expansion was likely driven by ongoing cell proliferation or the induction of *Avi-*

en in new territories. In contrast, *Avi-wnt1* expression appeared later than *Avi-en*, suggesting that *Avi-wnt1* is not essential for the initiation of *Avi-en* expression. This timing is consistent with RNA-seq data from *Platynereis dumerilii* transcriptomic studies (Chou et al., 2016; Pruitt et al., 2014).

Fig. 6. Schematic representation of *Avi-en* and *Avi-wnt1* expression, which patterns larval segments during early stages of *A. virens* development. Anterior is to the top. Black asterisks — stomodeum, black strokes — prototroch. Upper row — *Avi-en* expression, lateral view; middle row — *Avi-en* expression, ventral view, lower row — *Avi-wnt1* expression, ventral view.

The five groups of *Avi-en*-expressing cells represent the future metameric domains, which became clearly visible by the mid-trochophore stage (Fig. 1, 6). These domains corresponded to the molecular boundaries of future body segments: the anterior four were confined to larval segments, while the posterior-most was associated with the pygidium and the prospective growth zone.

The formation of the *Avi-en* metameric pattern occurred progressively. Initially, domains corresponding to the cryptic (0), 1st, and 2nd segments became distinct. Later, the prospective pygidial and 3rd segmental domains became individualized. Examination of the domain shapes for segments 1 and 2 (Fig. 1C) revealed differences in the degree of separation between dorsal and ventral regions: while the ventral parts remained connected, the dorsal parts had already separated. Eventually, the posterior medial *Avi-en* domain merged with the last band associated with the pygidium and prospective growth zone. The divergence of *Avi-en*-positive cells was likely driven by cell migration from the dorsal to the lateral sides, a process previously observed during somatic plate formation (Anderson, 1973; Wilson, 1892).

At the embryonic stages, the *Avi-en* signal was confined to the lateral sides, but by the mid-
trochophore stage, its expression expanded ventrally to include the neuroectoderm.
Rearrangements in neuroectodermal expression may have been caused by convergent
extension within the neuroectoderm (Steinmetz et al., 2007).
Avi-wnt1 expression was detected later, in cells at the posterior pole of
protrochophores, with metameric domains appearing only in trochophores. During early
larval development, the metameric bands of *Avi-wnt1* (except for the most posterior band)
were positioned more anteriorly than the corresponding *Avi-en* bands. This aligns with known
data showing that *wnt1* is expressed at the posterior border of body parts, while *engrailed* is
expressed at the anterior border, as observed in *P. dumerilii* (Prud'homme et al., 2003) and
other bilaterians (Janssen et al., 2010; Vellutini and Hejnol, 2016).
At the late metatrochophore stage, *Avi-en* metameric domains were found within the
territories of two adjacent segments (Fig. 2C), likely indicating the coexpression of *Avi-wnt1*
and *Avi-en* along the posterior segment border. Similar relationships were observed during
postlarval segment development: initially, *Avi-en* was expressed only at the anterior border of
the segment primordium. Later, its expression expanded across the furrow to reach the
posterior border of the adjacent anterior segment (Fig. 2H1), while *Avi-wnt1* remained
confined to the posterior border of the segment (Fig. 5L). During both trochophore and
metatrochophore stages of larval development, coexpression of *Avi-wnt1* and *Avi-en* was
observed in the presumptive growth zone (Fig. 3). However, this coexpression diminished as
Avi-wnt1 expression in the growth zone disappeared in nectochaetes.
The posterior medial domain of *Avi-wnt1* was confined to the proctodaeum (Fig. 5, 6).
Starting at the metatrochophore stage, this domain developed into two parts: a dorsal and a
ventral region (Fig. 5D). By the late metatrochophore stage, *Avi-wnt1* expression became
detectable internally, within the hindgut anlage (Fig. 5F), likely due to invagination during
hindgut morphogenesis, as previously noted (Kulakova et al., 2008).
Significant changes in *Avi-en* expression occurred in the most posterior domain during
the formation of the first postlarval segment. At the late metatrochophore stage, *Avi-en* was
expressed in a circular arrangement of superficial (ectodermal) and some inner (mesodermal)
cells. Subsequently, at the early nectochaete stage, this domain expanded along the AP axis,
forming two rows of superficial cells (Fig. 2D, 7). These rows in the pygidium eventually
separated due to cell proliferation, as demonstrated by the colocalization of *Avi-en* expression
signal with EdU labeling (Fig. 4). These processes reflect the sequential steps involved in the
formation of the first postlarval segment (Kairov and Kozin, 2023). During subsequent
postlarval growth, the emergence of new expression stripes of *Avi-en* and *Avi-wnt1*
consistently preceded the formation of the segment rudiment (Fig. 7).
Thus, the expression of *Avi-wnt1* and *Avi-en* accompanied the development of both
larval and postlarval segments. In both cases, the metameric *Avi-en* pattern emerged as a
result of the subdivision of an initially continuous expression domain. In contrast, *Avi-wnt1*
expression appeared later than *Avi-en* at the segment boundaries and exhibited a metameric
nature from its onset. The mechanisms inducing *Avi-en* and *Avi-wnt1* metameric expression
in the hyposphere remain the most challenging question.

Fig. 7. Schematic representation of *Avi-en* and *Avi-wnt1* expression during formation of the first postlarval
segment in *A. virens*. Anterior is to the top. Developmental stages are indicated above.

Evolutionary conserved and derived aspects of *Avi-en* and *Avi-wnt1* expression

The dynamics of *Avi-en* and *Avi-wnt1* expression reveal numerous developmental
traits in *A. virens*, some unique to nereidid polychaetes and others conserved across annelids.
In the embryos of *A. virens* and *P. dumerilii*, *en* and *wnt1* transcripts appear early in
development, specifically during the protrochophore stage. By the mid-trochophore stage, the
expression domains are clearly associated with the pygidium and future segments
(Prud'homme et al., 2003; Steinmetz et al., 2011). The dynamics of *en* expression in
nereidids resemble those in another polychaete with lecithotrophic larvae, *Capitella teleta*. In
all these species, *engrailed* mRNAs initially appear at the lateral sides of the hyposphere,
with expression domains subsequently expanding to the ventral and dorsal sides (Seaver and
Kaneshige, 2006). However, in *C. teleta*, *engrailed* expression begins only after the
differentiation of ciliary bands, whereas in *A. virens*, it occurs prior to ciliation. This
difference can be interpreted as heterochrony, implying accelerated pattern formation in
nereidids. In the basal polychaete *Chaetopterus* sp., *engrailed* expression begins in the L4
larva at the lateral sides of the body. This pattern later resolves into metameric domains and
expands dorsally (Seaver et al., 2001), but only after the formation of morphological segment
boundaries. Another common feature across these species is the detection of *engrailed*
mRNAs in the mesoderm of the prospective growth zone. In *A. virens* and other studied
polychaetes (*C. teleta*, *Chaetopterus* sp., and *Hydroides elegans*), *engrailed* expression first
appears in superficial cells, followed by expression in deeper mesodermal cells. This pattern
may arise from various processes, such as the hypothetical migration of cells from the surface
to the interior (Seaver and Kaneshige, 2006) or as a result of inductive interactions. This

feature of expression appears to be conserved among annelids but is absent in leeches. In the latter, *engrailed* expression is detected in the mesodermal descendants of teloblasts (Lans et al., 1993), but not in the teloblasts themselves, which together constitute the embryonic growth zone.

In the sedentary polychaete with planktotrophic larvae *H. elegans*, *engrailed* mRNA is detected starting only from the trochophore stage, first in the ectoderm, and then metamerically in deep cells (future bristle-bearing sacs) (Seaver and Kaneshige, 2006). By the time of morphological segmentation, the signal is observed in the mesoderm. A comparison of the dynamics of *engrailed* expression shows that in nereidids and *C. teleta* (polychaetes with lecithotrophic larvae), expression occurs earlier than in species with planktotrophic larvae. Moreover, in the echiurid *Urechis unicinctus*, transcripts of *engrailed* and other genes associated with segmentation are up-regulated quite early in development (Hou et al., 2019). All of this confirms the results of the recent study, which proposes that species with lecithotrophic larvae develop a trunk earlier than species with planktotrophic larvae (Martín-Zamora et al., 2023).

Engrailed genes are known to participate in the specification of many serial structures, such as CNS ganglia, nephridia, and chaetal sacs. All studied annelids showed *engrailed* expression in the nervous system (Bely and Wray, 2001; Kairov and Kozin, 2023; Patel et al., 1989; Prud'homme et al., 2003; Seaver et al., 2001; Seaver and Kaneshige, 2006). Moreover, neural expression is identified in other bilaterian animals, such as vertebrates, arthropods, and echinoderms (Byrne et al., 2005; Condron et al., 1994; Omi and Nakamura, 2015). These data suggest that the hypothesis proposed for arthropods about the ancestral expression of *engrailed* in the metameric elements of the nervous system and subsequent co-option in the process of segment establishment and patterning may also apply to annelids (Chipman, 2010; Patel et al., 1989).

The presence of *wnt1* at the posterior end of the body is a conserved feature of bilaterians (Loh et al., 2016; Martin and Kimelman, 2009; Petersen and Reddien, 2009). The corresponding *Avi-wnt1* expression domain is the earliest one in *A. virens*. Among annelids, *wnt1* expression in the hindgut anlage was characterized in *C. teleta* and the leech *Helobdella robusta* (Cho et al., 2010; Seaver and Kaneshige, 2006). There is metameric *wnt1* expression in their trunk, ectodermal in the leech, and mesodermal in *C. teleta*. Other *wnt* paralogs are expressed metamerically too, forming ectodermal stripes (e.g., *wnt16a* and *wnt16b* in *H. robusta*). Nereidid polychaetes are also characterized by metameric *wnt* expression both in larval and postlarval development (Janssen et al., 2010; Kozin et al., 2019; Prud'homme et al., 2003; Pruitt et al., 2014). Widely distributed metameric Wnt activity in annelids suggests that Wnt signaling was involved in segmentation and body patterning in the annelid ancestor.

Comparison of segmentation gene activity in annelids and arthropods provides rich material for evolutionary interpretations. Despite the similarities in *wnt1* and *engrailed* patterns between nereidids and other annelids, only nereidids have a complementary and striped arrangement of *wnt1* and *engrailed* domains of expression resembling arthropods. Virtually all arthropods and related to them onychophorans and tardigrades have metameric distribution of *wnt1* and *engrailed* mRNAs during parasegment development (Eriksson et al., 2009; Gabriel and Goldstein, 2007; Williams and Nagy, 2017). But the most studied annelids do not exhibit such a pattern. Leeches have metameric *engrailed* expression, but ablation of

engrailed-positive cells doesn't affect segmentation (Seaver and Shankland, 2001).
Therefore, nereidids appear to have gained a new mechanism of segment patterning similar to
arthropods but lacking in other annelids. However, unlike in arthropods, *engrailed* in
nereidids is expressed both at the anterior and posterior parts of the segment during both
larval (Fig. 2C, 5F) and postlarval stages (Fig. 2H1, 5K). The suggested coexpression of *Avi-*
en and *Avi-wnt1* raises questions about the function of these genes as segment polarity genes.
Furthermore, the temporal sequence of the appearance of the *wnt1* and *engrailed* metameric
bands is similar — in arthropods, during terminal growth, the *engrailed* band appears before
the *wnt1* band (Chesebro et al., 2013; Lim and Choe, 2020). In *A. virens*, the metameric
expression of *engrailed* appears much earlier than *wnt1* in the development of both larval and
postlarval segments. This indicates a fundamentally different interaction between *engrailed*
and Wnt signaling in the segment patterning of *A. virens* compared to arthropods.

Engrailed activity at the borders of segments and the pygidium in nereidids supports
the hypothesis of the ancestral role of *engrailed* in boundary formation (Vellutini and Hejnol,
2016). Our data align with the evolutionary scenario in which the expression of *engrailed*
was not initially associated with segmentation, and *engrailed* was likely co-opted into
segmentation independently in different phylogenetic lineages. The exact timing of this co-
option in nereidids remains unclear, as this expression pattern has been revealed only in this
family. Gene expression studies in other annelid families are essential to clarify this issue.

The expression of *engrailed* in the growth zone is characteristic of many studied
annelids (Bely and Wray, 2001; Prud'homme et al., 2003; Seaver and Kaneshige, 2006). In
Chaetopterus sp., expression of *engrailed* posterior to the C2 segment can be interpreted as
expression in the growth zone (Seaver et al., 2001). In both *C. teleta* and *Chaetopterus* sp.,
engrailed mRNA is detected in the growth zone only in pre-metamorphic larvae, whereas in
A. virens, the most posterior band of *engrailed* expression is evident as early as the
trochophore stage. Spatially, this domain corresponds to the future expression region of
multipotency markers (Kostyuchenko, 2022; Kozin and Kostyuchenko, 2015; Rebscher et al.,
2007). Based on this, we hypothesize that in *A. virens* (and other nereidids), there is an early
subdivision of the hyposphere into the segmental part and the future pygidium.

To summarize, *A. virens* exhibits both conserved features in the expression of
engrailed and *wnt1* (such as the presence of *engrailed* transcripts in the elements of the
nervous system, and in the ectoderm and mesoderm of the growth zone, as well as *wnt1*
expression in the hindgut and metameric domains), as well as traits unique to nereidid
polychaetes. These include the early induction of *engrailed* expression in the growth zone
and segmental rudiments, as well as the complementary arrangement of the *engrailed* and
wnt1 expression domains.

Relationships between larval and postlarval segmentation

The revealed differences in *engrailed* expression during the larval period across
various annelids prompt us to refer to Ivanov's theory on the primary heteronomy of
segments (Giangrande and Gambi, 1998; Ivanova-Kazas, 1978, p. 978; Iwanoff, 1928;
Schroeder and Hermans, 1975). The similar pattern of *engrailed* expression in polychaete
larvae—characterized by the subdivision of a continuous longitudinal domain into distinct

metameric spots, expanding from the lateral sides to the dorsal and ventral sides (Fig. 6)—
may indicate a more conservative mode of larval segment patterning. However, no such
similarity was observed in the molecular patterning of postlarval segments, which may
support the idea that larval segment development represents an ancestral characteristic. The
exact ancestral mechanisms involved in segment specification in annelids remain
controversial. In clitellates (oligochaetes and leeches), germ-band cells acquire their identity
upon the division of teloblasts. These nascent primary blast cells proliferate stereotypically
and assume specific positions within a segment, a process known as lineage-driven
segmentation (Weisblat and Kuo, 2014; Weisblat and Winchell, 2020). Whether this process
applies to non-clitellate annelids remains uncertain. In *P. dumerilii* early development,
teloblastic activity resembling that of clitellate annelids was observed in mesodermal bands,
but not in the ectoderm (Fischer and Arendt, 2013; Özpolat et al., 2017). In *C. teleta* and *H.*
elegans, no teloblasts have been described in either the ectoderm or mesoderm (Irvine and
Seaver, 2006; Seaver et al., 2005). However, in *C. teleta*, *nanos* expression was detected in a
distinct ring of ectodermal cells anterior to the telotroch, which may suggest the presence of
ectodermal teloblasts (Dill and Seaver, 2008). Morphologically distinct teloblasts have been
described in the larva of *Scoloplos armiger* (Anderson, 1959), but there is no molecular or
functional data to confirm their identity.

The question of the origin of the cellular material for segments is crucial in the
evolutionary analysis of segmentation (Minelli, 2005). At least in nereidids, it can be
confidently stated that larval segments arise from widespread proliferation and migration of
cells in the hyposphere, leading to its convergent extension (Demilly et al., 2013; Denes et
al., 2007; Steinmetz et al., 2007). These processes could explain the separation of the
originally united lateral longitudinal domains of *engrailed* expression (Fig. 1, 6). In contrast,
postlarval segments derive from the proliferation of subterminal growth zone cells and the
cellular material of each new segment anlage. In clitellate annelids, these differences do not
exist, as all segments are formed by local cell proliferation. By analogy with clitellates, some
authors have suggested that the source of larval segments in non-clitellate annelids is the
growth zone of the trochophore (Anderson, 1973; Shimizu and Nakamoto, 2001), but this
speculation is inconsistent with modern observations.

Inductive interactions between the ectoderm and mesoderm also play a critical role in
segment specification. In clitellate embryos, the mesoderm governs segment morphogenesis
and the axial identity of the ectoderm (Blair, 1982; Shimizu et al., 2001). Whether such
interactions are present in non-clitellate annelids remains unknown, but the metameric pattern
of *Avi-twist* expression in mesodermal bands (Kozin et al., 2016) forms even earlier than *Avi-*
en expression in the ectoderm (Fig. 1C). This is strikingly different from arthropods, where
the ectoderm determines mesoderm segmentation (Azpiazu et al., 1996; Hannibal et al.,
2012). To assess the autonomy of specification, it would be necessary to limit external
signaling influences on the region undergoing segmentation, a task that could be
accomplished through explantation or complete inhibition of signaling interactions.

Several studies on the inhibition of intracellular processes have revealed segmentation
abnormalities. Notably, interesting results were obtained in *A. virens* embryos treated with
antibiotics (actinomycin D and sibiromycin) during gastrulation and trochophore formation.
These treatments led to the formation of unsegmented larvae instead of segmented

metatrochophores (Dondua, 1975). This phenotype appears incompatible with a lineage-
driven mode and may thus support the inductive specification of larval segments. In line with
this, inhibition of Hedgehog signaling in *P. dumerilii* (Dray et al., 2010) also led to a
reduction in ectodermal striped expression of *hedgehog*, *engrailed*, *wnt1*, and *lbx*. Notably,
the reduction in segmental grooves in larval segments was more pronounced than in
postlarval segments. The authors concluded that Hedgehog is not necessary to establish early
segment patterns but is required to maintain them. Additionally, segmentation of regenerating
posterior body ends was impaired due to modulation of Wnt (Niwa et al., 2013; Ribeiro and
Aguado, 2021) and FGF signaling pathways (Shalaeva et al., 2021; Shalaeva and Kozin,
2024). In contrast, Delta/Notch signaling does not appear to be involved in segmental pattern
formation during either the larval or postlarval periods (Gazave et al., 2017; Rivera and
Weisblat, 2009; Thamm and Seaver, 2008).

Sequential segmentation and the growth zone organization

The organization of the growth zone (also referred to as the segment addition zone, or
SAZ, in arthropods) is central to understanding the origin of segmentation in bilaterians.
Among annelids, the growth zone has been well studied in clitellates, particularly in leeches,
where it consists of several large stem cells (teloblasts) that give rise to all body segments
(Weisblat and Kuo, 2014; Zattara and Weisblat, 2020). However, based on the limited
distribution of teloblasts and the existing hypotheses regarding the evolution of annelid
ontogeny, we propose that a teloblastic SAZ is not the ancestral trait of annelids (Kuo, 2017;
Minelli, 2005; Scholtz, 2002). Instead, it is likely a secondary gain, meaning clitellates are
not the most appropriate model for comparative analysis.
Currently, we lack a clear understanding of how the growth zone functions in non-
clitellate annelids. Most of the available data comes from nereidid polychaetes. Studies on *P.*
dumerilii have proposed the existence of a teloblastic growth zone, although teloblasts do not
differ in size from other cells (Balavoine, 2014; Gazave et al., 2013). However, there are no
signs of stereotypical division patterns in this zone, except for the longitudinal orientation of
mitoses in clusters of ectodermal growth zone cells during regeneration in *Perinereis nuntia*
(Niwa et al., 2013). The diffuse model of SAZ, characteristic of arthropods and vertebrates,
which involves periodic gene expression (e.g., *hairy* in vertebrates or pair-rule genes in
arthropods) (Bénazéraf and Pourquié, 2013; Williams and Nagy, 2017), is not applicable to
nereidids. The diffuse SAZ model designates a wide area of unsegmented tissue along the AP
axis, where the posterior shift of the wavefront results in a gradual decrease in differentiation
potency, leading to segment specification. This pattern is referred to as boundary-driven
segmentation (Weisblat and Kuo, 2014). However, no oscillations in gene expression have
been detected in annelids to date.
One of the most critical mechanisms for segment addition is cell proliferation. In
arthropods and vertebrate embryos, discrete regions of cell proliferation are observed (Nagy
and Williams, 2020; Venters et al., 2008). In particular, cell divisions in short germ-band
arthropods occur in the posterior part of the growth zone and in the last specified metamer
(Fig. 8 A-C). Annelids, in contrast, exhibit ubiquitous proliferation in the hyposphere of the
trochophore (Demilly et al., 2013) and in a continuous, posterior subterminal region,

extending from the compact growth zone to the newly formed segments (Fig. 4, Fig. 8 A'-C').
This pattern of cell division appears to be unique to the segment-producing tissues in
annelids, both clitellate (Bissen and Weisblat, 1989) and non-clitellate. Consequently, the
discrete pattern of cell divisions in a diffuse SAZ is incompatible with the mechanisms
forming both larval and postlarval segments in annelids.

664
Fig. 8. Comparison of sequential segment formation between Arthropods with the diffuse SAZ (upper row) and
A. virens (lower row). Dashed area — specifying metamer (parasegment). Blue areas — *wg/wnt1* expression,
red areas — *engrailed* expression. Charts below each diagram represent EdU incorporation level along the AP
axis (Nagy and Williams, 2020). (A) Cell accumulation in the anterior growth zone. (B) Gene expression
oscillations (pair-rule genes or Notch-signaling components) promote parasegment specification. (C)
Determination of parasegment borders (segment polarity genes expression). (A') Expansion of the posterior *Av-*
en domain indicates definition of the new segment's anterior border. (B') Cell divisions gradually produce more
posterior territories of the specifying segment. (C') *Avi-wnt1* expression in front of segment-pygidium furrow
indicates determination of the whole new segment anlage.

Our results suggest several key features of sequential segment addition in nereidid
annelids (Fig. 8). We distinguish three stages of postlarval segment development: elongation
of the AP axis, metamer specification and separation (which ultimately determines segmental

borders). Unlike arthropods, nereidids exhibit *engrailed* expression in the growth zone (Fig. 8A'), which resolves into two rows by the nectochaete stage (Fig. 8B'). According to our data and previous studies (Niwa et al., 2013), *wnt1* expression at the posterior border of the developing metamer represents the final step in the segmentation program (Fig. 8C'). Thus, axial elongation and segment specification occur simultaneously in nereidids. This contrasts with parasegment formation in arthropods, where these processes are strictly sequential (Williams and Nagy, 2017). These differences cast doubt on the homology of segments among bilaterians, as proposed by numerous authors (Balavoine, 2014; Balavoine and Adoutte, 2003; Couso, 2009; de Rosa et al., 2005; Malakhov et al., 2019; Prud'homme et al., 2003; Saudemont et al., 2008; Scholtz, 2002; Shcherbakov, 2023). Our data suggest that the involvement of *engrailed* and *wnt1* in segmentation may be independent in annelids and other taxa (Bleidorn et al., 2015; Chipman, 2010; Ferrier, 2012; Kairov and Kozin, 2023; Seaver, 2022, 2003; Zattara and Weisblat, 2020), which is also supported by *engrailed* patterns in diverse metazoans (Vellutini and Hejnol, 2016). This scenario is further reinforced by the fact that segmental grooves in arthropods are formed posteriorly from the *engrailed* expression band, while in nereidid annelids, they are formed anteriorly from the *engrailed* band and posteriorly from the *wnt1* band (Fig. 8). Annelid segments have been homologized with parasegments in arthropods (Janssen et al., 2010; Prud'homme et al., 2003), since parasegmental furrows are formed between the *wnt1* and *engrailed* expression bands. However, these parasegmental furrows may have convergent evolutionary origins and are not considered to be a conserved feature across arthropods (Janssen et al., 2022).

Thus, the model of sequential segmentation in nereidids involves the gradual formation of a segment along the AP axis (Kairov and Kozin, 2023). The specification of a new segment begins as *engrailed* expression in the growth zone expands through cell divisions. This widened ring of *engrailed*-positive cells presumably give rise to the anterior border of the new metamer and to the maintaining growth zone posteriorly. Subsequent proliferation splits the *engrailed* expression bands, which is morphologically manifested by pygidium growth. Segment specification is complete when the *wnt1* expression band appears, marking the posterior boundary of the segment.

The previous model of gradual segment specification was first proposed for regenerating *P. nuntia* (Niwa et al., 2013). In this model, the growth zone is not a population of bona fide multipotent stem cells but rather a region where the anterior-most pygidial cells undergo reiterative transdifferentiation. However, this does not align with the consistent *engrailed* expression at the anterior border of the pygidium, which is observed not only in nereidids but also in *C. teleta* and *Pristina leidyi* (Bely and Wray, 2001; Seaver and Kaneshige, 2006). The model proposed by Niwa and colleagues (2013) suggests that Wnt signaling induces segment precursors, as Wnt hyperactivation alters the size and number of emerging segments, which is proved in *P. nuntia* and *Sillys malaquini* (Niwa et al., 2013; Ribeiro and Aguado, 2021). The *Avi-wnt* expression patterns in *A. virens* are consistent with the hypothesis that Wnt signaling plays a crucial role in the patterning of both larval and postlarval segments.

Further interpretation of segmentation traits in nereidids leads us to hypothesize that *Avi-wnt* expression in the proctodaeum/hindgut is required for the induction and maintenance of growth zone function, whereas Wnt activity in specified segments is involved in their

polarization and in regulating the pace of segment addition, as reflected in the size of the newly forming segment. The terminal posterior Wnt signaling center is highly conserved across bilaterians (Loh et al., 2016; Martin and Kimelman, 2009; Petersen and Reddien, 2009), and it plays a critical role in axial elongation in both segmented (Bolognesi et al., 2008; Chesebro et al., 2013; Dunty et al., 2008; Shimizu et al., 2005) and non-segmented animals (Fritzenwanker et al., 2019). Posterior Wnt activity has been confirmed in all annelids studied (Cho et al., 2010; Janssen et al., 2010; Kozin et al., 2019; Niwa et al., 2013; Nyberg et al., 2012; Prud'homme et al., 2003; Pruitt et al., 2014; Ribeiro and Aguado, 2021; Seaver et al., 2001; Seaver and Kaneshige, 2006), whereas Wnt expression in stripes along segmental borders has only been reported in nereidids and *C. teleta* (Cho et al., 2010). To clarify how segmented body plans evolved within annelids, future studies should primarily focus on Wnt-mediated segmentation mechanisms across different phylogenetic lineages of non-clitellate annelids.

Acknowledgements

This research was funded by the RSF grant 23-74-10046, https://rscf.ru/en/project/23-74-10046/. We are grateful to the research resource centers "Microscopy and Microanalysis", "Molecular and Cell Technologies", "Chromas", "Culture Collection of Microorganisms", and the Marine Biological Station of St. Petersburg State University for their technical support. We thank Roman P. Kostyuchenko for plasmids with fragments of the cloned cDNA of *Avi-wnt1* and *Av-en*.

References

1. Anderson, D.T., 1973. Embryology and phylogeny in annelids and arthropods, 1st ed. ed, International series of monographs in pure and applied biology. Division: zoology. Pergamon Press, Oxford, New York.
2. Anderson, D.T., 1959. The Embryology of the Polychaete Scoloplos armiger. Q. J. Microsc. Sci. 100, 89–166.
3. Azpiazu, N., Lawrence, P.A., Vincent, J.P., Frasch, M., 1996. Segmentation and specification of the Drosophila mesoderm. Genes Dev. 10, 3183–3194. https://doi.org/10.1101/gad.10.24.3183
4. Balavoine, G., 2014. Segment formation in Annelids: patterns, processes and evolution. Int J Dev Biol 58, 469–483.
5. Balavoine, G., Adoutte, A., 2003. The Segmented Urbilateria: A Testable Scenario. Integr Comp Biol 43, 137–147.
6. Bastin, B.R., Chou, H.-C., Pruitt, M.M., Schneider, S.Q., 2015. Structure, phylogeny, and expression of the frizzled-related gene family in the lophotrochozoan annelid Platynereis dumerilii. EvoDevo 6, 37. https://doi.org/10.1186/s13227-015-0032-4
7. Bastin, B.R., Meha, S.M., Khindurangala, L., Schneider, S.Q., 2023. Cooption of regulatory modules for tektin paralogs during ciliary band formation in a marine annelid larva. Dev. Biol. 503, 95–110. https://doi.org/10.1016/j.ydbio.2023.07.006
8. Bely, A.E., Wray, G.A., 2001. Evolution of regeneration and fission in annelids: insights from engrailed- and orthodenticle-class gene expression. Development 128, 2781–2791. https://doi.org/10.1242/dev.128.14.2781

9. Bénazéraf, B., Pourquié, O., 2013. Formation and Segmentation of the Vertebrate
Body Axis. Annu Rev Cell Dev Biol 29, 1–26.

10. Bissen, S.T., Weisblat, D.A., 1989. The durations and compositions of cell cycles in
embryos of the leech, *Helobdella triserialis*. Development 106, 105–118.
https://doi.org/10.1242/dev.106.1.105

11. Blair, S.S., 1982. Interactions between mesoderm and ectoderm in segment formation
in the embryo of a glossiphoniid leech. Dev. Biol. 89, 389–396.
https://doi.org/10.1016/0012-1606(82)90327-X

12. Bleidorn, C., Helm, C., Weigert, A., Aguado, M.T., 2015. Annelida, in: Evolutionary
Developmental Biology of Invertebrates 2: Lophotrochozoa (Spiralia). Springer
Vienna, Vienna, pp. 193–230.

13. Bolognesi, R., Farzana, L., Fischer, T.D., Brown, S.J., 2008. Multiple Wnt Genes Are
Required for Segmentation in the Short-Germ Embryo of Tribolium castaneum. Curr.
Biol. 18, 1624–1629. https://doi.org/10.1016/j.cub.2008.09.057

14. Byrne, M., Cisternas, P., Elia, L., Relf, B., 2005. Engrailed is expressed in larval
development and in the radial nervous system of Patiriella sea stars. Dev. Genes Evol.
215, 608–617. https://doi.org/10.1007/s00427-005-0018-7

15. Chesebro, J.E., Pueyo, J.I., Couso, J.P., 2013. Interplay between a Wnt-dependent
organiser and the Notch segmentation clock regulates posterior development in
Periplaneta americana. Biol. Open 2, 227–237. https://doi.org/10.1242/bio.20123699

16. Chipman, A.D., 2010. Parallel evolution of segmentation by co-option of ancestral
gene regulatory networks. BioEssays 32, 60–70.
https://doi.org/10.1002/bies.200900130

17. Cho, S.-J., Valles, Y., Giani, V.C., Seaver, E.C., Weisblat, D.A., 2010. Evolutionary
Dynamics of the wnt Gene Family: A Lophotrochozoan Perspective. Mol. Biol. Evol.
27, 1645–1658. https://doi.org/10.1093/molbev/msq052

18. Chou, H.-C., Pruitt, M.M., Bastin, B.R., Schneider, S.Q., 2016. A transcriptional
blueprint for a spiral-cleaving embryo. BMC Genomics 17, 552.
https://doi.org/10.1186/s12864-016-2860-6

19. Condron, B.C., Pate, N.H., Zinn, K., 1994. engrailed Controls Glial/Neuronal Cell
Fate Decisions at the Midline of the Central Nervous System. Neuron 13, 541–554.

20. Constantinou, S.J., Duan, N., Nagy, L.M., Chipman, A.D., Williams, T.A., 2020.
Elongation during segmentation shows axial variability, low mitotic rates, and
synchronized cell cycle domains in the crustacean, Thamnocephalus platyurus.
EvoDevo 11, 1. https://doi.org/10.1186/s13227-020-0147-0

21. Couso, J.P., 2009. Segmentation, metamerism and the Cambrian explosion. Int. J.
Dev. Biol. 53, 1305–1316. https://doi.org/10.1387/ijdb.072425jc

22. de Rosa, R., Prud'homme, B., Balavoine, G., 2005. caudal and even-skipped in the
annelid Platynereis dumerilii and the ancestry of posterior growth. Evol. Dev. 7, 574–
587. https://doi.org/10.1111/j.1525-142X.2005.05061.x

23. Demilly, A., Steinmetz, P., Gazave, E., Marchand, L., Vervoort, M., 2013.
Involvement of the Wnt/β-catenin pathway in neurectoderm architecture in
Platynereis dumerilii. Nat. Commun. 4, 1915. https://doi.org/10.1038/ncomms2915

24. Denes, A.S., Jékely, G., Steinmetz, P.R.H., Raible, F., Snyman, H., Prud'homme, B.,
Ferrier, D.E.K., Balavoine, G., Arendt, D., 2007. Molecular Architecture of Annelid
Nerve Cord Supports Common Origin of Nervous System Centralization in Bilateria.
Cell 129, 277–288. https://doi.org/10.1016/j.cell.2007.02.040

25. Dill, K.K., Seaver, E.C., 2008. Vasa and nanos are coexpressed in somatic and germ
line tissue from early embryonic cleavage stages through adulthood in the polychaete

Capitella sp. I. Dev. Genes Evol. 218, 453–463. https://doi.org/10.1007/s00427-008-0236-x

26. Dondua, A.K., 1975. Effect of actinomycin D and sibiromycin on the embryonic and larval development of Nereis virens (Sars.). Ontogenez 6, 475–484.

27. Dray, N., Tessmar-Raible, K., Le Gouar, M., Vibert, L., Christodoulou, F., Schipany, K., Guillou, A., Zantke, J., Snyman, H., Béhague, J., Vervoort, M., Arendt, D., Balavoine, G., 2010. Hedgehog Signaling Regulates Segment Formation in the Annelid *Platynereis*. Science 329, 339–342. https://doi.org/10.1126/science.1188913

28. Dunty, W.C., Biris, K.K., Chalamalasetty, R.B., Taketo, M.M., Lewandoski, M., Yamaguchi, T.P., 2008. Wnt3a/β-catenin signaling controls posterior body development by coordinating mesoderm formation and segmentation. Development 135, 85–94. https://doi.org/10.1242/dev.009266

29. Eriksson, B.J., Tait, N.N., Budd, G.E., Akam, M., 2009. The involvement of engrailed and wingless during segmentation in the onychophoran Euperipatoides kanangrensis (Peripatopsidae: Onychophora) (Reid 1996). Dev. Genes Evol. 219, 249–264. https://doi.org/10.1007/s00427-009-0287-7

30. Ferrier, D.E.K., 2012. Evolutionary crossroads in developmental biology: annelids. Development 139, 2643–2653. https://doi.org/10.1242/dev.074724

31. Fischer, A.H., Henrich, T., Arendt, D., 2010. The normal development of Platynereis dumerilii (Nereididae, Annelida). Front. Zool. 7, 31. https://doi.org/10.1186/1742-9994-7-31

32. Fischer, A.H.L., Arendt, D., 2013. Mesoteloblast-Like Mesodermal Stem Cells in the Polychaete Annelid Platynereis dumerilii (Nereididae). J. Exp. Zoolog. B Mol. Dev. Evol. 320, 94–104. https://doi.org/10.1002/jez.b.22486

33. Fritzenwanker, J.H., Uhlinger, K.R., Gerhart, J., Silva, E., Lowe, C.J., 2019. Untangling posterior growth and segmentation by analyzing mechanisms of axis elongation in hemichordates. Proc. Natl. Acad. Sci. 116, 8403–8408. https://doi.org/10.1073/pnas.1817496116

34. Gabriel, W.N., Goldstein, B., 2007. Segmental expression of Pax3/7 and Engrailed homologs in tardigrade development. Dev. Genes Evol. 217, 421–433. https://doi.org/10.1007/s00427-007-0152-5

35. Gazave, E., Béhague, J., Laplane, L., Guillou, A., Préau, L., Demilly, A., Balavoine, G., Vervoort, M., 2013. Posterior elongation in the annelid Platynereis dumerilii involves stem cells molecularly related to primordial germ cells. Dev. Biol. 382, 246–267. https://doi.org/10.1016/j.ydbio.2013.07.013

36. Gazave, E., Guillou, A., Balavoine, G., 2014. History of a prolific family: the Hes/Hey-related genes of the annelid Platynereis. EvoDevo 5, 29. https://doi.org/10.1186/2041-9139-5-29

37. Gazave, E., Lemaître, Q.I.B., Balavoine, G., 2017. The Notch pathway in the annelid Platynereis: insights into chaetogenesis and neurogenesis processes. Open Biol. 7, 160242. https://doi.org/10.1098/rsob.160242

38. Giangrande, A., Gambi, M.C., 1998. Metamerism and life□style within polychaetes: Morpho□functional aspects and evolutionary implications. Ital. J. Zool. 65, 39–50. https://doi.org/10.1080/11250009809386725

39. Hannibal, R.L., Patel, N.H., 2013. What is a segment? Evodevo 17;4, 35.

40. Hannibal, R.L., Price, A.L., Patel, N.H., 2012. The functional relationship between ectodermal and mesodermal segmentation in the crustacean, Parhyale hawaiensis. Dev. Biol. 361, 427–438. https://doi.org/10.1016/j.ydbio.2011.09.033

41. Hou, X., Wei, M., Li, Q., Zhang, T., Zhou, D., Kong, D., Xie, Y., Qin, Z., Zhang, Z., 2019. Transcriptome Analysis of Larval Segment Formation and Secondary Loss in

the Echiuran Worm Urechis unicinctus. Int. J. Mol. Sci. 20, 1806. https://doi.org/10.3390/ijms20081806

42. Irvine, S., Martindale, M., 1996. Cellular and molecular mechanisms of segmentation in annelids. Semin. Cell Dev. Biol. 7, 593–604. https://doi.org/10.1006/scdb.1996.0073

43. Irvine, S.Q., Seaver, E.C., 2006. Early Annelid Development, A Molecular Perspective, in: Reproductive Biology and Phylogeny of Annelida. CRC Press.

44. Ivanova-Kazas, O.M., 1978. The modern state of the theory of primary heteronomy of the segments (To the centenary of birth of prof. P. P. Ivanov). Zool. Zhurnal 57, 1605–1617.

45. Iwanoff, P.P., 1928. Die entwicklung der larvalsegmente bei den anneliden. Z. Für Morphol. Ökol. Tiere 10, 62–161. https://doi.org/10.1007/BF00419279

46. Janssen, R., Le Gouar, M., Pechmann, M., Poulin, F., Bolognesi, R., Schwager, E.E., Hopfen, C., Colbourne, J.K., Budd, G.E., Brown, S.J., Prpic, N.-M., Kosiol, C., Vervoort, M., Damen, W.G., Balavoine, G., McGregor, A.P., 2010. Conservation, loss, and redeployment of Wnt ligands in protostomes: implications for understanding the evolution of segment formation. BMC Evol. Biol. 10, 374. https://doi.org/10.1186/1471-2148-10-374

47. Janssen, R., Turetzek, N., Pechmann, M., 2022. Lack of evidence for conserved parasegmental grooves in arthropods. Dev. Genes Evol. 232, 27–37. https://doi.org/10.1007/s00427-022-00684-5

48. Jekely, G., Arendt, D., 2007. Cellular resolution expression profiling using confocal detection of NBT/BCIP precipitate by reflection microscopy. Biotechniques 42, 751–755.

49. Kairov, A.I., Kozin, V.V., 2023. Expression of the engrailed Homologue in Larvae and Juveniles of the Annelid Alitta virens Characterizes the Formation of Segments from the Growth Zone. Russ. J. Dev. Biol. 54, 177–185. https://doi.org/10.1134/S1062360423030037

50. Kostyuchenko, R.P., 2022. Nanos Is Expressed in Somatic and Germline Tissue during Larval and Post-Larval Development of the Annelid Alitta virens. Genes 13, 270. https://doi.org/10.3390/genes13020270

51. Kozin, V.V., Borisenko, I.E., Kostyuchenko, R.P., 2019. Establishment of the Axial Polarity and Cell Fate in Metazoa via Canonical Wnt Signaling: New Insights from Sponges and Annelids. Biol. Bull. 46, 14–25. https://doi.org/10.1134/S1062359019010035

52. Kozin, V.V., Filimonova, D.A., Kupriashova, E.E., Kostyuchenko, R.P., 2016. Mesoderm patterning and morphogenesis in the polychaete Alitta virens (Spiralia, Annelida): Expression of mesodermal markers Twist, Mox, Evx and functional role for MAP kinase signaling. Mech. Dev. 140, 1–11. https://doi.org/10.1016/j.mod.2016.03.003

53. Kozin, V.V., Kostyuchenko, R.P., 2015. Vasa, PL10, and Piwi gene expression during caudal regeneration of the polychaete annelid Alitta virens. Dev. Genes Evol. 225, 129–138. https://doi.org/10.1007/s00427-015-0496-1

54. Kulakova, M.A., Cook, C.E., Andreeva, T.F., 2008. ParaHox gene expression in larval and postlarval development of the polychaete Nereis virens(Annelida, Lophotrochozoa). BMC Dev. Biol. 8, 61. https://doi.org/10.1186/1471-213X-8-61

55. Kuo, D.-H., 2017. The polychaete-to-clitellate transition: An EvoDevo perspective. Dev. Biol. 427, 230–240. https://doi.org/10.1016/j.ydbio.2017.01.016

56. Lans, D., Wedeen, C.J., Weisblat, D.A., 1993. Cell lineage analysis of the expression of an *engrailed* homolog in leech embryos. Development 117, 857–871. https://doi.org/10.1242/dev.117.3.857

57. Lim, J., Choe, C.P., 2020. Functional analysis of engrailed in Tribolium segmentation. Mech. Dev. 161, 103594. https://doi.org/10.1016/j.mod.2019.103594

58. Loh, K.M., van Amerongen, R., Nusse, R., 2016. Generating Cellular Diversity and Spatial Form: Wnt Signaling and the Evolution of Multicellular Animals. Dev. Cell 38, 643–655. https://doi.org/10.1016/j.devcel.2016.08.011

59. Malakhov, V.V., Bogomolova, E.V., Kuzmina, T.V., Temereva, E.N., 2019. Evolution of Metazoan Life Cycles and the Origin of Pelagic Larvae. Russ. J. Dev. Biol. 50, 303–316. https://doi.org/10.1134/S1062360419060043

60. Martin, B.L., Kimelman, D., 2009. Wnt Signaling and the Evolution of Embryonic Posterior Development. Curr. Biol. 19, R215–R219. https://doi.org/10.1016/j.cub.2009.01.052

61. Martín-Zamora, F.M., Liang, Y., Guynes, K., Carrillo-Baltodano, A.M., Davies, B.E., Donnellan, R.D., Tan, Y., Moggioli, G., Seudre, O., Tran, M., Mortimer, K., Luscombe, N.M., Hejnol, A., Marlétaz, F., Martín-Durán, J.M., 2023. Annelid functional genomics reveal the origins of bilaterian life cycles. Nature 615, 105–110. https://doi.org/10.1038/s41586-022-05636-7

62. Minelli, A., 2005. A morphologist's perspective on terminal growth and segmentation. Evol. Dev. 7, 568–573. https://doi.org/10.1111/j.1525-142X.2005.05060.x

63. Minelli, A., Fusco, G., 2004. Evo-devo perspectives on segmentation: model organisms, and beyond. Trends Ecol. Evol. 19, 423–429. https://doi.org/10.1016/j.tree.2004.06.007

64. Mundaca-Escobar, M., Cepeda, R.E., Sarrazin, A.F., 2022. The organizing role of Wnt signaling pathway during arthropod posterior growth. Front. Cell Dev. Biol. 10, 944673. https://doi.org/10.3389/fcell.2022.944673

65. Nagy, L.M., Williams, T.A., 2020. Cell Division, Movement, and Synchronization in Arthropod Segmentation, in: Cellular Processes in Segmentation. Taylor & Francis Group, pp. 39–70.

66. Nielsen, C., 2004. Trochophora Larvae: Cell-Lineages, Ciliary Bands,and Body Regions. 1. Annelida and Mollusca. J. Exp. Zool. MOL DEV EVOL 302, 1–102.

67. Niwa, N., Akimoto-Kato, A., Sakuma, M., Kuraku, S., Hayashi, S., 2013. Homeogenetic inductive mechanism of segmentation in polychaete tail regeneration. Dev. Biol. 381, 460–470. https://doi.org/10.1016/j.ydbio.2013.04.010

68. Nyberg, K.G., Conte, M.A., Kostyun, J.L., Forde, A., Bely, A.E., 2012. Transcriptome characterization via 454 pyrosequencing of the annelid Pristina leidyi, an emerging model for studying the evolution of regeneration. BMC Genomics 13, 287. https://doi.org/10.1186/1471-2164-13-287

69. Omi, M., Nakamura, H., 2015. Engrailed and tectum development. Dev. Growth Differ. 57, 135–145. https://doi.org/10.1111/dgd.12197

70. Özpolat, B.D., Handberg-Thorsager, M., Vervoort, M., Balavoine, G., 2017. Cell lineage and cell cycling analyses of the 4d micromere using live imaging in the marine annelid Platynereis dumerilii. eLife 6, e30463. https://doi.org/10.7554/eLife.30463

71. Patel, N.H., Martin-Blanco, E., Coleman, K.G., Poole, S.J., Ellis, M.C., Kornberg, B., Goodman, C.S., 1989. Expression of engrailed Proteins in Arthropods, Annelids, and Chordates. Cell 58, 955–968.

72. Paulus, T., Müller, M.C.M., 2006. Cell proliferation dynamics and morphological differentiation during regeneration in *Dorvillea bermudensis* (Polychaeta, Dorvilleidae). J. Morphol. 267, 393–403. https://doi.org/10.1002/jmor.10280

73. Petersen, C.P., Reddien, P.W., 2009. Wnt Signaling and the Polarity of the Primary Body Axis. Cell 139, 1056–1068. https://doi.org/10.1016/j.cell.2009.11.035

74. Prud'homme, B., de Rosa, R., Arendt, D., Julien, J.-F., Pajaziti, R., Dorresteijn, A.W.C., Adoutte, A., Wittbrodt, J., Balavoine, G., 2003. Arthropod-like Expression Patterns of engrailed and wingless in the Annelid Platynereis dumerilii Suggest a Role in Segment Formation. Curr. Biol. 13, 1876–1881. https://doi.org/10.1016/j.cub.2003.10.006

75. Pruitt, M.M., Letcher, E.J., Chou, H.-C., Bastin, B.R., Schneider, S.Q., 2014. Expression of the wnt gene complement in a spiral-cleaving embryo and trochophore larva. Int. J. Dev. Biol. 58, 563–573. https://doi.org/10.1387/ijdb.140084ss

76. Rebscher, N., Zelada-González, F., Banisch, T.U., Raible, F., Arendt, D., 2007. Vasa unveils a common origin of germ cells and of somatic stem cells from the posterior growth zone in the polychaete Platynereis dumerilii. Dev. Biol. 306, 599–611. https://doi.org/10.1016/j.ydbio.2007.03.521

77. Ribeiro, R.P., Aguado, M.T., 2021. Effects of GSK3β inhibition in the regeneration of Syllis malaquini (Syllidae, Annelida). Dev. Genes Evol. 231, 141–146. https://doi.org/10.1007/s00427-021-00681-0

78. Ribeiro, R.P., Egger, B., Ponz-Segrelles, G., Aguado, M.T., 2021. Cellular proliferation dynamics during regeneration in Syllis malaquini (Syllidae, Annelida). Front. Zool. 18, 27. https://doi.org/10.1186/s12983-021-00396-y

79. Rivera, A.S., Weisblat, D.A., 2009. And Lophotrochozoa makes three: Notch/Hes signaling in annelid segmentation. Dev Genes Evol 219, 37–43.

80. Saudemont, A., Dray, N., Hudry, B., Le Gouar, M., Vervoort, M., Balavoine, G., 2008. Complementary striped expression patterns of NK homeobox genes during segment formation in the annelid Platynereis. Dev. Biol. 317, 430–443. https://doi.org/10.1016/j.ydbio.2008.02.013

81. Scholtz, G., 2002. The Articulata hypothesis□? or what is a segment? Org. Divers. Evol. 2, 197–215. https://doi.org/10.1078/1439-6092-00046

82. Schroeder, P., Hermans, 1975. Annelida: Polychaeta. In "Reproduction of Marine Invertebrates." Academic Press, New York.

83. Seaver, E., 2022. Chapter Fourteen: Sifting through the mud: A tale of building the annelid Capitella teleta for EvoDevo studies, in: Emerging Model Systems in Developmental Biology. Curr Top Dev Biol, pp. 401–432.

84. Seaver, E.C., 2003. Segmentation: mono- or polyphyletic? Int J Dev Biol 47, 583–595.

85. Seaver, E.C., Kaneshige, L.M., 2006. Expression of 'segmentation' genes during larval and juvenile development in the polychaetes Capitella sp. I and H. elegans. Dev. Biol. 289, 179–194. https://doi.org/10.1016/j.ydbio.2005.10.025

86. Seaver, E.C., Paulson, D.A., Irvine, S.Q., Martindale, M.Q., 2001. The Spatial and Temporal Expression of Ch-en, the engrailed Gene in the Polychaete Chaetopterus, Does Not Support a Role in Body Axis Segmentation. Dev. Biol. 236, 195–209. https://doi.org/10.1006/dbio.2001.0309

87. Seaver, E.C., Shankland, M., 2001. Establishment of segment polarity in the ectoderm of the leech Helobdella. Development 128, 1629–1641.

88. Seaver, E.C., Thamm, K., Hill, S.D., 2005. Growth patterns during segmentation in the two polychaete annelids, Capitella sp. I and Hydroides elegans: comparisons at distinct life history stages. Evol. Dev. 7, 312–326.

89. Shalaeva, A.Y., Kostyuchenko, R.P., Kozin, V.V., 2021. Structural and Functional Characterization of the FGF Signaling Pathway in Regeneration of the Polychaete Worm Alitta virens (Annelida, Errantia). Genes 12, 788. https://doi.org/10.3390/genes12060788

90. Shalaeva, A.Y., Kozin, V.V., 2024. Annelid perspectives into the role of FGF signaling in caudal regeneration. bioRxiv. https://doi.org/10.1101/2024.11.26.625069

91. Shalaeva, A.Y., Kozin, V.V., 2023. Cell Proliferation Indices in Regenerating Alitta virens (Annelida, Errantia). Cells 12, 1354. https://doi.org/10.3390/cells12101354

92. Shcherbakov, D.E., 2023. The Main Line of the Evolution of Articulata—From Polychaetes to Insects. Paleontol. J. 57, 1286–1297. https://doi.org/10.1134/S0031030123110126

93. Shimizu, T., Bae, Y.-K., Muraoka, O., Hibi, M., 2005. Interaction of Wnt and caudal-related genes in zebrafish posterior body formation. Dev. Biol. 125–141.

94. Shimizu, T., Kitamura, K., Arai, A., Nakamoto, A., 2001. Pattern formation in embryos of the oligochaete annelid Tubifex: cellular basis for segmentation and specification of segmental identity. Hydrobiologia 463, 123–131. https://doi.org/10.1023/A:1013147523271

95. Shimizu, T., Nakamoto, A., 2001. Segmentation in Annelids: Cellular and Molecular Basis for Metameric Body Plan. Zoolog. Sci. 18, 285–298. https://doi.org/10.2108/zsj.18.285

96. Simsek, M.F., Özbudak, E.M., 2023. A design logic for sequential segmentation across organisms. FEBS J. 290, 5086–5093. https://doi.org/10.1111/febs.16899

97. Starunov, V.V., Dray, N., Belikova, E.V., Kerner, P., Vervoort, M., Balavoine, G., 2015. A metameric origin for the annelid pygidium? BMC Evol. Biol. 15, 25. https://doi.org/10.1186/s12862-015-0299-z

98. Steinmetz, P.R.H., Kostyuchenko, R.P., Fischer, A., Arendt, D., 2011. The segmental pattern of otx, gbx, and Hox genes in the annelid Platynereis dumerilii. Evol. Dev. 13, 72–79. https://doi.org/10.1111/j.1525-142X.2010.00457.x

99. Steinmetz, P.R.H., Zelada-Gonzáles, F., Burgtorf, C., Wittbrodt, J., Arendt, D., 2007. Polychaete trunk neuroectoderm converges and extends by mediolateral cell intercalation. Proc. Natl. Acad. Sci. 104, 2727–2732. https://doi.org/10.1073/pnas.0606589104

100. Sveshnikov, V., 1978. Morfologiya lichinok polikhet (Morphology of Polychaete Larvae). Nauka, Moscow.

101. Thamm, K., Seaver, E.C., 2008. Notch signaling during larval and juvenile development in the polychaete annelid Capitella sp. I. Dev. Biol. 320, 304–318. https://doi.org/10.1016/j.ydbio.2008.04.015

102. Vellutini, B.C., Hejnol, A., 2016. Expression of segment polarity genes in brachiopods supports a non-segmental ancestral role of engrailed for bilaterians. Sci. Rep. 6, 32387. https://doi.org/10.1038/srep32387

103. Venters, S.J., Hultner, M.L., Ordahl, C.P., 2008. Somite cell cycle analysis using somite-staging to measure intrinsic developmental time. Dev. Dyn. 237, 377–392. https://doi.org/10.1002/dvdy.21424

104. Weisblat, D.A., Kuo, D.-H., 2014. Developmental biology of the leech Helobdella. Int. J. Dev. Biol. 58, 429–443. https://doi.org/10.1387/ijdb.140132dw

105. Weisblat, D.A., Winchell, C.J., 2020. Segmentation in Leeches, in: Cellular Processes in Segmentation. Taylor & Francis Group, pp. 153–181.

106. Williams, T.A., Nagy, L.M., 2017. Linking gene regulation to cell behaviors in the posterior growth zone of sequentially segmenting arthropods. Arthropod Struct. Dev. 46, 380–394. https://doi.org/10.1016/j.asd.2016.10.003

107.   Wilson, E.B., 1892. The cell-lineage of Nereis. A contribution to the cytogeny
of the annelid body. J. Morphol. 6, 361–480.
https://doi.org/10.1002/jmor.1050060301
108.   Zattara, E.E., Bely, A.E., 2013. Investment Choices in Post☐Embryonic
Development: Quantifying Interactions Among Growth, Regeneration, and Asexual
Reproduction in the Annelid Pristina leidyi. J Exp Zool Mol Dev Evol 320, 471–488.
109.   Zattara, E.E., Weisblat, D.A., 2020. Cellular and Molecular Mechanisms of
Segmentation in Annelida An Open Question, in: Cellular Processes in Segmentation.
Taylor & Francis Group, pp. 71–97.
1069