Behavior Outpaces Form when Linking Traits to Ecological Responses within Populations: A Meta-Analysis

Thibaut Rota[1,2,3*], Allan Raffard[4,5], Iris Lang[1,6], Quentin Petitjean[1,7], Lisa Jacquin[1,8], Olivier Dézerald[9], Simon Blanchet[10], Andrew P. Hendry[11] and Régis Céréghino[1]

1 Centre de Recherche sur la Biodiversité et l'Environnement (CRBE), Université de Toulouse, CNRS, IRD, Toulouse INP, Université Toulouse 3 – Paul Sabatier (UT3), Toulouse, France
2 Institute of Microbiology, University of Applied Sciences and Arts of Southern Switzerland, Mendrisio, Switzerland
3 Forest Entomology, Swiss Federal Research Institute WSL, Birmensdorf, Switzerland
4 Université catholique de Louvain, Earth and Life Institute, Biodiversity Research Centre, Louvain-la-Neuve, Belgium
5 Univ. Savoie Mont Blanc, INRAE, CARRTEL, 74200 Thonon-les-Bains, France
6 Conservatoire d'espaces naturels d'Occitanie, 26 allées de Mycènes, 34000 Montpellier, France
7 INRAE Nouvelle-Aquitaine Bordeaux Centre, UR EABX, 50 Avenue de Verdun, 33612, Cestas Cedex, Nouvelle-Aquitaine, France
8 Institut Universitaire de France (IUF), Paris, France
9 DECOD, L'Institut Agro, IFREMER, INRAE, 35000 Rennes, France
10 Centre National de la Recherche Scientifique, Station d'Ecologie Théorique et Expérimentale, UAR 2029, 2 route du CNRS, 09200 Moulis, France
11 Redpath Museum and Department of Biology, McGill University, 859 Sherbrooke St. W., H3A 0C4 Montréal, Québec, Canada
* Corresponding author: Dr. Thibaut Rota (thibaut.rota@supsi.ch)

Abstract

Intraspecific trait variability (ITV) is an important aspect of community ecology, but we still have a poor understanding of what drives the magnitude of relationships between ITV and ecological dynamics. In order to guide ecologists interested in unravelling the ecological implications of ITV, we asked whether relationships' magnitudes differ (*i*) between intraspecific levels (among- *vs*. within- populations), (*ii*) among ecological responses spanning levels of biological organization (from individual performances to ecosystem functioning), and (*iii*) among trait categories (morphology, physiology and behavior). We performed a meta-analysis synthesizing a thousand effect sizes from nearly two-hundred studies, encompassing relationships between traits measured among individuals or populations of animals and a broad range of ecological responses. The average effect size was $|r| = 0.24$ [0.20 – 0.27] (95% confidence intervals). Within populations, relationships' magnitudes were higher for ecological

performance (foraging, diet), community and ecosystem responses than for fitness

(reproduction). Relationships were 28% stronger for physiological and behavioral traits than

for morphological traits. Our meta-analysis confirms the central aspect of ITV within

populations of animals to drive ecological dynamics, and emphasizes physiological and

behavioral traits to enhance the integration between the individual component of diversity and

processes at community and ecosystem-levels.

Key words: *Eco-evolutionary dynamics; functional traits; individual-based approach; phenotypic traits; trait-based ecology; intraspecific trait variation (ITV).*

Introduction

The question 'What Do Species Do in Ecosystems?' raised three decades ago by J. H.

Lawton (1994) has been foundational to trait-based ecology (TBE). Trait-based ecologists

revolve around organisms' functional traits (Violle et al., 2007) to describe and predict how

organisms interact altogether and with their environment. Trait-based approaches to community

ecology brought the promise to improve our understanding of the mechanisms driving

community assembly and ecosystem processes, in being complementary to, or even outpace

taxonomic approaches in that regard (Enquist et al. 2015). Traits can be expressed in physical

dimensions (e.g., mass, length, volume, energy, time), and can meaningfully express the fit of

organisms with their environment (Arnold 1983; Violle et al. 2007). Inferences drawn from

TBE are thus, ideally, transferable to other ecosystems and life forms (Funk et al. 2017).

General patterns include allometric scalings between body size and metabolic rate (Kleiber

1932), the leaf economic spectrum (Wright et al., 2004), *r*- and *K*- life history strategies (Pianka

1970). Trait-based approaches do not have borders. Most traits are cheap and easy to measure,

fostering inclusion among researchers from different horizons, as they integrate from individual

to ecosystem levels, and across various ecosystems (Carmona et al. 2016).

Including Intraspecific Trait Variation (ITV) in Trait-Based Ecology (TBE)

Early research in TBE led to rethink community ecology (McGill et al. 2006; Petchey and Gaston 2006). Yet, an important limitation is that studies focused heavily on species' mean trait values, assuming that all conspecifics shared identical trait values (Albert 2015). This assumption was paradoxal, as intraspecific trait variation (henceforth 'ITV') was essential to the development of natural selection (Darwin and Wallace 1858), niche (Elton 1927; Hutchinson 1957), and coexistence theories (Macarthur & Levins, 1967). In fact, ITV is as large as mean trait differences between species (Lecerf and Chauvet 2008; Siefert et al. 2015; Rota et al. 2022), and matters for species coexistence (e.g., Hart et al., 2016), predator–prey (Toscano and Griffen 2014), and consumer–resource interactions (Raffard et al. 2017; Rota et al. 2018). ITV is now recognized to be an important facet of ecology (Bolnick et al. 2011; Violle et al. 2012), provoking discussions on how it affects our understanding of food webs (e.g., behavior; Moran et al., 2017) and community assembly (e.g., metabolic traits; Brandl et al., 2022). However, the inclusion of the intraspecific component remains marginal, with only 4% of TBE studies acknowledging ITV (Green et al., 2022). Intraspecific variation is, however, fundamental in community ecology. For all individualizable organisms, ecological interactions (competition, predation etc.) are occuring among conspecific or heterospecific individuals, whom their traits differ or not. A trait-based approach at the individual level may improve to understand those interactions (Bolnick et al., 2011). Ignoring ITV can lead to inferential bias (Wong and Carmona 2021) and to miss important information about diversity. For instance, not acknowledging for intraspecific variability could distort relationships between species richness and functional diversity (Cianciaruso et al. 2009). On the other hand, acknowledging the processes affecting ITV around species' averages, such as evolution in response to competition regimes, can contribute to a mechanistic understanding of diversity–function relationships (Barabás et al. 2022).

Why Assessing Relationships' Magnitude between ITV and Ecological Dynamics?

Important insights emerged from a renewed interest for ITV. First, ITV links to community assembly patterns (e.g., species richness and diversity) and ecosystem functioning (e.g., productivity or nutrient cycling) as much as mean trait variation between species (Des Roches et al. 2018; Raffard et al. 2019). Second, ITV is changing fast with human-driven disturbances (Alberti et al. 2017), including population loss (Ceballos et al. 2017), erosion of genetic diversity (Exposito-Alonso et al. 2022), and changes in phenotypic variation (Sanderson et al. 2023). There is thus now a form of urgency in studying the primary role of ITV in ecological processes, such as answering how much and when ITV matters. Conservationists can take advantage of this research agenda, to better identifying and monitoring particular facets of ITV with particular implications for ecological functions (Blanchet et al. 2020) and nature's contribution to people (Des Roches et al. 2021).

We need to improve our knowledge about what drives the magnitude of relationships between ITV and ecological dynamics. This would also help understand how evolution, by affecting trait distributions, affects ecological dynamics (i.e., eco-evolutionary dynamics; Pelletier et al., 2009; Hendry, 2017). Most research has focused on natural selection (Kingsolver et al. 2012), heritability (Mousseau and Roff 1987), and evolvability so far (Wheelwright et al. 2014), in providing quantitative estimates on evolutionary processes ('*evo*'). In contrast, the relationships between ITV and ecological responses ('*eco*') did not receive yet a detailed analysis, despite that evolutionary and ecological processes are often congruent (Harmon et al. 2009). Previous studies compared the ecological impacts of populations varying in key traits (e.g., gill raker's morphology in anadromous or landlocked fish populations; Post et al., 2008), but there are still very few studies having focused at both the individual and population levels (Gibert et al. 2015). Hence, it remains unclear if relationships between ITV and ecological responses show similar magnitudes at the population and individual levels. On one hand, we could expect population-

level ITV to show stronger relationships with ecological dynamics than individual-level ITV,

because local adaptation can drive divergence in among-population trait values and trait

matching with local ecological processes (Post et al. 2008). On the other hand, ITV among

individuals has been shown to be similar or even larger than ITV among populations (Messier

et al. 2010; Rota et al. 2024). It thus remains possible that ITV among individuals express

similar or stronger effects on ecological processes than ITV among populations.

Predicting how much traits link to responses across levels of biological organization remains a

challenging issue (Violle et al., 2007; Enquist et al., 2015; Chacón-Labella et al., 2022). There

is still a debate about how the magnitude of relationships between ITV and ecological responses

may vary across levels of biological organization, from individuals to ecosystem processes.

Some predict stronger links with responses at low levels of biological organization (e.g., growth

rate) than with responses at higher levels (e.g., ecosystem functioning; Bailey et al., 2009).

Others showed an opposite pattern (Des Roches et al. 2018; Raffard et al. 2019).

Assessing Traits' Functionality

To generate meaningful insights, we assume more or less implicitly in TBE that the traits used

are functional in linking to the ecological performances or to the fitness of organisms (Arnold

1983), or more generally, to ecological functions (Violle et al. 2007). Hence, the *functionality*

of traits is the matter of discussions, as it influences the choice of traits by trait-based ecologists

(Mlambo 2014; Lefcheck et al. 2015; Dawson et al. 2021). In biological terms, all traits are

functional since they all link to one or several biological functions, making phenotypes

functional living entities (Bertossa 2011). Therefore, the question may not be whether a trait is

functional or not in absolute terms (for different viewpoints across disciplines, see Dawson et

al., 2021 and Sobral, 2021). Without entering into this debate, we propose to investigate *traits'*

functionality that we define as the magnitude of relationships between a trait and a large array

of ecological responses. Asking *how much* traits are functional may help to increase the general

quality of ecological inferences at the intraspecific level in guiding further investigators in their

trait choice.

A good example of high trait functionality is body size. Body size is used in the vast majority

of TBE studies (Green et al. 2022). It spans several orders of magnitude across and within taxa

and integrates many biological and ecological processes with often a great predictive power

(Peters 1993). Nevertheless, functional diversity is multi-dimensional (de Bello et al., 2021),

and it stays poorly known which traits are the most linked to ecological dynamics when we

consider individuals within a same cohort or size class (Brandl et al. 2022). As several authors

argue (Gordon 2011; Toscano et al. 2016; Moran et al. 2017; Brandl et al. 2022; Schleuning et

al. 2023), we need to look beyond body size to improve our understanding of the implications

of ITV (e.g., morphology, physiology, behavior varying among individuals).

Morphology, physiology and behavior are different facets depicting an individual phenotype.

Morphology reflects the abilities of an individual to perform in its environment given

biomechanical constraints (Van Valen 1965; Arnold 1983). For instance, the relative mass of

metabolically-active organs is used as an energetic proxy (Careau et al. 2008). Yet, energetic

currencies such as metabolic rates (i.e., energy loss) should link closer to ecological

performances such as feeding rates (i.e., energy intakes) and growth rates (i.e., energy not

allocated to maintenance nor reproduction; Biro & Stamps, 2008; Careau & Garland, 2012).

Through trophic and ecological interactions, it is expected that metabolic traits are involved in

community assembly patterns and ecosystem functioning (Brandl et al., 2022; fig. 1A). Some

examples suggest that behavioral traits have a close link to ecological interactions (Gordon,

2011; Toscano et al., 2016; for chosen examples from our literature review, see table S1).

Behavioral variation among individuals can strongly affect the outcomes of predator-prey

interactions (McGhee et al. 2013). Behavioral traits such as locomotion activity,

aggressiveness, or boldness are involved in energetic trade-offs (Careau et al., 2008), and hormones regulate metabolism and behavior (Ricklefs & Wikelski, 2002). Physiological and behavioral traits such as those, as well as other energy and matter currencies (e.g., stoichiometry; Elser et al., 2000) – perhaps as opposed to more structural, morphological traits – are therefore expected to exert strong links with ecological performances, fitness, and hence ecological dynamics (Gordon 2011; Moran et al. 2017; Brandl et al. 2022). However, to date, no study synthesized the functionality of morphological, behavioral and physiological traits on various ecological responses.

Questions and Hypotheses

We aimed filling this gap with a meta-analysis exploring the magnitude of relationships between ITV and various ecological responses (fig. 1B) across a large diversity of traits, systems, and species in the animal realm. We tested three main hypotheses:

(*Q1/H1*) Does the strength of relationships between ITV and ecological responses (hereafter 'effect sizes') differ within populations (*i_a*) versus among populations (*i_b*). We expected that *i_a* < *i_b*, because individuals should show stronger phenotypic divergences among- than within-populations. An alternative expectation is that *i_a* = *i_b*, if for instance the populations (*i_a*) have large sizes or are inter-connected populations, making them phenotypically representative of the entire meta-population (*i_b*) (Messier *et al.*, 2010; Rota *et al.*, 2024).

(*Q2/H2*) How do effect sizes vary among ecological responses at different levels of biological organization? We ranked ecological responses from (*ii_a*) proximal responses to the focal organism (e.g., foraging performance, growth, diet, and fitness) to (*ii_b*) distal responses as community assembly (further abbreviated "community") and ecosystem functioning (further abbreviated "ecosystem"; see fig. 1B). We expected that effect sizes would decrease from

proximal to distal responses ($ii_a > ii_b$; as the width of grey arrows in fig. 1B), as the strength of

relationships between traits and responses should dissipate as the number of intermediate

relationships increases (black arrows in fig. 1B; see the algebraic rationale in Bailey et al.,

2009). Effect sizes could on the contrary, be stronger for distal than for proximal ecological

responses ($ii_a < ii_b$) if intermediate relationships' strength shows additivity or multiplicativity

through biological scales (Bailey *et al.*, 2009).

(*Q3/H3*) How do effect sizes vary among trait types? Effect sizes were divided into (iii_m)

morphology (e.g., body shape, relative length of functional body parts), (iii_p) physiology (e.g.,

metabolic rate, hormone levels, elemental stoichiometry), and (iii_b) behavior (e.g., activity,

boldness, sociability). We hypothesized that behavioral and physiological traits would show the

strongest correlations with ecological responses (i.e., $iii_m < iii_p \sim iii_b$). However, it could be

relaxed when morphological differences occur between morphotypes having diverged from

hundreds to thousands of generations, and for which large phenotypic variation can match

ecological differences (e.g.; ecological radiation between limnetic and benthic eco-morphs

within a same lake population; Harmon *et al.*, 2009).

Figure 1: (A) Diagram showing how ITV could relate to various ecological responses in animals. (B) The magnitude of relationships between ITV and ecological responses is expected to change across levels of biological organization. Black arrows depict intermediate relationships propagating ITV effects across biological levels of organization. Following hypothesis 2 (see main text), the magnitude of relationships between ITV and ecological responses could decrease along higher levels of biological organization (Bailey et al. 2009), from responses located on the organism itself to community and ecosystem levels (grey arrows). All the relationships depicted by grey arrows in panel B are assessed in this meta-analysis, excepting the one for populations (*). Examples of relationships are depicted in table S1, with numbers corresponding with fig. 1A.

Methods

Search Strings and Selection Criteria

We searched for thirty keyword combinations in Web of Science ('all databases'), Scopus and Google Scholar (table S2), and obtained a total of 6904 studies. After having sorted out the duplicates manually in Excel (n = 335), we obtained 6569 studies, from which 613 studies followed our scope after having scrutinized titles and abstracts. Our general scope was studies investigating relationships between animal's trait variability at the intraspecific level and various ecological responses. Exclusion criteria at this stage were studies focusing on other groups than animals, investigating other questions than traits – ecological responses, or studies at the interspecific level.

After a full-text examination of those 613 studies, 121 studies met our criteria. We considered empirical studies (*i*) on wild animals, with observations conducted either in the field, microcosms or mesocosms, (*ii*) Studies reporting quantitative relationships or comparison statistics between a phenotypic trait measured on several individuals, or averaged at the level of a group of individuals (i.e., population or colonies of eusocial animals), and a measured ecological response, (*iii*) morphological, physiological, or behavioral traits are categorical or continuous, (i*v*) we did not include ecological responses reflecting population dynamics or population persistence since those are long-term phenomena for which relationships with ITV are difficult to estimate with empirical studies, (*v*) studies focused on the 'identity' of phenotypes (i.e., one phenotype versus another) not 'diversity' (i.e., statistics from treatments including mixtures of phenotypes were excluded), (*vi*) we did not define life history traits as "trait types", since they were categorized as ecological responses (e.g., growth, survival and reproduction), and (*vii*) we did not include body size since we aimed to focus on often neglected traits beyond body size (see justification in the Introduction section). We therefore ensured that

effect sizes were the least influenced by body size (for a detailed description, see Box1 in

Supplementary Materials).

We finally scrutinized the references of these 121 papers and added 49 additional studies

meeting our selection criteria. We further added 22 studies from our personal libraries (fig. S1).

Studies reporting undetailed statistics (e.g., only *P*-values) or for which notes of concerns have

been recently published were excluded from the final dataset. We kept the effect sizes (*r* values)

from Bolnick & Paull (2009) despite they retracted their paper, since the error was an incorrect

calculation of the *P*-values, not the *r* coefficients, hypotheses, nor methods.

237   *Intraspecific Level, Trait Types, Ecological Responses, and Covariates*

We noted the intraspecific level from which the relationships belonged (i.e., among individuals,

among populations, among colonies of eusocial animals). We considered the following seven

types of ecological response: '*Foraging efficiency*' (resource consumption or feeding rates);

'*trophic niche*' (trophic niche position, trophic level, and degree of specialization inferred from

either stable isotopes or diet assessments); '*growth*' (size-standardized growth rates over time

based on body length or body mass); '*survival'* (survival rates obtained from capture-mark-

release-recapture in the wild or from controlled experiments); *'reproduction'* (number of mates,

egg numbers, clutch sizes, or sibling survival); '*community*' (abundances, biomasses, diversity

and structures of communities of organisms of at least two species interacting with the focal

phenotypes); '*ecosystem*' (standing stocks or dynamics of basal resources such as primary

producers or detritus, as well as energy and matter cycling as C and N cycling). We then

categorized traits into morphology, physiology, and behavior. Morphological traits included

body shape, trophic apparatus shape, and linear morphometrics on specific features, locomotion

traits, coloration, body condition, reproductive organ relative size, brain relative size, and

morphotypes. Physiological traits were either assimilation, energy reserves, excretion,

metabolic traits, elemental stoichiometry, hormone levels, immunity, or physiological syndromes. Behavioral traits reflected differences in aggressiveness, exploration, boldness, sociability, activity or cognitive abilities. Finally, we considered the trophic level (i.e., primary consumers feeding on basal resources or predators feeding on other animals), the ecosystem type (i.e., freshwater and/or marine or terrestrial), the methodological approach (i.e., field observations, mesocosm, or microcosms), and the year of publication as covariates.

Calculation of Effect Sizes

We collected correlation coefficients (*r*) or statistics (e.g., *F* or *t* statistics) corresponding to each trait-to-ecological response relationship. The original estimates were transformed to correlation coefficients *r* and then to Fischer's *Z*-transformed correlation coefficients *Zr,* following established procedures (Nakagawa & Cuthill, 2007; see table S3). Since we aimed to quantify the magnitude of ITV – ecological response relationships, and to perform comparisons among trait and ecological response categories, the direction of the relationship was not meaningful here. We used absolute values of correlation coefficients |*Zr*| in further analyses (Bailey et al. 2009). For each effect size |*Zr*|, we computed a sampling variance (*vi*) that we used in our models (detailed below) (Nakagawa and Cuthill 2007; Nakagawa et al. 2022).

Statistical Analyses

We performed statistical analyses using R, version 4.3.2. (R Development Core Team, 2023). We used hierarchical multi-level models accounting for phylogeny (function '*rma.mv()*' in the R package '*metafor*'; Viechtbauer, 2010, see the package's website: https://www.metafor-project.org/doku.php/metafor). These models take random-effects accounting for between- and within-study variance, as well as a matrix of phylogenetic relatedness among species in the error structure. They acknowledge the three usual sources of non-independence of meta-analyses in ecology and evolution (Noble et al., 2017; see also a guide by S. Nakagawa and M.

Lagisz (2016) here: https://environmentalcomputing.net/statistics/meta-analysis/meta-

analysis-3/). In addition, this approach allowed us to account for a variance – covariance matrix

reflecting the non-independent structure of sampling variances (*vi*) in our dataset, and to obtain

cluster-robust point and variance estimates regarding selective reporting bias (see Yang et al.

2024*a*).

We computed phylogenetic relatedness among species following Moran et al. (2021). We

extracted phylogenetic and taxonomic information from the Open Tree of Life

(https://tree.opentreeoflife.org/; Hinchliff et al., 2015). We then resolved any polytomies by

randomisation in the R package '*rotl*' (Michonneau et al. 2016). Branch lengths were estimated

using Grafen's method (Grafen 1989) in the R package '*ape*' (Paradis and Schliep 2019), and

we added a residual error term accounting for the computed phylogenetic correlation matrix in

our meta-analytic models. We accounted for within-study non-independence by adding a

within-study random effect term (nested with *studyID*), attributing an identifier to all effect

sizes (see Noble et al. 2017; Yang et al. 2024*a*). Selective reporting bias and sources of non-

independence inherent to the structure of the data can be accounted for by modelling an

appropriate sampling variance – covariance matrix (*varcov*), and by following a two-step

cluster-robust estimation procedure (Yang et al. 2024*a*). We computed *varcov* with the function

vcalc() of the package '*metafor*', accounting for the basal level of our observations (*effect*

sizeID), the cluster of the model (*studyID*), and a subgrouping variable regarding trait-to-

response combinations (n= 468) (subgroup argument), in setting covariation among subgroups

as ρ=0.6. This *varcov* ensured that the residual errors of the model accounted for the non-

independence inherent to trait-to-response combinations (n= 468). This accounted for the fact

that traits have shared means and errors within those categories but that those means and errors

are expected to show larger differences with other traits and responses measured on other pairs

of individuals, and species, in line with the multi-level structure in the data. For each model

(see fixed-effect structures below), we re-ran the output from the function *rma.mv()* to get a

cluster-robust estimation of effect size estimates and their variance using the function *robust()*

of the package '*metafor*' (i.e., two-step procedure described in fig. 7 in Yang et al. 2024*a*).

Sampling variance (vi) and mean-centered year of publication are indicators of small study

effects and publication time-lag bias, respectively (Nakagawa et al., 2022). We assessed and

accounted for these two bias in adding these terms as fixed-effects in our models, so effect size

estimates are given for vi = zero, and for the average year in our dataset which was 2012.5

(Nakagawa et al., 2022). Using the same error and random-effects terms model structure as

mentioned above, we first estimated the grand-mean effect size with an intercept-only model.

We then setted intraspecific level as a fixed effect (*H1*). Here we tested if effect sizes obtained

among individuals vs. among populations differed, in merging 'populations' and 'colonies of

eusocial animals', to reduce the unbalancing of effect sizes between 'individual-level' (n=952)

and 'population-level' (n=56). We then tested our hypotheses regarding differences among trait

types and ecological response types (*H2* and *H3*) on data at the individual level, with n=952

effect sizes (94% of the dataset), to avoid any confounding effects of the intraspecific level on

those two hypotheses. We also tested H_2 and H_3 using all the dataset, to see if it would change

our conclusions (see table S4). In both cases, we built a model including ecological response

type as a fixed effect and adding trait type as a random effect (*H2*). We then tested for

differences between trait types, in adding ecological response type as a random effect (*H3*).

The conversion of correlation coefficients *Zr* to their absolute values |*Zr*| can cause |*Zr*| to be

biased upward, since when *Zr* is close to zero, any estimation error will increase |*Zr*| (Morrissey

2016). Moreover, since |*Zr*| $\geq$ 0, we cannot directly test its departure from zero. We therefore

generated null effect sizes |*Zr*/$_{null}$ (Raffard et al. 2019). For each 1008 effect size in our dataset,

we randomly simulated a distribution of 1000 *t* statistics based on sample sizes *N* of each effect

size under the null of no correlation. We then converted these *t* statistics into absolute |*Zr*| values

and averaged them for each effect size. Using the same model structure as above, we estimated

equivalent $|Zr|_{null}$ effect sizes. We obtained unbiased $|Zr|$ estimates and their 95% confidence

intervals, and subtracted the estimated $|Zr|_{null}$ from the modelled estimates of $|Zr|$. We back-

transformed $|Zr|_{unbiased}$ to $|r|_{unbiased}$ effect sizes for reporting magnitude estimates in the main text

using a popular statistic in empirical ecology (correlation coefficient).

As developed above, we accounted for selective reporting bias with a cluster-robust estimation.

We also assessed publication bias with a funnel plot on Zr from the intercept-only model

including all the dataset. We corrected effect sizes estimates by their null equivalent, therefore

accounting at our maximum for known sources of bias. We computed total heterogeneity and

tested its significance with a Q-test of total heterogeneity, as we computed between-, within-

study, and phylogenetic heterogeneity (I^2) using Viechtbauer's method (Viechtbauer, 2010; see

R code).

For all hypotheses, we considered statistical significance among groups using comparison tests

and P-values accounting for the multi-level structure of the dataset, with an alpha threshold of

5%. We visualized effect size raw distributions with orchard plots (package '*OrchaRd2.0*';

(Nakagawa et al. 2023), but we modified the function so that lower prediction and confidence

intervals at 95% were bounded to zero, as by definition, absolute effect sizes cannot extend

below zero. Our alignment with the PRISMA guidelines for ecology and evolutionary biology

meta-analysis reporting (O'Dea et al., 2021) is shown in table S5.

Results

347   Our dataset comprised 1008 effect sizes from 188 studies, covering 126 animal species across

348   seven major taxonomic groups and a vast range of ecological contexts. Morphology,

349   physiology, and behavior were represented by 351, 204, and 453 effect sizes, respectively (fig.

2). Effect sizes expressed ecological relationships among individuals (94%), among populations (4%) or among colonies of eusocial animals (2%; fig. 2B).

Heterogeneity and Covariates

Total heterogeneity in our meta-analysis was high (87%; $Q_{df=1005} = 6238.7$; $P < .0001$). Phylogenetic relatedness among species accounted for 4.8% of the total heterogeneity in effect sizes. Most of the heterogeneity was within studies (45.7%), rather than among studies (36.5%). We did not observe any statistical difference among categories of covariates, and accordingly, distributions were undistinguishable between systems (aquatic *vs*. terrestrial; $z = 0.2$; $P = .841$), trophic levels (consumers *vs*. predators; $z = 1.58$; $P = .114$), and methodological approaches (field observations, mesocosms, and microcosms; $z < 1.18$; $P > .239$; fig. S2–S4). It is noteworthy that the two levels 'colonies of social animals' and 'populations' that we merged to test for differences among intraspecific levels (among-population vs. within-populations) shared very similar effect sizes in average (fig. S5).

Figure 2: (A) Phylogenetic tree of species in our dataset, with species' names and taxonomic groups, with silhouettes of iconic taxa within each group. (B) Pie charts of main factors accounted for in the meta-analysis, with respective sample sizes of |*Zr*| effect sizes.

 Publication Bias

The funnel plot of *Zr* effect sizes was roughly symmetric (fig. S6), suggesting minor publication

bias. Raw effect sizes indicated that the most extreme relationships varied between $-3 \leq Zr \geq$

3, which is similar to $-0.99 \leq r \geq 0.99$. Effect sizes $|Zr|$ being comprised between 1 and 2 or

equivalently $|r|$ between 0.76 and 0.96 were not rare, while most part of effect sizes *Zr* were

comprised between -1 and 1, or equivalently $|r|$ between 0 and 0.76 (fig. S6). We did not

observe temporal bias ($t = -1.200$; $P = .230$), but as we worked with absolute effect sizes, the

effect of sampling variance *vi* was strongly positive ($t = 5.24$; $P < .0001$).

Estimated Effect Sizes

The global effect size of relationships between traits and ecological responses was non-null

($|r|_{unbiased} = 0.24$) as its 95% confidence intervals (further abbreviated '95% CI') excluded zero

([0.20 – 0.27] 95% CI; table 1). Effect sizes among populations were twice higher than among

individuals (fig. S7), but it was not significant (table 1).

Effect sizes for reproduction were twice lower than for foraging performance and trophic niche

(fig. 3, table 1). Effect sizes for community were twice higher than effect sizes for reproduction

(fig. 3, table 1). The same analysis performed on the whole dataset led to similar estimates, at

the difference that responses at the ecosystem-level appeared significantly higher than

reproduction (table S4).

Figure 3: Orchard plot of model estimates of |*Zr*| effects sizes estimated for each ecological response at the among-individual level (n=952 effect sizes). The size of each point is proportional to the precision of the effect size (1/SE). Thick and thin error bars give 95% confidence and prediction intervals, respectively. Sample sizes (k) and number of studies (in brackets) are given for each category of ecological responses. Model estimates are reported in table 1.

Physiological and behavioral traits showed effect sizes 28% higher than morphological ones (fig. 4, table 1). The difference between physiology and morphology was not significant when tested on the whole dataset (i.e., including both individual and among-population observations; table S4). Differences among trait type categories were not reflected in the relative frequencies of trait types included in the first quartile range (25th percentile) of effect sizes distribution (morphological: 37% physiological: 19%, and behavioral traits: 44%; fig. 4), but effect sizes with large magnitudes in the dataset were corroborating statistical results, as morphological (17%) and physiological traits (17%) were less represented than behavioral traits (66%) in the third quartile (75th percentile) of the effect sizes distribution (fig. 4).

Figure 4: Orchard plot of model estimates of |*Zr*| effects sizes estimated for each trait type category at the among-individual level (n=952 effect sizes). The size of each point is proportional to the precision of the effect size (1/SE). Thick and thin error bars give 95% confidence and prediction intervals, respectively. Sample sizes (*k*) and number of studies (in brackets) are given for each trait type category. Model estimates are reported in table 1. Categories are shown in blue, yellow, and orange for morphological, physiological and behavioral traits, respectively.

Reproductive organ relative size (n=1), immunity (n=3), morphotypes (n=75), hormone levels (n=11), and assimilation (n=4) were the three sub-trait categories maintaining the strongest relationships (fig. 5). Effect size with large values and estimated with sufficient sample sizes were then activity (n=133), metabolism (n=89), boldness (n=91), stoichiometry (n=52), exploration (n=92), and aggressiveness (n=57) (fig. 5). Some sample sizes were small and expressed large 95% CIs. We therefore advocate for caution when interpreting those results, as we refrained to perform direct statistical comparisons among sub-trait categories. In ranking effect sizes by their magnitudes, we instead aimed at a semi-quantitative analysis.

Figure 5: Orchard plot of model estimates of |*Zr*| effects sizes for each sub-trait type, ranked from highest (top) to lowest estimates (bottom). The size of bubbles is proportional to the precision of the effect sizes (1/SE). Thick and thin error bars give 95% confidence and prediction intervals, respectively. Sample sizes (k) and number of studies in brackets are given for each trait type category. Categories are shown in blue, yellow, and orange for morphological, physiological and behavioral traits, respectively.

Table 1: Mean effect sizes and 95% confidence intervals for (A) the intercept-only model
(global estimate, H_0), and the model for intraspecific levels (among- and within populations;
H_1) for all the dataset. (B) Estimates given for response types (H_2) and trait types (H_3), are based
on a subset of the data, including only effect sizes at the individual level (n=952; 94% of the
dataset). Raw estimated effect sizes are given as |Zr| and |r|, as well as with their unbiased
estimates (|Zr| – |Zr| null). Confidence intervals at 95% are given in brackets. For each model,
(†) indicates the category with the lowest estimated effect size, and the categories shown in
bold (*) are those with significantly higher estimates compared to (†). Pairwise comparison
statistics (z- and P- values) are given accordingly.

Parameters (/model)	|Zr|	|r|	|Zr|null	|Zr|unbiased	|r|unbiased	z	P
(A) (/Intercept-only)							
Global estimate	0.30 (0.25 – 0.34)	0.29 (0.25 – 0.33)	0.05 (0.05 – 0.06)	0.25 (0.20 – 0.28)	0.24 (0.20 – 0.27)	–	–
(/Intraspecific levels)							
Within-populations (†) *(individuals)*	0.28 (0.24 – 0.33)	0.28 (0.24 – 0.32)	0.06 (0.05 – 0.06)	0.22 (0.19 – 0.27)	0.22 (0.19 – 0.25)	–	–
Among-populations	0.51 (0.28 – 0.75)	0.47 (0.27 – 0.64)	0.04 (0.03 – 0.05)	0.47 (0.25 – 0.70)	0.43 (0.24 – 0.59)	1.90	.0572
(B) (/Trait types model)							
Morphology (†)	0.25 (0.21 – 0.30)	0.25 (0.20 – 0.29)	0.05 (0.05 – 0.06)	0.20 (0.17 – 0.24)	0.20 (0.16 – 0.23)	–	–
Physiology (*)	**0.32 (0.27 – 0.38)**	**0.31 (0.26 – 0.36)**	**0.06 (0.04 – 0.07)**	**0.26 (0.23 – 0.31)**	**0.26 (0.22 – 0.29)**	**2.42**	**.0157**
Behavior (*)	**0.32 (0.27 – 0.37)**	**0.31 (0.26 – 0.36)**	**0.06 (0.05 – 0.07)**	**0.26 (0.22 – 0.30)**	**0.25 (0.21 – 0.29)**	**2.55**	**.0108**
(/Response types)							
Foraging (*)	**0.34 (0.26 – 0.42)**	**0.33 (0.25 – 0.40)**	**0.07 (0.05 – 0.08)**	**0.27 (0.21– 0.34)**	**0.26 (0.20 – 0.32)**	**2.32**	**.0204**
Trophic niche (*)	**0.35 (0.25 – 0.45)**	**0.34 (0.25 – 0.42)**	**0.06 (0.05 – 0.07)**	**0.29 (0.20 – 0.38)**	**0.28 (0.20 – 0.35)**	**2.19**	**.0284**
Growth	0.31 (0.23 – 0.39)	0.30 (0.23 – 0.37)	0.06 (0.05 – 0.07)	0.25 (0.18 – 0.32)	0.24 (0.18 – 0.30)	1.80	.0717
Survival	0.27 (0.20 – 0.35)	0.26 (0.19 – 0.33)	0.05 (0.04 – 0.06)	0.22 (0.16 – 0.29)	0.22 (0.15 – 0.27)	1.21	.2271
Reproduction (†)	0.22 (0.15 – 0.29)	0.21 (0.14 – 0.28)	0.05 (0.04 – 0.06)	0.17 (0.11 – 0.23)	0.16 (0.10 – 0.22)	–	–
Community (*)	**0.37 (0.23 – 0.50)**	**0.35 (0.22 – 0.47)**	**0.08 (0.07 – 0.09)**	**0.29 (0.16 – 0.41)**	**0.27 (0.16 – 0.37)**	**2.00**	**.0466**
Ecosystem	0.33 (0.20 – 0.46)	0.32 (0.19 – 0.43)	0.08 (0.07 – 0.09)	0.25 (0.13 – 0.37)	0.24 (0.13 – 0.33)	1.53	.1264

433
434

Discussion

With this meta-analysis, we analyzed four decades of studies in ecology and evolution with the

aim to improve our knowledge of the magnitude of relationships between ITV and ecological

dynamics (Des Roches et al. 2018; Raffard et al. 2019). Not only our analysis emphasized that

ITV matters for ecology, but also that the intraspecific level (among populations vs. among

individuals), the type of ecological response, and the type of trait (morphology, physiology, and

behavior) are three important factors driving the strength of relationships between traits and

ecological responses within species of animals. Our synthesis focusing largely on trait differences among individuals provides the first broad sense estimates for the magnitude of relationships between traits and ecological responses. This opens new avenues to improve our knowledge of the links between individual traits and ecosystem processes, which we will distill through this discussion.

Global Effect Sizes for Relationship Magnitudes between ITV and Ecological Responses

The global effect size we report ($|r|_{unbiased}$ = 0.24 [0.20 – 0.27] 95% CI; table 1) might be considered small to moderate in absolute terms (Nakagawa and Cuthill 2007). However, it falls in the upper range of effect sizes reported in ecology and evolution (*ca.* 95% of $|r|$ values are between 0.14 and 0.26; Møller & Jennions, 2002). This remains true when emphasizing only among-individual variability ($|r|_{unbiased}$ = 0.22 [0.19 – 0.25] 95% CI; table 1).

Effect sizes among populations and groups of eusocial animals were strong in absolute terms ($|r|_{unbiased}$ = 0.43 [0.24 – 0.59] 95% CI; table 1), and twice stronger than effect sizes collected among individuals. However, this difference was not significant (P = .0572), as our test surely suffered from a too low sample size in the among-population category (fig. S7). While more data are needed to reach a conclusion, the twofold difference tends to support our first hypothesis. Populations whom phenotypically diverged, or expressed local trait matching with ecological processes show links between their traits and ecological responses of a greater magnitude. However, the alternative hypothesis that relationships assessed among individuals could be as large as the ones obtained among populations, remains a possibility as well. In a world where genetic and phenotypic diversity are eroding and changing fast (Exposito-Alonso et al. 2022; Sanderson et al. 2023), ITV has important implications for the conservation and management of intraspecific diversity (Blanchet et al. 2020; Des Roches et al. 2021). Management and conservation plans should consider that both ITV within and among

populations are relevant biological levels, as these two levels are nested, meaning that changes in ITV within populations are ultimately changing among population ITV, with overlooked consequences for ecological dynamics.

The heterogeneity in our dataset was large (87%), as were the prediction intervals at 95% around our estimates (fig. 3–5). Despite the extensive scope of our meta-analysis, the 95% confidence intervals around point estimates stays small, meaning that the multi-level structure of our models accounted well for this heterogeneity. Accordingly, all the covariates that we tested were not significant, suggesting that our results may generalize across years of publication, aquatic and terrestrial realms, consumers and predators, as well as methodological approaches (field, mesocosms, microcosms). Meta-analyses in ecology and evolution often consider that large total heterogeneities (often inherent to both disciplines) should inevitably lower the generality of findings, but this might not be always the case in multi-level meta-analyses. The within-study heterogeneity can comprise a substantial part of total heterogeneity (as in the present meta-analysis), which would need to be accounted for when evaluating the generality of findings, which lies more on the between-study heterogeneity (Yang et al. 2024*b*).

How ITV Relates to Ecological Responses across Biological Levels?

We expected that the overall strength of a cascade of indirect relationships between ITV and an ecological response would decrease from low levels to high levels of biological organization. Effects of ITV could dissipate due to the accumulation of intermediate relationships among distal relationships, for instance between a given phenotypic trait and a response at the community or ecosystem level (*H2*; see Bailey et al., 2009). Our results did not support this hypothesis. We rather observed large magnitudes for organismal performance and community-level responses, while relationships with fitness (survival and reproduction) displayed the

smallest magnitudes. The same analysis performed on the whole dataset showed that both

community and ecosystem-level responses were higher than reproduction.

Effect sizes were larger for foraging and trophic niche responses than for reproduction. This

can be explained since reproduction outcomes are still complex to predict, as they are

influenced by intrinsic (genetic and phenotypic), and extrinsic processes such as varying

environmental conditions and random events (Merilä & Sheldon, 2000). Then, effect sizes for

community-level responses were larger than for reproduction. This could arise because of

methodological differences. In our dataset, most studies investigating fitness took place in

nature (*ca.* 70%), whereas most studies investigating responses at the community and

ecosystem levels were conducted in mesocosms (*ca.* 75%). A dampening of effect size

magnitude can occur because of a greater complexity in nature compared to controlled

conditions (Hendry 2019). Nevertheless, we did not find differences between estimates from

field, mesocosm or microcosm studies. Thus, we are left with the result already reported by

others (Des Roches et al., 2018; Raffard et al., 2019) that the magnitude of the links between

ITV and ecological responses are substantial for community dynamics, even when they arise

from phenotypic differences among individuals of similar body sizes (this study).

The effect sizes for fitness proxies reported here are of similar nature to linear estimates of

directional selection (Hendry 2017). Indeed, our estimates for reproduction ($|r|_{\text{unbiased}} = 0.16$ for

among-individuals and 0.16 for the whole dataset) matched accurately with those reported

elsewhere for linear directional selection ($|\beta| = 0.15$; see Kingsolver et al., 2012). One potential

reason for the relatively small effect sizes of ITV on fitness related responses is that past

adaptation of individuals to the local environment has already shaped ITV near their optima

(i.e., stabilising selection). Another mechanism can be fluctuating selection regimes averaging

over time (Wright et al., 2019). As such, the estimated selection on traits is no longer strong:

"*selection erases its traces*" (Haller and Hendry 2014). By contrast, ITV effects on community

and ecosystem dynamics do not reflect adaptations *per se* and are not under selection to eliminate such effects, which might explain why the effect sizes we observed at these levels were larger than on reproduction. Yet, this does not mean that selection is weaker than the links between ITV and community/ ecosystem responses. Subtle rates of changes in trait distributions, when scaled up to population dynamics and higher numbers of generations, can lead to large eco-evolutionary dynamics (Hansen et al. 2011).

Other factors may explain why effect sizes were the largest at the community level. First, many relationships in ecology are non-linear. Because of Jensen's inequality (Bolnick et al. 2011), concave-up intermediate relationships can amplify the effects of ITV at high levels of biological organization (see Bailey et al., 2009). Second, large effects of ITV can occur through direct effects on community and ecosystem responses (i.e., 'trait-mediated effects'; Schmitz et al., 2004), which could be particularly strong as they may not dissipate through many intermediate relationships (fig. 1a). This can be particularly true when focal animals are keystone (Paine 1980), foundation (Ellison 2019), engineer species (Romero et al. 2015), or are prone to niche construction (Laland et al. 1999). A great example comes from the Yellowstone, where a small subset of wolf individuals within a pack are specialized on beavers (i.e., some individuals are hunting ~200% more beavers; Bump et al., 2022), and therefore, are contributing disproportionately to wetland appearance/loss dynamics (Gable et al. 2020).

Do Trait Types Showed Different Relationship Magnitudes with Ecological Responses?

We hypothesized that physiological and behavioral traits would maintain stronger ecological relationships than morphological ones (*H3*). Accordingly, physiological and behavioral trait variation among individuals showed 28% stronger relationships with ecological responses than did morphological traits. This confirms that traits reflecting energetic tradeoffs and interactions among individuals matter for our understanding of ecological processes. For instance,

functional responses of predator crabs (*Panopeus herbstii*) on mussel prey (*Brachidontes exustus*) are mediated by consistent variation among individuals in their activity levels, suggesting strong ecosystem-level effects given that mussels are famous ecosystem engineers (Toscano and Griffen 2014). Physiological traits showed strong relationship strengths across ecological responses (fig. 4&5). An example is how variations in corticosterone levels among individuals predict long-term reproductive performances in wandering albatross (*Diomedea exulans*; Angelier et al., 2010). This is consistent with the fact that variations in metabolic, immune, hormonal, and stoichiometric traits are all believed to be strongly related with fitness, community and ecosystem responses (Elser et al. 2000; Ricklefs and Wikelski 2002; Brandl et al. 2022).

Understanding how different phenotypic dimensions relate to ecological differences among individuals or populations is a priori challenging, and this motivated the present work. From a general standpoint, morphology could express a *potential* for ecological effects, where physiological and behavioral traits might be closer to *realized* effects (Wainwright and Reilly 1994). Our results are in agreement with this general view, albeit the average difference among large categories of traits were rather small in absolute (table 1). Morphological traits that showed the largest effect sizes were also the most *a priori* functionally-relevant (fig. 5). The category with the highest effect size and sufficient sample size was '*Morphotype*', which refers to dimorphic populations, or ecotypes which diverged ecologically hundreds to thousands of generations ago (Harmon et al., 2009). '*Locomotion*' designates morphological traits linked to the capacity for an individual to perform locomotion ('*activity*'), the behavioral sub-category showing the strongest effect size (fig. 5). The trait coined '*body condition*' is a proxy for physiological status (energy reserve). Morphological traits with the less general functional relevance for a broad range of ecological responses, such as linear morphometric traits (labelled '*specific morphometric traits*') exhibited unexpectedly, lower effect sizes (fig. 5).

Our results at the intraspecific level are contrasting with the strong reliance on morphological

traits in TBE (Dawson et al. 2021). This popularity for morphology may hold into that those

traits are easily measured on immobilized or dead animals. Another reason for the popularity

of morphology could be its low flexibility (i.e., low intra-individual variation) and high

heritability compared to physiology and behavior (Mousseau and Roff 1987). Although

morphological traits do not maintain the strongest ecological relationships in average (this

meta-analysis), since they are less labile and more heritable in comparison to physiological or

behavioral traits, they could maintain more persistent eco-evolutionary dynamics than

physiology and behavior. This idea would need to be investigatred. However, the lability of

physiological and behavioral traits does not mean randomness, as it should not prevent their use

in TBE. Those trait types show moderate to high repeatability among individuals (Bell et al.

2009), and in some cases, large heritabilities as well (Dochtermann et al. 2014). Moreover,

recent advances show that the residual intra-individual variation for those traits can be

structured, with predictable *vs.* unpredictable individuals (Stamps et al. 2012). Finally, if we

aim at quantifying the propensity of traits to be involved in eco-evolutionary dynamics, it would

be promising to compare the evolvabilities (Hansen et al. 2011) of those trait types with metrics

expressing their propensity to alter ecological dynamics.

General formulations have been suggested to explore theoretically the links between traits,

abundances, and ecosystem functions (Violle et al. 2007). Our trait functionality estimates can

help to parametrize eco-evolutionary or trait-based models intended to assess how shifts in trait

distributions through evolution, selection, patch dynamics, etc. can affect various ecological

responses from ecological performances to ecosystem functioning (Govaert et al. 2019).

Therefore, we invite to investigate how changes in the moments of traits distribution (mean,

variance, skewness and kurtosis), for instance under selection (Kingsolver et al. 2012) and

various ecological functionality estimates available for different traits (our approach), can

change the outcomes on ecological processes (Norberg et al. 2001; Barabás et al. 2022). In

addition, we invite to continue refining empirical estimates for trait selection rates, evolutionary

potentials, and trait's functionality for different trait categories, and to assess how those metrics

covary or not. This would help identify particular facets of ITV showing both large evolutionary

change potential, as well as large potential to link to ecosystem functions (Blanchet et al. 2020;

Des Roches et al. 2021).

Concluding Remarks

Ideal traits to move forward TBE of animals should meet two criteria: (*i*) link to a vast range of

ecological functions operating at different levels, and (*ii*) be 'universal', i.e., can be measured

across intra- and interspecific levels of various animal taxonomic groups (Carmona et al. 2016).

Here, several continuous traits showed relatively strong links with ecological responses, as they

can meet the second criterion (e.g., hormone levels, activity, metabolism, boldness,

stoichiometry, excretion, sociability, body condition, and cognitive abilities; fig. 5).

Technological developments should facilitate the measurement of such traits at the individual

level, allowing multi-trait approaches. Telemetry and camera traps, coupled with artificial

intelligence (AI) algorithms, are improving the study of movements, activity and stereotypic

behaviors of vertebrates in the field (Suraci et al. 2017; Salama AbdELminaam et al. 2020).

Flow cytometry, (-thermal) imagery, and respiration chambers are facilitating the measurement

of shape and size, metabolism, and behavior on aquatic invertebrates (Réveillon and Becks

2022). In a similar way, the continuous miniaturization of telemetry allows collecting

behavioral data on insects as small as bumblebees (Hagen et al. 2011).

We advocate the assessment of the functionality of traits used in TBE. This effort should

parallel their harmonisation towards a universal set of continuous individual-based traits and

their inclusion in databases. In the face of global changes, this information should help towards

a better appraisal of the implications of biodiversity changes at the levels of individuals, populations, species, communities and ecosystems. To unravel the ecological implications of animal intraspecific diversity, we encourage ecologists not to neglect physiological and behavioral traits. Future studies at the intraspecific level will gain in predictive abilities by addressing simultaneously all the facets of a phenotype and their covariations. Multiple morphological, physiological, and behavioral traits are integrated into one phenotype, and acknowledging this would inform about ecological and fitness trade-offs (Arnold 1983; Violle et al. 2007). Such an approach at the individual level will improve our integration of biological scales, evolutionary and ecological processes, as well as predictions of the effects of global changes on ecosystems.

Acknowledgements

We are grateful to the researchers who contributed to initial stages of the present work, especially J. Cucherousset and L. Závorka. We thank B. Corbara for the discussion about our results. We also thank A. Bruder, R. Oester, A. M. Roth, L. Govaert, S. J. Brandl, and J. O'Brien for their help in improving the manuscript before its submission to the journal. This work was funded by a French National Research grant called 'RESILIENCE' [ANR-18-CE02-0015], and a Swiss National Foundation for Science grant called 'MULTIBEF' [315230_204998].

Authorship

T.R. assembled the first ideas. T.R. designed the study with the guidance of A. R., S. B., A. P. H., and R. C.. T. R. and A. R. performed the systematic review of the literature in parallel, deleted duplicates and validated selected studies. T.R. extracted the data from the selected papers, did the statistical analyses and wrote a first draft of the manuscript. All co-authors then contributed to the discussion of the results and to the editing of the manuscript until its publication.

Data accessibility

The data and R code supporting the results and their reproducibility can be accessed through a

public repository (Figshare) [https://doi.org/10.6084/m9.figshare.27377529.v1].

Conflicts of interest

The authors have no conflict of interest to declare.

Literature Cited

Albert, C. H. 2015. Intraspecific trait variability matters. Journal of Vegetation Science 26:7–8.

Alberti, M., C. Correa, J. M. Marzluff, A. P. Hendry, E. P. Palkovacs, K. M. Gotanda, V. M. Hunt, et al. 2017. Global urban signatures of phenotypic change in animal and plant populations. Proceedings of the National Academy of Sciences 114:8951–8956.

Angelier, F., J. C. Wingfield, H. Weimerskirch, and O. Chastel. 2010. Hormonal correlates of individual quality in a long-lived bird: a test of the 'corticosterone–fitness hypothesis.' Biology Letters 6:846–849.

Arnold, S. J. 1983. Morphology, Performance and Fitness. American Zoologist 23:347–361.

Bailey, J. K., J. A. Schweitzer, F. Ubeda, J. Koricheva, C. J. LeRoy, M. D. Madritch, B. J. Rehill, et al. 2009. From genes to ecosystems: a synthesis of the effects of plant genetic factors across levels of organization. Philosophical Transactions of the Royal Society B: Biological Sciences 364:1607–1616.

Barabás, G., C. Parent, A. Kraemer, F. Van de Perre, and F. De Laender. 2022. The evolution of trait variance creates a tension between species diversity and functional diversity. Nature Communications 13:2521.

Bell, A. M., S. J. Hankison, and K. L. Laskowski. 2009. The repeatability of behaviour: a meta-analysis. Animal Behaviour 77:771–783.

Bertossa, R. C. 2011. Morphology and behaviour: functional links in development and evolution. Philosophical Transactions of the Royal Society B: Biological Sciences 366:2056–2068.

Biro, P. A., and J. A. Stamps. 2008. Are animal personality traits linked to life-history productivity? Trends in Ecology & Evolution 23:361–368.

Blanchet, S., J. G. Prunier, I. Paz-Vinas, K. Saint-Pé, O. Rey, A. Raffard, E. Mathieu-Bégné, et al. 2020. A river runs through it: The causes, consequences, and management of intraspecific diversity in river networks. Evolutionary Applications 13:1195–1213.

Bolnick, D. I., P. Amarasekare, M. S. Araújo, R. Bürger, J. M. Levine, M. Novak, V. H. W. Rudolf, et al. 2011. Why intraspecific trait variation matters in community ecology. Trends in Ecology & Evolution 26:183–192.

Brandl, S. J., J. S. Lefcheck, A. E. Bates, D. B. Rasher, and T. Norin. 2022. Can metabolic traits explain animal community assembly and functioning? Biological Reviews.12892.

Bump, J., T. Gable, S. Johnson-Bice, A. Homkes, D. Freund, S. Windels, and S. Chakrabarti. 2022. Predator personalities alter ecosystem services. Frontiers in Ecology and the Environment 20:275–277.

Careau, V., and T. Garland. 2012. Performance, Personality, and Energetics: Correlation, Causation, and Mechanism. Physiological and Biochemical Zoology 85:543–571.

Careau, V., D. Thomas, M. M. Humphries, and D. Réale. 2008. Energy metabolism and animal personality. Oikos 117:641–653.

Carmona, C. P., F. de Bello, N. W. H. Mason, and J. Lepš. 2016. Traits Without Borders: Integrating Functional Diversity Across Scales. Trends in Ecology & Evolution 31:382–394.

Ceballos, G., P. R. Ehrlich, and R. Dirzo. 2017. Biological annihilation via the ongoing sixth mass extinction signaled by vertebrate population losses and declines. Proceedings of the National Academy of Sciences 114.

Chacón-Labella, J., C. Hinojo-Hinojo, T. Bohner, M. Castorena, C. Violle, V. Vandvik, and B. J. Enquist. 2022. How to improve scaling from traits to ecosystem processes. Trends in Ecology & Evolution S0169534722002749.

Cianciaruso, M. V., M. A. Batalha, K. J. Gaston, and O. L. Petchey. 2009. Including intraspecific variability in functional diversity. Ecology 90:81–89.

Darwin, C., and A. Wallace. 1858. On the Tendency of Species to form Varieties; and on the Perpetuation of Varieties and Species by Natural Means of Selection. Journal of the Proceedings of the Linnean Society of London. Zoology 3:45–62.

Dawson, S. K., C. P. Carmona, M. González-Suárez, M. Jönsson, F. Chichorro, M. Mallen-Cooper, Y. Melero, et al. 2021. The traits of "trait ecologists": An analysis of the use of trait and functional trait terminology. Ecology and Evolution 11:16434–16445.

de Bello, F., Z. Botta-Dukát, J. Lepš, and P. Fibich. 2021. Towards a more balanced combination of multiple traits when computing functional differences between species. Methods in Ecology and Evolution 12:443–448.

Des Roches, S., L. H. Pendleton, B. Shapiro, and E. P. Palkovacs. 2021. Conserving intraspecific variation for nature's contributions to people. Nature Ecology & Evolution 5:574–582.

Des Roches, S., D. M. Post, N. E. Turley, J. K. Bailey, A. P. Hendry, M. T. Kinnison, J. A. Schweitzer, et al. 2018. The ecological importance of intraspecific variation. Nature Ecology & Evolution 2:57–64.

Dochtermann, N. A., T. Schwab, and A. Sih. 2014. The contribution of additive genetic variation to personality variation: heritability of personality. Proceedings of the Royal Society B: Biological Sciences 282:20142201–20142201.

Ellison, A. M. 2019. Foundation Species, Non-trophic Interactions, and the Value of Being Common. iScience 13:254–268.

Elser, J. j., R. w. Sterner, E. Gorokhova, W. f. Fagan, T. a. Markow, J. b. Cotner, J. f. Harrison, et al. 2000. Biological stoichiometry from genes to ecosystems. Ecology Letters 3:540–550.

Elton, C. S. 1927. Animal ecology, by Charles Elton; with an introduction by Julian S. Huxley. Macmillan Co., New York,.

Enquist, B. J., J. Norberg, S. P. Bonser, C. Violle, C. T. Webb, A. Henderson, L. L. Sloat, et al. 2015. Scaling from Traits to Ecosystems. Pages 249–318 *in*Advances in Ecological Research (Vol. 52). Elsevier.

Exposito-Alonso, M., T. R. Booker, L. Czech, L. Gillespie, S. Hateley, C. C. Kyriazis, P. L. M. Lang, et al. 2022. Genetic diversity loss in the Anthropocene. Science 377:1431–1435.

Funk, J. L., J. E. Larson, G. M. Ames, B. J. Butterfield, J. Cavender-Bares, J. Firn, D. C. Laughlin, et al. 2017. Revisiting the Holy Grail: using plant functional traits to understand ecological processes. Biological Reviews 92:1156–1173.

Gable, T. D., S. M. Johnson-Bice, A. T. Homkes, S. K. Windels, and J. K. Bump. 2020. Outsized effect of predation: Wolves alter wetland creation and recolonization by killing ecosystem engineers. Science Advances 6:eabc5439.

Gibert, J. P., A. I. Dell, J. P. DeLong, and S. Pawar. 2015. Scaling-up Trait Variation from Individuals to Ecosystems. Pages 1–17 *in*Advances in Ecological Research (Vol. 52). Elsevier.

Gordon, D. M. 2011. The fusion of behavioral ecology and ecology. Behavioral Ecology 22:225–230.

Govaert, L., E. A. Fronhofer, S. Lion, C. Eizaguirre, D. Bonte, M. Egas, A. P. Hendry, et al. 2019. Eco-evolutionary feedbacks—Theoretical models and perspectives. Functional Ecology 33:13–30.

Grafen, A. 1989. The phylogenetic regression. Philosophical Transactions of the Royal Society of London. B, Biological Sciences 326:119–157.

Green, S. J., C. B. Brookson, N. A. Hardy, and L. B. Crowder. 2022. Trait-based approaches to global change ecology: moving from description to prediction. Proceedings of the Royal Society B: Biological Sciences 289:20220071.

Hagen, M., M. Wikelski, and W. D. Kissling. 2011. Space Use of Bumblebees (Bombus spp.) Revealed by Radio-Tracking. PLoS ONE 6:e19997.

Haller, B. C., and A. P. Hendry. 2014. Solving the paradox of stasis: squashed stabilizing selection and the limits of detection: selection and the limits of detection. Evolution 68:483–500.

Hansen, T. F., C. Pélabon, and D. Houle. 2011. Heritability is not Evolvability. Evolutionary
Biology 38:258–277.

Harmon, L. J., B. Matthews, S. Des Roches, J. M. Chase, J. B. Shurin, and D. Schluter. 2009.
Evolutionary diversification in stickleback affects ecosystem functioning. Nature 458:1167–
1170.

Hart, S. P., S. J. Schreiber, and J. M. Levine. 2016. How variation between individuals affects
species coexistence. Ecology Letters 19:825–838.

Hendry, A. P. 2017. Eco-evolutionary dynamics. Princeton University Press, Princeton.

———. 2019. A critique for eco-evolutionary dynamics. Functional Ecology 33:84–94.

Hinchliff, C. E., S. A. Smith, J. F. Allman, J. G. Burleigh, R. Chaudhary, L. M. Coghill, K. A.
Crandall, et al. 2015. Synthesis of phylogeny and taxonomy into a comprehensive tree of life.
Proceedings of the National Academy of Sciences 112:12764–12769.

Hutchinson, G. E. 1957. Concluding Remarks. Cold Spring Harbor Symposia on Quantitative
Biology 22:415–427.

Kingsolver, J. G., S. E. Diamond, A. M. Siepielski, and S. M. Carlson. 2012. Synthetic analyses
of phenotypic selection in natural populations: lessons, limitations and future directions.
Evolutionary Ecology 26:1101–1118.

Kleiber, M. 1932. Body size and metabolism. Hilgardia 6:315–353.

Laland, K. N., F. J. Odling-Smee, and M. W. Feldman. 1999. Evolutionary consequences of
niche construction and their implications for ecology. Proceedings of the National Academy of
Sciences 96:10242–10247.

Lawton, J. H. 1994. What Do Species Do in Ecosystems? Oikos 71:367.

Lecerf, A., and E. Chauvet. 2008. Intraspecific variability in leaf traits strongly affects alder
leaf decomposition in a stream. Basic and Applied Ecology 9:598–605.

Lefcheck, J. S., V. A. G. Bastazini, and J. N. Griffin. 2015. Choosing and using multiple traits
in functional diversity research. Environmental Conservation 42:104–107.

Macarthur, R., and R. Levins. 1967. The Limiting Similarity, Convergence, and Divergence of
Coexisting Species. The American Naturalist 101:377–385.

McGhee, K. E., L. M. Pintor, and A. M. Bell. 2013. Reciprocal Behavioral Plasticity and
Behavioral Types during Predator-Prey Interactions. The American Naturalist 182:704–717.

McGill, B., B. Enquist, E. Weiher, and M. Westoby. 2006. Rebuilding community ecology
from functional traits. Trends in Ecology & Evolution 21:178–185.

Merilä, J., and B. C. Sheldon. 2000. Lifetime Reproductive Success and Heritability in Nature.
The American Naturalist 155:301–310.

Messier, J., B. J. McGill, and M. J. Lechowicz. 2010. How do traits vary across ecological
scales? A case for trait-based ecology: How do traits vary across ecological scales? Ecology
Letters 13:838–848.

Michonneau, F., J. W. Brown, and D. J. Winter. 2016. rotl: an R package to interact with the
Open Tree of Life data. Methods in Ecology and Evolution 7:1476–1481.

Mlambo, M. C. 2014. Not all traits are 'functional': insights from taxonomy and biodiversity-
ecosystem functioning research. Biodiversity and Conservation 23:781–790.

Møller, A., and M. D. Jennions. 2002. How much variance can be explained by ecologists and
evolutionary biologists? Oecologia 132:492–500.

Moran, N. P., A. Sánchez-Tójar, H. Schielzeth, and K. Reinhold. 2021. Poor nutritional
condition promotes high-risk behaviours: a systematic review and meta-analysis. Biological
Reviews 96:269–288.

Moran, N. P., B. B. M. Wong, and R. M. Thompson. 2017. Weaving animal temperament into
food webs: implications for biodiversity. Oikos 126:917–930.

Morrissey, M. B. 2016. Meta-analysis of magnitudes, differences and variation in evolutionary
parameters. Journal of Evolutionary Biology 29:1882–1904.

Mousseau, T. A., and D. A. Roff. 1987. Natural selection and the heritability of fitness
components. Heredity 59:181–197.

Nakagawa, S., and I. C. Cuthill. 2007. Effect size, confidence interval and statistical
significance: a practical guide for biologists. Biological Reviews 82:591–605.

Nakagawa, S., M. Lagisz, M. D. Jennions, J. Koricheva, D. W. A. Noble, T. H. Parker, A.
Sánchez-Tójar, et al. 2022. Methods for testing publication bias in ecological and evolutionary
meta-analyses. Methods in Ecology and Evolution 13:4–21.

Nakagawa, S., M. Lagisz, R. E. O'Dea, P. Pottier, J. Rutkowska, A. M. Senior, Y. Yang, et al.
2023. orchaRd 2.0: An R package for visualising meta-analyses with orchard plots. Methods in
Ecology and Evolution 14:2003–2010.

Noble, D. W. A., M. Lagisz, R. E. O'dea, and S. Nakagawa. 2017. Nonindependence and
sensitivity analyses in ecological and evolutionary meta-analyses. Molecular Ecology 26:2410–
2425.

Norberg, J., D. P. Swaney, J. Dushoff, J. Lin, R. Casagrandi, and S. A. Levin. 2001. Phenotypic
diversity and ecosystem functioning in changing environments: A theoretical framework.
Proceedings of the National Academy of Sciences 98:11376–11381.

O'Dea, R. E., M. Lagisz, M. D. Jennions, J. Koricheva, D. W. A. Noble, T. H. Parker, J.
Gurevitch, et al. 2021. Preferred reporting items for systematic reviews and meta-analyses in
ecology and evolutionary biology: a PRISMA extension. Biological Reviews 96:1695–1722.

Paine, R. T. 1980. Food Webs: Linkage, Interaction Strength and Community Infrastructure.
The Journal of Animal Ecology 49:666.

Paradis, E., and K. Schliep. 2019. ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. Bioinformatics 35:526–528.

Pelletier, F., D. Garant, and A. P. Hendry. 2009. Eco-evolutionary dynamics. Philosophical
Transactions of the Royal Society B: Biological Sciences 364:1483–1489.

Petchey, O. L., and K. J. Gaston. 2006. Functional diversity: back to basics and looking forward.
Ecology Letters 9:741–758.

Peters, R. H. 1993. The ecological implications of body size. Univ. Press, Cambridge [u.a.]

Pianka, E. R. 1970. On r- and K-Selection. The American Naturalist 104:592–597.

Post, D. M., E. P. Palkovacs, E. G. Schielke, and S. I. Dodson. 2008. Intraspecific variation in
a predator affects community structure and cascading trophic interactions. Ecology 89:2019–
2032.

Raffard, A., A. Lecerf, J. Cote, M. Buoro, R. Lassus, and J. Cucherousset. 2017. The functional
syndrome: linking individual trait variability to ecosystem functioning. Proceedings of the
Royal Society B: Biological Sciences 284:20171893.

Raffard, A., F. Santoul, J. Cucherousset, and S. Blanchet. 2019. The community and ecosystem
consequences of intraspecific diversity: a meta-analysis: The ecological effects of intraspecific
diversity. Biological Reviews.

Réveillon, T., and L. Becks. 2022. *Consistency in defence and competitiveness trade-off in a*
planktonic predator-prey system (preprint). Ecology.

Ricklefs, R. E., and M. Wikelski. 2002. The physiology/life-history nexus. Trends in Ecology
& Evolution 17:462–468.

Romero, G. Q., T. Gonçalves-Souza, C. Vieira, and J. Koricheva. 2015. Ecosystem engineering
effects on species diversity across ecosystems: a meta-analysis: Ecosystem engineering effects
across ecosystems. Biological Reviews 90:877–890.

Rota, T., J. Jabiol, E. Chauvet, and A. Lecerf. 2018. Phenotypic determinants of inter-individual
variability of litter consumption rate in a detritivore population. Oikos.

Rota, T., J. Jabiol, S. Lamothe, D. Lambrigot, É. Chauvet, and A. Lecerf. 2024. Phenotypes of
predator individuals underpin contrasting ecosystem effects. Freshwater Biology.14242.

Rota, T., A. Lecerf, É. Chauvet, and B. Pey. 2022. The importance of intraspecific variation in
litter consumption rate of aquatic and terrestrial macro-detritivores. Basic and Applied Ecology
63:175–185.

Salama AbdELminaam, D., A. M. Almansori, M. Taha, and E. Badr. 2020. A deep facial
recognition system using computational intelligent algorithms. PLOS ONE 15:e0242269.

Sanderson, S., D. I. Bolnick, M. T. Kinnison, R. E. O'Dea, L. D. Gorné, A. P. Hendry, and K.
M. Gotanda. 2023. Contemporary changes in phenotypic variation, and the potential
consequences for eco-evolutionary dynamics. Ecology Letters 26.

Schleuning, M., D. García, and J. A. Tobias. 2023. Animal functional traits: Towards a trait-
based ecology for whole ecosystems. Functional Ecology 37:4–12.

Schmitz, O. J., V. Krivan, and O. Ovadia. 2004. Trophic cascades: the primacy of trait-mediated
indirect interactions: Primacy of trait-mediated indirect interactions. Ecology Letters 7:153–
163.

Siefert, A., C. Violle, L. Chalmandrier, C. H. Albert, A. Taudiere, A. Fajardo, L. W. Aarssen,
et al. 2015. A global meta-analysis of the relative extent of intraspecific trait variation in plant
communities. Ecology Letters 18:1406–1419.

Sobral, M. 2021. All Traits Are Functional: An Evolutionary Viewpoint. Trends in Plant
Science 26:674–676.

Stamps, J. A., M. Briffa, and P. A. Biro. 2012. Unpredictable animals: individual differences in
intraindividual variability (IIV). Animal Behaviour 83:1325–1334.

Suraci, J. P., M. Clinchy, B. Mugerwa, M. Delsey, D. W. Macdonald, J. A. Smith, C. C.
Wilmers, et al. 2017. A new Automated Behavioural Response system to integrate playback
experiments into camera trap studies. Methods in Ecology and Evolution 8:957–964.

Toscano, B. J., N. J. Gownaris, S. M. Heerhartz, and C. J. Monaco. 2016. Personality, foraging
behavior and specialization: integrating behavioral and food web ecology at the individual
level. Oecologia 182:55–69.

Toscano, B. J., and B. D. Griffen. 2014. Trait-mediated functional responses: predator
behavioural type mediates prey consumption. Journal of Animal Ecology 83:1469–1477.

Van Valen, L. 1965. Morphological Variation and Width of Ecological Niche. The American
Naturalist 99:377–390.

Viechtbauer, W. 2010. Conducting Meta-Analyses in *R* with the **metafor** Package. Journal of
Statistical Software 36.

Violle, C., B. J. Enquist, B. J. McGill, L. Jiang, C. H. Albert, C. Hulshof, V. Jung, et al. 2012.
The return of the variance: intraspecific variability in community ecology. Trends in Ecology
& Evolution 27:244–252.

Violle, C., M.-L. Navas, D. Vile, E. Kazakou, C. Fortunel, I. Hummel, and E. Garnier. 2007.
Let the concept of trait be functional! Oikos 116:882–892.

Wainwright, P. C., and S. M. Reilly, eds. 1994. Ecological morphology: integrative organismal
biology. University of Chicago Press, Chicago.

Wheelwright, N. T., L. F. Keller, and E. Postma. 2014. The effect of trait type and strength of
selection on heritability and evolvability in an island bird population. Evolution 68:3325–3336.

Wong, M. K. L., and C. P. Carmona. 2021. Including intraspecific trait variability to avoid
distortion of functional diversity and ecological inference: Lessons from natural assemblages.
Methods in Ecology and Evolution.

Wright, I. J., P. B. Reich, M. Westoby, D. D. Ackerly, Z. Baruch, F. Bongers, J. Cavender-
Bares, et al. 2004. The worldwide leaf economics spectrum. Nature 428:821–827.

Wright, J., G. H. Bolstad, Y. G. Araya-Ajoy, and N. J. Dingemanse. 2019. Life-history
evolution under fluctuating density-dependent selection and the adaptive alignment of pace-of-
life syndromes: Pace-of-life syndromes. Biological Reviews 94:230–247.

Yang, Y., M. Lagisz, C. Williams, D. W. A. Noble, J. Pan, and S. Nakagawa. 2024*a*. Robust
point and variance estimation for meta-analyses with selective reporting and dependent effect
sizes. Methods in Ecology and Evolution 15:1593–1610.

Yang, Y., D. W. A. Noble, A. M. Senior, M. Lagisz, and S. Nakagawa. 2024*b*. Interpreting
prediction intervals and distributions for decoding biological generality in meta-analyses.
(Preprint)

Literature Cited in The Meta-Analysis (*)

*ADAMS, C.E. & HUNTINGFORD, F.A. (2002) The functional significance of inherited
differences in feeding morphology in a sympatric polymorphic population of Arctic charr.
Evolutionary Ecology **16**, 15–25.

*AKÇAY, Ç., CAMPBELL, S.E. & BEECHER, M.D. (2015) The fitness consequences of honesty:
Under-signalers have a survival advantage in song sparrows. *Evolution* **69**, 3186–3193.

*ALLGEIER, J.E., CLINE, T.J., WALSWORTH, T.E., WATHEN, G., LAYMAN, C.A. & SCHINDLER,
D.E. (2020) Individual behavior drives ecosystem function and the impacts of harvest. *Science
Advances* **6**, eaax8329.

*ÁLVAREZ, D. & NICIEZA, A.G. (2005) Is metabolic rate a reliable predictor of growth and
survival of brown trout (*Salmo trutta*) in the wild? *Canadian Journal of Fisheries and Aquatic
Sciences* **62**, 643–649.

*ANGELIER, F., WINGFIELD, J.C., WEIMERSKIRCH, H. & CHASTEL, O. (2010) Hormonal
correlates of individual quality in a long-lived bird: a test of the 'corticosterone–fitness
hypothesis'. *Biology Letters* **6**, 846–849.

*ARMITAGE, K.B. (1986) Individuality, Social Behavior, and Reproductive Success in Yellow-
Bellied Marmots. *Ecology* **67**, 1186–1193.

*ARMITAGE, K.B. & VAN VUREN, D.H. (2003) Individual differences and reproductive success
in yellow-bellied marmots. *Ethology Ecology & Evolution* **15**, 207–233.

*AUER, S.K., SALIN, K., RUDOLF, A.M., ANDERSON, G.J. & METCALFE, N.B. (2015a) The
optimal combination of standard metabolic rate and aerobic scope for somatic growth depends
on food availability. *Functional Ecology* **29**, 479–486.

*AUER, S.K., SALIN, K., RUDOLF, A.M., ANDERSON, G.J. & METCALFE, N.B. (2015b)
Flexibility in metabolic rate confers a growth advantage under changing food availability.
Journal of Animal Ecology **84**, 1405–1411.

*BAKER, J.A., CRESKO, W.A., FOSTER, S.A. & HEINS, D.C. (2005) Life-history differentiation
of benthic and limnetic ecotypes in a polytypic population of threespine stickleback
(Gasterosteus aculeatus). *Evolutionary Ecology Research* **7**, 121–131.

*BANKS, P.B., NORRDAHL, K. & KORPIMÄKI, E. (2002) Mobility decisions and the predation
risks of reintroduction. *Biological Conservation* **103**, 133–138.

*BEAUCHAMP, G. (2006) Phenotypic Correlates of Scrounging Behavior in Zebra Finches: Role
of Foraging Efficiency and Dominance. *Ethology* **112**, 873–878.

*BERCHTOLD, A.E., COLBORNE, S.F., LONGSTAFFE, F.J. & NEFF, B.D. (2015)
Ecomorphological patterns linking morphology and diet across three populations of
pumpkinseed sunfish (*Lepomis gibbosus*). *Canadian Journal of Zoology* **93**, 289–297.

*BLACKMER, A.L., MAUCK, R.A., ACKERMAN, J.T., HUNTINGTON, C.E., NEVITT, G.A. &
WILLIAMS, J.B. (2005) Exploring individual quality: basal metabolic rate and reproductive
performance in storm-petrels. *Behavioral Ecology* **16**, 906–913.

*BLAKE, C.A. & GABOR, C.R. (2014) Effect of prey personality depends on predator species.
Behavioral Ecology **25**, 871–877.

*BLANCKENHORN, W.U. (1991a) Fitness consequences of food-based territoriality in water
striders, *Gerris remigis*. *Animal Behaviour* **42**, 147–149.

*BLANCKENHORN, W.V. (1991b) Fitness consequences of foraging success in water striders
(*Gerris remigis*; Heteroptera: Gerridae). *Behavioral Ecology* **2**, 46–55.

*BLIGHT, O., VILLALTA, I., CERDÁ, X. & BOULAY, R. (2016) Personality traits are associated
with colony productivity in the gypsy ant *Aphaenogaster senilis*. *Behavioral Ecology and
Sociobiology* **70**, 2203–2209.

*BOLNICK, D.I. & ARAÚJO, M.S. (2011) Partitioning the relative fitness effects of diet and
trophic morphology in the threespine stickleback. *Evolutionary Ecology Research* **13**, 439–459.

*BOLNICK, D.I. & PAULL, J.S. (2009) Morphological and dietary differences between
individuals are weakly but positively correlated within a population of threespine stickleback.
Evolutionary Ecology Research **11**, 1217–1233.

*BOON, A.K., RÉALE, D. & BOUTIN, S. (2008) Personality, habitat use, and their consequences
for survival in North American red squirrels *Tamiasciurus hudsonicus*. *Oikos* **117**, 1321–1328.

*BORATYŃSKI, Z., KOSKELA, E., MAPPES, T. & SCHRODERUS, E. (2013) Quantitative genetics
and fitness effects of basal metabolism. *Evolutionary Ecology* **27**, 301–314.

*BORATYŃSKI, Z. & KOTEJA, P. (2010) Sexual and natural selection on body mass and
metabolic rates in free-living bank voles: Selection on metabolic rate. *Functional Ecology* **24**,
1252–1261.

*BOTH, C., DINGEMANSE, N.J., DRENT, P.J. & TINBERGEN, J.M. (2005) Pairs of extreme avian
personalities have highest reproductive success. *Journal of Animal Ecology* **74**, 667–674.

*BOULTON, K., WALLING, C.A., GRIMMER, A.J., ROSENTHAL, G.G. & WILSON, A.J. (2018)
Phenotypic and genetic integration of personality and growth under competition in the
sheepshead swordtail, *Xiphophorus birchmanni*. *Evolution* **72**, 187–201.

*BREMNER-HARRISON, S., PRODOHL, P.A. & ELWOOD, R.W. (2004) Behavioural trait
assessment as a release criterion: boldness predicts early death in a reintroduction programme
of captive-bred swift fox (*Vulpes velox*). *Animal Conservation* **7**, 313–320.

*BRODIN, T. & JOHANSSON, F. (2004) Conflicting selection pressures on the growth/predation-
risk trade-off in a damselfly. *Ecology* **85**, 2927–2932.

*CALSBEEK, R. (2009) Sex-specific adult dispersal and its selective consequences in the brown
anole, *Anolis sagrei*. *Journal of Animal Ecology* **78**, 617–624.

*CARLSTEAD, K., MELLEN, J. & KLEIMAN, D.G. (1999) Black rhinoceros (*Diceros bicornis*) in
U.S. zoos: I. Individual behavior profiles and their relationship to breeding success. *Zoo Biology*
18, 17–34.

*CARTER, A.J., GOLDIZEN, A.W. & TROMP, S.A. (2010) Agamas exhibit behavioral syndromes:
bolder males bask and feed more but may suffer higher predation. *Behavioral Ecology* **21**, 655–
661.

*CHISM, J. & ROGERS, W. (2010) Male Competition, Mating Success and Female Choice in a
Seasonally Breeding Primate (*Erythrocebus patas*). *Ethology* **103**, 109–126.

*CLOBERT, J., OPPLIGER, A., SORCI, G., ERNANDE, B., SWALLOW, J.G. & GARLAND, T. (2000)
Trade-offs in phenotypic traits: endurance at birth, growth, survival, predation and
susceptibility to parasitism in a lizard, *Lacerta vivipara*: *Trade-offs in phenotypic traits*.
Functional Ecology **14**, 675–684.

*COLE, E.F. & QUINN, J.L. (2012) Personality and problem-solving performance explain
competitive ability in the wild. *Proceedings of the Royal Society B: Biological Sciences* **279**,
1168–1175.

*CORNWELL, T.O., MCCARTHY, I.D. & BIRO, P.A. (2020) Integration of physiology, behaviour
and life history traits: personality and pace of life in a marine gastropod. *Animal Behaviour* **163**,
155–162.

*COTE, J., DREISS, A. & CLOBERT, J. (2008) Social personality trait and fitness. *Proceedings of*
the Royal Society B: Biological Sciences **275**, 2851–2858.

*CUCHEROUSSET, J., ACOU, A., BLANCHET, S., BRITTON, J.R., BEAUMONT, W.R.C. & GOZLAN,
R.E. (2011) Fitness consequences of individual specialisation in resource use and trophic
morphology in European eels. *Oecologia* **167**, 75–84.

*CUTTS, C.J., ADAMS, C.E. & CAMPBELL, A. (2001) Stability of physiological and behavioural
determinants of performance in Arctic char (*Salvelinus alpinus*). *Canadian Journal of Fisheries*
and Aquatic Sciences **58**, 961–968.

*DAVID, M., AUCLAIR, Y. & CÉZILLY, F. (2011) Personality predicts social dominance in
female zebra finches, *Taeniopygia guttata*, in a feeding context. *Animal Behaviour* **81**, 219–
224.

*DAVID, M., AUCLAIR, Y., GIRALDEAU, L.-A. & CÉZILLY, F. (2012) Personality and body
condition have additive effects on motivation to feed in Zebra Finches *Taeniopygia guttata*:
Personality and feeding motivation. *Ibis* **154**, 372–378.

*DAY, T. & MCPHAIL, J.D. (1996) The effect of behavioural and morphological plasticity on
foraging efficiency in the threespine stickleback (*Gasterosteus sp.*). *Oecologia* **108**, 380–388.

*DENOEL, M. (2004) Feeding Performance in Heterochronic Alpine Newts is Consistent with
Trophic Niche and Maintenance of Polymorphism. *Ethology* **110**, 127–136.

*DERTING, T.L. & MCCLURE, P.A. (1989) Intraspecific Variation in Metabolic Rate and Its
Relationship with Productivity in the Cotton Rat, *Sigmodon hispidus*. *Journal of Mammalogy*
70, 520–531.

*DES ROCHES, S., SHURIN, J.B., SCHLUTER, D. & HARMON, L.J. (2013) Ecological and
Evolutionary Effects of Stickleback on Community Structure. *PLoS ONE* **8**, e59644.

*DEWSBURY, D.A. (1984) Aggression, Copulation, and Differential Reproduction of Deer Mice
(*Peromyscus Maniculatus*) in a Semi-Natural Enclosure. *Behaviour* **91**, 1–23.

*DIAZ PAULI, B., EDELINE, E. & EVANGELISTA, C. (2020) Ecosystem consequences of multi-
trait response to environmental changes in Japanese medaka, *Oryzias latipes*. *Conservation*
Physiology **8**, coaa011.

*DINGEMANSE, N.J., BOTH, C., DRENT, P.J. & TINBERGEN, J.M. (2004) Fitness consequences
of avian personalities in a fluctuating environment. *Proceedings of the Royal Society of London.*
Series B: Biological Sciences **271**, 847–852.

*DUGATKIN, L.A. (1992) Tendency to inspect predators predicts mortality risk in the guppy
(*Poecilia reticulata*). *Behavioral Ecology* **3**, 124–127.

*EHLINGER, T.J. & WILSON, D.S. (1988) Complex foraging polymorphism in bluegill sunfish.
Proceedings of the National Academy of Sciences **85**, 1878–1882.

*Einum, S., Fossen, E.I.F., Parry, V. & Pélabon, C. (2019) Genetic Variation in Metabolic Rate and Correlations with Other Energy Budget Components and Life History in Daphnia magna. *Evolutionary Biology* **46**, 170–178.

*Fernández-Reiriz, M.J., Irisarri, J. & Labarta, U. (2016) Flexibility of Physiological Traits Underlying Inter-Individual Growth Differences in Intertidal and Subtidal Mussels Mytilusgalloprovincialis. *PLOS ONE* **11**, e0148245.

*Fisher, D.N., James, A., Rodríguez-Muñoz, R. & Tregenza, T. (2015) Behaviour in captivity predicts some aspects of natural behaviour, but not others, in a wild cricket population. *Proceedings of the Royal Society B: Biological Sciences* **282**, 20150708.

*Foote, A.D., Newton, J., Piertney, S.B., Willerslev, E. & Gilbert, M.T.P. (2009) Ecological, morphological and genetic divergence of sympatric North Atlantic killer whale populations. *Molecular Ecology* **18**, 5207–5217.

*Formica, V.A., Wood, C.W., Larsen, W.B., Butterfield, R.E., Augat, M.E., Hougen, H.Y. & Brodie, E.D. (2012) Fitness consequences of social network position in a wild population of forked fungus beetles (*Bolitotherus cornutus*). *Journal of Evolutionary Biology* **25**, 130–137.

*Foster, W.C., Armstrong, C.M., Chism, G.T. & Pruitt, J.N. (2017) Smaller and bolder prey snails have higher survival in staged encounters with the sea star *Pisaster giganteus*. *Current Zoology*, zow116.

*Gagnon, M.-C. & Turgeon, J. (2011) Sexual conflict in Gerris gillettei (Insecta: Hemiptera): intraspecific intersexual correlated morphology and experimental assessment of behaviour and fitness: Sexual conflict in *G. gillettei* populations. *Journal of Evolutionary Biology* **24**, 1505–1516.

*Galeotti, P., Sacchi, R., Pellitteri-Rosa, D., Bellati, A., Cocca, W., Gentilli, A., Scali, S. & Fasola, M. (2013) Colour polymorphism and alternative breeding strategies: effects of parent's colour morph on fitness traits in the common wall lizard. *Evolutionary Biology* **40**, 385–394.

*Gangloff, E.J., Vleck, D. & Bronikowski, A.M. (2015) Developmental and Immediate Thermal Environments Shape Energetic Trade-Offs, Growth Efficiency, and Metabolic Rate in Divergent Life-History Ecotypes of the Garter Snake *Thamnophis elegans*. *Physiological and Biochemical Zoology* **88**, 550–563.

*Gifford, M.E., Clay, T.A. & Careau, V. (2014) Individual (Co)variation in Standard Metabolic Rate, Feeding Rate, and Exploratory Behavior in Wild-Caught Semiaquatic Salamanders. *Physiological and Biochemical Zoology* **87**, 384–396.

*Giménez, J.O. (2015) Determinants of phenotypic variation in the iberian wall lizard species complex (*Podarcis hispanicus*). Master thesis.

*Glon, M.G., Larson, E.R. & Pangle, K.L. (2016) Connecting laboratory behavior to field function through stable isotope analysis. *PeerJ* **4**, e1918.

*Griffen, B.D. & Mosblack, H. (2011) Predicting diet and consumption rate differences between and within species using gut ecomorphology: Predicting diet and consumption rate differences. *Journal of Animal Ecology* **80**, 854–863.

*Hromada, M., Kuczyński, L., Krištín, A. & Tryjanowski, P. (2003) Animals of different phenotype differentially utilise dietary niche - The case of the Great Grey Shrike *Lanius excubitor*. *Ornis Fennica* **80**, 71–78.

*Hsu, Y.-C., Shaner, P.-J., Chang, C.-I., Ke, L. & Kao, S.-J. (2014) Trophic niche width increases with bill-size variation in a generalist passerine: a test of niche variation hypothesis. *Journal of Animal Ecology* **83**, 450–459.

*HULTHÉN, K., CHAPMAN, B.B., NILSSON, P.A., HANSSON, L.-A., SKOV, C., BRODERSEN, J., VINTERSTARE, J. & BRÖNMARK, C. (2017) A predation cost to bold fish in the wild. *Scientific Reports* **7**, 1239.

*INGRAM, T., SVANBÄCK, R., KRAFT, N.J.B., KRATINA, P., SOUTHCOTT, L. & SCHLUTER, D. (2012) Intraguild predation drives evolutionary niche shift in threespine stickleback: evolutionary response to intraguild predation. *Evolution* **66**, 1819–1832.

*IOANNOU, C.C., PAYNE, M. & KRAUSE, J. (2008) Ecological consequences of the bold–shy continuum: the effect of predator boldness on prey risk. *Oecologia* **157**, 177.

*JABLONSZKY, M., SZÁSZ, E., KRENHARDT, K., MARKÓ, G., HEGYI, G., HERÉNYI, M., LACZI, M., NAGY, G., ROSIVALL, B., SZÖLLŐSI, E., TÖRÖK, J. & GARAMSZEGI, L.Z. (2018) Unravelling the relationships between life history, behaviour and condition under the pace-of-life syndromes hypothesis using long-term data from a wild bird. *Behavioral Ecology and Sociobiology* **72**, 52.

*JOHNSON, J.C. & SIH, A. (2005) Precopulatory sexual cannibalism in fishing spiders (*Dolomedes triton*): a role for behavioral syndromes. *Behavioral Ecology and Sociobiology* **58**, 390–396.

*JOLLES, J.W., MANICA, A. & BOOGERT, N.J. (2016) Food intake rates of inactive fish are positively linked to boldness in three-spined sticklebacks *Gasterosteus aculeatus*. *Journal of Fish Biology* **88**, 1661–1668.

*KAIN, M.P. & MCCOY, M.W. (2016) Anti-predator behavioral variation among *Physa acuta* in response to temporally fluctuating predation risk by Procambarus. *Behavioural Processes* **133**, 15–23.

*KARPESTAM, E. & FORSMAN, A. (2011) Dietary differences among colour morphs of pygmy grasshoppers revealed by behavioural experiments and stable isotopes. *Evolutionary Ecology Research* **13**, 461–477.

*KATANO, O. (2011) Effects of individual differences in foraging of pale chub on algal biomass through trophic cascades. *Environmental Biology of Fishes* **92**, 101–112.

*KAUSE, A., OSSIPOV, V., HAUKIOJA, E., LEMPA, K., HANHIMÄKI, S. & OSSIPOVA, S. (1999) Multiplicity of biochemical factors determining quality of growing birch leaves. *Oecologia* **120**, 102–112.

*KEISER, C.N., INGLEY, S.J., TOSCANO, B.J., SCHARF, I. & PRUITT, J.N. (2018) Habitat complexity dampens selection on prey activity level. *Ethology* **124**, 25–32.

*KEISER, C.N., SLYDER, J.B., CARSON, W.P. & PRUITT, J.N. (2015) Individual differences in predators but not producers mediate the magnitude of a trophic cascade. *Arthropod-Plant Interactions* **9**, 225–232.

*KERR, N.R. (2017) Links between personality and individual niche in the freshwater fish *Gobiomorphus cotidianus*. In pp. 1–76. Master Thesis, Otago, New Zealand.

*KHELIFA, R., ZEBSA, R., AMARI, H., MELLAL, M.K. & MAHDJOUB, H. (2019) Field estimates of fitness costs of the pace-of-life in an endangered damselfly. *Journal of Evolutionary Biology* **32**, 943–954.

*KOBLER, A., KLEFOTH, T., MEHNER, T. & ARLINGHAUS, R. (2009) Coexistence of behavioural types in an aquatic top predator: a response to resource limitation? *Oecologia* **161**, 837–847.

*KORHONEN, H., NIEMELÄ, P. & SIIRILÄ, P. (2001) Temperament and reproductive performance in farmed sable. *Agricultural Food Science Finland* **10**, 91–98.

*KORHONEN, H.T., JAUHIAINEN, L. & REKIL&AUML, T. (2002) Effect of temperament and behavioural reactions to the presence of a human during the pre-mating period on reproductive

performance in farmed mink (*Mustela vison*). *Canadian Journal of Animal Science* **82**, 275–282.

*Kralj-Fišer, S., Hebets, E.A. & Kuntner, M. (2017) Different patterns of behavioral variation across and within species of spiders with differing degrees of urbanization. *Behavioral Ecology and Sociobiology* **71**, 125.

*Kruuk, L.E.B., Clutton-Brock, T.H., Slate, J., Pemberton, J.M., Brotherstone, S. & Guinness, F.E. (2000) Heritability of fitness in a wild mammal population. *Proceedings of the National Academy of Sciences* **97**, 698–703.

*Lang, I., Paz-Vinas, I., Cucherousset, J. & Loot, G. (2021) Patterns and determinants of phenotypic variability within two invasive crayfish species. *Freshwater Biology* **66**, 1782–1798.

*Lapiedra, O., Schoener, T.W., Leal, M., Losos, J.B. & Kolbe, J.J. (2018) Predator-driven natural selection on risk-taking behavior in anole lizards. *Science* **360**, 1017–1020.

*Larivee, M. (2010) The fitness consequences of variation in resting metabolic rate in juvenile North American red squirrels (*Tamiasciurus hudsonicus*). University of Alberta Libraries.

*Lattanzio, M.S. & Miles, D.B. (2016) Trophic niche divergence among colour morphs that exhibit alternative mating tactics. *Royal Society Open Science* **3**, 150531.

*Laurila, A. & Kujasalo, J. (1999) Habitat duration, predation risk and phenotypic plasticity in common frog (*Rana temporaria*) tadpoles. *Journal of Animal Ecology* **68**, 1123–1132.

*Lavin, P.A. & McPhail, J.D. (1986) Adaptive Divergence of Trophic Phenotype among Freshwater Populations of the Threespine Stickleback (*Gasterosteus aculeatus*). *Canadian Journal of Fisheries and Aquatic Sciences* **43**, 2455–2463.

*Lepetz, V., Massot, M., Chaine, A.S. & Clobert, J. (2009) Climate warming and the evolution of morphotypes in a reptile. *Global Change Biology* **15**, 454–466.

*Luiz, O.J., Crook, D.A., Kennard, M.J., Olden, J.D., Saunders, T.M., Douglas, M.M., Wedd, D. & King, A.J. (2019) Does a bigger mouth make you fatter? Linking intraspecific gape variability to body condition of a tropical predatory fish. *Oecologia* **191**, 579–585.

*Maldonado, K., Pablo, S., Piriz, G., Bogdanovich, J.M., Nespolo, R.F. & Bozinovic, F. (2016) Is maximum food intake in endotherms constrained by net or factorial aerobic scope? lessons from the leaf-eared mouse. *Frontiers in physiology* **7**.

*Marshall, H.H., Sanderson, J.L., Mwanghuya, F., Businge, R., Kyabulima, S., Hares, M.C., Inzani, E., Kalema-Zikusoka, G., Mwesige, K., Thompson, F.J., Vitikainen, E.I.K. & Cant, M.A. (2016) Variable ecological conditions promote male helping by changing banded mongoose group composition. *Behavioral Ecology* **27**, 978–987.

*Maskrey, D.K., White, S.J., Wilson, A.J. & Houslay, T.M. (2018) Who dares does not always win: risk-averse rockpool prawns are better at controlling a limited food resource. *Animal Behaviour* **140**, 187–197.

*Matthews, B., Aebischer, T., Sullam, K.E., Lundsgaard-Hansen, B. & Seehausen, O. (2016) Experimental evidence of an eco-evolutionary feedback during adaptive divergence. *Current Biology* **26**, 483–489.

*Matthews, B., Marchinko, K.B., Bolnick, D.I. & Mazumder, A. (2010) Specialization of trophic position and habitat use by sticklebacks in an adaptive radiation. *Ecology* **91**, 1025–1034.

*McCollum, S.A. & Van Buskirk, J. (1996) Costs and benefits of a predator-induced polyphenism in the gray treefrog *Hyla chrysoscelis*. *Evolution* **50**, 583–593.

*MCPHEE, M.V. & QUINN, T.P. (1998) Factors affecting the duration of nest defense and
reproductive lifespan of female sockeye salmon, *Oncorhynchus nerka*. *Environmental Biology*
of Fishes **51**, 369–375.

*METCALFE, N.B., TAYLOR, A.C. & THORPE, J.E. (1995) Metabolic rate, social status and life-
history strategies in Atlantic salmon. *Animal Behaviour* **49**, 431–436.

*MØLLER, A.P. (2010) Brain size, head size and behaviour of a passerine bird. *Journal of*
Evolutionary Biology **23**, 625–635.

*MONCEAU, K., DECHAUME-MONCHARMONT, F.-X., MOREAU, J., LUCAS, C., CAPODURO, R.,
MOTREUIL, S. & MORET, Y. (2017) Personality, immune response and reproductive success: an
appraisal of the pace-of-life syndrome hypothesis. *Journal of Animal Ecology* **86**, 932–942.

*MONTIGLIO, P., WEY, T.W., CHANG, A.T., FOGARTY, S. & SIH, A. (2017) Correlational
selection on personality and social plasticity: morphology and social context determine
behavioural effects on mating success. *Journal of Animal Ecology* **86**, 213–226.

*MORALES, J.A., CARDOSO, D.G., DELLA LUCIA, T.M.C. & GUEDES, R.N.C. (2013) Weevil x
Insecticide: Does 'Personality' Matter? *PLoS ONE* **8**, e67283.

*NAKAYAMA, S., RAPP, T. & ARLINGHAUS, R. (2016) Fast–slow life history is correlated with
individual differences in movements and prey selection in an aquatic predator in the wild.
Journal of Animal Ecology.

*NANNINI, M.A., PARKOS, J. & WAHL, D.H. (2012) Do behavioral syndromes affect foraging
strategy and risk-taking in a juvenile fish predator? *Transactions of the American Fisheries*
Society **141**, 26–33.

*NIEMELÄ, P.T., LATTENKAMP, E.Z. & DINGEMANSE, N.J. (2015) Personality-related survival
and sampling bias in wild cricket nymphs. *Behavioral Ecology* **26**, 936–946.

*NIEMELÄ, P.T., NIEHOFF, P.P., GASPARINI, C., DINGEMANSE, N.J. & TUNI, C. (2019) Crickets
become behaviourally more stable when raised under higher temperatures. *Behavioral Ecology*
and Sociobiology **73**, 81.

*NOBLE, D.W.A., WECHMANN, K., KEOGH, J.S. & WHITING, M.J. (2013) Behavioral and
Morphological Traits Interact to Promote the Evolution of Alternative Reproductive Tactics in
a Lizard. *The American Naturalist* **182**, 726–742.

*NYQVIST, M.J., CUCHEROUSSET, J., GOZLAN, R.E. & BRITTON, J.R. (2018) Relationships
between individual movement, trophic position and growth of juvenile pike (*Esox lucius*).
Ecology of Freshwater Fish **27**, 398–407.

*OUYANG, J.Q., SHARP, P.J., DAWSON, A., QUETTING, M. & HAU, M. (2011) Hormone levels
predict individual differences in reproductive success in a passerine bird. *Proceedings of the*
Royal Society B: Biological Sciences **278**, 2537–2545.

*VAN OVERVELD, T., ADRIAENSEN, F. & MATTHYSEN, E. (2015) No evidence for correlational
selection on exploratory behaviour and natal dispersal in the great tit. *Evolutionary Ecology* **29**,
137–156.

*PALKOVACS, E.P. & POST, D.M. (2009) Experimental evidence that phenotypic divergence in
predators drives community divergence in prey. *Ecology* **90**, 300–305.

*PALKOVACS, E.P., WASSERMAN, B.A. & KINNISON, M.T. (2011) Eco-evolutionary trophic
dynamics: Loss of top predators drives trophic evolution and ecology of prey. *PLoS ONE* **6**,
e18879.

*PARSONS, K.J. & ROBINSON, B.W. (2007) Foraging performance of diet-induced morphotypes
in pumpkinseed sunfish (Lepomis gibbosus) favours resource polymorphism. *Journal of*
Evolutionary Biology **20**, 673–684.

*PATRICK, S.C. & WEIMERSKIRCH, H. (2014) Personality, Foraging and Fitness Consequences in a Long Lived Seabird. *PLoS ONE* **9**, e87269.

*PEIMAN, K.S., BIRNIE-GAUVIN, K., LARSEN, M.H., COLBORNE, S.F., GILMOUR, K.M., AARESTRUP, K., WILLMORE, W.G. & COOKE, S.J. (2017) Morphological, physiological and dietary covariation in migratory and resident adult brown trout (*Salmo trutta*). *Zoology* **123**, 79–90.

*PETITJEAN, Q., JEAN, S., CÔTE, J., LAMARINS, A., LEFRANC, M., SANTOS, R., PERRAULT, A., LAFFAILLE, P. & JACQUIN, L. (2020) Combined effects of temperature increase and immune challenge in two wild gudgeon populations. *Fish Physiology and Biochemistry* **46**, 157–176.

*PINTOR, L.M., MCGHEE, K.E., ROCHE, D.P. & BELL, A.M. (2014) Individual variation in foraging behavior reveals a trade-off between flexibility and performance of a top predator. *Behavioral Ecology and Sociobiology* **68**, 1711–1722.

*PINTOR, L.M., SIH, A. & BAUER, M.L. (2008) Differences in aggression, activity and boldness between native and introduced populations of an invasive crayfish. *Oikos* **117**, 1629–1636.

*PINTOR, L.M., SIH, A. & KERBY, J.L. (2009) Behavioral correlations provide a mechanism for explaining high invader densities and increased impacts on native prey. *Ecology* **90**, 581–587.

*PIQUET, J.C., LÓPEZ-DARIAS, M., VAN DER MAREL, A., NOGALES, M. & WATERMAN, J. (2018) Unraveling behavioral and pace-of-life syndromes in a reduced parasite and predation pressure context: personality and survival of the Barbary ground squirrel. *Behavioral Ecology and Sociobiology* **72**, 147.

*POST, D.M., PALKOVACS, E.P., SCHIELKE, E.G. & DODSON, S.I. (2008) Intraspecific variation in a predator affects community structure and cascading trophic interactions. *Ecology* **89**, 2019–2032.

*POWERS, D.R. & NAGY, K.A. (1988) Field Metabolic Rate and Food Consumption by Free-Living Anna's Hummingbirds (*Calypte anna*). *Physiological Zoology* **61**, 500–506.

*PRUITT, J.N., KEISER, C.N., BANKA, B.T., LIEDLE, J.S., BROOKS, A.J., SCHMITT, R.J. & HOLBROOK, S.J. (2018) Collective aggressiveness of an ecosystem engineer is associated with coral recovery. *Behavioral Ecology*.

*PRUITT, J.N., RIECHERT, S.E. & JONES, T.C. (2008) Behavioural syndromes and their fitness consequences in a socially polymorphic spider, *Anelosimus studiosus*. *Animal Behaviour* **76**, 871–879.

*QUEVEDO, M., SVANBÄCK, R. & EKLÖV, P. (2009) Intrapopulation niche partitioning in a generalist predator limits food web connectivity. *Ecology* **90**, 2263–2274.

*QUINN, J.L., PATRICK, S.C., BOUWHUIS, S., WILKIN, T.A. & SHELDON, B.C. (2009) Heterogeneous selection on a heritable temperament trait in a variable environment. *Journal of Animal Ecology* **78**, 1203–1215.

*RAFFARD, A., CUCHEROUSSET, J., MONTOYA, J.M., RICHARD, M., ACOCA-PIDOLLE, S., POÉSY, C., GARREAU, A., SANTOUL, F. & BLANCHET, S. (2021) Intraspecific diversity loss in a predator species alters prey community structure and ecosystem functions. *PLOS Biology* **19**, e3001145.

*RAFFARD, A., CUCHEROUSSET, J., SANTOUL, F., DI GESU, L. & BLANCHET, S. (2023) Climate and intraspecific variation in a consumer species drive ecosystem multifunctionality. *Oikos*.

*RAFFARD, A., LECERF, A., COTE, J., BUORO, M., LASSUS, R. & CUCHEROUSSET, J. (2017) The functional syndrome: linking individual trait variability to ecosystem functioning. *Proceedings of the Royal Society B: Biological Sciences* **284**, 20171893.

*RAFFARD, A., SANTOUL, F., BLANCHET, S. & CUCHEROUSSET, J. (2020) Linking intraspecific variability in trophic and functional niches along an environmental gradient. *Freshwater Biology* **65**, 1401–1411.

*RAINE, N.E. & CHITTKA, L. (2005) Colour preferences in relation to the foraging performance
and fitness of the bumblebee *Bombus Terrestris*. *Uludag Bee Journal* **5**, 145–150.

*RÉALE, D. & FESTA-BIANCHET, M. (2003) Predator-induced natural selection on temperament
in bighorn ewes. *Animal Behaviour* **65**, 463–470.

*RÉALE, D., GALLANT, B.Y., LEBLANC, M. & FESTA-BIANCHET, M. (2000) Consistency of
temperament in bighorn ewes and correlates with behaviour and life history. *Animal Behaviour*
60, 589–597.

*REID, D., ARMSTRONG, J.D. & METCALFE, N.B. (2011) Estimated standard metabolic rate
interacts with territory quality and density to determine the growth rates of juvenile Atlantic
salmon: SMR, density, territory and growth. *Functional Ecology* **25**, 1360–1367.

*REID, D., ARMSTRONG, J.D. & METCALFE, N.B. (2012) The performance advantage of a high
resting metabolic rate in juvenile salmon is habitat dependent: RMR, dominance, habitat and
growth in salmon. *Journal of Animal Ecology* **81**, 868–875.

*ROBINSON, B.W. (2000) Trade offs in Habitat-Specific Foraging Efficiency and the Nascent
Adaptive Divergence of Sticklebacks in Lakes. Behaviour. *Behaviour* **137**, 865–888.

*ROTA, T., JABIOL, J., CHAUVET, E. & LECERF, A. (2018) Phenotypic determinants of inter-
individual variability of litter consumption rate in a detritivore population. *Oikos* **127**, 1670–
1678.

*ROTA, T., JABIOL, J., LAMOTHE, S., LAMBRIGOT, D., CHAUVET, É. & LECERF, A. (2024)
Phenotypes of predator individuals underpin contrasting ecosystem effects. *Freshwater*
Biology. https://doi.org/10.1111/fwb.14242.

*RUDMAN, S.M., RODRIGUEZ-CABAL, M.A., STIER, A., SATO, T., HEAVYSIDE, J., EL-SABAAWI,
R.W. & CRUTSINGER, G.M. (2015) Adaptive genetic variation mediates bottom-up and top-
down control in an aquatic ecosystem. *Proceedings of the Royal Society B: Biological Sciences*
282, 20151234.

*RUDMAN, S.M. & SCHLUTER, D. (2016) Ecological Impacts of Reverse Speciation in
Threespine Stickleback. *Current Biology* **26**, 490–495.

*SAMPAIO, A.L.A., PAGOTTO, J.P.A. & GOULART, E. (2013) Relationships between
morphology, diet and spatial distribution: testing the effects of intra and interspecific
morphological variations on the patterns of resource use in two Neotropical Cichlids.
Neotropical Ichthyology **11**, 351–360.

*SANTICCHIA, F., DANTZER, B., VAN KESTEREN, F., PALME, R., MARTINOLI, A., FERRARI, N. &
WAUTERS, L.A. (2018) Stress in biological invasions: Introduced invasive grey squirrels
increase physiological stress in native Eurasian red squirrels. *Journal of Animal Ecology* **87**,
1342–1352.

*SANTOSTEFANO, F., WILSON, A.J., NIEMELÄ, P.T. & DINGEMANSE, N.J. (2017) Indirect genetic
effects: a key component of the genetic architecture of behaviour. *Scientific Reports* **7**, 10235.

*SARNO, R.J. & FRANKLIN, W.L. (1999) Maternal expenditure in the polygynous and
monomorphic guanaco: suckling behavior, reproductive effort, yearly variation, and influence
on juvenile survival. *Behavioral Ecology* **10**, 41–47.

*SCHAACK, S. & CHAPMAN, L.J. (2003) Interdemic variation in the African cyprinid *Barbus*
neumayeri : correlations among hypoxia, morphology, and feeding performance. *Canadian*
Journal of Zoology **81**, 430–440.

*SCHUETT, W., DALL, S.R.X., KLOESENER, M.H., BAEUMER, J., BEINLICH, F. & EGGERS, T.
(2015) Life-history trade-offs mediate 'personality' variation in two colour morphs of the pea
aphid, *Acyrthosiphon pisum*. *Journal of Animal Ecology* **84**, 90–101.

*SCHULTE-HOSTEDDE, A.I. & MILLAR, J.S. (2004) Intraspecific variation of testis size and
sperm length in the yellow-pine chipmunk (*Tamias amoenus*): implications for sperm
competition and reproductive success. *Behavioral Ecology and Sociobiology* **55**, 272–277.

*SHACKLETON, M.A., JENNIONS, M.D. & HUNT, J. (2005) Fighting success and attractiveness
as predictors of male mating success in the black field cricket, *Teleogryllus commodus*: the
effectiveness of no-choice tests. *Behavioral Ecology and Sociobiology* **58**, 1–8.

*SHORT, K.H. & PETREN, K. (2008) Boldness underlies foraging success of invasive
Lepidodactylus lugubris geckos in the human landscape. *Animal Behaviour* **76**, 429–437.

*SMITH, T.B. (1987) Bill size polymorphism and intraspecific niche utilization in an African
finch. *Nature* **329**, 717–719.

*SNOWBERG, L.K. & BOLNICK, D.I. (2008) Assortative mating by diet in a phenotypically
unimodal but ecologically variable population of stickleback. *The American Naturalist* **172**,
733–739.

*SNOWBERG, L.K., HENDRIX, K.M. & BOLNICK, D.I. (2015) Covarying variances: more
morphologically variable populations also exhibit more diet variation. *Oecologia* **178**, 89–101.

*SOARES, A.O. & SCHANDERL, D.C.H. (2001) Fitness of two phenotypes of *Harmonia axyridis*
(Coleoptera: Coccinellidae). **98**, 287–293.

*SONG, Z.-G. & WANG, D.-H. (2003) Metabolism and thermoregulation in the striped hamster
Cricetulus barabensis. *Journal of Thermal Biology* **28**, 509–514.

*SPEAKMAN, J.R., KRÓL, E. & JOHNSON, M.S. (2004) The functional significance of individual
variation in basal metabolic rate. *Physiological and Biochemical Zoology* **77**, 900–915.

*SPRITZER, M.D., MEIKLE, D.B. & SOLOMON, N.G. (2005) Female choice based on male spatial
ability and aggressiveness among meadow voles. *Animal Behaviour* **69**, 1121–1130.

*START, D. & GILBERT, B. (2017) Predator personality structures prey communities and trophic
cascades. *Ecology Letters* **20**, 366–374.

*START, D. & GILBERT, B. (2019) Trait variation across biological scales shapes community
structure and ecosystem function. *Ecology* **100**.

*STEINMEYER, C., MUELLER, J.C. & KEMPENAERS, B. (2013) Individual variation in sleep
behaviour in blue tits *Cyanistes caeruleus*: assortative mating and associations with fitness-
related traits. *Journal of Avian Biology* **44**, 159–168.

*SVANBACK, R. & EKLOV, P. (2004) Morphology in perch affects habitat specific feeding
efficiency. *Functional Ecology* **18**, 503–510.

*TOSCANO, B.J. & GRIFFEN, B.D. (2014) Trait-mediated functional responses: predator
behavioural type mediates prey consumption. *Journal of Animal Ecology* **83**, 1469–1477.

*VAN BUSKIRK, J. & MCCOLLUM, S.A. (2000) Influence of tail shape on tadpole swimming
performance. *Journal of Experimental Biology* **203**, 2149–2158.

*VAN BUSKIRK, J., MCCOLLUM, S.A. & WERNER, E.E. (1997) Natural selection for
environmentally induced phenotypes in tadpoles. *Evolution* **51**, 1983–1992.

*VARGAS, R., MACKENZIE, S. & REY, S. (2018) 'Love at first sight': The effect of personality
and colouration patterns in the reproductive success of zebrafish (*Danio rerio*). *PLOS ONE* **13**,
e0203320.

*VERCKEN, E., MASSOT, M., SINERVO, B. & CLOBERT, J. (2007) Colour variation and
alternative reproductive strategies in females of the common lizard *Lacerta vivipara*. *Journal
of Evolutionary Biology* **20**, 221–232.

*WALTON, M. (1988) Relationships among Metabolic, Locomotory, and Field Measures of Organismal Performance in the Fowler's Toad (*Bufo woodhousei fowleri*). *Physiological Zoology* **61**, 107–118.

*WARD, A.J.W., THOMAS, P., HART, P.J.B. & KRAUSE, J. (2004) Correlates of boldness in three-spined sticklebacks (*Gasterosteus aculeatus*). *Behavioral Ecology and Sociobiology* **55**, 561–568.

*WEBSTER, M.M., WARD, A.J.W. & HART, P.J.B. (2009) Individual boldness affects interspecific interactions in sticklebacks. *Behavioral Ecology and Sociobiology* **63**, 511–520.

*WELLBORN, G.A. (2000) Selection on a sexually dimorphic trait in ecotypes within the *Hyalella azteca* species complex (Amphipoda: Hyalellidae). *The American Midland Naturalist* **143**, 212–225.

*WHITE, J.R., MEEKAN, M.G., McCORMICK, M.I. & FERRARI, M.C.O. (2013) A comparison of measures of boldness and their relationships to survival in young fish. *PLoS ONE* **8**, e68900.

*WHITEMAN, H.H., WISSINGER, S.A. & BROWN, W.S. (1996) Growth and foraging consequences of facultative paedomorphosis in the tiger salamander, *Ambystoma tigrinum nebulosum*. *Evolutionary Ecology* **10**, 433–446.

*WIELEBNOWSKI, N.C. (1999) Behavioral differences as predictors of breeding status in captive cheetahs. *Zoo Biology* **18**, 335–349.

*WILSON, A.D.M., GODIN, J.-G.J. & WARD, A.J.W. (2010) Boldness and reproductive fitness correlates in the eastern mosquitofish, *Gambusia holbrooki*. *Ethology* **116**, 96–104.

*WILSON, D.S., COLEMAN, K., CLARK, A.B. & BIEDERMAN, L. (1993) Shy-bold continuum in pumpkinseed sunfish (*Lepomis gibbosus*): An ecological study of a psychological trait. *Journal of Comparative Psychology* **107**, 250–260.

*WRAY, M.K., MATTILA, H.R. & SEELEY, T.D. (2011) Collective personalities in honeybee colonies are linked to colony fitness. *Animal Behaviour* **81**, 559–568.

*YAMAMOTO, T., UEDA, H. & HIGASHI, S. (1998) Correlation among dominance status, metabolic rate and otolith size in masu salmon. *Journal of Fish Biology* **52**, 281–290.

*ZAVORKA, L., ALDVEN, D., NASLUND, J., HOJESJO, J. & JOHNSSON, J.I. (2015) Linking lab activity with growth and movement in the wild: explaining pace-of-life in a trout stream. *Behavioral Ecology* **26**, 877–884.

*ZÁVORKA, L., KOECK, B., CUCHEROUSSET, J., BRIJS, J., NÄSLUND, J., ALDVÉN, D., HÖJESJÖ, J., FLEMING, I.A. & JOHNSSON, J.I. (2017) Co-existence with non-native brook trout breaks down the integration of phenotypic traits in brown trout parr. *Functional Ecology* **31**, 1582–1591.

*ZENG, L.-Q., ZHANG, A.-J., KILLEN, S.S., CAO, Z.-D., WANG, Y.-X. & FU, S.-J. (2017) Standard metabolic rate predicts growth trajectory of juvenile crucian carp (*Carassius auratus*) under changing food availability. *Biology Open*, bio.025452.

*ZHAO, T., VILLÉGER, S., LEK, S. & CUCHEROUSSET, J. (2014) High intraspecific variability in the functional niche of a predator is associated with ontogenetic shift and individual specialization. *Ecology and Evolution* **4**, 4649–4657.

*ZUB, K., BOROWSKI, Z., SZAFRAŃSKA, P.A., WIECZOREK, M. & KONARZEWSKI, M. (2014) Lower body mass and higher metabolic rate enhance winter survival in root voles, *Microtus oeconomus*. *Biological Journal of the Linnean Society* **113**, 297–309.