Gene regulatory mechanisms guiding bifurcation of inhibitory and excitatory neuron lineages in the anterior brainstem

Sami Kilpinen*, Lassi Virtanen, Silvana Bodington-Celma, Amos Bonsdorff, Heidi Heliölä, Kaia Achim*, Juha Partanen*

Molecular and Integrative Biosciences Research Programme, Faculty of Biological and Environmental Sciences, University of Helsinki, Finland

* Corresponding author

Abstract

Selector transcription factors control choices of alternative cellular fates during development. The ventral rhombomere 1 of the embryonic brainstem produces neuronal precursors that can differentiate into either inhibitory GABAergic or excitatory glutamatergic neurons important for the control of behaviour. Transcription factors (TFs) *Tal1*, *Gata2* and *Gata3* are required for adopting the GABAergic neuronal identity and inhibiting the glutamatergic identity. Here, we asked how these selector TFs are activated and how they control the identity of the developing brainstem neurons. We addressed these questions by analysing chromatin accessibility at putative gene regulatory elements active during GABAergic and glutamatergic neuron lineage bifurcation, combined with studies of transcription factor expression and DNA-binding. Our results show that the *Tal1*, *Gata2* and *Gata3* genes are activated by highly similar mechanisms, with connections to regional patterning, neurogenic cell cycle exit and general course of neuronal differentiation. After activation, *Tal1*, *Gata2* and *Gata3* are linked by auto- and cross-regulation as well as regulatory interactions with transcription factors of the glutamatergic branch. Predicted targets of these selector TFs include genes expressed in GABAergic neurons, glutamatergic neurons, or both. Unlike genes specific to the glutamatergic branch, the genes expressed in GABAergic neurons appear to be under combinatorial control of *Tal1*, *Gata2* and *Gata3*. Understanding gene regulatory interactions affecting the anterior brainstem GABAergic and glutamatergic neuron differentiation may give genetic and mechanistic insights into neurodevelopmental traits and disorders.

Keywords: Neuron, GABAergic, glutamatergic, brainstem, pons, selector gene, transcription factor, Tal1, Gata, enhancer, transcriptome, chromatin accessibility, gene regulatory network

Introduction

The anterior brainstem is a complex neuronal mosaic controlling mood, motivation and movement. These neurons include diverse inhibitory GABAergic and excitatory glutamatergic neurons, some of which are associated with the dopaminergic and serotonergic systems.

An important source of these neurons is the ventrolateral rhombomere 1 (r1), where GABA- and glutamatergic neuron precursors are generated from a neuroepithelial region expressing a homeodomain TF *Nkx6-1*(Lahti et al., 2016; Waite et al., 2012). We here refer to these precursors as rhombencephalic V2 domain (rV2), reflecting its molecular similarity to the V2 domain in the developing spinal cord. The rV2 derived precursors give rise to GABAergic neurons in the posterior Substantia Nigra pars reticulata, Ventral Tegmental Area, Rostromedial Tegmental Nucleus and Ventral Tegmental Nucleus, all important for various aspects of behavioural control (Lahti et al., 2016; Morello et al., 2020a). A failure in the differentiation of the rV2 GABAergic neurons in the embryonic mouse brain, caused by mutation of the *Tal1* gene, results in robust behavioral changes in postnatal mice, including hyperactivity, impulsivity, hypersensitivity to reward, and learning deficits (Morello et al., 2020b).

The *Nkx6-1* positive proliferative progenitors of rV2 give rise to postmitotic V2b-like GABAergic precursors expressing selector genes *Tal1*, *Gata2* and *Gata3*, and V2a-like glutamatergic precursors expressing *Vsx1* and *Vsx2* (Achim et al., 2014; Joshi et al., 2009; Karunaratne et al., 2002a; Muroyama et al., 2005). In the absence of *Tal1*, *Gata2* and *Gata3* function, differentiation of all the rV2 precursors is directed to glutamatergic neuron types (Achim et al., 2012; Lahti et al., 2016). *Tal1*, *Gata2* and *Gata3* also retain their expression in the mature GABAergic neurons. Thus, *Tal1*, *Gata2* and *Gata3* TFs operate as fate-selectors, responsible for the induction and maintenance of the GABAergic identity in the anterior ventrolateral brainstem.

It has remained unclear how the selector TFs are activated during development and how they guide neuronal differentiation in the anterior brainstem. To address these questions, we have characterized the accessibility and TF binding in putative gene regulatory elements and correlated these events with changes in gene expression during differentiation of GABAergic and glutamatergic neurons in the anterior brainstem. Our results suggest that the GABAergic selector TF genes are activated by interconnected mechanisms reflecting developmental regionalization and timing, and that their functions converge to promote GABAergic and suppress glutamatergic neuron differentiation.

Materials and Methods

Data and code availability

The E12.5 mouse rhombomere 1 scRNAseq and E12.5 scATACseq reads generated in our study are deposited in SRA. The BioProject IDs of the E12.5 R1 scRNAseq data are: GEO GSE157963; BioProject PRJNA663556 E12.5 R1 samples SAMN16134208, SAMN16134206, SAMN16134205, SAMN16134207; and E12.5 R1 scATACseq: BioProject PRJNA929317, samples SAMN32957134, SAMN32957135.

84 All original code with used parameter values will be deposited in GitHub at
85 (https://github.com/ComputationalNeurogenetics/rV2_bifurcation)
86 Any additional information required to reanalyze the data reported in this paper is available
87 upon request.
88
The exact cell number, read yields and read quality for each sample are found in
(Supplementary Methods 1 and
https://github.com/ComputationalNeurogenetics/rV2_bifurcation/seq_quality_reports). Batch
effect was evaluated not to be meaningful based on the distribution of cells from biological
replicates in the UMAPs of both r1 E12.5 scRNA-seq and scATAC-seq (Supplementary
Methods 1).
95

96 scRNAseq samples
97 scRNAseq sample preparation is described in detail in (Morello et al., 2020a). The GEO and
98 BioProject IDs of the data are listed above.
99

100 Processing of scRNAseq data
101 scRNAseq data processing per brain region is described in detail in
102 (https://github.com/ComputationalNeurogenetics/rV2_bifurcation) with main steps outlined
103 below.
104

scRNAseq data was read into *Seurat* objects from 10x *filtered_feature_bc* matrix by using
Ensembl gene IDs as features. Replicates per each brain region, or all replicates in the later
analysis steps were merged as *Seurat* objects. After merging, quality control (QC) was
performed by calculating *mt-percentage and filtering based on nFeature_RNA* and
percent.mt. Data was then log-normalized, top 3000 most variable features were detected,
finally data was scaled and *mt-percentage*, *nFeature_RNA*, *nCount_RNA* were set to be
regressed out.
112

113 Data was then run through PCA using n variable features, and *Seurat RunUMAP* with
114 selected top PCA dimensions was used to generate Uniform Manifold Approximation and
115 Projection (UMAP) for further analysis. *Seurat* functions *FindNeighbors* and *FindClusters*
116 were applied for neighborhood and community detection-based clustering.
117

scATACseq samples
Embryonic brains were isolated from E12.5 wildtype mouse embryos (n=2), and ventral R1
pieces (E12.5) were separated. Nuclei were isolated using the 10X Genomics Assay for
Transposase Accessible Chromatin Nuclei Extraction from fresh tissues protocol CG000212.
Chromium Single Cell Next GEM ATAC Solution (10X Genomics, 1000162) was used for
the capture of nuclei and the accessible chromatin capture. In chip loading, the nuclei yield
was set at 5000 nuclei per sample. Chromium Single Cell Next GEM ATAC Reagent
chemistry v1.1 was used for the DNA library preparation. Sample libraries were sequenced
on Illumina NovaSeq 6000 system using read lengths: 50bp (Read 1N), 8bp (i7 Index), 16bp

(i5 Index) and 50bp (Read 2N). The reads were quality filtered and mapped against the *Mus*
musculus genome *mm10* using Cell Ranger ATAC v1.2.0 pipeline.
129
Processing scATACseq data
In this study we defined feature space as described previously (Kilpinen et al., 2023), except
that we used both E14.5 (Kilpinen et al., 2023) and E12.5 scATACseq samples to further
increase resolution to detect features from rare cell population and cellular clade cut height
was defined based on smallest clade so that none of the clades contained less than 100 cells
(h=8).
136
These jointly defined features were then used to process the E12.5 rhombomere 1
scATACseq data. Data was subjected to QC, binarization and LSA, steps described in more
detail in
(https://github.com/ComputationalNeurogenetics/rV2_bifurcation) largely following steps
previously described (Stuart et al., 2021, 2019). Similarly, scATACseq data was integrated
with scRNAseq using an approach already described (Stuart et al., 2019).
143
144   The UMAP projection based on scATACseq data was done with *Seurat* using SVD
145   components 2-30 for cells with a reliability score from scRNAseq integration >0.5 (Figure
146   2A). The first SVD component was omitted due to the association with read depth.
147   Neighborhood and community detection were made with Seurat functions
148   *FindNeighbors*(dims=2:30) and *FindClusters*(resolution=1, algorithm=4).
149
Subsetting and subclustering the rV2 lineage
The rV2 lineage was subset from clustered E12.5 R1 scRNAseq data (Supplementary Figure
1A) as well as from E12.5 R1 scATACseq data (Supplementary Figure 2A). Subsetting was
based on the expression levels of known marker genes (Ccnb1, Nkx6-1, Vsx2, Slc17a6, Tal1,
Gad1). scATACseq clusters 1, 6, 8, 12, 13, 19, 20, 24 and 26 were considered to consist of
rV2 lineage. Clusters 13 and 19 were considered to be progenitors and were named PRO1
and PRO2, respectively. Cluster 26 was considered to consist of common precursors and was
named CO. Clusters tentatively belonging to common precursors, the GABAergic (branches
1, 6, 20 and 24), and the glutamatergic branches (8 and 12) were further subjected to Seurat
FindSubClusters to reveal higher resolution cell types. This resulted in the splitting of CO
into clusters CO1 and CO2, the GABAergic branch into six clusters named GA1 – GA6, and
the glutamatergic branch into five clusters GL1 – GL5.
162
163   rV2 cluster correspondence between scATACseq and scRNAseq
164   To evaluate the overlap of marker genes between scRNAseq and scATACseq clusters
165   (Supplementary Table 1, 2), we performed a hypergeometric test and visualised the results
166   using a heatmap (Figure 1G).
167
168   The scRNAseq and scATACseq cluster markers were filtered with adj. p-value < 0.05 and
169   top 25 based on log2FC. An intersection of unique gene names was identified between

170   scRNAseq and scATACseq objects and its size, N, was used as the total gene pool for the
171   hypergeometric test.
172

VIA pseudotime analysis
For VIA (Stassen et al., 2021) based pseudotime analysis rV2-lineage embeddings
coordinates from dimensions 2:30, as well as cluster identities as colors, were passed from
Seurat object to Python based VIA. Calculations in both for scRNAseq and for scATACseq,
gene expression levels were used to calculate the pseudotime value. In the case of
scATACseq, these were the interpolated values from integration with the scRNAseq (Stuart
et al., 2019). In the case of scATACseq PRO1 was chosen as the root cell population, and for
scRNAseq cluster 22 was chosen as root cell population.
181

Sequence conservation
Conservation data is derived from UCSC database at
(https://hgdownload.cse.ucsc.edu/goldenPath/mm10/phastCons60way/), which contains
compressed phastCons scores for multiple alignments of 59 vertebrate genomes to the mouse
mm10 genome (Siepel et al., 2005). Nucleotide was considered conserved if phast-score was
at least 0.5 (range 0-1).
188

189   The scATACseq features and other selected regions of study were examined for conservation
190   level in comparison to the whole genome, using Fisher's exact test. Conservation was also
191   calculated for exon and intron regions within all chromosomes to demonstrate the accuracy of
192   this method.
193

194   For sequence conservation at the TFBS-motif locations, positional weight matrix (PWM) of
195   the motif was used to define weight of each nucleotide based on max probability at that
196   nucleotide location in PWM and that weight was used in calculation of the weighted mean of
197   conservation score over the nucleotides at TFBS locations.
198

199   TAD
200   Topologically associated domains (TADs) of the chromatin (Singh and Berger, 2021) were
201   used to limit the search for cCREs. We used *tadmap* (Singh and Berger, 2021) for TAD
202   definition.
203

Feature accessibility over pseudotime
Features within TAD of the selector gene were considered to be candidate cis-regulatory
element (cCRE) if they had linkage correlation with expression of the gene with
abs(zscore)>2 and p-value <0.05. Accessibility changes of cCREs were visualized as spline
smoothed heatmaps over the pseudotime. Spline smoothing was calculated over entire rV2
cell groups in order of (PRO1-2, CO1-2, GA1-2, GA3-4, GA5-6, GL1-2, GL3-4, GL5).
210

 ### TF-footprinting from scATACseq reads
TF-footprinting of scATACseq data was performed by using TOBIAS-framework (Bentsen
et al., 2020) with Snakemake pipeline. As motif collection we used Hocomoco version 12
(Vorontsov et al., 2023). The motif for INSM1 was added to the Hocomoco v12 collection
from Hocomoco v11 collection as it was seen relevant in earlier research. Suitability of
mixing motifs from the two versions of the Hocomoco was checked in correspondence with
Hocomoco team.
218
For the purpose of TF-footprinting we combined cell groups as follows to increase signal to
noise ratio: PRO1 and PRO2 (PRO1-2), CO1 and CO2 (CO1-2), GA1 and GA2 (GA1-2),
GA3 and GA4 (GA3-4), GA5 and GA6 (GA5-6), GL1 and GL2 (GL1-2), GL3 and GL4
(GL3-4). GL5 was kept as is. E12R1 BAM files were merged, and PCR duplicates removed
with Picard (https://broadinstitute.github.io/picard/). BAM file was then split into BAM file
per combined cell group with Sinto (https://timoast.github.io/sinto/).
225
226 TOBIAS was run with parameters given in
227 (https://github.com/ComputationalNeurogenetics/rV2_bifurcation) and the cell group and
228 genomic location specific footprint results were imported into an SQLite database (total of
229 49,390,084 records).
230
231 Footprinting data was visualized through R ggplot based dotplots per selected features, with
232 filtering for TFBS-motifs having >0.5 weighted average conservation value, accessibility of
233 the feature > 0.06 and expression of the TF in question > 1.2 (log1p) in any of the PRO1-2,
234 CO1-2, GA1-2 or GL1-2 groups.
235

RNAscope® *in situ* hybridization
E12.5 wildtype mouse embryos were dissected and fixed in 4% paraformaldehyde (PFA;
Sigma-Aldrich, Cat#P6148) in 1× PBS for 36 hours at room temperature. The fixed brains
were embedded in paraffin (Histosec™ Pastilles (without DMSO, Cat#115161)), and
sectioned on Leica microtome (Leica RM2255) at 5 µm. RNAscope® Multiplex Fluorescent
Reagent Kit v2 (ACD, Cat#323270) was used to detect RNA transcripts. The paraffin
sections were deparaffinized in xylene and ethanol series, treated with RNAscope®
Hydrogen Peroxide (ACD, Cat#322381) for 10 minutes, and subjected to antigen retrieval in
1× Target Retrieval Solution (ACD, Cat#322000) for 15 minutes at 99–102°C. After rinsing
and dehydration, hydrophobic barriers were drawn using an ImmEdge Pen (ACD,
Cat#310018), and slides were dried for 30 minutes or overnight. Protease Plus (ACD,
Cat#322381) was applied for 30 minutes at 40°C, followed by hybridization of probes at
40°C for 2 hours. Probes included mouse Ebf1 (ACD, Cat#433411, C2), E2f1 (ACD,
Cat#431971, C1), Insm1 (ACD, Cat#43062, C1), Sox4 (ACD, Cat#471381, C2), Tal1 (ACD,
Cat#428221, C4), and Tead2 (ACD, Cat#42028, C3). Positive (ACD, Cat#320881) and
negative (ACD, Cat#320871) control probes were included. C1 probes were used undiluted;
C2, C3, and C4 probes were diluted 1:50 in C1 probe or Probe Diluent (ACD, Cat#300041).
Slides were washed in Wash Buffer (ACD, Cat#310091) and stored in 5× saline-sodium
citrate (SSC) buffer overnight. Amplification was conducted using AMP1, AMP2, and AMP3

reagents (30 min, 30 min, 15 min) at 40°C. Signal detection was performed with HRP
channels (C1, C2, C3) and TSA Vivid Fluorophores: TSA 520 (ACD, Cat#323271),TSA 570
(ACD, Cat#323272), and TSA 650 (ACD, Cat#323273), diluted 1:1500 in RNAscope®
Multiplex TSA Buffer (ACD, Cat#322809). Each HRP step included blocking and washing.
Slides were counterstained with DAPI (30 seconds), mounted with ProLong Gold Antifade
Reagent (Thermo Fisher, Cat#P36934), and imaged using a Zeiss AxioImager 2 microscope
with Apotome 2 structured illumination. Images were processed in ImageJ, and figures
prepared in Adobe Illustrator CC 2020. Slides were stored at 4°C and imaged within two
weeks.

RNAscope image segmentation and downstream analysis
We trained a Stardist (Schmidt et al., 2018; Weigert et al., 2020) model to segment the
RNAScope images, for this purpose we manually painted cell nucleus onto 10 representative
microscopy images of E12.5 rV2 area based on DAPI channel.
268
The trained model was then used to segment the analysis images and to return ROIs as
ImageJ ROI coordinates. Regions of interest (ROIs) were imported into R for analysis.
Fluorescence signals were quantified for three channels in each image. The boundary of the
ventricular zone (VZ) was defined manually using three points, and calculating circle
boundary passing these points (Supplementary Figure 3), and the distances of individual cells
from the VZ were calculated in units of average cell diameter (avg. across all images).
Fluorescence signals were filtered to remove background noise using a quantile threshold of
0.25. Raw fluorescence intensity was measured as the number of fluorescence dots per cell.
To analyze patterns across biological replicates, fluorescence data from three independent
experiments were included in the same aggregate analysis. Fluorescence intensities were
normalized across replicates for each channel to account for variations in signal strength and
aligned to ensure uniform spatial scaling. Aggregated data were visualized by calculating a
smoothed sliding mean (window size=12) fluorescence trend for each channel, highlighting
consistent spatial patterns of *Tal1*, *Tead2*, and *E2f1* fluorescence intensities across biological
replicates.
284
CUT&Tag
Embryonic brains were isolated from E12.5 wildtype (ICR) mouse embryos, and ventral R1
pieces (E12.5) were separated. The ventral R1 pieces from the embryos of the same litter
were pooled and dissociated. The cells were then divided between the samples. Embryos
from different E12.5 mouse litters were used as biological replicates in CUT&Tag. The
samples were processed following the CUT&Tag-It Assay Kit protocol (Active Motif, cat #
53170). The following primary antibodies were used: rabbit anti-Gata2 (Abcam, ab109241),
1:80 rabbit anti-Gata3 (Boster, M00593), rabbit anti-Tal1 (Abcam, ab75739), rabbit anti-
Vsx2 (BioSite, 25825-1-AP-20), rabbit anti-Ebf1 (Boller et al., 2016) rabbit anti-Insm1
(BioSite, ASJ-IO4DE3-50), rabbit anti-Tead2 (Biorbyt, orb382464), rabbit anti-IgG (Cell
Signalling, 86652/66362) and rabbit anti-H3K4me3 (Cell Signalling, 86652/9751). All
primary antibodies were used at 1:80 dilution. Sample and sequencing library preparation
was done following the recommended protocols of the assay. The libraries were multiplexed

at 10 nM equimolar ratios and sequenced on NextSeq 500 or Aviti. 75 bp paired-end
sequences were obtained.
300
301   The demultiplexed reads were quality filtered and aligned against the *Mus musculus* genome
302   *mm10*, using the nf-core (Ewels et al., 2020) pipeline nf-core/cutandrun.
303

304   Analysis of the TFs interacting with the selector gene cCREs
305   To search for the regulators of selector genes, we first searched for TFs per each selector
306   gene per cell group with conditions as follows:

        1) The TF has a footprint in any of the cCREs linked to the selector gene with LinkPeaks abs(z-score)>2 and p-value<0.05, and
        2) The weighted mean of the conservation of the motif location is >0.5, and
        3) The TF is expressed > 1.2 (log1p) on average in the cell group in question.

Defining the common regulators between the selector genes was done by comparison of the
regulators found for individual selectors based on their intersects. Statistical significance
levels reported for individual regulators are defined as minimum p-value for cCRE to gene
linkage for each regulator per each selector gene and maximum p-value over all selectors in
the intersect. To assess whether the observed overlap in regulators among the three selector
genes was greater than expected by chance, we performed a permutation-based test. First, we
determined the actual set of regulators for each of the three selector genes and calculated the
size of their intersection. Next, to construct a null model, we repeatedly and randomly
selected three sets of regulators (of the same sizes as the original sets) from the complete
universe of candidate regulators, recalculating the intersection size each time. With 1 000 000
iterations we obtained a distribution of intersection sizes expected under random selection.
The empirical p-value was then estimated as the proportion of these randomized trials in
which the intersection size was equal to or larger than the observed intersection. If no
randomized trial matched or exceeded the observed intersection size, we reported the p-value
as less than an upper bound (e.g., $<1/(N+1)$), reflecting the rarity of the observed overlap
under the assumed null hypothesis.
327

Selector gene target analysis
To uncover general functional roles of the selector genes at the bifurcation point we
conducted a gene set enrichment (GSEA) (Mootha et al., 2003) analysis of the putative target
genes of *Tal1*, *Gata2*, *Gata3* and *Vsx2* over the axis of avg_log2FC of mouse genes in
between GA1-2 and GL1-2. For the GSEA axis only genes having an adjusted p-value <0.01
in the DEG test (Seurat FindMarkers with Wilcoxon) and an expression proportion of at least
0.1 in either group were taken into account.
335

336   Target genes were defined according to following conditions:
        1) The selector gene (TF) motif is associated with a footprint in either GA1-2 or GL1-2, and
        2) the motif situates in a scATACseq feature which is linked to the target gene with LinkPeaks abs(z-score)>2 and p-value<0.01, and
        3) the weighted mean of the conservation of the motif location is >0.5, and
        4) there is a CUT&Tag peak of the selector gene overlapping the feature.

342
As our GSEA axis should be considered relevant from both ends, one end signifying targets
overexpressed in GA1-2 and another in GL1-2, we additionally looked for leading edge
(traditionally called leading and trailing edges in GSEA analysis) genes from both ends. We
defined GA1-2 leading edge genes to be the ones before the GSEA enrichment score gains its
maximum, and GL1-2 leading edge genes to be ones after the GSEA enrichment score gains
its minimum.
349
350 Comparison of selector gene targets and cell type markers
351 The selector gene target list was subset, selecting the genes with exp_avg_GL1-2>0.5 (log1p)
352 OR exp_avg_GA1-2>0.5 (log1p). The resulting gene list was used as input in online Enrichr
353 tool (Xie et al., 2021). The results against CellMarker database (Hu et al., 2023) were
354 extracted.
355
Overlaps of the cell-type specific target genes of selectors.
Cell-type specific genes were defined as log2FC>0.5; exp_avg_GL1-2<0.5 (log1p);
exp_avg_GA1-2>0.5 (log1p) for GABAergic neuron specific genes and log2FC<-0.5;
exp_avg_GL1-2>0.5 (log1p); exp_avg_GA1-2<0.5 (log1p) for glutamatergic neuron specific
genes. The Tal1, Gata2 and Gata3 target gene lists were subset using these criteria. Overlaps
of the resulting gene lists were found using venn diagram app as shown on Figure 6.
362
ChromVAR analysis with Hocomoco motifs
ChromVAR analysis (Schep et al., 2017) was performed using Hocomoco (Vorontsov et al.,
2023) v12 motifs and an INSM1 motif lifted from Hocomoco v11 by using functions built
into Signac package. The motifs whose chromatin accessibility correlated with the RNA
expression of their respective genes (Spearman correlation > 0.5 and p-value < 0.01) were
identified, and their ChromVAR z-scores across rV2 lineage cell groups were aggregated by
mean to a level of TF and then visualised using ComplexHeatmap R package applying
clustering method "complete" and the clustering distance as "euclidean".
371
372 Results
373 *Transcriptomic dynamics and chromatin accessibility in differentiating rV2 GABAergic and*
374 *glutamatergic neurons*
375
To understand the sequence of gene expression changes that correlate with the bifurcation of
the rV2 GABAergic and glutamatergic cell lineages, we first studied the transcriptome and
the genome accessibility in developing rV2 neurons. We used data from the scRNAseq of the
ventral r1 region of E12.5 embryos (Morello et al., 2020a), where the rV2 lineages were
identified based on the expression of markers of common proliferative progenitors (*Ccnb1*,
Nkx6-1) and post-mitotic precursor branches of the rV2 by markers of GABAergic (*Tal1*,
Gad1) and glutamatergic neurons (*Nkx6-1*, *Vsx2*, *Slc17a6*) (Figure 1A, C, Supplementary
Table S1). Using pseudotime method VIA (Stassen et al., 2021), we found that the rV2 cells
were arranged on a pseudotemporal axes representing differentiating GABAergic or
glutamatergic lineages (Figure 1B). The GABAergic and glutamatergic selector TF genes

(*Tal1*, *Gata2*, *Gata3*, *Vsx1*, *Vsx2*, *Lhx4*) were strongly expressed in the post-mitotic
precursors at lineage bifurcation point and at the start of the lineage branches (Supplementary
Figure 1B).
389
390
To find the gene regulatory elements in the rV2 cell types, we applied scATACseq in E12.5
ventral r1 cells. We then clustered the r1 scATACseq cells based on their open chromatin
regions (called features below; Supplementary Figure 2A), followed by integration with the
E12.5 scRNAseq data (Stuart et al., 2019) that provides an estimate of the transcript levels for
scATACseq measured cells. From the scATACseq data, we readily identified the cell clusters
of the rV2 lineage, expressing TFs specific for GABAergic and glutamatergic cell types
(Supplementary Figure 2). We identified two progenitor cell groups (PRO1, PRO2), rV2
common precursors (CO), GABAergic neuron branch consisting of 4 cell groups and
glutamatergic branch of 2 cell groups (Supplementary Figure 2A). Further clustering of the
isolated rV2 cell groups yielded two groups of common precursors (CO1, CO2), six cell
groups in the GABAergic branch (GA1-GA6), and five cell groups in the glutamatergic
branch (GL1-GL5) (Supplementary figure 2B, Figure 1D, F,). The placement of the
scATACseq r1 cells in the pseudotemporal axis (Figure 1E) shows that, as expected, the early
state in pseudotime is assigned to cells expressing markers of proliferative progenitors and
early post-mitotic common precursors, while late pseudotime state cells express markers of
differentiating GABAergic and glutamatergic precursors. Cell clusters have consistently
increasing average pseudotime from PRO1 to the end of both GABAergic and glutamatergic
branches (Figure 1E). We next compared the scRNAseq and scATACseq cell cluster markers
across the rV2 lineages (top 25 marker genes per cluster, similarity testing using
hypergeometric distribution) (Supplementary Table 1, Supplementary Table 2, Figure 1G).
The similarity scores indicate discrete matches between the scATACseq and scRNAseq
clusters, with the exception of PRO1 and PRO2 largely missing a counterpart in the
scRNAseq clusters, as the rV2 progenitor cells did not clearly segregate in the scRNAseq
data (Supplementary Figure 1A, progenitor markers *Nes* and *Phgdh*) and were therefore
excluded from the rV2 lineage analyses. In conclusion, open chromatin features largely
reflected the gene expression in the ventral r1 and allowed placing the developing rV2
neurons in cell groups and in a pseudotemporal order.
418
419 *Chromatin accessibility profiles in genomic regions around selector genes in rV2-lineage*
420 *subclusters*
421
Tal1, Gata2 and Gata3 are required for GABAergic fate selection in rV2 and their expression
is activated in the GABAergic post-mitotic precursors, whereas Vsx2 is expressed in the
glutamatergic precursors soon after rV2 lineage bifurcation (Supplementary Figure S1B,
Supplementary Figure S2B) (Achim et al., 2013; Lahti et al., 2016; Morello et al., 2020a). To
understand the mechanisms behind activation of the expression of these lineage-specific TFs,
we analysed the accessibility of chromatin features in *Tal1, Gata2, Gata3* and *Vsx2* loci in
the rV2 cell groups. To find selector gene-associated chromatin features, we tested all

features within the topologically associating domains (TAD) around the transcription start sites (TSS) of the selector TF genes for the correlation between the feature accessibility and the target gene expression. We defined the features linked to the target gene with LinkPeaks z-score below -2 or above 2 and p-value <0.05 as candidate cis-regulatory elements (cCRE) of the gene (Supplementary Table 3) (Stuart et al., 2021)).

The *Tal1* locus, +/-50 kb of the TSS, contains five genes (*Cmpk1, Stil, Tal1, Pdzk1ip1* and *Cyp4x1*) and 20 scATAC features (Figure 2A). *Tal1* TAD contains 107 features. Some of the features at the *Tal1* locus were accessible in all the cell groups, including the progenitors (PRO1, PRO2, Figure 2A-B). These include features at the promoter regions of *Cmpk1* and *Stil* (Supplementary Table 3). Other features become accessible in both GABAergic and glutamatergic precursors after the bifurcation of the lineages, like the *Tal1* +77.4 kb cCRE located at the promoter region of the neuronally expressed *Cyp4x1* gene (Bylund et al., 2002). Accessibility of 15 out of the 107 features in the TAD containing the *Tal1* gene were linked to the *Tal1* expression (p-value < 0.05) (Supplementary Table 3, *Tal1*). Notably, the accessibility of *Tal1* cCREs at +0.7 and +40 kb correlated closely with the expression level of *Tal1* mRNA (Figure 2B), strongly increasing in the earliest GABAergic precursors (GA1) and maintained at a lower level in the more mature GABAergic precursor groups (GA2-GA6), as well as *Pdzk1ip1*, which is activated transiently in the earliest GABAergic precursors (GA1) (Figure 2A, RNA expression). Other features within the *Tal1* gene (+7 kb) as well as in more distal intergenic area (+15.1, +16, +16.4 and +23 kb cCREs) were also found to correlate with the activation of the *Tal1* mRNA expression (Figure 2B). Remarkably, the identified *Tal1* promoter feature (*Tal1* +0.7kb cCRE) contains *Tal1* midbrain and hindbrain-spinal cord enhancers characterized earlier (Bockamp et al., 1997; Sinclair et al., 1999). The *Tal1* +40 kb cCRE corresponds to the previously characterized enhancer region driving expression in the midbrain and hindbrain (Ogilvy et al., 2007). Enhancers at -3.8 kb and +15.1/16/16.4 kb of *Tal1* have been shown to drive *Tal1* expression in hematopoietic stem cells and erythrocytes (Göttgens et al., 2004) (Supplementary Table 4, Supplementary Figure 4A).

Similarly to *Tal1* locus, the TADs of *Gata2* and *Gata3* genes contained features where accessibility increased specifically in the GABAergic precursors, correlating with the activation of *Gata2* and *Gata3* mRNA expression (Supplementary Table 3, Supplementary Figure 5). Two *Gata2* cCREs, at -3.7 kb and +0 kb, are located in the proximal and promoter regions, which have been earlier shown to function as *Gata2* mid- and hindbrain expression enhancers (Nozawa et al., 2009), as well as critical *Gata2* autoregulation elements in hematopoietic cells (Kobayashi-Osaki et al., 2005) (Supplementary Table 4, Supplementary Figure 4B). We also identified an intronic *Gata2* cCRE at +11.6 kb that may mediate the regulation of *Gata2* expression in the spinal cord V2 interneurons by Gata2:Tal1 co-binding (Joshi et al., 2009). Four additional previously uncharacterized *Gata2* cCREs were identified downstream of the *Gata2* gene (+22, +25.3, +32.8 and 47.5 kb cCREs, Supplementary Figure 5A-B) and five more further downstream within the TAD (Supplementary Figure 5B, Supplementary Figure 4B, Supplementary Table 3).

At the *Gata3* locus, the GABAergic precursor –specific features included several intragenic
features (*Gata3* +11.1 kb, +17.2 kb, +18.8 kb cCREs), and five features upstream the gene at
-2.7 kb to -7.7 kb of *Gata3* TSS (Supplementary Table 3; Supplementary Figure 5C-D). The
Gata3 +0.6 kb, +2.2 kb and the five upstream cCREs locate within the previously identified
Gata3 CNS enhancer region (Lakshmanan et al., 1999), and two of those also match Gata3
lens expression enhancers (Martynova et al., 2018) (Supplementary Table 4, Supplementary
Figure 4C). The *Gata3* +11.1 kb cCRE corresponds to the *Gata3* proximal enhancer driving
expression in PNS regions (Lieuw et al., 1997). The *Gata3* +17.2 kb cCRE is located in the
intron 3-4 of *Gata3* gene and may have enhancer activity in spinal cord V2b interneurons
(Joshi et al., 2009) (Supplementary Table 4, Supplementary Figure 4C).
482
Importantly, accessibility of the cCREs of the GABAergic selector TF genes changed with
different kinetics during neuronal differentiation. For example, the accessibility of *Tal1* +23
kb and *Tal1* +40 kb cCREs was robustly detected before and at the onset of GABAergic
differentiation and declined thereafter (Figure 2B). The accessibility pattern of the *Tal1* +23
kb cCRE is particularly interesting. The feature is accessible at 3' position early, and gains
accessibility at 5' positions concomitant with GABAergic differentiation (Figure 2A,
accessibility). Detailed feature analysis later indicated the binding of Nkx6-1 and Ascl1 that
are expressed in the rV2 neuronal progenitors, at 3' positions, and binding of Insm1 and Tal1
TFs that are activated in early precursors, at 5' positions (Described below, see Figure 3C).
The accessibility of the Tal1 +0.7 kb cCRE largely reflected the *Tal1* mRNA expression
(Figure 2B). Similarly, the *Gata2* +22 kb, +25.3 kb and +32.8 kb cCREs were accessible in
the early precursors, before the opening of the *Gata2* -3.7 kb cCRE in GA1 and *Gata2* +0 kb
cCRE in GA3-GA6 (Supplementary Figure 5B). Thus, distinct regulatory elements of the
selector TFs may be responsible for the activation and for the maintenance of the lineage-
specific expression of selector TF gene.
498
Vsx2 gene expression was detected in the common precursors CO1 and increased in CO2 and
the glutamatergic precursors (GL1-GL5) after bifurcation of the rV2 lineages (Figure 2C,
RNA expression). At the *Vsx2* locus, an intronic feature (*Vsx2* +20.5 kb) and the distal
features at -61 kb and -21.3 kb of *Vsx2* TSS were robustly opened concomitant with its
expression activation, whereas accessibility of the *Vsx2* promoter region (*Vsx2* +1.1 kb
cCRE) and +27 kb cCRE increased later in the glutamatergic precursors (Figure 2D).
The information about the enhancers of Vsx2 brain expression is scarce, but we found that
our *Vsx2* r1 CREs at -18.7 kb, -21.3 kb, +1.1 kb and +20.5 kb match previously characterized
enhancers of *Vsx2* expression in photoreceptor cells and retinal bipolar neurons, and contain
putative autoregulatory Vsx2 biding sites occupied in developing spinal cord motor neuron
and interneuron lineages (Supplementary Table 4, Supplementary Figure 4D).
510
Developmentally important gene regulatory elements are often conserved. We therefore
asked if features detected by scATACseq could present more evolutionary conserved
sequences than random sequences of mouse genome. To that aim, we analyzed nucleotide
conservation in scATACseq features, using phastCons study data over 60 vertebrates from
UCSC database (Siepel et al., 2005). We calculated statistical enrichment of conserved

nucleotides within exons, introns and our scATACseq features (Supplementary Table 5).
Systematically, odds ratios per chromosome were highest for exons and lowest for introns,
odds ratios for features were between these two. All of the odds ratios showed a significant
difference from what one would expect if there were no relationship between the nucleotides
within features and conservation (p=< 2.43e-33, Fisher's Exact test, Supplementary Table 5).
The conservation of chromatin features is consistent with a putative role as regulatory sites.
522
In summary, we identified the genomic features that are linked to GABAergic and
glutamatergic fate selector genes, and where chromatin accessibility changes specifically
during rV2 neuron differentiation. Those features could represent regulatory elements of the
selector genes. Furthermore, literature research revealed that the previously characterized
enhancers, and especially the nervous system expression enhancers of *Tal1*, *Gata2*, *Gata3*
and *Vsx2* mostly are found within the ATAC-features identified here (63.6 - 100% of gene
enhancers overlapping ATAC-features, Supplementary Figure 6).
530
531 *Transcription factor activity at the putative regulatory elements of the rV2 cell fate selector*
532 *genes*
533
Next, we wanted to identify the TFs associated with the putative regulatory elements of *Tal1*,
Gata2, *Gata3* and *Vsx2* genes. TF binding to chromatin affects its accessibility, leaving a
footprint which can be detected in scATACseq data (Bentsen et al., 2020). We applied the
footprint analysis across the rV2 cell groups using the known TF binding sites (TFBS) from
HOCOMOCO collection (Vorontsov et al., 2023). As validation, we analyzed chromatin
interactions of selected candidate TFs in E12.5 r1 using CUT&Tag-sequencing (CUT&Tag)
(Kaya-Okur et al., 2019). As a complementary method, we used ChromVAR (Schep et al.,
2017) to calculate genome-wide accessibility score for each TFBS we used in the study.
While ChromVAR is based on scATACseq data, this analysis is independent of the
footprinting and thus provides a robust genome-wide validation of the most active TFBS.
544
Our TF footprinting analysis identified global TF activity patterns correlating with GABA-
and glutamatergic neuron differentiation (Supplementary Figure 7). To find the TFs that may
regulate the expression of *Tal1*, *Gata2*, *Gata3* and *Vsx2*, we identified the TFBSs at the
conserved genomic locations (weighted average conservation at TFBS > 0.5) within each
selector gene cCRE. The results of the footprint analysis are shown for selected features of
Tal1 (Figure 3A-D) and *Vsx2* (Supplementary Figure 8A,C-E), showing the TFs expressed
(average RNA expression > 1.2 (log1p)) in any of the four cell groups from PRO1-2, CO1-2,
GA1-2 and GL1-2. We observed both common and distinct patterns of TF binding in the
cCREs of *Tal1* and *Vsx2* in the rV2 common precursors and early GABA- and glutamatergic
cells. Both *Tal1* and *Vsx2* cCREs contain conserved GC-rich areas, E-boxes, TCF class
binding sites of TCF and Ebf TFs, regions of Sox TF binding elements 5'TTGT, NF I binding
elements TGGCA and binding sites for various homeodomain (HD) TF-s (Figure 3B-D,
Supplementary Figure 8C-E). The analysis of TFBSs at the footprint regions suggested that
each region may have affinity to several different TFs belonging to a TF class. For example,
four Sox TFs are expressed in early rV2 lineage cells (*Sox2*, *Sox3*, *Sox4* and *Sox11*), which all

may compete for the same binding sites in *Tal1* +0.7 kb cCRE or *Vsx2* -61 kb cCRE (Figure
3C, Supplementary Figure 8D).
To further understand the mechanisms of the activation of lineage-specific selector
expression, we asked which TFs commonly and specifically regulate the three GABAergic
selector TFs: *Tal1*, *Gata2*, and *Gata3*. Search for the common regulators of GABAergic fate
selectors revealed 21 TFs in CO1-2 and 22 TFs in the GA1-2 cell groups (Figure 3E).
Interestingly, the common regulators between the stages are overlapping partially, with the
TFs associated with the neurogenic cell cycle exit predominant in the common precursors
(CO) and the postmitotic differentiation and the autoregulation network TFs showing activity
later, in the GABAergic precursors.
We next asked which upstream regulators are common to both rV2 GABAergic (*Tal1*, *Gata2*,
Gata3) and glutamatergic (*Vsx1* and *Vsx2)* fate selector genes. The analysis of common
regulators revealed 12 TFs, including *Ascl1*, *E2f1*, *Nkx6-1* and *Vsx1* may regulate both
GABA- and glutamatergic selectors in the CO1-2 cell group (Figure 3E and Supplementary
Figure 8F). The distinct regulators of GABA-and glutamatergic fate selectors, aside the
lineage-specific autoregulatory functions of the selectors themselves, include *Lhx1* and *En2*
in GABAergic sublineage (*Lhx1* binding *Tal1* +0.7 kb feature, Figure 3B), and *Lhx4*, *Shox2*
and *Nhlh2* regulating *Vsx2* expression in the glutamatergic lineage (Supplementary Figure
8C-E). Footprints at Nkx6-1 TFBS were detected in the cCREs of all selectors in common
precursors, but after onset of differentiation become restricted to the cCREs of *Vsx2* and *Vsx1*
in the glutamatergic cells. Interestingly, the *Vsx2* -68.8 kb cCRE contains conserved GATA
TF binding sites (Supplementary Figure 8E). Conversely, *Vsx1* and *Vsx2* binding sites were
found in the cCREs of *Gata2* and *Gata3*, suggesting a mutual cross-regulation between the
GABA-and glutamatergic selector genes. Putative *Vsx1* binding sites were detected in all
three GABAergic fate selector genes, *Tal1*, *Gata2* and *Gata3*, in the rV2 common precursors
(Figure 3E, CO, CO1-2).
586
The proneural bHLH TFs appeared to contribute to the activation of the selector genes.
Interestingly, *Ascl1* and *Neurog2* TFBS are found at the cCREs of both *Tal1* (Figure 3C, D)
and *Vsx2* (Supplementary Figure 8C, D). In addition, *Vsx2* is regulated by *Neurog1* in the rV2
common precursors (Supplementary Figure 8C, D).
Interestingly, several cell cycle-associated TFs may contribute to the regulation of selector
gene expression. Those include *Sox2*, *Sox21*, *Insm1, Ebf1, Ebf3, and E2f1*. *E2f1* and *Ebf1-3*
binding sites are located within the GC-rich areas (Figure 3B-D, Supplementary Figure 8C-
E). The expression of *E2f1* declines after differentiation, while *Ebf1-3* are expressed in
common precursors and maintained in differentiating neurons of both GABA- and
glutamatergic sublineages (Figure 3B-D, Supplementary Figure 8C-E).
597
In addition to the footprint analysis, we analyzed the chromatin interactions of *Tal1, Gata2,*
Gata3, Vsx2, Ebf1 and *Insm1* in the E12.5 r1 by CUT&Tag (Supplementary Figure 9, Figure
3A, Supplementary Figure 10, Supplementary Figure 11). The binding of *Ebf1* and *Insm1*, as
well as *Tal1*, *Gata2*, and *Gata3* at the *Tal1*-linked features was confirmed by CUT&Tag
(Figure 3A, Supplementary Figures 9-11), consistent with the autoregulatory activity of the
GABAergic fate selector genes. Interestingly, in addition to the *Tal1* +0.7 kb and +40 kb

cCREs, we detected *Tal1*, *Gata2* and *Gata3* binding at the *Tal1* +23 kb cCRE, where
scATACseq-footprinting did not indicate binding at conserved sites (Figure 3A, C).
606
In summary, the regulatory feature analysis revealed several mechanisms that may contribute
to the timing of activation and the maintenance of selector TF expression. Firstly, we found
the binding sites of cell cycle progression or progenitor identity-associated TFs *E2f1* and
Insm1, proneural TFs *Ascl1, Neurog1, Neurog2, Neurod1* and *Neurod2*, as well as TFs
expressed after cell cycle exit, such as *Sox4* and *Ebf3* (*Coe3*) at the putative distal regulatory
elements. Conserved binding sites for other *Ebf* TFs, *Ebf1* and *Ebf2* are also found in the
cCREs of *Tal1* and *Gata2*, but not *Gata3*. In addition, the proximal elements and the
promoter regions of *Gata2* and *Tal1* genes contained footprints at conserved GATA::TAL
motif in the GABAergic neurons, indicating autoregulation of the selector TFs. Similarly,
conserved *Vsx1* and *Vsx2* motifs in *Vsx2* cCREs were associated with ATAC-footprints in the
glutamatergic neurons.
618
619

620   *Pattern of expression of the putative regulators of the selector TF expression*
621

Our analysis of TFBSs in the selector gene cCREs indicated that GABAergic selector TF
expression may be regulated by E2f1 and Ebf1, TFs involved in cell cycle progression and
neuronal differentiation. In addition, we found mediators of Hippo and Notch signalling
pathways, *Tead2* and *Insm1*, among the possible regulators of *Tal1*.
To test whether the *E2f1*, *Ebf1*, *Sox4*, *Insm1* and *Tead2* genes are expressed in the *Tal1*+
lineage precursors, we analysed their pattern of expression in the E12.5 mouse r1 by RNA *in*
situ hybridisation (RNAscope). In the embryonic brain, the progression of neuronal
differentiation follows the apical-basal axis, as the newly born precursors exit the ventricular
zone and migrate to intermediate and mantle zones. *Tal1* is expressed shortly after the cell
cycle exit. The expression of *Insm1, E2f1, Ebf1* and *Tead2* RNA overlapped with *Tal1* RNA
expression in the intermediate zone of the ventrolateral r1 at E12.5 (Figure 4A-B, E-F, I-J).
We quantified the fluorescence signal intensity along the apical-to-basal axis of the
neuroepithelium, showing that the maximum intensity of *Insm1, E2f1, Ebf1* and *Tead2* probe
signal is reached slightly closer to the apical surface than that of *Tal1* (Figure 4C, G, K). The
expression of *Sox4* RNA (Figure 4A-B) was detected in the neuronal cells at more basal
position in the neuroepithelium, where the differentiated neurons reside. The expression
patterns observed along the apical-basal axis in the E12.5 r1 neuroepithelium matched with
the expression levels detected by scRNAseq in the respective cell groups (Figure 4C-D, G-H,
K-L).
641
642   In conclusion, *Insm1, E2f1, Ebf1* and *Tead2* are expressed in the early rV2 precursors,
643   including the *Tal1* expressing GABAergic cell lineage. The expression of these TFs precedes
644   the expression of *Tal1* on the differentiation axis, consistent with a putative role in the
645   initiation of *Tal1* gene expression.
646
647

Regulatory signature of developing rV2 GABAergic neurons

To shed light on the genome-wide activity of TFs in the differentiating rV2 neurons, we applied ChromVAR (Schep et al., 2017). Consistent with the footprint analysis in cCREs, ChromVAR analysis indicated that accessible genomic regions in rV2 progenitor and precursor cells are enriched in *Sox*, *NeuroD1* and *Nfia/b/x* TF binding motifs (Figure 5A). In addition, *Nhlh1/2*, *NeuroD2*, *Bhlhe22*, *Ebf1-3*, *Pou3f4* and *Lhx5* motifs are found accessible in common precursors, shortly preceding the gain of accessibility in *Gata2/3* and *Tal1* motifs in GABAergic neurons (Figure 5A, square). We also analyzed the activity of *Tal1*, *Gata2* and *Gata3* in the rV2 neurons more closely. For those TFs, the accessibility at most of their TFBSs closely follows the TF RNA expression (Figure 5B-C). Interestingly, there are three TFBS motifs of *Tal1* in the HOCOMOCO v12 resource. Of these *Tal1* TFBSs, the activity pattern of the dual GATA:TAL binding motif (Figure 5C, TAL1.H12CORE.0.P.B) was most specific to the GABAergic lineage, similar to the GATA2 and GATA3 TFBSs (Figure 5C). The other TFBSs of *Tal1* showed lower activity (TAL1.H12CORE.1.P.B) or an earlier peak of activity in common precursors with a decline after differentiation (TAL1.H12CORE.2.P.B) (Figure 5C).

Targets of the GABAergic fate selectors in the developing rhombomere 1

We next aimed to describe the roles of GABAergic fate selector genes in the determination of the GABA- vs glutamatergic phenotype in rV2 cell lineages in more detail. Using the combined evidence of TF footprints, TFBS conservation and *in vivo* CUT&Tag experiments (Figure 6A), we identified 3948 target genes of *Tal1*, 780 target genes of Gata2, and 721 target genes of *Gata3* (Figure 6B). About half of the identified target genes are differentially expressed in GABA- or glutamatergic neurons (statistically significant differential expression between in GA1-2 and GL1-2 groups with p-value <0.01 in Wilcoxon's test). However, most of the identified target genes of *Tal1*, *Gata2*, and *Gata3* are lowly expressed in the GABA- and glutamatergic neurons, with only 30% of the genes being expressed above RNA avg(exp)>0.5 (log1p) in GA1-2 or GL1-2 cell groups (Supplementary Table 6). GSEA (Subramanian et al., 2005) showed that *Tal1* target genes are expressed in both GABA- and glutamatergic neurons (Figure 6C), while *Gata2* and *Gata3* target genes are overexpressed in GABAergic over glutamatergic precursors (Figure 6 H, M). To further characterize the identified targets, we compared the numbers of target features associated with the target gene by a positive or negative feature-to-gene expression link (Figure 6 D, I, N). The positive links outnumbered negative links by roughly factor of 2, which may reflect transcriptional activator function of the studied TFs, but also the complexity of gene regulation as there could be multiple, positive and negative links to each gene. We also calculated the amount of target genes considering their closest linked feature with distance bins of 0-5kb, 5-50kb and >50kb (Figure 6 E, J, O). The results showed largely expected distribution where category of longest links contained most target genes. This is likely due to the category being limited by TAD which are often large, yielding considerable amounts of distal regulatory elements for many genes. We next calculated the expression variability of the target genes, stratified by the distance to closest selector-bound linked feature. The target genes with proximal selector-

bound features (0-5 kb) show the most cell type-specific patterns (Figure 6 F, K, P). For
example, the variability of proximal feature-linked *Tal1* target genes in rV2 cell groups was
0.509 - 0.423, and 0.282 - 0.191 for distal feature-linked genes. Thus, regardless of the large
amounts of potential distal regulatory elements for many targets, the targets having a
proximal selector-bound regulatory element show most drastic changes in expression over the
rV2 lineage cell states. However, a few cell type-specific genes also appeared to be regulated
by distal features at >50 kb from TSS (*St18*, *Nkx6-1*, *Nhlh1*) or 5-50 kb from TSS (*Zfpm1*,
Lmo1) in the case of Tal1 targets (Supplementary Table 6). Finally, we also investigated cell
type associations of the target genes (exp>0.5 (log1p) in GA1-2 or GL1-2 cell groups) by
calculating scores against CellMarker 2024 database (Chen et al., 2013; Hu et al., 2023),
revealing clear enrichment of inhibitory interneuron markers among the Tal1, Gata2 and
Gata3 target genes (Figure 6 G, L, Q).
704
Finally, we analyzed the overlap of the cell type-specific target gene lists, aiming to find
genes regulated by all the GABAergic fate selectors (common target genes) or by any
selector TF specifically. No common target genes were found among the glutamatergic
neuron specific genes (Figure 6R). The genes commonly regulated by Tal1, Gata2 and Gata3
are exclusively overexpressed in the GABAergic lineage and contain genes comprising the
autoregulatory and GATA co-factor networks (*Gata2*, *Tal1*, *Zfpm1*, *Lmo1*). The GABAergic
phenotype (*Gad1*, *Gad2*, *Slc32a1*) is regulated by Gata2 and Gata3 (Figure 6S). Tal1 seems
to be involved in the activation or modulation of Notch signaling pathway genes (*Pdzk1ip1*,
Hes5). Interestingly, Tal1 also interacts with genes specific to rV2 glutamatergic cells (*Lhx4*,
Nkx6-1), likely repressing their expression in the developing GABAergic precursors.
However, as *Tal1* is expressed already in the common precursors of rV2 lineage, the
possibility of the activation of *Lhx4* by Tal1 is not excluded. We also suggest the cross-
regulation of GABA-and glutamatergic selector genes in the developing rV2 neuronal
precursors: Gata3 may repress the *Vsx2* expression in the GABAergic cells, while Vsx2 may
repress *Lmo1* in glutamatergic cells, establishing the initiated cell type-specific gene
expression programs (Figure 6T).
721
722
723 Discussion
724
Extensive studies during the past decades have identified several TFs instrumental for
neuronal fate decisions in the embryonic brain. How these TFs are regulated and how they
guide neuronal differentiation remains less understood. Differentiation of GABAergic and
glutamatergic neurons in the ventrolateral r1 is a good model system for TF guided cell
differentiation, as in this region the activation of Tal and Gata TFs is both spatially and
temporally controlled, and these TFs guide a binary selection between the two alternative cell
fates. In this work, we integrated analyses of gene expression, chromatin accessibility and TF
occupancy during bifurcation of GABAergic and glutamatergic lineages in the ventrolateral
r1. Our results suggest shared mechanisms connecting selector TF activation to

developmental processes and demonstrate combinatory functions of *Tal1*, *Gata2* and *Gata3*
TFs in the GABAergic fate selection.

Mechanisms behind selector TF activation. The homeodomain TF code, established by
intercellular communication in the neuronal progenitors, defines the competence of
differentiation into specific lineages. In the ventrolateral r1, the *Tal1*, *Gata2*, *Gata3* and *Vsx2*
TFs are activated in the region where the proliferative progenitor cells express the
homeodomain TF *Nkx6-1* (Achim et al., 2012; Lahti et al., 2016). Our result suggests that
Nkx6-1 directly regulates the spatial expression pattern of the selector TFs important for
acquisition of GABAergic and glutamatergic neuron phenotypes. *Nkx6-1* is not sufficient for
selector TF expression, which is activated only when the proliferative progenitor cells exit the
cell cycle and give rise to postmitotic precursors. Our results suggest that TFs *E2f1*, *Ebf1* and
Insm1 temporally control the selector TFs and connect their expression to cell-cycle exit. The
expression of *E2f1*, *Ebf1* and *Insm1* is not spatially restricted, but is upregulated in early
postmitotic precursors across the whole r1 neuroepithelium. Our results show that the
expression of these TFs precedes activation of *Tal1*. *E2f1* is best known for its function as a
positive regulator of cell cycle progression. However, there also is evidence suggesting a
more complex role for *E2f1* in cell cycle regulation. It has been suggested that the effect of
E2f1 is dependent on its level of expression, low levels promoting cell cycle progression but
elevated levels cell cycle exit (Radhakrishnan et al., 2004; Shats et al., 2017; Wang et al.,
2007). Our results are consistent with this observation. *E2f1* is expressed at a low level in the
proliferative progenitors, and its expression peaks concomitant to cell cycle exit. Also, *Ebf1*
and *Insm1* have earlier been associated with cell cycle exit and delamination of neuronal
progenitors, as well as differentiation and migration of post-mitotic precursors (Faedo et al.,
2017; Garcia-Dominguez et al., 2003; Monaghan et al., 2017; Tavano et al., 2018). Thus, the
mechanisms that guide cell cycle exit and other cell biological events of neurogenesis may
also contribute to activation of neuronal selector TF expression and adoption of the neuronal
phenotype.

Selector TFs directly autoregulate themselves and cross-regulate each other. Positive
feedback loops are commonly found in the gene regulatory networks involving
developmental selector TFs. The positive feedback loops can ensure stable cell fate decisions.
Tal and Gata TFs have been shown to autoregulate their own gene expression in
hematopoietic cells. Our results suggest that autoregulation also is a prominent feature of the
control of *Tal1*, *Gata2* and *Gata3* expression in the developing brain. At maintaining their
expression by positive feedback, these TFs appear to work together, potentially as a physical
complex involving Lmo1 (Ono et al., 1998), promoting the expression of its components. In
addition to the positive cross-regulation of *Tal1*, *Gata2* and *Gata3*, these TFs also interact
with the genes of glutamatergic lineage-specific TFs, such as *Vsx2* and *Lhx4*, expressed in the
glutamatergic precursors. We envision negative feedback allows cross-repression between the
Tal1, Gata2, Gata3, Lmo1 network and Vsx2 in the cells where both alternative fate selectors
are expressed. Auto- and cross-regulation would ensure robust patterns of selector TF
expression, also after the initial inputs for their activation have ceased.

TFs often have context-dependent activating or repressive effects on their targets, and our
data on chromatin occupancy cannot distinguish between the two possible outcomes.
However, according to our scRNAseq and earlier studies in the developing spinal cord V2
domain (Francius et al., 2016; Karunaratne et al., 2002b), the time of co-expression of
Gata2/3 and *Vsx1/2* is short. In the rV2, *Vsx1* is expressed in the rV2 common precursors,
where *Tal1*, *Gata2* and *Gata3* expression is still low. Parsimonious explanations would be the
negative cross-regulation and differential regulation of *Lmo1*. Furthermore, Vsx1 has been
shown to repress the expression of *Tal1* in the zebrafish spinal cord V2a interneurons (Zhang
et al., 2020).
787
Targets of the selector TFs. In addition to establishing cell type-specific patterns of selector
TF expression, the selector TFs are thought to be responsible for realization of neuronal
identities. Our results show that a combinatory function of the selector TFs is important for
adopting a GABAergic identity. This is supported by similar fate transformations observed in
Tal1 and *Gata2/Gata3* mutant embryos. Importantly, our results here highlight differences
between *Tal1*, *Gata2* and *Gata3* target genes. Most prominently, the activity of *Tal1* is not as
clearly biased towards genes expressed in the GABAergic branch as are the genes regulated
by *Gata2* and *Gata3*. In general, the *Tal1*, *Gata2* and *Gata3* selector TFs appear to regulate
not only GABAergic subtype–specific genes, but also genes expressed more broadly. This
suggests that the function of the fate selectors is not restricted to regulation of the neuronal
subtype properties such as the neurotransmitter phenotype, but they also participate in
enabling the overall cellular function.
800
Concluding remarks. Our work provides insights into the complex gene regulatory networks
regulating spatially and temporally controlled cell fate choices in the developing mouse
brainstem. In addition to addressing principles of cell differentiation, our work may also
contribute to understanding of neuropsychiatric traits and disorders. The anterior brainstem is
an important center for behavioral control, and many of the behavioral disorders have a
developmental basis. We have shown that, in mice, a failure in development of *Tal1*-
dependent neurons results in severe ADHD-like behavioral changes (Morello et al., 2020a). It
is therefore possible that variants in the genomic regions important for gene regulation during
brainstem neuron differentiation can lead to altered brain development and predisposition to
neuropsychiatric disorders.
811
812 *Caveats of the study*. Partly due to technical limitations, some of the TF – chromatin
813 interactions are inferred only based on ATAC-footprint. Some of the footprints may signify
814 the binding sites of several different TFs, although mostly of the same TF family, and the
815 exact interacting TF may not be identified. In addition, chromatin interaction is only
816 suggestive of a regulatory function.
817

818 ## Acknowledgements
819 We thank Outi Kostia for expert technical assistance and Iiris Chrons for help with the
820 bioinformatic analyses. We thank Arto Viitanen for his guidance with methods of image

segmentation. We acknowledge the services of FIMM Single Cell Analysis Unit, University
of Helsinki. This work was supported by grants from the Academy of Finland, the Sigrid
Juselius Foundation, and the Magnus Ehrnrooth Foundation.

Figures

Figure 1. Differentiation of the E12.5 mouse rV2 GABAergic and glutamatergic neurons based on transcriptome and chromatin accessibility

A. UMAP projection of rV2 lineage cells based on scRNA profiles.

B. Pseudotime (VIA) scores shown on the scRNA UMAP projection.

C. Expression of marker genes of progenitors and postmitotic precursors of GABAergic and glutamatergic neurons shown on the scRNA UMAP projection.

D. UMAP projection of rV2 lineage cells based on scATAC profiles.

E. Pseudotime (VIA) scores shown on the scATAC UMAP projection.

F. Expression of the marker genes of progenitors and postmitotic precursors of GABAergic and glutamatergic neurons shown on the scATAC UMAP projection (interpolated expression values).

G. Heatmap of scATAC and scRNA cluster similarity scores.

Figure 2. Accessibility of chromatin features in the *Tal1* and *Vsx2* loci during GABAergic and glutamatergic development.

A. Chromatin accessibility per cell group (normalized signal), Ensembl gene track (Genes), scATACseq features (Feat.), feature linkage to gene (Links with the LinkPeaks abs(zscore) >2) and nucleotide conservation (Cons.) within +/-50 kbp region around *Tal1* TSS. Violin plots on the right show the expression levels of *Tal1* and *Pdzk1ip1* per cell group.

B. Spline-smoothed z-score transformed heatmaps of chromatin accessibility at *Tal1*-linked scATACseq features (*Tal1* cCREs) in the single cells of rV2 GABAergic and glutamatergic lineages with RNA expression (sliding window mean(width=6) smoothed) of *Tal1*, *Gata2*,

Gad1 and *Slc17a6* as column covariable (top). Cells on the x-axis are first grouped per cell
group and then ordered by the pseudotime (bottom) within each group.
C. Same as in A, but for the *Vsx2* locus.
D. Accessibility of *Vsx2* cCREs in the rV2 GABAergic and glutamatergic lineages, shown as
in (B). The expression levels of *Vsx2*, *Tal1*, *Gad1*, and *Slc17a6* are shown above the
heatmaps.
857

858
Figure 3. **TF binding in putative regulatory elements of *Tal1* and the overlap between**
TFs interacting with other GABAergic fate selector genes.
A. ATAC features within +/-50 kb of the *Tal1* gene. *Tal1* cCREs are shown in blue.
CUT&Tag, CUT&Tag consensus peaks indicating Tal1, Gata2, Gata3, Vsx2, Ebf1, Insm1,
and Tead2 binding in the E12.5 mouse r1.
B-D. Footprinting based TF binding activity in the features at +1 kb, +23 kb and +40 kb of
the *Tal1* TSS. Feature linkage p to *Tal1* is indicated by asterisks. Footprint scores at
conserved TFBSs are shown for progenitors (PRO1-2), common precursors (CO1-2),
GABAergic precursors (GA1-2) and glutamatergic precursors (GL1-2). In each dotplot, the
strength of footprint at TFBS in the feature is shown as colour (Footprint score, average of
cell group) and the expression of the TF gene in dot size (log1p). Average feature
accessibility in cell groups is shown at the right. TFBS names (Hocomoco v12) are shown at
the top and the TF gene names (mouse) are shown at the bottom of the dotplot.

The red arrowhead in (B) indicates the conserved Gata2 TFBS at –37 bp position required for the neural expression of Tal1 (see also Supplementary Table 4).

E. Overlap of the TFs with footprints in the cCREs of *Tal1*, *Gata2*, and *Gata3* in the common precursors of rV2 lineages (CO1-2) and in the rV2 GABAergic precursors (GA1-2). Venn diagrams show the number of TFs with a footprint detected in *Tal1*, *Gata2*, or *Gata3* -linked features (Fp in cCRE = 1) and with their gene expression > 1.2 (log1p) in the analysed cell group. The TFs associated with the cCREs of all three selector genes in common precursors (CO1-2) and GABAergic precursors (GA1-2) are listed. Blue text: 19 TFs interact with cCREs of *Tal1*, *Gata2* and *Gata3* in the CO1-2 cells. Some of these TFs continue to be expressed and interact with the *Tal1*, *Gata2* and *Gata3* genes in the GA1-2 cells. 22 TFs interact with *Tal1*, *Gata2* and *Gata3* in GA1-2 cells. Green text, TFs found associated with two selector genes in CO1-2 (green in the Venn diagram of CO1-2) and associated with all three selector genes in the GA1-2. Black text, TFs co-regulating the selector TFs in GA1-2 and not expressed in the CO1-2 cells.

§ The probability of finding n overlapping genes considering all mouse genes equally is p<1e-6.

*,** The collective minimum statistical significance of feature to gene links for selector genes *Tal1, Gata2* and *Gata3* cCREs for the given TF is shown as: *p-value<0.05; **p-value<0.01 (with LinkPeaks z-score above 2 or below -2).

Figure 4. Expression patterns of TFs associated with Tal1 cCREs in the developing anterior brainstem.

A. RNAscope® *In Situ* Hybridization with *Insm1*, *Sox4*, and *Tal1* RNA probes on paraffin sections of E12.5 wild-type mouse embryos. Transversal sections of r1 are shown. Dashed

lines indicate the ventricular zone (VZ) border. Numbered boxes indicate areas shown on the
close-up images in (B), numbering order is from apical to basal. Scale bar: 50 μm.
B. Close-up images showing co-expression of *Insm1*, *Sox4*, and *Tal1* with DAPI.
C. Signal intensity of *Insm1*, *Sox4*, and *Tal1* probes along the apical-basal axis. Average
intensity line is based on three replicates. The 95% confidence interval of fluorescence
intensity is highlighted in gray. The y-axis shows average fluorescence intensity per cell, and
the x-axis shows the number of average cell diameters (μm) from the VZ.
D. Violin plots showing *Insm1*, *Sox4*, and *Tal1* mRNA expression detected by single-cell
RNA-seq in the rV2 lineage. Cell clusters are aligned by pseudotime order on x-axis,
showing first the differentiated cell clusters of GABAergic lineage, and then glutamatergic
lineage.
E. RNAscope® *In Situ* Hybridization with *E2f1*, *Tead2*, and *Tal1* RNA probes, images
shown and labelled as in (A). Transversal sections of E12.5 r1 are shown. Scale bar: 50 μm.
F. Close-up images showing co-expression of *E2f1*, *Tead2*, and *Tal1* with DAPI in areas
indicated in (E).
G. Signal intensity of *E2f1*, *Tead2*, and *Tal1* probes along the apical-basal axis, with the lines
showing the average of three replicates.
H. Violin plots showing *E2f1*, *Tead2*, and *Tal1* mRNA expression in rV2 neuronal
populations.
I. RNAscope® *In Situ* Hybridization with *Ebf1*, *Tead2*, and *Tal1* RNA probes, images shown
and labelled as in (A) and (E). Transversal sections of E12.5 r1 are shown. Scale bar: 50 μm.
J. Close-up images showing co-expression of *Ebf1*, *Tead2*, and *Tal1* with DAPI in areas
indicated in (I).
K. Signal intensity of *Ebf1*, *Tead2*, and *Tal1* probes along the apical-basal axis, with the lines
showing the average of three replicates.
L. Violin plots showing *Ebf1*, *Tead2*, and *Tal1* mRNA expression in rV2 neuronal
populations.
924

Figure 5: Dynamic accessibility of TF binding sites suggests a genome-wide function for Gata2, Tal1 and Gata3 in the selection of the GABAergic fate.

A. Genome-wide enrichment of TF binding sites in the accessible regions in GABAergic and glutamatergic lineages. Heatmap of chromVAR z-scores (avg(TFBS motifs) per TF) for the expressed TFs in rV2 cell groups. All TFs of which a TFBS-motif chromVAR score was among top 10 scores in any cell group are shown.

B. Violin plots of the TF gene expression (average scRNA expression) for *Tal1*, *Gata2*, and
Gata3; and motif accessibility (chromVAR score) of *Tal1*, *Gata2*, and *Gata3* TF-motifs in
the rV2 cell clusters. HOCOMOCO v12 contains three *Tal1* motifs, two *Gata2* motifs and
two *Gata3* motifs. Ordering of cell groups is the same in all violin plots.

Figure 6. Genomic targets of the GABAergic selector TFs.
A. Schematic explaining the strategy of identifying the targets of *Tal1*, *Gata2* and *Gata3*
selector TFs. Within the TAD containing a gene, a feature overlapping a CUT&Tag peak for
the selector TF, and a footprint for the TF at a position with weighted mean conservation
score > 0.5 are found. Genes linked to features fulfilling these conditions are considered
target genes. For linkage, the Spearman correlation-based LinkPeaks score between the
feature targeted by the selector TF and expression of the gene is required to be >2 (positive
effect link) or <-2 (negative effect link) with a p-value <0.01.
B. Number of target genes and the overlap between *Gata2*, *Gata3* and *Tal1* target genes.
C. GSEA of *Tal1* targets. Mouse genes are ranked by the difference in the expression in
GA1-2 vs GL1-2 cell groups (log2 avg FC). Tal1 target genes are indicated with black lines.
In leading edges, the GA1-2-enriched target genes are highlighted in blue and GL1-2-
enriched genes with red. Scatterplots show the expression of the target genes in both edges.
D-F. Characterization of *Tal1* target genes.
D. Count of target features by the positive and negative effect link.
E. Count of target genes by the nearest linked feature, stratified by the nearest feature
distance bins as indicated.
F. The variability of target gene expression in rV2 lineage cell clusters, stratified by the
nearest feature distance bins.
G. Top terms in CellMarker gene set database using the list of *Tal1* target genes with exp>0.5
(log1p) in GA1-2 or GL1-2 cell groups or both.
H. GSEA of Gata2 targets, as in (C).
I-L. Characterization of Gata2 target genes, as in (D-G).
M. GSEA of Gata3 targets, as in (C).
N-Q. Characterization of Gata3 target genes, as in (D-G).
R. Venn diagram of selector TF target genes that show specificity to rV2 glutamatergic
neurons (defined as avg(exp in GA)<0.5 (log1p) AND avg(exp in GL)>0.5 (log1p), and see
Methods). Genes in every overlap category are listed. Vsx2 target genes that show specificity
to rV2 glutamatergic neurons (n=3) are separately listed.
S. Venn diagram of selector TF target genes that show specificity to rV2 GABAergic neurons
(defined as avg(exp in GL)<0.5 (log1p) AND avg(exp in GA)>0.5 (log1p)). Genes in every
overlap category are listed. Vsx2 target genes that show specificity to rV2 GABAergic
neurons (n=2) are separately listed.
T. Proposed gene regulatory network guiding the GABA- vs glutamatergic fate selection in
the rV2. Arrows represent positive regulation and were drawn when the regulator and the
target expression were co-expressed in the same cell group. Blunt arrows represent negative
regulation and were drawn when the regulator and the target expression were found in
different cell groups (GA1-2 vs GL1-2) and not in the same cell group together.
975

Supplementary Figure 1. **E12.5 mouse rhombomere 1 single-cell RNAseq**.

A. UMAP plot of scRNAseq clusters (left), and the expression level of *Nkx6-1, Tal1, Slc17a6* and *Gad1* RNA in the clusters (right).

B. The expression of GABA- and glutamatergic neuron markers in the UMAP of the ventrolateral r1 lineage after lineage subsetting (E12.5 rV2 scRNAseq clusters).

rV2 progenitors (rV2 pro), GABAergic neuron (rV2 GABA) and glutamatergic neuron branches (rV2 Glut) are labelled and indicated with arrows.

Supplementary Figure 2. E12.5 mouse rhombomere 1 single-cell ATACseq.
A. UMAP plot of scATACseq clusters (left), and the inferred expression level of *Nkx6-1, Tal1, Slc17a6* and *Gad1* RNA in the clusters (right).
B. The expression of GABA- and glutamatergic neuron markers in the UMAP of the ventrolateral R1 lineage clusters after lineage subsetting (E12.5 rV2 scATACseq clusters). Arrows indicate the clusters of rV2 lineage progenitors (rV2 PRO), common precursors (rV2 CO), GABAergic neuron (rV2 GABA) and glutamatergic neuron branches (rV2 Glut).

996
997 **Supplementary Figure 3. RNAscope image segmentation and VZ distance calculation**
998 **schematics**
999 A. Typical dissection image with VZ visible on the upper right corner.
1000 B. Same image with cells segmented with StarDist.
1001 C. Schematics of VZ boundary distance -based ordering of the cells.
1002

1003

Supplementary Figure 4. Comparison of the previously characterized enhancers of
***Tal1*, *Gata2*, *Gata3* and *Vsx2* with the scATAC features identified in this study.**
On the genomic loci of the genes, the previously characterised enhancers (details and
references in the Supplementary Table 5) are indicated in colour. The features identified in
this study are shown on the ATAC features (E12.5 R1) track. The gene-linked features are
labelled in blue text.

1010

1011

Supplementary Figure 5. Genomic features and feature accessibility at *Gata2* and *Gata3*
gene loci.

A. Normalized scATACseq signal in +/-50 kb region of Gata2 (A) TSS, in the rV2 lineage
single-cell clusters at top. Below, the Ensembl gene models (Genes), the linkage of features
to *Gata2* or *Gata3* (Links), the scATAC features (Feat.; cCREs are shown in blue), and by-

nucleotide conservation of DNA across vertebrate species (Cons.) is shown. Violin plots show the distribution of *Gata2* (A) RNA expression (log1p) in single cell clusters.

B. Smoothed heatmaps of the accessibility of the *Gata2* cCREs in the rV2 GABAergic and glutamatergic cell lineages (GABA, GLUT). Cluster identities are shown on top of heatmaps. Cells are ordered by the pseudotime value (Pseudotime). cCREs within the TAD bounds are shown on feature heatmaps. RNA expression (log1p) of *Gata2*, *Vsx2*, *Gad1* and *Slc17a6* is shown above the heatmaps.

C. scATACseq signal in the rV2 lineage single-cell clusters and genomic features in +/-50 kb region of *Gata3* TSS, similar to (A). Violin plots show the *Gata3* RNA expression (log1p) in single cells of rV2 cell clusters. *Gata3* gene ATG is located +11 kb from the TSS, the position is indicated in Genes view.

D. Smoothed heatmaps of the accessibility of the *Gata3* cCREs in the rV2 GABAergic and glutamatergic cell lineages (GABA, GLUT). RNA expression (log1p) of *Gata3*, *Tal1*, *Gad1* and *Slc17a6* is shown above the heatmaps.

Supplementary Figure 6. Comparison of the previously characterized enhancers of *Tal1*, *Gata2*, *Gata3* and *Vsx2* with the scATAC features identified in this study.

A. Stacked bar chart showing the count (n) of scATAC features overlapping enhancers and enhancers overlapping scATAC features, separated by the linkage to selector gene.

1037   B. Bar chart showing the percentage of gene-linked features from all features in gene TAD,
1038   the percentage of scATAC features overlapping enhancers and enhancers overlapping all
1039   scATAC features and gene-linked features.
1040

A

1041

Supplementary Figure 7. TF signatures found by scATACseq-footprinting.

A. Heatmap of the TF footprint scores (as avg(TFBS footprint scores) per TF) over the cell groups. Y-axis has been hierarchically clustered with Euclidean distance and complete linkage.

Supplementary Figure 8. Regulatory features of the glutamatergic fate selector *Vsx2* and common regulators of *Vsx2* and *Vsx1*.

A. *Vsx2* gene with its protein-coding regions and associated chromatin features.

B. CUT&Tag. Consensus peaks of CUT&Tag signal with Gata2, Gata3, Tal1, Vsx2, Ebf1, Insm1 and Tead2 antibodies in E12.5 mouse r1.

C-E. TF binding activity in the *Vsx2*-associated features at +20.5 kb, -61 kb and –68.8 kb of Vsx2 TSS according to ATAC-footprinting analysis. In each feature, the strength of footprint (averaged over the cell group shown at the side) at TFBS in the feature is shown as colour

and the expression of the TF gene in dot size. Feature accessibility in cell groups is shown at
the right (Accessibility dotplot). TFBS names are shown at the top and the TF gene names
(mouse genes) are shown under the dotplot.
F. Overlap of the TFs interacting with the cCREs of *Vsx1* and *Vsx2* in the common precursors
of rV2 lineages (CO) and in the rV2 glutamatergic precursors (GL). Common regulators *Vsx1*
and *Vsx2* in CO (n=18) and in GL (n=21) are listed next to venn diagrams. The 18 TFs
regulating both *Vsx1* and *Vsx2* in CO cells are listed in blue text.
§ The probability of finding n overlapping genes considering all mouse genes equally is p<1e-
6.
*,** The collective minimum statistical significance of feature-to-gene links for selector
genes *Vsx1 and Vsx2* cCREs for the given TF is shown as: *p-value<0.05; **p-value<0.01
(with LinkPeaks z-score above 2 or below -2).
1068

1069
Supplementary Figure 9. CUT&Tag of *Tal1* and *Vsx2* -associated chromatin in E12.5
mouse R1, and positive and negative controls.
A. Left panel, Heatmaps and profile plots of the CUT&Tag signal intensity along the mouse
genes in each replicate. Genes are scaled and the areas of 3 kb before transcription start site
(TSS) and 3 kb after transcription end site (TES) are shown. Next to heatmaps, genome views

of *Tal1*, *Lmo1* and *Gata2* gene loci showing the CUT&Tag signal and detected peaks in all

replicates (Re1 - Re4) and consensus peaks (Cons peaks).

B. Genome views of the CUT&Tag signal with the anti-H3K4me3 (positive control) and the

anti-IgG antibody (negative control, Re1-Re4). Consensus peaks are not shown. Only two

consensus peaks were detected between the Re1-Re4 of IgG treated samples.

1080

1081

Supplementary Figure 10. CUT&Tag of Gata2 and Gata3 -associated chromatin in

E12.5 mouse R1.

Heatmaps and profile plots of the CUT&Tag signal intensity along the mouse genes (gene +/-

3kb) are shown for each replicate. Genome views of *Tal1*, *Lmo1* and *Gata2* gene loci show

the CUT&Tag signal, the detected peaks in all replicates (Re1 - Re4) and consensus peaks

(Cons peaks). Correlation heatmaps in the right panel show the CUT&Tag sample correlation

by reads, and by peaks. Venn diagrams show the peak count and peak overlap between the

replicates.

1090

1091

1092

Supplementary Figure 11. Overview of the CUT&Tag for the candidate regulators of

Tal1 genes in E12.5 mouse R1.

CUT&Tag of *Ebf1*, *Insm1* and *Tead2* -associated chromatin in E12.5 mouse R1, and the

signal from the negative control (IgG) and positive control (H3K4me3) samples. Heatmaps

and profile plots show the CUT&Tag signal intensity along the mouse gwenes (gene +/- 3kb)

in each replicate. Genome views show the CUT&Tag signal and the detected peaks in *Tal1*,

Lmo1 and *Gata2* (*Ebf1*, *Insm1* CUT&Tag) or *Vgll3* (*Tead2* CUT&Tag) gene loci from all

replicates (Re1 - Re2) and the defined consensus peaks (Cons peaks).

Correlation heatmaps show the CUT&Tag sample correlation by peaks. Venn diagrams show

the peak count and peak overlap between the replicates.

1103

1104

1105

1106

1107

1108

1109

1110

1111

1112 References

1113 Achim K, Peltopuro P, Lahti L, Li J, Salminen M, Partanen J. 2012. Distinct

1114 developmental origins and regulatory mechanisms for GABAergic neurons

1115 associated with dopaminergic nuclei in the ventral mesodiencephalic region.

1116 *Development* **139**:2360–2370. doi:10.1242/DEV.076380

Achim K, Peltopuro P, Lahti L, Tsai HH, Zachariah A, Åstrand M, Salminen M, Rowitch D, Partanen J. 2013. The role of Tal2 and Tal1 in the differentiation of midbrain GABAergic neuron precursors. *Biol Open* **2**:990–997. doi:10.1242/BIO.20135041

Achim K, Salminen M, Partanen J. 2014. Mechanisms regulating GABAergic neuron development. *Cell Mol Life Sci* **71**:1395–1415. doi:10.1007/s00018-013-1501-3

Bentsen M, Goymann P, Schultheis H, Klee K, Petrova A, Wiegandt R, Fust A, Preussner J, Kuenne C, Braun T, Kim J, Looso M. 2020. ATAC-seq footprinting unravels kinetics of transcription factor binding during zygotic genome activation. *Nat Commun* **11**. doi:10.1038/S41467-020-18035-1

Bockamp EO, McLaughlin F, Göttgens B, Murrell AM, Elefanty AG, Green AR. 1997. Distinct mechanisms direct SCL/tal-1 expression in erythroid cells and CD34 positive primitive myeloid cells. *Journal of Biological Chemistry* **272**:8781–8790. doi:10.1074/jbc.272.13.8781

Boller S, Ramamoorthy S, Akbas D, Nechanitzky R, Burger L, Murr R, Schübeler D, Grosschedl R. 2016. Pioneering Activity of the C-Terminal Domain of EBF1 Shapes the Chromatin Landscape for B Cell Programming. *Immunity* **44**:527–541. doi:10.1016/J.IMMUNI.2016.02.021

Bylund J, Zhang C, Harder DR. 2002. Identification of a novel cytochrome P450, CYP4X1, with unique localization specific to the brain. *Biochem Biophys Res Commun* **296**. doi:10.1016/S0006-291X(02)00918-X

Chen EY, Tan CM, Kou Y, Duan Q, Wang Z, Meirelles G V., Clark NR, Ma'ayan A. 2013. Enrichr: Interactive and collaborative HTML5 gene list enrichment analysis tool. *BMC Bioinformatics* **14**:1–14. doi:10.1186/1471-2105-14-128/FIGURES/3

Ewels PA, Peltzer A, Fillinger S, Patel H, Alneberg J, Wilm A, Garcia MU, Di Tommaso P, Nahnsen S. 2020. The nf-core framework for community-curated bioinformatics pipelines. *Nature Biotechnology 2020 38:3* **38**:276–278. doi:10.1038/s41587-020-0439-x

Faedo A, Laporta A, Segnali A, Galimberti M, Besusso D, Cesana E, Belloli S, Moresco RM, Tropiano M, Fucà E, Wild S, Bosio A, Vercelli AE, Biella G, Cattaneo E. 2017. Differentiation of human telencephalic progenitor cells into MSNs by inducible expression of Gsx2 and Ebf1. *Proc Natl Acad Sci U S A* **114**:E1234–E1242. doi:10.1073/PNAS.1611473114

Francius C, Hidalgo-Figueroa M, Debrulle S, Pelosi B, Rucchin V, Ronellenfitch K, Panayiotou E, Makrides N, Misra K, Harris A, Hassani H, Schakman O, Parras C, Xiang M, Malas S, Chow RL, Clotman F. 2016. Vsx1 transiently defines an early intermediate V2 interneuron precursor compartment in the mouse developing spinal cord. *Front Mol Neurosci* **9**:222932. doi:10.3389/FNMOL.2016.00145/BIBTEX

Garcia-Dominguez M, Poquet C, Garel S, Charnay P. 2003. Ebf gene function is required for coupling neuronal differentiation and cell cycle exit. *Development* **130**:6013–6025. doi:10.1242/DEV.00840

Göttgens B, Broccardo C, Sanchez M-J, Deveaux S, Murphy G, Göthert JR, Kotsopoulou E, Kinston S, Delaney L, Piltz S, Barton LM, Knezevic K, Erber WN, Begley CG, Frampton J, Green AR. 2004. The scl +18/19 stem cell enhancer is not required for hematopoiesis: identification of a 5' bifunctional hematopoietic-endothelial enhancer bound by Fli-1 and Elf-1. *Mol Cell Biol* **24**:1870–1883. doi:10.1128/MCB.24.5.1870-1883.2004

Hu C, Li T, Xu Y, Zhang X, Li F, Bai J, Chen J, Jiang W, Yang K, Ou Q, Li X, Wang P, Zhang Y. 2023. CellMarker 2.0: an updated database of manually curated cell

markers in human/mouse and web tools based on scRNA-seq data. *Nucleic Acids Res* **51**. doi:10.1093/nar/gkac947

Joshi K, Lee S, Lee B, Lee JW, Lee SK. 2009. LMO4 controls the balance between excitatory and inhibitory spinal V2 interneurons. *Neuron* **61**:839–851. doi:10.1016/J.NEURON.2009.02.011

Karunaratne A, Hargrave M, Poh A, Yamada T. 2002a. GATA proteins identify a novel ventral interneuron subclass in the developing chick spinal cord. *Dev Biol* **249**:30–43. doi:10.1006/dbio.2002.0754

Karunaratne A, Hargrave M, Poh A, Yamada T. 2002b. GATA Proteins Identify a Novel Ventral Interneuron Subclass in the Developing Chick Spinal Cord. doi:10.1006/dbio.2002.0754

Kaya-Okur HS, Wu SJ, Codomo CA, Pledger ES, Bryson TD, Henikoff JG, Ahmad K, Henikoff S. 2019. CUT&Tag for efficient epigenomic profiling of small samples and single cells. *Nature Communications 2019 10:1* **10**:1–10. doi:10.1038/s41467-019-09982-5

Kilpinen S, Heliölä H, Achim K. 2023. Range of chromatin accessibility configurations are permissive of GABAergic fate acquisition in developing mouse brain. *BMC Genomics* **24**. doi:10.1186/s12864-023-09836-x

Kobayashi-Osaki M, Ohneda O, Suzuki N, Minegishi N, Yokomizo T, Takahashi S, Lim K-C, Engel JD, Yamamoto M. 2005. GATA motifs regulate early hematopoietic lineage-specific expression of the Gata2 gene. *Mol Cell Biol* **25**:7005–7020. doi:10.1128/MCB.25.16.7005-7020.2005

Lahti L, Haugas M, Tikker L, Airavaara M, Voutilainen MH, Anttila J, Kumar S, Inkinen C, Salminen M, Partanen J. 2016. Differentiation and molecular heterogeneity of inhibitory and excitatory neurons associated with midbrain dopaminergic nuclei. *Development (Cambridge)* **143**:516–529. doi:10.1242/DEV.129957/256855/AM/DIFFERENTIATION-AND-MOLECULAR-HETEROGENEITY-OF

Lakshmanan G, Lieuw KH, Lim K-C, Gu Y, Grosveld F, Engel JD, Karis A. 1999. Localization of Distant Urogenital System-, Central Nervous System-, and Endocardium-Specific Transcriptional Regulatory Elements in the GATA-3 Locus. *Mol Cell Biol* **19**:1558. doi:10.1128/MCB.19.2.1558

Lieuw KH, Li G long, Zhou Y, Grosveld F, Engel JD. 1997. Temporal and spatial control of murine GATA-3 transcription by promoter- proximal regulatory elements. *Dev Biol* **188**:1–16. doi:10.1006/dbio.1997.8575

Martynova E, Bouchard M, Musil LS, Cvekl A. 2018. Identification of Novel Gata3 Distal Enhancers Active in Mouse Embryonic Lens. *Dev Dyn* **247**:1186–1198. doi:10.1002/DVDY.24677

Monaghan CE, Nechiporuk T, Jeng S, McWeeney SK, Wang J, Rosenfeld MG, Mandel G. 2017. REST corepressors RCOR1 and RCOR2 and the repressor INSM1 regulate the proliferation-differentiation balance in the developing brain. *Proc Natl Acad Sci U S A* **114**:E406–E415. doi:10.1073/PNAS.1620230114

Mootha VK, Lindgren CM, Eriksson KF, Subramanian A, Sihag S, Lehar J, Puigserver P, Carlsson E, Ridderstråle M, Laurila E, Houstis N, Daly MJ, Patterson N, Mesirov JP, Golub TR, Tamayo P, Spiegelman B, Lander ES, Hirschhorn JN, Altshuler D, Groop LC. 2003. PGC-1α-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. *Nature Genetics 2003 34:3* **34**:267–273. doi:10.1038/ng1180

Morello F, Borshagovski D, Survila M, Tikker L, Sadik-Ogli S, Kirjavainen A, Estartús N, Knaapi L, Lahti L, Törönen P, Mazutis L, Delogu A, Salminen M, Achim K,

Partanen J. 2020a. Molecular Fingerprint and Developmental Regulation of the
Tegmental GABAergic and Glutamatergic Neurons Derived from the Anterior
Hindbrain. *Cell Rep* **33**:108268. doi:10.1016/J.CELREP.2020.108268
Morello F, Voikar V, Parkkinen P, Panhelainen A, Rosenholm M, Makkonen A,
Rantamäki T, Piepponen P, Aitta-aho T, Partanen J. 2020b. ADHD-like behaviors
caused by inactivation of a transcription factor controlling the balance of inhibitory
and excitatory neuron development in the mouse anterior brainstem. *Transl*
Psychiatry **10**. doi:10.1038/S41398-020-01033-8
Muroyama Y, Fujiwara Y, Orkin SH, Rowitch DH. 2005. Specification of astrocytes by
bHLH protein SCL in a restricted region of the neural tube. *Nature* **438**.
doi:10.1038/nature04139
Nozawa D, Suzuki N, Kobayashi-osaki M, Pan X, Engel JD, Yamamoto M. 2009.
GATA2-dependent and region-specific regulation of Gata2 transcription in the
mouse midbrain. *Genes Cells* **14**:569–582. doi:10.1111/J.1365-2443.2009.01289.X
Ogilvy S, Ferreira R, Piltz SG, Bowen JM, Göttgens B, Green AR. 2007. The SCL +40
enhancer targets the midbrain together with primitive and definitive hematopoiesis
and is regulated by SCL and GATA proteins. *Mol Cell Biol* **27**:7206–7219.
doi:10.1128/MCB.00931-07
Ono Y, Fukuhara N, Yoshie O. 1998. TAL1 and LIM-Only Proteins Synergistically Induce
Retinaldehyde Dehydrogenase 2 Expression in T-Cell Acute Lymphoblastic Leukemia
by Acting as Cofactors for GATA3. *Mol Cell Biol* **18**:6939–6950.
doi:10.1128/MCB.18.12.6939
Radhakrishnan SK, Feliciano CS, Najmabadi F, Haegebarth A, Kandel ES, Tyner AL,
Gartel AL. 2004. Constitutive expression of E2F-1 leads to p21-dependent cell
cycle arrest in S phase of the cell cycle. *Oncogene 2004 23:23* **23**:4173–4176.
doi:10.1038/sj.onc.1207571
Schep AN, Wu B, Buenrostro JD, Greenleaf WJ. 2017. chromVAR: inferring
transcription-factor-associated accessibility from single-cell epigenomic data.
Nature Methods 2017 14:10 **14**:975–978. doi:10.1038/nmeth.4401
Schmidt U, Weigert M, Broaddus C, Myers G. 2018. Cell detection with star-convex
polygonsLecture Notes in Computer Science (Including Subseries Lecture Notes in
Artificial Intelligence and Lecture Notes in Bioinformatics). doi:10.1007/978-3-
030-00934-2_30
Shats I, Deng M, Davidovich A, Zhang C, Kwon JS, Manandhar D, Gordân R, Yao G,
You L. 2017. Expression level is a key determinant of E2F1-mediated cell fate.
Cell Death Differ **24**:626–637. doi:10.1038/CDD.2017.12
Siepel A, Bejerano G, Pedersen JS, Hinrichs AS, Hou M, Rosenbloom K, Clawson H,
Spieth J, Hillier LDW, Richards S, Weinstock GM, Wilson RK, Gibbs RA, Kent
WJ, Miller W, Haussler D. 2005. Evolutionarily conserved elements in vertebrate,
insect, worm, and yeast genomes. *Genome Res* **15**. doi:10.1101/gr.3715005
Sinclair AM, Göttgens B, Barton LM, Stanley ML, Pardanaud L, Klaine M, Gering M,
Bahn S, Sanchez MJ, Bench AJ, Fordham JL, Bockamp EO, Green AR. 1999.
Distinct 5' SCL enhancers direct transcription to developing brain, spinal cord, and
endothelium: Neural expression is mediated by GATA factor binding sites. *Dev*
Biol **209**:128–142. doi:10.1006/dbio.1999.9236
Singh R, Berger B. 2021. Deciphering the species-level structure of topologically
associating domains. *bioRxiv* 2021.10.28.466333. doi:10.1101/2021.10.28.466333
Stassen S V., Yip GGK, Wong KKY, Ho JWK, Tsia KK. 2021. Generalized and
scalable trajectory inference in single-cell omics data with VIA. *Nat Commun* **12**.
doi:10.1038/S41467-021-25773-3

Stuart T. n.d. Sinto. *https://github.com/timoast/sinto*.

Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, Hao Y, Stoeckius M, Smibert P, Satija R. 2019. Comprehensive Integration of Single-Cell Data. *Cell* **177**:1888-1902.e21. doi:10.1016/j.cell.2019.05.031

Stuart T, Srivastava A, Madad S, Lareau CA, Satija R. 2021. Single-cell chromatin state analysis with Signac. *Nat Methods* **18**. doi:10.1038/s41592-021-01282-5

Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP. 2005. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**:15545–15550. doi:10.1073/PNAS.0506580102

Tavano S, Taverna E, Kalebic N, Haffner C, Namba T, Dahl A, Wilsch-Bräuninger M, Paridaen JTML, Huttner WB. 2018. Insm1 Induces Neural Progenitor Delamination in Developing Neocortex via Downregulation of the Adherens Junction Belt-Specific Protein Plekha7. *Neuron* **97**:1299-1314.e8. doi:10.1016/J.NEURON.2018.01.052

Vorontsov IE, Eliseeva IA, Zinkevich A, Nikonov M, Abramov S, Boytsov A, Kamenets V, Kasianova A, Kolmykov S, Yevshin IS, Favorov A, Medvedeva YA, Jolma A, Kolpakov F, Makeev VJ, Kulakovskiy IV. 2023. HOCOMOCO in 2024: a rebuild of the curated collection of binding models for human and mouse transcription factors. *Nucleic Acids Res* gkad1077. doi:10.1093/nar/gkad1077

Waite MR, Skaggs K, Kaviany P, Skidmore JM, Causeret F, Martin JF, Martin DM. 2012. Distinct populations of GABAergic neurons in mouse rhombomere 1 express but do not require the homeodomain transcription factor PITX2. *Molecular and Cellular Neuroscience* **49**:32–43. doi:10.1016/J.MCN.2011.08.011

Wang L, Wang R, Herrup K. 2007. E2F1 works as a cell cycle suppressor in mature neurons. *J Neurosci* **27**:12555–12564. doi:10.1523/JNEUROSCI.3681-07.2007

Weigert M, Schmidt U, Haase R, Sugawara K, Myers G. 2020. Star-convex polyhedra for 3D object detection and segmentation in microscopyProceedings - 2020 IEEE Winter Conference on Applications of Computer Vision, WACV 2020. doi:10.1109/WACV45572.2020.9093435

Xie Z, Bailey A, Kuleshov M V., Clarke DJB, Evangelista JE, Jenkins SL, Lachmann A, Wojciechowicz ML, Kropiwnicki E, Jagodnik KM, Jeon M, Ma'ayan A. 2021. Gene Set Knowledge Discovery with Enrichr. *Curr Protoc* **1**. doi:10.1002/CPZ1.90

Zhang Q, Xu H, Zhao W, Zheng J, Sun L, Luo C. 2020. Zygotic Vsx1 Plays a Key Role in Defining V2a Interneuron Sub-Lineage by Directly Repressing tal1 Transcription in Zebrafish. *Int J Mol Sci* **21**. doi:10.3390/IJMS21103600

A E12.5 rV2

scRNA-seq clusters

B

C RNA expression (scRNA clusters)

Ccnb1 Vsx2 Tal1

Nkx6−1 Slc17a6 Gad1

D scATAC-seq clusters

E

F Inferred RNA expression (scATAC clusters)

Ccnb1 Vsx2 Tal1

Nkx6−1 Slc17a6 Gad1

G Cluster similarity

A Genomic features +/- 50 kb of Tal1 in rV2 cell groups

B Accessibility of Tal1-linked features

C Genomic features +/- 50 kb of Vsx2 in rV2 cell groups

D Accessibility of Vsx2-linked features

A Analysis of regulatory elements of Tal1 gene

B Tal1 +0.7kb** chr4-115056171-115058012 (1841 bp)

C Tal1 +23kb* chr4-115078530-115080241 (1711 bp)

D Tal1 +40 kb** chr4-115095346-115097412 (2066 bp)

E Common regulators of GABAergic fate selectors

Exp (TF) >1.2
Fp in cCRE = 1
TFBS cons > 0.5

§overlap p<1e-06

cCRE linkage
**p<0.01
*p<0.05

A Genome-wide accessibility of TFs (with motifs averaged) per cell group

B
RNA expression level

Tal1 *Gata2* *Gata3*

C Motif accessibility (chromvar)

TAL1.H12CORE.0.P.B GATA2.H12CORE.0.PSM.A GATA3.H12CORE.0.PS.A

TAL1.H12CORE.1.P.B GATA2.H12CORE.1.P.B GATA3.H12CORE.1.SM.B

TAL1.H12CORE.2.P.B

A Transcription factor target genes

Within TAD

Links score > 2 or < -2,
p-value <0.01
(Acc. to Exp.)

Footprint in scATAC

ATAC feature — Gene

N target genes

Cut&Tag peak

Mean Cons. > 0.5

B All targets overlaps

Gata2 n=780   Gata3 n=721

70   50   218

343   317   136

3152

Tal1 n=3948

+/-

RE T
Target-linked features by effect

dist

RE T
Target genes by nearest feature

C Tal1 target enrichments in GA/GL DEG

enrichment score

rank

GA1_2 Expression log2(avg+1)
GL1_2 Expression log2(avg+1)

Rps8, Rps24, Rps15a, Rps2, Rpl8, Rpl27a, Srsf3, Rpl36, Rpl38, Rpl7a, Lmo1, Elavl2, Pbx1, Snrpe, Rps19, Rpl12, Sox2, Ube2s, Id4, Nme1, Rps15, Rnd2, Csrp2, Stpg, Gem1, Atp5a1, Vdp, Lhx5, Tal1, Irx1, Cfl1, Homc, Zfand5, Pcbp4, Cbx3, Hes5, St18, Luc7l3, Sumo1, Atp5c1, Gata3, Qk, Vim, Cmpk1, Ppdpf, Optc

Malat1
H3f3b, Basp1, Tmsb10, Dynlt1, Pantr1, Tubb3, Myl6, Gm1673, Jund, Stmn2, Mif2, Atp6v0b, Cirbp, Uqcr10, Scand1, Dnajb6, Ddah2, Cdc42, Npdc1, Tecr, Podxl2, Bcl11b, Ebf2, Igfbpl1, Nkx6-1, Lhx4, Nhlh2

D Target-linked features by effect
Count
Pos. link   Neg. link

E Count
0-5 kbp   5-50 kbp   > 50 kbp

F Expression Variability (Standard Deviation)
0-5 kbp   5-50 kbp   > 50 kbp

G CellMarker (2024)   combined score

Tal1 targets (exp>0.5)

Cell type	combined score
Progenitor Cell Kidney Human	166.38
Interneuron Brain Human	166.38
Transit Amplifying (TA) Cell Intestinal Crypt Mouse	147.61
Alpha Cell Pancreas Mouse	115.05
Goblet Cell Intestinal Crypt Mouse	101.81
Mid Radical Glial Cell Brain Human	80.5
Enteroendocrine Precursor Cell Intestinal Crypt Mouse	73.67
Neural Stem Cell Brain Mouse	72.46
Müller Glial Cell Retina Mouse	69.17
Neural Stem Cell Hypothalamus Mouse	69.17

H Gata2 target enrichments in GA/GL DEG

enrichment score

rank

GA1_2 Expression log2(avg+1)
GL1_2 Expression log2(avg+1)

Rps8, Rpl27a, Lmo1, Snrpe, Cbx3, Crabp2, Tal1, Gad2, St18, Runx1t1, Gata2, Nrxn3, Nr2f2, Zfbm1, Elsrz, Cmpk1, Lhx5, Sox13, Crt2, Klhl7, Shd, Pabpn1, Pdzk1ip1, Gm12324, Il17, Eif2s1, Nfit, Rad23a, Med8, Coq5, Ccdc15, Lyt1, Elmo2, Rps34, Rbm15, Nrtn, Lsamp, Slc25a39

Tubb3
Dynll1
Ralgds, Cxxc5, Tcf25, Trappc2l, Arl16, Coro1c, Zcchc11, Enl3, Tla3, Appb1, Rnf206, Sharpin, Nfu1, Shf, Mast1, Sh2d3c, Sorcs1, Speg, Gmppa, Kit26a, Rem2, Hdac5

I Target-linked features by effect
Count
Pos. link   Neg. link

J Count
0-5 kbp   5-50 kbp   > 50 kbp

K Expression Variability (Standard Deviation)
0-5 kbp   5-50 kbp   > 50 kbp

L

Gata2 targets (exp>0.5)

Cell type	score
Inhibitory Neuron Brain Human	612.44
Interneuron Brain Human	434.35
Endocardial Cell Embryo Mouse	434.35
Inhibitory Neuron Hippocampus Human	264.02
Red Blood Cell (Erythrocyte) Bone Marrow Mouse	217.53
GABAergic Neuron Brain Mouse	125.37
Endothelial Cell Muscle Mouse	108.14
Hematopoietic Stem Cell Spleen Human	100.06
Inhibitory Neuron Brain Mouse	100.06
Interneuron Cortex Human	100.06

M Gata3 target enrichments in GA/GL DEG

enrichment score

rank

GA1_2 Expression log2(avg+1)
GL1_2 Expression log2(avg+1)

Rps8, Sox11, Lmo1, Gata3, Id4, Elavl2, Snrpe, Slc2a1, Gad2, Zfpm1, Nrxn3, Sox2, En1, Runx1t1, Cbx3, Pdzrn4, Sox9, Tie4, Ddx6, Kin, Zfpm2, Lrx1, Lhx5, Ppdpf, Maby, Hrh1, Atp1b1, Pabpn1, Otx1, Plekha1, Scrt2, Khlf7, Kit2a, Sap30l, Mtap3, Mpdr8, Lyt1, Btbd3, Sox13, Gep78, Gpn1, Nrtn, Fam210b, Chst1, Cdh4, Rbms1

Tubb3
Rnd3
Tecr, Smc3, Cxxc5, Tcf25, Trappc2l, Orc6, Tla3, C13007C03Rik, Rem2, Tle3, Neurg1, Kif26a, Flrt3, Slc2a1, Pou3f1, Tle1, Vsx2

N Target-linked features by effect
Count
Pos. link   Neg. link

O Target genes by nearest feature
Count
0-5 kbp   5-50 kbp   > 50 kbp

P Expression Variability (Standard Deviation)
0-5 kbp   5-50 kbp   > 50 kbp

Q

Gata3 targets (exp>0.5)

Cell type	score
Inhibitory Neuron Brain Human	1963.74
GABAergic Neuron Brain Mouse	1386.52
Interneuron Brain Human	1233.11
Inhibitory Neuron Hippocampus Human	672.3
Inhibitory Neuron Brain Mouse	522.52
Interneuron Cortex Human	522.52
Pan-trophectoderm Cell Embryo Human	522.52
Ectoderm Cell Embryo Mouse	522.52
Somatic Cell Testis Human	522.52
Stem Cell Bladder Human	522.52

p-val<0.0001   p-val<0.05

R Target genes in glutamatergic neuron-specific genes

Gata2   Gata3
0   0   2
0   0   2
12

Tal1

Gata3 (2)
Vsx2
Tle1

Gata3, Tal1 (2)
Gm17750
Kif26a

Vsx2 targets (3)
Vsx2
Dlk1
Gm17750

Tal1 (12)
Nkx6-1
Skor1
Igfbpl1
Nhlh1
Tmem163
Zfhx4
Tppp3
Ebf2
Dlk1
2300009A05Rik
Lhx4
Nhlh2

S Target genes in GABAergic neuron-specific genes

Gata2   Gata3
0   3   9
0   7   2
3

Tal1

Gata2, Gata3 (3)
Gad2
Nrxn3
Tal2

Common targets (7)
Btbd17
Zfpm1
Pdzk1ip1
Tal1
Lmo1
Cbfa2t3
Gata2

Vsx2 targets (2)
Btbd17
Lmo1

Gata3, Tal1 (2)
9230102O04Rik
Sox9

Tal1 (3)
Hes5
Qk
Fam46a

Gata3 (9)
Kirrel3
Slc32a1
Otx1
Pdzrn4
Zfpm2
Plekha1
Gad1
Gata3
Nrip1

T Gene regulatory interactions in the rV2 GABAergic neuron fate determination

Nhlh1
Nhlh2
Nkx6-1
Lhx4

Hes5
Dlk1
St18
Lmo1

Tal1   Vsx2

Gata2
Gata3

Gad1
Gad2
Slc32a1