**Non-Visual Light Sensing Enhances Behavioral Memory and Drives Gene Expression in *C. elegans***

Zhijian Ji[1, §], Bingying Wang[1, §], Rashmi Chandra[1, §], Junqiang Liu[1, §], Supeng Yang[1, 2], Yong Long[3], Michael Egan[1, 2], Noelle L'Etoile[4], Dengke K. Ma[1, 5, 6, *]

[1]Cardiovascular Research Institute, University of California San Francisco, San Francisco, CA 94158, USA.

[2]Department of Molecular Cell Biology, University of California, Berkeley, Berkeley, CA 94720, USA.

[3]Key Laboratory of Breeding Biotechnology and Sustainable Aquaculture, Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan 430072, China.

[4]Department of Cell and Tissue Biology, University of California San Francisco, San Francisco, CA 94158, USA.

[5]Department of Physiology, University of California San Francisco, San Francisco, CA 94158, USA.

[6]Innovative Genomics Institute, University of California, Berkeley, Berkeley, CA 94720, USA.

[§]Equal contribution

*Correspondence: Dengke.Ma@ucsf.edu (D.K.M.)

**Abstract**

Visible light influences a range of physiological processes, yet how animals respond to it

independently of the visual system remains largely unknown. Here, we uncover a

previously undescribed light-induced transcriptional pathway that modulates behavioral

plasticity in *C. elegans*, a roundworm without eyes. We demonstrate that ambient visible

light or controlled-intensity visible-spectrum LED activates an effector gene *cyp-14A5* in

non-neuronal tissues through the bZIP transcription factors ZIP-2 and CEBP-2. Light

induction of *cyp-14A5* is more prominent at shorter wavelengths but is independent of

the known blue light receptors LITE-1 and GUR-3 in *C. elegans*. This bZIP-dependent

genetic pathway in non-neuronal tissues enhances behavioral adaptability and olfactory

memory, suggesting a body-brain communication axis. Furthermore, we use the light-

responsive *cyp-14A5* promoter to drive ectopic gene expression, causing synthetic light-

induced sleep and rapid aging phenotypes in *C. elegans*. These findings advance our

understanding of light-responsive mechanisms outside the visual system and offer a

new genetic tool for visible light-inducible gene expression in non-neuronal tissues.

**Introduction**

Visible light is crucial for image formation and regulating various physiological

processes through the visual system, yet how animals respond to ambient light

independently of sight remains poorly understood. Recent studies have uncovered

diverse non-visual photoreception mechanisms that modulate a range of biological

processes, from circadian rhythms to stress responses and metabolic homeostasis[1–4].

These mechanisms often involve specialized light-sensitive proteins, such as opsins

and cryptochromes, widely expressed in body locations, including the skin, brain, and

peripheral organs. For example, mammalian melanopsin-expressing retinal ganglion

cells play critical roles in systemic light responses largely independent of image

formation[5–8]. In the nematode roundworm *C. elegans*, blue light photoreception requires

the light-activated ion channels LITE-1 and GUR-3 in specific neurons, influencing

aversive behaviors and cellular physiology[9–13]. Visible light irradiation can also cause

photo-oxidative reactive oxygen species in animals[14–16]. Despite these advances, the

molecular pathways and physiological outcomes of non-visual light sensing and

responses remain largely unexplored, raising intriguing questions about the mechanistic

basis and functional implications of light as an environmental cue beyond vision.

We previously studied how genes encoding cytochrome P450 (CYP) proteins respond

to and mediate effects of exposure to environmental stresses in *C. elegans*[17,18]. Among

various transcriptional reporters we generated for CYP-encoding genes to monitor

environmental regulation, we serendipitously discovered that the *cyp-14A5* promoter-

driven GFP expression is particularly sensitive to ambient visible light exposure.

Building upon this initial finding, we conducted transcriptome profiling studies to identify

light-inducible genes in addition to *cyp-14A5*, determine key transcriptional regulators of

*cyp-14A5*, and show that the light-inducible CYP-14A5 promotes behavioral plasticity

and olfactory memory in *C. elegans*. The findings also provide a genetic tool to use

light-inducible *cyp-14A5* promoter to flexibly and ectopically drive gene expression.

**Results**

**Light activates expression of *cyp-14A5* and other genes in *C. elegans***

Cytochrome P450 proteins comprise a highly conserved superfamily of heme-containing

monooxygenases critical for metabolizing endogenous and xenobiotic compounds[19].

We constructed transcriptional reporters for genes encoding CYP in *C. elegans* and

found that *cyp-14A5p::GFP* was drastically up-regulated by bright-field transmission

light from a microscope inadvertently left on overnight. Using controlled light versus dark

conditions, we confirmed the finding from an integrated *cyp-14A5p::GFP* reporter and

observed its robust widespread GFP expression in many tissues induced by moderate-

intensity (500-3000 Lux, 16-48 hr duration) LED light exposure (**Fig. 1A**). The level of

GFP expression increased proportionally with both light intensity and duration, the

condition of which does not impact ambient temperature (**Fig. S1**), indicating that *cyp-*

*14A5p::GFP* expression is finely tuned to ambient light conditions (**Fig. 1B**).

To determine CYP-14A5 protein expression pattern, we constructed a translational GFP

(*cyp-14A5p::cyp-14A5::GFP*) reporter and observed robust light-induced expression of

CYP-14A5::GFP in many of the non-neuronal tissues, including the pharynx, hypoderm

and intestine (**Fig. 1C**). The transcriptional reporter exhibited similar patterns of non-

neuronal GFP induction by light (**Fig. 1A**). The translational reporter reveals CYP-

14A5::GFP patterns indicative of the endoplasmic reticulum structure (**Fig. 1C**),

consistent with the known subcellular localization of most CYPs to ER membranes[20].

To explore how eyeless *C. elegans* responds to ambient visible light independently of a

visual system, we conducted transcriptomic profiling by RNA-seq in *C. elegans* exposed

to controlled light or dark conditions. We found that defined light exposure (1500 Lux, 24

hr duration) to a synchronized population of young adults (24 hrs post L4) triggered a

robust genome-wide transcriptional response, including 7902 genes differentially

regulated (adjusted *P* value < 0.05, **Fig. 1D** and **Table S1**). Among these, *cyp-14A5*

was one of the most strongly upregulated genes (**Fig. 1E**). Gene ontology (GO) analysis

of the light-induced transcriptome reveals their significant enrichment in several

pathways, including transmembrane signaling, pathogen and stress responses, protein

phosphorylation, cellular homeostasis and metabolisms (**Fig. S2**).

**ZIP-2 and CEBP-2 mediate light-induced transcriptional responses**

We next investigated the molecular regulators driving *cyp-14A5* activation in response

to light. To test if it requires previously identified blue-light receptors LITE-1 or GUR-3,

we crossed the *cyp-14A5p::GFP* reporter with *lite-1* and *gur-3* double loss-of-function

mutants. Interestingly, light-induced GFP expression was preserved in *lite-1gur-3*

double mutants, indicating that *cyp-14A5* activation operates through an alternative,

non-visual light-sensing mechanism (**Fig. 2A**). Prolonged photon exposure may also

cause photo-oxidation of DNA and genotoxicity, leading to DNA damage and ATM

protein-dependent check points and transcriptional responses[15,21,22]. However, loss of

the DNA damage sensor ATM-1 did not apparently affect light-induced GFP expression

(**Fig. 2A**). These findings underscore the existence of a novel light-responsive pathway

in *C. elegans*, distinct from previously characterized photoreceptive systems.

To identify transcriptional regulators driving *cyp-14A5* activation in response to light, we

adopted an RNAi-based screening strategy, focusing on approximately 400 genes

encoding transcription factors, including those responding to various types of stresses.

Knockdown of the expression of genes encoding TF from a previously assembled RNAi

library or selected for mediating various common stress responses (hypoxia, oxidative

stress, heat shock etc.) did not appear to affect *cyp-14A5* activation in response to light

(**Fig. 2B**). Surprisingly, we identified from such screen two pathogen-responding bZIP

transcription factors, ZIP-2 and CEBP-2, as critical mediators of light-induced *cyp-14A5*

transcription (**Fig. 2B**). Knockdown or genetic ablation of either *zip-2* or *cebp-2*

abolished light-induced *cyp-14A5p::GFP* expression (**Fig. 2B, 2C**). ZIP-2 and CEBP-2

have been previously identified[23–25] to cooperate in a regulatory complex and mediate

transcriptional responses to the bacterial pathogen *Pseudomonas aeruginosa* PA14. In

these studies, the *irg-1*p::GFP transcriptional reporter has been shown to be robustly

activated by PA14 as a well-established target for ZIP-2 and CEBP-2. Interestingly, we

found *irg-1*p::GFP was not activated by the same light condition (1500 Lux, 24 hrs) that

reliably induced *cyp-14A5*p::GFP (**Fig. 2D**). Although ZIP-2 can be activated by

pathogen stresses through ribosomal inhibition[23–25], our results highlight the specific

roles of ZIP-2 in mediating light-induced *cyp-14A5* but not *irg-1* reporter expression,

suggesting the involvement of additional stress-specific factors in these processes.

We further explored conditions and mechanisms leading to light-induced *cyp-*

*14A5p::GFP*. To test potential effects of ultraviolet (UV) irradiation from our visible light

LED, we used a UV-masking shield to block UV irradiation. However, this did not affect

visible LED light-induced *cyp-14A5p::GFP* expression (**Fig. 2E**). In addition, we found

that the LED light exposure of equal intensities (1500 Lux, 24 hrs) but at different

wavelengths (red, green, blue) led to differential *cyp-14A5p::GFP* expression (**Fig. 2F,**

**2G**), indicating stronger effects of shorter wavelengths in the visible light spectrum. A

pseudo-open reading frame (uORF) in the 5' untranslated region (UTR) of *zip-2* mRNA

inhibits the ribosomal translation of the ZIP-2 main open reading frame (mORF) in the

context of PA14 pathogen exposure[25]. However, constitutive expression of *zip-2* uORF

by the *rpl-28* promoter did not affect light-induced *cyp-14A5p::GFP* (**Fig. 2H**).

Furthermore, a CRISPR phospho-site knock-in mutation of *eif-2alpha(S49A)* did not

affect global translation[26], yet abolished light-induced *cyp-14A5p::GFP* (**Fig. 2I**). As the

eukaryotic eIF2alpha complex facilitates translational switch from uORF to mORF upon

stress-induced ribosomal stall at uORF[27–29], these results suggest that it is the *zip-2*

uORF translational inhibition, not the uORF protein product function, that mediates

visible light-induced ZIP-2 activation and *cyp-14A5p::GFP* reporter expression (**Fig. 2J**).

**Light-induced CYP-14A5 enhances behavioral memory**

Although EIF-2alpha and ZIP-2/CEBP-2 functions appear essential for light-induced up-

regulation of *cyp-14A5*, the *zip-2, cebp-2* or *cyp-14A5* loss-of-function null mutants

show no apparent body-size, morphological, feeding, defecation or developmental

defects under dark or LED light treatment (1500 Lux for 16 or 24 hrs) conditions (**Fig.**

**S3**). These data suggest that transient visible light exposure or light-induced *cyp-14A5*

activation by ZIP-2 does not broadly affect development or physiology, unlike long-term

visible light exposure, which has been shown to shorten lifespan in *C. elegans*[15].

The lack of obvious morphological and basal behavioral defects led us to explore

whether light exposure influences other aspects of *C. elegans* biology, particularly

behavioral plasticity and associative memory formation that might require integration of

body physiology. Specifically, we chose a learning paradigm in which animals learn to

avoid an innately attractive odor after it is paired with starvation[30,31]. They can

consolidate this learning into a long-lasting memory if the training is followed by sleep

within two hours[30]. Using this conditioning protocol (**Fig. 3A**), in which *C. elegans* learns

to avoid butanone (an innately attractive odor associated with nutritious bacteria) and

maintain the memory for up to 16 hours[30,31], we observed that animals exposed to

ambient light (approximately 500–1000 Lux) during olfactory associative learning and

recovery exhibited significantly enhanced memory retention compared to those

maintained in darkness (**Fig. 3B**). To pinpoint the critical period for light exposure, we

deprived animals of light in two-hour intervals immediately post-learning. Remarkably,

light deprivation during the first 2–4 hours post-learning resulted in markedly impaired

memory retention (**Fig. 3C**). These results suggest that environmental light exposure

enhances aversive cue association post learning and is not required for learning itself

but is required immediately post learning for consolidation of memory.

Does the light-modulated behavioral memory consolidation require light activation of the

ZIP-2 pathway? To address this question, we examined the behavioral memory of two

independent *zip-2* deletion mutants as compared to wild type (*cebp-2* mutants are

pleiotropically sick and thus not included). We found that both the *ok3730* and *tm4246*

deletion mutations of *zip-2* caused a significantly impaired memory consolidation but not

learning (**Fig. 3D**). Strikingly, the loss-of-function mutation of *cyp-14A5*, but not *F43C1.7*

(another ZIP-2 target gene induced by visible light), also impaired memory as *zip-2* (**Fig.**

**3E**), indicating a crucial role of the ZIP-2/CYP-14A5 regulatory axis in mediating light-

modulated memory consolidation. To further delineate the role of CYP-14A5, we

performed tissue-specific rescue experiments in the behavioral memory assay.

Hypoderm-specific expression of *cyp-14A5* restored the behavioral memory in the *cyp-*

*14A5* mutant (**Fig. 3F**). We observed similar degrees of rescue by two independently

derived lines expressing hypoderm-specific *dpy-7p::cyp-14A5* transgenes. These

findings strongly suggest that hypodermal induction of CYP-14A5 by ZIP-2 plays a

central role in mediating light-modulated behavioral memory.

**The *cyp-14A5* promoter as a versatile tool for light-inducible gene expression**

The light-responsive nature of the *cyp-14A5* promoter prompted us to explore its

potential as a tool for controlling gene expression. We generated synthetic constructs

driving the expression of diverse effectors under the *cyp-14A5* promoter to confer

striking organismal phenotypes. We previously found that *zip-10* expression promotes

organismal phenoptosis[32,33]. Driven by the *cyp-14A5* promoter, light-induced *zip-10*

expression indeed caused a robust light-dependent rapid aging or phenoptosis-like

phenotype with markedly shortened median and maximal lifespans (**Fig. 4A-4D**). We

confirmed that LED light exposure successfully induced *zip-10* expression, as

evidenced by robust ZIP-10-tagging mCherry fluorescence in major non-neuronal

tissues of transgenic animals, only after light (1500 Lux, 24 hrs) exposure (**Fig. 4B**).

To test organismal behavioral outcomes, we expressed *nlp-22* under the *cyp-14A5*

promoter. *nlp-22* was previously identified as a sleep-promoting neuropeptide[34–36],

overexpression of which can cause drastic reduction of pumping and locomotion speed,

characteristic of sleep behaviors in *C. elegans*. We found that *cyp-14A5p::nlp-22* can

indeed trigger striking behavioral quiescence upon light exposure (1500 Lux, 24 hrs), as

quantified by pumping rates, bending frequencies and locomotion speed (**Fig. 4E-4G,**

**S4**). These proof-of-concept studies demonstrate that the *cyp-14A5* promoter enables

light-dependent ectopic induction of gene expression, offering a flexible tool for probing

gene function, studies of organismal biology and synthetic physiology applications.

## Discussion

Our findings uncover a previously unknown light-induced transcriptional pathway in *C.*

*elegans* that operates independently of known visual light receptors. Our study also

establishes a functional link between ambient light and behavioral plasticity through a

ZIP-2/CYP regulatory axis. The discovery of this pathway and its organismal functions

opens exciting avenues for understanding body-brain communication and how

environmental cues such as light can shape physiological and behavioral processes.

The specificity of this pathway is particularly intriguing, as *cyp-14A5* is robustly induced

by light at wavelength and intensity that do not apparently alter ambient temperature

(**Fig. S1**). Previous studies identified *cyp-14A5* as one of many genes moderately

regulated by bacterial pathogens[37–39], yet the classic pathogen-inducible gene reporter

*irg-1p::GFP* was not activated by light. This specificity raises important questions about

how ZIP-2 and *cyp-14A5* are selectively activated by light. Given the crucial roles of

ZIP-2 and eIF2alpha we find for light-induced expression of *cyp-14A5* and the

established role of the uORF at the 5' untranslated region of *zip-2* RNA in ZIP-2

regulation[25], it is plausible that light may regulate ZIP-2 translation by *zip-2* RNA photo-

oxidation at specific sites, eIF2alpha phosphorylation and specialized ribosomal

signaling[40–42]. Further investigation into the upstream signaling events and the

molecular sensors linking light exposure to ZIP-2/CEBP-2 activation is warranted.

Our behavioral assays demonstrate that light exposure enhances olfactory associative

memory, providing direct evidence for the functional relevance of light-induced *cyp-*

*14A5* expression. Interestingly, light exposure appears to exert its effects during a

specific temporal window immediately following learning, suggesting that light-induced

transcriptional changes post learning play a key role in memory consolidation. The

discovery of *cyp-14A5* as a key effector in this pathway also provides new insights into

how non-neuronal tissues contribute to behavioral plasticity. Our findings suggest that

light-induced CYP-14A5 and CYP-dependent metabolic or signaling changes in the

hypoderm may communicate with the nervous system to influence behavioral memory.

This body-brain communication axis highlights the importance of systemic integration in

mediating complex physiological and behavioral responses to environmental cues[43–46].

Beyond its biological significance, the light-inducible *cyp-14A5* promoter offers a useful

new tool for gene expression studies in *C. elegans*. The ability to drive ectopic gene

expression in response to light provides a versatile system for temporally controlled

genetic manipulations. Our demonstration of light-induced sleep and mortality

phenotypes through ectopic gene expression illustrates the potential applications of this

tool in studying diverse biological processes in synthetic biology and physiology.

Previous studies have used heat shock or drug-inducible promoters for temporally

controlled gene expression in *C. elegans*[47–49]. The light-inducible *cyp-14A5* promoter

provides an alternative, simple-to-implement approach and might be particularly useful

when the drug-inducible system is cumbersome, or heat shock effects are undesirable.

While our study uncovers a novel light-responding mechanism with functional

consequences in *C. elegans*, several limitations exist. First, the precise molecular

mechanism by which visible light activates ZIP-2 and/or CEBP-2 remains unclear, as

does the upstream signaling cascade linking light exposure to transcriptional activation.

Second, although we demonstrate a functional connection between light-induced *cyp-*

*14A5* expression and behavioral outcomes, the exact molecular interplay between

peripheral transcriptional changes and neural plasticity requires further exploration.

Finally, while the *cyp-14A5* promoter serves as a useful genetic tool, it does not confer

tissue specificity, and its ectopic effector expression requires control for light effects.

These limitations provide fertile ground for future research to build upon our findings.

**Materials & Methods**

*C. elegans* **strains**

*C. elegans* strains were grown on nematode growth media (NGM) plates seeded with

*Escherichia coli* OP50 at 20 °C with laboratory standard procedures unless otherwise

specified. The N2 Bristol strain was used as the reference wild type[50]. Mutants and

integrated transgenes were back-crossed at least 5 times. Genotypes of strains used are

as follows: *dmaIs156 IV [cyp-14A5p:: cyp-14A5::GFP; unc-54p::mCherry], agIs17 IV [irg-*

*1p::gfp], dmaEx [dpy-7p::cyp-14A5; myo-2p::mCherry], dmaEx [cyp-14A5p::zip-*

*10::mCherry; myo-2p::mCherry], dmaEx [cyp-14A5p::nlp-22; myo-2p::mCherry], cebp-2*

*(tm5421) I, eif-2alpha(rog3) I, zip-2(tm4248) III, zip-2(ok3730) III, cyp-14A5(gk152) V.*

PCR fusion constructs were used to generate transgenes[51], using primer sequences:

DM1310_*cyp-14A5*Pro c5p TCAACCACATCTTCCGATCA;

DM1311_*cyp-14A5*Pro to GFP c3p

CGACCTGCAGGCATGCAAGCTgatctttgttggacagaatagtttt;

DM2857_*dpy-7*p to *cyp-14A5* codutr Forward TGTCTCTGACGCCTGTGAGT;

DM2858_*dpy-7*p to *cyp-14A5* codutr Reverse

GATAAAGCAACGATGAAAACGCTCATTTTGTTTTCACAGAGCGGTAGA;

DM2944_*zip-2*uORF to *rpl-28*p-GOI-mCherry-5utr fusion F

CATCATAAAATAATTTATTTCCAGGTAAAATGTATCACGCAAAGACAACCACCG;

DM2945_*zip-2*uORF to *rpl-28*p-GOI-mCherry-5utr fusion

catgttatcttcttcaccctttgaggagccAAGCTCCCGTGGGAAGCTTGTG;

DM2948_*zip-10* to *cyp-14A5*p-GOI-mCherry-5utr fusion F

aaaactattctgtccaacaaagatcaaaATGACAACAATGACTAATTCTCTTATTTC;

DM2949_*zip-10* to *cyp-14A5*p-GOI-mCherry-5utr fusion R

catgttatcttcttcaccctttgaggagccGGAATGGTTGATTTGATTATTGAGTTG

DM2952_*nlp-22*cod::3utr to *cyp-14A5*p-GOI fusion

aaaactattctgtccaacaaagatcaaaATGCGTTCCATAATCGTCTTCATCG;

DM2953_*nlp-22*cod::3utr to *cyp-14A5*p-GOI fusion R   cggttccactttctcatgagt

**Fluorescence microscopy and imaging**

SPE confocal (Leica) and epifluorescence microscopes were used to capture

fluorescence images. Animals were randomly picked at the same stage and treated with

1 mM levamisole in M9 solution (31742-250MG, Sigma-Aldrich), aligned on a 2% agar

pad on a slide for imaging. Identical setting and conditions were used to compare

experimental groups with control. For quantification of GFP fluorescence, animals were

outlined and quantified by measuring gray values using the ImageJ software. The data

were plotted and analyzed by using GraphPad Prism10.

For light-induced reporter imaging, reporter animals (synchronized young adults, 24 hrs

post L4) were exposed to white light (1500 Lux for 16 or 24 hrs, by Viribright 12-Watt, 800

Lumen, LED Desk Lamp Dimmable Office Lamp). For blue, green (SPE confocal, Leica),

and red light (HQRP 660 nm 14w LED pure red) conditions, animals of the same stage

were exposed to the same intensities (1500 Lux, 16 or 24 hours). Control groups from the

same batch of animals were maintained in darkness by opaque shields. Light intensities

and temperature were quantitatively measured by digital light meters and thermometers.

**RNA sequencing**

A synchronized population of wild-type young adult (24 hrs post L4) animals were

exposed to LED light (1500 lux, 24 hr duration, Viribright 12-Watt, 800 Lumen, LED Desk

Lamp Dimmable Office Lamp). Control groups from the same batch of animals were

maintained in darkness by opaque light shields. Four independent biological replicates

were used for both light-treated and control groups. For sample collection, the animals

were washed down from NGM plates using M9 solution and bacteria-cleaned with M9

washing in centrifuge tubes, homogenized by tissue disruptors and subjected to RNA

extraction using the RNeasy Mini Kit from Qiagen. 1 mg total RNA from each sample was

used for sequencing library construction. The libraries were constructed and sequenced

for paired end 150 bp by DNBseq (Innomics). The cleaned RNAseq reads were mapped

to the genome sequence of *C. elegans* using hisat2 and the mapped reads were assigned

to the genes using featureCounts[52,53]. The abundance of genes was expressed as RPKM

(Reads per kilobase per million mapped reads) and identification of differentially

expressed genes was performed using the DESeq2 package[54].

***C. elegans* behavioral assays**

The olfactory behavioral memory assay was as described previously with modification[30,31].

Briefly, one day old adult worms (24 hrs post L4) were washed with S basal buffer (0.1M

NaCl, 0.05M $K_3PO_4$, pH 6.0) off 10 cm NGM plates and into microfuge tubes, where they

were washed three times with S basal buffer. The animals were split in two groups; one

group was added to a microfuge tube of S basal and the other group was added to a

microfuge tube of 1:10,000 dilution of butanone in S basal. The microfuge tubes were

then rotated for 80 minutes. The odor training includes three 80-minute cycles of training

with odor, or a control buffer interspersed with two 30-minute periods of feeding with OP50

*E. coli* bacteria. For chemotaxis assay, 1 μL of (1 M) $NaN_3$ was pipetted onto the odor

and diluent spots in 10 cm plastic petri dishes. 1 μL of 200 proof ethanol was added to

the diluent spot and 1 μL of 1:1000 butanone was added to the odor spot, while S basal

or butanone-trained worms were dropped onto the middle of 10 cm plastic petri dishes.

The recovery period was either under darkness or ambient light (600 Lux) for 16 hrs or

were kept under darkness for 2 or 4 hrs periods followed by light exposure and

chemotaxis to assay behavioral memory.

For sleep analysis induced by *cyp-14A5p::nlp-22*, the bending angles, moving average

speed, and track length of *C. elegans* after 48 hours of exposure to either light or dark

conditions were analyzed using WormLab. In such experiment, a synchronized population

of young adult (24 hrs post L4) animals of indicated genotype or transgene expression

were used. After light exposure or control dark treatment, they were transferred to a fresh

NGM plate seeded with a small OP50 bacterial lawn and allowed to settle for at least ten

minutes to recover at room temperature. After the recovery period, a one-hour recording

session was conducted using WormLab. Bending angles were calculated as described in

the referenced method as a metric for sleep behaviors[30]. Moving average speed was

determined by tracking the displacement of the worms over time.

**Statistics**

Numerical data were analyzed using GraphPad Prism 10 Software (Graphpad, San

Diego, CA) and presented as means ± S.D. unless otherwise specified, with *P* values

calculated by unpaired two-tailed t-tests (comparisons between two groups), one-way

ANOVA (comparisons across more than two groups) and two-way ANOVA (interaction

between genotype and treatment), with post-hoc Tukey and Bonferroni's corrections.

The lifespan assay was plotted and quantified using Kaplan–Meier lifespan analysis,

and *P* values were calculated using the log-rank test.

**Acknowledgment**

Some strains were provided by the *Caenorhabditis* Genetics Center (CGC), which is

funded by the NIH Office of Research Infrastructure Programs (P40 OD010440), and by

Dr. E. Troemel. We also thank the *C. elegans* Reverse Genetics Core Facility

(University of British Columbia), National Bioresource Project (S. Mitani, Tokyo

Women's Medical University, Tokyo, Japan), Wormbase.org (NIH grant #U24

HG002223 to P. Sternberg), Wormatlas.org (NIH grant #OD010943 to D.H. Hall.), Aging

Atlas (Dr. M. Wang) and CenGen (cengen.org) for their immensely helpful resources.

The work was supported by NIH grants (R35GM139618 to D.K.M.), BARI Investigator

Award (D.K.M.), and UCSF PBBR New Frontier Research Award (D.K.M.)

**Author Contributions**

D.K.M. designed and analyzed the *C. elegans* experiments, contributed to project

conceptualization and wrote the manuscript. B.W. made the initial observation on *cyp-*

*14A5* induction by light and performed RNAi screens and reporter imaging. Z.J.

prepared RNA-seq samples and characterized light effects on the *cyp-14A5* reporter,

cellular, behavioral and physiological phenotypes of various mutants with assistance

from W.Y. and M.E. Y.L. analyzed the RNA-seq data. R.C. designed, performed and

analyzed the behavioral memory assays. J.L. designed, performed and analyzed the

light-induced sleep and phenoptotic aging assays. N.L., Y.L., B.W., Z.J., R.C., J.L.

contributed to research materials, project conceptualization and editing manuscript.

**Competing Interest Statement:** The authors declare no competing interests.

    **Figures and figure legends**

**Figure 1 Light exposure activates the CYP-encoding gene *cyp-14A5* in a genetic**

**program in *C. elegans*.** A, Representative epifluorescence and brightfield images

showing *cyp-14A5*p::GFP induction by light exposure (1500 Lux, 24 hrs), in

synchronized young adults (24 hrs post L4). Scale bar: 50 µm. B, Dot plot showing fold

induction of *cyp-14A5*p::GFP as a function of light intensity (Lux) and duration of light

(hours of light_dark indicated in Y axis). C, Schematic of *cyp-14A5* transcriptional and

translational GFP reporters. The translational reporter shows non-neuronal (major

tissues indicated by arrows) induction of CYP-14A5::GFP by light (1500 Lux, 24 hrs).

Scale bar: 50 µm. D, Volcano plot showing genes differentially regulated by light (1500

Lux, 24 hrs), with *cyp-14A5* highlighted, in synchronized young adults (24 hrs post L4).

E, Heat map of top-ranking visible light-regulated genes (top 30 including *cyp-14A5*).

**Figure 2 Light induction of *cyp-14A5*p::GFP requires the transcription factors ZIP-**

**2 and CEBP-2**. A, Representative epifluorescence images showing light-induced *cyp-*

*14A5*p::GFP expression in *lite-1 gur-3* double and *atm-1* single loss-of-function mutants.

Scale bar: 50 µm. B, Summary of RNAi screens identifying ZIP-2 and CEBP-2 as

essential transcriptional regulators of light-induced *cyp-14A5p::GFP* expression. C,

Representative epifluorescence images showing light-induced *cyp-14A5p::GFP*

expression in wild type, *zip-2* and *cebp-2* loss-of-function mutants. Scale bar: 50 µm. D,

Representative epifluorescence images showing light-induced *cyp-14A5p::GFP*

expression and no light-induced *irg-1p::GFP* expression in wild type animals. Scale bar:

50 µm. E, Representative epifluorescence images showing light-induced *cyp-*

*14A5p::GFP* expression unaffected by a UV shield. Scale bar: 50 µm. F, Representative

epifluorescence images showing light-induced *cyp-14A5p::GFP* expression by red,

green, blue LED light sources of equal intensities. Scale bar: 50 µm. G, Quantification of

E and F. *** indicates $P < 0.001$, n.s., non-significant. H, Representative

epifluorescence images showing light-induced *cyp-14A5p::GFP* expression unaffected

in *rpl-28p::zip-2uORF* transgenic animals. Scale bar: 50 µm. I, Representative

epifluorescence images showing light-induced *cyp-14A5p::GFP* expression abolished in

*eif-2alpha* mutants. Scale bar: 50 µm. J, Schematic model for light-induced transition of

*zip-2* mRNA from translating uORF to mORF, leading to increased ZIP-2 and

subsequent increased transcriptional *cyp-14A5* expression in cooperation with CEBP-2.

uORF and mORF are separated for clarity.

**Figure 3 Light promotes sleep behavior and memory consolidation via a ZIP-2/CYP axis from hypodermis**. A, Schematic of behavioral setup to test effects of dark on olfactory memory. B, Wild-type learning and memory after 16 hours of dark exposure. 7 trials, 50-200 animals per trial/condition. Two-way ANOVA shows significant differences in chemotaxis (CI) under ambient light conditions (approximately 600 Lux) but not when the assay plates were placed in the dark. Learning (LI) indices reflect the differences between buffer- and butanone-treated animals, abolished under dark (one-way ANOVA). Pairwise t-tests of the amount of memory retention under light and dark recovery reveal the degree of memory loss under dark. C, Dark exposure timeline shows that exposing animals to dark immediately after training (0-2 hr, Dark) shows hampered memory retention, whereas dark conditions for 2-4 hours period after two hours after ambient light recovery is less sufficient to induce memory loss (two-way ANOVA). The lack of differences between buffer and butanone-trained animals is reflected in the respective LIs (one-way ANOVA). D, Two different loss-of-function alleles of *zip-2(tm4246)* or *zip-2(ok3730)* both showed impaired memory, but learning remained intact (two-way ANOVA). 5-10 trials, 50-200 animals per trial/condition. E, Memory impairment of *cyp-14A5(gk152)* but not *F43C1.7* mutant animals (two-way ANOVA for CIs and One-way ANOVA for LIs. 7-14 trials, 50-200 animals per trial/condition. F, Memory defects of *cyp-14A5(gk152)* mutants are rescued by hypodermal expression of wild-type *cyp-14A5*. Two independent transgenic lines show similar results with comparison of pairwise differences in LIs and the amount of memory rescued by hypodermal *cyp-14A5* (Cis: Two-way ANOVA; Lis: One-way ANOVA). *

445 indicates *P* < 0.05, ** indicates *P* < 0.01, *** indicates *P* < 0.001, **** indicates *P* <

446 0.0001, n.s., non-significant.

**Figure 4 Light-induced gene expression drives organismal phenotypes, including**

**sleep and shortened lifespans**. A, Schematic of synthetic constructs for light-inducible

*nlp-22* and *zip-10::mCherry* using the *cyp-14A5* promoter. B, Representative compound

epifluorescence images showing light-induced *cyp-14A5p::zip-10::mCherry* activation.

Scale bar: 50 µm. C, Representative confocal fluorescence images showing light-

induced *cyp-14A5p::zip-10::mCherry* activation (1500 Lux, 48 hrs starting at 24 hrs post

L4) in major non-neuronal tissues (hypoderm, intestinal cells indicated by arrows). Scale

bar: 50 µm. D, Representative lifespan curves showing that light-induced *zip-10* can

markedly shorten lifespan. **** indicates *P* < 0.0001. E, Representative bright field

images showing quiescent sleep behaviors by light-induced *cyp-14A5p::nlp-22*

expression. F, Quantification of population bending frequencies for light-treated (1500

Lux, 48 hrs starting at 24 hrs post L4) control wild type and *cyp-14A5p::nlp-22* animals.

G, Quantification of population track lengths for control wild type and *cyp-14A5p::nlp-22*

animals with light (1500 Lux, 48 hrs starting at 24 hrs post L4) or darkness treatments.

*** indicates *P* < 0.001, **** indicates *P* < 0.0001, n.s., non-significant.

**Figure S1 Light for *cyp-14A5p::GFP* induction does not alter ambient**

**temperature.** A, Schematic of the setup for measurements of light intensity and

temperature at a plane (in parallel to LED light sources) where animals are exposed to

LED light in NGM plates. B, Measurements of temperature at the 600 Lux light intensity

showing no change of temperature over 24 hrs. C, Measurements of temperature at the

1500 Lux light intensity showing no change of temperature over 24 hrs. D,

Measurements of temperature at the 3000 Lux light intensity showing no change of

temperature over 24 hrs.

**A**

Gene ontology analysis
by WormCat:

**B**  Gene ontology analysis
by WormEnrichr:

**GO Cellular Component 2018**  ⓘ

membrane raft (GO:0045121)
microbody (GO:0042579)
collagen and cuticulin-based cuticle extracel
peroxisome (GO:0005777)
integral component of mitochondrial outer

**GO Molecular Function 2018**  ⓘ

non-membrane spanning protein tyrosine k
acyl-CoA oxidase activity (GO:0003997)
oxidoreductase activity, acting on the CH-CH
UDP-glycosyltransferase activity (GO:000819
protein tyrosine kinase activity (GO:0004713

**GO Biological Process 2018**  ⓘ

peptidyl-tyrosine autophosphorylation (GO:
enzyme linked receptor protein signaling pa
peptidyl-serine modification (GO:0018209)
transmembrane receptor protein tyrosine ki
defense response to bacterium (GO:0042743

**Anatomic Associations WormBase 2018**  ⓘ

male gonad(WBbt:0006794)
sperm(WBbt:0006798)
hermaphrodite gonad(WBbt:0005178)
uterus(WBbt:0006760)
cuticle(WBbt:0005755)

**Coexpression Predicted GO Biological Process**  ⓘ

sodium ion transmembrane transport (GO:0
protein autophosphorylation (GO:0046777)
eye pigment biosynthetic process (GO:00067
ocellus pigment biosynthetic process (GO:00
peptidyl-S-diacylglycerol-L-cysteine biosynth

**Coexpression Predicted GO Cellular Component**  ⓘ

pseudopodium (GO:0031143)
integral component of mitochondrial memb
chloride channel complex (GO:0034707)
intrinsic component of mitochondrial outer
integral component of mitochondrial outer

**Figure S2 Gene ontology analysis of light-induced transcriptomic changes.** A,

Starburst plot of gene ontology analysis of light-induced genes using WormCat

(http://www.wormcat.com/). B, Table summary of gene ontology analysis of light-

induced genes using WormEnrichr (https://maayanlab.cloud/WormEnrichr/).

**Figure S3 No apparent effect of short-term visible light exposure on development,**

**morphology and simple behaviors.** A, Representative brightfield images showing no

apparent effects of visible light exposure (1500 Lux, 24 hrs) on the body lengths and

gross morphology of wild type, *cebp-2*, and *zip-2* loss-of-function mutants. Scale bar: 50

μm. B. Representative images showing the movement tracks of wild type, *cyp-14A5*,

and *zip-2* loss-of-function mutants under dark and light (1500 Lux, 24 hrs) conditions. C,

Quantification of track lengths of wild type, *cyp-14A5*, and *zip-2* loss-of-function mutants

under dark and light (1500 Lux, 24 hrs) conditions. D, Quantification of pumping rates of

wild type, *cyp-14A5*, and *zip-2* loss-of-function mutants under dark and light (1500 Lux,

24 hrs) conditions. E, Quantification of defecation behaviors of wild type, *cyp-14A5*, and

*zip-2* loss-of-function mutants under dark and light (1500 Lux, 24 hrs) conditions.

**Figure S4 WormLab analysis reveals sleep bouts caused by light-induced *cyp-***

***14A5p::nlp-22* expression.** (A) Representative tracking of moving angles for body

posture and average speed for control animals under dark conditions (N=15). (B)

Representative tracking of moving angles for body posture and average speed for

control animals after light exposure conditions (1500 Lux, 24 hrs, N=13). (C)

Representative tracking of moving angles for body posture and average speed for *cyp-*

*14A5p::nlp-22* transgenic animals under dark conditions (N=12). (B) Representative

tracking of moving angles for body posture and average speed for *cyp-14A5p::nlp-22*

transgenic animals after light exposure conditions (1500 Lux, 24 hrs, N=13).

## References

**References**

1. Andrabi, M., Upton, B.A., Lang, R.A., and Vemaraju, S. (2023). An Expanding Role for Nonvisual Opsins

in Extraocular Light Sensing Physiology. Annu Rev Vis Sci *9*, 245–267.

https://doi.org/10.1146/annurev-vision-100820-094018.

2. Cronin, T.W., and Johnsen, S. (2016). Extraocular, Non-Visual, and Simple Photoreceptors: An

Introduction to the Symposium. Integr Comp Biol *56*, 758–763. https://doi.org/10.1093/icb/icw106.

3. Van Gelder, R.N. (2008). Non-visual photoreception: sensing light without sight. Curr Biol *18*, R38-39.

https://doi.org/10.1016/j.cub.2007.11.027.

4. Do, M.T.H., and Yau, K.-W. (2010). Intrinsically photosensitive retinal ganglion cells. Physiol Rev *90*,

1547–1581. https://doi.org/10.1152/physrev.00013.2010.

5. Shi, Y., Zhang, J., Li, X., Han, Y., Guan, J., Li, Y., Shen, J., Tzvetanov, T., Yang, D., Luo, X., et al. (2024).

Non-image-forming photoreceptors improve visual orientation selectivity and image perception.

Neuron, S0896-6273(24)00840-7. https://doi.org/10.1016/j.neuron.2024.11.015.

6. Lucas, R.J., Hattar, S., Takao, M., Berson, D.M., Foster, R.G., and Yau, K.-W. (2003). Diminished

pupillary light reflex at high irradiances in melanopsin-knockout mice. Science *299*, 245–247.

https://doi.org/10.1126/science.1077293.

7. Berson, D.M., Dunn, F.A., and Takao, M. (2002). Phototransduction by retinal ganglion cells that set

the circadian clock. Science *295*, 1070–1073. https://doi.org/10.1126/science.1067262.

8. Hattar, S., Lucas, R.J., Mrosovsky, N., Thompson, S., Douglas, R.H., Hankins, M.W., Lem, J., Biel, M.,

Hofmann, F., Foster, R.G., et al. (2003). Melanopsin and rod-cone photoreceptive systems account for

all major accessory visual functions in mice. Nature *424*, 76–81.

https://doi.org/10.1038/nature01761.

9. Bhatla, N., and Horvitz, H.R. (2015). Light and hydrogen peroxide inhibit C. elegans Feeding through

gustatory receptor orthologs and pharyngeal neurons. Neuron *85*, 804–818.

https://doi.org/10.1016/j.neuron.2014.12.061.

10.    Gong, J., Yuan, Y., Ward, A., Kang, L., Zhang, B., Wu, Z., Peng, J., Feng, Z., Liu, J., and Xu, X.Z.S.

(2016). The C. elegans Taste Receptor Homolog LITE-1 Is a Photoreceptor. Cell *167*, 1252-1263.e10.

https://doi.org/10.1016/j.cell.2016.10.053.

11.    Hanson, S.M., Scholüke, J., Liewald, J., Sharma, R., Ruse, C., Engel, M., Schüler, C., Klaus, A.,

Arghittu, S., Baumbach, F., et al. (2023). Structure-function analysis suggests that the photoreceptor

LITE-1 is a light-activated ion channel. Curr Biol *33*, 3423-3435.e5.

https://doi.org/10.1016/j.cub.2023.07.008.

12.    Edwards, S.L., Charlie, N.K., Milfort, M.C., Brown, B.S., Gravlin, C.N., Knecht, J.E., and Miller, K.G.

(2008). A novel molecular solution for ultraviolet light detection in Caenorhabditis elegans. PLoS Biol

*6*, e198. https://doi.org/10.1371/journal.pbio.0060198.

13.    Bhatla, N., and Horvitz, H.R. (2015). Light and hydrogen peroxide inhibit C. elegans Feeding

through gustatory receptor orthologs and pharyngeal neurons. Neuron *85*, 804–818.

https://doi.org/10.1016/j.neuron.2014.12.061.

14.    Mahmoud, B.H., Ruvolo, E., Hexsel, C.L., Liu, Y., Owen, M.R., Kollias, N., Lim, H.W., and Hamzavi,

I.H. (2010). Impact of long-wavelength UVA and visible light on melanocompetent skin. J Invest

Dermatol *130*, 2092–2097. https://doi.org/10.1038/jid.2010.95.

15.    De Magalhaes Filho, C.D., Henriquez, B., Seah, N.E., Evans, R.M., Lapierre, L.R., and Dillin, A.

(2018). Visible light reduces C. elegans longevity. Nat Commun *9*, 927.

https://doi.org/10.1038/s41467-018-02934-5.

16.    Liebel, F., Kaur, S., Ruvolo, E., Kollias, N., and Southall, M.D. (2012). Irradiation of skin with

visible light induces reactive oxygen species and matrix-degrading enzymes. J Invest Dermatol *132*,

1901–1907. https://doi.org/10.1038/jid.2011.476.

17.    Keller, J., Ellieva, A., Ma, D.K., Ju, J., Nehk, E., Konkel, A., Falck, J.R., Schunck, W.-H., and Menzel,

R. (2014). CYP-13A12 of the nematode Caenorhabditis elegans is a PUFA-epoxygenase involved in

behavioural response to reoxygenation. Biochem. J. *464*, 61–71.

https://doi.org/10.1042/BJ20140848.

18.    Ma, D.K., Rothe, M., Zheng, S., Bhatla, N., Pender, C.L., Menzel, R., and Horvitz, H.R. (2013).

Cytochrome P450 drives a HIF-regulated behavioral response to reoxygenation by C. elegans. Science

*341*, 554–558. https://doi.org/10.1126/science.1235753.

19.    Denisov, I.G., Makris, T.M., Sligar, S.G., and Schlichting, I. (2005). Structure and chemistry of

cytochrome P450. Chem Rev *105*, 2253–2277. https://doi.org/10.1021/cr0307143.

20.    Neve, E.P.A., and Ingelman-Sundberg, M. (2010). Cytochrome P450 proteins: retention and

distribution from the endoplasmic reticulum. Curr Opin Drug Discov Devel *13*, 78–85.

21.    Ciccia, A., and Elledge, S.J. (2010). The DNA damage response: making it safe to play with knives.

Mol Cell *40*, 179–204. https://doi.org/10.1016/j.molcel.2010.09.019.

22.    Schuch, A.P., Moreno, N.C., Schuch, N.J., Menck, C.F.M., and Garcia, C.C.M. (2017). Sunlight

damage to cellular DNA: Focus on oxidatively generated lesions. Free Radic Biol Med *107*, 110–124.

https://doi.org/10.1016/j.freeradbiomed.2017.01.029.

23.    Estes, K.A., Dunbar, T.L., Powell, J.R., Ausubel, F.M., and Troemel, E.R. (2010). bZIP transcription

factor zip-2 mediates an early response to Pseudomonas aeruginosa infection in Caenorhabditis

elegans. Proc Natl Acad Sci U S A *107*, 2153–2158. https://doi.org/10.1073/pnas.0914643107.

24.    Reddy, K.C., Dunbar, T.L., Nargund, A.M., Haynes, C.M., and Troemel, E.R. (2016). The C. elegans

CCAAT-Enhancer-Binding Protein Gamma Is Required for Surveillance Immunity. Cell Rep *14*, 1581–

1589. https://doi.org/10.1016/j.celrep.2016.01.055.

25.    Dunbar, T.L., Yan, Z., Balla, K.M., Smelkinson, M.G., and Troemel, E.R. (2012). C. elegans detects

pathogen-induced translational inhibition to activate immune signaling. Cell Host Microbe *11*, 375–

386. https://doi.org/10.1016/j.chom.2012.02.008.

26.    Ma, Z., Horrocks, J., Mir, D.A., Cox, M., Ruzga, M., Rollins, J., and Rogers, A.N. (2023). The

integrated stress response protects against ER stress but is not required for altered translation and

lifespan from dietary restriction in Caenorhabditis elegans. Front Cell Dev Biol *11*, 1263344.

https://doi.org/10.3389/fcell.2023.1263344.

27.    Costa-Mattioli, M., and Walter, P. (2020). The integrated stress response: From mechanism to

disease. Science *368*, eaat5314. https://doi.org/10.1126/science.aat5314.

28.    Brito Querido, J., Díaz-López, I., and Ramakrishnan, V. (2024). The molecular basis of translation

initiation and its regulation in eukaryotes. Nat Rev Mol Cell Biol *25*, 168–186.

https://doi.org/10.1038/s41580-023-00624-9.

29.    Mir, D.A., Ma, Z., Horrocks, J., and Rogers, A. (2024). Stress-Induced Eukaryotic Translational

Regulatory Mechanisms. J Clin Med Sci *8*, 1000277.

30.    Chandra, R., Farah, F., Muñoz-Lobato, F., Bokka, A., Benedetti, K.L., Brueggemann, C., Saifuddin,

M.F.A., Miller, J.M., Li, J., Chang, E., et al. (2023). Sleep is required to consolidate odor memory and

remodel olfactory synapses. Cell *186*, 2911-2928.e20. https://doi.org/10.1016/j.cell.2023.05.006.

31.    Kauffman, A.L., Ashraf, J.M., Corces-Zimmerman, M.R., Landis, J.N., and Murphy, C.T. (2010).

Insulin signaling and dietary restriction differentially influence the decline of learning and memory

with age. PLoS Biol *8*, e1000372. https://doi.org/10.1371/journal.pbio.1000372.

32.    Wang, C., Long, Y., Wang, B., Zhang, C., and Ma, D.K. (2023). GPCR signaling regulates severe

stress-induced organismic death in Caenorhabditis elegans. Aging Cell *22*, e13735.

https://doi.org/10.1111/acel.13735.

33.    Pandey, T., and Ma, D.K. (2022). Stress-Induced Phenoptosis: Mechanistic Insights and

Evolutionary Implications. Biochemistry (Mosc) *87*, 1504–1511.

https://doi.org/10.1134/S0006297922120082.

34.    Bringmann, H. (2018). Sleep-Active Neurons: Conserved Motors of Sleep. Genetics *208*, 1279–

1289. https://doi.org/10.1534/genetics.117.300521.

35.    Van der Auwera, P., Frooninckx, L., Buscemi, K., Vance, R.T., Watteyne, J., Mirabeau, O.,

Temmerman, L., De Haes, W., Fancsalszky, L., Gottschalk, A., et al. (2020). RPamide neuropeptides

NLP-22 and NLP-2 act through GnRH-like receptors to promote sleep and wakefulness in C. elegans.

Sci Rep *10*, 9929. https://doi.org/10.1038/s41598-020-66536-2.

36.    Nelson, M.D., Trojanowski, N.F., George-Raizen, J.B., Smith, C.J., Yu, C.-C., Fang-Yen, C., and

Raizen, D.M. (2013). The neuropeptide NLP-22 regulates a sleep-like state in Caenorhabditis elegans.

Nat Commun *4*, 2846. https://doi.org/10.1038/ncomms3846.

37.    Troemel, E.R., Chu, S.W., Reinke, V., Lee, S.S., Ausubel, F.M., and Kim, D.H. (2006). p38 MAPK

regulates expression of immune response genes and contributes to longevity in C. elegans. PLoS

Genet *2*, e183. https://doi.org/10.1371/journal.pgen.0020183.

38.    Wong, D., Bazopoulou, D., Pujol, N., Tavernarakis, N., and Ewbank, J.J. (2007). Genome-wide

investigation reveals pathogen-specific and shared signatures in the response of Caenorhabditis

elegans to infection. Genome Biol *8*, R194. https://doi.org/10.1186/gb-2007-8-9-r194.

39.    Sinha, A., Rae, R., Iatsenko, I., and Sommer, R.J. (2012). System wide analysis of the evolution of

innate immunity in the nematode model species Caenorhabditis elegans and Pristionchus pacificus.

PLoS One *7*, e44255. https://doi.org/10.1371/journal.pone.0044255.

40.    Genuth, N.R., and Barna, M. (2018). Heterogeneity and specialized functions of translation

machinery: from genes to organisms. Nat Rev Genet *19*, 431–452. https://doi.org/10.1038/s41576-

018-0008-z.

41.    D'Orazio, K.N., and Green, R. (2021). Ribosome states signal RNA quality control. Mol Cell *81*,

1372–1383. https://doi.org/10.1016/j.molcel.2021.02.022.

42.    Sinha, N.K., McKenney, C., Yeow, Z.Y., Li, J.J., Nam, K.H., Yaron-Barir, T.M., Johnson, J.L.,

Huntsman, E.M., Cantley, L.C., Ordureau, A., et al. (2024). The ribotoxic stress response drives UV-

mediated cell death. Cell *187*, 3652-3670.e40. https://doi.org/10.1016/j.cell.2024.05.018.

43.    Zhang, B., Gong, J., Zhang, W., Xiao, R., Liu, J., and Xu, X.Z.S. (2018). Brain-gut communications

via distinct neuroendocrine signals bidirectionally regulate longevity in C. elegans. Genes Dev *32*,

258–270. https://doi.org/10.1101/gad.309625.117.

44.    Fukuda, A., Sato, K., Fujimori, C., Yamashita, T., Takeuchi, A., Ohuchi, H., Umatani, C., and Kanda,

S. (2025). Direct photoreception by pituitary endocrine cells regulates hormone release and

pigmentation. Science *387*, 43–48. https://doi.org/10.1126/science.adj9687.

45.    Liu, Y., Zhou, J., Zhang, N., Wu, X., Zhang, Q., Zhang, W., Li, X., and Tian, Y. (2022). Two sensory

neurons coordinate the systemic mitochondrial stress response via GPCR signaling in C. elegans. Dev

Cell *57*, 2469-2482.e5. https://doi.org/10.1016/j.devcel.2022.10.001.

46.    Aghayeva, U., Bhattacharya, A., Sural, S., Jaeger, E., Churgin, M., Fang-Yen, C., and Hobert, O.

(2021). DAF-16/FoxO and DAF-12/VDR control cellular plasticity both cell-autonomously and via

interorgan signaling. PLoS Biol *19*, e3001204. https://doi.org/10.1371/journal.pbio.3001204.

47.    Stringham, E.G., Dixon, D.K., Jones, D., and Candido, E.P. (1992). Temporal and spatial

expression patterns of the small heat shock (hsp16) genes in transgenic Caenorhabditis elegans. Mol

Biol Cell *3*, 221–233. https://doi.org/10.1091/mbc.3.2.221.

48.    Monsalve, G.C., Yamamoto, K.R., and Ward, J.D. (2019). A New Tool for Inducible Gene

Expression in Caenorhabditis elegans. Genetics *211*, 419–430.

https://doi.org/10.1534/genetics.118.301705.

49.    Wei, X., Potter, C.J., Luo, L., and Shen, K. (2012). Controlling gene expression with the Q

repressible binary expression system in Caenorhabditis elegans. Nat Methods *9*, 391–395.

https://doi.org/10.1038/nmeth.1929.

50.     Brenner, S. (1974). The genetics of Caenorhabditis elegans. Genetics *77*, 71–94.

51.     Hobert, O. (2002). PCR fusion-based approach to create reporter gene constructs for expression

analysis in transgenic C. elegans. Biotechniques *32*, 728–730. https://doi.org/10.2144/02324bm01.

52.     Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose program

for assigning sequence reads to genomic features. Bioinformatics *30*, 923–930.

https://doi.org/10.1093/bioinformatics/btt656.

53.     Kim, D., Langmead, B., and Salzberg, S.L. (2015). HISAT: a fast spliced aligner with low memory

requirements. Nat. Methods *12*, 357–360. https://doi.org/10.1038/nmeth.3317.

54.     Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550. https://doi.org/10.1186/s13059-

014-0550-8.

660