The 3D genomics of lampbrush chromosomes highlights the role of active transcription in chromatin organization

T. Lagunov[1,2]†, M. Gridina[1,2]†, A. Nurislamov[1,2,5], T. Kulikova[3], A. Maslova[3], V. Konstantinov[1,2], A. Popov[1,4], A. Krasikova[3,*], V. Fishman[1,2,4,5,*]

1 - Institute of Cytology and Genetics, Novosibirsk, Russia
2 - Novosibirsk State University, Novosibirsk, Russia
3 - Laboratory of Cell Nucleus Structure and Dynamics, Saint-Petersburg State University, Saint-Petersburg, Russia
4 - Artificial Intelligence Research Institute, Moscow, Russia
5 - Sirius University of Science and Technology, Sirius, Russia
†-equal contribution
*-correspondence should be sent to Veniamin Fishman (minja-f@yandex.com) and Alla Krasikova (alla.krasikova@gmail.com)

Abstract

Lampbrush chromosomes are giant meiotic bivalents in growing oocyte nuclei that have served as a classic model system for studying chromatin organization and RNA synthesis for over a century. Despite their importance, the molecular mechanisms underlying lampbrush chromosomes formation and their distinctive chromomere-loop architecture have remained poorly understood. Here, we provide the first comprehensive genomic, cytological, and biophysical analysis of lampbrush chromosome organization by integrating single-cell Hi-C, RNA-seq, NOMe-seq, FISH mapping, and chromatin simulations. Single-nucleus Hi-C analysis revealed CTCF-independent contact domains with stable boundaries defined by transcription units in a convergent orientation. Contact domains identified through Hi-C analysis correspond to insulated chromomeres in lampbrush chromosomes. Small transcriptionally inactive contact domains surrounded by transcription units in the diverged orientation form 'chromatin knots', which are often detached from the chromosome axis. Transcription loops frequently manifest as a 'cross' pattern with reduced contacts within chromatin domains. Integrative analysis of the whole-genome data uncovers the mechanisms underlying lampbrush chromosome structure, revealing how hypertranscription modulates chromatin stiffness and repositions SMC complexes to establish the distinctive chromomere-loop organisation. Biophysical modeling through polymer simulation reproduces key features of lampbrush chromosomes, including transcription loop formation, chromomere compaction, and insulation patterns. These findings offer a unifying framework for understanding the remarkable chromatin architecture of lampbrush chromosomes and their transcription-dependent organization.

Highlights

- First integration of single-cell Hi-C, RNA-seq, NOMe-seq and microscopy methods uncovers molecular mechanisms underlying lampbrush chromosome architecture.
- Hi-C reveals contact patterns corresponding to lampbrush chromomeres and transcription loops, validated through BAC-based FISH mapping.
- Lampbrush chromosomes are segmented into contact domains formed via a CTCF-independent mechanism, with boundaries coinciding with convergently oriented gene pairs.
- Hypertranscription shapes lampbrush chromosome through multiple mechanisms, increasing stiffness and decreasing compaction of transcribed units, generating outward pressure, pushing transcription loops away from the chromosome axis, and repositioning SMC complexes to form transcription-dependent domains with stable boundaries.
- Polymer simulations including these mechanisms replicate key features of lampbrush chromosomes organization, supporting the proposed model of their formation.

Key words

Chromatin architecture, chromatin domain, chromomere, cohesion, contact domain, hypertranscription, lampbrush chromosomes, loop extrusion, oocyte nucleus, polymer simulation, single-cell Hi-C, transcription loops

59   **Graphical Abstract**

Main

Chromosomes are folded non-randomly in the nuclear space. Their spatial organization is essential for several genomic processes ([1–3]), such as regulation of gene expression ([4–7]), DNA replication ([8–10]) and repair ([11]), and mechanical properties of the nucleus ([12,13]). Consistent with this pivotal role in genome regulation, spatial organization of chromatin was shown to be conserved across large evolutionary distances of hundreds of million years ([14–17]).

There are several essential mechanisms responsible for chromatin organization in animal cells. First, DNA is a polymer organized in the nuclear space according to the principles of polymer physics. This implies that frequency of contacts between loci decreases with increase of genomic distance, following power law. For the majority of the studied cell types, scaling of contacts with genomic distance is concordant with the "crumpled" states of the chromatin ([18]), characterized by largely unknotted conformations with a fractal dimension of ~3 ([19]). Second, multivalent homotypic interactions between proteins cause segregation of the chromatin into compartments with distinct epigenetic states ([20–24]). Moreover, active and inactive chromatin is directed towards different nuclear landmarks ([2]). Inactive chromatin is often associated with nuclear lamina and nucleolar periphery, while active chromatin is in proximity to the interchromatin space comprising nuclear speckles ([25–27]). Third, ATP-dependent extrusion of DNA by SMC-complexes cohesin and condensin results in chromosome compaction and increased probability of long-range interactions ([1,28,29]). The extrusion process can be modulated by several factors. In the majority of vertebrate cell types, CTCF protein blocks cohesin-mediated extrusion, therefore CTCF binding sites demarcate compact self-interacting genomic regions, often referred to as topologically associating domains (TADs) ([1,30–33]). Interactions of cohesin with other proteins, including replication-specific MCM ([34]) or transcription-associated RNA-polymerase II complexes ([35–38]) also may modulate extrusion, although details of these mechanisms are less studied. Moreover, interaction of cohesin with MCM and RNA-polymerase have less impact on genomic contacts compared to cohesin-CTCF interactions, and their effect becomes pronounced only when artificially modulated ([37]) or in analysis of specific cell cycle stages ([34]).

Although chromatin contacts in the majority of vertebrate cells can be explained by above mentioned mechanisms with relatively high accuracy ([39,40]), specific cell types or cell cycle stages show distinct arrangement of chromatin ([2]). This includes early stages of development, prior to or immediately after zygotic genome activation, and germ cells ([41]). Importantly, chromatin architecture is uniquely reorganized during the oocyte-to-zygote transition. In mammals, features of genomic organization at diplotene stage include compartments, TADs, and loops ([41,42]). Although these chromatin domains can be detected when averaged over the genome, the presence of each type of domain at a locus varies between cells ([42]). TAD boundaries in murine oocytes are located concordant with somatic cells, demarcated by CTCF, and associated with cohesin-mediated loop extrusion ([42]). Compartments in murine diplotene oocytes manifest as plaid-pattern of Hi-C contacts and demonstrate strong association with distribution of H3K27me3 mark ([41]). Compared to somatic cells, in oocytes interactions within compartments are restricted to the regions located less than 10 Mb away from each other, presenting a "local

compartmentalization" pattern ([41]). Similar depletion of long-range interactions and local compartmentalization was also observed during meiotic division of sperm cells ([43,44]). Strength of both TADs and compartments gradually decreases with oocyte transition from early to late diplotene stage, and these structures disappear at metaphase-II (MII) stage ([41,42]).

Previous whole-genomic studies focused on chromatin organization in oocytes did not include representatives of vertebrate taxa in which chromatin undergoes lampbrush chromosome state. Lampbrush chromosomes (LBCs) represent a specific form of nuclear chromatin that emerges at the diplotene stage of oogenesis in the majority of vertebrate species, including amphibians, fish, reptiles, birds (Figure 1B), monotremes, and certain insects. In the 3D space of the giant oocyte nucleus, lampbrush chromosomes do not contact each other, are not tethered to the nuclear periphery, and do not have a preferred radial position ([45,46]). Lampbrush chromosomes consist of separate chromatin globules connected by a thin thread and protruding lateral loops, together forming chromomere-loop complexes ([47–49]). The unusually enormous size of LBCs allows for chromomere-loop structure visualization using conventional light microscopy, making diplotene oocytes a preferred model for microscopy-based studies of chromatin architecture.

The phenomenon of LBC organization is attributed to hypertranscriptional activity of the oocyte nucleus ([50,51]). Hypertranscription results in formation of lateral loops tightly covered by elongating RNA polymerases and nascent transcripts ([49,52,53]) and containing single or multiple genes essential for maternal products biosynthesis ([54–57]). Another structural unit of LBC are lampbrush chromomeres – compact insulated chromatin globules present along LBC axes ([48]). Chromomeres do not clearly correspond to any of the known spatial chromatin domains of the interphase nucleus. Genomic loci of neighbouring somatic TADs can be detected as a part of a single chromomere, while genomic regions belonging to one somatic TAD can be detected as a part of two adjacent chromomeres ([55,58]). There is, however, partial correspondence between gene-poor heterochromatic regions of constitutive B-compartments identified in somatic cells and clusters of large DAPI-positive chromomeres, which are also enriched for inactive chromatin marks ([58–62]).

Overall, chromatin organization in LBCs drastically differs from somatic cells, asking whether there are unique mechanisms of genome architecture that are active at this stage.

Here, we present the first Hi-C analysis of lampbrush chromosomes, supplemented by transcriptomic and epigenomic profiling. We show that chromatin contacts in LBCs display unique features, including restriction of long-range and inter-chromosomal interactions and formation of CTCF-independent chromatin domains and loops. We show strong correlation between location and orientation of transcription units and formation of chromatin domain boundaries. Based on our observations, validated by FISH and other complementary analysis, we develop a physical model of LBCs explaining LBC architecture via interplay between SMC and RNA-polymerase complexes. We show that the polymer model can qualitatively explain observed Hi-C patterns genome-wide and accurately reproduce spatial configuration of individual loci. Finally, the developed model

highlights the importance of the transcription-mediated ribonucleoprotein matrix accumulation and nucleosome removal as factors determining spatial structure of LBC.

Results

Isolation and quality control of chicken diplotene oocytes

To investigate chromatin organization in avian lampbrush chromosomes, we selected chicken oocytes at the lampbrush stage based on follicle size. Individual nuclei were isolated and processed using a previously established single-cell Hi-C protocol ([63]) (Figure 1A). Multiple quality control measures were conducted to validate the integrity and reliability of the resulting data.

To confirm that formaldehyde fixation preserves the lampbrush chromosome structure, we stained fixed nuclei with SYTO 61 dye and visualized them using confocal microscopy. The analysis demonstrated the characteristic lampbrush morphology of the captured chromosomes (Figure 1C, Figure S1C), consistent with previous findings ([45]).

Next, we ensured that oocyte genomic material was not contaminated by somatic cells. Chicken follicles include various cell types, such as erythrocytes, which are significantly smaller than oocyte nuclei. To exclude contamination, nine randomly selected oocyte nuclei were stained with DAPI post-isolation and visually inspected. Microscopy analysis revealed no DAPI signals outside oocyte chromosomes. As a control, erythrocytes were manually added to oocyte samples, confirming that erythrocyte nuclei are detectable during visual inspection (Supplementary Figure S1A, B).

To further validate the absence of contamination, we leveraged the fact that a single oocyte nucleus contains no more than four copies of each genomic fragment. Consequently, no more than eight unique interactions should be detected per *Dpn*II restriction fragment (two ends per fragment, [64]). Genomic fragments exhibiting more than eight interactions would indicate contamination or Hi-C artifacts. Our analysis showed that the vast majority of fragments exhibit fewer than nine interactions (Supplementary Figure S1E). These results confirm that the isolated oocyte nuclei were free of contamination from other chicken cell types.

Single-nucleus Hi-C experiments were performed on a total of 96 oocyte nuclei. Initial library quality was evaluated through shallow sequencing (~200,000 reads per sample), using metrics such as the fraction of unmapped reads, duplicate reads, intra-chromosomal reads, and intra-fragment reads (see "Methods"). Based on these criteria, 25 single-cell oocyte libraries were selected for deeper sequencing (~3–40 million read pairs per nucleus) and downstream analysis (Supplementary Table S1).

The resulting Hi-C maps achieved >80% bin coverage at 4 Kb resolution for most samples (80% of the 4 Kb bins had at least one read or more), with a few exceptions where one or more chromosomes were completely absent but the remaining genome showed consistent coverage (Supplementary Figure S1D). Deeply sequenced Hi-C maps displayed a high degree of similarity across individual oocyte nuclei (Figure 3A, B), confirming the high quality of the obtained data.

Figure 1. Experimental workflow preserves structural integrity of chicken lampbrush chromosomes.

A) Experimental workflow, including oocyte dissection, oocyte separation by size (lampbrush chromosome stage oocyte: 1–2 mm diameter; post-lampbrush chromosome stage oocyte: 4–7 mm diameter), microsurgical isolation of nuclei, and preparation of single-nucleus Hi-C libraries. Selected oocytes and nuclei were randomly chosen for quality control.

B) Chromomere-loop structure of isolated chicken lampbrush chromosome 1. Elongating RNA polymerase II is labeled with antibodies (green), and chromatin is counterstained with DAPI (blue).

C) Lampbrush chromosomes preserve their 3D-structure after formaldehyde fixation. The entire nucleus is visualized after fixation and staining with SYTO 61.

Lampbrush chromosomes form elongated structures with minimal inter-chromosomal interactions

Lampbrush chromosomes at the diplotene stage of meiosis are known to form elongated structures that are spatially separated within the giant oocyte nucleus ([45]). This spatial arrangement is also preserved after fixation, as shown in Figure 1C and Figure S1C. Hi-C data analysis further confirms these unique features of LBCs.

Our results show a strikingly low frequency of inter-chromosomal (trans) contacts, with only 0.2–1% of spatial interactions occurring between non-homologous chromosomes

(Figure 2C, Supplementary Table S1). This is markedly lower than the ~10% reported for chicken fibroblasts ([15], Figure 2C) and the 10–30% typical of somatic cells in various species ([14,65], Figure 2C). These data underscore the exceptional rarity of inter-chromosomal interactions in lampbrush-stage oocytes (Figure 2C, Supplementary Table S1).

Intra-chromosomal (cis) interactions in lampbrush chromosomes exhibit an abrupt drop in contact frequency beyond ~1 Mb (Figure 2 A-B, and Supplementary Figure S2), consistent with the elongated, rope-like morphology of these chromosomes (Figure 1B). The depletion of both long-range and trans-chromosomal interactions in LBC Hi-C maps results in the absence of the plaid-pattern of contacts typically observed in interphase nucleus Hi-C data, which is associated with chromatin compartmentalization ([66], Figure S3; see "Methods").

Figure 2. Chromatin organization of lampbrush chromosomes at different genomic distances. A) Contact probability versus the genomic distance for single and merged oocyte data (lampbrush chromosome stage, LBC) and chicken embryonic fibroblasts (see "Methods"). The genomic distances are cut to display only representative contacts of oocytes. See extended version in Figure S2.

B) Hi-C contact maps comparing pseudo-bulk chicken oocytes and bulk fibroblast cells for chromosome 11. The fibroblast map was downsampled to match oocytes data (both maps sum to 19,946 contacts within the depicted region). Each bin represents 32 Kb. See extended version in Figure S2.

C) Fraction of inter-chromosomal contacts across different cell types and species (see "Methods"). Chicken oocytes at the lampbrush chromosome stage exhibit a markedly reduced proportion of inter-chromosomal contacts, consistent with cytological observations of spatial chromatin organisation.

Chromatin of lampbrush chromosomes forms domains with stable boundaries shared across individual oocytes

In contrast to the absence of a plaid-pattern, chromatin domains are clearly visible on LBC Hi-C maps at scales of 0.5–2 Mb (Figure 2B and 3B-D). To understand whether the domain organization observed in chicken LBCs represents a general feature of meiotic chromatin or a unique, potentially LBC-specific mechanism, we compared our results with mouse diplotene oocytes. Available Hi-C maps for individual nuclei of mouse diplotene oocytes, obtained by a similar approach ([42]), provide an opportunity to compare contact chromatin maps in organisms with different types of oogenesis.

In mouse diplotene oocytes at the non-surrounded nucleolus (NSN) stage, chromatin interaction analysis reveals TADs with highly variable boundary locations across individual cells ([42]). In contrast, single-nucleus Hi-C maps from chicken lampbrush-stage oocytes demonstrate striking consistency in domain boundary positions (Figure 3A-C), with a mean F1 score of 0.48 compared to 0.23 for mouse NSN oocytes (Figure 3A). This stability in domain boundaries across chicken LBCs suggests fundamental differences in the mechanisms underlying chromatin domain formation between these two systems.

Contact domains in LBCs can be classified into two types based on their length (Figure 3D,E): 1) Large Unstructured Domains (LUDs), defined as domains longer than 1.3 Mb (N50 of domains length, see "Methods") and 2) Small Squared Domains (SSDs), shorter than 1.3 Mb. Visually, LUDs have more dispersed contacts that are connecting loci up to ~2 Mb from each other (Figure S7A,B). Almost all LUDs are located on macro (chr1-chr5) and Z chromosomes (Figure 3H).

To characterize chromatin states in LUDs and SSDs, we performed a single cell NOMe-seq experiment. NOMe-seq measures open chromatin via deposition of exogenous GpC methylation, enabling simultaneous analysis of chromatin accessibility and CpG methylation levels in individual nuclei. We augmented NOME-seq data with CpG methylation from [[67]] (see "Methods"). We also quantified transcription levels, reanalyzing RNA-seq data from the previous chicken oocyte nucleus transcriptome study ([57]) and supplementing it with additional replicas generated in this work. Analysis of these data shows that LUDs have lower median gene density (Supplementary Figure S4D) and lower normalized expression levels than SSDs (Figure 3G). We found that SSDs have significantly higher levels of chromatin openness (measured in NOME-seq experiment via GpC methylation), indicating that shorter domains contain more accessible chromatin (Supplementary Figure S4F), while there is almost no difference in CpG methylation levels between SSDs and LUDs (Figure 3F, Figure S4E), as well as only a weak

correlation between promoter methylation and gene expression levels (Supplementary Figure S14).

Our findings indicate that Large Unstructured Domains correspond to gene-poor, transcriptionally inactive chromatin regions predominantly found on macrochromosomes and the Z chromosome. In contrast, Small Squared Domains are associated with gene-rich, transcriptionally active chromatin regions distributed across all chromosomes.

Figure 3. Chromatin domains in chicken lampbrush chromosomes.
A) Pairwise F1 scores for domain boundary concordance across individual oocytes. Left panel: chicken lampbrush-stage oocytes; right panel: mouse NSN oocytes ([42]). Oocytes are sorted by the number of Hi-C contacts. Higher F1 scores indicate greater concordance in boundary positions. The mean F1 score

is 0.48 for chicken oocytes and 0.23 for mouse oocytes. See Supplementary Figures S4–S8 and methods for additional details.

B) Domain annotations for individual cells and merged Hi-C data for chicken (left) and mouse (right) oocytes. Note the higher concordance of domain boundaries in chicken oocytes compared to mouse oocytes.

C) Aggregated Hi-C contacts and domain annotations for 14 chicken oocytes (left) and 34 mouse oocytes (right).

D) Merged Hi-C data from chicken oocytes with domain annotations overlaid. Domains were manually refined after automatic annotation (see "Methods"). The dashed line indicates the N50 domain size (~1.3 Mb).

E) Each horizontal segment represents a single chromatin domain, with the Y-position of the line indicating the domain's average characteristic (CpG methylation, GpC methylation, or AT fraction, see "Methods"). The dashed horizontal line represents the genome-wide average for each characteristic. This visualization highlights the distinct properties of Large Unstructured Domains (LUDs) and Small Squared Domains (SSDs). The genome-wide CpG and GpC methylation distributions can be found in Supplementary Figure S4E. The genome-wide AT fraction distribution can be found in Supplementary Figure S9D.

F) CpG and GpC methylation levels statistically compared between SSDs and LUDs (see "Methods"). As a baseline, domains were randomly redistributed across the genome while preserving the number of intervals and their length distributions (indicated as 'random'). SSDs exhibit significantly higher GpC methylation levels, reflecting increased chromatin accessibility. The genome-wide CpG and GpC methylation distributions can be found in Supplementary Figure S4E.

G) The transcription units (TUs) density and activity distribution show that the SSDs contain significantly more TUs, and TUs with higher activity, than LUDs.

H) Distribution of LUDs and SSDs lengths on chicken chromosomes.

Hi-C patterns align with chromatin units of lampbrush chromosomes identified by microscopy analysis

Lampbrush chromomeres are compact chromatin structures enriched in self-interactions and insulated from neighboring regions. Based on this characteristic, we propose that SSDs and LUDs in single-nucleus Hi-C maps correspond to chromomeres in LBCs. Additionally, given their properties, we speculate that LUDs represent more compact, DAPI-positive chromomeres, while SSDs correspond to less compact, DAPI-negative chromomeres. Supporting this hypothesis, statistical analysis of the AT fraction in domains—reflecting the genomic proportion susceptible to DAPI staining—shows a weak but significant correlation between AT fraction and domain length (Spearman's R = 0.55, compared to R = 0.28 for randomly shuffled domains, Supplementary Figure S9D).

To further validate the correlation between domain types and DAPI staining in chromomeres, we analyzed sequence markers with known positions in cytological maps of chicken LBCs, primarily derived from DNA- and RNA-FISH experiments using specific BAC clones (Figure 4A,B). The distribution of LUDs and SSDs correlated with the arrangement of DAPI-positive and DAPI-negative chromomeres at certain regions of chromosomes. For instance, we observed multiple LUDs between markers W030P07 and W030B21, coinciding with a region enriched in DAPI-positive chromomeres (Figure 4A,B). However, the number of cytologically visible chromomeres and contact domains on Hi-C maps does not always match exactly, probably due to the limited resolution of microscopy analysis and/or errors in domain callers. Together with sparse availability of sequence markers distinguishing chromomeres and resolution of cytological chromomere-loop maps this limited a genome-wide analysis of the relationship between SSDs, LUDs, and chromomeres.

A Chromosome 1 domains

W030P07   W030B21   WAG31B10   WAG67J15   P1H9   CH261-191J12   CH261-120O20
CH261-87J17   WAG13E20   CH261-162E14   CH261-54J7
CH261-54H10   WAG69C11   CH261-33I15
WAG53E23

SSDs
LUDs

0 MB   20 MB   40 MB   60 MB   80 MB   100 MB   120 MB   140 MB   160 MB   180 MB   200 MB

B Chromosome 1 domains

SSDs
LUDs

CH261-191J12   CH261-120O20
CH261-54J7

145 MB   160 MB   175 MB   190 MB

C Resolution 4 KB

59.5 MB

59.75 MB

60.0 MB

RNA-seq reads   288   0
Hi-C reads   61   0

59.5 MB   59.75 MB   60.0 MB
chr1

D

-6 gene len   0   +6 gene len

1.5
1
0.7

Forward transcription
Reverse transcription
Bin coverage

Figure 4. Concordance between microscopically visible chromatin structures and Hi-C map patterns along chicken lampbrush chromosomes

A) Comparison of Hi-C contact domains with chromatin domains identified through microscopy visualization of lampbrush chromosomes. From top to bottom: LBC Hi-C map for chicken chromosome 1, showing domain boundaries and DNA marker positions; Contact domain segmentation, with each triangle representing a single domain. LUDs and SSDs are distinguished by triangle colors, and black lines indicate DNA marker locations; Cytological map of chromomeres distribution along chicken lampbrush chromosome 1 ([68]), with DAPI-positive chromomeres shown as black circles and DAPI-negative chromomeres as white circles; the average lateral loop length in a region is shown by rectangles; DAPI-stained image of chicken lampbrush chromosome 1 (a fire look-up table corresponding to the intensity of fluorescence is used) (scale bar: 20 µm).

B) Zoomed region of a Hi-C map for a fragment of lampbrush chromosome 1 presented on panel A.

C) Hi-C map displaying an example of a "cross" pattern. The "cross" region overlaps with a high RNA-seq signal and shows no reduction in overall Hi-C coverage (denoted as "bin coverage").

D) Rescaled observed-over-expected Hi-C contact map averaged around the longest (top 50% by length) and most transcriptionally active (top 50% by FPKM) genes. A characteristic cross-shaped contact pattern is observed, extending approximately ±6 gene lengths from the gene end. Data represent aggregated signals from multiple genes at 2 kb resolution.

Transcription loops are a hallmark of lampbrush chromosomes. These loops represent stretched fibers of the actively transcribed chromatin emanating from the LBC axis. Given their spatial separation from each other and the LBC axis, we hypothesize that loci corresponding to transcription loops lack interactions with the rest of the genome.

Visual inspection of Hi-C maps reveals loci with a distinct "cross" pattern, where interactions are limited to nearest neighbors (Figures 4C,D). These loci compensate lack of long-range interactions by increasing frequency of proximal interactions, thus maintaining consistent coverage levels (Figure 4C). "Cross" patterns frequently occur within chromatin domains and almost always coincide with actively transcribed genes. Supporting this observation, aggregated Hi-C maps centered on long, highly expressed genes (top 50% by length and TPM) show a pronounced "cross" pattern (Figure 4D). These findings suggest that "cross" patterns in Hi-C maps represent the contact signatures of transcription loops in LBCs.

To align lampbrush chromatin contact patterns identified by Hi-C analysis with chromatin structures observable via microscopy, we conducted FISH mapping using probes specific to chromatin domain boundaries, internal domain regions, and loci forming "cross" patterns. A total of 29 BAC clones were hybridized in various combinations on lampbrush chromosomes 1, 4, and 13 ("Methods", Supplementary Table S3). The large size of lampbrush chromosomes enables detailed microscopy analysis, allowing us to distinguish chromomeres and transcription loops, as well as determine the orientation of transcription units.

The locus analyzed on chromosome 1 contains six Hi-C domains. Regions 4, 6, 7, 9, and 12 coincide with "cross" patterns observed in Hi-C maps and the corresponding actively transcribed genes (Figure 5A). Based on the annotated transcription units' location and orientation, we anticipated transcription loops with forward-oriented transcription for regions 4, 9, and 12, and reverse-oriented transcription for regions 6 and 7. FISH mapping

of LBC1 using BAC clone-based probes with a DNA+RNA hybridization protocol confirmed these predictions. Long lateral loops with a prominent RNP matrix were observed for BACs 4, 9, and 12, with transcription co-directional to the chromosome (Figure 5B, subpanels 2a, 3a-b). Similarly, genomic regions 6, 7, and 9, covered by three different BACs, correspond to transcription units forming one or two pairs of lateral loops (Figure 5B, subpanels 2c–e). BAC 5 doesn't correspond to a region with an obvious "cross" pattern according to the Hi-C data, however it covers the transcriptionally active region (in reverse direction), thus it can form a transcription loop (Figure 5B subpanels 2a-b). In our FISH experiments, BAC 5 marks a reverse strand transcription loop anchored in the chromomere (Figure 5A,C). Additionally, RNA-FISH experiments without chromosomal DNA denaturation for several transcribed regions confirmed that BAC clone-based probes hybridize with nascent transcripts on the lateral loops. These findings confirm that "cross" patterns observed on Hi-C maps of avian LBCs correspond to the microscopically visible lampbrush lateral loops containing actively transcribed genes.

The intra-domain regions 1, 2, 3, 8, 10, and 14 correspond to five of the six contact domains identified on LBC Hi-C maps, with BACs 2 and 3 covering the same domain. Consequently, we anticipated observing five insulated chromomeres along the chromosome axis in microscopy images, one of which would be associated with two BACs. FISH experiments confirmed this, with BAC clone-based probes mapping to five distinct chromomeres (Figure 5B, subpanels 1a–b). The chromomeres covered by BAC 2 and BAC 3 are closely spaced but remain distinguishable (Figure 5B, subpanels 1c–d).

Interestingly, the compact chromatin domain covered by BAC 8, which lacks transcription, exhibits variable positioning. This chromatin domain can either anchor directly to the chromosome axis or remain detached from it (indicated by white arrows in Figure 5B panels 1a-e and 3a-c), forming a compact "knot" on a lateral loop emanating from the chromosomal axis. These chromatin knots, located within transcriptionally active lateral loops, have been previously described as small genomic regions devoid of transcribed genes (Figure 6 in [57]).

Together, these results demonstrate that (1) chromatin contact domains identified through Hi-C analysis correspond to insulated chromomeres in lampbrush chromosomes, and (2) small regions between actively transcribed genes can form microdomains (knots) that are optionally tethered to the chromosomal axis.

Region 11 exhibits partial transcriptional activity and is positioned near the boundaries of chromatin domains (Figure 5A). Based on this, we anticipated that BAC 11 would cover a portion of chromatin spanning two adjacent chromomeres, one of which is covered by BAC 10. FISH mapping confirmed this prediction, showing partial colocalization of BACs 10 and 11 within a prominent DAPI-positive chromomere (Figure 5B, subpanels 1a–d and 3d-e).

Similarly, BAC 13, located near the boundary of contact domain, produced a signal spatially close to that of BAC 14, which maps to the same domain. While the signals from BACs 13 and 14 are closely positioned, they remain distinct and do not intermix.

These results confirm that boundaries between domains identified by Hi-C are concordant with LBC chromomere boundaries (see additional evidence supporting this statement in the "Supplementary Note 2").

On lampbrush chromosome 4, we mapped nine BAC clone-based probes across two regions of interest. In region A (51–52 Mb), Hi-C data revealed the formation of a contact domain with interactions at the top of the domain triangle and "cross" patterns within domain, indicative of chromatin loop formation (Figure 5D). Nuclear RNA-seq data showed that several genes within region A are transcribed in varying orientations. FISH mapping confirmed that this genomic region corresponds to a long transcription loop with multiple RNP matrix gradients (Figure 5E,F), consistent with the transcriptional activity detected. Detailed results for region B are provided in "Supplementary Note 2" and Supplementary Figure S10.

We integrated the FISH data to reconstruct the chromatin organization of the analyzed region on chromosome 1 (Figure 5C), accounting for two alternative states of chromatin knots discussed above. This reconstruction demonstrates a strong concordance between microscopically visible chromatin structures—such as chromomeres and transcription loops—and the contact domains and "cross" patterns observed in Hi-C maps. The reconstruction of the region A on chromosome 4 also confirmed the correspondence between the transcription loops and the "cross" patterns (Figure 5D,E).

Additionally, similar analyses of two other regions on chromosomes 4 and 13, which included BACs mapped to Hi-C domain boundaries, intra-domain fragments, and "cross" patterns, yielded consistent results (Supplementary Figure S10; "Supplementary Note 2"). These findings validate the Hi-C maps of chicken oocyte nuclei and confirm the interpretation of Hi-C patterns as representations of lampbrush chromosome chromomeres and lateral loops.

Figure 5. Comparison of Hi-C map patterns with microscopically identified chromatin domains and loops in lampbrush chromosomes 1 and 4

A) Hi-C map and associated data for a fragment of chromosome 1. From top to bottom: combined Hi-C map of the oocyte nuclei, with colored highlights indicating interactions in regions covered by BAC clone-based probes; Total stranded RNA-seq data for lampbrush-stage oocyte nuclei ([57]); Annotation of transcription units based on *de novo* transcriptome analysis (see "Methods"); BAC clone numbers and names (see Supplementary Table S3).

B) FISH mapping images for different combinations of BAC clone-based probes on lampbrush chromosome 1. All panels except a1 include DAPI signals (blue or grayscale). Panels a1, e2, a3 and d3 merge fluorescent images with phase contrast images. White arrows in panels 1a-e and 3a-c indicate a compact chromatin domain without transcriptional activity (covered by BAC 8) that may either anchor to the chromosome axis or remain insulated from the axis forming an isolated knot. Arrowheads in panels 3d-e indicate partial colocalization of genomic regions covered by BACs 10 and 11 within a compact chromomere.

C) Schematic representation of chromomere-loop organization for a fragment of lampbrush chromosome 1, derived from FISH mapping data. Numbers correspond to the locations of BAC clone-based FISH probes aligned on chromosome 1.

D) Hi-C map for region A on chromosome 4. The data is presented similarly as in panel A above.

E) Schematic representation of chromomere-loop organization for region A on lampbrush chromosome 4, based on FISH mapping data. Numbers correspond to the locations of BAC clone-based FISH probes aligned on region A.

F) FISH mapping images for two BAC clone-based probes in region A on lampbrush chromosome 4. Fluorescent images are merged with DAPI-stained chromatin (grayscale) or phase contrast images. Scale bars: 10 µm.

Chromatin domains in lampbrush chromosomes are formed by a CTCF-independent mechanism

In the majority of vertebrate cell types, including chicken fibroblasts ([15]), TAD boundaries are demarcated by CTCF binding sites ([1,66]). The moderate conservation of CTCF binding site distribution across cell types, combined with a strong correlation between CTCF binding and TAD boundaries, allows CTCF enrichment to be detected even when comparing Hi-C maps to CTCF ChIP-seq data from different cell types ([15]) and cell-type agnostic motif enrichment.

We show that LBC domain boundaries display almost no enrichment of CTCF motifs (Figure 6A). Concordantly, almost no insulation was detected when LBC Hi-C contacts were aggregated across CTCF motifs (Figure 6B). To provide a ground for comparison, we performed the same analysis for chicken fibroblasts data subsampled to the same depth. In fibroblasts, we detect strong insulation across CTCF motifs and enrichment of motifs at TAD boundaries (Figure 6A-B). We next replicated this analysis using CTCF ChIP-seq signal from chicken liver, which confirms the absence of CTCF-mediated insulation in LBC (Figure 6C). In contrast, strong enrichment of chicken liver CTCF ChIP-seq signal was observed at fibroblast TAD boundaries (Figure 6C), validating the approach and reinforcing the conclusion that chromatin contact domains in lampbrush chromosomes are formed independently of CTCF binding.

To understand which chromatin architecture proteins can contribute to the formation of contact domain boundaries in oocytes, we explored the chicken oocyte nucleus

transcriptome data (see "Methods"). Despite the absence of CTCF binding enrichment in the domain boundaries, in oocytes *CTCF* gene has the comparable transcript levels with fibroblasts. Genes for the other architectural proteins, such as *RAD21, STAG1, STAG2* (cohesin) and *NCAP-H, NCAP-D2, NCAP-G, NCAP-D3, NCAP-H2, NCAP-G2* (condensin) subunits also show comparable transcript levels in fibroblasts and oocytes (Figure 6D, Supplementary Table S4). Consistent with this, the presence of SMC1A, SMC3, Rad21, STAG1, and STAG2 on avian LBC axes in a punctate pattern has previously been shown by immunostaining, with an enrichment of cohesin subunits in the interchromomere regions and in chromomere cores ([69]). Among the condensin subunits, XCAPD2 was found to accumulate in chromomeres of Xenopus LBCs ([70]). However, the main cohesin subunits (SMC3, SMC1A) and the main condensin subunits (SMC2, SMC4) have about ten times weaker transcript levels in the LBC stage oocyte (see Figure 6D, Supplementary Table S4). In contrast, the meiosis specific cohesin subunit SMC1B in oocytes has an almost thousand times higher level of transcripts than in fibroblasts (Figure 6D, Supplementary Table S4). Thus, although components of extrusion complexes are present at LBC-stage, their activity might be modulated by quantity of proteins and usage of cell type specific subunits, potentially explaining why extrusion is not blocked by CTCF binding sites as in typical somatic cells. Moreover, the synaptonemal complex protein SYCP3 has been found at the axes of avian LBCs ([69]), which may affect the formation of structural chromatin loops.

Figure 6. The mechanism of chromatin contact domain formation in lampbrush chromosomes is CTCF independent, despite transcriptional activity of the *CTCF* gene.

A) Frequency distribution of CTCF-motifs at domain boundaries in lampbrush chromosome (LBC) and chicken embryonic fibroblast (CEF) maps. Both LBC and CEF boundaries show enrichment relative to the genome average, with peak enrichment factors of ~1.4 and ~2.3, respectively.
B) Aggregated Hi-C contact maps centered on CTCF-motifs with uniform orientation. While strong insulation is evident around CTCF-motifs in CEF, only weak insulation is observed in LBC contact maps. Resolution: 10 kb, fibroblast data subsampled to match number of Hi-C contacts in merged LBC.
C) Input-normalized CTCF ChIP-seq signal from chicken liver (SRR4068194, see "Methods") at oocyte contact domain boundaries and fibroblast contact domain boundaries. The absence of CTCF enrichment across LBC domain boundaries contrasts with the strong signal observed in fibroblast boundaries.
D) Comparison of transcript levels for cohesin subunits, condensin subunits, and *CTCF* (TPM fold-change relative to the somatic cell median). Oocytes exhibit reduced transcript levels for *SMC3, SMC1A, SMC2,* and *SMC4* genes*,* higher levels for *SMC1B* and *REC8*, and similar *CTCF* transcript levels compared to fibroblasts. Gene annotations were sourced from ncbiRefSeq, except for *STAG3*, which used ensGene annotations (see Supplementary Table S4). Green dots represent transcript levels from three chicken oocyte RNA-seq datasets, and black dots represent five replicates of chicken embryonic fibroblasts ([71]). Oocytes 1: Total RNA stranded RNA-seq data ([57]). Oocytes 2 and 3: Poly(A)-enriched stranded RNA-seq replicates generated in this study.

Transcriptional activity and orientation of transcription units define lampbrush chromosome chromatin domains

To investigate the genomic features underlying the formation of LBC domains, we
compared domain borders with position-dependent genomic elements. CpG methylation
levels and chromatin accessibility at domain boundaries were similar to the genomic
average (Supplementary Figure S9E, see "Methods"). However, boundaries showed
enriched GC-content relative to the background, suggesting a correlation with the
presence of genes.

Aggregation of contacts around exons, introns, and repetitive elements revealed no
insulatory potential for these features, arguing against their role in forming domain
boundaries at the lampbrush stage (Supplementary Figure S11). In contrast, we observed
a significant depletion of contacts across 3′-UTRs and, to a lesser extent, across 5′-UTRs
(Figure 7A). This suggests that the distribution and orientation of transcription units may
play a key role in LBC domain formation.

To explore the potential link between transcription and LBC domain boundaries, we
aligned RNA-seq data from poly(A)-enriched and total stranded RNA libraries ([57]) with Hi-
C maps and visually inspected these profiles. This analysis highlighted transcriptional
activity as a critical determinant of LBC chromatin domain organization.

Visual comparison of transcriptomic data with chromatin domain boundaries revealed
multiple transcription units absent from the current gene annotations in [[73]]
(Supplementary Figure S17). This discrepancy reflects the dependence of chicken
genome annotation on cell type and database, particularly for non-coding RNAs [[74]]. To
address this, we performed *de novo* transcriptome assembly using oocyte RNA-seq data,
generating a refined set of transcription units (TUs) active during the LBC stage (see
"Methods").

Analysis of this updated TU set confirmed a strong correlation between chromatin insulation and TU orientation (Figure 7D). Genomic regions flanked by TUs in a convergent (Forward-Reverse, FR) orientation exhibited the highest insulation scores compared to other TU pair orientations. Furthermore, FR-oriented TU pairs were significantly enriched at chromatin domain boundaries (Figure 7B).

Manual validation of domain boundaries revealed that nearly half of non-FR TU pairs initially identified at domain boundaries were artifacts of automatic domain annotation or transcriptome assembly (26 of 47 randomly selected for inspection boundaries; Supplementary Figure S18). Overall, convergent transcription accounts for at least 79% of domain boundaries, with 62% supported by automatic annotation and an additional 17% identified through manual correction.

Given the central role of transcription in defining LBC chromatin domains, we designate the domains detected in Hi-C maps as **T**ranscription-**D**ependent **D**omains (**TDDs**).

Although the majority of TDD boundaries are formed by TU pairs in a convergent (FR) orientation, only ~15% (~20% after manual curation) of FR-oriented TU pairs create boundaries, while the remainder are located within domains (Figure 7C). To investigate the properties of TU pairs associated with boundary formation, we compared transcript features of "border" versus "non-border" TU pairs in a convergent orientation (Figure 7E). No single feature was sufficient to distinguish "border" TU pairs from "non-border" pairs. However, we observed that TU pairs forming boundaries tend to have several distinguishing characteristics: they are, on average, longer, separated by a larger intergenic interval, and exhibit higher activity of the weaker TU of the pair. The activity of the stronger TU in the pair shows minimal difference between boundary-forming and non-boundary-forming pairs. These observations suggest that TDD boundary formation is governed by a complex interaction of TU features rather than a single determinant.

Figure 7. Aggregated 3D chromatin interaction profiles reveal genomic features underlying lampbrush chromosome chromatin domains

A) Aggregated contact maps around 5′ and 3′ UTRs show depletion of interactions across these regions. Insulation scores are displayed in the upper-left corner of each aggregated map triangle (see "Methods"). Resolution: 10 Kb

B) Pie charts displaying transcription unit (TU) orientations for all TU pairs and those forming domain boundaries. A significant enrichment of Forward-Reverse (FR, convergent) orientation is observed at domain boundaries.

C) Statistics of FR-oriented TU pairs. Despite their prevalence at domain boundaries, only ~15-20% of FR-oriented TU pairs form boundaries.

D) Aggregated contact maps centered between TUs in different orientations: Forward-Forward (FF, co-directional), Reverse-Reverse (RR, co-directional), Forward-Reverse (FR, convergent) and Reverse-Forward (RF, divergent). Insulation score, shown in the upper-left corner of each map, is highest for convergent (FR) TU pairs. RF pairs typically form a "cross" pattern, while FR, FF, and RR pairs display a stripe-like enrichment of contacts between TUs and upstream loci, with FF and RR stripes widening with distance from the diagonal. Insulation scores are displayed in the upper-left corner of each aggregated map triangle (see "Methods"). Resolution: 10 Kb

E) Comparison of TU features between border-forming and non-border FR-oriented TUs. Border-forming TUs are, on average, longer, more highly expressed, and separated by greater intergenic distances. However, the length of the shorter TU and the activity of the weaker TU in the pair show higher difference between border-forming and non-border FR-oriented TUs than the features of the longer or more strongly expressed TU. p-values obtained using Mann–Whitney U test.

465 *Interplay between chromatin extruders and RNA-polymerase complex explains formation*
466 *of transcription-dependent domains in lampbrush chromosomes*

The presence of cohesin and condensin subunits on LBCs, combined with the enrichment of TDD boundaries with convergent transcription units, lead us to the hypothesis that interactions between chromatin extrusion complexes and RNA polymerase II drive LBC domain formation. This model, previously proposed in the context of bacterial chromosomes [35,75] and mouse chromosomes under Wapl/CTCF-depleted conditions [37], suggests that RNA polymerase acts as a steric barrier for extruders.

Inspired by the RNA polymerase-extruders interaction model described in [37], we developed a coarse-grained model of LBC chromatin as beads-on-string, incorporating key adaptations to reflect unique LBC features (see "Methods"): 1) we connected two sister chromatids by passive cohesive cohesin, representing cohesin without extrusion activity; 2) DNA segments encoding actively transcribed genes are modeled with reduced linear compaction, reflecting their nucleosome-free state; 3) Locus-specific RNA polymerase density is incorporated based on observed TU FPKM values; 4) Bead sizes were adjusted, using the polymer model from [76], to account for the volume of ribonucleoprotein complexes associated with transcription; and 5) A simple square-distance repulsion force was applied to beads to simulate chromatin interactions. These modifications allow the model to capture critical features of transcription-driven LBC organization (see also Supplementary Note 3 and Figure S13).

We compared modeled and observed contact maps for single transcription units, stratified by gene length and expression levels. Aggregated Hi-C maps for long transcribed genes show insulatory boundaries at the 5′ UTR and a distinctive "cross" pattern formed by the TU body (Figure 8A). Our model successfully reproduces these patterns (Figure 8A).

While insulation caused by actively transcribed genes has been previously reported and modeled [37], prior studies did not observe the "cross" pattern centered around individual genes. We propose that this pattern arises from two factors: nucleosome removal and ribonucleoprotein matrix repulsion. Nucleosome removal linearly elongates the chromatin segment, while the RNP matrix repulsion spatially separates the active transcription loop from other chromatin regions. Even when a small part of the TU loop is attracted to the chromomere by loop-extruding factors (LEFs), the majority of the TU loop DNA remains separated from the chromomere due to RNP matrix repulsion. This mechanism effectively explains the observed "cross" patterns.

In addition to nucleosome removal and RNP matrix repulsion, our model incorporates modulation of LEFs speed due to interactions with RNA polymerase. This property makes TUs asymmetric, directionally permeable — that is, permeable in only one direction — barriers for extrusion complexes and gives rise to distinct chromatin interaction patterns for convergent, divergent, and co-directional TU pairs.

For example, in the case of convergent TU pairs, the extruder's "legs" are effectively "locked" between the TUs (Figure 8C-E), resulting in chromatin between the TUs predominantly interacting with itself. Furthermore, loci upstream and downstream of the convergent TU pair are isolated from each other, as one of the extruder's "legs" becomes "locked" upon encountering a TU. This mechanism provides a robust explanation for the characteristic insulation and interaction patterns observed in convergent TU pairs.

To validate the hypothesis that LEF speed decreases due to interactions with RNA polymerase, we compared Hi-C data across different TU pair orientations with predictions from our model (Figure 8B). For convergent TU pairs, the model accurately reproduces the stripe pattern observed at domain boundaries and matches the experimental insulation score. The model also explains why the activity of the weaker TU in a pair is the most critical feature: if the weaker TU is sufficiently active to act as a directional barrier, the pair can halt the extruder's "leg" and enforce insulation.

The "cross" pattern characteristic of divergent TU pairs is also captured by the model, albeit less prominently, and the model correctly reflects the lower insulation score of divergent TUs compared to convergent TUs (1.17 for divergent vs. 1.33 for convergent). As noted earlier, the stripe for co-directional TUs becomes thicker with increasing distance from the diagonal, and this feature is partially reproduced by the model. The insulation score for modeled co-directional TU pairs is unexpectedly higher than that for modeled convergent TU pairs at a resolution of 5 Kb (1.48 vs. 1.33, respectively; Figure 8B), although it is lower for other resolutions (Supplementary Figure S12A,B).

Overall, these results confirm that the model accurately reproduces the distinct interaction patterns of transcription unit pairs observed in Hi-C data. We conclude that only the convergent (FR) orientation of TUs can generate the strong insulation characteristic of transcription-dependent chromatin contact domains. This is because the FR orientation "locks" one or both extruder "legs" between the TUs, effectively insulating upstream chromatin from downstream regions.

Figure 8. Lampbrush chromosome modeling and comparison of simulated results with experimental data

A) Aggregated contact maps for single transcription units (TUs). All TUs were rescaled to occupy 1/12 of the figure width, supplemented with a window of ±6 TU lengths around the TU end. Resolution: 2 Kb.

B) Aggregated contact maps centered on the midpoint between neighboring TUs with different orientations. A window of ±500 Kb around the midpoint was used. Insulation scores are displayed in the upper-left corner of each aggregated map triangle (see "Methods"). For both A and B, the upper triangles represent experimental data, and the lower triangles show simulation results. The color bar values chosen for better contrasting the contact patterns. Resolution: 5 Kb.

C) Schematic representation of aggregated contact maps for a single TU, along with diagrams illustrating extruder velocities when moving "in" or "against" the transcription direction.

D) Schematic representation of aggregated contact maps centered on the midpoint between TUs with different orientations.

E) Extruders "legs" positions at different time points for each TU pair orientation

Fusion of transcription-dependent chromatin domains during oocyte growth and decreased transcriptional activity

Growth of chicken oocyte coincides with reduced transcriptional activity and gradual contraction of both chromosomes and lateral loops ([77]). To study the dynamics of this process, we isolated nuclei from the late-diplotene stage oocytes (hereinafter referred to as post-LBC) and performed a single-nucleus Hi-C experiment (Figure 1A, Supplementary Table S1,2). Overall, Hi-C patterns detected in the mid-diplotene (LBC) stage persist at the post-LBC stage. However, there were quantitative differences. Concordant with cytological evidence of LBC contraction, we observe an increase of long-range interactions at the post-LBC stage (Figure 9A,B; Supplementary Figure S9B,C). Moreover, the significant decrease of the loop sizes of the chicken post-LBC chromosomes ([77]) agrees well with the almost complete absence of the "cross" patterns (Figure 9F). This observation confirms our suggestions about the critical role of the TU loop repulsion in the "cross" pattern formation.

Despite the absence of "cross" patterns, domain patterns can be identified on the Hi-C maps of late-diplotene oocytes, and the majority of the boundaries coincide with TDD boundaries of LBC oocytes. However, post-LBC Hi-C maps contain fewer boundaries, resulting in increase of domain lengths (Figure 9C). Visual inspection shows that about half of the neighboring LBC TDDs were fused at the post-LBC stage (44 borders out of 97 examined on chromosome 3, Figure 9B). Genome-wide quantification shows that interaction frequency between neighboring domains in post-LBC oocytes are also increased (11% of contacts between domains in normal oocytes vs 22% of contacts for late oocytes). Additionally, analysis of the border of TU pairs shows that the activity of weakly expressed TUs decreases, as well as does the insulation (Figure 9D,E). Interestingly, despite the absence of the "cross" pattern of actively transcribed TUs, the insulation by the convergently oriented TUs is still present (Figure 9F).

Figure 9. Comparison of oocyte chromatin at the lampbrush and post-lampbrush stages.
A) Contact probability as a function of genomic distance for lampbrush (LBC) and post-lampbrush chromosome (post-LBC) stages. Post-LBC chromatin exhibits an increase in distant contacts, consistent with microscopy observations ([77]).
B) Pseudo-bulk Hi-C interaction maps for LBC (8719 contacts) and post-LBC (4949 contacts) chromatin. Grey lines indicate domain borders for each stage. Neighboring domains fuse in the post-LBC stage, as demonstrated in the shown examples.
C) Distribution of domain sizes across different chromosome types. The post-LBC stage shows an increase in mean domain size alongside a reduction in the total number of domains.

D) Transcriptional activity of TUs located at domain borders at LBC and post-LBC stages.

E) Distribution of insulation scores (see "Methods") for domain borders at LBC and post-LBC stages. Lower scores indicate stronger insulation.

F) Aggregated observed-over-expected Hi-C contact maps comparing lampbrush chromosome (LBC) stage (upper triangle) and post-LBC stage (lower triangle) chromatin. The characteristic cross-shaped contact enrichment associated with strong single transcription units (TUs) is markedly reduced in the post-LBC stage, whereas convergently oriented TUs continue to delineate insulating domain boundaries. The first two panels depict genes rescaled to span 1/12 of the figure width, with flanking regions extending ±6 TU lengths from gene ends (2 kb resolution). The third panel shows aggregated maps for convergently oriented gene pairs within a ±500 kb window around their midpoint (5 kb resolution). Insulation scores are indicated in the upper-left corner (see "Methods").

Discussion

559 *A unified model of lampbrush chromosomes formation.* Our study offers the first comprehensive, genome-wide analysis of chromatin architecture in chicken oocytes during the lampbrush chromosome stage. Despite lampbrush chromosomes being discovered over a century ago ([78,79]), our findings provide a new understanding of their architecture and function at unprecedented resolution and across the entire genome. By integrating these results with established knowledge and applying polymer physics, we propose a unified model elucidating the mechanisms underlying lampbrush chromatin organization (Figure 10).

567 Hypertranscription is the primary driver of lampbrush chromosome formation, which determines their chromomere-loop organisation ([57]). Here we highlight several mechanisms linking transcriptional activity to their distinctive organization. First, the accumulation of RNP complexes along the transcription axis increases chromatin stiffness ([51,76,80]) and generates pressure that pushes transcription units outward from the chromatin axis. Second, RNA-polymerases travel along a DNA template as a tightly spaced convoy, displacing DNA-bound proteins from transcription units ([81,82]). The removal of histones, combined with increased chromatin stiffness, extends the length of transcription loops.

576 Not all proteins, however, can be displaced by RNA polymerase. Cohesive cohesin, which topologically entraps DNA to hold sister chromatids together, most probably resists disassembly ([83]). Instead, RNA polymerase translocates SMC complexes along the DNA ([83–85]), functioning as a motor that slides cohesive cohesin toward the ends of transcription units ([83]). Similarly, cohesin with extrusion activity is translocated or disassembled during transcription ([36–38]). This creates directional barriers for the extrusion machinery at transcription unit ends, either through direct interactions with the transcription complex or indirectly via accumulated cohesive cohesin, which blocks reloading of dynamic cohesin extruders ([86]).

585 In lampbrush chromosomes, active transcription zones remain free of cohesive cohesins, locally decoupling sister chromatids. In contrast, downstream regions of transcription units enriched in cohesive cohesin remain tethered to the chromosomal axis (Figure 10).

Figure 10 Mechanisms underlying the formation of chromomere-loop complexes in lampbrush chromosomes.

Active transcription increases chromatin stiffness and generates pressure that pushes transcription units (TUs) outward from the chromatin axis. Elongating RNA-polymerases remove histones and translocate cohesive cohesin towards the ends of transcription units. In the case of convergent transcribed gene pairs, this results in accumulation of cohesive cohesin complexes between transcription units, anchoring two transcription loops to the chromosomal axes. In the case of divergent transcribed gene pairs, the depletion of cohesive cohesin rings between transcription units results in the detachment of the condensed chromatin knot from the chromatin axes. Loop-extruding SMC complexes active within transcriptionally-silent regions form compact chromatin domains located on chromosomal axis. Transcriptionally-silent regions separated by divergently oriented transcription units can be bridged by extruders moving co-directionally with transcription and/or by intrachromatid entrapment by non-extruding cohesin. However, separation by convergently oriented transcription units forms impenetrable barrier for extrusion due to cohesive cohesin accumulation, precluding interactions between flanking

> regions. These mechanisms underlie the segmentation of lampbrush chromosome into distinct chromomere-loop complexes.

A defining feature of LBCs is the presence of transcription loops, both monogenic and
multigenic, varying in length ([57]). Integration of Hi-C maps, oocyte nuclear RNA-seq data,
and FISH mapping reveals that transcription loops often appear as Hi-C "cross" patterns,
characterized by reduced contacts with adjacent regions. This pattern is consistent with
loops being histone-free transcription units extruded from the chromatin axis due to
ribonucleoprotein pressure, increased stiffness, and cohesive cohesin depletion (Figure
10).

Transcriptionally inactive regions are enriched with extrusion complexes and cohesive
cohesins, leading to their compaction and appearance as contact domains with long-
range interactions (Figure 10). FISH analysis shows that transcriptionally inactive regions
correspond to intrachromomeric loci (this study and [57,61]). Our model of inactive
chromatin organisation within chromomeres is based on previous models of lampbrush
chromomere structure that involve cohesive cohesin and condensin complexes in
maintaining lampbrush chromosome axes ([50,87–89]). In particular, cohesin complex
subunits have been revealed within chromomere cores and in interchromomeric regions
([69]), while condensin subunit has been found to be enriched within chromomeres ([70]).
Moreover, we have taken into account microscopic observations of the rosette structure
of discrete lampbrush chromomeres separated by interchromomeric fibre ([48]).

Our data reveal how transcription-dependent repositioning of SMC complexes
establishes boundaries between chromomere-loop complexes (Figure 10). A single
transcribed gene creates a unidirectional barrier at its termination site but does not fully
insulate flanking regions, as extrusion complexes can move co-directionally with
transcription. Similarly, clusters of closely spaced co-directional genes result in SMC
complex accumulation at the end of the last transcription unit, allowing residual
interactions between adjacent regions.

In contrast, convergent gene pairs concentrate cohesive cohesin complexes between
transcription units, forming an impenetrable barrier and anchoring two transcription loops
to the chromosomal axis (Figure 10). Consistent with our Hi-C and cytological data,
convergent gene pairs define boundaries between chromomere-loop complexes. Longer
genes in a convergent pair increase the likelihood of at least one cohesive cohesin
molecule being translocated to the intergenic region, reinforcing the boundary. Thus, the
probability of boundary formation scales with gene length.

Unlike intergenic regions between convergent gene ends, those separating divergent
gene starts are depleted of cohesive cohesin (Figure 10). Extrusion complexes and
histones loaded within these regions mediate their compaction. These regions are
tethered to the chromosomal axis if residual cohesive cohesin is present or remain
untethered and manifest as a part of lateral loop if it is absent (Figure 10). Consistent with
this, we observed small contact domains corresponding to cytologically visible chromatin

knots between adjacent transcription units. These knots frequently occur when divergent genes are transcribed in opposite directions from a compact chromatin domain ([57]), aligning with the described mechanisms.

Extrusion complexes loaded between divergent gene pairs can traverse the genes co-directionally with transcription, bridging flanking regions. Thus, a divergent gene pair can form a loop without strong insulation of neighboring loci. In contrast, as discussed above, convergent gene pairs block extrusion complexes due to cohesive cohesin accumulation between them (Figure 10). This mechanism explains the observed enrichment of convergent gene pairs at contact domain boundaries.

Simulations based on these mechanisms using a physical chromatin model accurately reproduce both the genome-wide statistical properties of chromatin contact distribution and the specific packaging observed at individual loci.

Opened questions. Although the obtained data unequivocally shows that interaction between RNA polymerases and SMC complexes is the major mechanism explaining LBC formation, nuances of this mechanism remain to be discovered.

It is not clear how many extrusion complexes are involved in LBC organization and what is their subunit composition. It is established that the XCAP-D2 (condensin I subunit) and SMC1A, SMC3, Rad21, STAG1, and STAG2 (cohesin subunits) are present on the lampbrush chromosomes ([69,70]). However, it remains unclear whether RAD21 functions solely within cohesive cohesin or also as a part of extrusive cohesin. This leaves the possibility of one to four types of extruders contributing, including cohesin-STAG1, cohesin-STAG2, condensin I, and condensin II, each potentially varying in lifetime, density, and velocity.

From the observed peak positions of the P(s) slope (Supplementary Figure S9A-C), we estimate an extruder density of approximately 1 per 100–200 Kb in our models. This is comparable to the density of condensin II in interphase or condensin I in prometaphase ([90]).

Another open question concerns the probability of an extruder traversing chromatin in the direction opposite to RNA polymerase. Banigan et al. ([37]) assumed the permeability of RNA polymerase to SMC with probability ~0.01 s^-1, but our analysis suggests that this value is too low to reproduce LBC Hi-C data. In LBCs, RNA polymerase complexes are tightly spaced ([82]). Given a polymerase velocity of ~100 bp/sec [[91]] and the extruder velocity of 675 bp/sec [[92]] results in the overstep probability of ~0.15. This oversteps probability indicates that the extruder would frequently become blocked within the gene body. Therefore, the permeability must exceed 0.15 for extruders moving against the direction of transcription to pass through the gene body effectively. This increased permeability would allow extruders to traverse tightly packed transcription units, aligning with the observed chromatin organization in LBCs.

Our model highlights the critical role of cohesive cohesin relocation in response to transcriptional activity, alongside interactions between RNA polymerase and extrusion complexes, in shaping LBC architecture. Interestingly, in yeast, transcription has been

shown to promote chromatin loop formation even in the absence of loop extrusion ability in the cohesin variants ([85]). The authors propose that cohesin can facilitate intrachromatid loop formation by topologically entrapping chromatin near transcription sites. Similarly, the hypertranscription characteristic of the lampbrush chromosome stage could contribute to chromatin loop formation through a transcription-dependent mechanism involving DNA loop capture by cohesin, independent of extrusion activity. This suggests that cohesive cohesins may partially supplement the role of extrusion complexes in establishing LBC architecture, emphasizing a combined mechanism of loop extrusion and transcription-dependent loop capture in shaping these remarkable chromatin structures (Figure 10).

Another unresolved question is why CTCF-mediated insulation, a hallmark of chromatin organization in typical somatic cells, does not shape LBC contact domains in oocytes of birds in contrast to mouse diplotene oocytes at the NSN (non-surrounded nucleolus) stage ([42]). One hypothesis is that alterations in cohesin subunits during LBC formation, such as unknown post-translational modifications, could result in complexes that lose their CTCF-interacting ability. An alternative explanation is that CTCF is unable to bind active chromatin in LBCs due to the high density of RNA polymerase complexes occupying these regions. This transcription-driven exclusion could prevent CTCF from establishing its characteristic insulatory boundaries. Further investigation is required to determine the precise mechanisms underlying this phenomenon.

Principles underlying lampbrush chromosome organization are universal across cell types. Although lampbrush chromosomes (LBCs) exhibit a unique type of chromatin organization, the mechanisms underlying their structure are universal and can be observed across various cell types. For instance, transcription loops formed by long, highly expressed genes in somatic interphase nuclei show extensive externalization and reduced cis-contacts, resembling LBC transcription loops ([80]). Similarly, loss of contacts or chromatin 'melting' was observed in the case of highly expressed long neuronal genes ([94]). LBC transcription loops also resemble actively transcribed genes in polytene chromosomes, which appear as puffs ([53]). Hi-C maps of *Drosophila* polytene chromosomes revealed absence of TADs or chromatin contact domains in the puffed regions ([95]). In addition, interbands in polytene chromosomes correspond to inter-TAD regions, which contain clusters of actively transcribed genes with a broad expression pattern ([93]) or the promoters and 5' ends of such genes ([96,97]). Thus, loss of chromatin contacts within LBC loops is a common feature of highly transcribed genes in different model species and cell types.

Looping of actively-transcribed genes, which do not interact with neighboring transcriptionally inactive chromatin but also do not form interactions with active loops from other domains, resembles the local compartments reported in oocytes ([41]) and meiotic sperm cells ([44]). Similar attenuation of long-range interactions within compartments has also been observed in Anopheles mosquitoes ([14]). Lack of compartments was also observed in the Drosophila polytene chromosomes ([95]). In all four cases, the chromosomes adopt a rod-like structure, which likely explains the absence of long-range homotopic interactions in these models.

Previous studies have demonstrated that RNA polymerase can act as a "moving barrier" to extruding complexes in bacteria ([35]), yeast ([36]), slime mold *Dictyostelium discoideum* ([38]) and even for vertebrate cohesin ([37]). However, in vertebrate cells, interactions between cohesin and RNA polymerase generally have a minor impact on chromatin architecture, becoming noticeable only in conditions that extend cohesin's extrusion activity, such as Wapl+CTCF double knockout, and through extensive data aggregation across the genome ([37]). In contrast, our findings suggest that extruder-polymerase interactions play a significantly greater role in the organization of lampbrush chromosomes, explaining the majority of short-range interactions in diplotene oocytes. These results suggest that transcription-mediated repositioning of SMC complexes represents an ancient, conserved mechanism shaping chromatin architecture across the tree of life.

Methods

Oocytes isolation

Chicken oocytes, with diameters ranging from 1 to 2 mm as lampbrush chromosome and from 4 to 7 mm as post-lampbrush chromosome stages, were dissected from the ovary of healthy chicken and placed in individual drops of cooled "5:1" medium (83 mM KCl, 17 mM NaCl, 6.5 mM Na2HPO4, 3.5 mM KH2PO4, 1 mM MgCl2, 1 mM DTT, pH 7.2). All animal experiments were approved by the bioethics committee of the Institute of Cytology and Genetics SB RAS (Protocol №66, October 9th, 2020). International guidelines were followed during experimental procedures ("Guide for the Care and Use of Laboratory Animals" [98]). Nuclei were isolated as described earlier ([45]), washed with "5:1" medium, and transferred to a dish containing 2% formaldehyde in PBS for fixation at room temperature for 15 minutes. The formaldehyde was quenched by adding glycine to a final concentration of 0.125 M for 15 minutes at room temperature. The integrity of the nuclei was assessed using an Olympus stereomicroscope, and the Hi-C protocol was initiated immediately.

Hi-C libraries preparation

Individual fixed nuclei were transferred to microplate wells containing 9 µL of ice-cold lysis buffer (10 mM Tris-HCl pH 8.0, 10 mM NaCl, 0.2% Igepal) for 20 minutes. The nuclei were then washed with NEB3.1 + 0.5% SDS and transferred to 0.2 mL PCR tubes with 3 µL of NEB3.1 + 0.5% SDS. 10 µL freshly prepared 1.2% Low Melting Point Agarose were added to each tube under a stereomicroscope. After solidification, 10 µL of NEB3.1 + 0.6% SDS was added, and the samples were incubated at 37 °C for 1 hour. SDS was quenched using 10 µL of 6% Triton X-100, and chromatin fragmentation was performed with 25 U of *Dpn*II at 37 °C overnight.

The digested chromatin ends were labeled using 5 U of the Klenow fragment in the presence of biotin-15-dCTP at 22 °C for 4 hours, followed by overnight ligation at 16 °C. Crosslinks were reversed by incubation at 65 °C overnight, and the Low Melting Point Agarose was digested with 0.4 U Agarase for 1 hour at 42 °C. The purified DNA was

fragmented with *Alu*I for 1 hour at 37 °C, and next-generation sequencing (NGS) libraries
were prepared using the KAPA HyperPrep kit with modifications. The volumes for the end
repair, adapter ligation, and PCR reactions were reduced by 5-fold, 5-fold, and 2-fold,
respectively, and adapters were diluted to 300 nM. A biotin pull-down was performed with
Dynabeads MyOne Streptavidin C1 beads after the adapter ligation step, followed by 25
cycles of PCR. Genomic libraries were sequenced using BGI DNBseq G400 machine in
2x150 paired-end mode.

Hi-C data analysis and quality control

The illumina adapters from raw sequences were cut with cutadapt ver-4.1 [99]. Data were
aligned with using juicer ver-1.5.6 [100] script against chicken telomere-to-telomere
genome assembly (ASM2420605v2, [73]). The read pairs without duplicates were filtered
by mapping quality >= 30. Quality statistics were computed using juicer output with slight
modifications in cis/trans ratio estimation described previously [101]. We performed deep
sequencing only for samples with duplicates percentage below 30% (estimated in test run
with less than 200k read pairs sequenced) and fraction of intra-fragment reads
percentage 20%.

We used juicer output to generate mcool files with cooler ver-0.10.2 [102]. Due to the low
number of contacts in the dataset (~500'000 per cell), we didn't use the cooler balance
function. Instead, we set weights for bins with zero coverage as "nan" and weights of bins
with non-zero coverage as 1. Aggregated plots were made with coolpuppy ver-1.1.0 [103]
with "maxshift" of 2 MB and "local" flag. The P(s) curves were calculated using cooltools
[104] gaussian smooth sigma value equal to 0.1.

The raw reads from [42] were aligned against GRCm39 genome assembly and processed
the same way (except cooler balance was used for mcool maps). The raw reads from [15]
were aligned against ASM2420605v2 genome assembly and processed the same way
(except cooler balance was used for mcool maps).

Domain boundaries annotation

To annotate domain boundaries for both chicken oocytes, mouse oocytes and chicken
somatic cells the lavaburst ver-0.2.0 was used (https://github.com/nezar-compbio/lavaburst).
This algorithm divides the genome into domains in such a way as to maximize a global
domain scoring function. The algorithm gets the gamma as a hyperparameter so it needs
to be found. We found the value of gamma separately for each single oocyte the same
way as in the [42] article: 1) the average domain size was calculated for each gamma
(taking into account only domains more than 200 Kb); 2) the gamma value that makes
the average domain size closest to 500 Kb for mouse NSN oocytes was used. For chicken
oocytes the preferable average domain size was 1 Mb.

The domains were divided into LUDs and SSDs by N50 threshold: all domains were
sorted by their size and the domain size that covers the half of the summed domain length
was set as threshold.

Hi-C map iInsulation score track

Insulation function from cooltools package ver 0.5.1 [104] has been used for calculating the Insulation scores on merged oocytes. Resolution size is 16 Kb and window size is 2 Mb.

Aggregated Hi-C maps analysis

The aggregated Hi-C maps were generated using the coolpuppy package (https://zenodo.org/badge/latestdoi/147190130). For normalization, the coolpuppy program utilized random shifting, a built-in feature designed to adjust positional biases in the data. Insulation scores for the aggregated maps were computed with this package by calculating the relative mean signal in the first and third quadrants compared to the second and fourth quadrants. Higher insulation scores indicate higher levels of spatial separation in this context.

Saddle strength analysis

To quantify A/B compartment interaction strength, we performed a saddle plot analysis based on the first principal eigenvector of the Hi-C contact matrix derived from chicken embryonic fibroblasts. This eigenvector was computed at 100 Kb resolution using the *cooltools* package (version 0.7.1) and served as a reference compartment annotation for all subsequent analyses. Eigenvector values were divided into 40 percentile-based bins. To minimize the influence of extreme regions with ambiguous compartment identity, bins corresponding to the highest and lowest mean eigenvector values were excluded from further analysis. Saddle plots were generated by calculating the observed-over-expected contact frequencies between each pair of bins. Compartment strength was defined as the ratio of mean intra-compartment interactions (AA + BB) to inter-compartment interactions (AB + BA). All computations were performed using functions implemented in *cooltools* (ver. 0.7.1).

Chicken liver ChIP-seq data preparation

The CTCF ChIP-seq data was taken from [72] as sequenced reads. The reads were aligned at ASM2420605v2 chicken genome assembly [73] and bamCoverage ver-3.5.4 were used to get bigwig files. The ChIP signal bigwig was divided on input signal bigwig for each bin and logarithm from results were taken.

NOMe-seq experiments and data analysis

Nuclei were isolated and washed as described above. Individual nuclei were transferred immediately to 3 μL of GC Reaction Buffer (NEB) with 0.2% Igepal for 20 minutes at RT. Reaction of methylation was performed in 1x GC Reaction Buffer supplemented with 160 μM S-adenosylmethionine with 6U GpC Methyltransferase at 37 °C 1 hour. Proteines were digested by proteinase K at 65 °C for 2 hours. Libraries for whole genome bisulfite sequencing were prepared using Pico Methyl-Seq Library Prep Kit (Zymo Research) according to the user manual. For control experiments we also supplemented chicken oocytes nuclei with human K562 cells.

Genomic libraries were sequenced using the BGI DNBseq G400 machine in 2x150 paired-end mode.

The illumina adapters from raw sequences were cut with cutadapt ver-4.1 [99]. To process
NOMe-seq reads, we used Bismark version 0.23.0 ([105,106]) in conjunction with Bowtie2
version 2.4.4 ([107–110]), setting the -t parameter to 10. Reads were aligned to the
ASM2420605v2 genome assembly ([73]), and any reads with MAPQ < 10 were excluded.
Methylation data from the aligned reads were then converted to the *.bedgraph* format
using the bismark-methylation-extractor.

The GpC methylation profile was obtained from the NOMe-seq experiment described
above. The CpG methylation profile was obtained from the NOMe-seq experiment and
merged with the profile from the Methyl-seq experiment from ([67]).

RNA-seq experiments and data analysis

Nuclei were isolated and washed as described above. Individual nuclei were frozen
immediately. Total RNA was isolated from the nuclei using TRIzol reagent (ThermoFisher
Scientific) according to previously described protocol ([57]). The RNA samples were sent to
BGI company to generate transcriptomic libraries using experimental workflow including
the following steps: 1) oligoT-enrichment; 2) RNA fragmentation; 3) first-strand synthesis
with random primers; 4) second strand synthesis with dUTP instead of dTTP; 5) end-
repair, 3'-adenylation and adaptors ligation; 6) Uracil-DNA-Glycosylase treatment
followed by PCR amplification and circularization; 7) Sequencing on DNBSEQ in PE100
paired-end mode.

Together with poly(A)-enriched stranded RNA-seq data generated in this study, the total
RNA stranded RNA-seq data from [57] were used. The illumina adapters from raw
sequences were cut with cutadapt ver-4.1 [99]. All data were aligned with hisat2 ver-2.2.1
([111]) on ASM2420605v2 genome assembly ([73]). Bam files were prepared using samtools
ver-1.15.1 ([112]) and bigwig files were made using bamCoverage ver-3.5.4
(https://github.com/BGI-shenzhen/BamCoverage). Each bam file was split on subfiles by
strand (the direction of transcription) and duplicates were marked with "samtools
markdup". The unique reads were merged from all experiments (still grouped by strand)
and *de novo* LBC transcript structure recovery was made with stringtie ver-2.2.2 ([113]). The
forward and reverse transcription files were separately used with stringtie and called as
"FORW" and "REV" respectively. Maximum gap allowed between read mappings was set
as 300 bp; minimum assembled transcript length = 1000 bp; minimum junction coverage
= 5; minimum anchor length for junctions = 10 bp; minimum isoform fraction = 0.05;
minimum reads per bp coverage to consider for single-exon transcript = 3.5; minimum
reads per bp coverage to consider for multi-exon transcript = 1.2.

For the expression analysis, stranded RNA-seq data of the oocyte nuclei from this work
and [57] was compared with the RNA-seq data of the chicken embryonic fibroblasts from
[71]. All raw reads were aligned on GRCg6a genome assembly. The TPMCalculator ver-
0.0.3 was used to calculate TPM values for genes of interest for ncbiRefSeq.2020-04-01,
ensGene.v101 and refGene annotations (see Supplementary Table S4).

FPKM calculation

The FPKM value was calculated for each gene and used as a measure of transcriptional activity in analysis. $FPKM\_i = 10^9 * N\_i / (N\_0 * L\_i)$, where i denotes the transcript index, N_i - number of reads mapped to transcript i, N_0 - total number of mapped reads, L_i - length of transcript i. To estimate the amount of nascent RNA connected to the elongating RNA polymerases, the values of intron FPKM and exon FPKM were calculated separately for each transcription unit annotated. The minimum transcript FPKM value was used for the further analysis.

Mitotic metaphase chromosomes preparation

Mitotic metaphase chromosomes were prepared from chicken embryonic fibroblast (CEF) culture. Cell cycle was arrested by 0.1 µg/ml colcemid (Biological Industries) for 5 hours, then the cells were fixed in cold 3:1 ethanol:acetic acid mixture. The CEF suspension was dropped onto the slides which had been heated in a water bath to 60°C and then air dried.

Lampbrush chromosomes preparation

Lampbrush chromosome spreads were prepared according to a well-described procedure ([114]). The experimental procedure was approved by the Ethics Committee for Animal Research of St. Petersburg State University (protocol #131–04-6 dated 25 March 2019). Nuclei were manually isolated from oocytes (1-2 mm), one by one, using tungsten needles under a stereomicroscope Leica S9D or M165C (Leica Microsystems) in the isolation medium "5:1" (83 mM KCl, 17 mM NaCl, 6.5 mM Na2HPO4, 3.5 mM KH2PO4, 1 mM MgCl2, 1 mM dithiothreitol, pH 7.2). The isolated nuclei were then washed in the hypotonic solution ("5:1" medium diluted 4 times with the addition of 0.1% formaldehyde) and transferred to the isolation chamber mounted on a microscope slide and filled with the hypotonic solution. Nuclear envelope was broken manually with a pair of thin tungsten needles to release the nuclear content and slides were left to disperse for 15-20 minutes at +4°C. Slides were then centrifuged at 1600 g for 30 minutes at +4°C in a cytology centrifuge (Hettich Universal 320R), fixed in 2% formaldehyde in PBS for 30 minutes and dehydrated in a series of ethanol solutions of increasing concentrations. Specimens were mounted in an antifade solution containing 1 mg/ml DAPI (Sigma), 55% glycerol, 10 mM Tris-HCl, 2% DABCO (Sigma) and were imaged prior to FISH procedure.

Preparation of DNA-probes

To obtain DNA probes for FISH mapping of the selected scHi-C chromatin domains, 29 BAC clones were selected: 14 BAC clones in a region (55-50 Mb) on chromosome 1; 2 and 7 BAC clones in two regions (50-53 Mb and 70-73 Mb) on chromosome 4, respectively; and 6 BAC clones in the region 3-7 Mb of chromosome 13 from the chicken BAC-library CHORI-261 (https://bacpacresources.org/chicken261.htm) (Supplementary Table S3). BAC DNA was isolated from night culture by standard alkaline lysis. DNA was labeled by nick-translation ([115]) with dUTP-bio (DNA-Synthesis), dUTP-dig (Jena Bioscience) and dUTP-Atto637 (Jena Bioscience) using DNA polymerase I (ABclonal) and DNASe I (New England BioLabs). Labeled DNA probes in various combinations (up to 10 probes in a single mixture) were precipitated with salmon sperm DNA and, in case of DNA probes to the region of chromosome 1, with the C0t-5 fraction of chicken genomic

DNA, and dissolved to final concentration of 20 ng/μl for each DNA probe in the hybridization buffer containing 50% formamide, 10% dextran sulfate and 2×SSC.

Fluorescent in situ hybridization

Chromosomal position of each BAC clone-based DNA probe was first verified by DNA/DNA FISH on CEF metaphase chromosomes: preparations were pretreated with RNAse A (Thermo Scientific) and pepsin (Sigma-Aldrich). For lampbrush chromosomes, the DNA/DNA+RNA FISH procedure was used, omitting RNAse treatment ([56,116]). Chromosome preparations were denatured in 70% formamide in 2×SSC at 70°C for 10 minutes, and dehydrated in a series of cold (-20°C) ethanol solutions of increasing concentration. Hybridization mixtures were denatured at 90°C and cooled on ice and applied to the slides with the denatured chromosomes. Hybridization was carried out overnight in a humidity chamber at 37°C, followed by a post-hybridization wash in 0.2×SSC at 60°C. Biotin was detected with Alexa488-conjugated streptavidin (Jackson ImmunoResearch Laboratories) and biotinylated anti-streptavidin antibody (Vectorlabs), and digoxigenin was detected with Cy3-conjugated mouse anti-digoxigenin antibody (Invitrogen) and Cy3-conjugated anti-mouse IgG (Jackson ImmunoResearch Laboratories).

Immunostaining

Lampbrush chromosome preparations were immunostained with mouse monoclonal antibodies against phosphorylated serine 5 of CTD repeat YSPTSPS of RNA polymerase II [H14] (Abcam, ab24759), followed by the detection with Alexa-488-conjugated goat anti-mouse IgG (Molecular Probes). Immunofluorescence staining procedure was described previously ([69,117]).

Microscopy and image processing

Chromosome preparations were analyzed with the fluorescence microscope Leica DM4000 (Leica Microsystems, GmbH) equipped with a monochrome CCD camera (1,3 Mp) and corresponding fluorescence filter cubes. LAS AF software (Leica Microsystems, GmbH) was used for image acquisition, and Adobe Photoshop CS2 (Adobe) and ImageJ (http://rsbweb.nih.gov/ij/) was used for image processing and layout. Schematic drawings were created in CorelDRAW X5 and represent an approximation of the FISH mapping of DNA probes to lampbrush chromatin domains seen in at least 4 micrographs for each DNA probe.

Polymer Modelling

We used polymer modeling based on the Polychrom tool [[118]] to gain insights into the roles of individual RNA polymerases in the folding of lampbrush chromosomes. The simulation was performed in the coarse-grained model approximation, where clusters of nucleosomes and individual polymerases acted as beads.

<u>1D</u>

The modeled object is a one-dimensional array of N beads of sizes ranging from 45 to 225 bp, with dynamically loaded-unloaded RNA polymerases and active extruders (loop extrusion factors, LEFs) and non-removable cohesins without the loop extruding activity (loaded with the density of 45 Kb).

The number of base pairs in a bead differs for TUs where RNA polymerases are located and for transcriptionally inactive regions with a dense fit of nucleosomes. The size of 45 bp per bead was used to represent one RNA polymerase (~15 nm, [119]). The regions between the TUs are condensed by the nucleosomes ~5 times (because ~140 bp entangle it and ~70 bp are between nucleosomes). Our estimation is also close to the estimation from [80]. Transcription units are placed according to the annotation obtained from RNA-seq data.

RNA polymerases are represented as objects with the following properties: they are loaded at the beginning of TUs with a probability proportional to the activity of the TU (estimated from FPKM metric), from where they begin movement at a speed of vp = 100 bp/sec [91] (the speed of LEFs is vL = 625 bp/sec for the naked DNA [92]) along the transcript. Nascent RNA fiber growth from each RNA polymerase at length *vp*dt* for the time step of *dt* seconds. The introns are excised from the pre-mRNA attached to the RNA polymerases. If the bead in the direction of movement is occupied by a LEF or cohesive cohesin (i.e. SMC), then RNA polymerase pushes the SMC [37] and all subsequent SMCs in the direction of movement until they collide with another RNA polymerase.

The mechanism of LEF action is as follows. It is randomly loaded onto a pair of adjacent beads that are not occupied by another LEF or RNA polymerase. Each time one of the two LEF retaining components can reattach to the next bead in the corresponding direction releasing the occupied one. The direction of movement changes every timestep (effectively two-way movement). Each part of the LEF can move independently of the other. If the next bead is occupied by another LEF then no movement occurs. If the next bead is occupied by RNA polymerase then the LEF can bypass the RNA polymerase with the stable probability *Pbypass*. The cohesive cohesin can be pushed by LEF if the next bead is not occupied. Extruding continues until the LEF is removed from the chromatin. The probability of removal is (*1 / lifetime)* for each timestep.

Some parameters for these LEFs were determined by using P(s) data from the experiment Hi-C data on lampbrush-stage oocyte nuclei (see "Supplementary methods 1. Modeling"): *lifetime* = 1 Mb (~60% of LEFs will pass this distance on naked DNA), *average_loop_size* = 90 Kb (determines the density of loaded condensins on chromatin), *Pbypass* = 0.01 (as in [37] article).

3D

The next step after forming 1D snapshots of the SMCs and RNA polymerase locations on the chromatin is to transfer the interactions into 3D. For this purpose all beads are grouped into bead clusters of 10 which include nucleosomes and transcription units. The sizes of the clusters vary from 450 to 2250 bp. New beads are placed into the starting conformation form of two parallel 2D fractals, corresponding to the two sister chromatids.

The distances between beads in 3D varied according to the fractal dimension of the site ("Supplementary methods 1. Modeling").

With the launch of a three-dimensional simulation, at each moment in time, a spring of a fixed size (~40 nm [120]) is stretched between each pair of beads connected by a given LEF, and the beads move according to a random Brownian motion.

RNA tails from 1D simulation are converted into increased bead sizes by the Marco and Siggia model [76]. The repulsive potential between beads is realized as *0.5*k*(r1+r2)^2*. A more detailed analysis of the model and parameters used is presented in the "Supplementary methods 1. Modeling".

In order to reproduce the patterns observed in LBC under microscopy, the original three-dimensional regions of the chromosome were flattened to 1 μm (see "Supplementary methods 1. Modeling").

LBC chromatin simulations were additionally validated using obtained FISH-mapping data in individual chromosomal regions (see "Supplementary Note 3").

Online supplementary files can be found via links (3D structure and Flattened structure) or be downloaded from git (https://github.com/genomech/LBC_model).

Simulated Hi-C

The resulting 3D polymers were used to construct contact maps similar to Hi-C maps. The beads of different sizes are correctly converted to Kb resolution that allow us to compare simulated and experimental contact maps of the specific chromatin slices. Since the contact map collected from one polymer conformation is essentially one conformation of an oocyte, the sum of contacts from one "*Dpn*II site" in a bead was normalized to 8 as in the experiment (see "Supplementary methods 1. Modeling" for the details). Simulated Hi-C maps were then stored in mcool format and the analyses that are the same with experiment Hi-C maps were performed.

References

1. Kabirova, E. *et al.* Function and evolution of the loop extrusion machinery in animals. *IJMS* **24**, 5017 (2023).

2. Misteli, T. The self-organizing genome: principles of genome architecture and function. *Cell* **183**, 28–45 (2020).

3. 4D Nucleome Consortium *et al.* An integrated view of the structure and function of the human 4D nucleome. Preprint at https://doi.org/10.1101/2024.09.17.613111 (2024).

4. Kabirova, E. *et al.* TAD border deletion at the Kit locus causes tissue-specific ectopic activation of a neighboring gene. *Nat Commun* **15**, 4521 (2024).

5. Salnikov, P. *et al.* Structural variants in the Epb41l4a locus: TAD disruption and Nrep gene misregulation as hypothetical drivers of neurodevelopmental outcomes. *Sci Rep* **14**, 5288 (2024).

6.   Rajderkar, S. *et al.* Topologically associating domain boundaries are required for normal genome function. *Commun Biol* **6**, 435 (2023).

7.   Lupiáñez, D. G. *et al.* Disruptions of Topological Chromatin Domains Cause Pathogenic Rewiring of Gene-Enhancer Interactions. *Cell* **161**, 1012–1025 (2015).

8.   Pope, B. D. *et al.* Topologically associating domains are stable units of replication-timing regulation. *Nature* **515**, 402–405 (2014).

9.   Klein, K. N. *et al.* Replication timing maintains the global epigenetic state in human cells. *Science* **372**, 371–378 (2021).

10.   Moindrot, B. *et al.* 3D chromatin conformation correlates with replication timing and is conserved in resting cells. *Nucleic Acids Research* **40**, 9470–9481 (2012).

11.   Piazza, A. *et al.* Cohesin regulates homology search during recombinational DNA repair. *Nat Cell Biol* **23**, 1176–1186 (2021).

12.   Strom, A. R. *et al.* HP1α is a chromatin crosslinker that controls nuclear and mitotic chromosome mechanics. *eLife* **10**, e63972 (2021).

13.   Uhler, C. & Shivashankar, G. V. Regulation of genome organization and gene expression by nuclear mechanotransduction. *Nat Rev Mol Cell Biol* **18**, 717–727 (2017).

14.   Lukyanchikova, V. *et al.* Anopheles mosquitoes reveal new principles of 3D genome organization in insects. *Nat Commun* **13**, 1960 (2022).

15.   Fishman, V. *et al.* 3D organization of chicken genome demonstrates evolutionary conservation of topologically associated domains and highlights unique architecture of erythrocytes' chromatin. *Nucleic Acids Research* **47**, 648–665 (2019).

16.   Krefting, J., Andrade-Navarro, M. A. & Ibn-Salem, J. Evolutionary stability of topologically associating domains is associated with conserved gene regulation. *BMC Biol* **16**, 87 (2018).

17.   Hoencamp, C. *et al.* 3D genomics across the tree of life reveals condensin II as a determinant of architecture type. *Science* **372**, 984–989 (2021).

18.   Polovnikov, K. E. *et al.* Crumpled Polymer with Loops Recapitulates Key Features of Chromosome Organization. *Phys. Rev. X* **13**, 041029 (2023).

19.   Mirny, L. A. The fractal globule as a model of chromatin architecture in the cell. *Chromosome Res* **19**, 37–51 (2011).

20.   Razin, S. V. & Gavrilov, A. A. The role of liquid–liquid phase separation in the compartmentalization of cell nucleus and spatial genome organization. *Biochemistry Moscow* **85**, 643–650 (2020).

21.   Rada-Iglesias, A., Grosveld, F. G. & Papantonis, A. Forces driving the three-dimensional folding of

1060   eukaryotic genomes. *Molecular Systems Biology* **14**, e8214 (2018).

1061   22.   Strom, A. R. & Brangwynne, C. P. The liquid nucleome – phase transitions in the nucleus at a glance. *Journal

1062   of Cell Science* **132**, jcs235093 (2019).

1063   23.   Cremer, T. *et al.* The Interchromatin Compartment Participates in the Structural and Functional Organization

1064   of the Cell Nucleus. *BioEssays* **42**, 1900132 (2020).

1065   24.   Li, H., Playter, C., Das, P. & McCord, R. P. Chromosome compartmentalization: causes, changes,

1066   consequences, and conundrums. *Trends in Cell Biology* **34**, 707–727 (2024).

1067   25.   Padeken, J. & Heun, P. Nucleolus and nuclear periphery: Velcro for heterochromatin. *Current Opinion in Cell

1068   Biology* **28**, 54–60 (2014).

1069   26.   Quinodoz, S. A. *et al.* Higher-Order Inter-chromosomal Hubs Shape 3D Genome Organization in the Nucleus.

1070   *Cell* **174**, 744-757.e24 (2018).

1071   27.   Solovei, I., Thanisch, K. & Feodorova, Y. How to rule the nucleus: divide et impera. *Current Opinion in Cell

1072   Biology* **40**, 47–59 (2016).

1073   28.   Dekker, J. & Mirny, L. A. The chromosome folding problem and how cells solve it. *Cell* **187**, 6424–6450

1074   (2024).

1075   29.   Davidson, I. F. & Peters, J.-M. Genome folding through loop extrusion by SMC complexes. *Nat Rev Mol Cell

1076   Biol* **22**, 445–464 (2021).

1077   30.   Szabo, Q., Bantignies, F. & Cavalli, G. Principles of genome folding into topologically associating domains.

1078   *Sci. Adv.* **5**, eaaw1668 (2019).

1079   31.   Wang, J. *et al.* Manipulation of TAD reorganization by chemical-dependent genome linking. *STAR Protocols

1080   2**, 100799 (2021).

1081   32.   Dixon, J. R. *et al.* Topological domains in mammalian genomes identified by analysis of chromatin

1082   interactions. *Nature* **485**, 376–380 (2012).

1083   33.   Nora, E. P. *et al.* Spatial partitioning of the regulatory landscape of the X-inactivation centre. *Nature* **485**,

1084   381–385 (2012).

1085   34.   Dequeker, B. J. H. *et al.* MCM complexes are barriers that restrict cohesin-mediated loop extrusion. *Nature

1086   606**, 197–203 (2022).

1087   35.   Brandão, H. B. *et al.* RNA polymerases as moving barriers to condensin loop extrusion. *Proc. Natl. Acad. Sci.

1088   U.S.A.* **116**, 20489–20499 (2019).

1089   36.   Rivosecchi, J. *et al.* RNA polymerase backtracking results in the accumulation of fission yeast condensin at

1090   active genes. *Life Sci. Alliance* **4**, e202101046 (2021).

37.   Banigan, E. J. *et al.* Transcription shapes 3D chromatin organization by interacting with loop extrusion. *Proc.*
Natl. Acad. Sci. U.S.A. **120**, e2210480120 (2023).

38.   Zhegalova, I. V. *et al.* Convergent gene pairs restrict chromatin looping in Dictyostelium discoideum, acting
as directional barriers for extrusion. Preprint at https://doi.org/10.1101/2024.06.12.598618 (2024).

39.   Belokopytova, P. & Fishman, V. Predicting Genome Architecture: Challenges and Solutions. *Front. Genet.*
11, 617202 (2021).

40.   Rowley, M. J. & Corces, V. G. Organizational principles of 3D genome architecture. *Nat Rev Genet* **19**, 789–
800 (2018).

41.   Du, Z. *et al.* Polycomb group proteins regulate chromatin architecture in mouse oocytes and early embryos.
Molecular Cell **77**, 825-839.e7 (2020).

42.   Flyamer, I. M. *et al.* Single-nucleus Hi-C reveals unique chromatin reorganization at oocyte-to-zygote
transition. *Nature* **544**, 110–114 (2017).

43.   Wang, Y. *et al.* Reprogramming of meiotic chromatin architecture during spermatogenesis. *Molecular Cell*
73, 547-561.e6 (2019).

44.   Cheng, G. *et al.* High resolution maps of chromatin reorganization through mouse meiosis reveal novel
features of the 3D meiotic structure. Preprint at https://doi.org/10.1101/2024.03.25.586627 (2024).

45.   Krasikova, A., Khodyuchenko, T., Maslova, A. & Vasilevskaya, E. Three-dimensional organisation of RNA-
processing machinery in avian growing oocyte nucleus. *Chromosome Res* **20**, 979–994 (2012).

46.   Maslova, A. V. & Krasikova, A. V. Spatial arrangement of macro-, midi-, and microchromosomes in
transcriptionally active nuclei of growing oocytes in birds of the order galliformes. *Cell Tiss. Biol.* **5**, 281–293
(2011).

47.   Vlad, M. & Macgregor, H. C. Chromomere number and its genetic significance in lampbrush chromosomes.
Chromosoma **50**, 327–347 (1975).

48.   Macgregor, H. C. Chromomeres revisited. *Chromosome Res* **20**, 911–924 (2012).

49.   Krasikova, A., Fishman, V. & Kulikova, T. Lampbrush chromosome studies in the post-genomic era.
BioEssays **45**, 2200250 (2023).

50.   Gall, J. G. Transcription in the *Xenopus* Oocyte Nucleus. in *Xenopus Development* (eds. Kloc, M. & Kubiak,
J. Z.) 3–15 (Wiley, 2014). doi:10.1002/9781118492833.ch1.

51.   Morgan, G. T. Imaging the dynamics of transcription loops in living chromosomes. *Chromosoma* **127**, 361–
374 (2018).

52.   Kaufmann, R., Cremer, C. & Gall, J. G. Superresolution imaging of transcription units on newt lampbrush
chromosomes. *Chromosome Res* **20**, 1009–1015 (2012).

53.   Mirny, L. A. & Solovei, I. Keeping chromatin in the loop(s). *Nat Rev Mol Cell Biol* **22**, 439–440 (2021).

54.   Keinath, M. C., Davidian, A., Timoshevskiy, V., Timoshevskaya, N. & Gall, J. G. Characterization of axolotl

1125   lampbrush chromosomes by fluorescence in situ hybridization and immunostaining. *Experimental Cell*

1126   *Research* **401**, 112523 (2021).

1127   55.   Kulikova, T., Maslova, A., Starshova, P., Rodriguez Ramos, J. S. & Krasikova, A. Comparison of the somatic

1128   TADs and lampbrush chromomere-loop complexes in transcriptionally active prophase I oocytes.

1129   *Chromosoma* **131**, 207–223 (2022).

1130   56.   Kulikova, T. & Krasikova, A. RNA-FISH—on Lampbrush Chromosomes: Visualization of Individual

1131   Transcription Units. in *Cytogenetics and Molecular Cytogenetics* 307–317 (CRC Press, Boca Raton, 2022).

1132   doi:10.1201/9781003223658-26.

1133   57.   Krasikova, A. *et al.* The first chicken oocyte nucleus whole transcriptomic profile defines the spectrum of

1134   maternal mRNA and non-coding RNA genes transcribed by the lampbrush chromosomes. *Nucleic Acids*

1135   *Research* **52**, 12850–12877 (2024).

1136   58.   Zlotina, A. *et al.* New insights into chromomere organization provided by lampbrush chromosome

1137   microdissection and high-throughput sequencing. *Front. Genet.* **11**, 57 (2020).

1138   59.   Zlotina, A., Kulikova, T., Kosyakova, N., Liehr, T. & Krasikova, A. Microdissection of lampbrush

1139   chromosomes as an approach for generation of locus-specific FISH-probes and samples for high-throughput

1140   sequencing. *BMC Genomics* **17**, 126 (2016).

1141   60.   Kulikova, T., Surkova, A., Zlotina, A. & Krasikova, A. Mapping epigenetic modifications on chicken

1142   lampbrush chromosomes. *Mol Cytogenet* **13**, 32 (2020).

1143   61.   Krasikova, A., Kulikova, T., Rodriguez Ramos, J. S. & Maslova, A. Assignment of the somatic A/B

1144   compartments to chromatin domains in giant transcriptionally active lampbrush chromosomes. *Epigenetics &*

1145   *Chromatin* **16**, 24 (2023).

1146   62.   Krasikova, A. V. & Kulikova, T. V. Distribution of heterochromatin markers in lampbrush chromosomes in

1147   birds. *Russ J Genet* **53**, 1022–1029 (2017).

1148   63.   Gridina, M. *et al.* Comparison and critical assessment of single-cell Hi-C protocols. *Heliyon* **8**, e11023 (2022).

1149   64.   Galitsyna, A. A. & Gelfand, M. S. Single-cell Hi-C data analysis: safety in numbers. *Briefings in*

1150   *Bioinformatics* **22**, bbab316 (2021).

1151   65.   Akgol Oksuz, B. *et al.* Systematic evaluation of chromosome conformation capture assays. *Nat Methods* **18**,

1152   1046–1055 (2021).

1153   66.   Rao, S. S. P. *et al.* A 3D Map of the Human Genome at Kilobase Resolution Reveals Principles of Chromatin

1154   Looping. *Cell* **159**, 1665–1680 (2014).

67.   Nurislamov, A., Lagunov, T., Gridina, M., Krasikova, A. & Fishman, V. Single-Cell DNA Methylation

Analysis of Chicken Lampbrush Chromosomes. *IJMS* **23**, 12601 (2022).

68.   Zlotina, A. *et al.* Centromere positions in chicken and Japanese quail chromosomes: de novo centromere

formation versus pericentric inversions. *Chromosome Res* **20**, 1017–1032 (2012).

69.   Krasikova, A., Barbero, J. L. & Gaginskaya, E. Cohesion proteins are present in centromere protein bodies

associated with avian lampbrush chromosomes. *Chromosome Res* **13**, 675–685 (2005).

70.   Beenders, B., Watrin, E., Legagneux, V., Kireev, I. & Bellini, M. Distribution of XCAP-E and XCAP-D2 in

the Xenopus oocyte nucleus. *Chromosome Research* **11**, 549–564 (2003).

71.   Li, D. *et al.* Comparative 3D genome architecture in vertebrates. *BMC Biol* **20**, 99 (2022).

72.   Jerković, I. *et al.* Genome-wide binding of posterior HOXA/D transcription factors reveals subgrouping and

association with CTCF. *PLoS Genet* **13**, e1006567 (2017).

73.   Huang, Z. *et al.* Evolutionary analysis of a complete chicken genome. *Proc. Natl. Acad. Sci. U.S.A.* **120**,

e2216641120 (2023).

74.   Degalez, F. *et al.* Enriched atlas of lncRNA and protein-coding genes for the GRCg7b chicken assembly and

its functional annotation across 47 tissues. *Sci Rep* **14**, 6588 (2024).

75.   Brandão, H. B., Ren, Z., Karaboja, X., Mirny, L. A. & Wang, X. DNA-loop-extruding SMC complexes can

traverse one another in vivo. *Nat Struct Mol Biol* **28**, 642–651 (2021).

76.   Marko, J. F. & Siggia, E. D. Polymer Models of Meiotic and Mitotic Chromosomes. *MBoC* **8**, 2217–2231

(1997).

77.   Maslova, A. & Krasikova, A. Nuclear actin depolymerization in transcriptionally active avian and amphibian

oocytes leads to collapse of intranuclear structures. *Nucleus* **3**, 300–311 (2012).

78.   Flemming, W. *Zellsubstanz, Kern Und Zelltheilung*. (Vogel, 1882).

79.   Rückert, J. *Zur Entwickelungsgeschichte Des Ovarialeies Bei Selachiern*. (1892).

80.   Leidescher, S. *et al.* Spatial organization of transcribed eukaryotic genes. *Nat Cell Biol* **24**, 327–339 (2022).

81.   Morgan, G. T., Jones, P. & Bellini, M. Association of modified cytosines and the methylated DNA-binding

protein MeCP2 with distinctive structural domains of lampbrush chromatin. *Chromosome Res* **20**, 925–942

(2012).

82.   Franke, W. W., Scheer, U., Trendelenburg, M. F., Spring, H. & Zentgraf, H. Absence of nucleosomes in

transcriptionally active chromatin. (1976).

83.   Ocampo-Hafalla, M., Muñoz, S., Samora, C. P. & Uhlmann, F. Evidence for cohesin sliding along budding

yeast chromosomes. *Open Biol.* **6**, 150178 (2016).

84.   Davidson, I. F. *et al.* Rapid movement and transcriptional re-localization of human cohesin on DNA. *The*

1187   *EMBO Journal* **35**, 2671–2685 (2016).

1188   85.   Guérin, T. M., Barrington, C., Pobegalov, G., Molodtsov, M. I. & Uhlmann, F. An extrinsic motor directs

1189   chromatin loop formation by cohesin. *EMBO J* **43**, 4173–4196 (2024).

1190   86.   Bastié, N. *et al.* Sister chromatid cohesion halts DNA loop expansion. *Molecular Cell* **84**, 1139-1148.e5

1191   (2024).

1192   87.   Gaginskaya, E., Kulikova, T. & Krasikova, A. Avian lampbrush chromosomes: a powerful tool for

1193   exploration of genome expression. *Cytogenet Genome Res* **124**, 251–267 (2009).

1194   88.   Krasikova, A. & Kulikova, T. *Hromosomy Tipa Lampovyh Shchetok: Sovremennye Predstavleniya i*

1195   *Perspektivy Issledovanij [Lampbrush Chromosomes: Current Insights and Research Perspectives].* (St

1196   Petersburg: St Petersburg University Publishing House, 2019).

1197   89.   Krasikova, A. Centromernye rajony hromosom i associirovannye s nimi struktury v yadrah rastushchih

1198   oocitov ptic [Centromeric regions of chromosomes and associated structures in the nuclei of growing avian

1199   oocytes]. Ph. D. Thesis. (Saint Petersburg State University, Saint Petersburg, 2007).

1200   90.   Samejima, K. *et al.* Rules of engagement for condensins and cohesins guide mitotic chromosome formation.

1201   Preprint at https://doi.org/10.1101/2024.04.18.590027 (2024).

1202   91.   Wang, M. D. *et al.* Force and velocity measured for single molecules of RNA polymerase. *Science* **282**, 902–

1203   907 (1998).

1204   92.   Kong, M. *et al.* Human condensin I and II drive extensive ATP-dependent compaction of nucleosome-bound

1205   DNA. *Molecular Cell* **79**, 99-114.e9 (2020).

1206   93.   Ulianov, S. V. *et al.* Active chromatin and transcription play a key role in chromosome partitioning into

1207   topologically associating domains. *Genome Res.* **26**, 70–84 (2016).

1208   94.   Winick-Ng, W. *et al.* Cell-type specialization is encoded by specific chromatin topologies. *Nature* **599**, 684–

1209   691 (2021).

1210   95.   Eagen, K. P., Hartl, T. A. & Kornberg, R. D. Stable chromosome condensation revealed by chromosome

1211   conformation capture. *Cell* **163**, 934–946 (2015).

1212   96.   Stadler, M. R., Haines, J. E. & Eisen, M. B. Convergence of topological domain boundaries, insulators, and

1213   polytene interbands revealed by high-resolution mapping of chromatin contacts in the early Drosophila

1214   melanogaster embryo. *eLife* **6**, e29550 (2017).

1215   97.   Kolesnikova, T. D. Banding pattern of polytene chromosomes as a representation of universal principles of

1216   chromatin organization into topological domains. *Biochemistry Moscow* **83**, 338–349 (2018).

98.   *Guide for the Care and Use of Laboratory Animals: Eighth Edition*. 12910 (National Academies Press,

Washington, D.C., 2011). doi:10.17226/12910.

99.   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet j.* **17**, 10

(2011).

100. Durand, N. C. *et al.* Juicer Provides a One-Click System for Analyzing Loop-Resolution Hi-C Experiments.

Cell Systems **3**, 95–98 (2016).

101. Gridina, M. *et al.* A cookbook for DNase Hi-C. *Epigenetics & Chromatin* **14**, 15 (2021).

102. Abdennur, N. & Mirny, L. A. Cooler: scalable storage for Hi-C data and other genomically labeled arrays.

Bioinformatics **36**, 311–316 (2020).

103. Flyamer, I. M., Illingworth, R. S. & Bickmore, W. A. *Coolpup.py:* versatile pile-up analysis of Hi-C data.

Bioinformatics **36**, 2980–2985 (2020).

104. Open2C *et al.* Cooltools: enabling high-resolution Hi-C analysis in Python. Preprint at

https://doi.org/10.1101/2022.10.31.514564 (2022).

105. Krueger, F., Kreck, B., Franke, A. & Andrews, S. R. DNA methylome analysis using short bisulfite

sequencing data. *Nat Methods* **9**, 145–151 (2012).

106. Krueger, F. & Andrews, S. R. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq

applications. *Bioinformatics* **27**, 1571–1572 (2011).

107. Ferragina, P. & Manzini, G. Opportunistic data structures with applications. in *Proceedings 41st Annual

Symposium on Foundations of Computer Science* 390–398 (IEEE Comput. Soc, Redondo Beach, CA, USA,

2000). doi:10.1109/SFCS.2000.892127.

108. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**, 357–359 (2012).

109. Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient alignment of short

DNA sequences to the human genome. *Genome Biol* **10**, R25 (2009).

110. Langmead, B., Wilks, C., Antonescu, V. & Charles, R. Scaling read aligners to hundreds of threads on

general-purpose processors. *Bioinformatics* **35**, 421–432 (2019).

111. Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome alignment and genotyping

with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**, 907–915 (2019).

112. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *GigaScience* **10**, giab008 (2021).

113. Shumate, A., Wong, B., Pertea, G. & Pertea, M. Improved transcriptome assembly using a hybrid of long and

short reads with StringTie. *PLoS Comput Biol* **18**, e1009730 (2022).

114. Solovei, I., Gaginskaya, E., Allen, T. & Macgregor, H. A novel structure associated with a lampbrush

chromosome in the chicken, *Gallus domesticus*. *Journal of Cell Science* **101**, 759–772 (1992).

115. Cremer, M. *et al.* Multicolor 3D Fluorescence In Situ Hybridization for Imaging Interphase Chromosomes. in

The Nucleus (ed. Hancock, R.) vol. 463 205–239 (Humana Press, Totowa, NJ, 2012).

116. Zlotina, A. & Krasikova, A. FISH in Lampbrush Chromosomes. in *Fluorescence In Situ Hybridization (FISH)*

(ed. Liehr, T.) 445–457 (Springer Berlin Heidelberg, Berlin, Heidelberg, 2017). doi:10.1007/978-3-662-

52959-1_45.

117. Krasikova, A., Kulikova, T., Saifitdinova, A., Derjusheva, S. & Gaginskaya, E. Centromeric protein bodies on

avian lampbrush chromosomes contain a protein detectable with an antibody against DNA topoisomerase II.

Chromosoma **113**, 316–323 (2004).

118. Imakaev, M., Goloborodko, A., & Hbbrandao. mirnylab/polychrom: v0.1.0. Zenodo

https://doi.org/10.5281/ZENODO.3579473 (2019).

119. Pradhan, B. *et al.* SMC complexes can traverse physical roadblocks bigger than their ring size. *Cell Reports*

41, 111491 (2022).

120. Ryu, J.-K. *et al.* Condensin extrudes DNA loops in steps up to hundreds of base pairs that are generated by

ATP binding events. *Nucleic Acids Research* **50**, 820–832 (2022).

1263

Acknowledgments

This manuscript is dedicated to the memory of Joseph G. Gall. This study was initiated
with support of RSF Project №20-64-46021 in 2020-2022. We acknowledge the collective
centers of Ministry of Science and Higher Education of the Russian Federation (state
project FWNR-2022-0019) for providing facilities for LBC isolation and genomic libraries
preparation and sequencing. Access to computational resources was provided by
Novosibirsk State University, supported by the Ministry of Education and Science of the
Russian Federation (FSUS-2024-0018). Microscopy analysis was performed using the
equipment of the Resource Center 'Molecular and Cell Technologies' (Saint-Petersburg
State University). Part of the bioinformatic analysis and manuscript preparation was
supported by grant of the state program of the "Sirius" Federal Territory "Scientific and
technological development of the "Sirius" Federal Territory" (agreement №26-03,
27/09/2024). The original manuscript text was composed by authors; proofreading was
conducted with the assistance of ChatGPT version 4, after that the text was corrected
and edited by the authors.

1279

Author contributions

1281   V.F. and A.K. conceptualized the study. M.G. conducted the genomics experiments with
1282   assistance from A.K., T.K., and A.M., who contributed to oocyte isolation, and A.N., who
1283   provided support for staining and NOMe-seq experiments. Cytological analyses, including
1284   FISH-mapping experiments, were performed by T.K. and A.K. with contribution from A.M.,
1285   who performed probe labeling. Bioinformatic analyses were carried out by T.L., with

contributions from V.F., A.P., A.N., and V.K. T.L. developed the methodology for polymer simulations and conducted the modeling experiments with input from V.K. T.L. and V.F. drafted the manuscript with contributions by A.K.; authors jointly developed mechanism explaining lampbrush chromosome organization, and A.K. suggested the final scheme of chromomere-loop complex formation. All authors reviewed, revised, and approved the final manuscript.

Data access

All raw sequencing data and proceed data are available under GEO accession GSE288415.

The program code used for lampbrush chromosome model simulation is freely and publicly available at https://github.com/genomech/LBC_model.