Climate and seed weight drive intraspecific variation in seed longevity in storage

Lea Klepka[1][*], Sascha Liepelt[1] & Anna Bucharova[1]

[1] Conservation Biology, Department of Biology, Philipps University Marburg, Karl-von-Frisch-Straße 8, 35043 Marburg, Germany

[*] Corresponding author: Lea Klepka, lea.klepka@biologie.uni-marburg.de

Running head

Intraspecific variation in seed longevity in storage

ABSTRACT

Premise: Conservation seed banks are essential for ex-situ plant conservation, but

stored seeds slowly deteriorate and lose viability. Seed longevity in storage is

determined by the initial seed viability and the rate of seed viability loss. The rate of

seed viability loss in storage varies between species, and there is possibly also some

variation between populations or even genotypes within species. However, the extent

of this intraspecific variability and its drivers remain unclear.

Methods: We investigated both inter- and intraspecific variability in seed longevity

and its predictors in 41 common grassland species and 188 seed accessions from

across Europe. We exposed the seeds to artificial ageing conditions (60% RH, 45°C)

and used probit analysis to obtain the rate of seed viability loss (σ) as a measure of

seed longevity. We then related σ to both accession- and species-specific factors.

Key results: Seed longevity (σ) varied significantly among accessions within 58% of

the species, and the probability of detecting such intraspecific differences increased

with the number of accessions available for a given species. This suggests that

within-species variation in seed longevity is widespread. Accession-specific

predictors explained only 14.4% of the within-species variability. Specifically, seed

longevity increased with the mean annual temperature at the accession origin and

decreased with the accession-specific seed weight. Across species, seed longevity

differed among plant families but was unrelated to seed weight or seed chemical

composition.

Conclusions. Our findings highlight substantial within-species variation in seed

longevity in storage, which, however, is difficult to predict.

KEYWORDS

INTRODUCTION

Biodiversity conservation is one of the critical global challenges, and *ex situ* methods play an increasingly important role in this task (Cohen et al., 1991). One of the most common ex-situ methods for plant conservation is seed banks, dedicated facilities that store seeds of wild plant species, particularly the vulnerable ones (Peres, 2016). The stored seeds represent a snapshot of past populations and are valuable resources for species reintroductions or ecosystem restoration, and also for scientific research (Cochrane et al., 2007; Etterson et al., 2016; Wambugu et al., 2023). Thus, the ultimate task of conservation seed banks is to keep the seeds viable for decades or even centuries.

Seed survival in storage depends on the type of seeds and storage conditions. Some types of seeds, so-called recalcitrant seeds, do not survive desiccation, making long-term storage impossible. Other types of seeds, called orthodox, can survive desiccation, and they can be stored for a couple of months up to centuries, depending on the species (Hay & Probert, 2013). Conservation seed banks aim to maintain seed viability as long as possible by storing the seeds under low temperatures and humidity to slow down internal chemical processes such as lipid peroxidation and DNA damage (Nagel et al., 2015; Walters et al., 2005a). The seed longevity also depends on the quality and viability of the seeds (Hay & Probert, 2013; Niedzielski et al., 2009), and thus, guidelines for seed banking recommend storing accessions with high initial viability (ENSCONET, 2009; Millennium Seed Bank, 2022). Yet, even highly viable seeds stored under optimal conditions slowly deteriorate and eventually die. This mortality is faster when the seeds are stored in suboptimal conditions, for example when storing seeds for large-scale ecosystem restoration in ambient conditions at a farm (Merritt & Dixon, 2011). To optimise the long-term effectiveness of seed storage – be it for conservation, research or ecosystem restoration purposes – we need to understand which factors predict seed longevity in storage.

Studying seed longevity presents unique challenges, as it requires comparing old and fresh seeds of the same seed lot, which would need time travel. To circumvent this obstacle, researchers have developed artificial ageing protocols which simulate natural ageing by exposing the seeds to high temperatures and high relative humidity

(Delouche & Baskin, 2021). These conditions accelerate the same or at least similar

chemical processes typical of natural ageing during long-term storage (Delouche &

Baskin, 2021). Although this method may not perfectly replicate natural ageing, the

longevity of seeds in artificial ageing correlates with the longevity of seeds stored in

seed banks both across and within species (Ninoles et al., 2022; Probert et al.,

2009). Nevertheless, artificial ageing is the best method we currently have to

compare seed longevity among accessions.

Seed survival in artificial ageing differs between species, and to some degree also

between higher taxa like plant families, although large plant families harbour

substantial variation in seed longevity (Probert et al., 2009; Walters et al., 2005a).

Studies aiming to detect general patterns in seed longevity beyond evolutionary

history yielded mixed results. Endospermic seeds survive shorter in storage than

non-endospermic ones (Probert et al., 2009). The influence of seed weight is less

clear; some studies suggest that seeds of small-seeded species survive longer

(Satyanti et al., 2018), while others report no effect of seed weight (Davies et al.,

2020; Merritt et al., 2014; Probert et al., 2009). Seed longevity is also affected by

climatic conditions at seed collection sites: Seeds from hot or dry environments

generally survive longer than those from cooler or wetter conditions (Merritt et al.,

2014; Probert et al., 2009; Zani & Müller, 2017). Furthermore, seed longevity should

hypothetically depend on the seed's chemical composition because high oil content

should reduce seed longevity due to ongoing lipid peroxidation in storage (Narayana

Murthy & Sun, 2000; Pritchard & Dickie, 2003), yet the results of empirical studies are

inconsistent (Nagel et al., 2015; Probert et al., 2009; Walters et al., 2005a).

In contrast to between-species comparison, there is less comprehensive information

on within-species variability on seed longevity in storage. Most data come from

cultivated crops or model species (e.g. Bizouerne et al., 2023; Franks et al., 2018;

Guzzon et al., 2021; Lee et al., 2019), with only a few studies addressing natural

populations of wild species (Genna et al., 2020; Kochanek et al., 2009; Mondoni et

al., 2011; White et al., 2023). These indicate that the variation in seed longevity within

a species can be as substantial as the variation between species from the same

ecosystem (Kochanek et al., 2009; Mondoni et al., 2011). The predictors of seed

longevity on the level of populations, genotypes, or varieties appear to be similar to

those driving between-species differences: In general, lighter seeds of the same
species typically survive longer than heavier seeds (Franks et al., 2019; Guzzon et
al., 2018; Lee et al., 2019, but see Mira et al., (2019)). Consequently, seeds of
genotypes or varieties with lighter seeds tend to survive longer in storage than
accessions with heavier seeds, although this effect is inconsistent across studies
(Guzzon et al., 2021; Schutte et al., 2008). The climate at the seeds' origin seems to
be also a critical factor, with populations from warmer and dryer environments
producing longer-lived seeds than populations from colder and more humid
environments (Mondoni et al., 2011, 2014, but see Kochanek et al. (2009)). The
differentiation in seed longevity among populations of the same species can be
driven by two, mutually not exclusive mechanisms: genetic differentiation (Mondoni et
al., 2014) or phenotypic plasticity, where parental growing conditions influence seed
longevity (Kochanek et al., 2010; Mondoni et al., 2014; Sinniah et al., 1998). While
the studies above provide valuable insights, we are still missing a comprehensive
assessment of intraspecific variability in seed longevity across wild species, its
magnitudes and predictors.

To fill this gap, we focused on more than 40 common grassland species represented
by up to 11 accessions from across Europe. We exposed the seeds to artificial
ageing, estimated their longevity, and analysed the data on two levels: Comparison
across accessions within the species and comparison across species.

We expect that (1) seed longevity under artificial ageing conditions differs between
species and accessions within species. The variation among species will be larger
than the variation within species. (2) Within species, longevity will increase with mean
annual temperature and annual precipitation at the seed origin and decrease with
seed mass. (3) Across species, longevity will decrease with seed weight and be
influenced by seed chemical composition, particularly oil content, and the plant
family.

MATERIALS AND METHODS

Study species and seed material

We focused on 41 species common in European grasslands, belonging to 11 families distributed across the phylogenetic tree of angiosperms (Table 1). They represent a wide variety of traits, some potentially associated with seed longevity, specifically seed weight, seed oil and protein content (Probert et al., 2009). All species have orthodox seeds, i.e. they survive desiccation and freezing during storage.

To test for within-species variability, most species were represented by multiple origins, further called accessions. The number of accessions per species was driven by seed availability and varied strongly between species. More than half of the species were represented by at least four origins, while for six species we managed to obtain only one accession (Table 1). In total, the experiment included 188 accessions.

The experiment required large amounts of seeds. Seeds of 14 accessions were obtained by direct collection in nature or by collecting at least 50 young plants in the wild and growing them in a cold greenhouse to produce seeds. However, most seeds come from on-farm propagation for restoration purposes. We are aware that on-farm propagation may cause unintended selection, which could alter seed traits (Conrady et al., 2023; Ensslin et al., 2018). In the seeds we used, this effect will be rather minor because most producers are relatively new in this business and the propagation did not take place for many generations (*personal communication*). Using farm-propagated seeds further has the advantage that the maternal plants grew under optimal conditions and the seeds were harvested when fully ripe. This reduces the variability in seed quality, a characteristic that has a massive effect on storage behaviour (Rahman & Ellis, 2019; Sano et al., 2016; Sinniah et al., 1998).

Artificial ageing

To artificially age the seeds, we exposed them to 45° C and 60% relative humidity in a climate-controlled cabinet (Rumed ®, Rubarth Apparate GmbH, Laatzen, Germany). These conditions enhance similar chemical processes that are typical of

- 6 -

natural ageing during long-term storage and accelerate seed deterioration (Merritt et
al., 2014; Probert et al., 2009).

We exposed the seeds to artificial ageing conditions for 0 (control group, fresh
seeds), 1, 5, 9, 15, 20, 30, 40, 57 and 72 days (adjusted after Probert et al. (2009)).
We put 3 replicates of 30 seeds of every accession and ageing duration into open 1.5
mL tubes in the ageing condition. The position of each accession in the ageing
condition was randomised within the ageing duration. Before the ageing treatment,
we manually scarified Fabaceae seeds with sandpaper to allow for moisture to be
imbibed (Baskin, 2003). In total, our experiment contained 188 accessions, each with
10 ageing treatments and 3 replicates, that is 5,640 replicates containing 30 seeds
each, that is 169,200 seeds.

Germination test

For germination tests, we used 6-well plates with a well-diameter of 39 mm (VWR
International GmbH, Darmstadt, Germany, catalogue number: 734-2323), because
they use space in the climate cabinet more efficiently than single petri dishes. For
species with large seeds (*Bupleurum rotundifolium*, *Centaurea cyanus*, *Bromus*
hordeaceus, *Arrhenatherum elatius*, *Lathyrus nissolia*, *Hordeum murinum*,
Hypochaeris radicata, *Agrostemma githago*, *Poa annua*) we used petri dishes of 6
cm diameter to avoid overcrowding. We wrapped Petri dishes with seeds of *Bromus*
hordeaceus in aluminium foil since light slightly inhibits germination (Andersson et al.,
2002). After the intended ageing duration, we removed the seeds from the ageing
conditions and transferred each replicate into one well or a petri dish laid out with two
filter papers in a fully randomised design. We watered the seeds with a 0.025%
Gibberellic acid solution to break dormancy and closed the plates with transparent
Parafilm® to avoid evaporation. Germination took place in a climate-controlled
cabinet (Rumed®, Rubarth Apparate GmbH, Laatzen, Germany) with a 14 h daily
photoperiod at 20° C and 10 h darkness at 10° C.

We checked for germination every seven days and added distilled water if necessary.
When a radicle was longer than 2 mm, we considered a seed viable and removed it
from the dish. For each replicate with viable seeds, we finished the trial when no

more seeds germinated for at least four weeks. If in a replicate no seeds germinated

at all for ten weeks, we considered the seeds not viable. If samples were affected by

mould, we added 1 ml of a 0.0835% Difenoconazol solution.

Seed traits and climatic data

For each accession, we determined seed weight by weighing 2 replicates of 50 seeds

and calculated the weight of a thousand seeds (further called seed weight for

simplicity). We also obtained the mean annual temperature and precipitation for each

accession location from WorldClim2 using the R package *raster* (Fick & Hijmans,

2017). In wild-collected seeds, we used the exact location of the seed collection. For

farm-propagated seeds, we used the location of the farm, as the initial seeds used to

establish the seed orchard were typically collected in the same region where the

seed producer is located (Bucharova et al., 2019, Figure 1). The oil and protein

content of the seeds was obtained from the Seed Information Database on species

level (Society for Ecological Restoration et al., 2023).

Statistical analyses

Before data analysis, we calculated the initial germination rate of each of the 188

seed accessions. In some species, initial germination rates were higher in short-

duration ageing treatments (one or five days) than in fresh seeds. This was likely

because short exposure to ageing treatment can break dormancy (Hay et al., 2021;

Mira et al., 2019). We thus defined "initial germination" for each accession as the

highest average germination rate of the 0, 1 or 5 days in the ageing treatment. We

removed accessions with initial germination below 50% and those particularly

affected by mould (n = 39).

To analyse seed survival, we used generalized linear models (GLMs) with a

quasibinomial error distribution (to account for overdispersion) and a probit link

function. The probit transformation converts percentage responses onto a linear

scale with a normal error distribution. We modelled the proportion of germinated

seeds per replicate as a function of the number of days in ageing (Wolkis et al.,

2025). The slope of the probit function directly describes the rate of seed viability

loss, and back transformation of the slope (-1/slope) yields σ, the number of days

needed for viability to drop by one probit (Hay et al., 2022; Klepka et al., 2025). We

are aware that most comparative studies of seed longevity in storage measured seed

longevity as p_{50}, the number of days needed for viability to decrease to 50%, yet this

parameter is heavily dependent on the initial seed viability (see Klepka et al. (2025)

for details). While the initial viability can be affected by the environmental effects, e.g.

during seed storage before the experiment, σ is assumed to be a genetic trait that

describes a species, population or genotype, and is thus the actual trait of interest

(Hay et al., 2022). We thus decided to use σ as a measure of seed longevity,

because it is independent of the initial seed viability of given accessions and directly

describes the slope of seed viability loss (Hay et al., 2022; Klepka et al., 2025).

Within-species variability. We obtained the seed longevity estimate (σ) for each

accession. To do this, we related seed germination in each replicate to the time in

ageing conditions in a separate probit model for each accession. If the model failed

to give significant estimates for the slope, or if σ was higher than the maximum

ageing duration in our experiment (72 days), we excluded the accession from further

analyses (n = 14), since we assumed that in these cases the data cover only a very

short portion of the survival curve and the estimated parameters are therefore

unprecise. These models yielded separate σ-values for each accession.

We then tested whether the seed longevity (σ) differs between accessions within

species. This was possible for 26 species for which we had multiple accessions that

passed the initial check (initial germination >50% and significant accession-specific

σ). For this, we related the seed germination in each replicate to the accession

identity, the ageing duration and their interaction as explanatory variables in a

separate probit model for each species. A significant interaction between accession

identity and ageing duration indicates significantly different slopes between

accessions within species and therefore different seed longevities (σ). We further

tested whether the probability of detecting significant differences in σ between

accessions within species depended on the number of accessions representing each

species in our data set. To do this, we related the presence of significant difference

between accessions in each species (binomial yes/no) to the number of accessions

per species in a generalised linear model with binomial error distribution.

To understand drivers of within-species variability in seed longevity in storage, we

related accession-specific longevity (σ) as the response variable to the accession-

specific seed weight, mean annual temperature and precipitation at the accession's

origin as explanatory variables in a multiple linear mixed model. We included the

species' identity as a random factor. For this analysis, we scaled the seed weight

within each species to standardise the variable across species, and included only

species that were represented by at least 2 accessions.

We quantified the proportion of within-species variance explained by the fixed effects

using marginal and conditional R^2 estimates from the MuMIn package. This allowed

us to calculate the contribution of the within-species predictors relative to the

variance remaining after accounting for species identity.

Between-species variability. We obtained the seed longevity estimate (σ) for each

species, common across accessions, regardless whether we detected significant

differences among accessions in the previous step. For this, we related seed

germination in each replicate (and from all accession of given species) to time in

ageing in a probit model for each species. For species with more than one accession,

we added accession identity as an explanatory variable to allow for different

intercepts (initial viability) in individual accessions. These models yielded one σ for

each species. To understand drivers of between-species variability in seed longevity

in storage, we related the log-transformed species-specific σ as the response

variable to the species' mean seed weight and plant family as explanatory variables

in a multiple linear regression model. As protein and oil contents were only available

for 15 study species, we evaluated their effects in a separate multiple linear

regression model.

All data analyses were performed using R version 4.4.2 (2024-10-31 ucrt) (R

Development Core Team, 2024). We log-transformed σ as the response variable to

ensure that model assumptions were met. For all models, we verified that model

assumptions were met by graphically evaluating the residuals (Zuur et al., 2010).

RESULTS

The estimated seed longevity (σ) significantly varied among accessions within 58%

of the species for which we had multiple accessions, specifically in 15 out of 26

species (Table S1). The probability of detecting significant differences among

accessions within species increased with the number of included accessions

(R^2=0.20, p=0.016, Figure S1). Within-species variation was the highest in *Prunella*

vulgaris, where σ varied more than 4-fold (Figure 2), specifically between 12.66 and

53.14 days. This within-species variability was hard to predict, as accession-specific

predictors explained only 14.4% of the variability among accessions, after correcting

for species identity. Specifically, seed longevity decreased with seed weight and

increased with the mean annual temperature at the accession origin, but was

unrelated to the mean annual precipitation (Table 2, Figure 3).

The estimated seed longevity also varied among species (Figure 2), and species s

identity explained 66% of the total variability among accessions. The average seed

longevity was shortest for *Thlaspi arvense* (σ = 5.04 days) and longest for *Lotus*

pedunculatus (σ = 66.70 days). The species seed longevity was unrelated to the

species-specific seed weight (Table 3A, Figure 4A) or the seed oil and protein

content (Table 3B, Figure 4B, C), but we found significant differences between plant

families (Table 3A, Figure 4D).

DISCUSSION

Effective conservation of seeds in ex-situ seed banks requires an understanding of seed storage behaviour not only for individual species but also for accessions within the very same species. We found, for multiple species common in European grasslands, that accessions of the same species commonly differ in seed longevity in storage. This intraspecific variability was partially explained by seed weight and mean annual temperature at the accession origin, but the predictive power of these variables was rather low (2.1% explained variability across all species). On the other hand, species identity was a strong predictor of seed longevity, explaining 66% of the variability among accessions. Our results thus suggest that seed longevity can be reliably predicted at the species level, but predicting accession-specific seed longevity in storage remains challenging.

Intraspecific variability in seed longevity

Seed longevity in storage varied considerably within-species, with accession-specific seed longevity differing significantly in 15 out of 26 species with multiple accessions. The probability of detecting significant differences among accessions within a species increased with the number of accessions included in the study. This suggests that with more accessions available, significant differences would likely be detected in even more species. Consequently, within-species differences in seed longevity in storage are likely very common.

Within-species, seed longevity was partially explained by the mean annual temperature at the collection site, a pattern that was reported previously (Mondoni et al., 2011). Other studies attributed such a positive correlation between temperature and seed longevity mainly to the species identity because they had species represented by single accessions (Merritt et al., 2014; Probert et al., 2009). We did not find any effect of the annual precipitation on seed longevity, which contrasts with previous studies, both within and across species (Merritt et al., 2014; Probert et al., 2009; White et al., 2023). This is likely because the mean annual precipitation was correlated with mean annual temperature (r=0.34), and as we used multivariate models, we were not able to tease apart the effect of both climate variables.

We found a negative relationship between the accession-specific seed weight and

seed longevity, with lighter seeds surviving significantly longer. Previous studies

primarily tested the relationship between seed weight and longevity across species,

yielding mixed results (Davies et al., 2020; Merritt et al., 2014; Pritchard & Dickie,

2003; Probert et al., 2009; Satyanti et al., 2018). Similarly, research on variability

within individual species has also produced inconsistent findings (Franks et al., 2018;

Genna et al., 2020; Guzzon et al., 2018, 2021; Mira et al., 2019; Schutte et al., 2008).

These inconsistent study findings underscore that seed mass remains an unreliable

predictor and highlight the complexity of seed longevity and its drivers. Nevertheless,

our results provide new evidence suggesting that, for a variety of wild grassland

species, lighter seeds may have a slight but consistent advantage over heavier seeds

of the same species in long-term storage survival.

The intraspecific variability in seed longevity may be driven by two mutually non-

exclusive mechanisms – phenotypic plasticity or genetic differentiation. For example,

Mondoni et al. (2011) detected that seeds collected in populations in warmer

environments survived longer in artificial ageing than conspecific seeds collected in

cooler environments. A large proportion of this effect was caused by phenotypic

plasticity, but the pattern was still significant, although weaker, in the second

generation grown in a common environment, which indicates that the effect is

partially heritable and thus genetic (Mondoni et al., 2014). Indeed, genes associated

with seed longevity in storage were identified in some crops and model species

(Bizouerne et al., 2023; Nagel et al., 2015; Raquid et al., 2021; Renard et al., 2020).

Our study worked with seeds that were collected in natural sites or produced in the

region of origin; we thus cannot discriminate whether the observed effects are plastic

or genetic.

After accounting for species identity, within-species predictors explained 14.4% of the

remaining variability among accessions. Still, species identity was the strongest

predictor of seed longevity and explained 66.9%. Our findings align with those of

Kochanek et al. (2009). On a continental scale, they collected data on seed longevity

in seven species represented by 2-8 populations, with populations within species

largely aggregated in a specific biome. Here, the species identity explained 69.1%

(adjusted R^2) of the variability among accessions, leaving 30% to variation among

accessions within species (own analysis of the data presented in Kochanek et al.

(2009)). In contrast, a study by Mondoni et al. found that the intraspecific variability in

seed longevity was much larger than the interspecific one. On a regional scale, they

examined the seed longevity of six ecologically similar species that were each

represented by two populations in contrasting environments, specifically high and low

altitudes. Here, species identity explained only 6% of variability, leaving an

astonishing 94% variation within species (own analysis of the data presented in

Mondoni et al. (2011)). In our study, we detected that species explained about two-

thirds of the variability in seed longevity among accessions, yet this number might be

underestimated because we excluded the species with the most long-lived seeds for

methodical reasons (see methods).

The comparison of our results with the findings of previous studies is complicated by

different measures of seed longevity. We used σ, the rate of seed viability loss, which

is independent of the initial seed viability and thus suitable for comparative studies

across accessions and species. The vast majority of previous studies used p_{50}, the

time till the seed viability decreases to half (e.g. Mondoni et al., 2011; Probert et al.,

2009; Satyanti et al., 2018), a measure that is partially determined by the rate of seed

viability loss (σ), but is heavily dependent on initial seed viability at the start of the

experiment (Klepka et al., 2025). For example, using seeds with an initial viability

ranging from 85 to 99.9%, but with the same rate of seed viability loss (σ), results in a

threefold variation in p_{50}. The p_{50} estimate thus does not appropriately represent

inherent differences between genotypes or populations (Hay et al., 2019). Direct

comparison between our study and many previous studies is thus complex, unless

raw viability data are available to calculate σ.

Interspecific variability in seed longevity

Across species, the only variable that significantly affected seed longevity was the

plant family, with the observed differences between families being roughly in line with

previous studies (Merritt et al., 2014; Mondoni et al., 2011; Probert et al., 2009;

Walters et al., 2005b): Fabaceae seeds were relatively long-lived, probably due to

their physical dormancy (Merritt et al., 2014), there was a large variability within

Asteraceae (Merritt et al., 2014; Walters et al., 2005a), and three of the six shortest-

lived species in our study belonged to the Brassicaceae family (*Thlaspi arvense*,

Cardamine hirsuta, *Barbarea vulgaris*).

Neither seed weight nor chemical composition influenced seed longevity in our

artificial ageing experiment. In general, species-specific seed weight seems to be an

unreliable predictor of seed longevity in storage, as the effects reported in previous

literature strongly vary from negative, over no effect to, rarely, positive (Davies et al.,

2020; Merritt et al., 2014; Pritchard & Dickie, 2003; Probert et al., 2009; Satyanti et

al., 2018). Similarly, the chemical composition of the seeds did not predict seed

longevity, which corresponds to previous empirical studies of wild species (Kochanek

et al., 2009; Walters et al., 2005b). The expected effect of the chemical composition

on seed longevity in storage, especially oil content, is based on theoretical

expectations and comparisons among oil crops, cereals, and legumes (Narayana

Murthy & Sun, 2000; Pritchard & Dickie, 2003 Nagel et al., 2015), but the empirical

evidence provided by others and us shows that it is likely not true for wild species.

CONCLUSIONS

Our study demonstrates that in the majority of species, there is substantial within-

species variation in the rate of seed viability loss in storage. Conservation seed

banks sometimes design viability monitoring of individual accessions based on

species identity and initial seed viability only, assuming that accessions of the same

species lose viability at the same rate when stored under the same conditions (Hay &

Whitehouse, 2017). Our data show that this assumption is not valid and support the

existing concerns that this method might be suboptimal.

ACKNOWLEDGEMENTS

We thank Johannes Kasper, Lena Lerbs, Christina Mengel, Sarah Paschen, Nadine

Pluquette, Annika Sommerfeld, and Helene Villhauer for technical assistance. This

project was funded by a scholarship of Marburg University Research Academy

(MARA) to LK.

AUTHOR CONTRIBUTIONS

Lea Klepka: Investigation, Formal analysis, Data Curation, Writing – original Draft,

Visualization, Funding Acquisition. Sacha Liepelt: Investigation, Writing –

Review&Editing, Anna Bucharova: Conceptualization, Methodology, Resources,

Writing – Review & Editing, Supervision

DATA AVAILABILITY STATEMENT

The data supporting the findings of this study are not publicly available at the time of

submission but will be deposited in an appropriate public repository upon acceptance

of the manuscript. Data will be made available to editors and reviewers upon

reasonable request.

REFERENCES

Andersson, L., Milberg, P., Schütz, W., & Steinmetz, O. (2002). Germination characteristics and emergence time of annual Bromus species of differing weediness in Sweden. *Weed Research*, *42*(2), 135–147. https://doi.org/10.1046/j.1365-3180.2002.00269.x

Baskin, C. C. (2003). Breaking physical dormancy in seeds – focussing on the lens. *New Phytologist*, *158*(2), 229–232. https://doi.org/10.1046/j.1469-8137.2003.00751.x

Bizouerne, E., Ly Vu, J., Ly Vu, B., Diouf, I., Bitton, F., Causse, M., Verdier, J., Buitink, J., & Leprince, O. (2023). Genetic Variability in Seed Longevity and Germination Traits in a Tomato MAGIC Population in Contrasting Environments. *Plants (Basel, Switzerland)*, *12*(20), 3632. https://doi.org/10.3390/plants12203632

Bucharova, A., Bossdorf, O., Hölzel, N., Kollmann, J., Prasse, R., & Durka, W. (2019). Mix and match: Regional admixture provenancing strikes a balance among different seed-sourcing strategies for ecological restoration. *Conservation Genetics*, *20*(1), 7–17. https://doi.org/10.1007/s10592-018-1067-6

Cochrane, J. A., Crawford, A. D., Monks, L. T., Cochrane, J. A., Crawford, A. D., & Monks, L. T. (2007). The significance of ex situ seed conservation to reintroduction of threatened plants. *Australian Journal of Botany*, *55*(3), 356–361. https://doi.org/10.1071/BT06173

Cohen, J. I., Williams, J. T., Plucknett, D. L., & Shands, H. (1991). Ex Situ Conservation of Plant Genetic Resources: Global Development and Environmental Concerns. *Science*, *253*(5022), 866–872. https://doi.org/10.1126/science.253.5022.866

Conrady, M., Lampei, C., Bossdorf, O., Hölzel, N., Michalski, S., Durka, W., & Bucharova, A. (2023). Plants cultivated for ecosystem restoration can evolve toward a domestication syndrome. *Proceedings of the National Academy of Sciences*, *120*(20), e2219664120. https://doi.org/10.1073/pnas.2219664120

Davies, R. M., Hudson, A. R., Dickie, J. B., Cook, C., O'Hara, T., & Trivedi, C. (2020). Exploring seed longevity of UK native trees: Implications for *ex situ* conservation. *Seed Science Research*, *30*(2), 101–111. https://doi.org/10.1017/S0960258520000215

Delouche, J., & Baskin, C. (2021). Accelerated Aging Techniques for Predicting the Relative Storability of Seed Lots. *All Articles*. https://scholarsjunction.msstate.edu/seedtechpapers/10

ENSCONET. (2009, June 16). *ENSCONET Curation Protocols & Recommendations*. http://ensconet.maich.gr/PDF/Curation_protocol_English.pdf

Ensslin, A., Van de Vyver, A., Vanderborght, T., & Godefroid, S. (2018). Ex situ cultivation entails high risk of seed dormancy loss on short-lived wild plant species. *Journal of Applied Ecology*, *55*(3), 1145–1154. https://doi.org/10.1111/1365-2664.13057

Etterson, J. R., Franks, S. J., Mazer, S. J., Shaw, R. G., Gorden, N. L. S., Schneider, H. E., Weber, J. J., Winkler, K. J., & Weis, A. E. (2016). Project Baseline: An unprecedented resource to study plant evolution across space and time. *American Journal of Botany*, *103*(1), 164–173. https://doi.org/10.3732/ajb.1500313

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: New 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology*, *37*(12), 4302–4315. https://doi.org/10.1002/joc.5086

Franks, S. J., Hamann, E., & Weis, A. E. (2018). Using the resurrection approach to understand contemporary evolution in changing environments. *Evolutionary Applications*, *11*(1), 17–28. https://doi.org/10.1111/eva.12528

Franks, S. J., Sekor, M. R., Davey, S., & Weis, A. E. (2019). Artificial seed aging reveals the invisible fraction: Implications for evolution experiments using the resurrection approach. *Evolutionary Ecology*, *33*(6), 811–824. https://doi.org/10.1007/s10682-019-10007-2

Genna, N. G., Walters, C., & Pérez, H. E. (2020). Viability and vigour loss during storage of Rudbeckia mollis seeds having different mass: An intra-specific perspective. *Seed Science Research*, *30*(2), 122–132. https://doi.org/10.1017/S0960258520000161

Guzzon, F., Gianella, M., Velazquez Juarez, J. A., Sanchez Cano, C., & Costich, D. E. (2021). Seed longevity of maize conserved under germplasm bank conditions for up to 60 years. *Annals of Botany*, *127*(6), 775–785. https://doi.org/10.1093/aob/mcab009

Guzzon, F., Orsenigo, S., Gianella, M., Müller, J. V., Vagge, I., Rossi, G., & Mondoni, A. (2018). Seed heteromorphy influences seed longevity in Aegilops. *Seed Science Research*, *28*(4), 277–285. https://doi.org/10.1017/S096025851800034X

Hay, F. R., Davies, R. M., Dickie, J. B., Merritt, D. J., & Wolkis, D. M. (2022). More on seed longevity phenotyping. *Seed Science Research*, *32*(3), 144–149. https://doi.org/10.1017/S0960258522000034

Hay, F. R., & Probert, R. J. (2013). Advances in seed conservation of wild plant species: A review of recent research. *Conservation Physiology*, *1*(1), cot030–cot030. https://doi.org/10.1093/conphys/cot030

Hay, F. R., Valdez, R., Lee, J.-S., & Sta. Cruz, P. C. (2019). Seed longevity phenotyping: Recommendations on research methodology. *Journal of Experimental Botany*, *70*(2), 425–434. https://doi.org/10.1093/jxb/ery358

Hay, F. R., & Whitehouse, K. J. (2017). Rethinking the approach to viability monitoring in seed genebanks. *Conservation Physiology*, *5*(1), cox009. https://doi.org/10.1093/conphys/cox009

Hay, F. R., Whitehouse, K. J., Ellis, R. H., Sackville Hamilton, N. R., Lusty, C., Ndjiondjop, M. N., Tia, D., Wenzl, P., Santos, L. G., Yazbek, M., Azevedo, V. C. R., Peerzada, O. H., Abberton, M., Oyatomi, O., de Guzman, F., Capilit, G., Muchugi, A., & Kinyanjui, Z. (2021). CGIAR genebank viability data reveal inconsistencies in seed collection management. *Global Food Security*, *30*, 100557. https://doi.org/10.1016/j.gfs.2021.100557

Klepka, L., Carta, A., Hay, F. R., & Bucharova, A. (2025). *Limitations of p50 as a measure of seed longevity and the way forward* (p. 2025.06.11.659081). bioRxiv. https://doi.org/10.1101/2025.06.11.659081

Kochanek, J., Buckley, Y. M., Probert, R. J., Adkins, S. W., & Steadman, K. J. (2010). Pre-zygotic parental environment modulates seed longevity. *Austral Ecology*, *35*(7), 837–848. https://doi.org/10.1111/j.1442-9993.2010.02118.x

Kochanek, J., Steadman, K. J., Probert, R. J., & Adkins, S. W. (2009). Variation in seed
longevity among different populations, species and genera found in collections
from wild Australian plants. *Australian Journal of Botany*, *57*(2), 123–131.
https://doi.org/10.1071/BT09023

Lee, J.-S., Velasco-Punzalan, M., Pacleb, M., Valdez, R., Kretzschmar, T., McNally, K.
L., Ismail, A. M., Cruz, P. C. S., Sackville Hamilton, N. R., & Hay, F. R. (2019).
Variation in seed longevity among diverse Indica rice varieties. *Annals of*
Botany, *124*(3), 447–460. https://doi.org/10.1093/aob/mcz093

Merritt, D. J., & Dixon, K. W. (2011). Restoration Seed Banks—A Matter of Scale.
Science, *332*(6028), 424–425. https://doi.org/10.1126/science.1203083

Merritt, D. J., Martyn, A. J., Ainsley, P., Young, R. E., Seed, L. U., Thorpe, M., Hay, F.
R., Commander, L. E., Shackelford, N., Offord, C. A., Dixon, K. W., & Probert,
R. J. (2014). A continental-scale study of seed lifespan in experimental
storage examining seed, plant, and environmental traits associated with
longevity. *Biodiversity and Conservation*, *23*(5), 1081–1104.
https://doi.org/10.1007/s10531-014-0641-6

Millennium Seed Bank. (2022). *Assessing a Population for Seed Collection*.
https://brahmsonline.kew.org/Content/Projects/msbp/resources/Training/02-
Assessing-population.pdf

Mira, S., Veiga-Barbosa, L., & Pérez-García, F. (2019). Seed dormancy and longevity
variability of *Hirschfeldia incana* L. during storage. *Seed Science Research*,
29(2), 97–103. https://doi.org/10.1017/S0960258519000072

Mondoni, A., Orsenigo, S., Donà, M., Balestrazzi, A., Probert, R. J., Hay, F. R., Petraglia,
A., & Abeli, T. (2014). Environmentally induced transgenerational changes in
seed longevity: Maternal and genetic influence. *Annals of Botany*, *113*(7),
1257–1263. https://doi.org/10.1093/aob/mcu046

Mondoni, A., Probert, R. J., Rossi, G., Vegini, E., & Hay, F. R. (2011). Seeds of alpine
plants are short lived: Implications for long-term conservation. *Annals of*
Botany, *107*(1), 171–179. https://doi.org/10.1093/aob/mcq222

Nagel, M., Kranner, I., Neumann, K., Rolletschek, H., Seal, C. E., Colville, L.,
Fernández-Marín, B., & Börner, A. (2015). Genome-wide association mapping
and biochemical markers reveal that seed ageing and longevity are intricately
affected by genetic background and developmental and environmental
conditions in barley. *Plant, Cell & Environment*, *38*(6), 1011–1022.
https://doi.org/10.1111/pce.12474

Narayana Murthy, U. M., & Sun, W. Q. (2000). Protein modification by Amadori and
Maillard reactions during seed storage: Roles of sugar hydrolysis and lipid
peroxidation. *Journal of Experimental Botany*, *51*(348), 1221–1228.
https://doi.org/10.1093/jexbot/51.348.1221

Niedzielski, M., Walters, C., Luczak, W., Hill, L. M., Wheeler, L. J., & Puchalski, J.
(2009). Assessment of variation in seed longevity within rye, wheat and the
intergeneric hybrid triticale. *Seed Science Research*, *19*(4), 213–224.
https://doi.org/10.1017/S0960258509990110

Ninoles, R., Planes, D., Arjona, P., Ruiz-Pastor, C., Chazarra, R., Renard, J., Bueso, E.,
Forment, J., Serrano, R., Kranner, I., Roach, T., & Gadea, J. (2022).
Comparative analysis of wild-type accessions reveals novel determinants of
Arabidopsis seed longevity. *PLANT CELL AND ENVIRONMENT*, *45*(9),
2708–2728. https://doi.org/10.1111/pce.14374

Peres, S. (2016). Saving the gene pool for the future: Seed banks as archives. *Studies in*
History and Philosophy of Science Part C: Studies in History and Philosophy
of Biological and Biomedical Sciences, *55*, 96–104.
https://doi.org/10.1016/j.shpsc.2015.09.002

Pritchard, H., & Dickie, J. (2003). *Predicting Seed Longevity: The use and abuse of seed*
viability equations (pp. 653–721).

Probert, R. J., Daws, M. I., & Hay, F. R. (2009). Ecological correlates of ex situ seed
longevity: A comparative study on 195 species. *Annals of Botany*, *104*(1), 57–
69. https://doi.org/10.1093/aob/mcp082

Rahman, S. M. A., & Ellis, R. H. (2019). Seed quality in rice is most sensitive to drought
and high temperature in early seed development. *Seed Science Research*,
29(4), 238–249. https://doi.org/10.1017/S0960258519000217

Raquid, R., Kohli, A., Reinke, R., Dionisio-Sese, M., Kwak, J., Chebotarov, D., Mo, Y., &
Lee, J.-S. (2021). Genetic factors enhancing seed longevity in tropical
japonica rice. *Current Plant Biology*, *26*, 100196.
https://doi.org/10.1016/j.cpb.2021.100196

Renard, J., Niñoles, R., Martínez-Almonacid, I., Gayubas, B., Mateos-Fernández, R.,
Bissoli, G., Bueso, E., Serrano, R., & Gadea, J. (2020). Identification of novel
seed longevity genes related to oxidative stress and seed coat by genome-
wide association studies and reverse genetics. *Plant, Cell & Environment*,
43(10), 2523–2539. https://doi.org/10.1111/pce.13822

Sano, N., Rajjou, L., North, H. M., Debeaujon, I., Marion-Poll, A., & Seo, M. (2016).
Staying Alive: Molecular Aspects of Seed Longevity. *Plant and Cell*
Physiology, *57*(4), 660–674. https://doi.org/10.1093/pcp/pcv186

Satyanti, A., Nicotra, A. B., Merkling, T., & Guja, L. K. (2018). Seed mass and elevation
explain variation in seed longevity of Australian alpine species. *Seed Science*
Research, *28*(4), 319–331. https://doi.org/10.1017/S0960258518000090

Schutte, B. J., Regnier, E. E., & Harrison, S. K. (2008). The association between seed
size and seed longevity among maternal families in Ambrosia trifida L.
populations. *Seed Science Research*, *18*(4), 201–211.
https://doi.org/10.1017/S0960258508082974

Sinniah, U. R., Ellis, R. H., & John, P. (1998). Irrigation and Seed Quality Development
in Rapid-cycling Brassica: Seed Germination and Longevity. *Annals of*
Botany, *82*(3), 309–314. https://doi.org/10.1006/anbo.1998.0748

Society for Ecological Restoration, International Network for Seed Based Restoration, &
Royal Botanical Gardens Kew. (2023, February). *Seed Information Database*
(SID). https://ser-sid.org/

Walters, C., Wheeler, L. M., & Grotenhuis, J. M. (2005a). Longevity of seeds stored in a
genebank: Species characteristics. *Seed Science Research*, *15*(1), 1–20.
https://doi.org/10.1079/SSR2004195

Walters, C., Wheeler, L. M., & Grotenhuis, J. M. (2005b). Longevity of seeds stored in a
genebank: Species characteristics. *Seed Science Research*, *15*(1), 1–20.
https://doi.org/10.1079/SSR2004195

Wambugu, P. W., Nyamongo, D. O., & Kirwa, E. C. (2023). Role of Seed Banks in
Supporting Ecosystem and Biodiversity Conservation and Restoration.
Diversity, *15*(8), Article 8. https://doi.org/10.3390/d15080896

White, F. J., Hay, F. R., Abeli, T., & Mondoni, A. (2023). Two decades of climate change
alters seed longevity in an alpine herb: Implications for ex situ seed
conservation. *Alpine Botany*, *133*(1), 11–20. https://doi.org/10.1007/s00035-
022-00289-8

Wolkis, D., Carta, A., Rezaei, S., & Hay, F. R. (2025). Seed longevity: Analysing post-
storage germination data in R to fit the viability equation. *Seed Science
Research*, 1–8. https://doi.org/10.1017/S0960258524000291

Zani, D., & Müller, J. V. (2017). Climatic control of seed longevity of *Silene* during the
post-zygotic phase: Do seeds from warm, dry climates possess higher
maturity and desiccation tolerance than seeds from cold, wet climates?
Ecological Research, *32*(6), 983–994. https://doi.org/10.1007/s11284-017-
1508-6

Zuur, A. F., Ieno, E. N., & Elphick, C. S. (2010). A protocol for data exploration to avoid
common statistical problems. *Methods in Ecology and Evolution*, *1*(1), 3–14.
https://doi.org/10.1111/j.2041-210X.2009.00001.x

Figure 1: Origins of the 135 seed accessions used in the study and included in the

analyses. Each point represents the location of a seed producer as a proxy for the

origins of the individual seed accessions and thus, is the origin of multiple species. In

three cases, the points represent the location of own seed collection.

Figure 2: Estimated seed longevity (σ) and 95% confidence intervals for 33 species
and 135 accessions. Blue points and bars represent species-level estimates; green
points and bars show accession-level estimates. Asterisks mark significant
differences between accessions within species. Numbers in brackets indicate the
number of accessions for species represented by multiple accessions.

Longevity (σ)

Figure 3: Relationship between seed longevity (σ) of each accession, accession-
specific seed traits, and environmental variables. Each point represents a seed
accession. Seed longevity was adjusted for species identity. (A) Mean annual
temperature, (B) annual precipitation at the origins of the accessions, and (C)
accession-specific seed mass (scaled); For full model results, see Table 2.

Figure 4: Relationship between species seed longevity (σ) and species-specific seed

traits. Each point represents one species. (A) Seed mass, average across

accessions of the same species; species-specific (B) protein and (C) oil content in

the seeds; and (D) plant families. The dashed lines indicate non-significant

relationships. For full model results, see Table 3.

Table 1: Species included in the study. The number of accessions is the number of

accessions used for the analyses, and in brackets is the initial number of accessions.

Seed weight is the weight of 1000 seeds, averaged across all accessions of the

same species. Oil and protein content are obtained from the Seed Information

Database (Society for Ecological Restoration et al., 2023). Life cycle: A = annual, P =

perennial, B = biennial. *: Information of conspecific

Species	Number of accessions [initial number]	Seed weight [g]	Family	Life cycle	Oil content [%]	Protein content [%]
Achillea millefolium	7 [10]	0.14	Asteraceae	P	27.03	33.1
Agrostemma githago	2 [2]	10.02	Caryophyllaceae	A	6.8	15.3
Ajuga reptans	0 [4]	1	Lamiaceae	P		
Anthemis arvensis	0 [1]	0.37	Asteraceae	A/B		
Arrhenatherum elatius	2 [5]	2.83	Poaceae	P	7.13	24.4
Barbarea vulgaris	3 [4]	0.39	Brassicaceae	A/B	31	15.6
Bromus hordeaceus	3 [3]	4.38	Poaceae	A		
Bupleurum rotundifolium	1 [1]	2.63	Apiaceae	A		
Capsella bursa-pastoris	2 [2]	0.15	Brassicaceae	A	30.53	28.23
Cardamine hirsuta	2 [2]	0.12	Brassicaceae	A		
Centaurea cyanus	1 [7]	3.94	Asteraceae	A	24.67	16.57
Centaurea jacea	6 [8]	1.72	Asteraceae	P		
Cerastium holosteoides	1 [1]	0.07	Caryophyllaceae	B/P		
Chelidonium majus	3 [5]	0.71	Papaveraceae	B		
Daucus carota	8 [9]	0.83	Apiaceae	A/B	19.26	24.38
Dianthus deltoides	8 [8]	0.2	Caryophyllaceae	P		
Galium wirtgenii	0 [1]	0.63	Rubiaceae	P		
Geranium pusillum	0 [1]	1	Geraniaceae	A		
Hordeum murinum	1 [1]	15.41	Poaceae	A	1.4	23.5
Hypochaeris radicata	7 [7]	0.74	Asteraceae	P		
Lamium album	0 [2]	1.57	Lamiaceae	P		
Lamium purpureum	0 [3]	0.69	Lamiaceae	A		
Lathyrus nissolia	1 [1]	7.13	Fabaceae	A		
Linaria vulgaris	5 [7]	0.15	Plantaceae	P		
Lotus corniculatus	7 [7]	1.33	Fabaceae	P	6.53	36.18
Lotus pedunculatus	1 [2]	0.52	Fabaceae	P		
Lychnis flos-cuculi	8 [8]	0.16	Caryophyllaceae	P		
Medicago lupulina	3 [6]	1.33	Fabaceae	A	5.45	35.57
Papaver rhoeas	0 [9]	0.1	Papaveraceae	A	43.47	24.53

Plantago lanceolata	11 [11]	1.49	Plantaginaceae	P	6.73	17.47
Plantago media	5 [6]	0.34	Plantaginaceae	P(/B/A)		
Poa annua	2 [2]	0.44	Poaceae	A/B		
Prunella vulgaris	9 [9]	0.77	Lamiaceae	P	21.7	20.2
Scandix pecten-veneris	0 [1]		Apiaceae	A		
Senecio vulgaris	2 [2]	0.17	Asteraceae	A		
Silene vulgaris	9 [9]	0.77	Caryophyllaceae	P	2.7	18.1
Stellaria media	2 [3]	0.46	Caryophyllaceae	A	5.9	17.8
Thlaspi arvense	1 [3]	1.18	Brassicaceae	A	31.2	24.9
Trifolium aureum	2 [2]	0.42	Fabaceae	A		
Trifolium campestre	3 [5]	0.31	Fabaceae	A/B		
Veronica chamaedrys	7 [8]	0.14	Plantaginaceae	P		

670 Table 2: **Intraspecific** variability in seed longevity in storage, results of the linear

671 mixed model testing the effect of accession-specific seed weight, mean annual

672 temperature, and annual precipitation on accession-specific seed longevity in storage

673 with species identity as random factor. Analysis of variance with error type III.

674 Significant effects (P < 0.05) are highlighted in bold.

	df	Chi Sq	p-value	$_{marg}R^2$	$_{cond}R^2$
				0.05	0.72
Accession-specific seed weight	1	4.23	**0.040**		
Mean annual temperature	1	7.30	**0.007**		
Annual precipitation	1	2.25	0.134		

676 Table 3: **Interspecific** variability in seed longevity in storage, results of the linear
677 model testing the effect of (A) the family, the species-specific seed weight, and (B)
678 the protein and oil content on species-specific seed longevity. Analysis of variance
679 with error type III. Significant effects ($p < 0.05$) are highlighted in bold.

		df	Resid.df	F	p	R^2
A			23			0.49
	Family	8		2.72	**0.029**	
	Species-specific seed weight	1		0.03	0.865	
B			12			0.22
	Protein content	1		0.77	0.125	
	Oil content	1		0.70	0.419	

Figure S1:

Relationship between the number of accessions per species and the probability of

detecting significant differences among accessions within species. The line shows a

logistic regression with the 95% confidence interval.

Table S1: Species-specific seed survival models. Bold type indicates a significant

effect of the parameter. A significant interaction between the accession identity and

the ageing duration means significantly different seed longevities between

accessions within species.

Species	Number of Accessions	Ageing Duration			Accession Identity			Accession × Ageing		
		Df	F-value	p-value	Df	F-value	p-value	Df	F-value	p-value
Achillea millefolium	7	1	20.05	**<0.001**	6	1.24	0.290	6	5.96	**<0.001**
Agrostemma githago	2	1	37.33	**<0.001**	1	0.24	0.623	1	0.07	0.788
Arrhenatherum elatius	2	1	66.81	**<0.001**	1	1.10	0.298	1	6.41	**0.014**
Barbarea vulgaris	3	1	124.86	**<0.001**	2	56.90	**<0.001**	2	5.52	**0.006**
Bromus hordeaceus	3	1	16.95	**<0.001**	2	11.07	**<0.001**	2	2.49	0.089
Capsella bursa-pastoris	2	1	30.59	**<0.001**	1	1.98	0.165	1	0.03	0.871
Cardamine hirsuta	2	1	194.27	**<0.001**	1	4.24	**0.044**	1	8.65	**0.005**
Centaurea jacea	6	1	50.24	**<0.001**	5	7.59	**<0.001**	5	9.85	**<0.001**
Chelidonium majus	3	1	101.61	**<0.001**	2	19.83	**<0.001**	2	5.23	**0.007**
Daucus carota	8	1	84.95	**<0.001**	7	11.50	**<0.001**	7	2.50	**0.017**
Dianthus deltoides	8	1	99.55	**<0.001**	7	5.94	**<0.001**	7	8.28	**<0.001**
Hypochaeris radicata	7	1	80.70	**<0.001**	6	10.98	**<0.001**	6	5.19	**<0.001**
Linaria vulgaris	5	1	23.02	**<0.001**	4	11.62	**<0.001**	4	2.09	0.086
Lotus corniculatus	7	1	26.03	**<0.001**	6	4.06	**0.001**	6	1.58	0.155
Lychnis flos-cuculi	8	1	82.17	**<0.001**	7	18.94	**<0.001**	7	1.98	0.059
Medicago lupulina	3	1	23.12	**<0.001**	2	9.84	**<0.001**	2	1.21	0.305
Plantago lanceolata	11	1	123.03	**<0.001**	10	12.22	**<0.001**	10	5.14	**<0.001**
Plantago media	5	1	56.65	**<0.001**	4	22.08	**<0.001**	4	5.59	**<0.001**
Poa annua	2	1	94.97	**<0.001**	1	0.02	0.877	1	2.47	0.122
Prunella vulgaris	9	1	37.96	**<0.001**	8	4.85	**<0.001**	8	6.50	**<0.001**
Senecio vulgaris	2	1	149.25	**<0.001**	1	2.91	0.093	1	0.81	0.372
Silene vulgaris	9	1	139.81	**<0.001**	8	12.98	**<0.001**	8	7.27	**<0.001**
Stellaria media	2	1	46.77	**<0.001**	1	1.25	**0.268**	1	0.31	0.579
Trifolium aureum	2	1	29.80	**<0.001**	1	5.75	**0.020**	1	0.12	0.732
Trifolium campestre	3	1	87.20	**<0.001**	2	2.68	0.074	2	4.83	**0.010**
Veronica chamaedrys	7	1	128.71	**<0.001**	6	10.05	**<0.001**	6	5.46	**<0.001**