Large interspecific and intraspecific variation in seed

longevity in storage

Lea Klepka[1*], Sascha Liepelt[1] & Anna Bucharova[1]

[1] Conservation Biology, Department of Biology, Philipps University Marburg, Karl-
von-Frisch-Straße 8, 35043 Marburg, Germany

* Corresponding author: Lea Klepka, lea.klepka@biologie.uni-marburg.de

Running head

Intraspecific variation in seed longevity in storage

Author contribution

Lea Klepka: Investigation, Formal analysis, Data Curation, Writing – original Draft,
Visualization, Funding Acquisition. **Sacha Liepelt**: Investigation, Writing –
Review&Editing, **Anna Bucharova**: Conceptualization, Methodology, Resources,
Writing – Review & Editing, Supervision

Acknowledgements

We thank Johannes Kasper, Lena Lerbs, Christina Mengel, Sarah Paschen, Nadine
Pluquette, Annika Sommerfeld, and Helene Villhauer for technical assistance. This
project was funded by a scholarship of Marburg University Research Academy
(MARA) to LK.

Declaration of Interest statement

Authors have no conflicts of interest.

ABSTRACT

Premise: Conservation seed banks are an essential tool for plant ex-situ

conservation. However, seeds stored for long periods of time slowly deteriorate and

lose viability. While seed longevity is known to vary between species, the extent of

variability within species and the factors driving it remain unclear.

Methods: We focused on both inter- and intraspecific variability in seed longevity and

its determinants. We studied 41 common grassland species and 188 seed

accessions from across Europe. We exposed the seeds to artificial ageing conditions

(60% RH, 45°C) and estimated the seed longevity (p_{50}), defined as the time required

for viability to decline to 50% of the initial viability. We then tested the relationships

between seed longevity and both species- and origin-specific factors.

Results: Variability in seed longevity was only slightly greater between species than

within species, with species identity explaining 51.9% of the total variation in seed

longevity among seed accessions, leaving 48.1% of the variation within species. In

interspecific species comparison, seed viability was predicted only by initial seed

viability, but unrelated to species-specific seed mass or seed chemical composition.

Within species, seed longevity increased with the initial viability of the seed accession

and mean annual temperature at the accession origin, but was unrelated to seed

mass or precipitation in the origin, and the residual variance was substantial.

Conclusion: Our findings highlight that seed longevity on the level of individual seed

accessions remains challenging to predict.

KEYWORDS

artificial seed ageing, conservation seed bank, species comparison, intraspecific

variability, seed longevity, seed storage, within-species variability

INTRODUCTION

Biodiversity conservation is one of the critical global challenges, and *ex situ* methods play an increasingly important role in this task (Cohen et al., 1991). One of the most common ex-situ methods for plant conservation are seed banks, dedicated facilities that store seeds of wild plant species, particularly the vulnerable ones (Peres, 2016). The stored seeds represent a snapshot of past populations and are valuable resources for species reintroductions or ecosystem restoration, and also for scientific research (Cochrane et al., 2007; Etterson et al., 2016; Wambugu et al., 2023). Thus, the ultimate task of conservation seed banks is to keep the seeds viable for decades or even centuries.

Seed survival in storage depends on the type of seeds and storage conditions. Some types of seeds, so-called recalcitrant seeds, do not survive desiccation, making long-term storage impossible. Other types of seeds, called orthodox, can survive desiccation, and they can be stored for a couple of months up to centuries, depending on the species (Hay & Probert, 2013). Conservation seed banks aim to maintain seed viability as long as possible by storing the seeds under low temperatures and humidity to slow down internal chemical processes such as lipid peroxidation and DNA damage (Nagel et al., 2015; Walters et al., 2005a). The seed longevity also depends on the quality and viability of the seeds (Hay & Probert, 2013; Niedzielski et al., 2009), and thus, guidelines for seed banking recommend storing accessions with high initial viability (ENSCONET, 2009; Millenium Seed Bank, 2022). Yet, even highly viable seeds stored under optimal conditions slowly deteriorate and eventually die. This mortality is of course faster when the seeds are stored in suboptimal conditions, for example when storing seeds for large-scale ecosystem restoration in ambient conditions at a farm (Merritt & Dixon, 2011). To optimise the long-term effectiveness of seed storage – be it for conservation, research or ecosystem restoration purposes – we need to understand which factors predict seed longevity in storage.

Studying seed longevity presents unique challenges, as it requires comparing old and fresh seeds of the same seed lot, which would need time travel. To circumvent this obstacle, researchers have developed artificial ageing protocols which simulate natural ageing by exposing the seeds to high temperatures and high relative humidity

(Delouche & Baskin, 2021). These conditions accelerate the same or at least similar

chemical processes typical of natural ageing during long-term storage (Delouche &

Baskin, 2021). Although this method may not perfectly replicate natural ageing, the

longevity of seeds in artificial ageing correlates with the longevity of seeds stored in

seed banks both across and within species (Ninoles et al., 2022; Probert et al.,

2009). Nevertheless, artificial ageing is the best method we currently have to study

seed longevity.

Seed survival in artificial ageing differs between species, and to some degree also

between higher taxa like plant families, although large plant families harbour

substantial variation in seed longevity (Probert et al., 2009; Walters et al., 2005a).

Studies aiming to detect general patterns in seed longevity beyond evolutionary

history yielded mixed results. Endospermic seeds survive shorter in storage than

non-endospermic ones (Probert et al., 2009). The influence of seed weight is less

clear; some studies suggest that seeds of small-seeded species survive longer

(Satyanti et al., 2018), while others report no effect of seed weight (Davies et al.,

2020; Merritt et al., 2014; Probert et al., 2009). Seed longevity is also affected by

climatic conditions at seed collection sites: Seeds from hot or dry environments

generally survive longer than those from cooler or wetter conditions (Merritt et al.,

2014; Probert et al., 2009; Zani & Müller, 2017). Furthermore, seed longevity should

hypothetically depend on the seed's chemical composition because high oil content

should reduce seed longevity due to ongoing lipid peroxidation in storage (Narayana

Murthy & Sun, 2000; Pritchard & Dickie, 2003), yet the results of empirical studies are

inconsistent (Nagel et al., 2015; Probert et al., 2009; Walters et al., 2005a).

In contrast to between-species comparison, there is less comprehensive information

on within-species variability on seed longevity in storage. Most data come from

cultivated crops or model species (e.g. Bizouerne et al., 2023; Franks et al., 2018;

Guzzon et al., 2021; Lee et al., 2019), with only a few studies addressing natural

populations of wild species (Genna et al., 2020; Kochanek et al., 2009; Mondoni et

al., 2011; White et al., 2023). These indicate that the variation in seed longevity within

a species can be as substantial as the variation between species from the same

ecosystem (Kochanek et al., 2009; Mondoni et al., 2011). The predictors of seed

longevity on the level of populations, genotypes, or varieties appear to be similar to

those driving between-species differences: In general, lighter seeds of the same

species typically survive longer than heavier seeds (Franks et al., 2019; Guzzon et

al., 2018; Lee et al., 2019, but see Mira et al., (2019)). Consequently, seeds of

genotypes or varieties with lighter seeds tend to survive longer in storage than

accessions with heavier seeds, although this effect is inconsistent across studies

(Guzzon et al., 2021; Schutte et al., 2008). The climate at the seeds' origin seems to

be also a critical factor, with populations from warmer and dryer environments

producing longer-lived seeds than populations from colder and more humid

environments (Mondoni et al., 2011, 2014, but see Kochanek et al., (2009)). The

differentiation in seed longevity among populations of the same species can be

driven by two, mutually not exclusive mechanisms: genetic differentiation (Mondoni et

al., 2014) or phenotypic plasticity, where parental growing conditions influence seed

longevity (Kochanek et al., 2010; Mondoni et al., 2014; Sinniah et al., 1998). While

the studies above provide valuable insights, we are still missing a comprehensive

assessment of intraspecific variability in seed longevity across wild species, its

magnitudes and predictors.

To fill this gap, we focused on more than 40 common grassland species represented

by up to 11 accessions from across Europe. We exposed the seeds to artificial

ageing, estimated their longevity, and analysed the data on two levels: Comparison

across species and comparison across accessions within the species.

We expect that (1) seed longevity under artificial ageing conditions differs between

species and accessions within species. The variation among species will be larger

than the variation within species. (2) Across species, longevity will increase with seed

quality (measured as initial viability) and decrease with seed weight. It will also be

influenced by seed chemical composition, particularly oil content, and the plant

family. (3) Within species, longevity will be higher in accessions with high initial

viability, lighter seeds and coming from from warmer and drier environments.

MATERIAL AND METHODS

MATERIAL AND METHODS

Study species and seed material

We focused on 41 species common in European grasslands, belonging to 11 families

distributed across the phylogenetic tree of angiosperms (Table 1). They represent a

wide variety of traits, some potentially associated with seed longevity, specifically

seed weight, seed oil and protein content (Probert et al., 2009). All species have

orthodox seeds, i.e. they survive desiccation and freezing during storage.

To test for within-species variability, most species were represented by multiple

origins, further called accessions. The number of accessions per species was driven

by seed availability and strongly varied between species. More than half of the

species were represented by at least four origins, while for six species we managed

to obtain only one accession (Table 1). In total, the experiment included 188

accessions.

The experiment required large amounts of seeds. Part of the seeds was obtained by

direct collection in nature, or collecting at least 50 small plants in the wild and

growing them in a cold greenhouse to produce seeds. However, most seeds come

from on-farm propagation for restoration purposes. We are aware that on-farm

propagation may cause unintended selection, which could alter seed traits (Conrady

et al., 2023; Ensslin et al., 2018). In the seeds we used, this effect will be rather

minor because most producers are relatively new in this business and the

propagation does not take place for many generations (*personal communication*).

Using farm-propagated seeds further has the advantage that the maternal plants

grew under optimal conditions and the seeds were harvested when fully ripe. This

reduces the variability in seed quality, a characteristic that has a massive effect on

storage behaviour (Rahman & Ellis, 2019; Sano et al., 2016; Sinniah et al., 1998).

Artificial ageing

To artificially age the seeds, we exposed them to 45° C and 60 % relative humidity in

a climate-controlled cabinet (Rumed ®, Rubarth Apparate GmbH, Laatzen,

Germany). These conditions enhance similar chemical processes that are typical of

natural ageing during long-term storage and accelerate seed deterioration (Merritt et
al., 2014; Probert et al., 2009).

We exposed the seeds to artificial ageing conditions for 0 (control group, fresh
seeds), 1, 5, 9, 15, 20, 30, 40, 57 and 72 days (adjusted after Probert et al. (2009)).
We put 3 replicates of 30 seeds of every accession and ageing duration into open 1.5
mL tubes in the ageing condition. The position of each accession in the ageing
condition was randomised within the ageing duration. Before the ageing treatment,
we manually scarified Fabaceae seeds with sandpaper to allow for moisture to be
imbibed (Baskin, 2003). In total, our experiment contained 188 accessions, each with
10 ageing treatments and 3 replicates, that is 5,640 replicates containing 30 seeds
each, that is 169,200 seeds.

Germination test

For germination tests, we used 6-well plates with a well-diameter of 39 mm (VWR
International GmbH, Darmstadt, Germany, catalogue number: 734-2323), because
they use space in the climate cabinet more efficiently than single petri dishes. For
species with large seeds (*Bupleurum rotundifolium*, *Centaurea cyanus*, *Bromus*
hordeaceus, *Arrhenatherum elatius*, *Lathyrus nissolia*, *Hordeum murinum*,
Hypochaeris radicata, *Agrostemma githago*, *Poa annua*) we used petri dishes of 6
cm diameter to avoid overcrowding. We wrapped Petri dishes with seeds of *Bromus*
hordeaceus in aluminium foil since light slightly inhibits germination (Andersson et al.,
2002). After the intended ageing duration, we removed the seeds from the ageing
conditions and transferred each replicate into one well or a petri dish laid out with two
filter papers in a fully randomised design. We watered the seeds with a 0.025 %
Gibberellic acid solution to break dormancy and closed the plates with transparent
Parafilm® to avoid evaporation. Germination took place in a climate-controlled
cabinet (Rumed®, Rubarth Apparate GmbH, Laatzen, Germany) with a 14 h daily
photoperiod at 20° C and 10 h darkness at 10° C.

We checked for germination every seven days and added distilled water if necessary.
When a radicle was longer than 2 mm, we considered a seed viable and removed it
from the dish. For each replicate with viable seeds, we finished the trial when no

more seeds germinated for at least four weeks. If in a replicate no seeds germinated

at all for ten weeks, we considered the seeds not viable. If samples were affected by

mould, we added 1 ml of a 0.0835 % Difenoconazol solution.

Seed traits and climatic data

For each accession, we determined seed weight by weighing 2 replicates of 50 seeds

and calculated the weight of a thousand seeds (further called seed weight for

simplicity). We also obtained the mean annual temperature and precipitation for each

accession location from WorldClim2 using the R package *raster* (Fick & Hijmans,

2017). In wild-collected seeds, we used the exact location of the seed collection. For

farm-propagated seeds, we used the location of the farm, as the initial seeds used to

establish the seed orchard was typically collected in the same region where the seed

producer is located (Bucharova et al., 2019, Figure 1). The oil and protein content of

the seeds was obtained from the Seed Information Database (SID) for each species

(Society for Ecological Restoration et al., 2023).

Statistical analyses

First, we calculated seed longevity for each accession as the number of days in seed

ageing needed to reduce the seed viability to 50 % of the initial viability. The

germination rate was sometimes higher in the short-duration ageing treatments (one

or five days) than in fresh seeds because short exposition to ageing treatment breaks

dormancy in some species (Hay et al., 2021; Mira et al., 2019). We thus defined

"initial viability" for each accession as the highest average germination rate of the 0, 1

or 5 days in the ageing treatment. We excluded accessions with poor initial viabilities

(below 0.33) and then fitted seed-survival curves for each remaining accession using

the *drc*-package for dose-response curves (Ritz et al., 2015). We selected two-

parameter log-logistic models based on the AIC criteria. To estimate the longevity

parameter p_{50}, we used the ED function of the *drc*-package. P_{50} indicates the time

after which the germination rate decreases to half compared to the initial viability and

we call p_{50} "seed longevity" in further text.

The ageing treatment took 72 days, and much longer seed longevity estimates are
strong extrapolations and therefore unreliable. For the following analyses, we thus
excluded 12 accessions with a p_{50}-value of more than 100 days. P_{50} estimates of
most *Papaver rhoeas* accessions were higher than 100, so after excluding them the
species' average longevity would have been underestimated. We therefore removed
this species from further analyses.

Second, we compared the seed longevity between species. For this purpose, we
averaged p_{50}-values across all the accessions of the same species. We then related
the log-transformed average seed longevity as response variable to the species'
mean initial viability, mean seed weight, and plant family as explanatory variables in a
multiple linear regression model. As protein and oil contents were only available for
16 study species, we evaluated the effect of these in a separate multiple linear
regression model, correcting for the initial viability.

Third, we analysed within-species variability in seed longevity. We included only
species for which we had at least 4 accessions. We related accession-specific
longevity (log-transformed p_{50}) as response variable to the species' identity (to
account for it), the accession-specific initial viability, accession-specific seed weight,
mean annual temperature and precipitation at the accession origin as explanatory
variables in a multiple linear regression model. We scaled the seed weight within
each species to standardise the variable across species. To understand the relative
importance of the predictors for seed longevity across accessions, we fitted a multiple
linear regression model containing only significant predictors of within-species
variability, and determined the proportion of variance explained by each of them
using the package *relaimpo* (Groemping & Lehrkamp, 2023).

All data analyses were performed using R version 2023.9.1.494 (R Development
Core Team, 2024). We log-transformed the p_{50} as response variable to ensure that
model assumptions were met. For all models, we verified that model assumptions
were met by graphically evaluating the residuals (Zuur et al., 2010).

RESULTS

The estimated seed longevity, that is the time until half of the viable seeds have died

(p_{50}), strongly varied among species (Figure 2). The average species seed longevity

was shortest for *Bupleurum rotundifolium* (p_{50} = 5.39 days) and longest for *Achillea*

millefolium (p_{50} = 62.88 days). The average species seed longevity significantly

increased with mean initial viability but was unrelated to the species-specific seed

weight (**Error! Reference source not found.**Table 2A, Figure 3), or the seed oil and

protein content (Table 2B, Figure 3). We also did not detect significant differences

between plant families (Table 2A, Figure 3).

The estimated seed longevity also strongly varied among accessions of the very

same species. This variation was the highest in *Daucus carota*, in which the p_{50}

varied more than 18-fold (Figure 2). In absolute values, the estimated seed longevity

varied most in *Dianthus deltoides*, with 60.28 days difference between the longest

and shortest-lived accession. After accounting for species identity, the accession-

specific seed longevity increased with the initial viability of the given seed accession

and mean annual temperature at the accession origin, but was unrelated to the

annual precipitation and accession-specific seed weight (Table 3, Figure 4).

Species identity was the strongest predictor of accession-specific seed longevity,

explaining 51.9% of the variation, followed by the accessions' initial viability (11.5%),

and the mean annual temperature at the seed origin (2.3%). A total of 34.2% of the

variance in seed longevity among accessions remained unexplained.

DISCUSSION

Effective conservation of seeds in ex-situ seed banks requires understanding of seed

storage behaviour not only for individual species but also for accessions within the

very same species. We found out that for multiple species common in European

grasslands, species identity explained only 51.9% of the among-accessions

variability in seed longevity. The residual variance was partly explained by initial seed

viability and mean annual temperature at the accession origin, yet 34.2% of the

among-accession variance remained unexplained. Across species, the seed

longevity was predicted by initial seed viability only. Our findings suggest that seed

longevity is hard to predict – both within and across species

Interspecific variability in seed longevity

In a comparison across species means, the only variable that predicted seed

longevity in ageing was the initial seed viability, where species with more viable

seeds survived longer in ageing. Lower initial viability might indicate that the seeds

have already experienced some deterioration and have reached a point at the

sigmoidal-shaped survival curve, where viability declines more rapidly, and p_{50} is

lower (Hay et al., 2022). Such previous deterioration is naturally more likely in

species with fast-deteriorating seeds. Alternatively, the seeds could have received

suboptimal post-harvest treatment or have low quality from the point of collection, for

example when the seeds were not fully ripe and consequently, had lower germination

and shorter longevity in storage (Hay & Probert, 1995; Probert et al., 2007). Both

previous deterioration and suboptimal post-harvest treatment are naturally more likely

in species whose seeds are more vulnerable to ageing or environmental conditions

during storage. Variability in initial viability might thus amplify the interspecific

difference in seed longevity in artificial ageing. Our findings support current

guidelines for conservation seed banks which recommend that conservation seed

banks preferably select and store seeds with high initial viability (ENSCONET, 2009).

Seed longevity in artificial ageing was affected neither by seed weight nor by the

chemical composition of the seeds. In general, seed weight seems to be an

unreliable predictor of seed longevity in storage, as the effects reported in previous

literature strongly vary from negative, over no effect to, rarely, positive (Davies et al.,

2020; Merritt et al., 2014; Pritchard & Dickie, 2003; Probert et al., 2009; Satyanti et

al., 2018). Similarly, the chemical composition of the seeds did not predict seed

longevity, which corresponds to previous empirical studies of wild species (Probert et

al., 2009, Walters et al 2005). The expected effect of chemical composition on seed

longevity in storage, especially oil content, is based on theoretical expectations and

comparisons among oil crops, cereals, and legumes (Narayana Murthy & Sun, 2000;

Pritchard & Dickie, 2003 Nagel et al., 2015), but the empirical evidence provided by

others and us shows that it is likely not true for wild species.

Surprisingly, we did not detect significant differences between plant families, which

contrasts with nearly all previous studies (Merritt et al., 2014; Mondoni et al., 2011;

Probert et al., 2009; Walters et al., 2005b). This is likely caused by the low number of

species per family (1-6), which resulted in a low power of the statistical tests.

Nevertheless, we observed similar trends as previous studies: Fabaceae seeds were

relatively long-lived, probably due to their physical dormancy (Merritt et al., 2014),

Apiaceae were shorter-lived and there was large variability within Asteraceae (Merritt

et al., 2014; Walters et al., 2005a).

Intraspecific variability in seed longevity

We detected substantial within-species variability in seed longevity in storage.

Specifically, species identity explained only 51.9% of the variation among the

accessions, leaving 48.1% to variation within species. Such high intraspecific

variability is in line with previous works. On a continental scale, Kochanek et al.

(2009) collected data on seed longevity in seven species represented by 2-8

populations, with populations within species largely aggregated in a specific biome.

Here, the species identity explained 70% of the variability among accessions, leaving

30% to variation among accessions within species (own analysis of the data

presented in Kochanek et al. (2009)). On a regional scale, Mondoni et al. (2011)

examined seed longevity of six ecologically similar species that were each

represented by two populations in contrasting environments, specifically high and low

altitude. Here, species identity explained only 6% of variability, leaving astonishing

94% to variation within species (own analysis of the data presented in Mondoni et al.

(2011). In our study, we detected that species explained about half of the variability in

seed longevity among accessions, yet this number might be underestimated because

we excluded the species with the most long-lived seeds for methodical reasons (see

methods). Nevertheless, the available evidence suggests that intraspecific variation

in seed longevity is nearly as substantial as between-species variability, at least in

species coming from the same type of habitat.

The intraspecific variation in seed longevity was predicted by initial seed viability and

climate in seed origin, specifically mean annual temperature. These variables

explained comparatively less variability in seed longevity than species identity,

specifically 11.5% and 2.3%, respectively. The effect of initial seed viability is in line

with the cross-species comparison, and is discussed above. The seed longevity of

the accession increased with the mean annual temperature at the collection site, a

pattern that was reported previously. Populations experiencing warmer climates are

more long-lived than seeds from populations experiencing colder climates (Mondoni

et al., 2011). Other studies attributed such a positive correlation between temperature

and seed longevity mainly to the species identity, because they had species

represented by single accessions (Merritt et al., 2014; Probert et al., 2009). We did

not find any effect of mean annual precipitation on seed longevity, which contrasts

with previous studies both within and across species (Merritt et al., 2014; Probert et

al., 2009; White et al., 2023). This is likely because the mean annual precipitation

was correlated with mean annual temperature (r=33), and as we used multivariate

models, we were not able to tear the effect of both climate variables apart.

The intraspecific variability in seed longevity might be driven by two, mutually non-

exclusive mechanisms – phenotypic plasticity, or genetic differentiation. For example,

Mondoni et al. (2011) detected that seeds collected in populations in warmer

environments survived longer in artificial ageing than conspecific seeds collected in

cooler environments. A large proportion of this effect was caused by phenotypic

plasticity, but the pattern was still significant, although weaker, in the second

generation grown in a common environment, which indicates that the effect is

partially heritable and thus genetic (Mondoni et al., 2014). Indeed, genes associated

with seed longevity in storage were detected in some crops and model species

(Bizouerne et al., 2023; Nagel et al., 2015; Raquid et al., 2021; Renard et al., 2020).

Our study worked with seeds that were collected in natural sites or produced in the
region of origin; we thus cannot discriminate whether the observed effects are plastic
or genetic.

CONCLUSION

Our results show that seed longevity in storage varies within species nearly as much
as between species. Seed samples from different populations of the same species
survive longer in storage if they have high initial seed viability and come from warmer
climates, but there is substantial residual variability. This finding highlights that seed
longevity remains challenging to predict. Conservation seed banks sometimes design
their monitoring of seed viability based on species identity, with the assumption that
accessions of the same species lose viability at the same rate when stored under the
same conditions. Our data support existing concerns that this method might be
suboptimal (Hay & Whitehouse, 2017).

References

Andersson, L., Milberg, P., Schütz, W., & Steinmetz, O. (2002). Germination
characteristics and emergence time of annual Bromus species of differing
weediness in Sweden. *Weed Research*, *42*(2), 135–147.
https://doi.org/10.1046/j.1365-3180.2002.00269.x

Baskin, C. C. (2003). Breaking physical dormancy in seeds – focussing on the lens.
New Phytologist, *158*(2), 229–232. https://doi.org/10.1046/j.1469-
8137.2003.00751.x

Bizouerne, E., Ly Vu, J., Ly Vu, B., Diouf, I., Bitton, F., Causse, M., Verdier, J.,
Buitink, J., & Leprince, O. (2023). Genetic Variability in Seed Longevity and
Germination Traits in a Tomato MAGIC Population in Contrasting Environments.
Plants (Basel, Switzerland), *12*(20), 3632. https://doi.org/10.3390/plants12203632

Bucharova, A., Bossdorf, O., Hölzel, N., Kollmann, J., Prasse, R., & Durka, W.
(2019). Mix and match: Regional admixture provenancing strikes a balance among
different seed-sourcing strategies for ecological restoration. *Conservation
Genetics*, *20*(1), 7–17. https://doi.org/10.1007/s10592-018-1067-6

Cochrane, J. A., Crawford, A. D., Monks, L. T., Cochrane, J. A., Crawford, A. D., &
Monks, L. T. (2007). The significance of ex situ seed conservation to reintroduction
of threatened plants. *Australian Journal of Botany*, *55*(3), 356–361.
https://doi.org/10.1071/BT06173

Cohen, J. I., Williams, J. T., Plucknett, D. L., & Shands, H. (1991). Ex Situ
Conservation of Plant Genetic Resources: Global Development and Environmental
Concerns. *Science*, *253*(5022), 866–872.
https://doi.org/10.1126/science.253.5022.866

Conrady, M., Lampei, C., Bossdorf, O., Hölzel, N., Michalski, S., Durka, W., &
Bucharova, A. (2023). Plants cultivated for ecosystem restoration can evolve
toward a domestication syndrome. *Proceedings of the National Academy of
Sciences*, *120*(20), e2219664120. https://doi.org/10.1073/pnas.2219664120

Davies, R. M., Hudson, A. R., Dickie, J. B., Cook, C., O'Hara, T., & Trivedi, C. (2020).
Exploring seed longevity of UK native trees: Implications for *ex situ* conservation.
Seed Science Research, *30*(2), 101–111.
https://doi.org/10.1017/S0960258520000215

Delouche, J., & Baskin, C. (2021). Accelerated Aging Techniques for Predicting the
Relative Storability of Seed Lots. *All Articles*.
https://scholarsjunction.msstate.edu/seedtechpapers/10

ENSCONET. (2009, June 16). *ENSCONET Curation Protocols & Recommendations*.
http://ensconet.maich.gr/PDF/Curation_protocol_English.pdf

Ensslin, A., Van de Vyver, A., Vanderborght, T., & Godefroid, S. (2018). Ex situ
cultivation entails high risk of seed dormancy loss on short-lived wild plant species.
Journal of Applied Ecology, *55*(3), 1145–1154. https://doi.org/10.1111/1365-
2664.13057

Etterson, J. R., Franks, S. J., Mazer, S. J., Shaw, R. G., Gorden, N. L. S., Schneider,
H. E., Weber, J. J., Winkler, K. J., & Weis, A. E. (2016). Project Baseline: An
unprecedented resource to study plant evolution across space and time. *American
Journal of Botany*, *103*(1), 164–173. https://doi.org/10.3732/ajb.1500313

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: New 1-km spatial resolution climate
surfaces for global land areas. *International Journal of Climatology*, *37*(12), 4302–
4315. https://doi.org/10.1002/joc.5086

Franks, S. J., Hamann, E., & Weis, A. E. (2018). Using the resurrection approach to
understand contemporary evolution in changing environments. *Evolutionary
Applications*, *11*(1), 17–28. https://doi.org/10.1111/eva.12528

Franks, S. J., Sekor, M. R., Davey, S., & Weis, A. E. (2019). Artificial seed aging
reveals the invisible fraction: Implications for evolution experiments using the
resurrection approach. *Evolutionary Ecology*, *33*(6), 811–824.
https://doi.org/10.1007/s10682-019-10007-2

Genna, N. G., Walters, C., & Pérez, H. E. (2020). Viability and vigour loss during
storage of Rudbeckia mollis seeds having different mass: An intra-specific
perspective. *Seed Science Research*, *30*(2), 122–132.
https://doi.org/10.1017/S0960258520000161

Groemping, U., & Lehrkamp, M. (2023). *relaimpo: Relative Importance of Regressors
in Linear Models* (Version 2.2-7) [Computer software]. https://cran.r-
project.org/web/packages/relaimpo/index.html

Guzzon, F., Gianella, M., Velazquez Juarez, J. A., Sanchez Cano, C., & Costich, D.
E. (2021). Seed longevity of maize conserved under germplasm bank conditions

for up to 60 years. *Annals of Botany*, *127*(6), 775–785.
https://doi.org/10.1093/aob/mcab009

Guzzon, F., Orsenigo, S., Gianella, M., Müller, J. V., Vagge, I., Rossi, G., & Mondoni,
A. (2018). Seed heteromorphy influences seed longevity in Aegilops. *Seed
Science Research*, *28*(4), 277–285. https://doi.org/10.1017/S096025851800034X

Hay, F. R., Davies, R. M., Dickie, J. B., Merritt, D. J., & Wolkis, D. M. (2022). More on
seed longevity phenotyping. *Seed Science Research*, *32*(3), 144–149.
https://doi.org/10.1017/S0960258522000034

Hay, F. R., & Probert, R. J. (1995). Seed Maturity and the Effects of Different Drying
Conditions on Desiccation Tolerance and Seed Longevity in Foxglove (Digitalis
purpurea L.). *Annals of Botany*, *76*(6), 639–647.
https://doi.org/10.1006/anbo.1995.1142

Hay, F. R., & Probert, R. J. (2013). Advances in seed conservation of wild plant
species: A review of recent research. *Conservation Physiology*, *1*(1), cot030–
cot030. https://doi.org/10.1093/conphys/cot030

Hay, F. R., & Whitehouse, K. J. (2017). Rethinking the approach to viability
monitoring in seed genebanks. *Conservation Physiology*, *5*(1), cox009.
https://doi.org/10.1093/conphys/cox009

Hay, F. R., Whitehouse, K. J., Ellis, R. H., Sackville Hamilton, N. R., Lusty, C.,
Ndjiondjop, M. N., Tia, D., Wenzl, P., Santos, L. G., Yazbek, M., Azevedo, V. C.
R., Peerzada, O. H., Abberton, M., Oyatomi, O., de Guzman, F., Capilit, G.,
Muchugi, A., & Kinyanjui, Z. (2021). CGIAR genebank viability data reveal
inconsistencies in seed collection management. *Global Food Security*, *30*, 100557.
https://doi.org/10.1016/j.gfs.2021.100557

Kochanek, J., Buckley, Y. M., Probert, R. J., Adkins, S. W., & Steadman, K. J. (2010).
Pre-zygotic parental environment modulates seed longevity. *Austral Ecology*,
35(7), 837–848. https://doi.org/10.1111/j.1442-9993.2010.02118.x

Kochanek, J., Steadman, K. J., Probert, R. J., & Adkins, S. W. (2009). Variation in
seed longevity among different populations, species and genera found in
collections from wild Australian plants. *Australian Journal of Botany*, *57*(2), 123–
131. https://doi.org/10.1071/BT09023

Lee, J.-S., Velasco-Punzalan, M., Pacleb, M., Valdez, R., Kretzschmar, T., McNally,
K. L., Ismail, A. M., Cruz, P. C. S., Sackville Hamilton, N. R., & Hay, F. R. (2019).
Variation in seed longevity among diverse Indica rice varieties. *Annals of Botany*,
124(3), 447–460. https://doi.org/10.1093/aob/mcz093

Merritt, D. J., & Dixon, K. W. (2011). Restoration Seed Banks—A Matter of Scale.
Science, *332*(6028), 424–425. https://doi.org/10.1126/science.1203083

Merritt, D. J., Martyn, A. J., Ainsley, P., Young, R. E., Seed, L. U., Thorpe, M., Hay,
F. R., Commander, L. E., Shackelford, N., Offord, C. A., Dixon, K. W., & Probert,
R. J. (2014). A continental-scale study of seed lifespan in experimental storage
examining seed, plant, and environmental traits associated with longevity.
Biodiversity and Conservation, *23*(5), 1081–1104. https://doi.org/10.1007/s10531-
014-0641-6

Millenium Seed Bank. (2022). *Assessing a Population for Seed Collection*.
https://brahmsonline.kew.org/Content/Projects/msbp/resources/Training/02-
Assessing-population.pdf

Mira, S., Veiga-Barbosa, L., & Pérez-García, F. (2019). Seed dormancy and longevity
variability of *Hirschfeldia incana* L. during storage. *Seed Science Research*, *29*(2),
97–103. https://doi.org/10.1017/S0960258519000072

Mondoni, A., Orsenigo, S., Donà, M., Balestrazzi, A., Probert, R. J., Hay, F. R.,
Petraglia, A., & Abeli, T. (2014). Environmentally induced transgenerational
changes in seed longevity: Maternal and genetic influence. *Annals of Botany*,
113(7), 1257–1263. https://doi.org/10.1093/aob/mcu046

Mondoni, A., Probert, R. J., Rossi, G., Vegini, E., & Hay, F. R. (2011). Seeds of
alpine plants are short lived: Implications for long-term conservation. *Annals of*
Botany, *107*(1), 171–179. https://doi.org/10.1093/aob/mcq222

Nagel, M., Kranner, I., Neumann, K., Rolletschek, H., Seal, C. E., Colville, L.,
Fernández-Marín, B., & Börner, A. (2015). Genome-wide association mapping and
biochemical markers reveal that seed ageing and longevity are intricately affected
by genetic background and developmental and environmental conditions in barley.
Plant, Cell & Environment, *38*(6), 1011–1022. https://doi.org/10.1111/pce.12474

Narayana Murthy, U. M., & Sun, W. Q. (2000). Protein modification by Amadori and
Maillard reactions during seed storage: Roles of sugar hydrolysis and lipid
peroxidation. *Journal of Experimental Botany*, *51*(348), 1221–1228.
https://doi.org/10.1093/jexbot/51.348.1221

Niedzielski, M., Walters, C., Luczak, W., Hill, L. M., Wheeler, L. J., & Puchalski, J.
(2009). Assessment of variation in seed longevity within rye, wheat and the
intergeneric hybrid triticale. *Seed Science Research*, *19*(4), 213–224.
https://doi.org/10.1017/S0960258509990110

Ninoles, R., Planes, D., Arjona, P., Ruiz-Pastor, C., Chazarra, R., Renard, J., Bueso,
E., Forment, J., Serrano, R., Kranner, I., Roach, T., & Gadea, J. (2022).
Comparative analysis of wild-type accessions reveals novel determinants of
Arabidopsis seed longevity. *PLANT CELL AND ENVIRONMENT*, *45*(9), 2708–
2728. https://doi.org/10.1111/pce.14374

Peres, S. (2016). Saving the gene pool for the future: Seed banks as archives.
Studies in History and Philosophy of Science Part C: Studies in History and
Philosophy of Biological and Biomedical Sciences, *55*, 96–104.
https://doi.org/10.1016/j.shpsc.2015.09.002

Pritchard, H., & Dickie, J. (2003). *Predicting Seed Longevity: The use and abuse of*
seed viability equations (pp. 653–721).

Probert, R. J., Adams, J., Coneybeer, J., Crawford, A., & Hay, F. (2007). Seed quality
for conservation is critically affected by pre-storage factors. *Australian Journal of*
Botany, *55*(3), 326–335. https://doi.org/10.1071/BT06046

Probert, R. J., Daws, M. I., & Hay, F. R. (2009). Ecological correlates of ex situ seed
longevity: A comparative study on 195 species. *Annals of Botany*, *104*(1), 57–69.
https://doi.org/10.1093/aob/mcp082

Rahman, S. M. A., & Ellis, R. H. (2019). Seed quality in rice is most sensitive to
drought and high temperature in early seed development. *Seed Science*
Research, *29*(4), 238–249. https://doi.org/10.1017/S0960258519000217

Raquid, R., Kohli, A., Reinke, R., Dionisio-Sese, M., Kwak, J., Chebotarov, D., Mo,
Y., & Lee, J.-S. (2021). Genetic factors enhancing seed longevity in tropical
japonica rice. *Current Plant Biology*, *26*, 100196.
https://doi.org/10.1016/j.cpb.2021.100196

Renard, J., Niñoles, R., Martínez‐Almonacid, I., Gayubas, B., Mateos‐Fernández, R.,
Bissoli, G., Bueso, E., Serrano, R., & Gadea, J. (2020). Identification of novel seed
longevity genes related to oxidative stress and seed coat by genome‐wide
association studies and reverse genetics. *Plant, Cell & Environment*, *43*(10),
2523–2539. https://doi.org/10.1111/pce.13822

Ritz, C., Baty, F., Streibig, J. C., & Gerhard, D. (2015). Dose-Response Analysis
Using R. *PLOS ONE*, *10*(12), e0146021.
https://doi.org/10.1371/journal.pone.0146021

Sano, N., Rajjou, L., North, H. M., Debeaujon, I., Marion-Poll, A., & Seo, M. (2016).
Staying Alive: Molecular Aspects of Seed Longevity. *Plant and Cell Physiology*,
57(4), 660–674. https://doi.org/10.1093/pcp/pcv186

Satyanti, A., Nicotra, A. B., Merkling, T., & Guja, L. K. (2018). Seed mass and
elevation explain variation in seed longevity of Australian alpine species. *Seed*
Science Research, *28*(4), 319–331. https://doi.org/10.1017/S0960258518000090

Schutte, B. J., Regnier, E. E., & Harrison, S. K. (2008). The association between
seed size and seed longevity among maternal families in Ambrosia trifida L.
populations. *Seed Science Research*, *18*(4), 201–211.
https://doi.org/10.1017/S0960258508082974

Sinniah, U. R., Ellis, R. H., & John, P. (1998). Irrigation and Seed Quality
Development in Rapid-cycling Brassica: Seed Germination and Longevity. *Annals*
of Botany, *82*(3), 309–314. https://doi.org/10.1006/anbo.1998.0748

Society for Ecological Restoration, International Network for Seed Based Restoration,
& Royal Botanical Gardens Kew. (2023, February). *Seed Information Database*
(SID). https://ser-sid.org/

Walters, C., Wheeler, L. M., & Grotenhuis, J. M. (2005a). Longevity of seeds stored
in a genebank: Species characteristics. *Seed Science Research*, *15*(1), 1–20.
https://doi.org/10.1079/SSR2004195

Walters, C., Wheeler, L. M., & Grotenhuis, J. M. (2005b). Longevity of seeds stored
in a genebank: Species characteristics. *Seed Science Research*, *15*(1), 1–20.
https://doi.org/10.1079/SSR2004195

Wambugu, P. W., Nyamongo, D. O., & Kirwa, E. C. (2023). Role of Seed Banks in
Supporting Ecosystem and Biodiversity Conservation and Restoration. *Diversity*,
15(8), Article 8. https://doi.org/10.3390/d15080896

White, F. J., Hay, F. R., Abeli, T., & Mondoni, A. (2023). Two decades of climate
change alters seed longevity in an alpine herb: Implications for ex situ seed
conservation. *Alpine Botany*, *133*(1), 11–20. https://doi.org/10.1007/s00035-022-
00289-8

Zani, D., & Müller, J. V. (2017). Climatic control of seed longevity of *Silene* during the
post‑zygotic phase: Do seeds from warm, dry climates possess higher maturity
and desiccation tolerance than seeds from cold, wet climates? *Ecological*
Research, *32*(6), 983–994. https://doi.org/10.1007/s11284-017-1508-6

Zuur, A. F., Ieno, E. N., & Elphick, C. S. (2010). A protocol for data exploration to
avoid common statistical problems. *Methods in Ecology and Evolution*, *1*(1), 3–14.
https://doi.org/10.1111/j.2041-210X.2009.00001.x

Figure 1: Origins of the 153 seed accessions used in the study and included in the

analyses. Each point represents the location of a seed producer as a proxy for the

origins of the individual seed accessions and thus, is the origin of multiple species. In

three cases, the points represent the location of own seed collection.

Figure 2: (a) Estimated seed longevity (p_{50}) across 35 species and 153 accessions.

Blue points represent species averages, green points accession estimates. (b)

Proportion of explained variability in seed longevity by significant predictors and

residuals (this analysis included only species represented by at least four

accessions).

Figure 3: Relationship between average seed longevity (p_{50}) of each species and
species-specific seed traits. Each point represents one species. (a) Initial seed
viability, average across accessions of the same species; (b) seed mass, average
across accessions of the same species; species-specific (c) protein and (d) oil
content in the seeds; and (e) plant families. Dashed line indicates non-significant
relationship. For full model results, see Table 2.

624
625

Figure 4: Relationship between seed longevity (p_{50}) of each accession, accession-
specific seed traits, and environmental variables. Each point represents a seed
accession. Seed longevity was adjusted for species identity. (a) Accession-specific
initial seed viability; (b) accession-specific seed mass (scaled); (c) mean annual
precipitation and (d) mean annual temperature at the origins of the accessions. For
full model results see Table 3.

Table 1: Species included in the study. The number of accessions is the number of

accessions used for the analyses and in brackets the initial number of accessions.

Seed weight is the weight of 1000 seeds, averaged across all accessions of the

same species. Oil and protein content are obtained from the Seed Information

Database (Society for Ecological Restoration et al., 2023). Life cycle: A = annual, P =

perennial, B = biennial. *: Information of conspecific

Species	Number of accessions [initial number]	Seed weight [g]	Family	Life cycle	Oil content [%]	Protein content [%]
Achillea millefolium	8 [10]	0.14	Asteraceae	P	27.03	33.1
Agrostemma githago	2 [2]	10.02	Caryophyllaceae	A	6.8	15.3
Ajuga reptans	0 [4]	1	Lamiaceae	P		
Anthemis arvensis	1 [1]	0.37	Asteraceae	A/B		
Arrhenatherum elatius	4 [5]	2.83	Poaceae	P	7.13	24.4
Barbarea vulgaris	4 [4]	0.39	Brassicaceae	A/B	31	15.6
Bromus hordeaceus	3 [3]	4.38	Poaceae	A		
Bupleurum rotundifolium	1 [1]	2.63	Apiaceae	A		
Capsella bursa-pastoris	2 [2]	0.15	Brassicaceae	A	30.53	28.23
Cardamine hirsuta	2 [2]	0.12	Brassicaceae	A		
Centaurea cyanus	3 [7]	3.94	Asteraceae	A	24.67	16.57
Centaurea jacea	8 [8]	1.72	Asteraceae	P		
Cerastium holosteoides	1 [1]	0.07	Caryophyllaceae	B/P		
Chelidonium majus	3 [5]	0.71	Papaveraceae	B		
Daucus carota	9 [9]	0.83	Apiaceae	A/B	19.26	24.38
Dianthus deltoides	7 [8]	0.2	Caryophyllaceae	P		
Galium wirtgenii	1 [1]	0.63	Rubiaceae	P		
Geranium pusillum	0 [1]	1	Geraniaceae	A		
Hordeum murinum	1 [1]	15.41	Poaceae	A	1.4	23.5
Hypochaeris radicata	6 [7]	0.74	Asteraceae	P		
Lamium album	0 [2]	1.57	Lamiaceae	P		
Lamium purpureum	0 [3]	0.69	Lamiaceae	A		
Lathyrus nissolia	1 [1]	7.13	Fabaceae	A		
Linaria vulgaris	7 [7]	0.15	Plantaceae	P		
Lotus corniculatus	6 [7]	1.33	Fabaceae	P	6.53	36.18
Lotus pedunculatus	0 [2]	0.52	Fabaceae	P		
Lychnis flos-cuculi	8 [8]	0.16	Caryophyllaceae	P		
Medicago lupulina	6 [6]	1.33	Fabaceae	A	5.45	35.57
Papaver rhoeas	0 [9]	0.1	Papaveraceae	A	43.47	24.53
Plantago lanceolata	11 [11]	1.49	Plantaginaceae	P	6.73	17.47

Plantago media	6 [6]	0.34	Plantaginaceae	P(/B/A)		
Poa annua	2 [2]	0.44	Poaceae	A/B		
Prunella vulgaris	9 [9]	0.77	Lamiaceae	P	21.7	20.2
Scandix pecten-veneris	0 [1]		Apiaceae	A		
Senecio vulgaris	2 [2]	0.17	Asteraceae	A		
Silene vulgaris	9 [9]	0.77	Caryophyllaceae	P	2.7	18.1
Stellaria media	3 [3]	0.46	Caryophyllaceae	A	5.9	17.8
Thlaspi arvense	2 [3]	1.18	Brassicaceae	A	31.2	24.9
Trifolium aureum	2 [2]	0.42	Fabaceae	A		
Trifolium campestre	3 [5]	0.31	Fabaceae	A/B		
Veronica chamaedrys	8 [8]	0.14	Plantaginaceae	P		

645 Table 2: **Interspecific** variability in seed longevity in storage, results of the linear
646 model testing the effect of (A) initial viability, the family, the species-specific seed
647 weight, and (B) and the protein and oil content on species-specific seed longevity.
648 Analysis of variance with error type III. Significant effects (p < 0.05) are highlighted in
649 bold.

	df	Resid.df	F	p	R^2
A		22			0.4
Initial viability	1		4.78	**0.04**	
Family	9		0.72	0.686	
Species-specific seed weight	1		1.07	0.312	
B		11			0.52
Initial viability	1		7.3	**0.021**	
Protein content	1		4.58	0.056	
Oil content	1		0.15	0.703	

652 Table 3: **Intraspecific** variability in seed longevity in storage, results of the linear
653 model testing the effect of species identity, initial viability, accession-specific seed
654 weight, mean annual temperature and mean annual precipitation on accession-
655 specific seed longevity in storage. Analysis of variance with error type III. Significant
656 effects (P < 0.05) are highlighted in bold.

	df	Resid.df	F	p	R^2
		94			0.67
Species	15		8.5	**< 0.001**	
Initial viability	1		25.6	**< 0.001**	
Accession-specific seed weight	1		0.89	0.348	
Mean annual temperature	1		7.09	**0.009**	
Annual precipitation	1		0.46	0.5	

657 .