Title

The shortest telomere in telomerase-negative cells triggers replicative senescence at a critical threshold length and also fuels genomic instability

Author list

Prisca Berardi[1,#], Veronica Martinez-Fernandez[1,#], Anaïs Rat[2], Fernando R. Rosas Bringas[3], Pascale Jolivet[1], Rachel Langston[1], Stefano Mattarocci[1], Alexandre Maes[1], Théo Aspert[4,5,6,7], Bechara Zeinoun[1], Karine Casier[1], Hinke G. Kazemier[3], Gilles Charvin[4,5,6,7], Marie Doumic[8], Michael Chang[3] and Maria Teresa Teixeira[1,*]

Author affiliations

[1] Sorbonne Université, CNRS, Laboratoire de Biologie Moléculaire et Cellulaire des Eucaryotes, LBMCE, F-75005 Paris, France
[2] Univ Brest, CNRS, UMR 6205, Laboratoire de Mathématiques de Bretagne Atlantique, 29200 Brest, France
[3] European Research Institute for the Biology of Ageing, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands
[4] Department of Developmental Biology and Stem Cells, Institut de Génétique et de Biologie Moléculaire et Cellulaire, Illkirch, France.
[5] CNRS, UMR7104, Illkirch, France.
[6] INSERM, U964, Illkirch, France.
[7] Université de Strasbourg, Illkirch, France.
[8] CMAP, Inria, IP Paris, Ecole polytechnique, CNRS, 91128 Palaiseau cedex, France.
[#] The authors contributed equally (by alphabetical order)
*Correspondence: teresa.teixeira@cnrs.fr

In the absence of telomerase, telomere shortening triggers the DNA damage checkpoint and replicative senescence, a potent tumor suppressor mechanism. Paradoxically, this same process is also associated with oncogenic genomic instability. Yet, the precise mechanism that connects these seemingly opposing forces remains poorly understood. To directly study the complex interplay between senescence, telomere dynamics and genomic instability, we developed a system in *Saccharomyces cerevisiae* to generate and track the dynamics of telomeres of precise length in the absence of telomerase. Using single-telomere and single-cell analyses combined with mathematical modeling, we identify a threshold length at which telomeres switch into dysfunction. A single shortest telomere below the threshold length is necessary and sufficient to trigger the onset of replicative senescence in a majority of cells. At population level, fluctuation assays establish that rare genomic instability arises predominantly in *cis* to the shortest telomere as non-reciprocal translocations that result in re-elongation of the shortest telomere and likely escape from senescence. The switch of the shortest telomere into dysfunction and subsequent processing in telomerase-negative cells thus serves as the mechanistic link between replicative senescence onset, genomic instability and the initiation of post-senescence survival, explaining the contradictory roles of replicative senescence in oncogenesis.

Telomeres, located at the ends of linear chromosomes in eukaryotes, comprise G/C-rich repeated DNA sequences–with the G-rich strand extending to form a 3' protruding end, called G-overhang, as well as specialized proteins and non-coding RNA. The primary function of telomeres is to shield chromosome ends from being recognized as DNA extremities resulting from double strand breaks (DSBs) and prevent DNA repair activities that might compromise genome stability[1,2]. During DNA replication, telomeres naturally undergo shortening in a phenomenon referred to as the DNA-end replication problem[3-5]. Telomerase, a specialized ribonucleoprotein complex, counteracts this shortening by synthesizing telomeric repeats *de novo* using its reverse transcriptase subunit and template RNA[6,7].

Telomerase plays a crucial role in the long-term proliferation of eukaryotic cells. Nonetheless, in many somatic cells of humans and some other species, telomerase activity is repressed. In these cells, as well as in experimental model systems, inactivation of telomerase leads to telomere erosion, eventually causing replicative senescence–a permanent cell cycle arrest triggered by the activation of the DNA-damage checkpoint[8-10]. For this reason, replicative senescence has been considered a potent tumor-suppressor mechanism. However, cancer cells often escape this mechanism through the acquisition of the ability to re-extend telomeres by either re-expressing telomerase or activating pathways collectively termed alternative lengthening of telomeres (ALT)[11,12]. ALT pathways, first observed in *Saccharomyces cerevisiae* cells lacking telomerase, involve homology-dependent repair mechanisms such as break-induced replication (BIR)[13-16]. Despite significant progress, the mechanisms by which short telomeres trigger replicative senescence and by which cells activate ALT pathways remain elusive, in part due to the lack of systems that allow direct testing of hypotheses.

Genomic instability also increases with telomere shortening. Accordingly, rare gross chromosomal rearrangements such as translocations, or more massive genomic catastrophes, have been attributed to presumably dysfunctional short telomeres[17-21]. Thus, when combined with genomic instability, ALT could theoretically enable the long-term proliferation of cells carrying genomic rearrangements that present a threat to organismal long-term survival. However, whether and how genomic instability is mechanistically linked to ALT is unknown. Moreover, the very nature of the initial telomere dysfunction at the origin of the genomic instability is unclear and difficult to assess. Several non-exclusive hypotheses have been put forward, including replication fork collapse when passing through hard-to-replicate telomeric repeated sequences, oxidation of telomeric DNA following mitochondrial dysfunction, and extreme telomere shortening[22,23]. However, none of these hypotheses provide a comprehensive explanation. Also, the study of replicative senescence presents challenges due to inherent variability across telomeres and cellular phenotypes associated with senescence[24]. Telomere

length varies not only between different telomeres within a single cell but also among cells. Additionally, the telomere shortening mechanism hides several layers of complexity, which include the asymmetry of leading and lagging replication machineries contributing to variations in telomere length distributions[4,25]. Variability also stems from significant inter-cellular differences in proliferation potential and gene expression[26]. Furthermore, senescence-specific genomic instability contributes to increased genetic diversity, further complicating the study of the mechanisms involved in senescence[18,19]. Therefore, it is particularly challenging to directly connect a specific molecular event at specific telomere(s) to the onset of senescence.

The heterogeneity observed in telomere biology is recapitulated in the budding yeast *Saccharomyces cerevisiae*, where telomeres consist of degenerated TG_{1-3} repeats spanning approximately 300 ± 75 bp[27,28]. Mutant yeast cells that lack telomerase experience gradual asymmetrical telomere shortening at an average rate of ~3-5 bp per cell division[4,29]. As telomeres shorten, an increasing but highly variable fraction of cells undergo replicative senescence[9,30]. Studies in this organism suggest a model in which telomerase-negative budding yeast cells arrest and activate DNA damage checkpoints as soon as one telomere reaches a critically short length[22,31-33]. Notably, microscopy tracking in microfluidics circuits of consecutive cell divisions over time, from telomerase inactivation to cell death, combined with mathematical modeling, have provided valuable insights into the intricate relationship between telomere shortening and replicative senescence in budding yeast telomerase-negative cells[25,30,34-37]. However, the hypothesis that a single telomere is sufficient to trigger senescence remains controversial and was not possible to be directly tested in a physiological system with native telomeres. In order to understand the mechanisms underlying the onset of replicative senescence and its escape, it is thus crucial to examine the structure and fate of the shortest telomere in budding yeast cells lacking telomerase.

In this work, we present a novel system called *FinalCut*, which enables the generation in budding yeast cells of a single short telomere with a defined length to control telomere length heterogeneity. *FinalCut* targets the ectopic endonuclease Cas9 to a short sequence within the TG_{1-3} repeats of a specific native telomere. Combining structural analysis of the rate of *FinalCut* telomere shortening by single-molecule sequencing, together with the monitoring of single consecutive cell cycles and mathematical modeling, we show that the shortest telomere in cells reaching a critical length threshold is required and sufficient to trigger replicative senescence in a large majority of telomerase-negative cells. Our data also demonstrate that genome instability upon telomerase inactivation occurs primarily in the vicinity of the critically short telomere, while the stability of the rest of the genome does not appear to be compromised. Furthermore, non-reciprocal translocations appear as

a mutational signature associated with the shortest telomere when telomerase is absent. Overall, our findings demonstrate that the shortest telomere in cells is both the primary cause of proliferation limit and the major threat to genome integrity.

The *FinalCut* system enables precise experimental control of telomere length

While telomere length appears as a critical determinant of telomere biology, telomere length variation often masks the understanding of underlying molecular mechanisms. Therefore, the precise experimental control of telomere length is a challenging but essential objective. Experimental systems that aim at modifying telomere length exist, but they exhibit limitations. They often involve alterations of subtelomeric elements, exposure of non-telomeric DNA ends, or lack precision in achieving the desired final telomere length. We thus undertook the development of a new tool we named *FinalCut*. By utilizing an inducible CRISPR/Cas9 approach, the *FinalCut* allows us to selectively truncate a single native telomere to a precise length with temporal control (Fig.1a).

To ensure a controlled Cas9 cleavage of the telomere, we chose the right end of the chromosome 6 (*TEL6R*) to integrate a construct containing a variable number of native 6R telomeric TG_{1-3} repeats, a unique 29 bp ectopic sequence containing a protospacer adjacent motif (PAM) sequence and Cas9 guide RNA (gRNA)-complementary target sequence (ttDNA), and a set of terminal TG_{1-3} repeats. *TEL6R* location was chosen because of the well-characterized nature of this telomere[38,39]. In the integrated context, the variable amount of telomeric sequence before the Cas9 binding site determines the length of the telomere after the Cas9 cut. To enable precise interrogation of the impact of telomere length, we generated a set of strains with *TEL6R* lengths of 0, 21, 31, 41, 51 or 70 bp of the native TG_{1-3} (*FCx*), which we named *FC0*, *FC20*, *FC30*, *FC40*, *FC50*, *FC70*, respectively (*noFC* refers to the original control strain harboring an untouched *TEL6R*).

To temporally control Cas9 cleavage, we use a plasmid-based galactose-inducible Cas9 enzyme. Further control of the telomere was achieved by constructing the *FinalCut* system in strains that placed the telomerase template RNA encoding gene, *TLC1*, under the control of a doxycycline repressible promoter (P_{tetO2}-*TLC1*) so that telomerase could be deactivated as appropriate[4].

We collected DNA samples from different time points of galactose induction of the *FinalCut* strains and analyzed by Southern blots and TeloPCR[40] for detection of the terminal restriction fragment (TRF) of the 6R chromosome end (Fig.1b, Extended Data Fig.1a). At t=0 prior to galactose induction, a smeary signal was identified, showing that the *FinalCut* construct is well maintained by telomerase to a variable length (Fig.1c, Extended Data Fig.1b-d), as described[4,41]. Upon galactose induction, this

signal was converted into a sharp band both in the presence and absence of telomerase activity. This demonstrated efficient cleavage of the *FinalCut* constructs by Cas9 to produce telomeres at 6R of the expected lengths. Detection of all telomeres using a CA_{1-3} probe showed that the *FinalCut* telomeres were the only ones cut in these conditions (Extended Data Fig.1e).

In the presence of telomerase, we observed that in the *FC30-70* strains, the Cas9-shortened 6R telomere fragments were rapidly elongated to longer fragments, whereas the signal in the *FC20* strain vanished (Extended Data Fig.1b). This likely reflects the ability of telomeric sequences above a threshold of ~30 bp to be elongated by telomerase[42], whereas below, they are rapidly degraded. We also observed a relative decrease in the quantity of telomerase-elongated *FC70* telomeres, since telomerase is less frequently recruited to relatively longer telomeres[43]. These observations suggest that the *FinalCut* telomeres are functional with respect to telomerase recruitment.

When telomerase was inactivated prior to Cas9 induction by doxycycline addition to the media to suppress *TLC1* expression, approximately 90% of the uncut smear signal was converted to a sharp cut band within 6-9 hours (Fig.1c-d, and Supplementary Table 6). We also found that the total signal intensity of *FinalCut* telomeres gradually decrease over time, with a more pronounced effect on shorter telomeres compared to longer telomeres. Indeed, while we observed a rapid disappearance of the 6R TRF signal in both *FC20* and *FC30* strains, the signal persisted for *FC40-70* strains, when the cut telomere was 40 bp or longer (Extended Data Fig.1f). Since no other signal at other lengths was detected (Extended Data Fig.1c), we could exclude the possibility of a systematic fusion or recombination of shorter *FinalCut* telomeres with another specific region of the genome. This indicates that the shorter *FinalCut* telomeres of *FC20* and *FC30* strains are likely degraded immediately, whereas the ones of *FC40* and *FC50* are initially stable prior to delayed degradation (Fig.1d, Extended Data Fig.1f). Probing at a more internal restriction fragment, 23 kb away from *TEL6R*, indicates that degradation at *FC20* and *FC30* may extend internally into chromosomal regions containing essential genes (Extended Data Fig.1f). We propose that the shift between telomere function to dysfunction manifests as end degradation progressing toward the interior of the chromosome. The structural nature of this shift may have remained directly undetected likely because it affects only a small fraction of the 32 telomeres in the budding yeast genome, as illustrated in the Extended Data Fig.1g, and a small fraction of cells in a population[30].

Temporal control of *FinalCut* Telomere generation reveals threshold for telomere dysfunction

To further improve the *FinalCut* system to enable evaluation of telomere stability, we decided to temporally restrain the generation of *FinalCut* telomeres before evaluating their stability over time. This eliminates the confounding variable of *FinalCut* telomeres being continuously generated as cells divide and telomeres shorten and/or degrade. In this enhanced system, we provided galactose for 4 hours, a period shown to be sufficient to allow a substantial production of *FinalCut* telomeres, and then changed the carbon source to glucose to repress the *GAL1* promoter driving Cas9 expression (Fig.2a). Additionally, to promote rapid degradation of the remaining Cas9, we used a version of the nuclease fused to the auxin-inducible degron (AID) and induced rapid degradation of Cas9 by addition of the auxin analogue 1-Naphthalenacetic acid (NAA) (Extended Data Fig.2a-b)[44,45].

Intriguingly, analysis of *FinalCut* TRFs across different time points upon Cas9 shut off showed that a telomere of 30 bp is rapidly degraded after the cut and hardly quantifiable (Extended Data Fig.2c-d). In contrast, the signal did not decrease in the case of a 40 bp or a 50 bp telomere until after a few additional rounds of cell divisions. Together with our earlier data suggesting that *FinalCut* telomeres enable telomerase elongation for telomeres longer than 30 bp (Extended Data Fig.1b), and enable protection from degradation in telomeres longer than 40 bp (Fig.1c-d, Extended Data Fig.1f), we speculate that 20-30 bp represents a threshold below which the telomere is targeted for degradation rather than elongation. In contrast, telomeres spanning 40-50 bp are above the deterministic threshold for dysfunction and therefore have a slower degradation profile. Overall, this difference in the length threshold for these two fundamental telomere characteristics (~30 bp for telomerase-elongation and ~40 bp for protection from degradation), suggests that the underlying respective mechanisms are distinct.

As *FC40* and *FC50* cells continuously divide in the above experiment, telomeres are expected to shorten due to the DNA end replication problem prior to dysfunction. According to the current model of telomere shortening[4], one out of two cells should inherit of a shorter telomere by the length of the overhang at each cell division (Extended Data Fig.2e). Therefore, our results would be compatible with half of *FC40* cells inheriting a short telomere below a deterministic threshold length at each population doubling (PD). For *FC50*, these events would be delayed by ~1-2 population doublings.

FinalCut combined with single-molecule sequencing recapitulates the current model of the DNA end replication problem

To test the idea that *FinalCut* telomeres undergo gradual shortening prior to dysfunction as proposed above, we next precisely traced their shortening at nucleotide resolution using single-molecule sequencing. The DNA end replication problem has traditionally been studied in telomere populations of varying lengths where the exact shortening path is difficult to identify. Examining telomere shortening in a context where we can generate telomeres of exact lengths therefore presents a unique opportunity to directly address the phenomenon.

We first examined when the *FinalCut* telomere acquired G-overhangs following Cas9 cleavage, by determining when they are recognized as authentic telomerase substrates in cells. This recognition requires the regeneration of a 3' overhang from the blunt ends created by Cas9 endonucleolytic cleavage[46]. We thus monitored the sequence of *FC50* telomeres over time in the presence of telomerase and developed a dedicated protocol to ensure read depth on individual telomeric sequences and subsequent accuracy (Extended Data Fig.2f). Inspection of *FC50* sequences at t=0, prior to Cas9 induction, showed that the majority of telomeres contain the *FinalCut* construct and display variable lengths, as expected (Extended Data Fig.2g). From centromere to chromosome end, they are composed of a common 6R subtelomeric sequence, followed by 50 bp of degenerated TG_{1-3} repeats (*FC50*), the Cas9 target ttDNA, a common stretch of degenerated TG_{1-3} repeats, followed by TG_{1-3} of variable length and divergent sequence among the telomere molecules. This profile is consistent with the expected composition of telomeres in budding yeast. The core region consists of a common TG_{1-3} sequence that is primarily propagated by the semi-conservative DNA replication machinery. The distal region, however, exhibited sequence divergence among the progeny due to independent, degenerated telomerase-mediated re-elongations of shorter telomeres, as part of the mechanism of telomere length maintenance[40,43].

When cells are incubated in the presence of Cas9 for 4 hours and then grown for an additional 6 hours in glucose, nearly half of the telomeres displayed sequence divergence starting near the expected Cas9 cleavage site, reflecting *de novo* telomere re-elongations (Extended Data Fig.2h). These are likely telomerase-dependent as they are detected in telomerase-positive conditions but not when telomerase is repressed, and fall in the range of telomerase elongations (Extended Data Fig.2i). We identified position 57 as the most common substrate for telomerase, corresponding to the predicted last nucleotide left by Cas9 double-stranded cleavage (Extended Data Fig.2j). However, a substantial number of telomerase-dependent elongations were also detected at position 58. This

is presumably due to the variability in Cas9 cleavage, which can create a 1-nt 5′-overhang that is filled by Pol4, as described for some *loci* in various species[47,48]. We concluded that our ttDNA drives staggered cleavage by Cas9, resulting in either a double-stranded blunt end or a 5′-overhang intermediate. Importantly, the data show that all these ends at positions 57 and 58 function as genuine substrates for telomerase, indicating that Cas9-generated ends are likely converted into single-stranded 3′ ends suitable for telomere elongation. Other positions, namely 51 and 46, were also found to share a last common nucleotide, indicating that a telomere shortening event subsequent to Cas9 cleavage must have occurred prior to telomerase elongation. When analyzing the presumed substrates for telomerase elongations, we see that 5′-TCCGGG-3′ is the most common sequence used by telomerase presumably because it is the most frequent extremity available right after Cas9 cleavage (position 57). Accordingly, annealing of this substrate to the 3′-CCC-5′ of the 3′-ACACACACCCACACCAC-5′ template RNA would be followed by the synthesis of 5′-TGTG-3′, which is exactly what we see (Extended Data Fig.2j). However, when it comes to shortened substrates, annealing of telomerase RNA template to 5′-GTG–3′ is the most common situation[49]. In this context, the substrates for telomerase elongation at positions 51 and 45 indicate possible shortening steps of ~6 bp from the original *FinalCut* telomere.

We next examined telomeres in cells lacking telomerase. Sequencing of *FC50* and *FC70* over time shows gradual shortening of telomeres at an average of 4.4 and 3.4 bp/PD, respectively, compatible with previous estimations[29] (Fig.2b). This suggests that the procedure to generate *FinalCut* telomeres does not affect their subsequent behavior in the cell as compared to native telomeres. Examination of sequence lengths at the individual telomere level revealed a non-uniform distribution (Fig.2b-c, Extended Data Fig.2k). For *FC50*, at 0 hours, corresponding to ~1.5 PD since the beginning of galactose induction of Cas9, 51.9% +/-4.7 of telomeres show expected lengths upon Cas9 staggered cleavage at positions 57 and 58, as above. Strikingly, nearly 23.8% +/-2.2 of *FinalCut* telomeres are shortened by 6 bp and end in position 51. At 2 hours corresponding to ~2.7 PD, the fraction of telomeres ending at positions 57 and 58 decreased by half, and the ones ending in position 51 is maintained at 30.9% +/-1.8 while a fraction of telomeres ending near positions 40 to 45 starts to emerge. At 4 hours, corresponding to 4 PDs, the fraction of telomeres ending at position 51 is now halved to 16.8% +/- 1.5. In sum, our results are compatible with telomeres ending at positions 57-58 being consecutively converted into telomeres ending at position 51, and subsequently into telomeres ending near positions 40-45. These positions correspond to positions re-elongated by telomerase (Extended Data Fig.2j and text above). We observed similar results with the longer *FC70*, except that the positions were shifted by approximately 20 bp (Extended Data Fig.2l). Taken

altogether, our data suggests that *FC50* shortens at a rate of ~6-11 bp on half of telomeres per cell division. This indicates that the overhang length would be ~6-11 bp, which would accommodate an average shortening of ~3-5 bp/cell division measured over a population of telomeres of heterogeneous lengths.

In conclusion, sequencing of *FinalCut* telomeres allows for the detection of telomerase-mediated telomere re-elongations at Cas9-derived cleavages, demonstrating the conversion of blunt ends into canonical 3′-overhangs characteristic of chromosome ends. We further recapitulate the DNA end replication problem with unprecedent precision in its finest detail by combining the *FinalCut* system with individual telomere sequencing. Therefore, in the absence of telomerase, the shortest telomere in cells undergo gradual telomere shortening in accordance to the current model of the DNA end replication problem, prior to their switch into dysfunction and degradation when reaching 30-40 bp.

The length of the shortest telomere in the cell limits the proliferation capacity in the absence of telomerase

Our data using *FinalCut* showed that a single telomere of fixed length shortens in an asymmetrical manner over time up to a threshold length between 30 and 40 bp, below which, it transitions from functional to dysfunctional telomere. However, the precise moment when cells activate the DNA damage checkpoint and enter replicative senescence in relation to telomere status is unclear. We thought to elucidate this event using our experimental system. Since each cell contains a single critically short telomere, this complexity may manifest at the single-cell level, and we therefore analyzed cell fates in the context of these evolving telomere states in single cells, in addition to bulk populations.

At a DSB, the DNA damage checkpoint signaling cascade involves the recruitment of the Mec1/ATR kinase along with its interacting partner Ddc2/ATRIP, where they bind to newly formed ssDNA[50,51]. This complex phosphorylates Rad9 (functionally related to 53BP1 in mammals), which then binds the fork-head associated (FHA) domains of the Rad53 and Chk1 kinase. Once bound to Rad9, Rad53 is phosphorylated by Mec1, which initiates auto-hyper-phosphorylation of Rad53, and which is required to fully activate and amplify the checkpoint signal, resulting in a cell cycle arrest in G2/M. As a first approach to determine the length of the *FinalCut* telomere activating the DNA damage checkpoint, we assessed Rad53 phosphorylation in bulk *FinalCut* cell populations upon continuous Cas9 induction as described in Fig.1. We observed that the DNA damage checkpoint was fully activated 3h after the generation of a *FC20* using *FinalCut*, in accordance with observed rapid degradation of

this chromosome end and arm (Extended Data Fig.3a). This is consistent with *FC20* being recognized as a non-repairable DSB in the cell. In contrast, the induction of the *noFC* control did not alter Rad53 phosphorylation status, as predicted for 1-2 days in culture without telomerase[52,53]. Generation of *FC40*, however, triggered an increased fraction of phosphorylated Rad53 starting at 6 hours after Cas9 induction. The extent of Rad53 phosphorylation paralleled the fraction of degraded chromosome ends (Fig.1c-d). Therefore, at a population level, both telomere degradation and the activation of the DNA damage checkpoint is gradual over time for *FC40*.

We next turned to experiments assessing DNA damage checkpoint activation at the level of single cells using microfluidics coupled to live-cell imaging, which allowed us to monitor consecutive divisions of single yeast cells starting from onset of telomerase inactivation to replicative senescence onset and cell death (Fig.3a)[34,54]. Using this experimental setting, we had previously established that part of the telomerase-negative cells undergo a highly variable number of normal cell divisions before undergoing an abrupt and irreversible transition signaled by 2-4 extremely prolonged cell cycles followed by cell death, a route to senescence dubbed in our previous study as *Type A* senescence.[30] We also had found that, shortly after telomerase inactivation, 40-60% of cell lineages undergo, frequent and reversible DNA damage checkpoint cell cycle arrests prior to the terminal 2-4 extremely prolonged cell cycles and death, a separate route to senescence we called *Type B* senescence (Extended Data Fig.3b)[34,54]. While consecutive, prolonged, checkpoint-dependent cell cycles have been linked to mutagenic adaptation to DNA damage–a process allowing budding yeast cells to bypass checkpoint-induced cell cycle arrest–*type B* cells may resume normal cycles after a reversible arrest[34,54]. Assessing replicative senescence at the single-cell level in a time-resolved manner has therefore uncovered layers of intercellular heterogeneity that were not detectable in bulk population assays[24].

We thus introduced the *FinalCut* strains into our custom microfluidics device. Induction of Cas9 generates at *TEL6R* the *FinalCut* telomere that is always the shortest telomere in the cell for all *FC* strains. Cells were allowed to invade the microfluidics circuit microcavity for 16 hours in a media containing raffinose as a carbon source, prior to telomerase inactivation *via* doxycycline. Following 6 hours induction of Cas9 expression by galactose addition to the media, the carbon source was switched back to raffinose. Cells were monitored by microscopy throughout the experiment, from telomerase inactivation up to cell death (Fig.3a). Control cells with active telomerase (*noFC*, noCas9 noDox, Extended Data Fig.3c) expanded in population without limit with an average of cell cycle duration of ~145 min in raffinose and ~140 min in galactose. In this context, only 4.3% (raffinose) and

4.2% (galactose) of cell cycles were considered "long", ie above 290 or 224 min for growth in raffinose or galactose, respectively (see Methods), and the rate of cell death was 0.5%.

Analysis of the *noFC* control strain grown along the same protocol in the presence of doxycycline to repress telomerase indicated that cells entered senescence with similar dynamics as when they were grown in glucose media, even though the cell cycle durations are longer due to slower growth in raffinose and galactose media (Extended Data Fig.3d)[34,54]. The overall median lifespan of ~30 generations we observed in the absence of telomerase was compatible with previous measurements; we also observed both *Type A* and *Type B* lineages consistent with previous measurements[34,54]. These observations indicate a comparable rate of telomere shortening and a consistent mechanism of replicative senescence for cells growing in different carbon sources.

In contrast, cells with shorter *FinalCut* telomeres lose most of the inter-lineage lifespan variation (Fig.3b). *FC0* cells, for example, divide for about ~2 cell divisions after Cas9 induction before undergoing 1-2 very prolonged cell cycles and dying (Extended Data Fig.3e), suggesting that loss of a single telomere and exposure of a non-telomeric DNA end causes immediate DNA damage checkpoint activation[55]. A similar profile was found when *FC20* and *FC30* telomeres were analyzed (Fig.3b), consistent with the immediate degradation that we observed after Cas9 cleavage to generate the *FinalCut* telomere (Fig.1d-e, Extended Data Fig.2c-d). Cells with longer *FinalCut* telomeres *FC40*, *FC50*, and *FC70* exhibited a median of approximately 3, 5, and 9 cell divisions, respectively, before the onset of very prolonged cell cycles and cell death (Fig.3b). Interestingly, we also observed that these longer *FinalCut* telomeres resulted in relatively greater inter-lineage lifespan variation, mirroring the telomere length heterogeneity acquired in each cell division due to the DNA end replication problem (Extended Data Fig.2e). Taken together, these data provide compelling evidence that the sudden transition from normal to prolonged cell cycles correlates with the length of the shortest telomere in a telomerase-negative cell and the time to reach a telomere length that triggers dysfunction. This is consistent with a model whereby the length of the shortest telomere in the cell determines the onset of senescence. However, because of the growing variability in telomere length and senescence phenotypes over time, establishing a cause-and-effect relationship using cells is not trivial.

A deterministic threshold model for telomere transition into dysfunction and activation of the DNA damage checkpoint

To complement our cell-based experiments and to more directly test the hypothesis that a sudden transition into dysfunction at a precise threshold telomere length triggers senescence, we turned to mathematical modeling. We previously developed a mathematical model of replicative senescence based on the idea that the length of the shortest telomere in cells drives senescence[30]. Our model was calibrated using quantitative data obtained in microfluidics experiments of cells depleted of telomerase (using an identical repressible P_{tetO2}-*TLC1* locus) grown in glucose-based media in the presence of doxycycline, and took into account both *Type A* and *Type B* routes to senescence (Extended Data Fig.3b). The model predicted that the onset of replicative senescence of *Type A* cells is mainly driven by a deterministic threshold length of the shortest telomere in telomerase-negative cells. In contrast, it predicted that the likelihood of transition to the *Type B* lineage gradually increases as the shortest telomere shortens but that the subsequent onset of replicative senescence in these lineages occurs uniformly randomly (Fig.3c)[30].

We set our mathematical model to accommodate the experimental conditions of the *FinalCut* system. The parameters concerning cell cycle durations were adapted for cell growth in raffinose and galactose-containing media. The telomere lengths were also adapted to account for the generation of a single telomere set at a given length, according to the kinetics of Cas9 cleavage, as in Fig.1d. We next simulated cell growth in the microfluidics device and scanned a range of threshold lengths that would trigger the activation of the DNA damage checkpoint at a precise telomere length for both *Type A* and *Type B* cells. We then compared the simulations to our experimental results. We found a best fit between the experimental data and the simulations for a threshold telomere length of 33-39 bp, which defines the law of probability of transition of *Type A* cells into senescence ($p_{sen,A}$) (Fig.3d, Extended Data Fig.3f-g and Supplementary Table 7). For *Type B* cells, parameters were kept unchanged compared to the general model[30]. *FC70* contained some lineages growing longer than expected (Fig.3d). While we do not know their exact origin, we suspect that they may represent precursors of post-senescence survivors since their emergence is not included in the mathematical model[30] (see Discussion).

The close corroboration between our experimental results with the predictions of the mathematical model provide validation for the model's accuracy and assumptions to effectively capture the underlying dynamics of replicative senescence. Taken altogether, our results provide strong *in silico* and *in vivo* support that a threshold length exists to activate the DNA damage checkpoint and shift

cells suddenly to the last prolonged cell cycles of *Type A* cells (accounting for most cells in the population) and subsequent cell death.

Telomeres of 30-40 bp are also detected in a context independent of an experimentally induced critically short telomere

Taken altogether, our experimental and mathematical modeling data obtained with the *FinalCut* system show a threshold length below 40 bp at which cell division halts. This is significantly shorter than previous approximations, which estimated a longer telomere length limit of about 70-75 bp from genome sequencing data[14,56]. These latter estimations were derived from sequencing data of telomerase-negative cell populations. We thus wished to test whether telomere threshold lengths we established for telomere protection, clearly below 70-75 bp, could be detected in native telomeres not modulated by *FinalCut*. Of course, cells carrying a critically short telomere are expected to die and be immediately outgrown by proliferating cells, so that the critically short telomere in these cells becomes highly diluted within a population (by 1/32, and in an arrested cell mixed with growing ones) and thus challenging to detect. To circumvent this challenge, we sought to enrich a cell population with short telomeres without compromising cell growth by using the DNA damage checkpoint gene *RNR3* promoter[57] to control the telomerase activity through regulation of the telomerase RNA component, TLC1. We reasoned that in this strain, P_{RNR3}-*TLC1*, telomerase activity is restricted enough that cells should perpetually be on the verge of senescence (Extended Data Fig.3h). Because telomerase adds degenerated telomere repeats in budding yeast, telomerase-dependent additions can be identified by comparing the sequence of a given telomere from different cells in a population and identifying the divergence point where telomerase added *de novo* telomeric sequences as above (see Extended Data Fig.2h for an example). In the P_{RNR3}-*TLC1* strain, we expect this divergence point to reflect the minimum length of the telomere immediately before telomerase induction for extension, and therefore to reflect the telomere length reached by a given chromosome immediately before senescence regardless of whether or not it is the shortest telomere in the cell.

Consistent with our expectations that P_{RNR3}-*TLC1* cells would perpetually be on the verge of senescence, we found that these cells grew poorly, forming heterogenous-sized colonies on agar plates, which suggested a recurrent activation of the DNA damage checkpoint. Sequencing of telomeres of these cells revealed that they can shorten to a length that activates the DNA damage response yet is still extendable by telomerase. We observed a wide distribution of telomere lengths with a peak of ~80 bp (Extended Data Fig.3i-j). Notably, telomeres below 70 bp accounted for 27% of sequenced telomeres. These data show that fully native telomeres can reach lengths shorter than

70-75 bp without triggering cell death. This is compatible with our conclusion that the lower limit of telomere signaling cell cycle arrest and senescence is below 40 bp.

Genomic instability initiates at the shortest telomere in cells

Our data supports a model where a functional telomere gradually shortens due to the DNA end replication problem before transitioning to a dysfunctional state characterized by degradation along the chromosome arm (Fig.1, Extended Data Fig.1). During this process, rare genomic instability can arise, a conserved phenomenon between mammals and yeast that prompts oncogenesis in mammalian contexts[18,19,58]. In budding yeast, telomere-proximal regions have been shown to be involved in gross chromosomal rearrangements in telomerase negative populations at a frequency of $\sim 10^{-6}-10^{-4}$ [54,59,60]. We therefore asked whether the shortest telomere had a greater chance of being involved in this phenomenon.

To assess this, we mapped and quantified the genomic instability following telomerase inactivation with respect to the genomic location of a *FinalCut* telomere displaying a known length. We use fluctuation assay as a proxy for genetic instability, and hypothesized that if the telomere shortening causes genetic instability broadly across the genome, we would observe genetic instability *in trans* to the shortest telomere induced by *FinalCut*. In contrast, if genetic instability is more directly connected to dysfunction at the shortest telomere, we would expect to see instability *in cis* to the *FinalCut* telomere on the same chromosome.

We generated strains with a classical reporter cassette containing *CAN1* and *URA3* genes, which allow positive selection of mutants that arise within a population from genetic changes using canavanine- or 5-fluoroorotic acid (5FOA)-containing media. This cassette was inserted at multiple genomic locations in *cis* to the subtelomeric region adjacent to the *FinalCut* telomere, or in *trans* at another subtelomere (Supplementary Table 8, Fig.4a-b, Extended Data Fig.4a-b). The genomic locations were non-essential in haploid cells as they were positioned downstream of the last essential gene on each respective chromosome arm. When the cassette was inserted in *cis* to the *FinalCut* telomere, we generated two additional sets of strains to evaluate different scenarios: one with a homologous region for *CAN1* present on another chromosome (where the endogenous *can1-100* is present in chromosome 5, hereafter referred to as the 'donor sequence', Extended Data Fig.4a-b) and one where this region was absent (*can1-Δ*, Fig.4b).

Mutation rates per cell division were determined by scoring canavanine-resistant (Can^R) colonies and applying fluctuation analysis. These mutants were sampled from multiple independent cultures

grown for 24-hour in doxycycline-containing galactose media, and then plated onto canavanine-containing media without doxycycline to re-express telomerase to ensure their survival (Fig.4c). We chose a strain with *FC70* because it allowed gradual telomere shortening for 24 hours in the absence of telomerase prior to FC70 dysfunction (~10 generations, see Fig.3b and Extended Data Fig.4c). As negative controls, we used strains without the *FinalCut* construct (Supplementary Tables 1,8). These strains showed no increase in mutation rate throughout the experiments, as 24 hours of growth in the absence of telomerase was not sufficient to cause an increase in the *CAN1* mutation rate (Fig.4d-e, Extended Data Fig.4d-e, Supplementary Table 9)[54,59].

We found that when the reporter cassette *CAN1-URA3* was located *in trans* near *TEL5L* (at 10 kb from the last essential gene of chromosome 5L arm), the *CAN1* mutation rate remained unchanged 24 h after the generation of the *FC70* telomere (Fig.4d). In contrast, when the *CAN1-URA3* cassette was placed *in cis* near *TEL6R* bearing the *FC70* construct (either 12.5 kb or 5 kb from the first essential gene on the 6R arm, Fig.4b, Extended Data Fig.4a-b), we observed a significant increase in the *CAN1* mutation rate 24 h after the cut, regardless of the presence or absence of a donor sequence (Fig.4e, Extended Data Fig.4d-e). These results show that genomic instability during replicative senescence occurs primarily in *cis,* in the vicinity of the *FC70* telomere.

Non-reciprocal translocations at subtelomere 6R serve as a signature of genomic instability at the shortest telomere in the absence of telomerase.

We next investigated whether mutations associated with critically short telomeres exhibited a distinctive pattern or mutational signature. To accomplish this, canavanine-resistant (Can^R) colonies from the preceding experiments, selected from multiple independent cultures, were replica-plated on media lacking uracil and on media containing 5-FOA. This allowed us to distinguish gross chromosomal rearrangements (GCRs) from point mutations. We reasoned that [Can^R 5-FOA^R ura^-] colonies likely correspond to GCR whereas [Can^R 5-FOA^s ura^+] colonies likely correspond to point mutations[61]. This analysis revealed that the mutations leading to Can^R colonies in the absence of a critically short telomere in cells and of a donor sequence, both in the *noFC* strains and prior to the generation of *FC70* (0 hours in galactose), predominantly corresponded to point mutations. In contrast, the increase in the *CAN1* mutation rate proximal to the *FinalCut* telomere was caused by an increase in GCR (24 hours in galactose, Fig.4d-e).

In the presence of donor sequence, a substantial portion of Can^R colonies corresponded to GCRs in all conditions (Supplementary Figures 4d-e). To gain insight into their karyotype, we separated yeast

chromosomes by pulsed-field gel electrophoresis (PFGE) of [Can^R 5-FOA^R ura$^-$] colonies obtained 24 h after the generation of the *FC70* when the cassette was located at 12.5 kb from the first essential gene (colonies obtained with strain yT1696). As controls, we established the karyotype of the initial strain [Can^S 5-FOA^S ura$^+$] and of a [Can^R 5-FOA^S ura$^+$] colony obtained at 0 hours of galactose. For the [Can^R 5-FOA^R ura$^-$] colonies, we detected a specific translocation involving chromosome 6, when compared to the [Can^S 5-FOA^S ura$^+$] and [Can^R 5-FOA^S ura$^+$] karyotypes (Extended Data Fig.5a). Subsequent Oxford Nanopore Technologies (ONT) whole genome sequencing of a [Can^R 5-FOA^R ura$^-$] colonies obtained 24 h after the cut indicated a specific nonreciprocal translocation involving the *CAN1* sequence of the cassette proximal to *FC70* at chromosome 6 and the donor sequence at chromosome 5 (Extended Data Fig.5b-f). The donor sequence (*can1-100* locus and its flanking regions) shares perfect homology across 2674 bp (ChrV:31457-34131) with the *CAN1-URA3* cassette, except for a single nucleotide causing the ochre mutation (AAA-to-TAA nonsense change at codon 47). We reasoned that the presence of those two highly homologous sequences, both located in non-essential regions, are responsible for this observed specific nonreciprocal translocation. Moreover, this high homology between the donor sequence and the cassette could explain the substantial portion of Can^R colonies corresponding to GCRs in all conditions (Supplementary Figures 4d-e).

We next analyzed the *cis* strains lacking the donor sequence (Supplementary Tables 1,8). Results demonstrated that deletion of the *can1-100* locus and its flanking regions (*can1-Δ*) prevented spontaneous GCRs, although those associated with *FC70* telomere at 24 hours in galactose were still observed (Fig.4b,e). ONT analysis of three [Can^R 5-FOA^R ura-] colonies obtained 24 hours upon induction of *FC70* telomere, revealed several nonreciprocal translocations involving the paralogous gene loci *HXK1* (subtelomere 6R) and *HXK2* (subtelomere 7L), which share 78.1% sequence identity (Fig.5a-c, Extended Data Fig.5g-h). Taken altogether, our results demonstrate that the shortest telomere in the cell is an "at-risk" factor that increases genomic instability and promotes GCR. The characteristic of GCR being non-reciprocal translocations result in the re-lengthening of telomeric sequences at *TEL6R* possibly enabling temporary senescence escape.

Discussion

Telomere shortening in the absence of telomerase limits cell proliferation by triggering cell cycle arrest and is considered a crucial barrier to oncogenic transformation. However, this same phenomenon promotes genomic instability on the path to cellular senescence. In this work, we use *S. cerevisiae* and a new experimental system of telomere length control to decisively demonstrate

that the shortest telomere in cells is pivotal to both the onset of replicative senescence and genomic instability. We demonstrate that the transition into dysfunction of the shortest telomere in a cell provides the mechanistic link between these two seemingly opposite but intimately related processes. Our results outline a step-by-step progression from gradual telomere shortening to dysfunction at a critical length. Telomere dysfunction is characterized by extensive inward chromosome degradation from the telomere and causes a DNA-damage response checkpoint-induced arrest in cell cycle progression. This telomere degradation enhances the likelihood of presenting subtelomeric elements as substrates for gross chromosomal rearrangements.

Our *FinalCut* system, in which we use genetically controlled Cas9 to generate a single near-native telomeres of precise length per cell, offers key advantages to previous approaches to generate and study short telomeres. For instance, in the HO-cut system, the 24 nt non-telomeric HO endonuclease binding site is introduced in a subtelomeric region downstream of the last essential gene[62,63], leaving a non-telomeric terminal end within a truncated subtelomeric region. The FRT or the Cre-LoxP systems, which achieve the production of short telomeres of variable length by excising internal telomeric repeats, generates shortening to a variable length[29,64]. In contrast, the *FinalCut* system exposes a defined length of nearly identical telomeric sequences within a native subtelomeric environment, making it the most accurate system for studying telomere dynamics and the impact of telomere shortening to date.

The *FinalCut* system allowed us to observe the DNA end replication problem with unprecedented accuracy, starting with the generation of a near-native telomere of a defined length. We demonstrated that this telomere rapidly converts into a telomerase-extendible 3'-overhang, enabling it to function as a native telomerase substrate. Using *FinalCut* coupled with single molecule sequencing, we provide a solution to a long-standing controversy regarding the number of shortened telomeres that are needed to drive onset of replicative senescence. Early studies have associated telomere length with cell proliferation capacity in eukaryotes and suggested that short telomere length must limit cell divisions[8,65-67]. Experiments in human primary cells cultured *ex vivo* have indicated that approximately five short telomeres are required to arrest the cell cycle at the Hayflick limit[68]. However, *in vivo*, old baboons accumulate dermal fibroblasts displaying a single focus containing both telomeric DNA and DNA damage activation markers[69,70], suggesting that a single telomere could be enough to trigger replicative senescence (although whether this single focus corresponds to the clustering of several telomeres or to a single short telomere remains unclear). These *in vivo* findings raise the possibility that the observations in the *ex vivo* experiments could be the result of strong selection bias for cells that have bypassed the first steps of replicative senescence.

In budding yeast, our previous work suggested that an induced short telomere could accelerate replicative senescence and that senescence heterogeneity would be compatible with a single short telomere being sufficient to trigger replicative senescence[25,30,31,34,36]. We now employ single-telomere and single-cell analyses, combined with mathematical modeling, to unequivocally demonstrate that in telomerase-negative budding yeast cells, the first telomere that reaches a deterministic threshold short length between 30 and 40 bps is both necessary and sufficient to trigger the onset of replicative senescence in a large majority of cells. Whether a single short telomere is necessary and sufficient also in other eukaryotes is a possibility that requires further careful testing.

Using single molecule sequencing, we were able to visualize telomere shortening with nucleotide resolution over approximately two rounds of replication fork passage, and directly demonstrate that the resulting telomere length distributions align with the current model's prediction of asymmetrical telomere shortening, where one telomere out of two shortens (Extended Data Fig.2e)[4]. Our findings also show that the length of the 3' overhang produced by telomerase at the *FinalCut* blunt end varies from 6 to ~11 nucleotides. According to the current model, the overhang length upon telomere processing results from two parameters related to C-strand synthesis: the positioning of the primase catalytic site within the CST-Polα-primase complex relative to the chromosome end and the length of the RNA primer of the last Okazaki fragment, which is eventually degraded by a yet unknown mechanism. Given that telomere shortening in budding yeast is minimal (~3-5 bp per cell division), these parameters must be tightly regulated. However, the detected variability in shortening steps suggests some flexibility. This flexibility could arise from either the CST-Polα-primase complex positioning the RNA primer's start at various sites or from variability in the RNA primer's length. Recent studies have highlighted the CST complex's role in addressing the DNA end replication problem and stabilizing the G-overhang[5,71]. We believe the technologies developed in this work will soon clarify additional unresolved aspects of the DNA end replication problem.

What happens at the specific threshold length we identified, below which degradation extends inward into the chromosome? One can hypothesize that the telomere structure, composed of DNA, proteins, and RNA, breaks down abruptly at the threshold length. Perhaps the higher-order molecular Velcro, composed of Rap1-Rif1 and Rif2[72,73], can no longer maintain its integrity below a certain telomere length and collapses, preventing the anti-checkpoint function of the telomeres[74]. It is also possible that multimerization of CST, required for telomere elongation, requires a certain amount of DNA to synergize effectively and maintain protection from nucleases in G1 or during the passage of the replication fork[71,75]. Future work to define the telomere composition before and after telomere dysfunction will be greatly facilitated by the *FinalCut* system. It will also be important to define DNA

damage checkpoint status in relation to telomere structure. In previous work, we and others have suggested a transition from a Tel1-bound state of the telomere, to a Mec1-bound state at replicative senescence onset[31,32,76,77]. We suggest that this transition corresponds to gradual telomere shortening, involving regulated yet consistent Tel1 activation at each replication fork passage, followed by abrupt degradation potentially characterized by transient accumulation of ssDNA capable of activating Mec1. While we observed in these studies the disappearance of the telomeric signal at the point of dysfunction on both strands, further investigation is needed to determine if this indicates differential or delayed in time degradation of one strand compared to the other.

In our experiments, we demonstrated that genome instability upon telomerase inactivation is initiated in the vicinity of the first critically short telomere, the *FinalCut* telomere, while the stability of the rest of the genome does not appear to be compromised. Furthermore, the analysis of individual Can[R] colonies that arose from mutations following *FinalCut* in *cis* revealed that the vast majority of mutations associated with the shortest telomere corresponded to GCRs. These findings align with previous investigations observing an increase in genomic instability during replicative senescence in budding yeast, generally proximal to chromosome ends in comparison to the centromere-proximal region[59,60]. We now propose that these previous genomic instabilities may correspond to the subset of cells in the population where the mutation reporter is located adjacent to the shortest telomere, which is different in different cells. This result is also reminiscent of observations made in human MRC5 fibroblast cells, where the relatively short telomere on chromosome 12p is frequently involved in rearrangement events during telomere crisis[21]. Additionally, the consistently short telomere on chromosome 17p has been associated with a higher risk of esophageal cancer in individuals[78-82]. We thus speculate that the mechanisms driving genomic instability during replicative senescence, as demonstrated in yeast, may also account for these correlations found in humans. For instance, the shortest telomere, such as the 17p in humans, would be at the greatest risk of genomic instability because it would also be the first to undergo a risky specific processing, required to trigger replicative senescence.

Whole genome sequencing of Can[R] colonies selected upon *FinalCut* induction revealed that the GCRs found in *cis* of *FinalCut* telomeres correspond to nonreciprocal translocations. These translocations are characterized by junctions at homologous regions within subtelomeric repeated elements, such as 5L or 7L, serving as a mutational signature. These observations now enable us to interpret similar events found in conditions where the shortest telomere was random and not identified, mentioned above[59,60], as likely having occurred at the shortest telomeres in the ancestral cells they derived from. Notably, in our case, the subtelomere of the 6R arm is systematically

truncated. The most parsimonious hypothesis is thus that genomic instability follows the chromosome degradation we detect upon telomere dysfunction at telomeres below the length threshold. We therefore speculate that the 5'-3' exonuclease Exo1 and the flap endonuclease Rad1, previously involved in GCR formation in replicative senescence[60], are involved in telomere degradation. We also speculate that the translocations result from break-induced replication (BIR), a mechanism also involved in the alternative lengthening of telomeres (ALT) detected in post-senescent survivors in budding yeast and in approximately 10-15% of human cancers associated with poor prognosis[83,84].

Given that the nonreciprocal translocations detected in this work involve the copy of another subtelomere and telomere, the resulting cell should deactivate the DNA damage checkpoint. Technically, this cell can potentially divide for some additional generations before the second shortest telomere eventually triggers replicative senescence later on. Therefore, cells having undergone through this type of GCR share the required features that characterize senescence escape, and could therefore constitute post-senescent survivor precursors maintained by ALT via BIR. We thus speculate that the shortest telomere in cells, in addition to being the mechanistic link between proliferation limitation and genomic instability, also serves as a basis for the initial events of ALT. However, much remains to be discovered, as the type of GCRs identified in this study likely represents only a small subset of all possible events, given the selection for the loss of the *CAN1* cassette at specific genomic locations.

In conclusion, since many of our findings in budding yeast may be conserved in human cells, this work offers a potential new paradigm for understanding the link between cellular aging and increased cancer risk, as well as the etiology of syndromes associated with short telomeres. The finding that the proliferative limit is set by the shortest telomere, and that genomic instability predominantly occurs at this telomere rather than impacting other chromosomes, could have significant implications for diagnosing predisposition to cancer and yet uncharacterized genetic disorders.

Methods

Yeast strains

Yeast strains used in this work are listed in Supplementary Table 1. All strains used in this work are based on yT1487, the *noFC* control. yT1487 is a derivative of G49 (*RAD5 ADE2* W303 strain obtained

from M. Lisby) in which *TLC1* is expressed from the repressible promoter P_{tetO2} by doxycycline[34] and *P_{ADH1}-AFB2* gene is inserted at the *trp1* locus by transformation of Ylp204-PADH1-AFB2 (a gift a from Helle Ulrich). *FinalCut* strains were constructed by first creating a Cas9-induced DSB at the junction between the 6R subtelomere and *TEL6R* followed by repair with a donor fragment containing a homologous sequence for the 6R sub-telomere, a stretch of 26 bp-insertion enabling the removal of the previous Cas9 target and the insertion of a *Hind* III restriction site, a variable number of telomeric repeats n(TG$_{1-3}$), the target DNA for Cas9 (ttDNA) and a stretch of telomeric repeats. In practice, yT1487 was transformed simultaneously by pT63 and *Sph* I – *Eco* RI digested-plasmid pT182, pT177, pT176, pT175, pT174, pT187 or pT188 (Supplementary Table 3). Transformants were selected on media lacking leucine, subcloned on YPD and checked for [leu⁻]. After loss of the Cas9 expressing plasmid, constructions were checked by Southern blot and regularly by PCR using oT1572-oT1881 right before their use in subsequent experiments. In yT1612 and yT1614, the selectable marker cassette *CAN1-URA3* was inserted by transformation at 4 kb or 12 kb from *TEL6R* (12.5 or 5 kb from the last essential gene on 6R chromosome arm), to generate the strains yT1696, yT1698, yT1710 and yT1712. The *CAN1-URA3* cassette was amplified by PCR from the pT196 plasmid using the oligos oT2126-oT2127 or oT2130-oT2131 for the insertion at 4 kb or 12 kb from *TEL6R*, respectively. Transformants were selected on media lacking uracil and the expression of the *CAN1* gene was evaluated by the absence of growth on plates containing canavanine. Insertion of the *CAN1-URA3* cassette at 4 kb or 12 kb from *TEL6R* was confirmed by PCR using oT1894-oT1324 and oT1894-oT2139 primer pairs, respectively. Strains lacking the *can1-100* locus were engineered by CRISPR-Cas9 (expressed from pT194), followed by repair with the ssODN oT2276 in yT1612 and yT1614 strains. Subsequently, the *CAN1-URA3* cassette was inserted at 12 kb from *TEL6R* as previously described. A list of primers used in this work is provided in Supplementary Table 4.

Plasmids

Plasmids are described in Supplementary Table 3. Cas9-expressing pT63 was constructed by cloning annealed oT1450 and oT1451 into *Bpl* I-digested bRA90, as described[85]. Cas9-expressing pT117 was constructed by NEBuilder® HiFi DNA Assembly of 4 fragments obtained by PCR of bRA77[85] using primer pairs oT1806-oT1667; oT1666-oT1750; oT1742-oT1749 and plasmid pHis–AID*–9myc[45] using oT1741-oT1807. pT122 was constructed by first cloning annealed oT1540-oT1541 into *Bpl* I-digested bRA77, which provided the PCR amplicons with oT1829-oT1832 and oT1831-oT1834 to assemble with the PCR amplicons of pRS305 with oT1833-oT1836 and oT1830-oT1835 using the NEBuilder® HiFi DNA Assembly. pT126 was constructed by digesting pT117 with *Bpl* I and ligating to annealed oT1550 and oT1551. Plasmids pT182, pT177, pT176, pT175, pT174, pT187 and pT188

were synthesized by Genecust based on derivatives of pURTel[86] and pT26[39] harboring a cloned fragment of subtelomere 6R followed by cloned duplex oT1713-oT1714 Hind III-Xho I fragment (replacing original *URA3* marker) and by *TEL6R* telomeric repeats. To create the P_{RNR3}-*TLC1* gene, the *RNR3* promoter was amplified from yeast genomic DNA using primers FW_RNR3pr+HindIII and REV_RNR3pr+BamHI and inserted into plasmid pRS305 digested with *Hin*dIII and *Bam*HI. *TLC1* was amplified from yeast genomic DNA using primers FW_TLC1+BamHI and REV_TLC1+XbaI and inserted into the pRS305-P_{RNR3} plasmid digested with *Bam*HI and *Xba*I to create plasmid pIK2. Linearized pIK2 can then be inserted into the yeast genome at the *leu2* locus. pT194 was constructed by cloning annealed oT2057and oT2058 into *Bpl* I digested bRA90. pT196 was constructed by NEBuilder® HiFi DNA Assembly of 3 fragments obtained by PCR of *CAN1* locus from a BY4741 background strain with oT2082 and oT2085 and PCRs of pRS306 with primers oT2028 and oT2083 on one hand and oT2084 and oT2029, on the other hand.

Galactose induced Cas9 cleavage of *FinalCut* telomeres

FinalCut strains transformed with pT126 with the one-step protocol[87] were inoculated in 10 ml of synthetic media lacking leucine containing 2% raffinose (S -LEU R), grown for 5 hours at 30°C at 220 rpm and diluted to OD_{600nm} = 0,01-0,02 in S -LEU R either with or without doxycyline at 30 µg/ml. Exponentially growing cells were collected 6h to overnight growth. Cells were sampled for time point 0 hours, washed and diluted to OD_{600nm} = 0,2 in synthetic media lacking leucine containing 2% raffinose and 3% galactose (S -LEU RG) either with or without doxycyline at 30 µg/ml, and grown for 3 hours. Sample collection, dilution to DO_{600nm} = 0,3 were repeated each 3 hours with 9-hour time point being the last one collected. In experiments in which Cas9 is switched off, cells were grown similarly with the following differences. At 4 hours after galactose addition, cells were washed and diluted to OD_{600nm} = 0,1 in YPD either with or without doxycyline at 30 µg/ml and 0,25 mM of 1-Naphthalenacetic acid (NAA). Samples were collected every two hours (t=4+2, 4+4, 4+6). In microfluidics experiments, *FinalCut* strains were transformed with integrative pT122 and selected on media lacking leucine. Exponentially growing cells were injected in the microfluidics circuit and allowed to divide in YPD to invade the cavities for 12–24 h before the medium was switched to YP-2% Raffinose either with or without doxycyline at 30 µg/ml. The induction of Cas9 was achieved by switching the medium to YP-2% Raffinose, 2% galactose either with or without doxycyline at 30 µg/ml for 6 hours before changing media back to YP 2% Raffinose either with or without doxycyline at 30 µg/ml. For each set of experiments, at least two independent original transformants for the *FinalCut* constructs were involved in triplicate experiments.

Southern blots

Genomic DNA was extracted from cultures using a standard phenol: chloroform:isoamyl (25:24:1) purification procedure and isopropanol precipitation. 500 ng of genomic DNA were digested with *Hpa* I (NEB), ethanol-precipitated, resuspended in gel loading dye (Purple 6x, NEB), and resolved on a 1% agarose gel for 16 h at 50 V. The gel was then soaked in 0,25 M HCl followed by 20 min incubation in 0.4 M NaOH, 1 M NaCl. Transfer was performed overnight by capillarity to a charged nylon membrane (Hybond XL, GE Healthcare or Hybond N+ cytiva #RPN303B). Membranes were pre-hybridized for 2 h at 65°C in rotating tubes before radioactive probes were added. Random priming reaction by Klenow Fragment (3'→5' exo-, NEB) using α-^{32}P-dATP was employed to label amplicons of gDNA with oT1959-oT1960 (TRF *FinalCut* probe), with oT2189-oT2190 (Chromosome arm 6R internal band) or with oT1963-oT1964 (loading control). The telomere-specific oligonucleotide probe oT2032 was 5'-labelled using T4 polynucleotide kinase (NEB) and γ-^{32}P- ATP. Hybridization was performed overnight at 65°C. Membranes were washed increasing stringency's buffers (Supplementary Table 5). Membranes were exposed to BAS Storage Phosphor Screens (GE Healthcare) for 2 days and imaged with a Typhoon FLA 9500 scanner (GE Healthcare). Band intensities were measured using ImageJ 1.53a[88]. The smears/bands corresponding to the 6R telomere were first normalized to the loading control band prior to other calculations. Uncut and cut quantities were obtained by a second normalization on the 0 hours "uncut" time point. Degraded quantities were estimated subtracting the sum of uncut and cut to the sum of uncut and cut at 0 hours. In experiments in which Cas9 is switched off after 4 h in galactose, the stability of the *FinalCut* telomere at different time points was estimated by normalizing to t = [4+0].

Western Blot

Protein analyses were performed as described[53]. In brief, 5 OD$_{600nm}$ of cells were lysed in NaOH 0,2 M for 10 minutes on ice and proteins precipitated at a final concentration of 0,5% trichloroacetic acid by incubation again on ice for 10 minutes. After centrifuging for 10 minutes at maximum speed at 4°C, the pellet was resuspended in 50 μL of Laemmli 4x buffer and denatured at 95°C for five minutes. Protein samples were electrophoresed on 7,5% of 37,5:1 Acrylamide:Bis-acrylamide (Sigma-Aldrich #A3699) for Rad53 detection or 4–15% Mini-PROTEAN® TGX™ precast polyacrylamide gels (BioRad) for Cas9 detection. Proteins were then transferred to a nitrocellulose membrane (Amersham Protran 0.45 NC, GE HealthCare), stained with Ponceau red (Sigma-Aldrich). Membranes were blocked with 5% milk and incubated with the appropriate primary antibody: Rad53 (1:1,000, Abcam #ab104252), c-Myc (1:2,000, Sigma-Aldrich #M5546), FLAG (1:2,000, Sigma Aldrich, SAB431135) and Pgk1

(1:20,000, Abcam #ab113687). After washes, the primary antibodies were recognized by the horseradish peroxidase-coupled secondary antibody (1:10,000, Sigma-Aldrich #A9169). The signals were revealed using an electrochemiluminescence reagent (ClarityWestern ECL, Biorad) and recorded using the ChemiDoc Imaging System (Biorad).

TeloPCR

Amplification of 6R specific telomeres was performed with 40-100 ng of gDNA as described[40]. DNA was denatured in a volume of 6 µl at 95°C during 5 min and end-tailed in 1x NEB4 buffer (NEB), 100 µM dCTP and 1 unit of Terminal Transferase (NEB #M0315) at 37°C 30 min, followed by a 10 min incubation at 65°C and 5 min at 94°C. PCR mix was added to obtain final concentrations of 200 µM of dNTPs, 0.5 µM of oT883, 0.75 µM of oT1310, 1x Taq ThermoPol Mg2+ free buffer (NEB #B9005) and 2,5 U of Taq polymerase (NEB #M0273) in a 40 µl final volume. PCR program was 95°C 3 min; followed by 45 cycles of 95°C 20 sec, 62°C 15 sec and 72°C 50 sec, followed by 72°C 5 min.

Individual telomere sequencing by Nanopore

6R telomeres were amplified by TeloPCR with the following modifications. Primers used were oT2132 and oT2158. PCR mix was composed of 1x Colorless GoTaq Flexi buffer (Promega, Cat M7806)) and 2,5 U of GoTaq G2 Flexi DNA polymerase. TeloPCR products were purified from a 1.5% agarose gel using the Macherey-Nagel NucleoSpin Gel and PCR Clean-up Kit (Macherey Nagel #740609.50), and cloned into pUC18 amplified with oT2162 and oT2163 using the NEBuilder HiFi DNA Assembly following supplier instructions. 2 µl of the ligation was transformed into Stable competent bacteria (NEB #C3040) according to supplier instructions. Plates containing 500-10,000 colonies were scraped and plasmids were extracted in batch with the NucleoSpin Plasmid kit (Macherey Nagel #740588.50). ~1 µg of plasmid DNA was linearized with *Eco* RI (NEB #R3101) and libraries of 12 samples were multiplexed using the Ligation sequencing gDNA - Native Barcoding Kit 24 V14 (Oxford Nanopore Technologies SQK-NBD114.24) and run on a MinION flowcell (v.10.4.1). For analyses, two distinct pipelines were followed. In Fig. 2c, Extended Data Fig. 2g-l, reads were basecalled and demultiplexed by Guppy v6.5.7. Nanopore adapter barcodes were removed and reads were demultiplexed by TelIndex. Resulting fastq reads displaying the same TelIndex were aligned using a custom Geneious Prime 2023.2.1 workflow (see Supplementary Pipeline 1). Resulting Consensus were aligned to eliminate residual minor sequencing errors manually and visualizations were made using Geneious Prime 2023.2.1. For Fig. 2b-c and Extended Data Fig. 2l was followed.

Microfluidics

Experiments were conducted as described[34] in [10-chamber chips](#) casted with polydimethylsiloxane (PDMS; Sylgard 184) and curing agent mixed in a 10:1 ratio, degassed with a vacuum pump for 30 min, and poured into the mold. The PDMS was cured by baking for 5 h at 70°C and then carefully removed from the mold. A biopsy puncher (1.5 mm; Harris Unicore) was used to create holes to connect the tubing. The PDMS and a glass coverslip (24×50 mm) were surface-activated by plasma (Diener Electronic) to covalently bond the two parts. YP 2% Raffinose or YP 2% Raffinose 3% Galactose dissolved in milliQ water were filtered using a 0.22-µm polyethersulfone filter (Millipore) and loaded into the device using a peristaltic pump (IPCN, Ismatec). Cells from a log-phase culture (OD_{600nm} = 0.5) were injected into the device using a 1-mL syringe. A constant medium flow (10 µL/min) was maintained throughout the experiment and media replaced every 48 hours. Cells were imaged using a fully motorized Axio Observer Z1 inverted microscope (Zeiss) with a 100× immersion objective, a Hamamatsu Orca R2 camera, and constant focus maintained with focus stabilization hardware (Definite focus, Zeiss). For phase contrast images we used light-emitting diode light sources (Colibri 2, Zeiss) with the following settings for standard microfluidic time-lapse experiments: 4.0 V for 70 msec. The temperature was maintained at 30°C with a controlled heating unit and an incubation chamber that held the entire microscope base, including the stage and the objectives. Images were acquired every 10 min using Zen software (Zeiss). All aspects of image acquisition were fully automated and controlled, including temperature, focus, stage position, and time-lapse imaging. Images were acquired for up to 576 h. Time-lapse images were analyzed using a Matlab R2021b software as described[53]. Lineages that experienced very long cell cycles before the galactose addition were excluded from the analysis. The threshold for prolonged cell cycles was set to [mean+ 2xSD] of the dataset, the mean and SD being computed from cell cycle durations in media containing raffinose only or raffinose and galactose. For each set of experiments, at least two independent original transformants for the *FinalCut* constructs were involved in triplicate experiments.

Mathematical modeling

The mathematical model is based on our previous work [30] and more details, original and FinalCut-adapted codes can be found [here](#). The following modifications have been implemented to encode the *FinalCut* experimental setting. Time 0 in the simulation still corresponds to telomerase inactivation, but we now also model galactose addition at time 6h. As soon as galactose is added, we forget the lineage's past history and reset cell generation so the first cell born under galactose has generation 0. From this time until time $t_{noCut} = t_{raf} + delay$, uncut cells have a certain probability at each

division to have one of their telomeres cut down to a defined length ℓ_{cut}. We chose *delay* = 9h and $t_{raf} = t_{gal}$ + 6 h. We then estimated the probability of cut from the cumulated distribution of cells where a telomere has been cut, fitted with the experimental cutting efficiency of *FC20* (Fig. 1e).

Despite adding a probability to be cut, the implementation of galactose and raffinose conditions also comes with new distributions of cell cycle duration times. We followed the methodology of [30] to construct them, using the data of strain *noFC* noCas9 noDOX (Extended Data Fig. 3c). As indicated above, the threshold for prolonged cell cycles was set to [mean+ 2xSD] of the dataset, the mean and SD being computed from cell cycle durations in media containing raffinose only or raffinose and galactose. Other parameters of the model required a few adjustments. We modified the parameter $\ell_{min,A}$ to $\ell_{min,A}$ = 36 bp in order to better fit, for all ℓ_{cut} values, the generations of senescence onset obtained with *FinalCut* experiments. Because the parameters of $p_{sen.A}$, $p_{sen.B}$ and p_{nta} are not independent, modifying $p_{sen.A}$ through $\ell_{min,A}$ cannot be done without slight corrections of the other parameters, in such a way that the previous experiments (simulated in [30]) are still well fitted (Extended Data Fig. 3a). These slight modifications are in accordance with our previous sensitivity simulations, which proved robustness of our fits to minor changes of the value of $\ell_{min,A}$.

Fluctuation assay for mutation rate estimation

Five independent transformants with pT126 plasmid of relevant strains (Table 1) were inoculated, in 50 mL of S -LEU R and 30 µg/mL doxycycline to obtain exponentially growing cultures after 16 h at 30°C. 200 cells were plated on YPD plates (control) and 10^6, 10^7 or 10^8 cells were plated on synthetic complete medium-containing 2% glucose (SD) plates lacking arginine and supplemented with 60 µg/mL canavanine. These cultures were considered as time zero. To induce Cas9-AID*-3MYC, time 0 cultures were diluted into 40 mL of S -LEU RG and 30 µg/mL doxycycline at an appropriate concentration to obtain an exponentially growing culture after 24h. Cultures were counted and plated as described above. Plates were incubated for 48h at 30°C, and colonies counted. The mutation rate was estimated as described in [54] using a Matlab code. For each set of experiments, at least two independent original transformants for the *FinalCut* constructs were involved in triplicate experiments (each involving 5 transformants of pT126).

Pulse-field gel electrophoresis (PFGE)

For separation of yeast chromosomes, 5×10^7 cells from exponential cultures were washed in 10 mM Tris-HCl and 50 mM EDTA (pH 8.0) and then cast in 0.45% low melting point agarose (MP Biomedicals) plugs (molds, Biorad) in Zymoliase buffer (50 mM potassium phosphate, 50 mM EDTA,

pH 7.0). Plugs were incubated in the Zymoliase buffer containing 10mM DTT and 0.4 mg/ml Zymoliase 20T (Seikagaku) overnight at 37°C. The solution was replaced by the proteinase K buffer (10mM Tris, pH 7.5, 50mM EDTA, 1% Sarkosyl) and 2 mg/ml proteinase K (NEB), and then incubated 24 h at 50 °C. After washing three times with 10 mM Tris-HCl and 50 mM EDTA (pH 8.0), plugs were then inserted into the wells of a 0.9 % agarose (Seakem Gold, Lonza) gel in 0.5x Tris-Borate-EDTA. Whole chromosomes were then separated in a rotating PFGE apparatus (Rotaphor 6.0, Biometra) with parameters chosen for *Saccharomyces cerevisia* according to manufacturer instructions. DNA was visualized by staining with 0.5 µg/mL ethidium bromide under UV illumination (Gel Doc, Bio-Rad). High molecular weight genomic DNA size was estimated through an 0.75% agarose gel in 0.5X TBE in a PFGE apparatus (Rotaphor 6.0, Biometra) following parameters: pulse time 4-2 s (decreasing linearly), a rotor angle of 120-110° (decreasing linearly) and voltage of 130-90 V (decreasing linearly) during 18h at 13°C.

Whole genome sequencing

High molecular weight genomic DNA was extracted from exponential phase cultures using MasterPure™ yeast purification kit (Biosearch technologies, #MPY80200) or Genomic DNA buffer set (Qiagen, #19060) with Qiagen Genomic-Tip 100/G (Qiagen, #10243), following the supplier instructions. DNA concentration was measured using Qubit 2.0 fluorometer with Qubit™ dsDNA BR Assay Kit (Invitrogen, #Q32853), while DNA purity was estimated by measuring the OD_{260}/OD_{280} using NanoDrop™ 2000. DNA length was determined by PFGE. Libraries were prepared according to the Ligation sequencing gDNA - Native Barcoding Kit 24 V14 (Oxford Nanopore Technologies SQK-NBD114.24) and run on a MinION flowcell (v.10.4.1). The signal was basecalled using Dorado v0.71 and high accuracy model. Reads were mapped on the W303 (ura-can) genome and CGR are reported using LAST software (lastal v.1548) and dnarrange respectively[89]. CGR were visualized using CIRCOS[90].

References

de Lange, T. Shelterin-Mediated Telomere Protection. *Annu Rev Genet* **52**, 223-247 (2018). https://doi.org:10.1146/annurev-genet-032918-021921

Jain, D. & Cooper, J. P. Telomeric strategies: means to an end. *Annu Rev Genet* **44**, 243-269 (2010). https://doi.org:10.1146/annurev-genet-102108-134841

Lingner, J., Cooper, J. P. & Cech, T. R. Telomerase and DNA end replication: no longer a lagging strand problem? *Science* **269**, 1533-1534 (1995).

Soudet, J., Jolivet, P. & Teixeira, M. T. Elucidation of the DNA end-replication problem in Saccharomyces cerevisiae. *Mol Cell* **53**, 954-964 (2014). https://doi.org:10.1016/j.molcel.2014.02.030

Takai, H., Aria, V., Borges, P., Yeeles, J. T. P. & de Lange, T. CST-polymerase α-primase solves a second telomere end-replication problem. *Nature* **627**, 664-670 (2024). https://doi.org:10.1038/s41586-024-07137-1

Vasianovich, Y. & Wellinger, R. J. Life and Death of Yeast Telomerase RNA. *Journal of molecular biology* **429**, 3242-3254 (2017). https://doi.org:10.1016/j.jmb.2017.01.013

Ha, T., Kaiser, C., Myong, S., Wu, B. & Xiao, J. Next generation single-molecule techniques: Imaging, labeling, and manipulation in vitro and in cellulo. *Mol Cell* **82**, 304-314 (2022). https://doi.org:10.1016/j.molcel.2021.12.019

Lundblad, V. & Szostak, J. W. A mutant with a defect in telomere elongation leads to senescence in yeast. *Cell* **57**, 633-643 (1989). https://doi.org:10.1016/0092-8674(89)90132-3

Enomoto, S., Glowczewski, L. & Berman, J. MEC3, MEC1, and DDC2 are essential components of a telomere checkpoint pathway required for cell cycle arrest during senescence in Saccharomyces cerevisiae. *Molecular biology of the cell* **13**, 2626-2638 (2002). https://doi.org:10.1091/mbc.02-02-0012

d'Adda, d. F. F. *et al.* A DNA damage checkpoint response in telomere-initiated senescence. *Nature* **426**, 194-198 (2003). https://doi.org:10.1038/nature02118

Shay, J. W. Role of Telomeres and Telomerase in Aging and Cancer. *Cancer discovery* **6**, 584-593 (2016). https://doi.org:10.1158/2159-8290.Cd-16-0062

Hanahan, D. & Weinberg, R. A. The hallmarks of cancer. *Cell* **100**, 57-70 (2000). https://doi.org:S0092-8674(00)81683-9 [pii]

Lundblad, V. & Blackburn, E. H. An alternative pathway for yeast telomere maintenance rescues est1- senescence. *Cell* **73**, 347-360 (1993). https://doi.org:10.1016/0092-8674(93)90234-h

Kockler, Z. W., Comeron, J. M. & Malkova, A. A unified alternative telomere-lengthening pathway in yeast survivor cells. *Mol Cell* **81**, 1816-1829 e1815 (2021). https://doi.org:10.1016/j.molcel.2021.02.004

Dilley, R. L. *et al.* Break-induced telomere synthesis underlies alternative telomere maintenance. *Nature* **539**, 54-58 (2016). https://doi.org:10.1038/nature20099

Lydeard, J. R., Jain, S., Yamaguchi, M. & Haber, J. E. Break-induced replication and telomerase-independent telomere maintenance require Pol32. *Nature* **448**, 820-823 (2007). https://doi.org:10.1038/nature06047

Feldser, D. M., Hackett, J. A. & Greider, C. W. Telomere dysfunction and the initiation of genome instability. *Nature reviews. Cancer* **3**, 623-627 (2003). https://doi.org:10.1038/nrc1142

Maciejowski, J. & de, L. T. Telomeres in cancer: tumour suppression and genome instability. *Nature reviews. Molecular cell biology* **18**, 175-186 (2017). https://doi.org:10.1038/nrm.2016.171

Henninger, E. & Teixeira, M. T. Telomere-driven mutational processes in yeast. *Current opinion in genetics & development* **60**, 99-106 (2020). https://doi.org:10.1016/j.gde.2020.02.018

Cleal, K. & Baird, D. M. Catastrophic Endgames: Emerging Mechanisms of Telomere-Driven Genomic Instability. *Trends in genetics : TIG* **36**, 347-359 (2020). https://doi.org:10.1016/j.tig.2020.02.001

Dewhurst, S. M. *et al.* Structural variant evolution after telomere crisis. *Nat Commun* **12**, 2093 (2021). https://doi.org:10.1038/s41467-021-21933-7

Teixeira, M. T. Saccharomyces cerevisiae as a Model to Study Replicative Senescence Triggered by Telomere Shortening. *Frontiers in oncology* **3**, 101 (2013). https://doi.org:10.3389/fonc.2013.00101

Thosar, S. A. *et al.* Oxidative guanine base damage plays a dual role in regulating productive ALT-associated homology-directed repair. *Cell reports* **43**, 113656 (2024). https://doi.org:10.1016/j.celrep.2023.113656

Xu, Z. & Teixeira, M. T. The many types of heterogeneity in replicative senescence. *Yeast* **36**, 637-648 (2019). https://doi.org:10.1002/yea.3433

Bourgeron, T., Xu, Z., Doumic, M. & Teixeira, M. T. The asymmetry of telomere replication contributes to replicative senescence heterogeneity. *Scientific reports* **5**, 15326 (2015). https://doi.org:10.1038/srep15326

Taherian, F. A. *et al.* Deconstructing heterogeneity of replicative senescence in human mesenchymal stem cells at single cell resolution. *GeroScience* **46**, 999-1015 (2024). https://doi.org:10.1007/s11357-023-00829-y

Wellinger, R. J. & Zakian, V. A. Everything you ever wanted to know about Saccharomyces cerevisiae telomeres: beginning to end. *Genetics* **191**, 1073-1105 (2012). https://doi.org:10.1534/genetics.111.137851

Kupiec, M. Biology of telomeres: lessons from budding yeast. *FEMS Microbiol Rev* **38**, 144-171 (2014). https://doi.org:10.1111/1574-6976.12054

Marcand, S., Brevet, V. & Gilson, E. Progressive cis-inhibition of telomerase upon telomere elongation. *The EMBO journal* **18**, 3509-3519 (1999). https://doi.org:10.1093/emboj/18.12.3509

Rat, A., Martinez Fernandez, V., Doumic, M., Teixeira, M. T. & Xu, Z. Mathematical model linking telomeres to senescence in Saccharomyces cerevisiae reveals cell lineage versus population dynamics. *Nat Commun* **16**, 1024 (2025). https://doi.org:10.1038/s41467-025-56196-z

Abdallah, P. *et al.* A two-step model for senescence triggered by a single critically short telomere. *NCB* **11**, 988-993 (2009). https://doi.org:10.1038/ncb1911

Khadaroo, B. *et al.* The DNA damage response at eroded telomeres and tethering to the nuclear pore complex. *NCB* **11**, 980-987 (2009). https://doi.org:10.1038/ncb1910

Xu, Z., Duc, K. D., Holcman, D. & Teixeira, M. T. The length of the shortest telomere as the major determinant of the onset of replicative senescence. *Genetics* **194**, 847-857 (2013). https://doi.org:10.1534/genetics.113.152322

Xu, Z. *et al.* Two routes to senescence revealed by real-time analysis of telomerase-negative single lineages. *Nat Commun* **6**, 7680 (2015). https://doi.org:10.1038/ncomms8680

Xie, Z. *et al.* Early telomerase inactivation accelerates aging independently of telomere length. *Cell* **160**, 928-939 (2015). https://doi.org:10.1016/j.cell.2015.02.002

Martin, H., Doumic, M., Teixeira, M. T. & Xu, Z. Telomere shortening causes distinct cell division regimes during replicative senescence in Saccharomyces cerevisiae. *Cell Biosciences* **11**, 180 (2021). https://doi.org:10.1186/s13578-021-00693-3

Eugene, S., Bourgeron, T. & Xu, Z. Effects of initial telomere length distribution on senescence onset and heterogeneity. *JTB* **413**, 58-65 (2017). https://doi.org:10.1016/j.jtbi.2016.11.010

Pasquier, E. & Wellinger, R. J. In vivo chromatin organization on native yeast telomeric regions is independent of a cis-telomere loopback conformation. *Epigenetics & chromatin* **13**, 23 (2020). https://doi.org:10.1186/s13072-020-00344-w

Graf, M. *et al.* Telomere Length Determines TERRA and R-Loop Regulation through the Cell Cycle. *Cell* **170**, 72-85 e14 (2017). https://doi.org:10.1016/j.cell.2017.06.006

Forstemann, K., Hoss, M. & Lingner, J. Telomerase-dependent repeat divergence at the 3' ends of yeast telomeres. *Nucleic acids research* **28**, 2690-2694 (2000). https://doi.org:10.1093/nar/28.14.2690

Bah, A., Gilson, E. & Wellinger, R. J. Telomerase is required to protect chromosomes with vertebrate-type T2AG3 3' ends in Saccharomyces cerevisiae. *The Journal of biological chemistry* **286**, 27132-27138 (2011). https://doi.org:10.1074/jbc.M111.220186

Strecker, J. *et al.* A sharp Pif1-dependent threshold separates DNA double-strand breaks from critically short telomeres. *eLife* **6:e23783** (2017). https://doi.org:10.7554/eLife.23783

Teixeira, M. T., Arneric, M., Sperisen, P. & Lingner, J. Telomere length homeostasis is achieved via a switch between telomerase- extendible and -nonextendible states. *Cell* **117**, 323-335 (2004). https://doi.org:10.1016/s0092-8674(04)00334-4

Nishimura, K., Fukagawa, T., Takisawa, H., Kakimoto, T. & Kanemaki, M. An auxin-based degron system for the rapid depletion of proteins in nonplant cells. *Nat Methods* **6**, 917-922 (2009). https://doi.org:10.1038/nmeth.1401

Morawska, M. & Ulrich, H. D. An expanded tool kit for the auxin-inducible degron system in budding yeast. *Yeast* **30**, 341-351 (2013). https://doi.org:10.1002/yea.2967

Lingner, J. & Cech, T. R. Purification of telomerase from Euplotes aediculatus: requirement of a primer 3' overhang. *Proc Natl Acad Sci U S A* **93**, 10712-10717 (1996). https://doi.org:10.1073/pnas.93.20.10712

Lemos, B. R. *et al.* CRISPR/Cas9 cleavages in budding yeast reveal templated insertions and strand-specific insertion/deletion profiles. *Proc Natl Acad Sci U S A* (2018). https://doi.org:10.1073/pnas.1716855115

Ben-Tov, D. *et al.* Uncovering the dynamics of precise repair at CRISPR/Cas9-induced double-strand breaks. *Nat Commun* **15**, 5096 (2024). https://doi.org:10.1038/s41467-024-49410-x

Forstemann, K. & Lingner, J. Molecular basis for telomere repeat divergence in budding yeast. *MCB* **21**, 7277-7286. (2001). https://doi.org:10.1128/MCB.21.21.7277-7286.2001

Cejka, P. & Symington, L. S. DNA End Resection: Mechanism and Control. *Annu Rev Genet* **55**, 285-307 (2021). https://doi.org:10.1146/annurev-genet-071719-020312

Waterman, D. P., Haber, J. E. & Smolka, M. B. Checkpoint Responses to DNA Double-Strand Breaks. *Annu. Rev. Biochem.* (2020). https://doi.org:10.1146/annurev-biochem-011520-104722

Ijpma, A. S. & Greider, C. W. Short telomeres induce a DNA damage response in Saccharomyces cerevisiae. *Molecular biology of the cell* **14**, 987-1001 (2003). https://doi.org:10.1091/mbc.02-04-0057

Zeinoun, B., Teixeira, M. T. & Barascu, A. Hog1 acts in a Mec1-independent manner to counteract oxidative stress following telomerase inactivation in Saccharomyces cerevisiae. *Communications Biology* **7**, 761 (2024). https://doi.org:10.1038/s42003-024-06464-3

Coutelier, H. *et al.* Adaptation to DNA damage checkpoint in senescent telomerase-negative cells promotes genome instability. *Genes Dev* **32**, 1499-1513 (2018). https://doi.org:10.1101/gad.318485.118

Sandell, L. L. & Zakian, V. A. Loss of a yeast telomere: arrest, recovery, and chromosome loss. *Cell* **75**, 729-739 (1993). https://doi.org:10.1016/0092-8674(93)90493-a

Sholes, S. L. *et al.* Chromosome-specific telomere lengths and the minimal functional telomere revealed by nanopore sequencing. *Genome Research* **32**, 616-628 (2021). https://doi.org:10.1101/gr.275868.121

Elledge, S. J. & Davis, R. W. Two genes differentially regulated in the cell cycle and by DNA-damaging agents encode alternative regulatory subunits of ribonucleotide reductase. *Genes Dev* **4**, 740-751 (1990). https://doi.org:10.1101/gad.4.5.740

Artandi, S. E. & DePinho, R. A. A critical role for telomeres in suppressing and facilitating carcinogenesis. *Current opinion in genetics & development* **10**, 39-46 (2000). https://doi.org:10.1016/s0959-437x(99)00047-7

Hackett, J. A., Feldser, D. M. & Greider, C. W. Telomere dysfunction increases mutation rate and genomic instability. *Cell* **106**, 275-286 (2001). https://doi.org:10.1016/s0092-8674(01)00457-3

Hackett, J. A. & Greider, C. W. End resection initiates genomic instability in the absence of telomerase. *MCB* **23**, 8450-8461 (2003). https://doi.org:10.1128/MCB.23.23.8450-8461.2003

Chen, C. & Kolodner, R. D. Gross chromosomal rearrangements in Saccharomyces cerevisiae replication and recombination defective mutants. *Nature Genetics* **23**, 81-85 (1999). https://doi.org:10.1038/12687

Haber, J. E. Mating-type gene switching in Saccharomyces cerevisiae. *Annu Rev Genet* **32**, 561-599 (1998). https://doi.org:10.1146/annurev.genet.32.1.561

Diede, S. J. & Gottschling, D. E. Telomerase-mediated telomere addition in vivo requires DNA primase and DNA polymerases alpha and delta. *Cell* **99**, 723-733 (1999). https://doi.org:10.1016/s0092-8674(00)81670-0

Marcand, S., Brevet, V., Mann, C. & Gilson, E. Cell cycle restriction of telomere elongation. *Current Biology* **10**, 487-490 (2000). https://doi.org:10.1016/s0960-9822(00)00450-4

Greider, C. W. Telomeres, telomerase and senescence. *BioEssays : news and reviews in molecular, cellular and developmental biology* **12**, 363-369 (1990). https://doi.org:10.1002/bies.950120803

Lendvay, T. S., Morris, D. K., Sah, J., Balasubramanian, B. & Lundblad, V. Senescence mutants of Saccharomyces cerevisiae with a defect in telomere replication identify three additional EST genes. *Genetics* **144**, 1399-1412 (1996). https://doi.org:10.1093/genetics/144.4.1399

Hemann, M. T., Strong, M. A., Hao, L. Y. & Greider, C. W. The shortest telomere, not average telomere length, is critical for cell viability and chromosome stability. *Cell* **107**, 67-77. (2001). https://doi.org:10.1016/s0092-8674(01)00504-9

Kaul, Z., Cesare, A. J., Huschtscha, L. I., Neumann, A. A. & Reddel, R. R. Five dysfunctional telomeres predict onset of senescence in human cells. *EMBO Rep* **13**, 52-59 (2011). https://doi.org:10.1038/embor.2011.227

Herbig, U., Ferreira, M., Condel, L., Carey, D. & Sedivy, J. M. Cellular Senescence in Aging Primates. *Science* **311**, 1257 (2006). https://doi.org:Cellular Senescence in Aging Primates

Jeyapalan, J. C., Ferreira, M., Sedivy, J. M. & Herbig, U. Accumulation of senescent cells in mitotic tissue of aging primates. *Mech Ageing Dev* **128**, 36-44 (2007). https://doi.org:10.1016/j.mad.2006.11.008

Coloma, J. *et al.* Molecular architecture and oligomerization of Candida glabrata Cdc13 underpin its telomeric DNA-binding and unfolding activity. *Nucleic acids research* **51**, 668-686 (2023). https://doi.org:10.1093/nar/gkac1261

Shi, T. *et al.* Rif1 and Rif2 shape telomere function and architecture through multivalent Rap1 interactions. *Cell* **153**, 1340-1353 (2013). https://doi.org:10.1016/j.cell.2013.05.007

Yang, C. W. *et al.* Telomere shortening triggers a feedback loop to enhance end protection. *Nucleic acids research* **45**, 8314-8328 (2017). https://doi.org:10.1093/nar/gkx503

Michelson, R. J., Rosenstein, S. & Weinert, T. A telomeric repeat sequence adjacent to a DNA double-stranded break produces an anticheckpoint. *Genes Dev* **19**, 2546-2559 (2005). https://doi.org:10.1101/gad.1293805

Runnberg, R., Narayanan, S., Itriago, H. & Cohn, M. Either Rap1 or Cdc13 can protect telomeric single-stranded 3' overhangs from degradation in vitro. *Scientific reports* **9**, 19181 (2019). https://doi.org:10.1038/s41598-019-55482-3

Hector, R. E. *et al.* Mec1p associates with functionally compromised telomeres. *Chromosoma* **121**, 277-290 (2012). https://doi.org:10.1007/s00412-011-0359-0

Fallet, E. *et al.* Length-dependent processing of telomeres in the absence of telomerase. *Nucleic acids research* **42**, 3648-3665 (2014). https://doi.org:10.1093/nar/gkt1328

Martens, U. M. *et al.* Short telomeres on human chromosome 17p. *Nat Genet* **18**, 76-80 (1998). https://doi.org:10.1038/ng0198-018

Xing, J. *et al.* Constitutive short telomere length of chromosome 17p and 12q but not 11q and 2p is associated with an increased risk for esophageal cancer. *Cancer Prev Res (Phila)* **2**, 459-465 (2009). https://doi.org:10.1158/1940-6207.Capr-08-0227

Karimian, K. *et al.* Human telomere length is chromosome end-specific and conserved across individuals. *Science* **384**, 533-539 (2024). https://doi.org:10.1126/science.ado0431

Schmidt, T. T. *et al.* High resolution long-read telomere sequencing reveals dynamic mechanisms in aging and cancer. *Nat Commun* **15**, 5149 (2023). https://doi.org:10.1101/2023.11.28.569082

Sanchez, S. E. *et al.* Digital telomere measurement by long-read sequencing distinguishes healthy aging from disease. *Nat Commun* **15**, 5148 (2024). https://doi.org:10.1038/s41467-024-49007-4

Dilley, R. L. & Greenberg, R. A. ALTernative Telomere Maintenance and Cancer. *Trends Cancer* **1**, 145-156 (2015). https://doi.org:10.1016/j.trecan.2015.07.007

Willis, N. A., Rass, E. & Scully, R. Deciphering the Code of the Cancer Genome: Mechanisms of Chromosome Rearrangement. *Trends Cancer* **1**, 217-230 (2015). https://doi.org:10.1016/j.trecan.2015.10.007

Anand, R., Beach, A., Li, K. & Haber, J. Rad51-mediated double-strand break repair and mismatch correction of divergent substrates. *Nature* **544**, 377-380 (2017). https://doi.org:10.1038/nature22046

Gottschling, D. E., Aparicio, O. M., Billington, B. L. & Zakian, V. A. Position effect at S. cerevisiae telomeres: reversible repression of Pol II transcription. *Cell* **63**, 751-762 (1990).

Chen, D.-C., Yang, B.-C. & Kuo, T.-T. One step transformation of yeast in stationary phase. *Current Genetics* **21**, 83-84 (1992). https://doi.org:10.1007/BF00318659

Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* **9**, 671-675 (2012). https://doi.org:10.1038/nmeth.2089

Shrestha, A. M. & Frith, M. C. An approximate Bayesian approach for mapping paired-end DNA reads to a reference genome. *Bioinformatics* **29**, 965-972 (2013). https://doi.org:10.1093/bioinformatics/btt073

Krzywinski, M. *et al.* Circos: an information aesthetic for comparative genomics. *Genome Research* **19**, 1639-1645 (2009). https://doi.org:10.1101/gr.092759.109

Data availability

ONT sequencing data for this study have been deposited in the European Nucleotide Archive (ENA) at EMBL-EBI under accession number PRJEB80380 (https://www.ebi.ac.uk/ena/browser/view/PRJEB80380). Source data is available here.

Code availability

Codes for mathematical modeling are available at https://github.com/anais-rat/telomeres. Other codes are available here.

Acknowledgements

We wish to thank E. Fabre and V. Borde for sharing technical advice, Z. Xu for first mathematical simulations and fluctuation assay code and M. Godinho Ferreira and C. Azzalin for fruitful discussion. We also acknowledge J. Haber, H. Ulrich and M. Lisby for strains and plasmids. We also thank the Teixeira lab, numerous trainees and the UMR8226 unit members for technical support and fruitful discussions. We also thank T. Weinert, D. Kappei, E. Fabre and M. Bao (Life Science Editors) for critical reading of the manuscript. During the preparation of this work, the authors used ChatGPT version 3.5 to enhance the clarity and fluency of the English writing. After using this tool, the authors reviewed and edited the content as needed and take full responsibility for the content of the publication. PB was supported by a fellowship from the Ligue Contre le Cancer (France), FRRB was supported by a "Consejo Nacional de Ciencia y Tecnología" (CONACYT) scholarship and KC was supported by a fellowship from "Fondation pour la Recherche Médicale" (FRM). Work in MC lab was supported by an Open Competition M-2 grant from the Dutch Research Council. Work in MTT lab was supported by the "Fondation de la Recherche Medicale" ("équipe labellisée" FRM), by the French National Research Agency (ANR) as part of the "Investissements d'Avenir" Program (LabEx Dynamo ANR-11-LABX-0011-01) and ANR-16-CE12-0026, and the "Institut National du Cancer (PLBIO20-312).

Author contributions

Conceptualization Ideas; formulation or evolution of overarching research goals and aims: PB (ExtFig.2), VMF (Fig.4-5, ExtFig.4-5), RL (Fig.1,3, ExtFig.1-3), SM (Fig1,3, ExtFig.1,3), MD (Fig.3, ExtFig.3), MC (ExtFig.3), MTT (Fig.1-5, ExtFig.1-5);Methodology Development or design of methodology; creation of models: PB (Fig.1,3, ExtFig.1-3), VMF (Fig.2, 4-5, ExtFig.2, 4-5), AR (Fig.3, ExtFig.3), FRB (ExtFig.3), PJ (Fig.2, ExtFig.2), RL (Fig.1,3, ExtFig.1-3), SM (Fig1,3, ExtFig.1-3), TA (Fig.3), BZ (ExtFig.3), KC (ExtFig.4), HGK (ExtFig.3), GC (Fig.3, ExtFig.3), MD (Fig.3, ExtFig.3), MC (ExtFig.3), MTT (Fig.1-2 , ExtFig.1-2); Investigation Conducting a research and investigation process, specifically performing the experiments, or data/evidence collection: PB (Fig.1-3 , ExtFig.1-3), VMF (Fig.2, 4-5, ExtFig.2, 4-5), AR (Fig.3, ExtFig.3), FRB (ExtFig.3), PJ (Fig.2, ExtFig.2), BZ (ExtFig.3), KC (ExtFig.4), HGK (ExtFig.3), MD (Fig.3, ExtFig.3), MTT (Fig.2, ExtFig.2); Data curation Management activities to annotate (produce metadata), scrub data and maintain research data (including software code, where it is necessary for interpreting the data itself) for initial use and later re-use: PB (Fig1-3, ExtFig.1-3), VMF (Fig.4-5, ExtFig.4-5), AR (Fig.3, ExtFig.3), FRB (ExtFig.3), PJ (Fig.2, ExtFig.2), BZ (ExtFig.3), MD (Fig.3, ExtFig.3), MTT (Fig.2, ExtFig.2); Formal Analysis Application of statistical, mathematical, computational, or other formal techniques to analyse or synthesize study data: PB (Fig1-3, ExtFig.1-3), VMF (Fig.4-5, ExtFig.4-5), AR (Fig.3, ExtFig.3), FRB (ExtFig.3), PJ (Fig.2, ExtFig.2), BZ (ExtFig.3), MD (Fig.3, ExtFig.3), MTT (Fig.2, ExtFig.2); Software Programming, software development; designing computer programs; implementation of the computer code and supporting algorithms; testing of existing code components: VMF (Fig.4-5, ExtFig.4-5), AR (Fig.3, ExtFig.3), PJ (Fig.2, ExtFig.2), AM (Fig.2,5, ExtFig.2,5), GC (Fig.3, ExtFig.3), MD (Fig.3, ExtFig.3), MTT (Fig.2, ExtFig.2); Validation Verification, whether as a part of the activity or separate, of the overall replication/reproducibility of results/experiments and other research outputs: PB (Fig1,3, ExtFig.1-3), VMF (Fig.4-5, ExtFig.4-5), AR (Fig.3, ExtFig.3), FRB (ExtFig.3), PJ (Fig.2, ExtFig.2), KC (ExtFig.4), MD (Fig.3, ExtFig.3), MTT (Fig.1-5, ExtFig.1-5); Visualization Preparation, creation and/or presentation of the published work, specifically visualization/data presentation. PB (Fig1,3, ExtFig.1-3), VMF (Fig.4-5, ExtFig.4-5), AR (Fig.3, ExtFig.3), PJ (Fig.2, ExtFig.2), AM (Fig5, ExtFig.5), MD (Fig.3, ExtFig.3), MC (ExtFig.3), MTT (Fig.ExtFig.2,3); Funding acquisition Acquisition of the financial support for the project leading to this publication. PB (Fig1-3, ExtFig.1-3), GC (Fig.3, ExtFig.3), MD (Fig.3, ExtFig.3), MC (ExtFig.3), MTT (Fig.1-5, ExtFig.1-5); Resources Provision of study materials, reagents, materials, patients, laboratory samples, animals, instrumentation, computing resources, or other analysis tools. VMF (Fig.4-5, ExtFig.4-5), AR (Fig.3, ExtFig.3), GC (Fig.3, ExtFig.3), MD (Fig.3, ExtFig.3), MC (ExtFig.3), MTT (Fig.1-5, ExtFig.1-5); Project administration Management and coordination responsibility for the research activity planning and

execution. VMF (Fig.4-5, ExtFig.4-5), RL (Fig.1-3, ExtFig.1-3), SM (Fig.1-3, ExtFig.1-3), GC (Fig.3, ExtFig.3), MD (Fig.3, ExtFig.3), MC (ExtFig.3), MTT (Fig.1-5, ExtFig.1-5); Supervision Oversight and leadership responsibility for the research activity planning and execution, including mentorship external to the core team. VMF (Fig.4-5, ExtFig.4-5), RL (Fig.1-3, ExtFig.1-3), SM (Fig.1-3, ExtFig.1-3), GC (Fig.3, ExtFig.3), MD (Fig.3, ExtFig.3), MC (ExtFig.3), MTT (Fig.1-5, ExtFig.1-5); Writing – original draft Preparation, creation and/or presentation of the published work, specifically writing the initial draft: PB, VMF, SM, MTT; Writing – review & editing Preparation, creation and/or presentation of the published work by those from the original research group, specifically critical review, commentary or revision – including pre- or post-publication stages. PB, VMF, AR, FRB, PJ, RL, SM, AM, TA, BZ, KC, HGK, GC, MD, MC, MTT.

Competing interests

The authors declare no competing interests.

Fig.1: the FinalCut system efficiently generates a unique telomere of defined length. a, *The* FinalCut *construct at telomere 6R: upon induction of Cas9 and association with gRNA, a single telomere in cell containing the complementary sequence of gRNA, ttDNA, is cut to generate a telomere of defined length FCx.* **b**, *Experimental procedure.* **c**, *Southern blot to detect the* Hpa *I-terminal restriction fragment of Chr6R. Loading control: internal locus in chromosome 11.* **d**, *Uncut and cut bands normalized to loading control and to uncut signal measured at t=0h. Bars correspond to average and error bars to SD of 3 independent experiments.*

***Fig.2: Rapid degradation of telomeres below 40 bp is preceded by progressive shortening by the DNA end replication problem. a**, Experimental design to study the outcome of FinalCut telomeres. FinalCut strains are precultured in Raffinose media containing or not Doxycycline (t=-4h) to control telomerase activity. Galactose is then added to induce Cas9-AID gene for 4 hours (t= 0 h). Cells are subsequently diluted in Glucose media containing NAA to degrade Cas9-AID. Samples are taken at indicated time points. **b**, Changes in lengths of individual indicated FinalCut telomeres over population doublings after Cas9 shut off. Data corresponds to 3 pooled independent experiment (FC50: N=2769, 4595, 1538 for t=0h; N=2589, 8840, 2954 for t=2h; N=4838, 6548, 7594 for t=4h; FC70: N=1119, 1109, 764 for t=0h; N=819, 831, 1416 for t=2h; N=1102, 1018, 745 for t=4h. Bars correspond to SD of pools. **c**, Magnified view of **b**, showing the distribution of individual telomere lengths at nucleotide resolution, over time and population doublings after Cas9 deactivation in the FC50 strain, without telomerase (+Dox). Dots represent the average from three independent experiments, with error bars indicating the standard deviation (SD) of the averages. Sequence of uncut FC50 telomere is indicated, as well as expected Cas9 cleavage.*

Fig.3: A length threshold of FinalCut telomere determines cell proliferation capacity. a, Microfluidics experimental setting. FinalCut strains are introduced in a microfluidics circuit and observed using microscopy. Cells are allowed to invade microcavities through cell divisions. Doxycycline is added to inactivate telomerase up to the end the experiment. Galactose is next added for 6 hours to induce Cas9. Durations between two cell divisions, namely between two budding events, are recorded, for consecutive cell divisions up to cell death. Colors indicate whether the cell cycle is normal (green) or abnormally long (red). **b**, Display of consecutive cell cycle durations of indicated FinalCut cell lineages. Generation 0 corresponds to the addition of Galactose. Each horizontal line corresponds to an individual cell lineage, and each segment to a cell cycle. Cell cycle duration (in minutes) is indicated by the color bar. X at the end of the lineage indicates cell death, whereas an ellipsis (...) indicates that the cell was alive at the end of the experiment. Data shown corresponds to the sum of at least two independent experiments. **c**, Mathematical model of telomerase-negative individual cells as grown in microfluidic experiments, defining Type A cells and Type B cells (Extended Data Fig.3b). Inset: Simplified tree diagram of the mathematical model. It indicates the several fates of each cell type and respective probability rates at each cell division. The colors of the squares indicate whether the cell cycle is normal (green) or transiently abnormally long (orange) or irreversibly abnormally long (red). The graph represents the best fits of indicated probability rates as a function of the length of the shortest telomere in the cell being the TEL6R telomere. Parameters are listed in Supplementary Table 7. **d**, Lifespan of individual cell lineages entering senescence upon telomerase inactivation, as assessed by microfluidics in **b** and 100 simulations of the same number of individual cell lineages with the mathematical model in **c**.

Fig.4: Genome instability in replicative senescence stems from the first telomere reaching a critical short length. a-b, *Relevant genomic features of main strains used to quantify genomic instability in* cis (**b**: yT1760) or trans (**a**: yT1690) *of* FC70 *telomere by fluctuation assays. Last essential gene of chromosome 5 left arm is* PCM1/YEL058W; *last essential gene of chromosome 6 right arm is* RPN12/YFR052W. *Not to scale.* **c**, *Experimental protocol for the fluctuation assay in cells harboring a FC70 telomere. Multiple cultures are grown in parallel in which telomerase is inactivated by doxycycline addition, followed by carbon source switch to galactose to induce FC70. Then, a portion of each culture is plated at t=0 h and t=24 h after Cas9 induction on canavanine-containing glucose media lacking doxycycline.This allows for Cas9 shut off, telomerase re-expression to stabilize the genome and selection of mutants at CAN1 locus. Can*[R] *colonies number distribution is used to estimate mutation rate according to fluctuation assay mathematical models. Can*[R] *cells are then replica-plated on plates containing 5-FOA to score the fraction of GCRs or point mutations.* **d-e**, *Mutation rate of* CAN1 *gene placed in* trans (**d**) *or in* cis (**e**) *of* FC70 *calculated from cells plated at t=0 h and 24 h after Cas9 induction (N=10-19). Data are represented as the mutation rate ±95% confidence interval. Mutation type distribution of Can*[R] *colonies obtained at t=0 h and t=24 h after Cas9 induction is indicated. See Supplementary Table 9 for statistical details. Strain used in* **d**: yT1690 *(FC70) and* yT1694 *(noFC). Strains used in* **e**: yT1760 *(FC70) and* yT1759 *(noFC).*

Fig.5: Genomic instability at the shortest telomere in the cell is characterized by non-reciprocal translocations enabling senescence escape. a, CIRCOS plot showing the genome of a representative Can^R ura$^-$ colony classified in this study as GCR, obtained after 24 h of Cas9 induction in the absence of telomerase of cells harboring a FC70 telomere, the CAN1-URA3 cassette at 5 kb from the last essential gene on chromosome 6R and in a can1-Δ backgound (initial strain: yT1760, Fig.4b; experiment Fig4e). The outer ring represents reference S.cerevisiae chromosomes. Subtelomeric regions are displayed in blue. Centromeres are represented by a black line. Chromosome coordinates are indicated in Mbp. The inner ring displays the read-depth, ranging from 0x to 30x. Extremities of splitted reads are shown as red bridged lines. **b**, Zoom of indicated region of (a) showing loci involved in the breakpoint region, including HXK1/2 in bold. The solid red line represents the longest splitted read. Red dots indicate the extremities of splitted reads. (*) Reads containing exclusively TG$_{1-3}$, which map telomeric regions indiscriminately. (**) Reads mapping a region of the CAN1-URA3 cassette (in grey) containing a short plasmid-derived sequence (60 bp), also present at the AFB2-TRP1 locus. Initial strain: yT1760. N=3. **c**, Proposed model for the chromosomal rearrangement involving chromosomes 6 and 7 of Can^R. See Extended Data Fig.5 for details.