Nemo-like kinase disrupts nuclear import and drives TDP43 mislocalization in ALS

Michael E. Bekier II[1#], Emile Pinarbasi[1,2#], Jack J. Mesojedec[1], Layla Ghaffari[3], Martina de Majo[3], Erik Ullian[4], Mark Koontz[3], Sarah Coleman[3], Xingli Li[1], Elizabeth M. H. Tank[1], Jacob Waksmacki[1], Sami Barmada[1]*

[1]Department of Neurology, University of Michigan, Ann Arbor, MI, 48109, United States

[2]Department of Pathology, Michigan Medicine, University of Michigan, Ann Arbor, MI, 48109, United States

[3]Synapticure, Chicago, IL, United States 60612

[4]Department of Ophthalmology & Physiology, University of California San Francisco, San Francisco, CA, 94143, United States

# contributed equally

*Correspondence to Sami Barmada: 109 Zina Pitcher Place, Ann Arbor, MI, 48019; 734-764-8425;

sbarmada@umich.edu

Conflict of Interest Statement:

S.J.B. serves on the advisory board for Neurocures, Inc., Symbiosis, Eikonizo Therapeutics, 15 Ninesquare Therapeutics, the Live Like Lou Foundation, Synapticure, and the Robert Packard

21   Center for ALS Research. S.J.B. has received research funding from Denali Therapeutics, Biogen,

22   Inc., Lysoway Therapeutics, Amylyx Therapeutics, Acelot Therapeutics, Meira GTX, Inc., Prevail

23   Therapeutics, Eikonizo Therapeutics, and Ninesquare Therapeutics.

Abstract

Cytoplasmic TDP43 mislocalization and aggregation are pathological hallmarks of amyotrophic lateral sclerosis (ALS). However, the initial cellular insults that lead to TDP43 mislocalization remain unclear. In this study, we demonstrate that Nemo-like kinase (NLK)—a proline-directed serine/threonine kinase—promotes the mislocalization of TDP43 and other RNA-binding proteins by disrupting nuclear import. NLK levels are selectively elevated in neurons exhibiting TDP43 mislocalization in ALS patient tissues, while genetic reduction of *NLK* reduces toxicity in human neuron models of ALS. Our findings suggest that NLK is a promising therapeutic target for neurodegenerative diseases.

Introduction

Amyotrophic lateral sclerosis (ALS) is a fatal and progressive neurodegenerative disease distinguished by loss of motor neurons in the cortex and spinal cord (1). The clinical presentation of ALS is heterogeneous, depending in large part upon the neuroanatomical pathways involved. Despite this, a single pathological hallmark — mislocalization and aggregation of the RNA binding protein TDP43 — is found in >95% of individuals (2).

TDP43 is crucial for several aspects of RNA processing, including RNA splicing, stability and transport (3-7). TDP43 is primarily nuclear in healthy cells, in contrast to the nuclear exclusion and cytosolic deposition of TDP43 that are characteristic of ALS (3). Although the origins of TDP43 mislocalization are unclear, deficiencies in nucleocytoplasmic transport machinery are increasingly implicated as a potential contributing factor to ALS pathology (8). Impaired nuclear import of TDP43 and other RNA binding proteins, together with dysfunction of the nuclear pore complex itself, are observed in human induced pluripotent stem cell (iPSC)-derived neurons from sporadic and familial ALS patients as well as post-mortem samples. Genetic screens in ALS model systems have repeatedly uncovered nuclear pore and nucleocytoplasmic transport factors as potent disease modifiers, confirming the significance of these pathways in disease pathogenesis (9-11).

Among the most consistent and dramatic disease modifiers to emerge from these screens is nemo-like kinase (NLK), a proline directed serine-threonine kinase that regulates cell differentiation (12), proliferation, and apoptosis (13, 14). NLK depletion mitigates disease phenotypes not just in *Drosophila* and murine models of ALS, but also murine models of spinobulbar muscular atrophy, and murine models of spinocerebellar ataxia (15-17). Conversely,

NLK overexpression reduces toxicity in Huntington's disease models, indicating context-dependent

effects of NLK in neurodegeneration (18).

NLK has several predicted substrates that function in nucleocytoplasmic transport, but whether

NLK itself may influence nuclear import and ALS pathogenesis remains unknown. Here, we

demonstrate that NLK is a pivotal regulator of nucleocytoplasmic transport. NLK accumulation

leads directly to the cytoplasmic accumulations of TDP43 and other RNA-binding proteins

associated with ALS, while reduction in NLK promotes the survival of iPSC-derived neurons

carrying ALS-associated mutations. Importantly, NLK is upregulated selectively in affected neurons

from ALS patients and correlates with TDP43 mislocalization, implicating NLK as a key determinant

of disease pathophysiology and highlighting NLK as a promising therapeutic target for ALS and

related TDP43 proteinopathies.

Results

NLK overexpression drives mislocalization of TDP43 and other ALS-linked RNA-binding

proteins

To determine if NLK is capable of directly influencing TDP43 localization, we transfected HEK293

cells with plasmids encoding FLAG-tagged wild-type (WT) NLK and examined TDP43 localization by

immunofluorescence (Figure 1A). Exogenous NLK, like endogenous NLK, is both cytoplasmic and

nuclear in distribution (19). As a control, we also transfected cells with plasmids encoding FLAG-

tagged kinase-negative (KN) NLK, which harbors a point mutation in the active site (K155M) that

abrogates kinase activity (19). NLK WT overexpression significantly decreased the nuclear-

cytoplasmic (NC) ratio of endogenous TDP43 compared to NLK KN (Figure 1B). The change in

TDP43 NC ratio was driven by both a reduction in nuclear TDP43 as well as a concordant increase

in cytoplasmic TDP43, consistent with nuclear-cytoplasmic TDP43 mislocalization (Figure 1C-D).

We next asked whether NLK WT solely affected TDP43 localization, or whether it also regulated the

distribution of other ALS-associated RNA binding proteins that undergo nucleocytoplasmic

transport, including HNRNPA2B1 and Matrin-3. We also examined the localization of FUS, an RNA

binding protein harboring a non-classical PY-nuclear localization signal (NLS) recognized by

importin-β_2, in contrast to the classical K/R-rich NLS in TDP43, HNRNPA2B1, and MATR3 that binds

importin-α (20-22). As before, HEK293 cells were transfected with KN or WT NLK, and the

subcellular distribution of each protein determined by immunofluorescence (Figure 1F-K).

Compared to NLK KN, NLK WT overexpression significantly reduced the NC ratio of both FUS and

HNRNPA2B1 but had no significant effect on Matrin-3 localization (Figure 1F-K). Together, these

findings suggest that WT NLK overexpression disrupts nuclear localization of multiple RNA-binding

proteins harboring classical as well as non-classical NLSs, in a kinase-dependent manner.

NLK overexpression disrupts nuclear import

At steady state, nuclear localization of TDP43 and other RNA-binding proteins is maintained

through two competing processes: active nuclear import and passive efflux through the nuclear

pore (23-25). To directly evaluate nuclear import, we co-expressed NLK together with YFP fusion

proteins containing NLS sequences from TDP43 (YFP-NLS[TDP]), FUS (YFP-NLS[FUS]), Matrin-3 (YFP-

NLS[MATR3]) and SV40 (YFP-NLS[SV40], a canonical classical NLS) followed by immunofluorescence for

TDP43, FUS, or Matrin-3 (Figures 2A-H). Notably, WT NLK expression drives mislocalization of YFP-

NLS[MATR3], but not endogenous Matrin-3 (Figure 1J, K), potentially due to the relatively large size of

Matrin-3 (95kDa, vs 27kDa for YFP-NLS[MATR3]) (23, 24). Compared to overexpression of NLK KN, NLK

WT significantly increased the NC ratio of all NLS-fusion proteins (Figure 2C). Similar to what we

observed for native RNA binding proteins (Figure 1), WT NLK affects reporters containing both

classical as well as non-classical NLS motifs (26). Collectively, these data indicate that WT NLK

overexpression disrupts global nuclear import through its kinase activity.

NLK-induced TDP43 mislocalization is independent of KPNA2 nuclear accumulation

Nuclear import relies on transport receptor proteins, such as KPNA2 and KPNB1, which recognize

and bind to NLS-containing proteins such as TDP43 (27). Thus, we examined the sub-cellular

localization of KPNB1 and KPNA2 by immunofluorescence after overexpression of either KN or WT

NLK. Overexpression of WT NLK significantly increased the NC ratio of both KPNA2 and KPNB11,

driven by a significant increase in their nuclear fractions without a corresponding decrease in their

cytoplasmic abundance (Figure 3A-D, S1A-B). Because of the critical importance of these factors

for nuclear import, we questioned whether the mislocalization of TDP43 and other RNA binding

proteins may be secondary to the observed nuclear accumulation of KPNA2 and/or KPNB1. To test

this hypothesis, we took advantage of previous data showing that nuclear accumulation of KPNA2

can be reversed by the expression of the E3 ubiquitin ligase FBXW7 (28). HEK293 cells were

transfected with plasmids encoding FLAG-tagged NLK WT and either mApple (negative control) or

FBXW7 followed by immunofluorescence for KPNA2 (Figure 3E, F). Compared to mApple, FBXW7-

V5 significantly reduced the NC ratio of KPNA2 in NLK-overexpressing cells (Figure 3G), due

primarily to reduction in the nuclear KPNA2signal (Figure 3D, S1C). Despite this, FBXW7-V5

coexpression failed to significantly correct TDP43 mislocalization in cells transfected with WT NLK

(Figure 3G, H, S1D). These results indicate that NLK-induced mislocalization of TDP43 does not

depend on the nuclear accumulation of KPNA2.

NLK overexpression promotes mislocalization of Ran, Ran-GAP1, and RanBP2

Nuclear localization of receptor-bound cargo is mediated by RanGAP1 and RanBP2, nuclear pore-

associated factors that regulate the Ran gradient (29)(Figure S2). Consistent with a previous screen

for kinase-interacting proteins (30), both RanGAP1 and RanBP2 co-immunoprecipitated with FLAG-

NLK in HEK293 cells (Figure 4A-C). NLK overexpression also promoted the accumulation of non-

sumoylated RanGAP1 (Figure 4A). RanGAP1 sumoylation is critical for nuclear envelope

localization and its interaction with RanBP2 (Figure 4 C, (31-33)). This prompted us to investigate

the impact of NLK overexpression on the subcellular localization of RanGAP1, RanBP2, and

ultimately Ran. Compared to KN NLK overexpression, WT NLK significantly disrupted the expected

nuclear envelope localization of RanGAP1 and RanBP2 as measured by the ratio of nuclear rim

density to cytoplasmic density (Figure 4D-G). Conversely, overexpression of NLK WT did not

significantly impact FG nucleoporins as detected by MAb414 (Figure 4H, I). WT NLK overexpression

also reduced the NC ratio of Ran, driven by both a decrease in nuclear Ran and an increase in

cytoplasmic Ran (Figure 4J, K). As such, NLK-induced disruption of the Ran gradient correlates with

disruption of the RanGAP1-RanBP2 complex and impaired nucleocytoplasmic transport.

NLK drives redistribution of mRNA and disassembles nuclear speckles

The localization of TDP43 and other RNA-binding proteins is heavily influenced not just by their NLS

motifs, but also by their cognate RNA substrates (34). Based on this, we examined whether TDP43

mislocalization upon WT NLK overexpression is RNA-dependent. Initially, we transfected HEK293

cells with an EGFP-tagged variant of TDP43 harboring two mutations within RRM1 that abolish RNA

binding, TDPF2L-EGFP (F147L/F149L)(35, 36) , together with WT or KN NLK (Figure 5A). TDPF2L-

EGFP formed phase-separated droplets that were largely restricted to the nucleus in cells co-

expressing KN NLK, as in prior studies (37). However, co-transfection with WT NLK resulted in the

appearance of cytosolic TDP43F2L-EGFP droplets, suggesting that TDP43 mislocalization upon WT

NLK overexpression is independent of RNA binding (Figure 5A and S3A).

We also investigated whether WT NLK overexpression could affect the distribution of

polyadenylated (polyA) mRNA. We first immunostained for NXF1, an mRNA export factor which

itself contains a PY-NLS (20) and saw that in NLK WT transfected cells, NXF1 accumulates in the

cytoplasm (Figure 5B and S3B). Next, we directly assessed mRNA distribution using fluorescence

in-situ hybridization (FISH). While untransfected and KN NLK-expressing cells displayed a punctate

pattern of polyA mRNA within the nucleus, WT NLK overexpression resulted in a more diffuse and

evenly distributed nuclear signal, and minimal changes in the polyA mRNA NC ratio (Figure 5C and

S4C). Given the enrichment of polyA mRNA within nuclear speckles, and the apparent loss of such

structures with WT NLK overexpression, we also immunostained WT or KN NLK-transfected cells

using the SC35 antibody, which recognizes SRRM2, a core component of nuclear speckles (38). WT

NLK-expressing cells, in contrast to untransfected or KN NLK-transfected cells, displayed a

dramatic reduction in nuclear speckles (Figure 5D). This effect appeared to be specific for nuclear

speckles, as we saw no change in other nuclear membraneless organelles such as paraspeckles

(marked by SFPQ (39); Figure 5E) or nucleoli (marked by nucleophosmin (40); Figure 5F) in WT NLK-

expressing cells.

NLK overexpression disrupts nuclear import in mammalian neurons

To examine the impact of NLK overexpression on RBPs and nuclear import factors in neurons, we

transfected rodent primary mixed cortical neurons with either SNAP-FLAG (SF; negative control) or

SNAP-FLAG-NLK WT (SF-NLK) before immunostaining for TDP43 and other factors affected by NLK.

Consistent with our results in HEK293 cells, expression of SF-NLK but not SF significantly impacted

the subcellular localization of TDP43, FUS, RanGAP1, RanBP2, and Ran (Figure 6A-D). As before,

the central channel of the NPC marked by FG nucleoporins was unaffected by SF-NLK expression

(Figure 6E), while exogenous reporters such as YFP-NLS[SV40] were mislocalized by SF-NLK but not SF

in transfected neurons (Figure S4).

Nucleocytoplasmic trafficking is essential for maintaining protein and RNA homeostasis; thus, we

predicted that NLK-induced disruption of nucleocytoplasmic transport would lead to substantial

toxicity. To assess this, we utilized automated longitudinal microscopy to track hundreds of rodent

primary mixed cortical neurons cultures prospectively for 10 days in culture. Neurons were

transfected with SF or SF-NLK, in combination with a survival marker (GFP) enabling us to

determine the time of cell death (Figure 6F, G). SF-NLK overexpression significantly increased the

cumulative risk of death in transfected neurons compared to SF alone (HR=1.61, p<0.001, Cox

proportional hazards analysis; Figure 6H). Since all data on survival are acquired from individual

neurons, and the abundance of fluorescently-tagged proteins is directly proportional to the

measured signal intensity (41), we investigated the relationship between SNAP-FLAG-NLK intensity

and the hazard of the event using a Cox proportional hazards model with penalized splines (42-44).

The Cox model revealed significant non-linear relationships between signal intensity and hazard,

as indicated by a Likelihood Ratio Test statistic of χ^2 (4.06) = 25.89 (p < 0.0001), suggesting that the

model fits the data well. The concordance index was 0.587, indicating a moderate ability of the

model to discriminate between different hazard levels. Hazard ratios were calculated for distinct

intensity segments derived from the spline model. For low signal intensities, the hazard ratio was

0.898 (95% CI: 0.378 - 2.132), indicating a slight decrease in hazard compared to the baseline. In

the medium intensity range, the hazard ratio increased to 1.555 (95% CI: 0.532 - 4.545), suggesting

a potential increase in risk. Notably, for high signal intensities, the hazard ratio rose significantly to

4.629 (95% CI: 0.701 - 30.584), reflecting a substantial elevation in hazard associated with

increased signal intensity (Figure 6I). Collectively, our data indicate that overexpression of NLK

significantly impairs the mechanisms that regulate nucleocytoplasmic transport, leading to the

mislocalization of pertinent RNA-binding proteins and ultimately neuron death.

Increased NLK levels correlate with TDP43 pathology in disease models and patients

Approximately 50% of individuals with frontotemporal lobar degeneration (FTLD) show TDP43

mislocalization as in ALS(3). In addition, up to half of people with ALS demonstrate cognitive

impairment reminiscent of FTLD, while ~1/3 of those with FTLD show motor neuron disease that is

indistinguishable from ALS (3, 45). These observations, as well as shared genetics underlying both

ALS and FTD, testify to the close overlap between ALS and FTLD with TDP43 pathology (FTLD-TDP)

(3, 45). Therefore, to determine if NLK dysregulation may be involved in ALS/FTLD-TDP disease

pathogenesis, we initially investigated *NLK* expression in *GRN* knockout mature brain organoids

(mbOrgs), an FTLD-TDP model that recapitulates key pathological features of disease, including

TDP43 mislocalization, phosphorylated TDP43, and characteristic missplicing of TDP43 substrate

RNAs (46). Neurogenin-2 inducible cortical neurons (iNeurons) and mature cortical astrocytes

(iAstrocytes) derived from isogenic wild-type (WT) or GRN-/- iPSCs were combined in fixed ratios to

form mbOrgs (Figure 7A). RNA-seq revealed significantly elevated normalized counts of *NLK* in

GRN-/- mbOrgs compared to WT controls (Figure 7B), a finding that was also confirmed by

quantitative RT-PCR (qRT-PCR) (Figure 7C).

To further explore the link between NLK changes and TDP43 pathology, we turned to a unique

dataset generated by Liu et al. in which neuronal nuclei were sorted from frontal cortices of FTLD-

TDP patients into two populations — those with and without nuclear TDP43 – prior to RNA

sequencing (RNA-seq) (30). Reanalysis of these data demonstrated a significant upregulation of

NLK mRNA in nuclei lacking TDP43 (Figure 7D). Using dual-immunohistochemistry, we confirmed

that spinal neurons from ALS spinal cord sections with TDP43 pathology (nuclear loss of TDP43

with cytosolic inclusions) exhibited more intense staining for NLK in comparison to unaffected

neurons present in the same section (Figure 7E and S5C). Dual staining for NLK and TDP43 was

also performed in sections from four control patients without spinal pathology (Figure S5D5),

showing no clear differences in NLK abundance. These results demonstrate a clear relationship

between elevated NLK, at both the mRNA and protein levels, and TDP43 mislocalization in FTLD-

TDP and ALS.

Together, these data show that NLK is upregulated in human patients and in disease models

featuring TDP43 pathology, in accord with our data demonstrating TDP43 mislocalization upon NLK

overexpression.

NLK reduction improves survival in iPSC-derived neuron models of ALS

Given the detrimental effects of NLK overexpression on nucleocytoplasmic transport (Figures 1-6)

and neuron survival (Figure 6), and the elevated *NLK* mRNA and protein observed in patients and

disease models (Figure 7A-E), we asked whether targeting *NLK* could ameliorate disease

phenotypes in ALS/FTLD-TDP models. First, we examined the survival of iPSC-derived neurons

carrying *C9ORF72* expansions, the most prevalent mutation underlying familial ALS and FTLD-TDP

in Europe and North America (47, 48). Non-disease and *C9ORF72* mutant neurons were

transduced with virus encoding non-targeting or *NLK* shRNA, resulting in a ~50% reduction in NLK

RNA levels compared to non-targeting shRNA (Figure 7F). Individual neurons were followed by

automated microscopy for 10 days, as before (Figure 7G), and differences in survival assessed via

Cox proportional hazards analysis. Three separate lines of *C9ORF72* mutant neurons exhibited

significantly higher cumulative risks of death compared to unrelated non-disease neurons (Figure

7H and S6A-C). Transduction with *NLK* shRNA significantly reduced the cumulative risk of death in

all three lines of *C9ORF72* neurons (HR= 0.403, p= 7.71e-50). To examine if NLK reduction is

neuroprotective outside of *C9ORF72* mutations, we utilized an isogenic pair of *TARDBP* mutant

iPSC-derived neurons that were created by CRISPR/Cas9 genome engineering (49). In comparison

to isogenic controls ("WT"), *TARDBP* mutant (M337V) neurons transduced with lentivirus

expressing non-targeting shRNA exhibited a significantly higher cumulative risk of death (Figure 7I).

As with *C9ORF72* mutant neurons, however, transduction with *NLK* shRNA-expressing virus

significantly extended the survival of M337V neurons (HR = 0.227, p= 0.0013). Collectively, these

data imply that *NLK* overexpression is toxic, while *NLK* reduction promotes neuronal survival in

models of ALS and FTLD-TDP.

Discussion

Our findings indicate that NLK overexpression disrupts the nuclear import and localization of

TDP43 and related RNA-binding proteins, including FUS and HNRNPA2B1, in a kinase-dependent

manner. These effects correlate with the mislocalization of the RanBP2-RanGAP1 complex and

collapse of the Ran gradient, both of which are crucial for functional nucleocytoplasmic transport.

As expected based on these observations, NLK dose-dependently increased the risk of death in

primary rodent neurons. Notably, we uncovered elevated NLK expression selectively in neurons

with TDP-43 pathology both at the RNA level (in FTLD-TDP frontal cortex) and at the protein level (in

ALS spinal cord). NLK was also upregulated in a brain organoid model of FTLD-TDP, while reducing

NLK in human iPSC-derived neurons carrying disease-associated mutations prolonged survival. These results suggest that NLK may contribute to the pathology of ALS-FTLD-TDP by impairing nucleocytoplasmic transport and promoting neurotoxicity. Targeting NLK protein levels or kinase activity could thus represent a novel therapeutic approach for ALS, FTLD-TDP and other TDP43 proteinopathies.

Disruption of the Ran gradient, mislocalization of nuclear pore components, and disturbance of nuclear pore architecture — all of which we observed upon NLK overexpression — have likewise been described in ALS/FTLD-TDP samples and disease models (9-11, 50). There is no predicted NLK phosphorylation site within TDP43 itself; rather, we suspect that NLK may interfere with the nucleocytoplasmic transport of several RNA binding proteins and other factors by acting on integral components of the nuclear pore. We and others noted a direct interaction between NLK and the RanGAP1-RanBP2 complex ((30); see Figure 4), which is critical for maintaining the Ran gradient and nuclear import/export (51). NLK overexpression prevented RanGAP1 sumoylation (Figure 4A) and reduced the amount of RanGAP1 in complex with RanBP2 (Figure 4C). In the absence of sumoylation, RanGAP1 is unable to bind RanBP2 or localize to the nuclear envelope (Figure 4C, (31-33). One possibility is that NLK directly phosphorylates RanGAP1, inhibiting its GAP activity (50) as well as its sumoylation, thereby disrupting its interaction with RanBP2 at the nuclear pore complex. Alternatively, NLK may negatively regulate Ubc9, a SUMO E3 ligase required for recruitment of RanGAP1 to the RanBP2 complex. None of these possibilities are mutually exclusive; NLK may impact nucleocytoplasmic transport through more than one overlapping mechanism.

At baseline, NLK is highly expressed in the CNS, and its expression is unregulated by oxidative and osmotic stress (52, 53), conditions associated with the cytosolic accumulation of TDP43 and other, predominantly nuclear, RNA binding proteins. NLK interacts with several proteins associated with

neurodegenerative diseases, including poly-Q expanded androgen receptor in spinobulbar

muscular atrophy (SBMA) and ataxin-1 in spinocerebellar ataxia 1 (SCA1) (16, 17); accumulations

of these proteins may also contribute to changes in NLK expression and/or activity. Although

previous studies have failed to detect NLK upregulation in disease, in most cases these

investigations are limited to evaluation of NLK levels (RNA or protein) in bulk tissue. In contrast, our

work revealed NLK upregulation solely within affected neurons displaying TDP43 pathology, in

association with evidence of disrupted nucleocytoplasmic trafficking in the same cells, suggesting

that NLK expression changes are restricted to cells with TDP43 redistribution. Outside of changes

in expression, NLK may also be inappropriately activated in disease via diverse stimuli.

Proinflammatory factors such as transforming growth factor ß (TGFß), interleukin-6 (IL6) and Wnt

all activate NLK through MAPK-dependent signaling (54). Notably, TGF-ß is upregulated in ALS (55),

and aberrant Wnt signaling is associated with TDP43 mislocalization in cellular and animal models

(56).

NLK is an essential kinase — homozygous *Nlk* deletion results in death *in utero* or postnatally,

depending on the mouse strain (57, 58). Conditional *Nlk* knockout in adult animals is well-

tolerated, however (59), and *Nlk* haploinsufficiency (*Nlk+/-*) is protective in several

neurodegenerative disease models, including TDP43-overexpressing mice (15-18). Similarly, partial

NLK knockdown in our investigations was associated with extended survival in *C9ORF72* and

TARDBP mutant human iPSC-derived neurons and was safe in control neurons. These data indicate

that even modest reductions in *NLK* may be sufficient for mitigating neurodegeneration in ALS and

FTLD-TDP.

Although prior interaction studies highlighted several potential partners for NLK, its substrates in

neurons and phosphorylation sites within these substrates remain largely unexplored. RanBP2 and

RanGAP1 are two potential targets for NLK with clear connections to nucleocytoplasmic transport,

but NLK is also likely to act on distinct substrates and disease-related pathways, including the

integrated stress response and autophagy (15). One advantage of therapeutic strategies that act on

upstream signaling factors such as NLK is the capacity to influence several neuroprotective

mechanisms simultaneously. Nevertheless, not all downstream events are likely to be

beneficial(18), emphasizing the need for further investigations into NLK targets and the impact of

NLK-mediated phosphorylation on their function and contribution to disease.

Methods

Sex as a biologic variable

Our study examined tissue from both male and female patients, and similar findings are reported

for both sexes.

HEK293 cell culture and transfection

Human embryonic kidney (HEK) 293T were cultured in DMEM (Gibco, 11995065) supplemented

with 10% FBS (Gibco, ILT10082147) at 37°C in 5% CO_2. Cells were transfected with Lipofectamine

2000 (Invitrogen, 11668027) according to the manufacturer's instructions. For experiments with

YFP-NLS reporters, the lipofectamine solution was divided toto ensure there would be both cells

co-transfected with NLK and reporter as well as cells transfected with the reporter alone. Of the

lipofectamine solution, 75% contained two plasmids (NLK and reporter) while the remaining 25%

contained reporter alone. The lipofectamine solution was then combined and briefly mixed before

adding directly to cells. After 24 hours, transfected cells were split in media containing poly D-

lysine (1:500, cat Millipore A-008-E) onto coverslips coated with laminin (1:100, Sigma L2020-

1MG). Cells were cultured an additional 24 hours then immunocytochemistry was performed as

detailed below.

Primary neuron cell culture and transfection

Cortices from embryonic day (E)19-20 Long-Evans rat embryos were dissected and disassociated,

and primary neurons were plated at a density of 6×10^5 cells/mL in 96-well plates or in 24 well plate

on coverslips. At *in vitro* day (DIV) 4, neurons were transfected using Lipofectamine 2000 as

previously described (49, 60). For experiments with the YFP-NLS reporter, the transfection mix was

split as described above for HEK cells. Following transfection, cells were placed in Neurobasal

Complete Media (Neurobasal (Gibco 21103-049), 1x B27 Supplement (Gibco, 17504-044), 1x

Glutamax, 100 units/mL Pen Strep). Cells were fixed for immunocytochemistry 48 hours after

transfection.

iPSC maintenance and differentiation

Creating Neural Progenitors: iPSCs were dissociated using Accutase, counted, and plated at

32,000 cells/mL in E8 media with ROCK inhibitor. Differentiation was then induced with

doxycycline for 48 hours (2 µg/mL; Sigma, #D3447). Differentiating neural progenitors (NP) were

dissociated using Accutase after two days of doxycycline induction and frozen for future

experiments.

Differentiating Neural Progenitors: Previously frozen neural progenitors were thawed in E8 media

with ROCK inhibitor and incubated for 24 hours. Subsequently, the media was changed to N2

media containing 1x N2 Supplement (Gibco, #17502-048), 1x NEAA Supplement (Gibco, #11140-

050), 10 ng/mL BDNF (Peprotech, #450-02), 10 ng/mL NT3 (Peprotech, #450-03), 0.2 µg/mL

Laminin (Sigma, #L2020), and 2 µg/mL doxycycline (Sigma, #D3447). On Day 2, the media was

replaced with transition media (1:1 full E8 media and DMEM/F12 (Gibco, #11320-033)). On Day 3,

the media was switched to B27 media, prepared with 1x B27 Supplement (Gibco, #17504-044), 1x

Glutamax Supplement (Gibco, #35050-061), 10 ng/mL BDNF, 10 ng/mL NT3, 0.2 µg/mL Laminin, 1x

CultureOne (Gibco, #A33202-01), and Neurobasal-A (Gibco, #12349-015). On Day 6, cells were

transduced with the virus (University of Michigan Vector Core) and maintained in the same media

for the remainder of the experiment.

iPSC cell lines

The cell lines 1021 and 793 (controls) and 883 and 321 (C9ORF) were generated and characterized

as previously described (7).. The CS52i and corrected cell lines were obtained from Cedars Sinai.

See table below for additional details.

	ID	Onset age	Biopsy age	Gender	Onset	Notes
C9orf72-fALS	C9b (883)	49	51	M	Lumbar	C9orf72 mutation +
	C9a (312)	52	54	M	Lumbar	C9orf72 mutation +
	C9c	46	47	M	Lumbar	C9orf72 mutation +

	(CS52)				
TDP43-fALS	TDP43 (M337V)		54	F		M337V and C-terminal Dendra2 inserted into ND1 (1021) by CRISPR/Cas9
	TDP43 (WT)		54	F		C-terminal Dendra2 inserted intoND1 (1021) by CRISPR/Cas9
Control	ND1 (1021)		54	F		Healthy
	ND2 (746)		58	M		Healthy
	C9c-corrected	46	47	M	Lumbar	C9orf72 mutation removed from CS52i by CRISPR/Cas9

mbOrg generation

Isogenic human iPSC line WTC11 (GRN+/+ and GRN−/−) were generated by Dr. Bruce R. Conklin, as

previously described (61). iPSCs were cultured and maintained in StemMACS™ iPS-Brew XF

(Miltenyi Biotec, 130-104-368) media on 6-well cell culture plates (GenClone, 25-105MP) coated

with Vitronectin (Gibco, A14700) in DPBS. iPSCs were dissociated and passaged using EDTA

(Invitrogen, AM9260G) in DPBS.

iA and iN were differentiated separately, plated in a 1:1 ratio, and aged for 4 weeks as previously

described (46). For iN differentiation, iPSCs were transduced with NGN2-expressing lentivirus

constructs and a previously published protocol was followed for differentiation (61). Briefly, iPSCs

are expanded, dissociated, and replated on Matrigel-coated plates. Cells are grown in specialized

iNeuron induction media (DMEM-F12 + Glutamax; Gibco, 10565-018), N-2 supplement (Gibco,

17502-048), MEM-NEAA (Gibco, 11140-050) containing doxycycline (Sigma, D3072) for 72 h, with

media changed every 24 h. Cells are then dissociated using Accutase and plated together with iA.

Cortical-like iA are generated as previously described (62). Human induced pluripotent stem cells

(GRN+/+ and GRN-/-) were grown on vitronectin coated tissue culture plates using StemMACS™

iPS-Brew XF (Miltenyi Biotec, 130-104-368) media. On day 0 of differentiation, iPSCs are

dissociated into small aggregates and transferred in neurosphere induction medium (NSIM)

(DMEM-F12/Neurobasal-A at 1:1) plus SMAD inhibitors SB431542 and DMH1. On day 7, spheroids

are transferred to Matrigel-coated tissue culture plates with NIM. On day 14, rosette clusters were

mechanically removed and transferred to tissue culture flasks with NIM plus FGFb (Peprotech,

100-18B). Media was changed every 72 hours. On day 20, spheroids were triturated into a single

cell suspension and transferred to a new untreated cell culture flask with astrocyte media (ASM)

[(DMEM-F12 (Gibco, 10565-018), N2 Supplement (Gibco, 17502-048), B27 -Vit.A Supplement

(Gibco, 12587-010), Heparin (Stemcell Tech, 07980)) plus Y27632 (Tocris, 1254)]. From Day 28 to

180, spheroid aggregates were maintained in suspension with ASM plus EGF and FGFb (Peprotech,

100-15 & 100-18B) with media changes every 4-5 days. Spheroid aggregates were triturated every

7-10 days and transferred to new untreated tissue culture flasks. Spheroids are triturated every 7–

10 days and transferred to new tissue culture flasks. In order to generate mbOrgs, iN and iA are

plated together at a 1:1 ratio on Matrigel-coated 24-well plates at a collective density of ~1x106

cells/well. Cells are maintained in BrainPhys Complete medium with a 50% medium change every

72 h. mbOrgs will be created on 384-microwell plates as described previously (46), resulting in

mbOrgs that are highly uniform in size (~500 m) after brief spin-down and 1–2 days of subsequent

culture. These 3D co-cultures were aged for 4 weeks.

Longitudinal microscopy

At in vitro day (DIV) 4, neurons were transfected with 100ng of a control fluorescent plasmid to

mark cell bodies and 100ng of an experimental construct using Lipofectamine 2000 as previously

described (43, 49, 62). Neurons were imaged as described previously (43, 49, 60, 63) using a Nikon

Eclipse Ti inverted microscope with PerfectFocus3a 20X objective lens and either an Andor iXon3

897 EMCCD camera or Andor Zyla 4.2 (+)sCMOS camera. A Lambda CL Xenon lamp (Sutter) with 5

mm liquid light guide (Sutter)was used to illuminate samples, and custom scripts written in

Beanshell for use in micromanager controlled all stage movements, shutters, and filters. For

automated analyses of primary neuron survival, custom ImageJ/FIJI macros and Python scripts

were used to identify neurons and draw cellular regions of interest (ROIs) based upon size,

morphology, and fluorescence intensity. Custom Python scripts were used to track ROIs over time,

and cell death marked a set of criteria that include rounding of the soma, loss of fluorescence and

degeneration of neuritic processes (63, 64). For iNeuron survival, brightfield and fluorescent

images for survival experiments started on Day 14 and continued for 10 days. For manual analysis

of iNeuron survival, image time-series were processed by flat-field correction and image

registration, followed by programmatic de identification for blinded analysis. Time-series images

were uploaded to a browser-based server where a trained user manually counted survival using the

point tracking mode. GFP-positive cells were identified at T1, and cell fate was tracked in brightfield

images, where neuron death (uncensored event) was recorded at the appropriate time point. Living

neurons at the final time point were considered right-censored. Cox proportional hazards model

was applied to the datasets, stratifying by biological replicates, and the corresponding hazard plots

were generated in R.

Immunocytochemistry

Cells were fixed with 4% paraformaldehyde (PFA; Sigma, P6148) for 10m, rinsed with PBS, and

permeabilized with 0.1% Triton X-100 (Bio-rad, 161-0407) for 10m. Cells were then blocked in 3%

bovine serum albumin (BSA; Fisher, BP9703-100) in PBS at RT for 1h before incubation O/N at 4°C

in primary antibody diluted in 3% BSA (See table for additional details). Cells were then washed 3

times in PBS and incubated at RT with secondary antibodies diluted 1:250 in 3% BSA for 1h.

Following 3 washes in PBS containing 1:10,000 Hoechst 33258 dye (Invitrogen, H3569), cells were

mounted on Superfrost Plus Microscope Slides (Fisher, 1255015) with Prolong Gold Antifade

Mounting Reagent (Fisher, P10144) and imaged as described below. Images were analyzed by

collecting measurements within cellular regions of interest using custom ImageJ/FIJI macros.

Graphs were made using R.

Fluorescence in situ hybridization

Cells were fixed with 4% paraformaldehyde (Cat #) for 10 m, rinsed with PBS and permeabilized

with 0.1% Triton X-100 (Bio-rad, 161-0407) for 10m. Cells were then washed with 2X SSC (2 x 5 m)

and incubated in hybridization buffer for 2 hours at 37 degrees (8.75% dextran, 1.75X SSC, 17.5%

formamide, 0.5 ug/ul yeast tRNA, 10 mmol RVC, 0.1% BSA, 0.002 ug/ul Cy3-oligo (dT) 30 probe).

Cells were then washed with successive SSC rinses (4X SSC 15 m, 2X SSC 15 m, 2X SSC 15m).

Then immunocytochemistry was performed as above, beginning with blocking in 3% BSA. All

solutions prior to hybridization were DEPC treated.

Immunohistochemistry

Duplex immunohistochemistry was performed on a Ventana Discovery Ultra stainer (Indianapolis,

IN). Slides were dewaxed, rehydrated and subjected to heat induced epitope retrieval on board the

stainer. Slides were then subjected to sequential incubation with NLK (Rabbit polyclonal antibody,

AbCam, Cambridge, MA, Ab26050, 1:250, 32 minutes) and polymer goat anti-rabbit IgG conjugated

to HRP (Ventana) and developed with Discovery Green chromogen (Ventana). After an additional

round of heat induced epitope retrieval to remove the NLK primary antibody-secondary antibody

complex, the slides were stained with TDP43 (Rabbit polyclonal antibody, Proteintech, Rosemont,

IL, 10782-2-AP, 1:2000, 20 minutes), polymer goat anti-rabbit IgG conjugated to HRP and

developed with Discovery Brown chromogen (Ventana). Slides were then countered stained with

hematoxylin and coverslipped.

RNA isolation for bulk RNAseq and RT-PCR

RNA extraction was performed using the Trizol/phenol-Chloroform method (Sigma, T9424) as

previously described (62) and according to manufacturer specifications. Each sample contained

~60 mbOrgs, totaling ~3 x10^6 of cells per sample. The extracted RNA was used as a template for the

synthesis of complementary DNA (cDNA) through reverse transcription, using iScript™ cDNA

Synthesis Kit (Bio-Rad Cat#1708891) according to the manufacturer's protocol.

Quantitative PCR

cDNA samples were treated for genomic DNA contamination using DNA-*free*™ DNA Removal Kit

(Invitrogen Cat#AM1906) per manufacturer's instructions. The cDNA was then diluted to a

concentration of 5 ng/μl and 4 μl of each sample (total of 20 ng) were aliquoted in a MicroAmp™

Optical 96-Well Reaction Plate (Thermo Scientific Cat#N8010560) in technical duplicates. Samples

were processed using 2X SYBR Green qPCR Master Mix Assay and quantitative PCR was run on

QuantStudio 6 Real-Time PCR system following manufacturer's instructions. Data analysis was

carried out applying the Pfaffl mathematical model for relative transcript quantification (65) using

GAPDH as a housekeeping gene.

RNA Sequencing

RNA sequencing and analysis of RNA integrity was analyzed on an Agilent 2100 Bioanalyzer using

RNA 6000 Nano kit (Agilent, 5067-1511). Only samples with an RNA integrity number (RIN) ≥ 9.4

were used to perform bulk RNA sequencing. Nugen Universal Plus (Tecan) was used as a library kit

and libraries were sequenced on a SP300 flow cell of the Illumina NovaSeq 6000 machine with a

paired end 150 bp sequencing strategy (average depth 90 million reads/sample) at UCSF Genomics

Core Facility. Genome was aligned to Ensembl Human.GRCh38.103. Kallisto 0.46.01 was used to

generate transcript abundance files for each sample. Transcript counts files for each sample were

generated using txImport and transcript differential analysis was performed using DESeq2 v1.24.0.

A total of 6 samples were spread across two conditions.

Confocal microscopy

Confocal images were taken on a Nikon AXR NSPARC confocal system with a 60x NA1.42 Oil/DIS

PLan-Apochromat Lambda D objective with a working distance of 1.5 mm, and a 40x CFI

Apochromat LWD Lambda S objective with a working distance of 0.30 mm.

Light microscopy

Whole-slide images were generated by the University of Michigan Digital Pathology group within the

Department of Pathology using an Aperio AT2 scanner (Leica Biosystems) equipped with a 20x NA

0.75 Plan Apochromat objective. 40x scanning is achieved using a 2x optical magnification

changer. Resolution is 0.25 µm/pixel for 40x scans. Focus during the scan was maintained using a

triangulated focus map built from individual focus points determined in a separate step before

scanning was started. Proprietary software was used for image processing during the acquisition.

TIFs were taken of the entire ventral horn of all cases, then a pathologist selected each neuron,

manually drew an ROI, and annotated the TDP status (pathology or no pathology) using custom Fiji

scripts. Then deconvolution and analysis were performed using custom Fiji scripts.

High quality images for figures were acquired on an Olympus BX51 light microscope equipped with

a UPlanSApo100x oil objective with a numerical aperture of 1.40 and a working distance of 0.12

mm. Image deconvolution was performed using Fiji.

Antibodies:

	Vendor	Catalog #	Concentration for ICC
Flag	AbCam	ab1170	1:200
TDP43	R&D	MAB7778	1:200
TDP43	Proteintech	10782-2-AP	1:200
FUS	Sigma	HPA008784	1:200

HNRNPA2B1	Proteintech	14813-1-AP	1:100
MATR3	Abcam	ab281927	1:200
V5	Abcam	ab53418	1:200
KPNA2	Proteintech	10819-1-AP	1:200
KPNB1	Abcam	AB2811	1:200
Ran	BD Transduction labs	610340	1:400
RanGAP	Abcam	AB92360	1:200
RanBP2	Abcam	AB64276	1:200
Mab414	Abcam	AB24609	1:200
NXF1	Abcam	AB129160	1:200
SC-35	Sigma	S4045-.2ML	1:200
SFPQ	Proteintech	15585-1-AP	1:200
Nucleophosmin	Millipore	MABE937	1:200

509 **Plasmids**

Plasmid	Source
FLAG-NLK-WT	Gift from T. Ishitani (66)
FLAG-NLK KN (K155M)	Gift from T. Ishitani (66)
EYFP2-SV40NLS-NES	Gift from Yuh Min Chook (67)
EYFP2-SV40NLS	This paper
EYFP2-TDPNLS	This paper
EYFP2-FUSNLS	This paper
EYFP-MATR3NLS	This paper
pGW1-TDP43(2F/L)-EGFP	Flores et al 2019 (36)
FUGW-SNAP-FLAG	This paper
FUGW-SNAP-FLAG-NLK	This paper

FLAG-NLK-WT and FLAG-NLK KN were a gift from T Ishitani . SNAP-FLAG and SNAP-FLAG NLK were

synthesized and subcloned into FUGW. All YFP reporter plasmids were derived from EYFP2-

SV40NLS-NES, a kind gift from Yuh Min Chook (67). Site-directed mutagenesis was used to add a

stop codon after the SV40NLS. The NLS of TDP43 (residues 82-98) and FUS (residues 495-526)

were PCR amplified using primers below, digested with XbaI and BglII, and cloned into the

corresponding sites in EYFP2-SV40NLS-NES. The NLS of MATR3 (residues 583-602) was purchased

as a Geneblock from Integrated DNA Technologies (IDT), and then digested with XbaI and BglII, and

cloned into the corresponding sites in EYFP2-SV40NLS-NES. All plasmids were verified by Sanger

sequencing.

Primers

Amplify TDP43 NLS	For: ctcagatctcgaAAAAGAAAAATGGAT
	Rev: ttatctagagatTCTTTTCACTTTCAC
Amplify FUS NLS	For: tcagatctCGAGGGGGCCGGGGT
	Rev: tatctagaATACGGCCTCTCCCT
Mutagenesis: introduce stop codon into EYFP2-SV40NLS-NES	For: ctcgaccaaaaaagaagagaaaggtataagctttagccttgaaa
	Rev: tttcaaggctaaagcttatacctttctcttcttttttggtcgag
qPCR NLK	For: GGGTCCTCATAAACAGCCATC
	Rev: AGACCAACATCCTGCAAAGG
qPCR GAPDH	For: GGG GTC ATT GAT GGC AAC AAT A
	Rev: ATG GGC AAG GTG AAG GTC G

Data analysis and statistics

Statistical information, including mean and statistical significance values is indicated in the figures

or figure legends. At least three biological replicates were used per experiment. For analysis of

immunocytochemistry images, ROIs for nucleus and cytoplasm were either generated using CVAT

(68) or manually drawn in ImageJ. Nuclear/cytoplasmic ratio was calculated and statistical

significance was determined with GraphPadP Prism 9 using the tests indicated in each figure. Data

were considered statistically significant at $*P < 0.05$, $**P < 0.01$, $***P < 0.001$ and $****P < 0.0001$.

Data from Liu et al. was analyzed with assistance from the TDP43 loss of function database (69);

DESeq2 P-values were calculated using Wald tests and corrected for multiple testing using the

Benjamini–Hochberg method. DEU events with Padj <0.05 were considered significantly changed.

Study approval

Skin samples for iPSC creation were collected and de- identified in collaboration with the Michigan

Institute for Clinical and Health Research (MICHR, UL1TR000433) through an institutional review

board (IRB)-approved protocol (HUM00028826).

Data availability

All correspondence regarding the materials and data presented in this paper should be addressed

to sbarmada@med.umich.edu.

Author Contributions

MB and SB designed the study. EP and MB designed experiments with SB. MB and EP collected

data for most experimental studies, analyzed the data, assembled the figures, and wrote the

manuscript. MB and EP performed immunocytochemistry. JM assisted with data collection and

analysis. MB performed survival assays in cultured rat neurons and iNeurons, wrote custom FIJI

scripts for quantitation of immunocytochemistry and dual IHC. EP made YFP reporter constructs,

supervised dual IHC (with the IHC core),and selected motor neurons for quantitation. LGG and MM

contributed to mbOrg concept, development, and data analysis. MK and SC cultured mbOrgs and

performed RNASeq and qPCR. EU contributed to original concept and development of mbOrgs,

and provided RNA-seq data from mbOrgs. XL isolated and cultured rat cortical neurons. EMHT

derived iPSC lines and developed protocols for their differentiation. JW developed original code for

neuron segmentation and image processing. SB provided resources, funding, and conceptual input

for experiments and supervised the research. All authors were involved throughout the research

process, agreed amongst themselves regarding roles and responsibilities, and contributed to the

review, editing, and approval of the manuscript.

Acknowledgements

This work was supported by: National Institutes of Health AWD012778 (EP), National Institutes of

Health (R01NS097542, R01NS113943 and 1R56NS128110-01 to SJB; R44NS124457 to MM;

P30AG072931 to the University of Michigan Brain Bank and Alzheimer's Disease Research

CenterNational Institute on Drug Abuse R44NS124457 (MDM and EMU), the family of Angela

Dobson and Lyndon Welch, the A. Alfred Taubman Medical Research Institute, the Danto Family,

Ann Arbor Active Against ALS, and the Robert Packard Center for ALS Research.

We thank all members of the Barmada laboratory for their advice and suggestions. We thank Dr.

Dafydd Thomas of the Rogel Cancer Center Tissue and Molecular Pathology Shared Resource

Laboratory at the University of Michigan (NIH P30 CA04659229). We thank the Michigan Institute for

Clinical and Health Research (MICHR, UL1TR000433), who collaborated to collect and de-identify

skin samples for iPSC creation. Finally, we thank the patients that donated tissue samples to make

this work possible.

Schematics were created in BioRender. Barmada, S. (2024) BioRender.com/z94b294

Figure 1

A FLAG-NLK KN FLAG-NLK WT **TDP43**

FLAG

TDP43

Merge w/ DNA

B N/C *** KN WT

C Nuclear Mean *** KN WT

D Cytoplasmic Mean *** KN WT

E Whole-cell Mean n.s. KN WT

F FLAG-NLK KN FLAG-NLK WT

FLAG

TDP43

FUS

Merge

H FLAG-NLK KN FLAG-NLK WT

FLAG

TDP43

HNRNPA2B1

Merge

J FLAG-NLK KN FLAG-NLK WT

FLAG

TDP43

MATR3

Merge

G FUS N/C *** KN WT

I HNRNPA2B1 N/C ** KN WT

K MATR3 N/C n.s. KN WT

Figure 1. Overexpression of NLK leads to cytoplasmic accumulation of TDP43 and ALS-FTD relevant RNA-binding proteins. (**A**) HEK cells were transfected with plasmids encoding either FLAG-NLK KN (KN; kinase-negative) or FLAG-NLK WT (WT; wild-type) followed by immunofluorescence using antibodies against FLAG and TDP4343 while DNA was stained with Hoechst. Scale bar = 10μm. (**B**) Superplot of Nuclear-Cytoplasmic ratio (N/C) of TDP43 from images in (A). Line = mean, error bar = standard error. *p<0.05, ** p<0.01, ***p<0.001. (unpaired T-test with Welch's correction). (**C**) Superplot of TDP43 nuclear intensity in cells that overexpress KN or WT NLK. Line = mean, error bar = standard error. (**D**) Superplot of TDP43 cytoplasmic intensity in cells that overexpress KN or WT NLK. Line = mean, error bar = standard error. (**E**) Superplot quantification of TDP43 whole-cell intensity in cells that overexpress KN or WT NLK. Line = mean, error bar = standard error. (**F**) HEK cells were transfected with the indicated FLAG-NLK plasmids as in (A) followed by immunofluorescence using antibodies against FLAG, TDP43, and FUS while DNA was stained with Hoechst. Scale bar = 10μm (**G**) Superplot of FUS localization in cells that overexpress KN or WT NLK. Line = mean, error bar = standard error. (**H**) HEK cells were transfected with the indicated FLAG-NLK plasmids as in (A) followed by immunofluorescence using antibodies against FLAG, TDP43, and HNRNPA2B1 while DNA was stained with Hoechst. Scale bar = 10μm. (**I**) Superplot of HNRNPA2B1 localization in cells that overexpress KN or WT NLK. Line = mean, error bar = standard error. (**J**) HEK cells were transfected with the indicated FLAG-NLK plasmids as in (A) followed by immunofluorescence using antibodies against FLAG, TDP43, and MATR3 while DNA was stained with Hoechst. Scale bar = 10μm. (**K**) HEK cells were transfected with the indicated FLAG-NLK plasmids as in (A) followed by immunofluorescence using antibodies against FLAG and MART3 while DNA was stained with Hoechst. Scale bar = 10μm. (**L**) Superplot of HNRNPA2B1 localization in cells that overexpress KN or WT NLK. Line = mean, error bar = standard error.

Figure 2

Figure 2. NLK overexpression disrupts NLS dependent nuclear import. (A-D) HEK cells were co-

transfected with plasmids encoding either FLAG-NLK KN or FLAG-NLK WT and either of the

following NLS-reporter eYFP-NLS^{TDP43} (A), eYFP-NLS^{FUS} (B), eYFP-NLS^{MATR3} (C), or eYFP-NLS^{SV40}.

Schematics created in BioRender. (D). Immunofluorescence was performed using antibodies

against FLAG and either TDP43, FUS, or Matrin-3 while DNA was stained with Hoechst. Scale bar 10

uM. **(E-H)** Superplot of NLS-reporter Nuclear-Cytoplasmic ratio (N/C) from images in A-D. Line =

606   mean, error bar= standard error. *** indicates p-value < 0.001 (unpaired t-test with Welch's

607   correction).

Figure 3

Figure 3. NLK-dependent mislocalization of TDP43 does not depend on nuclear accumulation

of KPNA2 and KPNB1. (**A**) HEK cells were transfected with plasmids encoding either FLAG-NLK KN

(KN; kinase-negative) or FLAG-NLK WT (WT; wild-type) followed by immunofluorescence using

antibodies against FLAG, TDP43, and KPNA2 or KPNB1 while DNA was stained with Hoechst. Scale

bar = 5μm. (**B**) Quantification of KPNA2 and KPNB1 localization in cells that overexpress NLK KN or

WT. Line = mean, error bar = standard error. Statistical significance indicated as such in all sub-

panels: n.s. = not significant, *p<0.05, ** p<0.01, ***p<0.001(unpaired t-test with Welch's

correction).(**C**) HEK cells were co- transfected with plasmids encoding FLAG-NLK WT and either

mApple (negative-control) or V5-FBXW7 followed by immunofluorescence using antibodies against

KPNA2, FLAG, and V5 while DNA was stained with Hoechst. Scale bar = 20μm. (**D**) Quantification of

KPNA2 localization in cells that overexpress FLAG-NLK WT and either mApple or V5-FBXW7. Line=

mean, error bar= standard error. *$p<0.05$, ** $p<0.01$, ***$p<0.001$ (unpaired t-test with Welch's

correction). (**E**) HEK cells were co-transfected with plasmids encoding FLAG-NLK WT and either

mApple or V5-FBXW7 followed by immunofluorescence using antibodies against TDP43, FLAG, and

V5 while DNA was stained with Hoechst. Scale bar = 20μm. (**F**) Quantification of TDP43 localization

in cells that overexpress FLAG-NLK WT and either mApple or V5-FBXW7. Line = mean, error bar=

standard error. *$p<0.05$, ** $p<0.01$, ***$p<0.001$ (unpaired t-test with Welch's correction). (G) HEK

cells were co-transfected as in E) followed by immunofluorescence using antibodies against

KPNA2, FLAG, and V5 while DNA was stained with Hoescht. Scale bar = 20 μm.(H) HEK cells were

co- transfected as in E) followed by immunofluorescence using antibodies against KPNA2, FLAG,

and V5 while DNA was stained with Hoechst. Scale bar = 20μm. (I) Quantification of KPNB1

localization in cells that overexpress NLK WT and either mApple orr V5-FBXW7. Line = mean, error

bar = standard error.

Figure 4

Figure 4. NLK overexpression mislocalizes RanBP2-RanGAP1 and disrupts the Ran gradient.

(**A**) Representative western blots from HEK cells transfected with either vector (-) or FLAG-NLK WT

(WT, +; wild-type). #'s to the left represent molecular weight in kDa. (**B**) Western blots after

immunoprecipitation (IP) using 1µg of the indicated antibodies in 1mg of lysates from (A). (**C**)

Western blots after RanBP2 co-IP from lysates in (A) were run on a 4-20% tris-glycine gel to

facilitate visualization of RanBP2 and validate NLK complex formation with RanBP2-RanGAP1-

SUMO. (**D**) HEK cells were transfected with plasmids encoding either FLAG-NLK KN (KN; kinase-

negative) or FLAG-NLK WT (WT; wild-type) followed by immunofluorescence using antibodies

against FLAG, TDP43, and either Ran, RanGAP1, RanBP2, or MAb414 (FG-nucleoporins) while DNA

was stained with Hoechst. Scale bar = 10µm. (**E**) Superplot of nuclear-cytoplasmic ratio (N/C) of

Ran. Line = mean, error bar= standard error. Statistical significance indicated as such in all sub-

panels: n.s. = not significant, *p<0.05, ** p<0.01, ***p<0.001. *p<0.05, ** p<0.01, ***p<0.001

(unpaired t-test with Welch's correction). (**F**) HEK cells were transfected as in (D) followed by

immunofluorescence using antibodies against FLAG, TDP43, and RANBP2 while DNA was stained

with Hoechst. Scale bar = 10µm. (**G**) Superplot of nuclear-rim-cytoplasm ratio (Nuc. Rim/Cyto) of

RANBP2. (**H**) HEK cells were transfected as in (D) followed by immunofluorescence using

antibodies against FLAG, TDP43, and MAB414 (FG-nucleoporins) while DNA was stained with

Hoechst. Scale bar = 10µm. (**I**) Superplot of nuclear-rim-cytoplasm ratio (Nuc. Rim/Cyto) of

MAB414. (**J**) HEK cells were transfected as in (D) followed by immunofluorescence using antibodies

against FLAG, TDP43, and RAN while DNA was stained with Hoechst. Scale bar = 10µm. (**K**)

Superplot quantification of nuclear-cytoplasm ratio (N/C) of RAN.

Figure 5

A FLAG-NLK KN   FLAG-NLK WT

FLAG

TDP F2L

FUS

Merge

B FLAG-NLK KN   FLAG-NLK WT

FLAG

NXF1

TDP43

Merge

C FLAG-NLK KN   FLAG-NLK WT

FLAG

PolyA

TDP43

Merge

D FLAG-NLK KN   FLAG-NLK WT

FLAG

SC35

TDR43

Merge

E FLAG-NLK KN   FLAG-NLK WT

FLAG

SFPQ

TDP43

Merge

F FLAG-NLK KN   FLAG-NLK WT

FLAG

Npm

TDP43

Merge

Figure 5. NLK overexpression drives dissolution of nuclear speckles. (**A**) HEK cells were co-

transfected with plasmids encoding either FLAG-NLK KN or FLAG-NLK WT and a GFP-tagged RNA-

binding mutant TDP43 (TDP43 F147/9L; F2L) followed by immunofluorescence using antibodies

against FLAG and FUS and direct visualization of tagged protein while DNA was stained with

Hoechst. Scale bar 10 μm. (**B**) HEK cells were transfected with plasmids encoding either FLAG-NLK

KN or FLAG-NLK WT followed by immunofluorescence using antibodies against FLAG, TDP43, and

NXF1. Scale bar 10 μm. (**C**) HEK cells were transfected with plasmids encoding either FLAG-NLK

KN or FLAG-NLK WT followed by polyA FISH and immunofluorescence for TDP43. Scale bar 10 μm.

(**D-F**) HEK cells were transfected with plasmids encoding either FLAG-NLK KN or FLAG-NLK WT

followed by immunofluorescence for TDP43 and markers of speckles (SC-35) (D), paraspeckles

(SFPQ) (E), or nucleolus (nucleophosmin, Npm) (F). Scale bar 10 μm.

Figure 6

Figure 6. NLK overexpression disrupts nuclear import and is toxic in primary rat neurons. (**A**)

Rodent primary cortical neurons were transfected with either SNAP-FLAG (SF; negative-control) or

SNAP-FLAG-NLK (SF-NLK) followed by immunofluorescence using antibodies against FLAG and

either TDP43, FUS, Ran, RanBP2, RanGAP1, and Mab414 while DNA was stained with

Hoechst.Scale bar = 10 μm. (**B**) Rodent primary cortical neurons were co-transfected with plasmids

encoding either SF or SF-NLK and EGFP (survival marker), treated with SNAP-647 dye at T1 to

visualize SNAP-positive neurons, and tracked by longitudinal microscopy to determine neuron fate

(see methods). Scale bar = 20μm. (**C**) Cumulative Hazard plot showing the relative risk of death in

neurons that express either SF or SF-NLK. Hazard ratio = 1.616. ***p=1.494e-59. (**D**) Cox

proportional hazards model predicting relative hazard associated with different intensities of

SNAP-FLAG-NLK. Solid lines indicate the estimated hazard, with color gradients corresponding to

intensity levels: grey (low intensity), light red (medium intensity), and red (high intensity). Dashed

lines represent 95% confidence intervals for hazard estimates.

Figure 7

Figure 7. TDP43 pathology is associated with NLK overexpression in human model systems and patient samples. (**A**) Schematic of generation of mbOrgs. (**B**) Normalized NLK counts from RNASeq performed on WT or GRN[-/-] mbOrgs. Line = mean, error bar = standard deviation. ** $p< 0.01$ (unpaired t-test with Welch's correction). (**C**) QPCR was performed on WT or GRN[-/-] mbOrgs (2 biological replicates, 3 technical replicates). Superplot of NLK levels normalized to GAPDH. Line = mean, error bar = standard deviation. * $p<0.05$ (unpaired t-test with Welch's correction). (**D**) NLK normalized counts from RNAseq performed on TDP positive and negative nuclei. Line = mean, error bar = standard deviation. **** $p<0.001$ (Wald tests, corrected for multiple testing using the Benjamini–Hochberg method). (**E**) Dual immunohistochemistry for NLK and TDP43, performed on spinal cord tissue from four patients with sporadic ALS. Images deconvolved using FIJI. Upper panels: scale bar 100 µm. Lower panels: scale bar 20 µm. (**F**) Quantitative real-time PCR of NLK mRNA levels in iPSC derived neurons after transduction with lentivirus virus encoding either non-targeting shRNA (shNT) or NLK shRNA (shNLK). Line = mean, error bar = standard error. *** $p< 0.001$ (unpaired t-test with Welch's correction). (**G**) Isogenic (WT) and mutant TDP43 (M337V) iPSC-derived neurons were transduced with virus encoding either non targeting (NT) shRNA or NLK shRNA followed by longitudinal microscopy to track neuron fate (see methods). Scale bar = 20 µm. (**H**) Cumulative Hazard plot showing the relative risk of death in ND and C9 neurons that express either shNT shRNA or shNLK (HR = 0.4, $p<0.001$). (**I**) Cumulative Hazard plot showing the relative risk of death in WT and M337V neurons that express either Scramble shRNA or NLKshRNA. (HR = 0.23, p=0.0013).

References

1.   Brown RH, and Al-Chalabi A. Amyotrophic Lateral Sclerosis. *N Engl J Med*.

2017;377(16):1602.

2.   Tan RH, Ke YD, Ittner LM, and Halliday GM. ALS/FTLD: experimental models and reality.

Acta Neuropathol. 2017;133(2):177-96.

3.   Neumann M, Sampathu DM, Kwong LK, Truax AC, Micsenyi MC, Chou TT, et al.

Ubiquitinated TDP-43 in frontotemporal lobar degeneration and amyotrophic lateral

sclerosis. *Science*. 2006;314(5796):130-3.

4.   Alami NH, Smith RB, Carrasco MA, Williams LA, Winborn CS, Han SSW, et al. Axonal

transport of TDP-43 mRNA granules is impaired by ALS-causing mutations. *Neuron*.

2014;81(3):536-43.

5.   Ling JP, Pletnikova O, Troncoso JC, and Wong PC. TDP-43 repression of nonconserved

cryptic exons is compromised in ALS-FTD. *Science*. 2015;349(6248):650-5.

6.   Polymenidou M, Lagier-Tourenne C, Hutt KR, Huelga SC, Moran J, Liang TY, et al. Long pre-

mRNA depletion and RNA missplicing contribute to neuronal vulnerability from loss of TDP-

43. *Nat Neurosci*. 2011;14(4):459-68.

7.   Tank EM, Figueroa-Romero C, Hinder LM, Bedi K, Archbold HC, Li X, et al. Abnormal RNA

stability in amyotrophic lateral sclerosis. *Nat Commun*. 2018;9(1):2845.

8.   Takeuchi R, Toyoshima Y, Tada M, Shiga A, Tanaka H, Shimohata M, et al. Transportin 1

accumulates in FUS inclusions in adult-onset ALS without FUS mutation. *Neuropathol Appl*

Neurobiol. 2013;39(5):580-4.

9.   Freibaum BD, Lu Y, Lopez-Gonzalez R, Kim NC, Almeida S, Lee KH, et al. GGGGCC repeat

expansion in C9orf72 compromises nucleocytoplasmic transport. *Nature*.

2015;525(7567):129-33.

10.   Jovičić A, Mertens J, Boeynaems S, Bogaert E, Chai N, Yamada SB, et al. Modifiers of

732   C9orf72 dipeptide repeat toxicity connect nucleocytoplasmic transport defects to FTD/ALS.

733   *Nat Neurosci*. 2015;18(9):1226-9.

734   11.   Zhang K, Donnelly CJ, Haeusler AR, Grima JC, Machamer JB, Steinwald P, et al. The C9orf72

735   repeat expansion disrupts nucleocytoplasmic transport. *Nature*. 2015;525(7567):56-61.

736   12.   Thorpe CJ, and Moon RT. nemo-like kinase is an essential co-activator of Wnt signaling

737   during early zebrafish development. *Development*. 2004;131(12):2899-909.

738   13.   Yasuda J, Tsuchiya A, Yamada T, Sakamoto M, Sekiya T, and Hirohashi S. Nemo-like kinase

739   induces apoptosis in DLD-1 human colon cancer cells. *Biochem Biophys Res Commun*.

740   2003;308(2):227-33.

741   14.   Emami KH, Brown LG, Pitts TE, Sun X, Vessella RL, and Corey E. Nemo-like kinase induces

742   apoptosis and inhibits androgen receptor signaling in prostate cancer cells. *Prostate*.

743   2009;69(14):1481-92.

744   15.   Tejwani L, Jung Y, Kokubu H, Sowmithra S, Ni L, Lee C, et al. Reduction of nemo-like kinase

745   increases lysosome biogenesis and ameliorates TDP-43-related neurodegeneration. *J Clin

746   Invest*. 2023;133(16).

747   16.   Ju H, Kokubu H, Todd TW, Kahle JJ, Kim S, Richman R, et al. Polyglutamine disease toxicity

748   is regulated by Nemo-like kinase in spinocerebellar ataxia type 1. *J Neurosci*.

749   2013;33(22):9328-36.

750   17.   Todd TW, Kokubu H, Miranda HC, Cortes CJ, La Spada AR, and Lim J. Nemo-like kinase is a

751   novel regulator of spinal and bulbar muscular atrophy. *Elife*. 2015;4:e08493.

752   18.   Jiang M, Zhang X, Liu H, LeBron J, Alexandris A, Peng Q, et al. Nemo-like kinase reduces

753   mutant huntingtin levels and mitigates Huntington's disease. *Hum Mol Genet*.

754   2020;29(8):1340-52.

755   19.   Brott BK, Pinsky BA, and Erikson RL. Nlk is a murine protein kinase related to Erk/MAP

756   kinases and localized in the nucleus. *Proc Natl Acad Sci U S A*. 1998;95(3):963-8.

757   20.   Lee BJ, Cansizoglu AE, Süel KE, Louis TH, Zhang Z, and Chook YM. Rules for nuclear

758   localization sequence recognition by karyopherin beta 2. *Cell*. 2006;126(3):543-58.

759   21.   Winton MJ, Igaz LM, Wong MM, Kwong LK, Trojanowski JQ, and Lee VM. Disturbance of

760   nuclear and cytoplasmic TAR DNA-binding protein (TDP-43) induces disease-like

761   redistribution, sequestration, and aggregate formation. *J Biol Chem*. 2008;283(19):13302-9.

762   22.   Hisada-Ishii S, Ebihara M, Kobayashi N, and Kitagawa Y. Bipartite nuclear localization signal

763   of matrin 3 is essential for vertebrate cells. *Biochem Biophys Res Commun*.

764   2007;354(1):72-6.

765   23.   Ederle H, Funk C, Abou-Ajram C, Hutten S, Funk EBE, Kehlenbach RH, et al. Nuclear egress

766   of TDP-43 and FUS occurs independently of Exportin-1/CRM1. *Sci Rep*. 2018;8(1):7084.

767   24.   Pinarbasi ES, Cağatay T, Fung HYJ, Li YC, Chook YM, and Thomas PJ. Active nuclear import

768   and passive nuclear export are the primary determinants of TDP-43 localization. *Sci Rep*.

769   2018;8(1):7083.

770   25.   Archbold HC, Jackson KL, Arora A, Weskamp K, Tank EM, Li X, et al. TDP43 nuclear export

771   and neurodegeneration in models of amyotrophic lateral sclerosis and frontotemporal

772   dementia. *Sci Rep*. 2018;8(1):4606.

773   26.   Dormann D, Rodde R, Edbauer D, Bentmann E, Fischer I, Hruscha A, et al. ALS-associated

774   fused in sarcoma (FUS) mutations disrupt Transportin-mediated nuclear import. *EMBO J*.

775   2010;29(16):2841-57.

776   27.   Görlich D, Vogel F, Mills AD, Hartmann E, and Laskey RA. Distinct functions for the two

777   importin subunits in nuclear protein import. *Nature*. 1995;377(6546):246-8.

28.   Xing Z, Zhen Y, Chen J, Du M, Li D, Liu R, et al. KPNA2 Silencing, Regulated by E3 Ubiquitin

Ligase FBXW7, Alleviates Endothelial Dysfunction and Inflammation Through Inhibiting the

Nuclear Translocation of p65 and IRF3: A Possible Therapeutic Approach for

Atherosclerosis. *Inflammation*. 2023;46(6):2071-88.

29.   Stewart M. Molecular mechanism of the nuclear protein import cycle. *Nat Rev Mol Cell Biol*.

2007;8(3):195-208.

30.   Varjosalo M, Keskitalo S, Van Drogen A, Nurkkala H, Vichalkovski A, Aebersold R, et al. The

protein interaction landscape of the human CMGC kinase group. *Cell Rep*. 2013;3(4):1306-

20.

31.   Mahajan R, Delphin C, Guan T, Gerace L, and Melchior F. A small ubiquitin-related

polypeptide involved in targeting RanGAP1 to nuclear pore complex protein RanBP2. *Cell*.

1997;88(1):97-107.

32.   Saitoh H, Pu R, Cavenagh M, and Dasso M. RanBP2 associates with Ubc9p and a modified

form of RanGAP1. *Proc Natl Acad Sci U S A*. 1997;94(8):3736-41.

33.   Matunis MJ, Coutavas E, and Blobel G. A novel ubiquitin-like modification modulates the

partitioning of the Ran-GTPase-activating protein RanGAP1 between the cytosol and the

nuclear pore complex. *J Cell Biol*. 1996;135(6 Pt 1):1457-70.

34.   Duan L, Zaepfel BL, Aksenova V, Dasso M, Rothstein JD, Kalab P, et al. Nuclear RNA binding

regulates TDP-43 nuclear localization and passive nuclear export. *Cell Rep*.

2022;40(3):111106.

35.   Buratti E, and Baralle FE. Characterization and functional implications of the RNA binding

properties of nuclear factor TDP-43, a novel splicing regulator of CFTR exon 9. *J Biol Chem*.

2001;276(39):36337-43.

36.   Flores BN, Li X, Malik AM, Martinez J, Beg AA, and Barmada SJ. An Intramolecular Salt Bridge

Linking TDP43 RNA Binding, Protein Stability, and TDP43-Dependent Neurodegeneration.

Cell Rep. 2019;27(4):1133-50.e8.

37.   Pérez-Berlanga M, Wiersma VI, Zbinden A, De Vos L, Wagner U, Foglieni C, et al. Loss of

TDP-43 oligomerization or RNA binding elicits distinct aggregation patterns. *EMBO J.*

2023;42(17):e111719.

38.   Ilik İ, Malszycki M, Lübke AK, Schade C, Meierhofer D, and Aktaş T. SON and SRRM2 are

essential for nuclear speckle formation. *Elife.* 2020;9.

39.   Imamura K, Imamachi N, Akizuki G, Kumakura M, Kawaguchi A, Nagata K, et al. Long

noncoding RNA NEAT1-dependent SFPQ relocation from promoter region to paraspeckle

mediates IL8 expression upon immune stimuli. *Mol Cell.* 2014;53(3):393-406.

40.   Mitrea DM, Cika JA, Guy CS, Ban D, Banerjee PR, Stanley CB, et al. Nucleophosmin

integrates within the nucleolus via multi-modal interactions with proteins displaying R-rich

linear motifs and rRNA. *Elife.* 2016;5.

41.   Arrasate M, Mitra S, Schweitzer ES, Segal MR, and Finkbeiner S. Inclusion body formation

reduces levels of mutant huntingtin and the risk of neuronal death. *Nature.*

2004;431(7010):805-10.

42.   Barmada SJ, Serio A, Arjun A, Bilican B, Daub A, Ando DM, et al. Autophagy induction

enhances TDP43 turnover and survival in neuronal ALS models. *Nat Chem Biol.*

2014;10(8):677-85.

43.   Barmada SJ, Ju S, Arjun A, Batarse A, Archbold HC, Peisach D, et al. Amelioration of toxicity

in neuronal models of amyotrophic lateral sclerosis by hUPF1. *Proc Natl Acad Sci U S A.*

2015;112(25):7821-6.

44.   Miller J, Arrasate M, Shaby BA, Mitra S, Masliah E, and Finkbeiner S. Quantitative

825   relationships between huntingtin levels, polyglutamine length, inclusion body formation,

826   and neuronal death provide novel insight into huntington's disease molecular

827   pathogenesis. *J Neurosci.* 2010;30(31):10541-50.

828   45.   Strong MJ, Abrahams S, Goldstein LH, Woolley S, Mclaughlin P, Snowden J, et al.

829   Amyotrophic lateral sclerosis - frontotemporal spectrum disorder (ALS-FTSD): Revised

830   diagnostic criteria. *Amyotroph Lateral Scler Frontotemporal Degener.* 2017;18(3-4):153-74.

831   46.   de Majo M, Koontz M, Marsan E, Salinas N, Ramsey A, Kuo YM, et al. Granulin loss of

832   function in human mature brain organoids implicates astrocytes in TDP-43 pathology. *Stem*

833   *Cell Reports.* 2023;18(3):706-19.

834   47.   DeJesus-Hernandez M, Mackenzie IR, Boeve BF, Boxer AL, Baker M, Rutherford NJ, et al.

835   Expanded GGGGCC hexanucleotide repeat in noncoding region of C9ORF72 causes

836   chromosome 9p-linked FTD and ALS. *Neuron.* 2011;72(2):245-56.

837   48.   Morita M, Al-Chalabi A, Andersen PM, Hosler B, Sapp P, Englund E, et al. A locus on

838   chromosome 9p confers susceptibility to ALS and frontotemporal dementia. *Neurology.*

839   2006;66(6):839-44.

840   49.   McMillan M, Gomez N, Hsieh C, Bekier M, Li X, Miguez R, et al. RNA methylation influences

841   TDP43 binding and disease pathogenesis in models of amyotrophic lateral sclerosis and

842   frontotemporal dementia. *Mol Cell.* 2023;83(2):219-36.e7.

843   50.   Boeynaems S, Bogaert E, Michiels E, Gijselinck I, Sieben A, Jovičić A, et al. Drosophila

844   screen connects nuclear transport genes to DPR pathology in c9ALS/FTD. *Sci Rep.*

845   2016;6:20877.

51.   Hutten S, Flotho A, Melchior F, and Kehlenbach RH. The Nup358-RanGAP complex is

required for efficient importin alpha/beta-dependent nuclear import. *Mol Biol Cell*.

2008;19(5):2300-10.

52.   Hong AW, Meng Z, Yuan HX, Plouffe SW, Moon S, Kim W, et al. Osmotic stress-induced

phosphorylation by NLK at Ser128 activates YAP. *EMBO Rep*. 2017;18(1):72-86.

53.   Yuan HX, Wang Z, Yu FX, Li F, Russell RC, Jewell JL, et al. NLK phosphorylates Raptor to

mediate stress-induced mTORC1 inhibition. *Genes Dev*. 2015;29(22):2362-76.

54.   Cargnello M, and Roux PP. Activation and function of the MAPKs and their substrates, the

MAPK-activated protein kinases. *Microbiol Mol Biol Rev*. 2011;75(1):50-83.

55.   Peters S, Zitzelsperger E, Kuespert S, Iberl S, Heydn R, Johannesen S, et al. The TGF-β

System As a Potential Pathogenic Player in Disease Modulation of Amyotrophic Lateral

Sclerosis. *Front Neurol*. 2017;8:669.

56.   Zhang N, Westerhaus A, Wilson M, Wang E, Goff L, and Sockanathan S. Physiological

regulation of neuronal Wnt activity is essential for TDP-43 localization and function. *EMBO

J*. 2024.

57.   Ke H, Masoumi KC, Ahlqvist K, Seckl MJ, Rydell-Törmänen K, and Massoumi R. Nemo-like

kinase regulates the expression of vascular endothelial growth factor (VEGF) in alveolar

epithelial cells. *Sci Rep*. 2016;6:23987.

58.   Kortenjann M, Nehls M, Smith AJ, Carsetti R, Schüler J, Köhler G, et al. Abnormal bone

marrow stroma in mice deficient for nemo-like kinase, Nlk. *Eur J Immunol*.

2001;31(12):3580-7.

59.   Liu R, Khalil H, Lin SJ, Sargent MA, York AJ, and Molkentin JD. Nemo-Like Kinase (NLK) Is a

Pathological Signaling Effector in the Mouse Heart. *PLoS One*. 2016;11(10):e0164897.

60.   Weskamp K, Tank EM, Miguez R, McBride JP, Gómez NB, White M, et al. Shortened TDP43

870   isoforms upregulated by neuronal hyperactivity drive TDP43 pathology in ALS. *J Clin Invest*.

871   2020;130(3):1139-55.

872   61.   Miyaoka Y, Chan AH, Judge LM, Yoo J, Huang M, Nguyen TD, et al. Isolation of single-base

873   genome-edited human iPS cells without antibiotic selection. *Nat Methods*. 2014;11(3):291-

874   3.

875   62.   Barmada SJ, Skibinski G, Korb E, Rao EJ, Wu JY, and Finkbeiner S. Cytoplasmic

876   mislocalization of TDP-43 is toxic to neurons and enhanced by a mutation associated with

877   familial amyotrophic lateral sclerosis. *J Neurosci*. 2010;30(2):639-49.

878   63.   Malik AM, Miguez RA, Li X, Ho YS, Feldman EL, and Barmada SJ. Matrin 3-dependent

879   neurotoxicity is modified by nucleic acid binding and nucleocytoplasmic localization. *Elife*.

880   2018;7.

881   64.   Weskamp K, Safren N, Miguez R, and Barmada S. Monitoring Neuronal Survival via

882   Longitudinal Fluorescence Microscopy. *J Vis Exp*. 2019(143).

883   65.   Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR.

884   *Nucleic Acids Res*. 2001;29(9):e45.

885   66.   Ishitani T, Hirao T, Suzuki M, Isoda M, Ishitani S, Harigaya K, et al. Nemo-like kinase

886   suppresses Notch signalling by interfering with formation of the Notch active

887   transcriptional complex. *Nat Cell Biol*. 2010;12(3):278-85.

888   67.   Fu SC, Fung HYJ, Cağatay T, Baumhardt J, and Chook YM. Correlation of CRM1-NES affinity

889   with nuclear export activity. *Mol Biol Cell*. 2018;29(17):2037-44.

890   68.   CVAT.ai, and Corporation. Computer Vision Annotation Tool (CVAT) (v2.11.2).

891   doi:10.5281/zenodo.10807509.

892   69.   Cao MC, and Scotter EL. Transcriptional targets of amyotrophic lateral

893   sclerosis/frontotemporal dementia protein TDP-43 - meta-analysis and interactive

894   graphical database. *Dis Model Mech*. 2022;15(9).