Pilidium lythri disrupts plant microbiome assembly under climate

warming: Evidence for niche-specific community reassembly

Dominika Siegieda[1]

Jacek Panek[1]

S. Emilia Hannula[2]

Magdalena Frąc[1]

[1]Institute of Agrophysics, Polish Academy of Sciences, Doświadczalna 4 street, 20-290 Lublin, Poland

[2]Leiden University, Institute of Environmental Sciences, Department of Environmental Biology, Leiden,

The Netherlands

Keywords

12 *Pilidium lythri*, emerging fungal phytopathogen, microbial community dynamics, plant holobiont,

13 temperature adaptability, biotic stress in plants, microbial assembly mechanisms

Abstract

Although ongoing global warming is an important driver of phytopathogen spread to new areas, little

is known on how fungal pathogens disrupt fragile balance in plant-microbe interactions. Here we

demonstrate that thermotolerant *Pilidium lythri* destabilizes microbial communities assembly in soil

and plant niches. For our exploratory investigation, we employed high-throughput marker sequencing

of bacterial/archeal 16S and fungal ITS markers from 72 samples, combined with mixed-effects

modelling, source-tracking, microbiota assembly mechanisms investigation and network analysis. Our

analysis revealed that *P. lythri* infection disrupted microbial community assembly processes,

transitioning from deterministic to stochastic dynamics in the rhizosphere and bulk soil, aligning with

the Anna Karenina Principle. Though bacterial and fungal microbiome diversity did not change under

infection, microbial migration patterns highlighted shifts in community interactions. Network analysis

underscored *P. lythri*'s role in destabilizing fungal community connectivity, enhancing antagonistic

interactions between taxa. This study provides valuable insights into how spreading phytopathogens

modulate microbial equilibrium in the plant holobiont, with implications of managing emerging threats

in agriculture.

Introduction

Fungal and bacterial microorganisms, together with their host plants, are interconnected through intricate, multi-level interactions that shape ecosystems and sustain plant health and productivity. Based on these relationships, the plant holobiont concept was formed - an integrated system where hosts and microbiota function as one. Plants produce root exudates to modulate microbial assembly of theirs niches and vicinity, while the microbiome, in turn, is able to influence and regulate plant health, improving its resilience against abiotic and biotic stressors (Berg, 2009; Baldrian, 2016; Trivedi *et al.*, 2020; Gupta *et al.*, 2021; Marco *et al.*, 2022; Michl *et al.*, 2023). These finely tuned interactions are widely studied and hold great potential for improving agricultural productivity in a sustainable manner (Vishwakarma *et al.*, 2020; Trivedi *et al.*, 2020; Frąc *et al.*, 2022).

However, this delicate balance between microbiomes and plants is being increasingly threatened by ongoing global warming. The rise of annual global temperature and changed precipitation patterns has created the opportunity for a spread and establishment of invasive phytopathogens into new regions (Bebber *et al.*, 2013; Helfer, 2014; Liu *et al.*, 2019). Despite this urgent challenges, limited attention has been given to understand how these emerging pathogens alter and interact with microbial communities. This issue is particularly concerning for perennial crops like strawberries (*Fragaria x ananassa* Duch.), which depend heavily on stable rhizosphere and soil microbial communities to thrive and are particularly vulnerable to such disruptions.

Pilidium lythri, emerging fungal phytopathogen, causing damage to strawberry plantations, was originally reported in countries located in warmer climate zones, such as India (Phatak & Payak, 1965), Iran (Arzanlou *et al.*, 2013; Sayami *et al.*, 2013) and China (Zhang *et al.*, 2008; Duan *et al.*, 2010). Notably, its occurrence had been recently identified in several temperate climate countries, including Belgium (Debode *et al.*, 2011), Croatia (Milicević *et al.*, 2014), Spain (Aguín *et al.*, 2018), France (Cardin *et al.*, 2009; Caesar *et al.*, 2012) and Poland, although in 2004 only morphologically, but in 2020 we confirmed its presence in symptomatic Polish strawberry fields with molecular methods (Gołębniak & Jarosz, 2004; Malarczyk *et al.*, 2020), and with new isolates collected in present study. Primary targets of this fungal phytopathogen are strawberry plants and fruits (Lopes *et al.*, 2010; Geng *et al.*, 2012; Ayoubi *et al.*, 2016), but it also affects other agriculturally significant plants, including *Olea europaea* L. (Arzanlou *et al.*, 2013), *Cornus mas* (Torbati *et al.*, 2019), *Prunus domestica* (Sayami *et al.*, 2013), *Eucalyptus* spp. (Maússe-Sitoe *et al.*, 2016), *Vitis vinifera* (Aguín *et al.*, 2018), and various ornamental flowers (Zhang *et al.*, 2008; Cardin *et al.*, 2009; Caesar *et al.*, 2012; Bruckart *et al.*, 2013; Garfinkel & Chastagner, 2019). Notably concerning is that *P. lythri* is not host-specific. Strains isolated from olives were able to cause disease symptoms on strawberry fruits (Karimi *et al.*, 2016), further increasing the

severity of the threat posed by this pathogen as it spreads to new regions. Studies conducted in Brazil,

Iran and Belgium showed incidence as high as 70% in high humidity and warm temperatures,

highlighting its threat in warming climate (Lopes *et al.*, 2010; Ayoubi *et al.*, 2016; Karimi *et al.*, 2016).

Pathogenicity test revealed rapid progression of the disease, with symptoms appearing on up to 100%

of fruits within days after inoculation (Debode *et al.*, 2011; Fernández-Ortuño *et al.*, 2014). The

pathogen is able to infect both wounded and intact grape berries, further underscoring its high

virulence (Aguín *et al.*, 2018). The pathogen's ability to adapt and spread under changing

environmental conditions makes it an important microorganism to explore relationships between

emerging fungal pathogens, and host plants microbiomes. Furthermore, its prompt proliferation and

versatile host range makes it an ideal research object to evaluate how emerging pathogens disrupt

microbial balance across diverse plant microbiomes.

To explore climate adaptation and microbial interaction in emerging fungal pathogens, we isolated

Pilidium lythri from symptomatic strawberry plants in a Polish nursery. We compared the metabolic

profiles of 3 *P. lythri* strains grown in 24°C and 30°C with BIOLOG PM21-25 plates, identifying its

sensitivity to chemical substances in response to different temperatures to imitate climate scenarios.

We also investigated the impact of pathogen's infection on the composition and assembly mechanisms

of the bacterial and fungal microbiomes of soil (bulk soil, rhizosphere) and strawberry (roots) niches.

Current work significantly advances our understanding the microbial dynamics associated with this

spreading phytopathogen, offering valuable insights into broader ecological implications of emerging

fungal pathogens.

Methods

Fungal isolates and molecular identification

Pure fungal isolates were obtained by placing symptomatic strawberry leaves and fruits (surface-

sterilized with detergent and 70% ethanol) on PDA medium and incubating them at 24°C in darkness

for 7 days. After initial screening, fungi with macroscopic and microscopic characteristics matching the

description of *Pilidium lythri* (Palm, 1991; Karimi *et al.*, 2019) were identified through amplification and

Sanger sequencing of the Internal Transcribed Spacer (ITS) regions of fungal rDNA, with ITS1 and ITS4

primers (White *et al.*, 1990), as previously described (Frąc *et al.*, 2015). The obtained sequences were

deposited in GenBank under accession numbers: PQ632564.1, PQ632565.1, PQ632566.1, PQ632567.1.

These sequences were then used to construct phylogenetic tree with other sequences available in

Genbank database from the *Pilidium* genus. Initial tree for the heuristic search was obtained

automatically by applying Neighbor-Join and BioNJ algorithms to a matrix of pairwise distances

estimated using the Tamura-Nei model, and then selecting the topology with superior log likelihood

value. The tree was drawn to scale, with branch lengths measured in the number of substitutions per

site. This analysis involved 57 nucleotide sequences and there were a total of 372 positions in the final

dataset. Evolutionary analyses were conducted in MEGA X (Kumar *et al.*, 2018). To fulfil Koch's

postulates, a spore suspension of 1.6×10^4 spores/mL was prepared, and surface-sterilized strawberry

leaves were briefly submerged in the solution. The leaves were then sealed in ziplock bags and

incubated at 24°C in darkness for 7 days. Sterile deionized water was used as the untreated control.

Metabolic profile arrays

Each cell of Phenotype Microarray 21-25 BIOLOG plates (Biolog, Hayward, CA, USA) was inoculated

with the fungal mycelium as recommended by the manufacturer and described before (Panek *et al.*,

2016). For each plate type, 3 *Pilidium lythri* strains in triplicates and, two temperatures (24 and 30°C)

and 4 concentrations were incubated for 96 hours in the darkness. Microplate reader (Biolog, Hayward,

CA, USA) was used to determine the optical density at 490 nm (to establish catabolic activity of fungal

strains) at the beginning of the experiment and every 24 hours. Obtained data was analysed

statistically and visualised in RStudio, using tidyverse (Wickham *et al.*, 2019) and lme4 (Bates *et al.*,

2015) packages. First, values obtained at 0 h for each read was divided from read values at 96 h to

establish growth of respiration rates, and the data was rescaled from 0 to 1. To establish which factors

had an influence on fungal activity, we used generalized linear mixed-effects model (GLLM) with

Gamma family, controlling for the plate and plate, and to establish that inhibit its respiration rates, we

controlled for plate, strain, temperature and concentration.

Pot experiment and isolation of microbial DNA

Strawberry plants, obtained from the nursery, were grown in pots containing sieved soil from a

certified organic strawberry plantation. After 20 days of initial growth of strawberry plants at 28°C and

70% humidity, half of the pots were inoculated with *P. lythri* spores using 10 mL of a $2 \cdot 10^4$ spores mL$^-$

^{1} solution per plant. Sterile, deionized water was used as a negative control. Samples were collected

from two pots in triplicates (6 samples per niche per time point) for healthy and pathogen-inoculated

samples in two timepoints: at the time of inoculation and again after 9 days of incubation. Bulk soil

was transferred into sterile 1.5 mL Eppendorf tubes and frozen at -20°C. The rhizosphere was collected

as described by Lundberg et al. (Lundberg *et al.*, 2012). Roots, flushed free of rhizosphere soil, were

also frozen at -20°C prior to DNA isolation which was conducted with the QIAQUBE instrument (Qiagen,

Hilden, Germany) and GeneMATRIX Soil DNA Purification Kit (EURx, Gdańsk, Poland) after

homogenization of the samples in the FastPrep-24 instrument (MP Biomedicals) at 4 m s^{-1} for 15 s.

DNA isolates were then frozen in -20°C until library construction and sequencing.

Metataxonomic sequencing, bioinformatics and statistical analysis

For the metataxonomic sequencing, microbial DNA ITS2 (Schoch *et al.*, 2012) and 16Sv3 regions (Klindworth *et al.*, 2013) were amplified, indexed and sequenced on Illumina MiSeq instrument (2x300). Raw data was deposited at SRA database under accession numbers: PRJNA1213298 for 16S and PRJNA1213307 for ITS. Raw data was then passed through the custom QIIME2 (Bolyen *et al.*, 2019) pipelines as reported before (Siegieda *et al.*, 2023, 2024). In brief, after primer trimming with Cutadapt (Martin, 2011) and ITSxpress (Rivers *et al.*, 2018), denoising, chimera removal, merging and Amplicon Sequence Variant (ASV) calling was made. The taxonomy was assigned based on UNITE v9 dynamic release (Nilsson *et al.*, 2019) and Silva 138 (Pruesse *et al.*, 2007) databases. The results were then analysed with R (v. 4.0.3), using libraries: phyloseq (McMurdie & Holmes, 2013), tidyverse (Wickham *et al.*, 2019), microeco (Liu *et al.*, 2021), microbiomeMarker (Cao *et al.*, 2022), NetCoMi (Peschel *et al.*, 2021). First, for ITS data, only ASVs identified to Fungi and Stramenopiles; and for 16S - Bacteria and Archaea. Then, ITS data was repeatedly rarefied at 6,000 and 16S at 11,000 reads using mirlyn package (McMurdie & Holmes, 2014; Cameron *et al.*, 2021). We employed Effective Number of Species (ENS) (Jost, 2006) on ASV level to evaluate the richness of microbiomes in each niche. We then used Linear Mixed-Effect Models with controlling for the plant to compare differences between control and infected samples. We used supervised machine learning model to evaluate the fungal biomarkers of each niche for healthy and infected samples with Supervised Machine Method (SVM), using Total Sum Scaling normalization method. SourceTracker was used for tracking migration between niches (Knights *et al.*, 2011). RCbray (Bray-Curtis dissimilarity based Raup-Crick metric) and βNTI (β Nearest Taxon) were used to evaluate the ecological processes that dominated the bacterial assemblies (Stegen *et al.*, 2013; Liu *et al.*, 2017). When βNTI fell between -2 and 2, microbial assembly was dominated by stochastic mechanisms, and below -2 and higher that 2 - by deterministic. Microbial networks were constructed with 200 most frequent taxa and SPIEC-EASI (Sparse Inverse Covariance Estimation for Ecological Association Inference) method (Kurtz *et al.*, 2015), using glasso (Graphical Least Absolute Shrinkage and Selection Operator) approach. The dataset was resampled 30 times and 20 penalty levels with 1e-4 as the smallest ratio, taking the average of symmetric edges. Fast greedy clustering method was used for network analysis.

Results

Fungal isolates

Fungal ITS sequences, obtained from symptomatic strawberry leaves (G320/23, G321/23, G322/23 and G323/23) showed 100% similarity to those of *Pilidium lythri* already present in the GenBank database and phylogenetic tree revealed clustering of the sequences with other representatives of *Pilidium lythri*

and *Pilidium concavum*. The strains collected after 7 days in 23°C in the darkness on Potato Dextrose

Agar were flat and granulose, cinnamon in colour (Karimi *et al.*, 2016; Marin-Felix *et al.*, 2017). The

presence of hyaline conidia with smooth walls and slightly pointed tips were confirmed under the

microscope. After the inoculation of the surface-sterilized strawberry leaves with the spore suspension

and incubation in 24°C in darkness for 7 days, 100% of the leaves showed brown necrotic lesions.

Leaves submerged in water did not show symptoms of the disease. Growth of the cinnamon-colour

fungal strains were observed after incubation (24°C for 7 days) of infested leaves on the Petri dishes

with the PDA medium, fulfilling he Koch's postulate (Figure 1).

Figure 1. A) Phylogenetic tree of the evolutionary history of *P. lythri* was inferred by using the Maximum Likelihood method

and Tamura-Nei model (Tamura & Nei, 1993) and the e tree with the highest log likelihood (-1144.51) is shown. The

percentage of trees in which the associated taxa clustered together is shown next to the branches. B,C) Symptoms of the

Pilidium lythri infection on the strawberry leaves and in the D) control leaf. E) Growth of *P. lythri* after 7 days of growth in 24

°C on PDA medium, F) conidia of *P. lythri* (40x), and G) growth on PDA medium with infected leaf.

Chemical sensitivity

To evaluate *Pilidium lythri*'s sensitivity to various chemicals, BIOLOG PM 21-25 plates were we

incubated with the spore and mycelium solution of three fungal strains at 24°C and 30°C for 96 hours

(in triplicates, thus 9 replications for each substrate in both temperatures). The plates are coated with

4 different concentrations of the chemical substances. The precise concentrations of the substances

used for microarrays are confidential and protected as proprietary information by the manufacturer,

thus, for the purposes of this article, we refer to them as concentration levels 1, 2, 3, and 4 to maintain

clarity. An evaluation of the model revealed that, as expected, the temperature growth was positively

correlated with fungal respiration growth, also substance concentration growth had negative

correlation with the fungal respiration. We observed no effect of replicate; and different response of

one of strains - G322/23 was characterised with higher respiration rates compared to other 2

(Supplementary Figure 1). We also evaluated substances that negatively impacted respiration rate

growth (Figure 2). Top inhibitors included thiourea, dodecyltrimethyl ammonium bromide, tetrazolium

violet, alexidine and fluorodeoxyuridine.

Figure 2. BIOLOG PM21-25 results of the substances that differently influence growth of *Pilidium lythri.* A) Volcano plot

visualizing biological and statistical effect for different substances. Gray circles represent no significant importance, light blue

- significant results. Dashed vertical lines represent p.value of 0.05 and estimate of -log10(0.05). B) Bar plot representing

estimate for each substance significantly inhibiting pathogens respiration rate in tested conditions.

Fungal and bacterial community changes after *P. lythri* inoculation of strawberry

A total of 72 samples from soil (24), rhizosphere (24), and roots (24) were amplified and sequenced.

Rarefaction curves plateaued for all samples, indicating sufficient sequencing depth. To evaluate the

presence of *Pilidium lythri* in infected samples, we employed a predictive approach to identify 15 most

important bacterial and fungal taxa that differentiate healthy and infected strawberry compartments.

The analysis revealed that the *Pilidium* fungi belonged to the most important taxa that differentiate

between healthy and infected plants and soil (Figure 3). This, along with the visible symptoms of the

disease, confirmed that the infection was successful and the pathogen migrated from the soil to

rhizosphere and roots.

Figure 3. Top 15 fungal taxa found more abundant in (orange) infected and in (blue) healthy soils and plants identified based on Supervised Vector Machine machine learning approach.

After evaluating that infection was successful, statistical analysis was conducted to compare the alpha diversity of bacterial and fungal communities between healthy and inoculated (infected) niches. No significant differences in richness were observed between healthy and infected niches at both sampling times (p > 0.05) for bacteria and fungi (Figure 4). Notably, bacterial communities showed a significant increase richness across all three compartments at the second sampling time, whereas fungal community richness remained stable across time points. Additionally, both bacterial and fungal communities exhibited a gradient of decreasing diversity from soil to rhizosphere and root at second sampling.

Figure 4. Comparison of alpha diversity (in Effective Number of Species) of bacterial (A) and fungal (B) communities of strawberry niches immediately after *P. lythri* inoculation (0 days) and after 9 days of incubation in 25°C and ~70% of humidity (two). LMER model was used for comparison of non-independent samples. Light blue color represents samples without pathogen inoculation, and orange - with inoculation.

To elucidate the influence of *Pilidium lythri* on microbial migration between the niches, we used

samples from sampling at 0 days as a source, and samples from sampling at 9 days as sinks. In bulk soil,

only bacterial community revealed significantly higher share of microorganisms in bulk soil and a lower

portion from unknown source in infected samples (Figure 5A). Furthermore, we observed migration of

bacteria from bulk soil to rhizosphere for both bacteria and fungi in infected communities, but also

from rhizosphere to rhizosphere for bacteria in infected communities. Finally, analysis revealed

increased migration from rhizosphere to roots in infected samples in bacterial community.

Figure 5. Source tracking analysis (A) of bacterial and fungal microbiomes between sampling one (source) and two (sink).

Comparison between healthy (light blue, left) and contaminated (light orange, right) samples. Values are shown as mean

standard error from bootstrap. (B) BetaNRI idenx analysis of bacterial community in second sampling. Significant

differences on both panels between healthy and contaminated samples are shown with asterisk.

We also evaluated the mechanisms involved in bacterial community assembly by calculating the

betaNRI index for bulk soil, rhizosphere, and root samples collected during the second sampling period

(Figure 4B). The results revealed a dominance of stochastic processes in both healthy and infected

rhizosphere and root samples of strawberry (betaNRI values between -2 and 2). Furthermore, we

observed differences in community assembly processes between healthy and infected bulk soil, with

a shift from deterministic processes (characterized by lower turnover) in healthy bulk soil to a

dominance of stochastic processes in infected soil.

Finally, to evaluate the influence of *Pilidium lythri* on the structure of fungal networks, we compared

the network structure in the rhizosphere niche between control and infected samples during the

second sampling (Figure 6A). Both networks showed similar characteristics in terms of the number of

components (1), clustering coefficient (0.995, 0.999), modularity (-0.007, -0.008), edge density (0.994,

0.999) and natural connectivity (0.375, 0.310), average dissimilarity and average path length (0.6, 0.6).

On the other hand, the control rhizosphere exhibited a higher percentage of positive edges (52.308%

vs. 43.344%) but lower vertex and edge connectivity (58, 71).

Figure 6. Fungal network analysis in A) control and B) infected rhizosphere samples of strawberry. Each node represents

individual fungal ASV, and node size is proportional to number of reads. Node colours represent the same clusters. Green

lines represent positive interactions and red- negative. The position of the same nodes is the same in both networks.

Notably, changes were observed in centrality of networks (Jaccard Index). Degree, betweenness and

hub taxa measures showed no overlap in most connected, keystone and key hub taxa, significant shift

was observed in closeness of most central nodes (Table 1).

Table 1. Centrality changes between networks of rhizosphere samples in control and *Pilidium* contaminated samples. Asterisk

shows statistically significant change.

	Degree	Betweenness	Closeness	Eigenvector	Hub taxa
Jaccard Index	0	0	0.071	0.209	0
p-value (P≤Jacc)	1	0.296	0.027*	0.055	1

Discussion

To close the knowledge gap regarding emerging fungal pathogens and theirs influence on microbiomes

of plants we used newly isolated strains of *Pilidium lythri*. We identified 3 of them, that were obtained

from symptomatic strawberry plants in Polish nursery, by sequencing of the ITS region. We then

evaluated the impact of temperature growth on inhibition of fungal development with 112 different

chemical substances in 4 concentrations. We then evaluated the impact of this phytopathogen on the

structure, assembly processes and association networks of bacterial and fungal metacommunities in

soil and plant niches (bulk soil, rhizosphere, roots) of strawberry.

In vitro evaluation of the impact of various chemical substances on the growth and development of

the tested phytopathogen at two temperatures (24 and 30 °C) using BIOLOG microarrays raised serious

concerns, as *Pilidium lythri* demonstrated overall higher respiration rates at higher temperatures.

Global warming has indeed been widely reported to contribute to the spread of fungal phytopathogens

to new areas (Ward *et al.*, 2007; Bebber *et al.*, 2013; Chaloner *et al.*, 2021). Among the wide variety of

chemical substances tested, some of them presented inhibition mechanism on pathogens respiration

rates. Top 5 analysed substances likely disrupted electron chain, destabilized cell membranes, and

inhibited DNA/RNA synthesis (Waldorf & Polak, 1983; Mamouei *et al.*, 2018; Ansari *et al.*, 2024; Yang

et al., 2025).

While the dissemination of fungal pathogens is widely reported (Bebber *et al.*, 2013), still little is known

about the effect of such invasive microorganisms on the microbial structure of the plant holobiont.

Recent studies have reported significant differences in the α-diversity of microbiome structure

between healthy and infected soil and plant niches (Lefcheck *et al.*, 2015; Delgado-Baquerizo *et al.*,

2016; Bakker *et al.*, 2018). In the present study, we compared the diversity of bacterial and fungal

metacommunities in healthy and *Pilidium lythri*-infected bulk soil, rhizosphere, and roots at two time

points: directly after pathogen inoculation and 9 days post-inoculation. The results revealed an

evidence of increased α-diversity in the bacterial community 9 days after inoculation in both healthy

and inoculated niches, suggesting that bacterial community diversity are less influenced by *Pilidium*

lythri inoculation and more are responsive to environmental factors. To test this, we analyzed the

mechanisms involved in bacterial microbiome assembly by calculating the βNRI index, treating the first

sampling as the source of microorganisms and the second sampling as the sink. This index measures

how the differences in species composition between the two samplings differ from what would be

expected by the null model of random assembly. The results indicated that composition of bacterial

communities in healthy and diseased rhizosphere and roots samples was shaped by stochastic

processes (i.e. random colonization, dispersal, and ecological drift). Interestingly, we observed a shift

from deterministic processes in healthy bulk soil, to stochastic processes in *Pilidium lythri*-infected

samples, indicating a disruption in microbial turnover. Infected soil showed random processes that

could have been triggered by disturbances in plant biology caused by the infection, described by the

Anna Karenina Principle (AKP) (Arnault *et al.*, 2023). The scenario described here can be explained by

the loss of plant control over microbiome assembly, resulting in a temporal AKP effect (Cheng *et al.*,

2013, 2019). Over time, this could lead to the recovery of the plant's control over microbial assembly

or the introduction of new deterministic processes to control bacterial assembly (Gao *et al.*, 2021).

Analysis of microbial migration suggests increased migration of bacteria from the soil in diseased

samples, as well as increased migration of bacteria from the rhizosphere to the roots in infected plant

samples. This inconsistency between migration (suggesting plant filtering) and microbial assembly

(suggesting random processes) could be caused by the smaller scale of deterministic processes being

masked by larger-scale microbial community assembly. Furthermore, these two mechanisms

represent different time scales, where βNRI focuses on broader, longer-term assembly processes

(Arnault *et al.*, 2023). Importantly, migration of microorganisms can be caused by random dispersal

(stochastic processes) but may appear deterministic as migration increases. Thus, these two processes
are not necessarily contradictory.

In the fungal microbiome, we observed observed increased migration of fungi from bulk soil to the
rhizosphere, highlighting the importance of this niche in fungal microbiome dynamics and the role of
plant in microbial recruitment under pathogen stress (Bais *et al.*, 2006; Berendsen *et al.*, 2012). To
further asses the influence of the pathogen in microbial metacommunity, we analyzed the fungal
networks in the rhizosphere. A number of methods for construction and analysis of co-occurrence
networks had been developed, and SPIEC-EASI is characterized with important advantages, such as
accounting for compositional type of data and mitigation of non-independence of samples. Also, an
implementation of glasso approach allowed for the analysis in high-dimensional setting with more
variables than samples (Kurtz *et al.*, 2015). Thus, we implemented these methods for the analysis in
current study. Comparison of healthy and infected samples revealed shift in microbial fungal networks
under *Pilidium lythri* influence. The fungal networks in infected rhizospheres exhibited lower
percentages of positive edges caused lower robustness of networks, suggesting restructuring of fungal
community in response to pathogen presence. These findings underline decrease in natural
connectivity percentage of positive edges, that suggest lowered positive interactions between fungal
taxa. While the presence of the pathogen did not significantly disrupt fungal networks, it did decrease
network connectivity, which indicates a shift toward increased competition and antagonism within the
fungal microbiome. This suggests adaptation of the fungal meta-community, making it more resistant
to further infections and environmental stressors (Hernandez *et al.*, 2021).

Concluding, in this study, we isolated and molecularly identified isolates of spreading fungal
phytopathogen from symptomatic strawberry plants - *Pilidium lythri*. We observed pathogen's low
sensitivity against various chemical substances in 30°C compared with 24°C, highlighting dangers of its
spread due to climate change. We also noted significant variability in inhibition caused by chemical
substances between strains. We used *P. lythri* to evaluate ecological impacts of spreading fungal
phytopathogen on microorganism metacommunities in bulk soil, rhizosphere and roots of strawberry
with high-throughput marker sequencing. We revealed that *Pilidium lythri* infection had a major effect
on both the structure and dynamics of the plant associated microbiome. Its ability to shift bacterial
microbial assembly mechanism from deterministic to stochastic in the bulk soil, underscores its
capacity to destabilize plant-associated microbiomes. Additionally, we observed that *P. lythri* shifted
the fungal microbiome dynamics in the strawberry rhizosphere towards reducing mutualistic
connections, while maintaining overall network stability. This underlined adaptive reorganization of
fungal community, potentially increasing resilience to secondary infections and other (a)biotic
stresses.

These findings underline the critical role of emerging fungal pathogens like *Pilidium lythri* in destabilizing plant-microbiome interactions. This is significantly important in the context of global warming, which accelerates spread and impact of such pathogens.

References

Aguín, O., Ferreiroa, V., González-Jartín, J. M., Alfonso, A., Botana, L. M., Mansilla, J. P., & Sainz, M. J. (2018). *Pilidium lythri* is associated with bunch rot of grapevine (Vitis vinifera). *American Journal of Enology and Viticulture*, *69*(4), 410–416. https://doi.org/10.5344/ajev.2018.17093

Arnault, G., Mony, C., & Vandenkoornhuyse, P. (2023). Plant microbiota dysbiosis and the Anna Karenina Principle. *Trends in Plant Science*, *28*(1), 18–30. https://doi.org/10.1016/j.tplants.2022.08.012

Arzanlou, M., Torbati, M., & Jafary, H. (2013). Fruit rot on olive caused by *Pilidium concavum* in Iran. *Australasian Plant Disease Notes*, *8*(1), 117–121. https://doi.org/10.1007/s13314-013-0111-0

Ayoubi, N., Soleimani, M. J., & Zare, R. (2016). Pilidium concavum, causing tan-brown rot on strawberry in Iran. *Journal of Plant Pathology*, *98*(3), 667–669. https://doi.org/10.4454/JPP.V98I3.063

Bais, H. P., Weir, T. L., Perry, L. G., Gilroy, S., & Vivanco, J. M. (2006). The role of root exudates in rhizosphere interactions with plants and other organisms. *Annual Review of Plant Biology*, *57*(1), 233–266. https://doi.org/10.1146/annurev.arplant.57.032905.105159

Bakker, P. A. H. M., Pieterse, C. M. J., de Jonge, R., & Berendsen, R. L. (2018). The Soil-Borne Legacy. *Cell*, *172*(6), 1178–1180. https://doi.org/10.1016/j.cell.2018.02.024

Baldrian, P. (2016). Forest microbiome: diversity, complexity and dynamics. *FEMS Microbiology Reviews*, fuw040. https://doi.org/10.1093/femsre/fuw040

Bebber, D. P., Ramotowski, M. A. T., & Gurr, S. J. (2013). Crop pests and pathogens move polewards in a warming world. *Nature Climate Change*, *3*(11), 985–988. https://doi.org/10.1038/nclimate1990

Berendsen, R. L., Pieterse, C. M. J., & Bakker, P. A. H. M. (2012). The rhizosphere microbiome and plant health. *Trends in Plant Science*, *17*(8), 478–486. https://doi.org/10.1016/j.tplants.2012.04.001

Berg, G. (2009). Plant–microbe interactions promoting plant growth and health: perspectives for controlled use of microorganisms in agriculture. *Applied Microbiology and Biotechnology*, *84*(1), 11–18. https://doi.org/10.1007/s00253-009-2092-7

Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., Alexander, H., Alm, E. J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J. E., Bittinger, K., Brejnrod, A., Brislawn, C.

J., Brown, C. T., Callahan, B. J., Caraballo-Rodríguez, A. M., Chase, J., … Caporaso, J. G. (2019). Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nature Biotechnology*, *37*(8), 852–857. https://doi.org/10.1038/s41587-019-0209-9

Bruckart, W. L., Eskandari, F. M., Coombs, E. M., Rossman, A. Y., & Palm, M. E. (2013). First report of Pilidium concavum causing leaf necrosis on Fallopia japonica in the United States. *Plant Disease*, *97*(1), 146–146. https://doi.org/10.1094/PDIS-07-12-0688-PDN

Caesar, A. J., Lartey, R. T., & Caesar-TonThat, T. (2012). First report of *Pilidium concavum* causing root lesions of meadow hawkweed in France. *Plant Disease*, *96*(12), 1830–1830. https://doi.org/10.1094/PDIS-05-12-0509-PDN

Cameron, E. S., Schmidt, P. J., Tremblay, B. J. M., Emelko, M. B., & Müller, K. M. (2021). Enhancing diversity analysis by repeatedly rarefying next generation sequencing data describing microbial communities. *Scientific Reports*, *11*(1), 1–13. https://doi.org/10.1038/s41598-021-01636-1

Cao, Y., Dong, Q., Wang, D., Zhang, P., Liu, Y., & Niu, C. (2022). microbiomeMarker: an R/Bioconductor package for microbiome marker identification and visualization. *Bioinformatics*, *38*(16), 4027–4029. https://doi.org/10.1093/bioinformatics/btac438

Cardin, L., Vincenot, L., & Balesdent, M. H. (2009). First report of Pilidium concavum on Bergenia crassifolia in France. *Plant Disease*, *93*(5), 548–548. https://doi.org/10.1094/PDIS-93-5-0548B

Chaloner, T. M., Gurr, S. J., & Bebber, D. P. (2021). Plant pathogen infection risk tracks global crop yields under climate change. *Nature Climate Change*, *11*(8), 710–715. https://doi.org/10.1038/s41558-021-01104-8

Cheng, C., Gao, X., Feng, B., Sheen, J., Shan, L., & He, P. (2013). Plant immune response to pathogens differs with changing temperatures. *Nature Communications*, *4*(1), 2530. https://doi.org/10.1038/ncomms3530

Cheng, Y. T., Zhang, L., & He, S. Y. (2019). Plant-Microbe Interactions Facing Environmental Challenge. *Cell Host & Microbe*, *26*(2), 183–192. https://doi.org/10.1016/j.chom.2019.07.009

Debode, J., Van Hemelrijck, W., Heungens, K., Maes, M., & Creemers, P. (2011). First Report of Pilidium concavum Causing Tan-Brown Rot on Strawberry Fruit in Belgium. *Plant Disease*, *95*(8), 1029–1029. https://doi.org/10.1094/PDIS-10-10-0752

Delgado-Baquerizo, M., Maestre, F. T., Reich, P. B., Jeffries, T. C., Gaitan, J. J., Encinar, D., Berdugo, M., Campbell, C. D., & Singh, B. K. (2016). Microbial diversity drives multifunctionality in terrestrial ecosystems. *Nature Communications*, *7*, 1–8. https://doi.org/10.1038/ncomms10541

Duan, Y. B., Kang, Y. B., & Yu, Z. Z. (2010). First report of *Pilidium concavum* on *Paeonia suffruticosa* in
China. *Plant Disease*, *94*(2), 271–271. https://doi.org/10.1094/PDIS-94-2-0271B

Fernández-Ortuño, D., Bryson, P. K., & Schnabel, G. (2014). First Report of Pilidium concavum Causing
Tan-brown Rot on Strawberry Nursery Stock in South Carolina. *Plant Disease*, *98*(7), 1010–1010.
https://doi.org/10.1094/PDIS-01-14-0048-PDN

Frąc, M., Hannula, E. S., Bełka, M., Salles, J. F., & Jedryczka, M. (2022). Soil mycobiome in sustainable
agriculture. *Frontiers in Microbiology*, *13*(November), 1–9.
https://doi.org/10.3389/fmicb.2022.1033824

Frąc, M., Jezierska-Tys, S., & Yaguchi, T. (2015). Occurrence, detection, and molecular and metabolic
characterization of heat-resistant fungi in soils and plants and their risk to human health.
Advances in Agronomy, *132*, 161–204.

Gao, M., Xiong, C., Gao, C., Tsui, C. K. M., Wang, M. M., Zhou, X., Zhang, A. M., & Cai, L. (2021). Disease-
induced changes in plant microbiome assembly and functional adaptation. *Microbiome*, *9*(1), 1–
18. https://doi.org/10.1186/s40168-021-01138-2

Garfinkel, A. R., & Chastagner, G. A. (2019). Survey reveals a broad range of fungal pathogens and an
oomycete on peonies in the United States. *Plant Health Progress*, *20*(4), 250–254.
https://doi.org/10.1094/PHP-09-19-0065-S

Geng, W. L., Hu, P., Ma, Z., Zhao, X. Y., & Wei, Y. M. (2012). First Report of Pilidium concavum Causing
Tan-Brown Leaf Spot on Strawberry in China. *Plant Disease*, *96*(9), 1377–1377.
https://doi.org/10.1094/PDIS-03-12-0310-PDN

Gołębniak, B., & Jarosz, A. (2004). First report of tan-brown rot (*Hainesia lythri*) on strawberry fruits in
Poland. *Phytopathogia Polonica*, *31*, 57–60.

Gupta, R., Anand, G., Gaur, R., & Yadav, D. (2021). Plant–microbiome interactions for sustainable
agriculture: a review. *Physiology and Molecular Biology of Plants*, *27*(1), 165–179.
https://doi.org/10.1007/s12298-021-00927-1

Helfer, S. (2014). Rust fungi and global change. *New Phytologist*, *201*(3), 770–780.
https://doi.org/10.1111/nph.12570

Hernandez, D. J., David, A. S., Menges, E. S., Searcy, C. A., & Afkhami, M. E. (2021). Environmental stress
destabilizes microbial networks. *ISME Journal*, *15*(6), 1722–1734.
https://doi.org/10.1038/s41396-020-00882-x

Jost, L. (2006). Entropy and diversity. *Oikos*, *113*(2), 363–375. https://doi.org/10.1111/j.2006.0030-
1299.14714.x

Karimi, K., Arzanlou, M., Babai-ahari, A., & Pertot, I. (2016). Biological and molecular characterisation
of *Pilidium lythri*, an emerging strawberry pathogen in Iran. *Phytopathologia Mediterranea*, *55*(3),
366–379. https://doi.org/10.14601/Phytopathol_Mediterr-18391

Karimi, K., Arzanlou, M., & Pertot, I. (2019). Development of novel species-specific primers for the
specific identification of *Colletotrichum nymphaeae* based on conventional PCR and LAMP
techniques. *European Journal of Plant Pathology*. https://doi.org/10.1007/s10658-019-01895-9

Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., & Glöckner, F. O. (2013).
Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation
sequencing-based   diversity   studies.   *Nucleic   Acids   Research*,   *41*(1),   e1–e1.
https://doi.org/10.1093/nar/gks808

Knights, D., Kuczynski, J., Charlson, E. S., Zaneveld, J., Mozer, M. C., Collman, R. G., Bushman, F. D.,
Knight, R., & Kelley, S. T. (2011). Bayesian community-wide culture-independent microbial source
tracking. *Nature Methods*, *8*(9), 761–765. https://doi.org/10.1038/nmeth.1650

Lefcheck, J. S., Byrnes, J. E. K., Isbell, F., Gamfeldt, L., Griffin, J. N., Eisenhauer, N., Hensel, M. J. S.,
Hector,   A.,   Cardinale,   B.   J.,   &   Duffy,   J.   E.   (2015).   Biodiversity   enhances   ecosystem
multifunctionality   across   trophic   levels   and   habitats.   *Nature   Communications*,   *6*.
https://doi.org/10.1038/ncomms7936

Liu, C., Cui, Y., Li, X., & Yao, M. (2021). microeco: an R package for data mining in microbial community
ecology. *FEMS Microbiology Ecology*, *97*(2). https://doi.org/10.1093/femsec/fiaa255

Liu, C., Yao, M., Stegen, J. C., Rui, J., Li, J., & Li, X. (2017). Long-term nitrogen addition affects the
phylogenetic turnover of soil microbial community responding to moisture pulse. *Scientific
Reports*, *7*(1), 1–11. https://doi.org/10.1038/s41598-017-17736-w

Liu, X., Ma, Z., Cadotte, M. W., Chen, F., He, J.-S., & Zhou, S. (2019). Warming affects foliar fungal
diseases more than precipitation in a Tibetan alpine meadow. *New Phytologist*, *221*(3), 1574–
1584. https://doi.org/10.1111/nph.15460

Lopes, U. P., Zambolim, L., Lopes, U. N., Pereira, O. L., & Costa, H. (2010). First report of *Pilidium
concavum* causing tan-brown rot in strawberry fruits in Brazil. *Plant Pathology*, *59*, 1171.
https://doi.org/10.1111/j.1365-3059.2010.02336.x

Lundberg, D. S., Lebeis, S. L., Paredes, S. H., Yourstone, S., Gehring, J., Malfatti, S., Tremblay, J.,
Engelbrektson, A., Kunin, V., Rio, T. G. Del, Edgar, R. C., Eickhorst, T., Ley, R. E., Hugenholtz, P.,
Tringe, S. G., & Dangl, J. L. (2012). Defining the core *Arabidopsis thaliana* root microbiome.
Nature, *488*(7409), 86–90. https://doi.org/10.1038/nature11237

Malarczyk, D. G., Panek, J., & Frąc, M. (2020). Triplex Real-Time PCR Approach for the Detection of
Crucial Fungal Berry Pathogens—*Botrytis* spp., *Colletotrichum* spp. and *Verticillium* spp.
International Journal of Molecular Sciences, *21*(22), 8469. https://doi.org/10.3390/ijms21228469

Marco, S., Loredana, M., Riccardo, V., Raffaella, B., Walter, C., & Luca, N. (2022). Microbe-assisted crop
improvement: a sustainable weapon to restore holobiont functionality and resilience.
Horticulture Research, *9*(October). https://doi.org/10.1093/hr/uhac160

Marin-Felix, Y., Groenewald, J. Z., Cai, L., Chen, Q., Marincowitz, S., Barnes, I., Bensch, K., Braun, U.,
Camporesi, E., Damm, U., de Beer, Z. W., Dissanayake, A., Edwards, J., Giraldo, A., Hernández-
Restrepo, M., Hyde, K. D., Jayawardena, R. S., Lombard, L., Luangsa-ard, J., … Crous, P. W. (2017).
Genera of phytopathogenic fungi: GOPHY 1. *Studies in Mycology*, *86*, 99–216.
https://doi.org/10.1016/j.simyco.2017.04.002

Martin, M. (2011). Cutadapt removes adapter sequences from hight-throughput sequencing reads.
EMBnet.Journal, *17*(1), 10–12. https://doi.org/https://doi.org/10.14806/ej.17.1.200

Maússe-Sitoe, S. N., Chen, S., Wingfield, M. J., & Roux, J. (2016). Diseases of eucalypts in the central
and northern provinces of Mozambique. *Southern Forests: A Journal of Forest Science*, *78*(3),
169–183. https://doi.org/10.2989/20702620.2015.1126780

McMurdie, P. J., & Holmes, S. (2013). Phyloseq: An R Package for Reproducible Interactive Analysis and
Graphics of Microbiome Census Data. *PLoS ONE*, *8*(4).
https://doi.org/10.1371/journal.pone.0061217

McMurdie, P. J., & Holmes, S. (2014). Waste Not, Want Not: Why Rarefying Microbiome Data Is
Inadmissible. *PLoS Computational Biology*, *10*(4), e1003531.
https://doi.org/10.1371/journal.pcbi.1003531

Michl, K., Berg, G., & Cernava, T. (2023). The microbiome of cereal plants: The current state of
knowledge and the potential for future applications. *Environmental Microbiome*, *18*(1), 28.
https://doi.org/10.1186/s40793-023-00484-y

Milicević, T., Kaliterna, J., Ivić, D., & Milović, M. (2014). Records of phytopathogenic fungal species on
native plants new to Croatia. *Natura Croatica*, *23*(1), 179–187.

Nilsson, R. H., Larsson, K. H., Taylor, A. F. S., Bengtsson-Palme, J., Jeppesen, T. S., Schigel, D., Kennedy,
P., Picard, K., Glöckner, F. O., Tedersoo, L., Saar, I., Kõljalg, U., & Abarenkov, K. (2019). The UNITE
database for molecular identification of fungi: Handling dark taxa and parallel taxonomic
classifications.   *Nucleic   Acids   Research*,   *47*(D1),   D259–D264.
https://doi.org/10.1093/nar/gky1022

Palm, M. E. (1991). Taxonomy and Morphology of the Synanamorphs *Pilidium concavum* and *Hainesia*
lythri (Coelomycetes). *Mycologia*, *83*(6), 787. https://doi.org/10.2307/3760436

Panek, J., Frąc, M., & Bilińska-Wielgus, N. (2016). Comparison of chemical sensitivity of fresh and long-
stored heat resistant *Neosartorya fischeri* environmental isolates using Biolog Phenotype
MicroArray system. *PLoS ONE*, *11*(1), 4–8. https://doi.org/10.1371/journal.pone.0147605

Peschel, S., Müller, C. L., Von Mutius, E., Boulesteix, A. L., & Depner, M. (2021). NetCoMi: Network
construction and comparison for microbiome data in R. *Briefings in Bioinformatics*, *22*(4), 1–18.
https://doi.org/10.1093/bib/bbaa290

Phatak, H. C., & Payak, M. M. (1965). *Hainesia lythri* on strawberry - a new record for India. *Indian*
Phytopathology, *18*, 237–239.

Pruesse, E., Quast, C., Knittel, K., Fuchs, B. M., Ludwig, W., Peplies, J., & Glöckner, F. O. (2007). SILVA:
A comprehensive online resource for quality checked and aligned ribosomal RNA sequence data
compatible   with   ARB.   *Nucleic   Acids   Research*,   *35*(21),   7188–7196.
https://doi.org/10.1093/nar/gkm864

Rivers, A. R., Weber, K. C., Gardner, T. G., Liu, S., & Armstrong, S. D. (2018). ITSxpress: Software to
rapidly trim internally transcribed spacer sequences with quality scores for marker gene analysis.
F1000Research, *7*(0), 1418. https://doi.org/10.12688/f1000research.15704.1

Sayami, M., Babaeizad, V., & Fathi, J. (2013). First report of *Pilidium concavum* as the causative agent
of tan-brown rot of *Prunus domestica* fruits. *Journal of Plant Pathology*, *95*(2), 451.

Schoch, C. L., Seifert, K. A., Huhndorf, S., Robert, V., Spouge, J. L., Levesque, C. A., Chen, W., Bolchacova,
E., Voigt, K., Crous, P. W., Miller, A. N., Wingfield, M. J., Aime, M. C., An, K. D., Bai, F. Y., Barreto,
R. W., Begerow, D., Bergeron, M. J., Blackwell, M., … Schindel, D. (2012). Nuclear ribosomal
internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proceedings*
of the National Academy of Sciences of the United States of America, *109*(16), 6241–6246.
https://doi.org/10.1073/pnas.1117018109

Siegieda, D., Panek, J., & Frąc, M. (2023). Plant and soil health in organic strawberry farms – Greater
importance of fungal trophic modes and networks than α -diversity of the mycobiome. *Applied*
Soil Ecology, *188*(April), 1–15. https://doi.org/10.1016/j.apsoil.2023.104925

Siegieda, D., Panek, J., & Frąc, M. (2024). Ecological processes of bacterial microbiome assembly in
healthy   and   dysbiotic   strawberry   farms.   *BMC   Plant   Biology*,   *24*(1).
https://doi.org/10.1186/s12870-024-05415-8

Stegen, J. C., Lin, X., Fredrickson, J. K., Chen, X., Kennedy, D. W., Murray, C. J., Rockhold, M. L., &
Konopka, A. (2013). Quantifying community assembly processes and identifying features that
impose them. *ISME Journal*, *7*(11), 2069–2079. https://doi.org/10.1038/ismej.2013.93

Tamura, K., & Nei, M. (1993). *Estimation of the Number of Nucleotide Substitutions in the Control*
Region of Mitochondrial DNA in Humans and. *10*(3).

Torbati, M., Arzanlou, M., Abed-Ashtiani, F., & Golmohammadi, H. (2019). Occurrence of fruit rot on
cornelian   cherry   caused   by   *Pilidium lythri*   in   Iran.   *Crop Protection*,   *125*,   104884.
https://doi.org/10.1016/j.cropro.2019.104884

Trivedi, P., Leach, J. E., Tringe, S. G., Sa, T., & Singh, B. K. (2020). Plant–microbiome interactions: from
community   assembly   to   plant   health.   *Nature Reviews Microbiology*,   *18*(11),   607–621.
https://doi.org/10.1038/s41579-020-0412-1

Vishwakarma, K., Kumar, N., Shandilya, C., Mohapatra, S., Bhayana, S., & Varma, A. (2020). Revisiting
Plant–Microbe Interactions and Microbial Consortia Application for Enhancing Sustainable
Agriculture: A   Review.   *Frontiers   in   Microbiology*,   *11*(December),   1–21.
https://doi.org/10.3389/fmicb.2020.560406

Ward, J. R., Kim, K., & Harvell, C. D. (2007). Temperature affects coral disease resistance and pathogen
growth. *Marine Ecology Progress Series*, *329*, 115–121. https://doi.org/10.3354/meps329115

White, T. J., Bruns, T., Lee, S., & Taylor, J. (1990). Amplification and direct sequencing of fungal
ribosomal RNA genes for phylogenetics. *PCR Protocols: A Guide to Methods and Applications*,
315–322.

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G., Hayes, A.,
Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K., Ooms, J., Robinson,
D., Seidel, D., Spinu, V., … Yutani, H. (2019). Welcome to the Tidyverse. *Journal of Open Source*
Software, *4*(43), 1686. https://doi.org/10.21105/joss.01686

Zhang, M., Li, H. L., Zhao, A. L., & Zhang, J. X. (2008). First Report of *Hainesia lythri* Causing Leaf Spots
of *Paeonia suffruticosa* in China. *Plant Disease*, *92*(3), 486–486. https://doi.org/10.1094/PDIS-92-
3-0486A

Acknowledgments

This research was funded in whole by National Science Centre, Poland, contract number: 2022/45/N/NZ9/02089. For the purpose of Open Access, the author has applied a CC-BY public copyright license to any Author Accepted Manuscript (AAM) version arising from this submission. The DS's 3-month visit at the Institute of Environmental Sciences, Leiden University, Leiden, Netherlands was financed by the Polish National Agency for Academic Exchange, contract number: BPN/BEK/2023/1/00125/U/00001.

Competing interests

The authors have declared that no competing interests exist.

Data availability

Metabolic profile data and R code used for ststistical analysis were deposited at gihub (https://github.com/dsiegieda/biolog_and_pot_Pilidium) for peer review purposes. After acceptance of the manuscript for publication, this data will be published in RepOD repository (https://repod.icm.edu.pl/), under permanent doi number. ITS sequences of 4 *Pilidium lythri* strains were deposited in GenBank under accession numbers: PQ632564.1, PQ632565.1, PQ632566.1, PQ632567.1. Metataxonomic sequencing data of soil and plant niches were deposited at SRA database under accession numbers: PRJNA1213298 for 16S and PRJNA1213307 for ITS.

CREdiT (https://credit.niso.org/)

Conceptualization: DS, MF, JP, Data curation: DS, JP, Formal analysis: DS, Funding acquisition: DS, Investigation: DS, JP, Methodology: DS, JP, MF, EH, Project administration: DS, Resources: MF, DS, EH, Software: DS, JP, EH, Supervision: DS, Validation: DS, Visualization: DS, Writing – original draft: DS, Writing – review & editing: DS, EH, JP, MF

A) bulk soil

	Importance score
Pilidium	
Wardomyces	
Ilyonectria	
Unclassified Chytridiomycota	
Rhizophydiales_gen_Incertae_sedis	
Unclassified Lecanoromycetes	
Onygenales_gen_Incertae_sedis	
GS16_gen_Incertae_sedis	
Lobulomycetales_gen_Incertae_sedis	
Pyronemataceae_gen_Incertae_sedis	
Entoloma	
Cladochytriales_gen_Incertae_sedis	
Preussia	
Ascomycota_gen_Incertae_sedis	
Pleotrichocladium	

B) rhizosphere

	Importance score
Pilidium	
Pleotrichocladium	
Eocronartium	
Sistotrema	
Trichoderma	
Paraphysoderma	
Solicoccozyma	
Unclassified Mortierellaceae	
Ramicandelaber	
Unclassified Lecanoromycetes	
Niesslia	
Verrucariales_gen_Incertae_sedis	
Ophiosphaerella	
Hypocreales_gen_Incertae_sedis	
Aphelidiomycetes_gen_Incertae_sedis	

C) roots

	Importance score
Peziza	
Pseudeurotium	
Unclassified Fungi	
Unclassified Pleosporales	
Leucoagaricus	
Mucor	
Parascedosporium	
Cladosporium	
Mortierella	
Sporobolomyces	
Kernia	
Fungi_gen_Incertae_sedis	
Ascomycota_gen_Incertae_sedis	
Phialophora	
Metarhizium	