Disruption without diversity loss: how spreading thermotolerant *Pilidium lythri* infection perturbs plant-soil microbiome stability

Dominika Siegieda[1*]

Jacek Panek[1]

S. Emilia Hannula[2]

Magdalena Frąc[1]

[1]Institute of Agrophysics, Polish Academy of Sciences, Doświadczalna 4 street, 20-290 Lublin, Poland

[2]Leiden University, Institute of Environmental Sciences, Department of Environmental Biology, Leiden, The Netherlands

correspondence: Dominika Siegieda (d.siegieda@ipan.lublin.pl)

Keywords

microbiome destabilization, emerging fungal phytopathogen, *Pilidium lythri*, plant holobiont, Anna Karenina Principle, microbial migration, thermotolerance, climate change adaptation

Abstract

While global warming accelerates phytopathogen spread, still little is known on how fungal pathogens disrupt fragile balance in plant-microbe interactions. Here, we demonstrate that the thermotolerant fungus *Pilidium lythri* combines broad chemical resistance with a remarkable ability to destabilize microbial communities in strawberry soil-plant systems. Using high-throughput sequencing (16S rRNA and ITS markers) across 72 samples, combined with mixed-effects modelling,

source-tracking, microbiota assembly mechanisms investigation and network analysis we

demonstrate that *P. lythri* disrupted bacterial community assembly processes, transitioning from

deterministic to stochastic dynamics in the bulk soil, aligning with the Anna Karenina Principle.

Despite stable dominance index (ENS) under infection bacterial and fungal communities, rhizosphere

fungal diversity increased under infection. Network analysis of this niche revealed decreased fungal

connectivity, with enhanced antagonistic interactions between taxa. We also noted that microbial

migration patterns between niches under pathogen inoculation highlighted shifts in community

interactions. Our work uncovers how emerging thermotolerant fungal phytopathogens destabilize

plant holobiont without reducing diversity, important for managing emerging agricultural threats in a

warming climate.

Introduction

Fungal and bacterial microorganisms, together with their host plants, are interconnected through

intricate, multi-level interactions that shape ecosystems and sustain plant health and productivity.

Based on these relationships, the plant holobiont concept was formed - an integrated system where

hosts and microbiota function as one. Plants produce root exudates to modulate microbial assembly

of their niches and vicinity, while the microbiome, in turn, is able to influence and regulate plant

health, improving its resilience against abiotic and biotic stressors (Berg, 2009; Baldrian, 2016; Trivedi

et al., 2020; Gupta *et al.*, 2021; Marco *et al.*, 2022; Michl *et al.*, 2023). These finely tuned interactions

are widely studied and hold great potential for improving agricultural productivity in a sustainable

manner (Trivedi *et al.*, 2020; Vishwakarma *et al.*, 2020; Frąc *et al.*, 2022).

However, this delicate balance between microbiomes and plants is being increasingly threatened by

ongoing global warming. The rise of annual global temperature and changed precipitation patterns

has created the opportunity for a spread and establishment of invasive phytopathogens into new

regions (Bebber *et al.*, 2013; Helfer, 2014; Liu *et al.*, 2019). Although substantial attention has been

given to beneficial plant–microbe interactions, far less is known about how invasive or

thermotolerant fungal pathogens perturb microbial community structure, niche interactions and

assembly rules. This knowledge gap is particularly concerning for perennial crops like strawberries

(*Fragaria x ananassa* Duch.), which depend heavily on stable rhizosphere and soil microbial

communities to thrive and are particularly vulnerable to such disruptions.

Ecological theory offers a conceptual lens for understanding such perturbations. The *Anna Karenina*

Principle (AKP) predicts that under biotic or abiotic stress, host-associated microbiomes become

more variable and stochastic and unpredictible. Increasing evidence suggests that pathogen-induced

disruption can generate divergent microbiome states rather than uniform shifts, but experimental

demonstrations in soil–plant systems remain limited (Arnault *et al.*, 2023).

Pilidium lythri is an emerging fungal phytopathogen with a versatile host range and limited genomic

data. This ascomycete was originally reported in countries located in warmer climate zones, such as

India (Phatak and Payak, 1965), Iran (Arzanlou *et al.*, 2013; Sayami *et al.*, 2013) and China (Zhang *et*

al., 2008; Duan *et al.*, 2010). Notably, its occurrence had been recently identified in several

temperate climate countries, including Belgium (Debode *et al.*, 2011), Croatia (Milicević *et al.*, 2014),

Spain (Aguín *et al.*, 2018), France (Cardin *et al.*, 2009; Caesar *et al.*, 2012) and Poland, although in

2004 only morphologically, but in 2020 we confirmed its presence in symptomatic Polish strawberry

fields with molecular methods (Gołębniak and Jarosz, 2004; Malarczyk *et al.*, 2020), and with new

isolates collected in present study. Primary targets of this fungal phytopathogen are strawberry

plants and fruits (Lopes *et al.*, 2010; Geng *et al.*, 2012; Ayoubi *et al.*, 2016), but it also affects other

agriculturally significant plants, including *Olea europaea* L. (Arzanlou *et al.*, 2013), *Cornus mas*

(Torbati *et al.*, 2019), *Prunus domestica* (Sayami *et al.*, 2013), *Eucalyptus* spp. (Maússe-Sitoe *et al.*,

2016), *Vitis vinifera* (Aguín *et al.*, 2018), and various ornamental flowers (Zhang *et al.*, 2008; Cardin *et*

al., 2009; Caesar *et al.*, 2012; Bruckart *et al.*, 2013; Garfinkel and Chastagner, 2019). Notably

concerning is that *P. lythri* is not host-specific. Strains isolated from olives were able to cause disease

symptoms on strawberry fruits (Karimi *et al.*, 2016), further increasing the severity of the threat

posed by this pathogen as it spreads to new regions. Studies conducted in Brazil, Iran and Belgium

showed incidence as high as 70% in high humidity and warm temperatures, highlighting its threat in

warming climate (Lopes *et al.*, 2010; Ayoubi *et al.*, 2016; Karimi *et al.*, 2016). Pathogenicity test

revealed rapid progression of the disease, with symptoms appearing on up to 100% of fruits within

days after inoculation (Debode *et al.*, 2011; Fernández-Ortuño *et al.*, 2014). The pathogen is able to

infect both wounded and intact grape berries, further underscoring its high virulence (Aguín *et al.*,

2018). Yet, no study has examined how *P. lythri* or other spreading, thermotolerant phytopathogens

influence microbial processes in soil–plant interfaces or whether its effects produce AKP-like

disruptions in microbial assembly.

Here, we hypothesised that infection by a thermotolerant fuganal pathogen destabilizes microbial

community assembly across soil-plany niches, increasing stochasticity and altering microbial

interaction patterns consistent with the Anna Karenina Principle. To test this, we (I) explored

chemical astress tolerance of *P. lythri* across temperatures, (II) quantified pathogen-induced changes

in bacterial and fungal communities in bulk soil, rhizosphere and roots using high-throughput

sequencing, and (III) assessed assembly mechanisms, source-sink dynamics and network stability

under infection. This approach contribute to a broader understanding how spreading pathogens

influence plant-microbiome systems in the face of environmental change.

Methods

Fungal isolates and molecular identification

Pure fungal isolates were obtained by placing symptomatic strawberry leaves and fruits (surface-

sterilized with detergent and 70% ethanol) on PDA medium and incubating them at 24°C in darkness

for 7 days. After initial screening, fungi with macroscopic and microscopic characteristics matching

the description of *P. lythri* (Palm, 1991; Marin-Felix *et al.*, 2017; Karimi *et al.*, 2019) were identified

through amplification and Sanger sequencing of the Internal Transcribed Spacer (ITS) regions of

fungal rDNA, with ITS1 and ITS4 primers (White *et al.*, 1990), as previously described (Frąc *et al.*,

2015). The obtained sequences were deposited in GenBank under accession numbers: PQ632564.1,

PQ632565.1, PQ632566.1, PQ632567.1. These sequences were then used to construct phylogenetic

tree with other sequences available in Genbank database from the *Pilidium* genus. Evolutionary

analyses were conducted in MEGA X (Kumar *et al.*, 2018). Initial tree for the heuristic search was

obtained automatically by applying Neighbor-Join and BioNJ algorithms to a matrix of pairwise

distances estimated using the Tamura-Nei model, and then selecting the topology with superior log

likelihood value. The tree was drawn to scale, with branch lengths measured in the number of

substitutions per site. This analysis involved 57 nucleotide sequences and there were a total of 372

positions in the final dataset. To fulfil Koch's postulates, a spore suspension of 1.6×10^4 spores/mL

was prepared, and surface-sterilized strawberry leaves were briefly submerged in the solution. The

leaves were then sealed in ziplock bags and incubated at 24°C in darkness for 7 days. Sterile

deionized water was used as the untreated control.

Metabolic profile arrays

Each cell of Phenotype Microarray 21-25 BIOLOG plates (Biolog, Hayward, CA, USA) was inoculated

with the fungal mycelium as recommended by the manufacturer and described before (Panek *et al.*,

2016). The plates are coated with 4 different concentrations of the chemical substances and precise

concentrations of the substances used for microarrays are confidential and protected as proprietary

information by the manufacturer, thus, for the purposes of this article, we refer to them as

concentration levels 1, 2, 3, and 4 to maintain clarity. For each of 120 tested chemical substances

(Supplementary Table 1), we incubated 3 *P. lythri* strains across four concentrations at two

temperatures (24 and 30°C), resulting in 72 replicates per substance (3 strains x 3 replicates x 4

concentrations x 2 temperatures). The temperature of 24°C reflects the optimal *in vitro* growth

conditions for *P. lythri*, while 30°C was selected to simulate elevated temperature scenarios

associated with ongoing climate warming. This design allowed us to assess the thermal plasticity of

the fungus' metabolic response under environmental shifts. All plates were incubated for 96 hours in

the darkness. Microplate reader (Biolog, Hayward, CA, USA) was used to determine the absorbance

at 490 nm at the beginning of the experiment and every 24 hours which reflects the reduction of

tetrazolium dye, portraying a metabolic activity of the analysed microorganism. Higher values

indicate robust redox activity and potential tolerance against tested chemical. Obtained data was

analysed statistically and visualised in RStudio, using tidyverse (Wickham *et al.*, 2019) and lme4

(Bates *et al.*, 2015) packages. First, values obtained at 0 h for each cell was divided from read values

at 96 h to establish growth of metabolic activity, and the data was rescaled from 0 to 1. To establish

which factors had an influence on fungal activity, we used generalized linear mixed-effects model

(GLLM) with Gamma family, controlling for the plate. And to establish whether the substance inhibit

fungus' respiration rates, we controlled for plate, strain, temperature and concentration.

Pot experiment and isolation of microbial DNA

Strawberry plants, obtained from the nursery, were grown in pots containing sieved soil from a

certified organic strawberry plantation. After 20 days of initial growth of strawberry plants at 29°C

and 70% humidity, half of the pots were inoculated with *P. lythri* G322/23 spores using 10 mL of a 2 ·

10^4 spores mL^{-1} solution per plant. Sterile, deionized water was used as a negative control. Samples

were collected from two pots in triplicates (6 samples per niche per time point) for healthy and

pathogen-inoculated samples in two timepoints: at the time of inoculation and again after 9 days of

incubation. Bulk soil was transferred into sterile 1.5 mL Eppendorf tubes and frozen at -20°C. The

rhizosphere was collected as described by Lundberg et al. (Lundberg *et al.*, 2012). Roots, flushed free

of rhizosphere soil, were also frozen at -20°C prior to DNA isolation which was conducted with the

QIAQUBE instrument (Qiagen, Hilden, Germany) and GeneMATRIX Soil DNA Purification Kit (EURx,

Gdańsk, Poland) after homogenization of the samples in the FastPrep-24 instrument (MP

Biomedicals) at 4 m s^{-1} for 15 s. DNA isolates were then frozen in -20°C until library construction and

sequencing.

Metataxonomic sequencing, bioinformatics and statistical analysis

For the metataxonomic sequencing, microbial ITS2 (Schoch *et al.*, 2012) and 16Sv3 regions (Klindworth *et al.*, 2013) were amplified, indexed and sequenced on Illumina MiSeq instrument (2x300). Raw data was deposited at SRA database under accession numbers: PRJNA1213298 for 16S and PRJNA1213307 for ITS. Raw data was then passed through the custom QIIME2 (Bolyen *et al.*, 2019) pipelines as reported before (Siegieda *et al.*, 2023, 2024). In brief, after primer trimming with Cutadapt (Martin, 2011) and ITSxpress (Rivers *et al.*, 2018), denoising, chimera removal, merging and Amplicon Sequence Variant (ASV) calling were performed. The taxonomy was assigned based on UNITE v9 dynamic release (Nilsson *et al.*, 2019) and Silva 138 (Pruesse *et al.*, 2007) databases. The results were then analysed with R (v. 4.0.3), using libraries: phyloseq (McMurdie and Holmes, 2013), tidyverse (Wickham *et al.*, 2019), microeco (Liu *et al.*, 2021), microbiomeMarker (Cao *et al.*, 2022), NetCoMi (Peschel *et al.*, 2021). First, for ITS data, we only used ASVs identified to Fungi and Stramenopiles; and for 16S - Bacteria and Archaea. Then, ITS data was repeatedly rarefied at 6,000 and 16S at 11,000 reads using mirlyn package (McMurdie and Holmes, 2014; Cameron *et al.*, 2021). We used supervised machine learning model to evaluate the fungal biomarkers of each niche for healthy and infected samples with Supervised Machine Method (SVM), using Total Sum Scaling normalization method in microbiomeMarker R library (Cao *et al.*, 2022). We employed Effective Number of Species (ENS) (Jost, 2006) and Simpson index on ASV level to evaluate the dominance of taxa and diversity in each niche. We then used Linear Mixed-Effect Models while controlling for the plant to compare differences between control and infected samples. For the beta-diversity analysis, we used PCoA ordination method based on the weihghthed UniFrac and Bray method for bacteria and fungi. We also used adonis2 (Oksanen *et al.*, 2007) and permutest to compare centroid localisation and dispersion between infection groups. To estimate the migration of the bacterial and fungal communities from the niches (sources) to other niches (sinks), we employed SourceTracker (Knights *et al.*, 2011). This Bayesian method models the proportion of a microbial community in a given sample (sink) that originates from predefined source environments based on shared microbial

taxa. We also used RCbray (Bray-Curtis dissimilarity based Raup-Crick metric) and βNTI (β Nearest

Taxon) to evaluate the ecological processes that dominated the bacterial assemblies (Stegen *et al.*,

2013; Liu *et al.*, 2017), using microeco (Liu *et al.*, 2021) R library. βNTI index measures how the

differences in species composition between the two samplings differ from what would be expected

by the null model of random assembly. When βNTI fell between |2|, microbial assembly was

dominated by stochastic mechanisms, and higher than |2|, by deterministic. We then compared the

βNTI indexes between control and pathogen-inoculated samples using Wilcoxon signed-rank test.

The phylogenetic tree is essential to perform this analysis, we did this only for bacterial communities,

as the ITS region is characterised with the variable length for different taxa. Finally, we constructed

microbial networks with 200 most frequent taxa and SPIEC-EASI (Sparse Inverse Covariance

Estimation for Ecological Association Inference) method (Kurtz *et al.*, 2015), using glasso (Graphical

Least Absolute Shrinkage and Selection Operator) approach within NetCoMi R library (Peschel *et al.*,

2021). The dataset was resampled 30 times and 20 penalty levels with 1e-4 as the smallest ratio,

taking the average of symmetric edges. Fast greedy clustering method was used for network

visualization.

Results

Fungal isolates

Fungal ITS sequences, obtained from symptomatic strawberry leaves (Figure 1B) showed 100%

similarity to those of *P. lythri* already present in the GenBank database and phylogenetic tree

revealed clustering of the sequences with other representatives of *P. lythri* and *Pilidium concavum*

(former name of the *P. lythri* (Marin-Felix *et al.*, 2017)) (Figure 1A). The strains collected after 7 days

in 23°C in the darkness on Potato Dextrose Agar were flat and granulose, cinnamon in colour (Karimi

et al., 2016; Marin-Felix *et al.*, 2017) (Figure 1E). The presence of hyaline conidia with smooth walls

and slightly pointed tips were confirmed under the microscope (Figure 1F). After the inoculation of

the surface-sterilized strawberry leaves with the spore suspension and incubation in 24°C in darkness

for 7 days, 100% of the leaves showed brown necrotic lesions. Leaves submerged in water did not

show symptoms of the disease (Figure 1C). Growth of the cinnamon-colour fungal strains were

observed after incubation (24°C for 7 days) of infested leaves on the Petri dishes with the PDA

medium, fulfilling he Koch's postulate (Figure G).

Figure 1. A) Phylogenetic tree of the evolutionary history of *P. lythri* was inferred by using the Maximum Likelihood method

and Tamura-Nei model (Tamura & Nei, 1993) and the e tree with the highest log likelihood (-1144.51) is shown. The

percentage of trees in which the associated taxa clustered together is shown next to the branches. B,C) Symptoms of the *P.*

lythri infection on the strawberry leaves and in the D) control leaf. E) Growth of *P. lythri* after 7 days of growth in 24 °C on

PDA medium, F) conidia of *P. lythri* (40x), and G) growth on PDA medium with infected leaf.

Chemical sensitivity

To evaluate *P. lythri*'s sensitivity to various chemicals, BIOLOG PM 21-25 plates were we incubated

with the spore and mycelium solution of three fungal strains (G321/23, G322/23, G323/23) at 24°C

and 30°C for 96 hours. An evaluation of the GLLM model revealed that, as expected, the temperature

increase was positively correlated with fungal metabolic activity growth, and the substance

concentration had negative correlation with the fungal metabolic activity. We observed different

response of one of strains - G322/23 that was characterised with higher metabolic activity compared

to other 2 (Supplementary Figure 1, Supplementary Table 2). We also evaluated substances that

negatively impacted metabolic activity (Figure 2). Top inhibitors included thiourea, dodecyltrimethyl

ammonium bromide, tetrazolium violet, alexidine and fluorodeoxyuridine, although the effect size

was very small: -0.09, underlining high tolerance of the pathogen to various chemicals.

Figure 2. BIOLOG PM21-25 results of the substances that differently influence growth of *P. lythri*. A) Volcano plot visualizing

biological and statistical effect for different substances. Gray circles represent no significant importance, light blue -

significant results. Dashed vertical lines represent p.value of 0.05 and estimate of -log10(0.05). B) Bar plot representing

estimate for each substance significantly inhibiting pathogens metabolic activity in tested conditions.

Fungal and bacterial community changes after *P. lythri* inoculation of

strawberry

Fungal and bacterial molecular markers for 72 samples from soil (24), rhizosphere (24), and roots (24)

were amplified and sequenced. Rarefaction curves plateaued for all samples, indicating sufficient

sequencing depth. To evaluate the presence of *P. lythri* in infected samples, we employed a

predictive approach to identify 15 most important bacterial (Supplementary Figure 3) and fungal

(Figure 3) taxa that differentiate healthy and infected strawberry compartments. The analysis

revealed that the *Pilidium* fungi belonged to the most important taxa that differentiate between

healthy and infected plants and soil (Figure 3). This, along with the visible symptoms of the disease,

confirmed that the infection was successful and the pathogen migrated from the soil to rhizosphere

and roots.

Figure 3. Top 15 fungal taxa found more abundant in (orange) pathogen-inoculated and in (blue) healthy soils and plants

identified based on Supervised Vector Machine machine learning approach.

After evaluating that infection was successful, we conducted statistical analysis (LMER modelling

while controlling for plant/pot) to compare the alpha diversity of bacterial and fungal communities

between healthy and pathogen-inoculated niches. No significant differences in dominance nor

richness were observed between healthy and pathogen-inoculated niches at both sampling times (p

> 0.05) for bacteria and fungi (Figure 4A, B and Supplementary Table 3). We only noted decreased

diversity of fungal microbiome in pathogen-inoculated rhizosphere samples in fungal

metacommunities at the second sampling time (p < 0.05). We also conducted beta-diversity analysis

of the bacterial (Figure 4 C) and fungal (Figure 4 F) communities. We did not see differences between

centroid localization for bacteria (PERMANOVA: r^2: 0.019, p > 0.05), but the groups were

characterized with different dispersions (permutest: F = 0.0011, p < 0.05), suggesting increased

variability in response of bacterial community to pathogen infection. In fungal metacommunities, the

analysis revealed no significant differences between infected group and the control (PERMANOVA: R^2

= 1.29, p > 0.05, permutest: F = 0.77, p > 0.05) (Supplementary Table 3).

Figure 4. Comparison of alpha and beta diversity of bacterial (A, B, C) and fungal (D, E, F) communities of strawberry niches

immediately after *P. lythri* inoculation (0 days) and after 9 days of (9 days). LMER model was used for comparison of non-

independent samples in alpha diversity. Light blue color represents samples without pathogen inoculation, and orange -

with inoculation.

To elucidate the influence of *P. lythri* on microbial migration between the niches, we used samples

from sampling at 0 days as a source, and samples from sampling at 9 days as sinks. In bulk soil, only

bacterial community revealed significantly higher share of microorganisms in bulk soil and a lower

portion from unknown source in infected samples (Figure 5A). Furthermore, we observed migration

of bacteria from bulk soil to rhizosphere for both bacteria and fungi in infected communities, but also

from rhizosphere to rhizosphere for bacteria in infected communities. Finally, analysis revealed

increased migration from rhizosphere to roots in infected samples in bacterial community.

Figure 5. Source tracking analysis (A) of bacterial and fungal microbiomes between sampling one (source) and two (sink).

Comparison between healthy (light blue, left) and contaminated (light orange, right) samples. Values are shown as mean

standard error from bootstrap. (B) BetaNRI index analysis of bacterial community in second sampling. Significant

differences on both panels between healthy and contaminated samples are shown with asterisk.

We also evaluated the mechanisms involved in bacterial community assembly by calculating the

betaNRI index for bulk soil, rhizosphere, and root samples collected during the second sampling

period (Figure 5B). The results revealed a dominance of stochastic processes in both healthy and

infected rhizosphere and root samples of strawberry (betaNRI values between -2 and 2).

Furthermore, we observed differences in community assembly processes between healthy and

infected bulk soil, with a shift from deterministic processes (characterized by lower turnover) in

healthy bulk soil to a dominance of stochastic processes in infected soil.

Finally, to evaluate the influence of *P. lythri* on the structure of fungal community, we compared the

network structure in the rhizosphere niche between control and infected samples during the second

sampling (Figure 6, Supplementary Figure 3, Supplementary Table 5). Both networks showed similar

characteristics in terms of the number of components (1), clustering coefficient (0.995, 0.999),

modularity (-0.007, -0.008), edge density (0.994, 0.999) and natural connectivity (0.375, 0.310),

average dissimilarity and average path length (0.6, 0.6). On the other hand, the control rhizosphere

exhibited a higher percentage of positive edges (52.308% vs. 43.344%) but lower vertex and edge

connectivity (58, 71).

Figure 6. Fungal network analysis in A) control and B) pathogen-inoculated rhizosphere samples of strawberry. Each node

represents individual fungal ASV, and node size is proportional to number of reads. Node colours represent the same

clusters. Green lines represent positive interactions and red- negative. The position of the same nodes is the same in both

networks.

Notably, changes were observed in centrality of networks (Jaccard Index). Degree, betweenness and

hub taxa measures showed no overlap in most connected, keystone and key hub taxa, significant

shift was observed in closeness of most central nodes (Table 1).

Table 1. Centrality changes between networks of rhizosphere samples in control and *P.* contaminated samples. Asterisk

shows statistically significant change.

	Degree	Betweenness	Closeness	Eigenvector	Hub taxa
Jaccard Index	0	0	0.071	0.209	0
p-value (P≤Jacc)	1	0.296	0.027*	0.055	1

293　Importantly, *Pilidium* was identified as a connector taxa in the control rhizosphere (betweenness =

294　1), loosing this status in pathogen-inoculated (betweenness = 0) samples (Supplementary Table 5).

Discussion

Here, we investigate the underexplored role of emerging thermotolerant fungal pathogen in shaping

plant microbiomes using newly isolated strains of *Pilidium lythri*. We assessed its thermal tolerance

by screening metabolic response to 120 different chemical substances and explored its impact on the

structure, assembly processes and association networks of bacterial and fungal metacommunities

across strawberry soil and plant niches (bulk soil, rhizosphere, roots).

In vitro evaluation of the impact of various chemical substances on the growth and development of

the tested phytopathogen at two temperatures (24 and 30 °C) using BIOLOG microarrays raised

serious concerns, as *P. lythri* demonstrated overall higher metabolic activity at higher temperatures.

This aligns with the growing evidence of global warming contributing to the spread of fungal

phytopathogens to new areas (Ward *et al.*, 2007; Bebber *et al.*, 2013; Chaloner *et al.*, 2021). Among

the 120 chemical substances tested, top five presented inhibition mechanism on pathogens

metabolic activity, likely by disruption of electron chain, destabilization of cell membranes, and

inhibition of DNA/RNA synthesis (Waldorf and Polak, 1983; Mamouei *et al.*, 2018; Ansari *et al.*, 2024;

Yang *et al.*, 2025). Unfortunately, although significant, the effect of the inhibition was modest. These

results underline high resilience of the pathogen that points to its high environmental tolerance and

potential to spread.

While the dissemination of fungal pathogens is widely reported (Bebber *et al.*, 2013), still little is

known about the effect of such invasive microorganisms on the microbial structure of the plant

holobiont. A number of studies have reported significant differences in the α-diversity of microbiome

structure between healthy and infected soil and plant niches (Lefcheck *et al.*, 2015; Delgado-

Baquerizo *et al.*, 2016; Bakker *et al.*, 2018). In the present study, we compared alpha and beta

diversity of bacterial and fungal metacommunities in healthy and *P. lythri*-infected bulk soil,

rhizosphere, and roots at two time points: directly after pathogen inoculation and 9 days post-

inoculation. For the bacteria, alpha diversity did not differ between inoculated and control samples,

but we observed increased community dispersion, that points to possible destabilization of bacterial

assemblages under biotic stress. To test this, we analyzed the mechanisms involved in bacterial

microbiome assembly. The results indicated that composition of bacterial communities in healthy

and diseased rhizosphere and roots samples was shaped by stochastic processes (i.e. random

colonization, dispersal, and ecological drift). Interestingly, we observed a shift from deterministic

processes in healthy bulk soil, to stochastic processes in *P. lythri*-infected samples, indicating a

disruption in microbial turnover. Infected soil showed random processes that could have been

triggered by disturbances in microbial community caused by the infection, described by the Anna

Karenina Principle (AKP) (Arnault *et al.*, 2023). Analysis of microbial migration suggests increased

migration of bacteria from the soil in pathogen-inoculated samples, as well as increased migration of

bacteria from the rhizosphere to the roots in infected plant samples. This inconsistency between

migration (suggesting plant filtering) and microbial assembly (suggesting random processes) could be

caused by the smaller scale of deterministic processes being masked by larger-scale microbial

community assembly. Furthermore, these two mechanisms represent different time scales, where

βNRI focuses on broader, longer-term assembly processes (Arnault *et al.*, 2023). Importantly,

migration of microorganisms can be caused by random dispersal (stochastic processes) but may

appear deterministic as migration increases. Thus, these two processes are not necessarily

contradictory.

In the fungal microbiome, we observed increased migration of fungi from bulk soil to the rhizosphere

in the inoculated samples compared to control, highlighting the importance of this niche in fungal

microbiome dynamics and the role of plant in microbial recruitment under pathogen stress (Bais *et*

al., 2006; Berendsen *et al.*, 2012). The alpha diversity results revealed an increased diversity of the

fungal community under pathogen influence in the strawberry rhizosphere, and we then compared

the microbial networks in this niche to further explore the influence of the pathogen. A number of

methods for construction and analysis of co-occurrence networks had been developed, and SPIEC-

EASI is characterized with important advantages, such as accounting for compositional type of data

and mitigation of non-independence of samples. Also, an implementation of glasso approach allowed

for the analysis in high-dimensional setting with more variables than samples (Kurtz *et al.*, 2015).

Thus, we implemented these methods for the analysis in current study. Comparison of healthy and

infected samples revealed shift in microbial fungal networks under *P. lythri* influence. The fungal

networks in infected rhizospheres exhibited lower percentages of positive edges compared to

control, and lower robustness of networks, suggesting restructuring of fungal community in response

to pathogen presence. These findings underline decrease in natural connectivity percentage of

positive edges, that suggest lowered positive interactions between fungal taxa. Infection with the

pathogen significantly disrupted fungal networks, by decreasing networks connectivity, which

indicates a shift toward increased competition and antagonism within the fungal microbiome. This

suggests adaptation of the fungal meta-community, making it more resistant to further infections

and environmental stressors (Hernandez *et al.*, 2021).

Concluding, in this study, we isolated and molecularly identified isolates of spreading fungal

phytopathogen from symptomatic strawberry plants - *P. lythri*. We observed pathogen's high

tolerance to various chemical substances in both, lower and higher temperature, highlighting it hight

thermal plasticity and resilience. We also noted significant variability in tolerance caused by chemical

substances between strains. We used *P. lythri* to evaluate ecological impacts of spreading fungal

phytopathogen on microorganism metacommunities in bulk soil, rhizosphere and roots of strawberry

with high-throughput marker sequencing. We revealed that *P. lythri* infection had a major effect on

both the structure and dynamics of the plant associated microbiome. Its ability to shift bacterial

microbial assembly mechanism from deterministic to stochastic in the bulk soil, underscores its

capacity to destabilize plant-associated microbiomes. Additionally, we observed that *P. lythri* shifted

the fungal microbiome dynamics in the strawberry rhizosphere towards reducing mutualistic

connections, while maintaining overall network stability. This underlined adaptive reorganization of

fungal community, potentially increasing resilience to secondary infections and other (a)biotic

stresses.

These findings underline the critical role of emerging, thermotolerant fungal pathogens like *Pilidium*

lythri in destabilizing soil microbiomes and plant-microbiome interactions. This is significantly

important in the context of global warming, which accelerates spread and impact of such pathogens.

Author Contribution

376 Conceptualization: DS, MF, JP, Data curation: DS, JP, Formal analysis: DS, Funding acquisition: DS,

377 Investigation: DS, JP, Methodology: DS, JP, MF, EH, Project administration: DS, Resources: MF, DS,

378 EH, Software: DS, JP, EH, Supervision: DS, Validation: DS, Visualization: DS, Writing – original draft:

379 DS, Writing – review & editing: DS, EH, JP, MF

Acknowledgments

This research was funded in whole by National Science Centre, Poland, contract number:

2022/45/N/NZ9/02089. For the purpose of Open Access, the author has applied a CC-BY public

copyright license to any Author Accepted Manuscript (AAM) version arising from this submission. The

DS's 3-month visit at the Institute of Environmental Sciences, Leiden University, Leiden, Netherlands

was financed by the Polish National Agency for Academic Exchange, contract number:

BPN/BEK/2023/1/00125/U/00001.

Conflict of Interest

388 The authors have declared that no competing interests exist.

Data Availability Statement

Metabolic profile data and R code used for ststistical analysis were deposited at gihub

(https://github.com/dsiegieda/biolog_and_pot_Pilidium) for peer review purposes. After acceptance

of the manuscript for publication, this data will be published in RepOD repository

(https://repod.icm.edu.pl/), under permanent doi number. ITS sequences of 4 *Pilidium lythri* strains

were deposited in GenBank under accession numbers: PQ632564.1, PQ632565.1, PQ632566.1,

PQ632567.1. Metataxonomic sequencing data of soil and plant niches were deposited at SRA

database under accession numbers: PRJNA1213298 for 16S and PRJNA1213307 for ITS. Benefits

generated: benefits from this research accrue from the sharing of our data and results on public

databases, as described above.

Declaration of generative AI and AI-assisted technologies in the writing process

401 During the preparation of original draft of this manuscript the author used ChatGPT in order to

402 improve language and readability. After using this tool, the author reviewed and edited the content

403 as needed and take full responsibility for the content of the publication.

References

Aguín, O., Ferreiroa, V., González-Jartín, J.M., Alfonso, A., Botana, L.M., Mansilla, J.P., and Sainz, M.J.
(2018) *Pilidium lythri* is associated with bunch rot of grapevine (Vitis vinifera). *Am J Enol Vitic*
69: 410–416.

Ansari, A., Kumar, D., Gupta, P., Poluri, K.M., Rai, N., Ameen, F., and Kumar, N. (2024) Mechanistic
insights into antifungal potential of Alexidine dihydrochloride and hexachlorophene in Candida
albicans: a drug repurposing approach. *Arch Microbiol* **206**: 383.

Arnault, G., Mony, C., and Vandenkoornhuyse, P. (2023) Plant microbiota dysbiosis and the Anna
Karenina Principle. *Trends Plant Sci* **28**: 18–30.

Arzanlou, M., Torbati, M., and Jafary, H. (2013) Fruit rot on olive caused by *Pilidium concavum* in Iran.
Australas Plant Dis Notes **8**: 117–121.

Ayoubi, N., Soleimani, M.J., and Zare, R. (2016) Pilidium concavum, causing tan-brown rot on
strawberry in Iran. *Journal of Plant Pathology* **98**: 667–669.

Bais, H.P., Weir, T.L., Perry, L.G., Gilroy, S., and Vivanco, J.M. (2006) The role of root exudates in
rhizosphere interactions with plants and other organisms. *Annu Rev Plant Biol* **57**: 233–266.

Bakker, P.A.H.M., Pieterse, C.M.J., de Jonge, R., and Berendsen, R.L. (2018) The Soil-Borne Legacy.
Cell **172**: 1178–1180.

Baldrian, P. (2016) Forest microbiome: diversity, complexity and dynamics. *FEMS Microbiol Rev*
fuw040.

Bates, D., Mächler, M., Bolker, B., and Walker, S. (2015) Fitting Linear Mixed-Effects Models Using
lme4. *J Stat Softw* **67**:.

Bebber, D.P., Ramotowski, M.A.T., and Gurr, S.J. (2013) Crop pests and pathogens move polewards in
a warming world. *Nat Clim Chang* **3**: 985–988.

Berendsen, R.L., Pieterse, C.M.J., and Bakker, P.A.H.M. (2012) The rhizosphere microbiome and plant
health. *Trends Plant Sci* **17**: 478–486.

Berg, G. (2009) Plant–microbe interactions promoting plant growth and health: perspectives for
controlled use of microorganisms in agriculture. *Appl Microbiol Biotechnol* **84**: 11–18.

Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith, G.A., et al. (2019)
Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat*
Biotechnol **37**: 852–857.

Bruckart, W.L., Eskandari, F.M., Coombs, E.M., Rossman, A.Y., and Palm, M.E. (2013) First report of
Pilidium concavum causing leaf necrosis on Fallopia japonica in the United States. *Plant Dis* **97**:
146–146.

Caesar, A.J., Lartey, R.T., and Caesar-TonThat, T. (2012) First report of *Pilidium concavum* causing
root lesions of meadow hawkweed in France. *Plant Dis* **96**: 1830–1830.

Cameron, E.S., Schmidt, P.J., Tremblay, B.J.M., Emelko, M.B., and Müller, K.M. (2021) Enhancing
diversity analysis by repeatedly rarefying next generation sequencing data describing microbial
communities. *Sci Rep* **11**: 1–13.

Cao, Y., Dong, Q., Wang, D., Zhang, P., Liu, Y., and Niu, C. (2022) microbiomeMarker: an
R/Bioconductor package for microbiome marker identification and visualization. *Bioinformatics*
38: 4027–4029.

Cardin, L., Vincenot, L., and Balesdent, M.H. (2009) First report of Pilidium concavum on Bergenia
crassifolia in France. *Plant Dis* **93**: 548–548.

Chaloner, T.M., Gurr, S.J., and Bebber, D.P. (2021) Plant pathogen infection risk tracks global crop
yields under climate change. *Nat Clim Chang* **11**: 710–715.

Debode, J., Van Hemelrijck, W., Heungens, K., Maes, M., and Creemers, P. (2011) First Report of
Pilidium concavum Causing Tan-Brown Rot on Strawberry Fruit in Belgium. *Plant Dis* **95**: 1029–
1029.

Delgado-Baquerizo, M., Maestre, F.T., Reich, P.B., Jeffries, T.C., Gaitan, J.J., Encinar, D., et al. (2016)
Microbial diversity drives multifunctionality in terrestrial ecosystems. *Nat Commun* **7**: 1–8.

Duan, Y.B., Kang, Y.B., and Yu, Z.Z. (2010) First report of *Pilidium concavum* on *Paeonia suffruticosa* in
China. *Plant Dis* **94**: 271–271.

Fernández-Ortuño, D., Bryson, P.K., and Schnabel, G. (2014) First Report of Pilidium concavum
Causing Tan-brown Rot on Strawberry Nursery Stock in South Carolina. *Plant Dis* **98**: 1010–
1010.

Frąc, M., Hannula, E.S., Bełka, M., Salles, J.F., and Jedryczka, M. (2022) Soil mycobiome in sustainable
agriculture. *Front Microbiol* **13**: 1–9.

Frąc, M., Jezierska-Tys, S., and Yaguchi, T. (2015) Occurrence, detection, and molecular and
metabolic characterization of heat-resistant fungi in soils and plants and their risk to human
health. *Advances in Agronomy* **132**: 161–204.

Garfinkel, A.R. and Chastagner, G.A. (2019) Survey reveals a broad range of fungal pathogens and an
oomycete on peonies in the United States. *Plant Health Prog* **20**: 250–254.

Geng, W.L., Hu, P., Ma, Z., Zhao, X.Y., and Wei, Y.M. (2012) First Report of Pilidium concavum Causing
Tan-Brown Leaf Spot on Strawberry in China. *Plant Dis* **96**: 1377–1377.

Gołębniak, B. and Jarosz, A. (2004) First report of tan-brown rot (*Hainesia lythri*) on strawberry fruits
in Poland. *Phytopathogia Polonica* **31**: 57–60.

Gupta, R., Anand, G., Gaur, R., and Yadav, D. (2021) Plant–microbiome interactions for sustainable
agriculture: a review. *Physiology and Molecular Biology of Plants* **27**: 165–179.

Helfer, S. (2014) Rust fungi and global change. *New Phytologist* **201**: 770–780.

Hernandez, D.J., David, A.S., Menges, E.S., Searcy, C.A., and Afkhami, M.E. (2021) Environmental
stress destabilizes microbial networks. *ISME Journal* **15**: 1722–1734.

Jost, L. (2006) Entropy and diversity. *Oikos* **113**: 363–375.

Karimi, K., Arzanlou, M., Babai-ahari, A., and Pertot, I. (2016) Biological and molecular
characterisation of *Pilidium lythri*, an emerging strawberry pathogen in Iran. *Phytopathol*
Mediterr **55**: 366–379.

Karimi, K., Arzanlou, M., and Pertot, I. (2019) Development of novel species-specific primers for the
specific identification of *Colletotrichum nymphaeae* based on conventional PCR and LAMP
techniques. *Eur J Plant Pathol.*

Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., and Glöckner, F.O. (2013)
Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation
sequencing-based diversity studies. *Nucleic Acids Res* **41**: e1–e1.

Knights, D., Kuczynski, J., Charlson, E.S., Zaneveld, J., Mozer, M.C., Collman, R.G., et al. (2011)
Bayesian community-wide culture-independent microbial source tracking. *Nat Methods* **8**: 761–
763.

Kumar, S., Stecher, G., Li, M., Knyaz, C., and Tamura, K. (2018) MEGA X: Molecular Evolutionary
Genetics Analysis across computing platforms. *Mol Biol Evol* **35**: 1547–1549.

Kurtz, Z.D., Müller, C.L., Miraldi, E.R., Littman, D.R., Blaser, M.J., and Bonneau, R.A. (2015) Sparse and
Compositionally Robust Inference of Microbial Ecological Networks. *PLoS Comput Biol* **11**:.

Lefcheck, J.S., Byrnes, J.E.K., Isbell, F., Gamfeldt, L., Griffin, J.N., Eisenhauer, N., et al. (2015)
Biodiversity enhances ecosystem multifunctionality across trophic levels and habitats. *Nat*
Commun **6**:.

Liu, C., Cui, Y., Li, X., and Yao, M. (2021) microeco: an R package for data mining in microbial
community ecology. *FEMS Microbiol Ecol* **97**:.

Liu, C., Yao, M., Stegen, J.C., Rui, J., Li, J., and Li, X. (2017) Long-term nitrogen addition affects the
phylogenetic turnover of soil microbial community responding to moisture pulse. *Sci Rep* **7**: 1–
11.

Liu, X., Ma, Z., Cadotte, M.W., Chen, F., He, J.-S., and Zhou, S. (2019) Warming affects foliar fungal
diseases more than precipitation in a Tibetan alpine meadow. *New Phytologist* **221**: 1574–1584.

Lopes, U.P., Zambolim, L., Lopes, U.N., Pereira, O.L., and Costa, H. (2010) First report of *Pilidium*
concavum causing tan-brown rot in strawberry fruits in Brazil. *Plant Pathol* **59**: 1171.

Lundberg, D.S., Lebeis, S.L., Paredes, S.H., Yourstone, S., Gehring, J., Malfatti, S., et al. (2012) Defining
the core *Arabidopsis thaliana* root microbiome. *Nature* **488**: 86–90.

Malarczyk, D.G., Panek, J., and Frąc, M. (2020) Triplex Real-Time PCR Approach for the Detection of
Crucial Fungal Berry Pathogens—*Botrytis* spp., *Colletotrichum* spp. and *Verticillium* spp. *Int J*
Mol Sci **21**: 8469.

Mamouei, Z., Alqarihi, A., Singh, S., Xu, S., Mansour, M.K., Ibrahim, A.S., and Uppuluri, P. (2018)
Alexidine Dihydrochloride Has Broad-Spectrum Activities against Diverse Fungal Pathogens.
mSphere **3**:.

Marco, S., Loredana, M., Riccardo, V., Raffaella, B., Walter, C., and Luca, N. (2022) Microbe-assisted
crop improvement: a sustainable weapon to restore holobiont functionality and resilience.
Hortic Res **9**:.

Marin-Felix, Y., Groenewald, J.Z., Cai, L., Chen, Q., Marincowitz, S., Barnes, I., et al. (2017) Genera of
phytopathogenic fungi: GOPHY 1. *Stud Mycol* **86**: 99–216.

Martin, M. (2011) Cutadapt removes adapter sequences from hight-throughput sequencing reads.
EMBnet J **17**: 10–12.

Maússe-Sitoe, S.N., Chen, S., Wingfield, M.J., and Roux, J. (2016) Diseases of eucalypts in the central
and northern provinces of Mozambique. *Southern Forests: a Journal of Forest Science* **78**: 169–
183.

McMurdie, P.J. and Holmes, S. (2013) Phyloseq: An R Package for Reproducible Interactive Analysis
and Graphics of Microbiome Census Data. *PLoS One* **8**:.

McMurdie, P.J. and Holmes, S. (2014) Waste Not, Want Not: Why Rarefying Microbiome Data Is
Inadmissible. *PLoS Comput Biol* **10**: e1003531.

Michl, K., Berg, G., and Cernava, T. (2023) The microbiome of cereal plants: The current state of
knowledge and the potential for future applications. *Environ Microbiome* **18**: 28.

Milicević, T., Kaliterna, J., Ivić, D., and Milović, M. (2014) Records of phytopathogenic fungal species
on native plants new to Croatia. *Natura Croatica* **23**: 179–187.

Nilsson, R.H., Larsson, K.H., Taylor, A.F.S., Bengtsson-Palme, J., Jeppesen, T.S., Schigel, D., et al.
(2019) The UNITE database for molecular identification of fungi: Handling dark taxa and parallel
taxonomic classifications. *Nucleic Acids Res* **47**: D259–D264.

Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Stevens, H., and Oksanen, M.J. (2007) The vegan
package. *Community ecology package* **10**: 719.

Palm, M.E. (1991) Taxonomy and Morphology of the Synanamorphs *Pilidium concavum* and *Hainesia*
lythri (Coelomycetes). *Mycologia* **83**: 787.

Panek, J., Frąc, M., and Bilińska-Wielgus, N. (2016) Comparison of chemical sensitivity of fresh and
long-stored heat resistant *Neosartorya fischeri* environmental isolates using Biolog Phenotype
MicroArray system. *PLoS One* **11**: 4–8.

Peschel, S., Müller, C.L., Von Mutius, E., Boulesteix, A.L., and Depner, M. (2021) NetCoMi: Network
construction and comparison for microbiome data in R. *Brief Bioinform* **22**: 1–18.

Phatak, H.C. and Payak, M.M. (1965) *Hainesia lythri* on strawberry - a new record for India. *Indian*
Phytopathol **18**: 237–239.

Pruesse, E., Quast, C., Knittel, K., Fuchs, B.M., Ludwig, W., Peplies, J., and Glöckner, F.O. (2007) SILVA:
A comprehensive online resource for quality checked and aligned ribosomal RNA sequence data
compatible with ARB. *Nucleic Acids Res* **35**: 7188–7196.

Rivers, A.R., Weber, K.C., Gardner, T.G., Liu, S., and Armstrong, S.D. (2018) ITSxpress: Software to
rapidly trim internally transcribed spacer sequences with quality scores for marker gene
analysis. *F1000Res* **7**: 1418.

Sayami, M., Babaeizad, V., and Fathi, J. (2013) First report of *Pilidium concavum* as the causative
agent of tan-brown rot of *Prunus domestica* fruits. *Journal of Plant Pathology* **95**: 451.

Schoch, C.L., Seifert, K.A., Huhndorf, S., Robert, V., Spouge, J.L., Levesque, C.A., et al. (2012) Nuclear
ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi.
Proc Natl Acad Sci U S A **109**: 6241–6246.

Siegieda, D., Panek, J., and Frąc, M. (2024) Ecological processes of bacterial microbiome assembly in
healthy and dysbiotic strawberry farms. *BMC Plant Biol* **24**: 1–14.

Siegieda, D., Panek, J., and Frąc, M. (2023) Plant and soil health in organic strawberry farms – Greater
importance of fungal trophic modes and networks than α -diversity of the mycobiome. *Applied*
Soil Ecology **188**: 1–15.

Stegen, J.C., Lin, X., Fredrickson, J.K., Chen, X., Kennedy, D.W., Murray, C.J., et al. (2013) Quantifying
community assembly processes and identifying features that impose them. *ISME Journal* **7**:
2069–2079.

Torbati, M., Arzanlou, M., Abed-Ashtiani, F., and Golmohammadi, H. (2019) Occurrence of fruit rot on
cornelian cherry caused by *Pilidium lythri* in Iran. *Crop Protection* **125**: 104884.

Trivedi, P., Leach, J.E., Tringe, S.G., Sa, T., and Singh, B.K. (2020) Plant–microbiome interactions: from
community assembly to plant health. *Nat Rev Microbiol* **18**: 607–621.

Vishwakarma, K., Kumar, N., Shandilya, C., Mohapatra, S., Bhayana, S., and Varma, A. (2020)
Revisiting Plant–Microbe Interactions and Microbial Consortia Application for Enhancing
Sustainable Agriculture: A Review. *Front Microbiol* **11**: 1–21.

Waldorf, A.R. and Polak, A. (1983) Mechanisms of action of 5-fluorocytosine. *Antimicrob Agents*
Chemother **23**: 79–85.

Ward, J.R., Kim, K., and Harvell, C.D. (2007) Temperature affects coral disease resistance and
pathogen growth. *Mar Ecol Prog Ser* **329**: 115–121.

White, T.J., Bruns, T., Lee, S., and Taylor, J. (1990) Amplification and direct sequencing of fungal
ribosomal RNA genes for phylogenetics. *PCR Protocols: A Guide to Methods and Applications*
315–322.

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., et al. (2019) Welcome to
the Tidyverse. *J Open Source Softw* **4**: 1686.

Yang, Z.H., Li, G.T., and Chen, Y. (2025) Synthesis and Antifungal Activity of Aldehydes-Thiourea
Derivatives as Promising Antifungal Agents Against Postharvest Gray Mold Disease. *Chem*
Biodivers.

Zhang, M., Li, H.L., Zhao, A.L., and Zhang, J.X. (2008) First Report of *Hainesia lythri* Causing Leaf Spots
of *Paeonia suffruticosa* in China. *Plant Dis* **92**: 486–486.

A) bulk soil

	Importance score
Pilidium	
Wardomyces	
Ilyonectria	
Unclassified Chytridiomycota	
Rhizophydiales_gen_Incertae_sedis	
Unclassified Lecanoromycetes	
Onygenales_gen_Incertae_sedis	
GS16_gen_Incertae_sedis	
Lobulomycetales_gen_Incertae_sedis	
Pyronemataceae_gen_Incertae_sedis	
Entoloma	
Cladochytriales_gen_Incertae_sedis	
Preussia	
Ascomycota_gen_Incertae_sedis	
Pleotrichocladium	

B) rhizosphere

	Importance score
Pilidium	
Pleotrichocladium	
Eocronartium	
Sistotrema	
Trichoderma	
Paraphysoderma	
Solicoccozyma	
Unclassified Mortierellaceae	
Ramicandelaber	
Unclassified Lecanoromycetes	
Niesslia	
Verrucariales_gen_Incertae_sedis	
Ophiosphaerella	
Hypocreales_gen_Incertae_sedis	
Aphelidiomycetes_gen_Incertae_sedis	

C) roots

	Importance score
Peziza	
Pseudeurotium	
Unclassified Fungi	
Unclassified Pleosporales	
Leucoagaricus	
Mucor	
Parascedosporium	
Cladosporium	
Mortierella	
Sporobolomyces	
Kernia	
Fungi_gen_Incertae_sedis	
Ascomycota_gen_Incertae_sedis	
Phialophora	
Metarhizium	