1 Disruption without diversity loss: how spreading thermotolerant

2 *Pilidium lythri* infection perturbs plant-soil microbiome stability

Dominika Siegieda[1*]

Jacek Panek[1]

S. Emilia Hannula[2]

Magdalena Frąc[1]

[1]Institute of Agrophysics, Polish Academy of Sciences, Doświadczalna 4 street, 20-290 Lublin, Poland

[2]Leiden University, Institute of Environmental Sciences, Department of Environmental Biology, Leiden,

The Netherlands

correspondence: Dominika Siegieda (d.siegieda@ipan.lublin.pl)

12 Keywords

13 microbiome destabilization, emerging fungal phytopathogen, *Pilidium lythri*, plant holobiont, Anna

14 Karenina Principle, microbial migration, thermotolerance, climate change adaptation

15 Abstract

While global warming accelerates phytopathogen spread, still little is known on how fungal pathogens

disrupt fragile balance in plant-microbe interactions. Here, we demonstrate that the thermotolerant

fungus *Pilidium lythri* combines broad chemical resistance with a remarkable ability to destabilize

microbial communities in strawberry soil-plant systems. Using high-throughput sequencing (16S rRNA

and ITS markers) across 72 samples, combined with mixed-effects modelling, source-tracking,

microbiota assembly mechanisms investigation and network analysis we demonstrate that *P. lythri*

disrupted bacterial community assembly processes, transitioning from deterministic to stochastic

dynamics in the bulk soil, aligning with the Anna Karenina Principle. Despite stable dominance index

(ENS) under infection bacterial and fungal communities, rhizosphere fungal diversity increased under

infection. Network analysis of this niche revealed decreased fungal connectivity, with enhanced

antagonistic interactions between taxa. We also noted that microbial migration patterns between

niches under pathogen inoculation highlighted shifts in community interactions. Our work uncovers

how emerging thermotolerant fungal phytopathogens destabilize plant holobiont without reducing

diversity, important for managing emerging agricultural threats in a warming climate.

Introduction

Fungal and bacterial microorganisms, together with their host plants, are interconnected through intricate, multi-level interactions that shape ecosystems and sustain plant health and productivity. Based on these relationships, the plant holobiont concept was formed - an integrated system where hosts and microbiota function as one. Plants produce root exudates to modulate microbial assembly of their niches and vicinity, while the microbiome, in turn, is able to influence and regulate plant health, improving its resilience against abiotic and biotic stressors [1–6]. These finely tuned interactions are widely studied and hold great potential for improving agricultural productivity in a sustainable manner [6–8].

However, this delicate balance between microbiomes and plants is being increasingly threatened by ongoing global warming. The rise of annual global temperature and changed precipitation patterns has created the opportunity for a spread and establishment of invasive phytopathogens into new regions [9–11]. Despite this urgent challenges, limited attention has been given to understand how these emerging pathogens alter and interact with microbial communities. This issue is particularly concerning for perennial crops like strawberries (*Fragaria x ananassa* Duch.), which depend heavily on stable rhizosphere and soil microbial communities to thrive and are particularly vulnerable to such disruptions.

Pilidium lythri is an emerging fungal phytopathogen with a versatile host range and limited genomic data. This ascomycete was originally reported in countries located in warmer climate zones, such as India [12], Iran [13, 14] and China [15, 16]. Notably, its occurrence had been recently identified in several temperate climate countries, including Belgium [17], Croatia [18], Spain [19], France [20, 21] and Poland, although in 2004 only morphologically, but in 2020 we confirmed its presence in symptomatic Polish strawberry fields with molecular methods [22, 23], and with new isolates collected in present study. Primary targets of this fungal phytopathogen are strawberry plants and fruits [24–26], but it also affects other agriculturally significant plants, including *Olea europaea* L. [13], *Cornus mas* [27], *Prunus domestica* [14], *Eucalyptus* spp. [28], *Vitis vinifera* [19], and various ornamental flowers [16, 20, 21, 29, 30]. Notably concerning is that *P. lythri* is not host-specific. Strains isolated from olives were able to cause disease symptoms on strawberry fruits [31], further increasing the severity of the threat posed by this pathogen as it spreads to new regions. Studies conducted in Brazil, Iran and Belgium showed incidence as high as 70% in high humidity and warm temperatures, highlighting its threat in warming climate [24, 26, 31]. Pathogenicity test revealed rapid progression of the disease, with symptoms appearing on up to 100% of fruits within days after inoculation [17, 32]. The pathogen is able to infect both wounded and intact grape berries, further underscoring its high virulence [19].

Given the limited knowledge about its ecological behaviour, *P. lythri* provides a valuable lens to explore

how understudied, spreading fungal pathogens shape microbial community assembly and niche-

specific interactions within the plant holobiont.

Building on this rationale, we isolated *P. lythri* from symptomatic strawberry plants in a Polish nursery

to explore how this pathogen responds to temperature pressures and host-associated microbiomes.

We characterised its metabolic phenotype at 24°C and 30°C with BIOLOG PM21-25 plates, which

allowed us to assess strains' tolerance to various chemical compounds under varying thermal

conditions, mimicking environmental shifts. In parallel, we examined the impact of *P. lythri* infection

on the bacterial/archaeal and fungal microbiomes in distinct niches: bulk soil, rhizosphere and

strawberry roots. This study significantly advances our understanding of ecological behaviour of an

emerging, thermotolerant fungal phytopathogen, presenting how it disrupts microbial dynamics in the

plant holobiont. These findings contribute to a broader understanding how spreading pathogens

influence plant-microbiome systems in the face of environmental change.

Methods

Fungal isolates and molecular identification

Pure fungal isolates were obtained by placing symptomatic strawberry leaves and fruits (surface-

sterilized with detergent and 70% ethanol) on PDA medium and incubating them at 24°C in darkness

for 7 days. After initial screening, fungi with macroscopic and microscopic characteristics matching the

description of *P. lythri* [33–35] were identified through amplification and Sanger sequencing of the

Internal Transcribed Spacer (ITS) regions of fungal rDNA, with ITS1 and ITS4 primers [36], as previously

described [37]. The obtained sequences were deposited in GenBank under accession numbers:

PQ632564.1, PQ632565.1, PQ632566.1, PQ632567.1. These sequences were then used to construct

phylogenetic tree with other sequences available in Genbank database from the *Pilidium* genus.

Evolutionary analyses were conducted in MEGA X [38]. Initial tree for the heuristic search was obtained

automatically by applying Neighbor-Join and BioNJ algorithms to a matrix of pairwise distances

estimated using the Tamura-Nei model, and then selecting the topology with superior log likelihood

value. The tree was drawn to scale, with branch lengths measured in the number of substitutions per

site. This analysis involved 57 nucleotide sequences and there were a total of 372 positions in the final

dataset. To fulfil Koch's postulates, a spore suspension of 1.6×10^4 spores/mL was prepared, and

surface-sterilized strawberry leaves were briefly submerged in the solution. The leaves were then

sealed in ziplock bags and incubated at 24°C in darkness for 7 days. Sterile deionized water was used

as the untreated control.

Metabolic profile arrays

Each cell of Phenotype Microarray 21-25 BIOLOG plates (Biolog, Hayward, CA, USA) was inoculated with the fungal mycelium as recommended by the manufacturer and described before [39]. The plates are coated with 4 different concentrations of the chemical substances and precise concentrations of the substances used for microarrays are confidential and protected as proprietary information by the manufacturer, thus, for the purposes of this article, we refer to them as concentration levels 1, 2, 3, and 4 to maintain clarity. For each of 120 tested chemical substances (Supplementary Table 1), we incubated 3 *P. lythri* strains across four concentrations at two temperatures (24 and 30°C), resulting in 72 replicates per substance (3 strains x 3 replicates x 4 concentrations x 2 temperatures). The temperature of 24°C reflects the optimal *in vitro* growth conditions for *P. lythri*, while 30°C was selected to simulate elevated temperature scenarios associated with ongoing climate warming. This design allowed us to assess the thermal plasticity of the fungus' metabolic response under environmental shifts. All plates were incubated for 96 hours in the darkness. Microplate reader (Biolog, Hayward, CA, USA) was used to determine the absorbance at 490 nm at the beginning of the experiment and every 24 hours which reflects the reduction of tetrazolium dye, portraying a metabolic activity of the analysed microorganism. Higher values indicate robust redox activity and potential tolerance against tested chemical. Obtained data was analysed statistically and visualised in RStudio, using tidyverse [40] and lme4 [41] packages. First, values obtained at 0 h for each cell was divided from read values at 96 h to establish growth of metabolic activity, and the data was rescaled from 0 to 1. To establish which factors had an influence on fungal activity, we used generalized linear mixed-effects model (GLLM) with Gamma family, controlling for the plate. And to establish whether the substance inhibit fungus' respiration rates, we controlled for plate, strain, temperature and concentration.

Pot experiment and isolation of microbial DNA

Strawberry plants, obtained from the nursery, were grown in pots containing sieved soil from a certified organic strawberry plantation. After 20 days of initial growth of strawberry plants at 29°C and 70% humidity, half of the pots were inoculated with *P. lythri* G322/23 spores using 10 mL of a $2 \cdot 10^4$ spores mL^{-1} solution per plant. Sterile, deionized water was used as a negative control. Samples were collected from two pots in triplicates (6 samples per niche per time point) for healthy and pathogen-inoculated samples in two timepoints: at the time of inoculation and again after 9 days of incubation. Bulk soil was transferred into sterile 1.5 mL Eppendorf tubes and frozen at -20°C. The rhizosphere was collected as described by Lundberg et al. [42]. Roots, flushed free of rhizosphere soil, were also frozen at -20°C prior to DNA isolation which was conducted with the QIAQUBE instrument (Qiagen, Hilden, Germany) and GeneMATRIX Soil DNA Purification Kit (EURx, Gdańsk, Poland) after homogenization of

128 the samples in the FastPrep-24 instrument (MP Biomedicals) at 4 m s^{-1} for 15 s. DNA isolates were then

129 frozen in -20°C until library construction and sequencing.

Metataxonomic sequencing, bioinformatics and statistical analysis

For the metataxonomic sequencing, microbial ITS2 [43] and 16Sv3 regions [44] were amplified, indexed

and sequenced on Illumina MiSeq instrument (2x300). Raw data was deposited at SRA database under

accession numbers: PRJNA1213298 for 16S and PRJNA1213307 for ITS. Raw data was then passed

through the custom QIIME2 [45] pipelines as reported before [46, 47]. In brief, after primer trimming

with Cutadapt [48] and ITSxpress [49], denoising, chimera removal, merging and Amplicon Sequence

Variant (ASV) calling were performed. The taxonomy was assigned based on UNITE v9 dynamic release

[50] and Silva 138 [51] databases. The results were then analysed with R (v. 4.0.3), using libraries:

phyloseq [52], tidyverse [40], microeco [53], microbiomeMarker [54], NetCoMi [55]. First, for ITS data,

we only used ASVs identified to Fungi and Stramenopiles; and for 16S - Bacteria and Archaea. Then,

ITS data was repeatedly rarefied at 6,000 and 16S at 11,000 reads using mirlyn package [56, 57]. We

used supervised machine learning model to evaluate the fungal biomarkers of each niche for healthy

and infected samples with Supervised Machine Method (SVM), using Total Sum Scaling normalization

method in microbiomeMarker R library [54]. We employed Effective Number of Species (ENS) [58] and

Simpson index on ASV level to evaluate the dominance of taxa and diversity in each niche. We then

used Linear Mixed-Effect Models while controlling for the plant to compare differences between

control and infected samples. For the beta-diversity analysis, we used PCoA ordination method based

on the weihghthed UniFrac and Bray method for bacteria and fungi. We also used adonis2 [59] and

permutest to compare centroid localisation and dispersion between infection groups. To estimate the

migration of the bacterial and fungal communities from the niches (sources) to other niches (sinks),

we employed SourceTracker [60]. This Bayesian method models the proportion of a microbial

community in a given sample (sink) that originates from predefined source environments based on

shared microbial taxa. We also used RCbray (Bray-Curtis dissimilarity based Raup-Crick metric) and

βNTI (β Nearest Taxon) to evaluate the ecological processes that dominated the bacterial assemblies

[61, 62], using microeco [53] R library. βNTI index measures how the differences in species composition

between the two samplings differ from what would be expected by the null model of random

assembly. When βNTI fell between |2|, microbial assembly was dominated by stochastic mechanisms,

and higher than |2|, by deterministic. We then compared the βNTI indexes between control and

pathogen-inoculated samples using Wilcoxon signed-rank test. The phylogenetic tree is essential to

perform this analysis, we did this only for bacterial communities, as the ITS region is characterised with

the variable length for different taxa. Finally, we constructed microbial networks with 200 most

frequent taxa and SPIEC-EASI (Sparse Inverse Covariance Estimation for Ecological Association

Inference) method [63], using glasso (Graphical Least Absolute Shrinkage and Selection Operator)

approach within NetCoMi R library [55]. The dataset was resampled 30 times and 20 penalty levels

with 1e-4 as the smallest ratio, taking the average of symmetric edges. Fast greedy clustering method

was used for network visualization.

Results

Fungal isolates

Fungal ITS sequences, obtained from symptomatic strawberry leaves (Figure 1B) showed 100%

similarity to those of *P. lythri* already present in the GenBank database and phylogenetic tree revealed

clustering of the sequences with other representatives of *P. lythri* and *Pilidium concavum* (former

name of the *P. lythri* [35]) (Figure 1A). The strains collected after 7 days in 23°C in the darkness on

Potato Dextrose Agar were flat and granulose, cinnamon in colour [31, 35] (Figure 1E). The presence

of hyaline conidia with smooth walls and slightly pointed tips were confirmed under the microscope

(Figure 1F). After the inoculation of the surface-sterilized strawberry leaves with the spore suspension

and incubation in 24°C in darkness for 7 days, 100% of the leaves showed brown necrotic lesions.

Leaves submerged in water did not show symptoms of the disease (Figure 1C). Growth of the

cinnamon-colour fungal strains were observed after incubation (24°C for 7 days) of infested leaves on

the Petri dishes with the PDA medium, fulfilling he Koch's postulate (Figure G).

Figure 1. A) Phylogenetic tree of the evolutionary history of *P. lythri* was inferred by using the Maximum Likelihood method

and Tamura-Nei model (Tamura & Nei, 1993) and the e tree with the highest log likelihood (-1144.51) is shown. The

percentage of trees in which the associated taxa clustered together is shown next to the branches. B,C) Symptoms of the *P.*

lythri infection on the strawberry leaves and in the D) control leaf. E) Growth of *P. lythri* after 7 days of growth in 24 °C on

PDA medium, F) conidia of *P. lythri* (40x), and G) growth on PDA medium with infected leaf.

Chemical sensitivity

To evaluate *P. lythri*'s sensitivity to various chemicals, BIOLOG PM 21-25 plates were we incubated with the spore and mycelium solution of three fungal strains (G321/23, G322/23, G323/23) at 24°C and 30°C for 96 hours. An evaluation of the GLLM model revealed that, as expected, the temperature increase was positively correlated with fungal metabolic activity growth, and the substance concentration had negative correlation with the fungal metabolic activity. We observed different response of one of strains - G322/23 that was characterised with higher metabolic activity compared to other 2 (Supplementary Figure 1, Supplementary Table 2). We also evaluated substances that negatively impacted metabolic activity (Figure 2). Top inhibitors included thiourea, dodecyltrimethyl ammonium bromide, tetrazolium violet, alexidine and fluorodeoxyuridine, although the effect size was very small: -0.09, underlining high tolerance of the pathogen to various chemicals.

Figure 2. BIOLOG PM21-25 results of the substances that differently influence growth of *P. lythri.* A) Volcano plot visualizing biological and statistical effect for different substances. Gray circles represent no significant importance, light blue - significant results. Dashed vertical lines represent p.value of 0.05 and estimate of -log10(0.05). B) Bar plot representing estimate for each substance significantly inhibiting pathogens metabolic activity in tested conditions.

Fungal and bacterial community changes after *P. lythri* inoculation of strawberry

Fungal and bacterial molecular markers for 72 samples from soil (24), rhizosphere (24), and roots (24) were amplified and sequenced. Rarefaction curves plateaued for all samples, indicating sufficient sequencing depth. To evaluate the presence of *P. lythri* in infected samples, we employed a predictive approach to identify 15 most important bacterial (Supplementary Figure 3) and fungal (Figure 3) taxa that differentiate healthy and infected strawberry compartments. The analysis revealed that the

Pilidium fungi belonged to the most important taxa that differentiate between healthy and infected plants and soil (Figure 3). This, along with the visible symptoms of the disease, confirmed that the infection was successful and the pathogen migrated from the soil to rhizosphere and roots.

Figure 3. Top 15 fungal taxa found more abundant in (orange) pathogen-inoculated and in (blue) healthy soils and plants identified based on Supervised Vector Machine machine learning approach.

After evaluating that infection was successful, we conducted statistical analysis (LMER modelling while controlling for plant/pot) to compare the alpha diversity of bacterial and fungal communities between healthy and pathogen-inoculated niches. No significant differences in dominance nor richness were observed between healthy and pathogen-inoculated niches at both sampling times ($p > 0.05$) for bacteria and fungi (Figure 4A, B and Supplementary Table 3). We only noted decreased diversity of fungal microbiome in pathogen-inoculated rhizosphere samples in fungal metacommunities at the second sampling time ($p < 0.05$). We also conducted beta-diversity analysis of the bacterial (Figure 4 C) and fungal (Figure 4 F) communities. We did not see differences between centroid localization for bacteria (PERMANOVA: r^2: 0.019, $p > 0.05$), but the groups were characterized with different dispersions (permutest: $F = 0.0011$, $p < 0.05$), suggesting increased variability in response of bacterial community to pathogen infection. In fungal metacommunities, the analysis revealed no significant differences between infected group and the control (PERMANOVA: $R^2 = 1.29$, $p > 0.05$, permutest: $F = 0.77$, $p > 0.05$) (Supplementary Table 3).

Figure 4. Comparison of alpha and beta diversity of bacterial (A, B, C) and fungal (D, E, F) communities of strawberry niches immediately after *P. lythri* inoculation (0 days) and after 9 days of (9 days). LMER model was used for comparison of non-independent samples in alpha diversity. Light blue color represents samples without pathogen inoculation, and orange - with inoculation.

To elucidate the influence of *P. lythri* on microbial migration between the niches, we used samples from sampling at 0 days as a source, and samples from sampling at 9 days as sinks. In bulk soil, only bacterial community revealed significantly higher share of microorganisms in bulk soil and a lower portion from unknown source in infected samples (Figure 5A). Furthermore, we observed migration of bacteria from bulk soil to rhizosphere for both bacteria and fungi in infected communities, but also from rhizosphere to rhizosphere for bacteria in infected communities. Finally, analysis revealed increased migration from rhizosphere to roots in infected samples in bacterial community.

Figure 5. Source tracking analysis (A) of bacterial and fungal microbiomes between sampling one (source) and two (sink). Comparison between healthy (light blue, left) and contaminated (light orange, right) samples. Values are shown as mean standard error from bootstrap. (B) BetaNRI index analysis of bacterial community in second sampling. Significant differences on both panels between healthy and contaminated samples are shown with asterisk.

We also evaluated the mechanisms involved in bacterial community assembly by calculating the

betaNRI index for bulk soil, rhizosphere, and root samples collected during the second sampling period

(Figure 5B). The results revealed a dominance of stochastic processes in both healthy and infected

rhizosphere and root samples of strawberry (betaNRI values between -2 and 2). Furthermore, we

observed differences in community assembly processes between healthy and infected bulk soil, with

a shift from deterministic processes (characterized by lower turnover) in healthy bulk soil to a

dominance of stochastic processes in infected soil.

Finally, to evaluate the influence of *P. lythri* on the structure of fungal community, we compared the

network structure in the rhizosphere niche between control and infected samples during the second

sampling (Figure 6, Supplementary Figure 3, Supplementary Table 5). Both networks showed similar

characteristics in terms of the number of components (1), clustering coefficient (0.995, 0.999),

modularity (-0.007, -0.008), edge density (0.994, 0.999) and natural connectivity (0.375, 0.310),

average dissimilarity and average path length (0.6, 0.6). On the other hand, the control rhizosphere

exhibited a higher percentage of positive edges (52.308% vs. 43.344%) but lower vertex and edge

connectivity (58, 71).

Figure 6. Fungal network analysis in A) control and B) pathogen-inoculated rhizosphere samples of strawberry. Each node

represents individual fungal ASV, and node size is proportional to number of reads. Node colours represent the same clusters.

Green lines represent positive interactions and red- negative. The position of the same nodes is the same in both networks.

Notably, changes were observed in centrality of networks (Jaccard Index). Degree, betweenness and

hub taxa measures showed no overlap in most connected, keystone and key hub taxa, significant shift

was observed in closeness of most central nodes (Table 1).

Table 1. Centrality changes between networks of rhizosphere samples in control and *P.* contaminated samples. Asterisk shows

statistically significant change.

	Degree	Betweenness	Closeness	Eigenvector	Hub taxa
Jaccard Index	0	0	0.071	0.209	0
p-value (P≤Jacc)	1	0.296	0.027*	0.055	1

268 Importantly, *Pilidium* was identified as a connector taxa in the control rhizosphere (betweenness = 1),

269 loosing this status in pathogen-inoculated (betweenness = 0) samples (Supplementary Table 5).

Discussion

Here, we investigate the underexplored role of emerging thermotolerant fungal pathogens in shaping

plant microbiomes using newly isolated strains of *Pilidium lythri*. We assessed their thermal tolerance

by screening metabolic response to 120 different chemical substances and explored its impact on the

structure, assembly processes and association networks of bacterial and fungal metacommunities

across strawberry soil and plant niches (bulk soil, rhizosphere, roots).

In vitro evaluation of the impact of various chemical substances on the growth and development of

the tested phytopathogen at two temperatures (24 and 30 °C) using BIOLOG microarrays raised serious

concerns, as *P. lythri* demonstrated overall higher metabolic activity at higher temperatures. This aligns

with the growing evidence of global warming contributing to the spread of fungal phytopathogens to

new areas [9, 64, 65]. Among the 120 chemical substances tested, top five presented inhibition

mechanism on pathogens metabolic activity, likely by disruption of electron chain, destabilization of

cell membranes, and inhibition of DNA/RNA synthesis [66–69]. Unfortunately, although significant, the

effect of the inhibition was modest. These results underline high resilience of the pathogen that points

to its high environmental tolerance and potential to spread.

While the dissemination of fungal pathogens is widely reported [9], still little is known about the effect

of such invasive microorganisms on the microbial structure of the plant holobiont. A number of studies

have reported significant differences in the α-diversity of microbiome structure between healthy and

infected soil and plant niches [70–72]. In the present study, we compared alpha and beta diversity of

bacterial and fungal metacommunities in healthy and *P. lythri*-infected bulk soil, rhizosphere, and roots

at two time points: directly after pathogen inoculation and 9 days post-inoculation. For the bacteria,

alpha diversity did not differ between inoculated and control samples, but we observed increased

community dispersion, that points to possible destabilization of bacterial assemblages under biotic

stress. To test this, we analyzed the mechanisms involved in bacterial microbiome assembly. The

results indicated that composition of bacterial communities in healthy and diseased rhizosphere and

roots samples was shaped by stochastic processes (i.e. random colonization, dispersal, and ecological

drift). Interestingly, we observed a shift from deterministic processes in healthy bulk soil, to stochastic

processes in *P. lythri*-infected samples, indicating a disruption in microbial turnover. Infected soil

showed random processes that could have been triggered by disturbances in microbial community

caused by the infection, described by the Anna Karenina Principle (AKP) [73]. Analysis of microbial

migration suggests increased migration of bacteria from the soil in pathogen-inoculated samples, as

well as increased migration of bacteria from the rhizosphere to the roots in infected plant samples.

This inconsistency between migration (suggesting plant filtering) and microbial assembly (suggesting

random processes) could be caused by the smaller scale of deterministic processes being masked by

larger-scale microbial community assembly. Furthermore, these two mechanisms represent different

time scales, where βNRI focuses on broader, longer-term assembly processes [73]. Importantly,

migration of microorganisms can be caused by random dispersal (stochastic processes) but may appear

deterministic as migration increases. Thus, these two processes are not necessarily contradictory.

In the fungal microbiome, we observed increased migration of fungi from bulk soil to the rhizosphere

in the inoculated samples compared to control, highlighting the importance of this niche in fungal

microbiome dynamics and the role of plant in microbial recruitment under pathogen stress [74, 75].

The alpha diversity results revealed an increased diversity of the fungal community under pathogen

influence in the strawberry rhizosphere, and we then compared the microbial networks in this niche

to further explore the influence of the pathogen. A number of methods for construction and analysis

of co-occurrence networks had been developed, and SPIEC-EASI is characterized with important

advantages, such as accounting for compositional type of data and mitigation of non-independence of

samples. Also, an implementation of glasso approach allowed for the analysis in high-dimensional

setting with more variables than samples [63]. Thus, we implemented these methods for the analysis

in current study. Comparison of healthy and infected samples revealed shift in microbial fungal

networks under *P. lythri* influence. The fungal networks in infected rhizospheres exhibited lower

percentages of positive edges compared to control, and lower robustness of networks, suggesting

restructuring of fungal community in response to pathogen presence. These findings underline

decrease in natural connectivity percentage of positive edges, that suggest lowered positive

interactions between fungal taxa. Infection with the pathogen significantly disrupted fungal networks,

by decreasing networks connectivity, which indicates a shift toward increased competition and

antagonism within the fungal microbiome. This suggests adaptation of the fungal meta-community,

making it more resistant to further infections and environmental stressors [76].

Concluding, in this study, we isolated and molecularly identified isolates of spreading fungal

phytopathogen from symptomatic strawberry plants - *P. lythri*. We observed pathogen's high tolerance

to various chemical substances in both, lower and higher temperature, highlighting it hight thermal

plasticity and resilience. We also noted significant variability in tolerance caused by chemical

substances between strains. We used *P. lythri* to evaluate ecological impacts of spreading fungal

phytopathogen on microorganism metacommunities in bulk soil, rhizosphere and roots of strawberry

with high-throughput marker sequencing. We revealed that *P. lythri* infection had a major effect on both the structure and dynamics of the plant associated microbiome. Its ability to shift bacterial microbial assembly mechanism from deterministic to stochastic in the bulk soil, underscores its capacity to destabilize plant-associated microbiomes. Additionally, we observed that *P. lythri* shifted the fungal microbiome dynamics in the strawberry rhizosphere towards reducing mutualistic connections, while maintaining overall network stability. This underlined adaptive reorganization of fungal community, potentially increasing resilience to secondary infections and other (a)biotic stresses.

These findings underline the critical role of emerging, thermotolerant fungal pathogens like *Pilidium lythri* in destabilizing soil microbiomes and plant-microbiome interactions. This is significantly important in the context of global warming, which accelerates spread and impact of such pathogens.

Author Contribution

Conceptualization: DS, MF, JP, Data curation: DS, JP, Formal analysis: DS, Funding acquisition: DS, Investigation: DS, JP, Methodology: DS, JP, MF, EH, Project administration: DS, Resources: MF, DS, EH, Software: DS, JP, EH, Supervision: DS, Validation: DS, Visualization: DS, Writing – original draft: DS, Writing – review & editing: DS, EH, JP, MF

Acknowledgments

This research was funded in whole by National Science Centre, Poland, contract number: 2022/45/N/NZ9/02089. For the purpose of Open Access, the author has applied a CC-BY public copyright license to any Author Accepted Manuscript (AAM) version arising from this submission. The DS's 3-month visit at the Institute of Environmental Sciences, Leiden University, Leiden, Netherlands was financed by the Polish National Agency for Academic Exchange, contract number: BPN/BEK/2023/1/00125/U/00001.

Conflict of Interest

The authors have declared that no competing interests exist.

Data Availability Statement

Metabolic profile data and R code used for ststistical analysis were deposited at gihub (https://github.com/dsiegieda/biolog_and_pot_Pilidium) for peer review purposes. After acceptance of the manuscript for publication, this data will be published in RepOD repository (https://repod.icm.edu.pl/), under permanent doi number. ITS sequences of 4 *Pilidium lythri* strains were deposited in GenBank under accession numbers: PQ632564.1, PQ632565.1, PQ632566.1, PQ632567.1. Metataxonomic sequencing data of soil and plant niches were deposited at SRA database

References

Aguín, O., Ferreiroa, V., González-Jartín, J. M., Alfonso, A., Botana, L. M., Mansilla, J. P., & Sainz, M. J.

(2018). *Pilidium lythri* is associated with bunch rot of grapevine (Vitis vinifera). *American Journal*

of Enology and Viticulture, *69*(4), 410–416. https://doi.org/10.5344/ajev.2018.17093

Arnault, G., Mony, C., & Vandenkoornhuyse, P. (2023). Plant microbiota dysbiosis and the Anna

Karenina   Principle.   *Trends   in   Plant   Science*,   *28*(1),   18–30.

https://doi.org/10.1016/j.tplants.2022.08.012

Arzanlou, M., Torbati, M., & Jafary, H. (2013). Fruit rot on olive caused by *Pilidium concavum* in Iran.

Australasian Plant Disease Notes, *8*(1), 117–121. https://doi.org/10.1007/s13314-013-0111-0

Ayoubi, N., Soleimani, M. J., & Zare, R. (2016). Pilidium concavum, causing tan-brown rot on strawberry

in Iran. *Journal of Plant Pathology*, *98*(3), 667–669. https://doi.org/10.4454/JPP.V98I3.063

Bais, H. P., Weir, T. L., Perry, L. G., Gilroy, S., & Vivanco, J. M. (2006). The role of root exudates in

rhizosphere interactions with plants and other organisms. *Annual Review of Plant Biology*, *57*(1),

233–266. https://doi.org/10.1146/annurev.arplant.57.032905.105159

Bakker, P. A. H. M., Pieterse, C. M. J., de Jonge, R., & Berendsen, R. L. (2018). The Soil-Borne Legacy.

Cell, *172*(6), 1178–1180. https://doi.org/10.1016/j.cell.2018.02.024

Baldrian, P. (2016). Forest microbiome: diversity, complexity and dynamics. *FEMS Microbiology*

Reviews, fuw040. https://doi.org/10.1093/femsre/fuw040

Bebber, D. P., Ramotowski, M. A. T., & Gurr, S. J. (2013). Crop pests and pathogens move polewards in

a warming world. *Nature Climate Change*, *3*(11), 985–988. https://doi.org/10.1038/nclimate1990

Berendsen, R. L., Pieterse, C. M. J., & Bakker, P. A. H. M. (2012). The rhizosphere microbiome and plant

health. *Trends in Plant Science*, *17*(8), 478–486. https://doi.org/10.1016/j.tplants.2012.04.001

Berg, G. (2009). Plant–microbe interactions promoting plant growth and health: perspectives for

controlled use of microorganisms in agriculture. *Applied Microbiology and Biotechnology*, *84*(1),

11–18. https://doi.org/10.1007/s00253-009-2092-7

Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., Alexander, H.,

Alm, E. J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J. E., Bittinger, K., Brejnrod, A., Brislawn, C.

J., Brown, C. T., Callahan, B. J., Caraballo-Rodríguez, A. M., Chase, J., … Caporaso, J. G. (2019).
Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nature*
Biotechnology, *37*(8), 852–857. https://doi.org/10.1038/s41587-019-0209-9

Bruckart, W. L., Eskandari, F. M., Coombs, E. M., Rossman, A. Y., & Palm, M. E. (2013). First report of
Pilidium concavum causing leaf necrosis on Fallopia japonica in the United States. *Plant Disease*,
97(1), 146–146. https://doi.org/10.1094/PDIS-07-12-0688-PDN

Caesar, A. J., Lartey, R. T., & Caesar-TonThat, T. (2012). First report of *Pilidium concavum* causing root
lesions of meadow hawkweed in France. *Plant Disease*, *96*(12), 1830–1830.
https://doi.org/10.1094/PDIS-05-12-0509-PDN

Cameron, E. S., Schmidt, P. J., Tremblay, B. J. M., Emelko, M. B., & Müller, K. M. (2021). Enhancing
diversity analysis by repeatedly rarefying next generation sequencing data describing microbial
communities. *Scientific Reports*, *11*(1), 1–13. https://doi.org/10.1038/s41598-021-01636-1

Cao, Y., Dong, Q., Wang, D., Zhang, P., Liu, Y., & Niu, C. (2022). microbiomeMarker: an R/Bioconductor
package for microbiome marker identification and visualization. *Bioinformatics*, *38*(16), 4027–
4029. https://doi.org/10.1093/bioinformatics/btac438

Cardin, L., Vincenot, L., & Balesdent, M. H. (2009). First report of Pilidium concavum on Bergenia
crassifolia in France. *Plant Disease*, *93*(5), 548–548. https://doi.org/10.1094/PDIS-93-5-0548B

Chaloner, T. M., Gurr, S. J., & Bebber, D. P. (2021). Plant pathogen infection risk tracks global crop
yields under climate change. *Nature Climate Change*, *11*(8), 710–715.
https://doi.org/10.1038/s41558-021-01104-8

Cheng, C., Gao, X., Feng, B., Sheen, J., Shan, L., & He, P. (2013). Plant immune response to pathogens
differs with changing temperatures. *Nature Communications*, *4*(1), 2530.
https://doi.org/10.1038/ncomms3530

Cheng, Y. T., Zhang, L., & He, S. Y. (2019). Plant-Microbe Interactions Facing Environmental Challenge.
Cell Host & Microbe, *26*(2), 183–192. https://doi.org/10.1016/j.chom.2019.07.009

Debode, J., Van Hemelrijck, W., Heungens, K., Maes, M., & Creemers, P. (2011). First Report of Pilidium
concavum Causing Tan-Brown Rot on Strawberry Fruit in Belgium. *Plant Disease*, *95*(8), 1029–
1029. https://doi.org/10.1094/PDIS-10-10-0752

Delgado-Baquerizo, M., Maestre, F. T., Reich, P. B., Jeffries, T. C., Gaitan, J. J., Encinar, D., Berdugo, M.,
Campbell, C. D., & Singh, B. K. (2016). Microbial diversity drives multifunctionality in terrestrial
ecosystems. *Nature Communications*, *7*, 1–8. https://doi.org/10.1038/ncomms10541

Duan, Y. B., Kang, Y. B., & Yu, Z. Z. (2010). First report of *Pilidium concavum* on *Paeonia suffruticosa* in
China. *Plant Disease*, *94*(2), 271–271. https://doi.org/10.1094/PDIS-94-2-0271B

Fernández-Ortuño, D., Bryson, P. K., & Schnabel, G. (2014). First Report of Pilidium concavum Causing
Tan-brown Rot on Strawberry Nursery Stock in South Carolina. *Plant Disease*, *98*(7), 1010–1010.
https://doi.org/10.1094/PDIS-01-14-0048-PDN

Frąc, M., Hannula, E. S., Bełka, M., Salles, J. F., & Jedryczka, M. (2022). Soil mycobiome in sustainable
agriculture.   *Frontiers   in   Microbiology*,   *13*(November),   1–9.
https://doi.org/10.3389/fmicb.2022.1033824

Frąc, M., Jezierska-Tys, S., & Yaguchi, T. (2015). Occurrence, detection, and molecular and metabolic
characterization of heat-resistant fungi in soils and plants and their risk to human health.
Advances in Agronomy, *132*, 161–204.

Gao, M., Xiong, C., Gao, C., Tsui, C. K. M., Wang, M. M., Zhou, X., Zhang, A. M., & Cai, L. (2021). Disease-
induced changes in plant microbiome assembly and functional adaptation. *Microbiome*, *9*(1), 1–
18. https://doi.org/10.1186/s40168-021-01138-2

Garfinkel, A. R., & Chastagner, G. A. (2019). Survey reveals a broad range of fungal pathogens and an
oomycete   on   peonies   in   the   United   States.   *Plant Health Progress*, *20*(4), 250–254.
https://doi.org/10.1094/PHP-09-19-0065-S

Geng, W. L., Hu, P., Ma, Z., Zhao, X. Y., & Wei, Y. M. (2012). First Report of Pilidium concavum Causing
Tan-Brown   Leaf   Spot   on   Strawberry   in   China.   *Plant   Disease*,   *96*(9),   1377–1377.
https://doi.org/10.1094/PDIS-03-12-0310-PDN

Gołębniak, B., & Jarosz, A. (2004). First report of tan-brown rot (*Hainesia lythri*) on strawberry fruits in
Poland. *Phytopathogia Polonica*, *31*, 57–60.

Gupta, R., Anand, G., Gaur, R., & Yadav, D. (2021). Plant–microbiome interactions for sustainable
agriculture: a review. *Physiology and Molecular Biology of Plants*, *27*(1), 165–179.
https://doi.org/10.1007/s12298-021-00927-1

Helfer, S. (2014). Rust fungi and global change. *New Phytologist*, *201*(3), 770–780.
https://doi.org/10.1111/nph.12570

Hernandez, D. J., David, A. S., Menges, E. S., Searcy, C. A., & Afkhami, M. E. (2021). Environmental stress
destabilizes   microbial   networks.   *ISME   Journal*,   *15*(6),   1722–1734.
https://doi.org/10.1038/s41396-020-00882-x

Jost, L. (2006). Entropy and diversity. *Oikos*, *113*(2), 363–375. https://doi.org/10.1111/j.2006.0030-
1299.14714.x

Karimi, K., Arzanlou, M., Babai-ahari, A., & Pertot, I. (2016). Biological and molecular characterisation
of *Pilidium lythri*, an emerging strawberry pathogen in Iran. *Phytopathologia Mediterranea*, *55*(3),
366–379. https://doi.org/10.14601/Phytopathol_Mediterr-18391

Karimi, K., Arzanlou, M., & Pertot, I. (2019). Development of novel species-specific primers for the
specific identification of *Colletotrichum nymphaeae* based on conventional PCR and LAMP
techniques. *European Journal of Plant Pathology*. https://doi.org/10.1007/s10658-019-01895-9

Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., & Glöckner, F. O. (2013).
Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation
sequencing-based diversity studies. *Nucleic Acids Research*, *41*(1), e1–e1.
https://doi.org/10.1093/nar/gks808

Knights, D., Kuczynski, J., Charlson, E. S., Zaneveld, J., Mozer, M. C., Collman, R. G., Bushman, F. D.,
Knight, R., & Kelley, S. T. (2011). Bayesian community-wide culture-independent microbial source
tracking. *Nature Methods*, *8*(9), 761–765. https://doi.org/10.1038/nmeth.1650

Lefcheck, J. S., Byrnes, J. E. K., Isbell, F., Gamfeldt, L., Griffin, J. N., Eisenhauer, N., Hensel, M. J. S.,
Hector, A., Cardinale, B. J., & Duffy, J. E. (2015). Biodiversity enhances ecosystem
multifunctionality across trophic levels and habitats. *Nature Communications*, *6*.
https://doi.org/10.1038/ncomms7936

Liu, C., Cui, Y., Li, X., & Yao, M. (2021). microeco: an R package for data mining in microbial community
ecology. *FEMS Microbiology Ecology*, *97*(2). https://doi.org/10.1093/femsec/fiaa255

Liu, C., Yao, M., Stegen, J. C., Rui, J., Li, J., & Li, X. (2017). Long-term nitrogen addition affects the
phylogenetic turnover of soil microbial community responding to moisture pulse. *Scientific
Reports*, *7*(1), 1–11. https://doi.org/10.1038/s41598-017-17736-w

Liu, X., Ma, Z., Cadotte, M. W., Chen, F., He, J.-S., & Zhou, S. (2019). Warming affects foliar fungal
diseases more than precipitation in a Tibetan alpine meadow. *New Phytologist*, *221*(3), 1574–
1584. https://doi.org/10.1111/nph.15460

Lopes, U. P., Zambolim, L., Lopes, U. N., Pereira, O. L., & Costa, H. (2010). First report of *Pilidium
concavum* causing tan-brown rot in strawberry fruits in Brazil. *Plant Pathology*, *59*, 1171.
https://doi.org/10.1111/j.1365-3059.2010.02336.x

Lundberg, D. S., Lebeis, S. L., Paredes, S. H., Yourstone, S., Gehring, J., Malfatti, S., Tremblay, J.,
Engelbrektson, A., Kunin, V., Rio, T. G. Del, Edgar, R. C., Eickhorst, T., Ley, R. E., Hugenholtz, P.,
Tringe, S. G., & Dangl, J. L. (2012). Defining the core *Arabidopsis thaliana* root microbiome.
Nature, *488*(7409), 86–90. https://doi.org/10.1038/nature11237

Malarczyk, D. G., Panek, J., & Frąc, M. (2020). Triplex Real-Time PCR Approach for the Detection of
Crucial Fungal Berry Pathogens—*Botrytis* spp., *Colletotrichum* spp. and *Verticillium* spp.
International Journal of Molecular Sciences, *21*(22), 8469. https://doi.org/10.3390/ijms21228469

Marco, S., Loredana, M., Riccardo, V., Raffaella, B., Walter, C., & Luca, N. (2022). Microbe-assisted crop
improvement: a sustainable weapon to restore holobiont functionality and resilience.
Horticulture Research, *9*(October). https://doi.org/10.1093/hr/uhac160

Marin-Felix, Y., Groenewald, J. Z., Cai, L., Chen, Q., Marincowitz, S., Barnes, I., Bensch, K., Braun, U.,
Camporesi, E., Damm, U., de Beer, Z. W., Dissanayake, A., Edwards, J., Giraldo, A., Hernández-
Restrepo, M., Hyde, K. D., Jayawardena, R. S., Lombard, L., Luangsa-ard, J., … Crous, P. W. (2017).
Genera of phytopathogenic fungi: GOPHY 1. *Studies in Mycology*, *86*, 99–216.
https://doi.org/10.1016/j.simyco.2017.04.002

Martin, M. (2011). Cutadapt removes adapter sequences from hight-throughput sequencing reads.
EMBnet.Journal, *17*(1), 10–12. https://doi.org/https://doi.org/10.14806/ej.17.1.200

Maússe-Sitoe, S. N., Chen, S., Wingfield, M. J., & Roux, J. (2016). Diseases of eucalypts in the central
and northern provinces of Mozambique. *Southern Forests: A Journal of Forest Science*, *78*(3),
169–183. https://doi.org/10.2989/20702620.2015.1126780

McMurdie, P. J., & Holmes, S. (2013). Phyloseq: An R Package for Reproducible Interactive Analysis and
Graphics of Microbiome Census Data. *PLoS ONE*, *8*(4).
https://doi.org/10.1371/journal.pone.0061217

McMurdie, P. J., & Holmes, S. (2014). Waste Not, Want Not: Why Rarefying Microbiome Data Is
Inadmissible. *PLoS Computational Biology*, *10*(4), e1003531.
https://doi.org/10.1371/journal.pcbi.1003531

Michl, K., Berg, G., & Cernava, T. (2023). The microbiome of cereal plants: The current state of
knowledge and the potential for future applications. *Environmental Microbiome*, *18*(1), 28.
https://doi.org/10.1186/s40793-023-00484-y

Milicević, T., Kaliterna, J., Ivić, D., & Milović, M. (2014). Records of phytopathogenic fungal species on
native plants new to Croatia. *Natura Croatica*, *23*(1), 179–187.

Nilsson, R. H., Larsson, K. H., Taylor, A. F. S., Bengtsson-Palme, J., Jeppesen, T. S., Schigel, D., Kennedy,
P., Picard, K., Glöckner, F. O., Tedersoo, L., Saar, I., Kõljalg, U., & Abarenkov, K. (2019). The UNITE
database for molecular identification of fungi: Handling dark taxa and parallel taxonomic
classifications. *Nucleic Acids Research*, *47*(D1), D259–D264.
https://doi.org/10.1093/nar/gky1022

Palm, M. E. (1991). Taxonomy and Morphology of the Synanamorphs *Pilidium concavum* and *Hainesia*
lythri (Coelomycetes). *Mycologia*, *83*(6), 787. https://doi.org/10.2307/3760436

Panek, J., Frąc, M., & Bilińska-Wielgus, N. (2016). Comparison of chemical sensitivity of fresh and long-
stored heat resistant *Neosartorya fischeri* environmental isolates using Biolog Phenotype
MicroArray system. *PLoS ONE*, *11*(1), 4–8. https://doi.org/10.1371/journal.pone.0147605

Peschel, S., Müller, C. L., Von Mutius, E., Boulesteix, A. L., & Depner, M. (2021). NetCoMi: Network
construction and comparison for microbiome data in R. *Briefings in Bioinformatics*, *22*(4), 1–18.
https://doi.org/10.1093/bib/bbaa290

Phatak, H. C., & Payak, M. M. (1965). *Hainesia lythri* on strawberry - a new record for India. *Indian*
Phytopathology, *18*, 237–239.

Pruesse, E., Quast, C., Knittel, K., Fuchs, B. M., Ludwig, W., Peplies, J., & Glöckner, F. O. (2007). SILVA:
A comprehensive online resource for quality checked and aligned ribosomal RNA sequence data
compatible with ARB. *Nucleic Acids Research*, *35*(21), 7188–7196.
https://doi.org/10.1093/nar/gkm864

Rivers, A. R., Weber, K. C., Gardner, T. G., Liu, S., & Armstrong, S. D. (2018). ITSxpress: Software to
rapidly trim internally transcribed spacer sequences with quality scores for marker gene analysis.
F1000Research, *7*(0), 1418. https://doi.org/10.12688/f1000research.15704.1

Sayami, M., Babaeizad, V., & Fathi, J. (2013). First report of *Pilidium concavum* as the causative agent
of tan-brown rot of *Prunus domestica* fruits. *Journal of Plant Pathology*, *95*(2), 451.

Schoch, C. L., Seifert, K. A., Huhndorf, S., Robert, V., Spouge, J. L., Levesque, C. A., Chen, W., Bolchacova,
E., Voigt, K., Crous, P. W., Miller, A. N., Wingfield, M. J., Aime, M. C., An, K. D., Bai, F. Y., Barreto,
R. W., Begerow, D., Bergeron, M. J., Blackwell, M., … Schindel, D. (2012). Nuclear ribosomal
internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proceedings*
of the National Academy of Sciences of the United States of America, *109*(16), 6241–6246.
https://doi.org/10.1073/pnas.1117018109

Siegieda, D., Panek, J., & Frąc, M. (2023). Plant and soil health in organic strawberry farms – Greater
importance of fungal trophic modes and networks than α -diversity of the mycobiome. *Applied*
Soil Ecology, *188*(April), 1–15. https://doi.org/10.1016/j.apsoil.2023.104925

Siegieda, D., Panek, J., & Frąc, M. (2024). Ecological processes of bacterial microbiome assembly in
healthy and dysbiotic strawberry farms. *BMC Plant Biology*, *24*(1).
https://doi.org/10.1186/s12870-024-05415-8

Stegen, J. C., Lin, X., Fredrickson, J. K., Chen, X., Kennedy, D. W., Murray, C. J., Rockhold, M. L., &
Konopka, A. (2013). Quantifying community assembly processes and identifying features that
impose them. *ISME Journal*, *7*(11), 2069–2079. https://doi.org/10.1038/ismej.2013.93

Tamura, K., & Nei, M. (1993). *Estimation of the Number of Nucleotide Substitutions in the Control*
Region of Mitochondrial DNA in Humans and. *10*(3).

Torbati, M., Arzanlou, M., Abed-Ashtiani, F., & Golmohammadi, H. (2019). Occurrence of fruit rot on
cornelian cherry caused by *Pilidium lythri* in Iran. *Crop Protection*, *125*, 104884.
https://doi.org/10.1016/j.cropro.2019.104884

Trivedi, P., Leach, J. E., Tringe, S. G., Sa, T., & Singh, B. K. (2020). Plant–microbiome interactions: from
community assembly to plant health. *Nature Reviews Microbiology*, *18*(11), 607–621.
https://doi.org/10.1038/s41579-020-0412-1

Vishwakarma, K., Kumar, N., Shandilya, C., Mohapatra, S., Bhayana, S., & Varma, A. (2020). Revisiting
Plant–Microbe Interactions and Microbial Consortia Application for Enhancing Sustainable
Agriculture: A Review. *Frontiers in Microbiology*, *11*(December), 1–21.
https://doi.org/10.3389/fmicb.2020.560406

Ward, J. R., Kim, K., & Harvell, C. D. (2007). Temperature affects coral disease resistance and pathogen
growth. *Marine Ecology Progress Series*, *329*, 115–121. https://doi.org/10.3354/meps329115

White, T. J., Bruns, T., Lee, S., & Taylor, J. (1990). Amplification and direct sequencing of fungal
ribosomal RNA genes for phylogenetics. *PCR Protocols: A Guide to Methods and Applications*,
315–322.

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G., Hayes, A.,
Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K., Ooms, J., Robinson,
D., Seidel, D., Spinu, V., … Yutani, H. (2019). Welcome to the Tidyverse. *Journal of Open Source*
Software, *4*(43), 1686. https://doi.org/10.21105/joss.01686

Zhang, M., Li, H. L., Zhao, A. L., & Zhang, J. X. (2008). First Report of *Hainesia lythri* Causing Leaf Spots

of *Paeonia suffruticosa* in China. *Plant Disease*, *92*(3), 486–486. https://doi.org/10.1094/PDIS-92-

3-0486A