Pilidium lythri: A Climate-Driven Disruptor of Plant-Associated

Microbiomes and Ecological Stability

Dominika Siegieda[1]

Jacek Panek[1]

S. Emilia Hannula[2]

Magdalena Frąc[1]

[1]Institute of Agrophysics, Polish Academy of Sciences, Doświadczalna 4 street, 20-290 Lublin, Poland

[2]Leiden University, Institute of Environmental Sciences, Department of Environmental Biology, Leiden,

The Netherlands

Keywords

12 *Pilidium lythri*, emerging fungal phytopathogen, microbial community dynamics, plant holobiont,

13 temperature adaptability, biotic stress in plants, microbial assembly mechanisms

Abstract

Although ongoing global warming is an important driver of phytopathogen spread to new areas, little

is known on how fungal pathogens disrupt fragile balance in plant-microbe interactions. After

assessing high adaptability of emerging *Pilidium lythri* to growing temperatures, we analysed its impact

on microbiome structure of soil (bulk soil, rhizosphere) and strawberry plant (roots) niches. For this,

we employed high-throughput marker sequencing and various statistical models. Our results revealed

that *P. lythri* infection disrupted microbial community assembly processes, transitioning from

deterministic to stochastic dynamics in the rhizosphere and bulk soil, aligning with the Anna Karenina

Principle. Fungal microbiome diversity increased locally in infected rhizospheres, while microbial

migration patterns highlighted shifts in community interactions. Network analysis underscored *P.*

lythri's role in destabilizing fungal community connectivity, enhancing antagonistic interactions, and

occupying a central ecological niche. This study provides valuable insights into how spreading

phytopathogens modulate microbial equilibrium in the plant holobiont, with implications of managing

emerging threats in agriculture.

Introduction

Fungal and bacterial microorganisms, together with their host plants, are interconnected through intricate, multi-level interactions that shape ecosystems and sustain plant health and productivity. Based on these relationships, the plant holobiont concept was formed - an integrated system where hosts and microbiota function as one. Plants produce root exudates to modulate microbial assembly of theirs niches and vicinity, while the microbiome, in turn, is able to influence and regulate plant health, improving its resilience against abiotic and biotic stressors (Baldrian, 2016; Berg, 2009; Gupta et al., 2021; Marco et al., 2022; Michl et al., 2023; Trivedi et al., 2020). These finely tuned interactions are widely studied and hold great potential for improving agricultural productivity in a sustainable manner (Frąc et al., 2022; Trivedi et al., 2020; Vishwakarma et al., 2020).

However, this delicate balance between microbiomes and plants is being increasingly threatened by ongoing global warming. The rise of annual global temperature and changed precipitation patterns has created the opportunity for a spread and establishment of invasive phytopathogens into new regions (Bebber et al., 2013; Helfer, 2014; X. Liu et al., 2019). Despite this urgent challenges, limited attention has been given to understand how these emerging pathogens alter and interact with native microbial communities. This issue is particularly concerning for perennial crops like strawberries (*Fragaria x ananassa* Duch.), which depend heavily on stable rhizosphere and soil microbial communities to thrive and are particularly vulnerable to such disruptions.

Pilidium lythri, emerging fungal phytopathogen, causing damage to strawberry plantations, was originally reported in countries located in warmer climate zones, such as India (Phatak and Payak 1965), Iran (Arzanlou et al., 2013; Sayami et al., 2013) and China (Duan et al., 2010; Zhang et al., 2008). Notably, its occurrence had been recently identified in several temperate climate countries, including Belgium (Lopes et al., 2010), Croatia (Milicević et al., 2014), Spain (Aguín et al., 2018), France (Caesar et al., 2012; Cardin et al., 2009) and Poland, although in 2004 only morphologically, but in 2020 we confirmed its presence in symptomatic Polish strawberry fields with molecular methods (Gołębniak & Jarosz, 2004; Malarczyk et al., 2020), and with new isolates collected in present study. Primary targets of this fungal phytopathogen are strawberry plants and fruits (Ayoubi et al., 2016; Geng et al., 2012; Lopes et al., 2010), but it also affects other agriculturally significant plants, including *Olea europaea* L. (Arzanlou et al., 2013), *Cornus mas* (Torbati et al., 2019), *Prunus domestica* (Sayami et al., 2013), *Eucalyptus* spp. (Maússe-Sitoe et al., 2016), *Vitis vinifera* (Aguín et al., 2018), and various ornamental flowers (Bruckart et al., 2013; Caesar et al., 2012; Cardin et al., 2009; Garfinkel & Chastagner, 2019; Zhang et al., 2008). Notably concerning is that *P. lythri* is not host-specific. Strains isolated from olives were able to cause disease symptoms on strawberry fruits (Karimi et al., 2016), further increasing the

severity of the threat posed by this pathogen as it spreads to new regions. Studies conducted in Brazil,
Iran and Belgium showed incidence as high as 70% in high humidity and warm temperatures,
highlighting its threat in warming climate (Ayoubi et al., 2016; Karimi et al., 2016; Lopes et al., 2010).
Pathogenicity test revealed rapid progression of the disease, with symptoms appearing on up to 100%
of fruits within days after inoculation (Debode et al., 2011; Fernández-Ortuño et al., 2014). The
pathogen is able to infect both wounded and intact grape berries, further underscoring its high
virulence (Aguín et al., 2018). The pathogen's ability to adapt and spread under changing
environmental conditions makes it an ideal model organism to explore relationships between
emerging fungal pathogens, theirs host plants and native microbiomes. Furthermore, its prompt
proliferation and versatile host range makes it an ideal research object to evaluate how emerging
pathogens disrupt microbial balance across diverse plant microbiomes.

To explore climate adaptation and microbial interaction in emerging fungal pathogens, *Pilidium lythri*
was used as a model organism. We isolated pure fungal strains from symptomatic strawberry plants in
a Polish nursery. We compared the metabolic profiles of 3 *P. lythri* strains grown in 24°C and 30°C with
BIOLOG PM21-25 plates, identifying its sensitivity to chemical substances in response at different
temperatures to imitate climate scenarios. We also investigated the impact of pathogen's infection on
the composition and assembly mechanisms of the bacterial and fungal microbiomes of soil (bulk soil,
rhizosphere) and strawberry (roots) niches. Current work significantly advances our understanding the
microbial dynamics associated with this spreading phytopathogen, offering valuable insights into
broader ecological implications of emerging fungal pathogens.

Methods

Fungal isolates and molecular identification

Pure fungal isolates were obtained by placing symptomatic strawberry leaves and fruits (surface-
sterilized with detergent and 70% ethanol) on PDA medium and incubating them at 24°C in darkness
for 7 days. After initial screening, fungi with macroscopic and microscopic characteristics matching the
description of *Pilidium lythri* (Karimi et al., 2019; Palm, 1991) were identified through amplification and
Sanger sequencing of the Internal Transcribed Spacer (ITS) regions of fungal rDNA, with ITS1 and ITS4
primers (White et al., 1990), as previously described (Frąc et al., 2015). The obtained sequences were
deposited in GenBank under accession numbers: PQ632564.1, PQ632565.1, PQ632566.1, PQ632567.1.
These sequences were then used to construct phylogenetic tree with other sequences available in
Genbank database from the *Pilidium* genus. Initial tree for the heuristic search was obtained
automatically by applying Neighbor-Join and BioNJ algorithms to a matrix of pairwise distances
estimated using the Tamura-Nei model, and then selecting the topology with superior log likelihood

value. The tree was drawn to scale, with branch lengths measured in the number of substitutions per

site. This analysis involved 57 nucleotide sequences and there were a total of 372 positions in the final

dataset. Evolutionary analyses were conducted in MEGA X (Kumar et al. 2018). To fulfil Koch's

postulates, a spore suspension of $1.6 \cdot x\ 10^4$ spores/mL was prepared, and surface-sterilized strawberry

leaves were briefly submerged in the solution. The leaves were then sealed in ziplock bags and

incubated at 24°C in darkness for 7 days. Sterile deionized water was used as the untreated control.

Metabolic profile arrays

Each cell of Phenotype Microarray 21-25 BIOLOG plates (Biolog, Hayward, CA, USA) was inoculated

with the fungal mycelium as recommended by the manufacturer and described before (Panek et al.,

2016). For each plate type, 3 *Pilidium lythri* strains in triplicates and in two temperatures (24 and 30°C)

were incubated for 96 hours in the darkness. Microplate reader (Biolog, Hayward, CA, USA) was used

to determine the optical density at 490 nm at the beginning and every 24 hours. Obtained data was

analysed statistically and visualised in RStudio, using tidyverse package (Wickham et al., 2019). First,

values obtained at 0h for each replicate was divided from read values at 96h, and values <0 were

assigned as 0. To compare differences in fungal growth between different temperatures for each

concentration we performed Wilcoxon test with False Discovery Rate method to correct for multiple

comparisons. Next, to differentiate growth of different strains, we used Kruskal-Wallis method with

post-hoc pairwise comparison with Wilcoxon test.

Pot experiment and isolation of microbial DNA

Strawberry plants, obtained from the nursery, were grown in pots containing sieved soil from a

certified organic strawberry plantation. After 20 days of initial growth of strawberry plants at 28°C and

70% humidity, half of the pots were inoculated with *P. lythri* spores using 10 mL of a $2 \cdot 10^4$ spores mL^{-1}

^{1} solution per plant. Sterile, deionized water was used as a negative control. Samples were collected

from two pots in triplicate (6 samples per niche per time point) at the time of inoculation and again

after 9 days of incubation. Bulk soil was transferred into sterile 1.5 mL Eppendorf tubes and frozen at

-20°C. The rhizosphere was collected as described by Lundberg et al. (Lundberg et al., 2012). Roots,

flushed free of rhizosphere soil, were also frozen at -20°C prior to DNA isolation which was conducted

with the QIAQUBE instrument (Qiagen, Hilden, Germany) and GeneMATRIX Soil DNA Purification Kit

(EURx, Gdańsk, Poland) after homogenization of the samples in the FastPrep-24 instrument (MP

Biomedicals) at 4 m s^{-1} for 15 s. DNA isolates were then frozen in -20°C until library construction and

sequencing.

 Metataxonomic sequencing, bioinformatics and statistical analysis

For the metataxonomic sequencing, microbial DNA ITS2 (Schoch et al., 2012) and 16Sv3 regions

(Klindworth et al., 2013) were amplified, indexed and sequenced on Illumina MiSeq instrument

(2x300), and raw data was then passed through the custom QIIME2 (Bolyen et al., 2019) pipelines as

reported before (Siegieda et al., 2023, 2024). In brief, after primer trimming with Cutadapt (Martin,

2011) and ITSxpress (Rivers et al., 2018), denoising, chimera removal, merging and Amplicon Sequence

Variant (ASV) calling was made. The taxonomy was assigned based on UNITE v9 dynamic release

(Nilsson et al., 2019) and Silva 138 (Pruesse et al., 2007) databases. The results were then analysed

with R (v. 4.0.3), using libraries: phyloseq (McMurdie & Holmes, 2013), tidyverse (Wickham et al.,

2019), microeco (C. Liu et al., 2021), microbiomeMarker (Cao et al., 2022), NetCoMi (Peschel et al.,

2021). First, for ITS data, only ASVs identified to Fungi and Stramenopiles; and for 16S - Bacteria and

Archaea. Then, ITS data was repeatedly rarefied at 6,000 and 16S at 11,000 reads using mirlyn package

(Cameron et al., 2021; McMurdie & Holmes, 2014). We employed Effective Number of Species (ENS)

(Jost, 2006) on ASV level to evaluate the richness of microbiomes in each niche. We used supervised

machine learning model to evaluate the fungal biomarkers of each niche for healthy and infected

samples with Supervised Machine Method (SVM), using Total Sum Scaling normalization method.

SourceTracker was used for tracking migration between niches (Knights et al., 2011). RCbray (Bray-

Curtis dissimilarity based Raup-Crick metric) and βNTI (β Nearest Taxon) were used to evaluate the

ecological processes that dominated the bacterial assemblies (C. Liu et al., 2017; Stegen et al., 2013).

When βNTI fell between -2 and 2, microbial assembly was dominated by stochastic mechanisms, and

below -2 and higher that 2 - by deterministic. Microbial networks were constructed with 1,000 most

frequent taxa (using Pearson correlation measure) and fast greedy clustering method.

Results

Fungal isolates

Fungal ITS sequences, obtained from symptomatic strawberry leaves (G320/23, G321/23, G322/23 and

G323/23) showed 100% similarity to those of *Pilidium lythri* already present in the GenBank database

and phylogenetic tree revealed clustering of the sequences with other representatives of *Pilidium*

lythri. The strains collected after 7 days in 23°C in the darkness on Potato Dextrose Agar were flat and

granulose, cinnamon in colour (Karimi et al., 2016; Marin-Felix et al., 2017). The presence of hyaline

conidia with smooth walls and slightly pointed tips were confirmed under the microscope. After the

inoculation of the surface-sterilized strawberry leaves with the spore suspension and incubation in

24°C in darkness for 7 days, 100% of the leaves showed brown necrotic lesions. Leaves submerged in

water did not show symptoms of the disease. Growth of the cinnamon-colour fungal strains were

observed after incubation (24°C for 7 days) of infested leaves on the Petri dishes with the PDA medium,

fulfilling he Koch's postulate (Figure 1).

Figure 1. A) Phylogenetic tree of the evolutionary history of *P. lythri* was inferred by using the Maximum Likelihood method
and Tamura-Nei model (Tamura & Nei, 1993) and the e tree with the highest log likelihood (-1144.51) is shown. The
percentage of trees in which the associated taxa clustered together is shown next to the branches. B,C) Symptoms of the
Pilidium lythri infection on the strawberry leaves and in the D) control leaf. E) Growth of *P. lythri* after 7 days of growth in 24
°C on PDA medium, F) conidia of *P. lythri* (40x), and G) growth on PDA medium with infected leaf.

Chemical sensitivity

To evaluate *Pilidium lythri*'s sensitivity to various chemicals, BIOLOG PM 21-25 plates were we

incubated of three strains at 24°C and 30°C for 96 hours. These plates are coated with 4 different

concentrations of the chemical substances. The precise concentrations of the substances used for

microarrays are confidential and protected as proprietary information by the manufacturer. For the

purposes of this article, we refer to them as concentration levels 1, 2, 3, and 4 to maintain clarity. The

substances were grouped in 8 groups, based on their structure and function: antibiotics,

antimicrobials, chelators, drugs (used in medicine), membrane function compounds, nitrogen

compounds, salts and others (Panek et al., 2016). Overall, strains exhibited lower sensitivity to

chemical compounds at higher temperature (30 °C) in concentration 3 and 4 (p-values < 0.01,

Supplementary Figure 1). We also observed variation between strains in higher temperature

(Supplementary Figure 2), where the strain G321/23 was characterized with the highest sensitivity,

and G322/23 the lowest. Increase in substance concentration showed decrease in median fungal

growth in both temperatures (excluding zeros) (Supplementary Table 1). Alarmingly, only Niaproof (in

level 4 concentration) effectively inhibited growth off all 3 strains at both temperatures (Figure 2,

Supplementary Figure 3, Supplementary Figure 4, Supplementary Figure 5).

Figure 2. BIOLOG PM21-25 results of the cells with highest concentration of chemical substances (concentration level 4). Shapes represent different strains of *Pilidium lythri*. Gray symbols represent no growth, red - growth of the strain after 96 hours. Dashed vertical grey line represents average response for each chemical substrate in each temperature (excluding zeros).

Fungal and bacterial community changes after *P. lythri* inoculation of strawberry

A total of 72 samples from soil (2x12), rhizosphere (2x12), and roots (2x12) were amplified and sequenced. Rarefaction curves plateaued for all samples, indicating sufficient sequencing depth. To evaluate the presence of *Pilidium lythri* in infected samples, we employed a predictive approach to identify 15 most important bacterial and fungal taxa that differentiate healthy and infected strawberry compartments. The analysis revealed that the *Pilidium* fungi belonged to the most important taxa that differentiate between healthy and infected plants and soil (Figure 3). This, along with the visible symptoms of the disease, confirmed that the infection was successful and the pathogen migrated from the soil to rhizosphere and roots.

Figure 3. Top 15 fungal taxa found more abundant in (orange) infected and in (blue) healthy soils and plants identified based on Supervised Vector Machine machine learning approach.

After evaluating that infection was successful, statistical analysis was conducted to compare the alpha diversity of bacterial and fungal communities between healthy and inoculated (infected) niches. No significant differences in richness were observed between healthy and infected niches at both sampling times (p > 0.05) (Figure 4). Only fungal community in the infected rhizosphere at the second sampling displayed higher richness (p < 0.05). Notably, bacterial communities showed a significant increase richness across all three compartments at the second sampling time (p < 0.05), whereas fungal community richness remained stable across time points (p > 0.05). Additionally, both bacterial and fungal communities exhibited a gradient of decreasing diversity from soil to rhizosphere and root at second sampling.

Figure 4. Comparison of alpha diversity (in Effective Number of Species) of bacterial (A) and fungal (B) communities of strawberry niches immediately after *P. lythri* inoculation (one) and after 9 days of incubation in 25°C and ~70% of humidity (two). Higher result represent full model comparison, and lower - pairwise comparisons within each sampling. Light blue color represents samples without pathogen inoculation, and orange - with inoculation.

To elucidate the influence of *Pilidium lythri* on microbial migration between the niches, we used samples from sampling at 0 days as a source, and samples from sampling at 9 days as sinks, and did this both fungal and bacterial communities. In bulk soil, only bacterial community revealed significantly higher share of microorganisms in bulk soil and a lower portion from unknown source in infected samples (Figure 4A). Furthermore, we observed migration of bacteria from bulk soil to rhizosphere for both bacteria and fungi in infected communities, but also from rhizosphere to rhizosphere for bacteria in infected communities. Finally, analysis revealed increased migration from rhizosphere to roots in infected samples in bacterial community.

Figure 5. Source tracking analysis (A) of bacterial and fungal microbiomes between sampling one (source) and two (sink). Comparison between healthy (light blue, left) and contaminated (light orange, right) samples. Values are shown as mean standard error from bootstrap. (B) BetaNRI idenx analysis of bacterial community in second sampling. Significant differences on both panels between healthy and contaminated samples are shown with asterisk.

We also evaluated the mechanisms involved in bacterial community assembly by calculating the betaNRI index for bulk soil, rhizosphere, and root samples collected during the second sampling period (Figure 4B). The results revealed a dominance of stochastic processes in both healthy and infected rhizosphere and root samples of strawberry (betaNRI values between -2 and 2). Furthermore, we observed differences in community assembly processes between healthy and infected bulk soil, with a shift from deterministic processes (characterized by lower turnover) in healthy bulk soil to a dominance of stochastic processes in infected soil.

Finally, to evaluate the influence of *Pilidium lythri* on the structure of fungal networks, we compared the network structure in the rhizosphere niche between control and infected samples during the second sampling (Figure 6A). Both networks showed similar characteristics in terms of the number of components (1), natural connectivity (0.339, 0.302), and vertex and edge connectivity (72, 69). On the other hand, the control rhizosphere exhibited a higher percentage of positive edges (50.15% vs. 46.55%) but lower average dissimilarity and average path length (0.644 vs. 0.670). Notably, the ASV

240　identified as *Pilidium* demonstrated higher values of betweenness (0 vs. 3) and closeness centrality

241　(1170 vs. 154) in the infected samples.

242　Figure 6. Fungal network analysis in A) control and B) infected rhizosphere samples of strawberry. Each node represents

243　individual fungal ASV, and node size is proportional to number of reads. Node colours represent the same clusters. Green

244　lines represent positive interactions and red- negative. The black asterisk shows *Pilidium*.

Discussion

To close the knowledge gap regarding emerging fungal pathogens and theirs influence on native plants

we used newly isolated *Pilidium lythri* as a model organism. We identified 3 strains obtained from

symptomatic strawberry plants in Polish nursery by sequencing of the ITS region and evaluated the

impact of temperature growth on inhibition of fungal development with 112 different chemical

substances. We then evaluated the impact of this phytopathogen on the structure of bacterial and

fungal metacommunities in soil and plant niches (bulk soil, rhizosphere, roots) of strawberry.

In vitro evaluation of the impact of various chemical substances on the growth and development of

the tested phytopathogen at two temperatures using BIOLOG microarrays raised serious concerns, as

Pilidium lythri demonstrated lower susceptibility to chemicals at higher temperatures, with most of

the tested compounds showing limited or no inhibitory effects on fungal growth. Notably, increased

temperature allowed the pathogen to grow even in the presence of these substances, highlighting the

high potential for environmental adaptability of this fungus. Global warming has indeed been widely

reported to contribute to the spread of fungal phytopathogens to new areas (Bebber et al., 2013;

Chaloner et al., 2021; Ward et al., 2007). Among the wide variety of chemical substances tested, only

Niaproof showed inhibition at both temperatures, and could be further tested to evaluate their

antifungal effects. Since Niaproof (tetradecyl sodium sulfate) is an anionic surface-active agent, its

antifungal mechanism may involve disrupting the fungal cell membrane, thus preventing the pathogen

from surviving in the environment.

As the dissemination of fungal pathogens, is widely reported (Bebber et al., 2013), still little is known

about the effect of such invasive microorganisms on the native microbial structure of the plant

holobiont. Recent studies have reported significant differences in the α-diversity of microbiome

structure between healthy and infected soil and plant niches (Bakker et al., 2018; Delgado-Baquerizo

et al., 2016; Lefcheck et al., 2015). In the present study, we compared the diversity of bacterial and

fungal metacommunities in healthy and *Pilidium lythri*-infected bulk soil, rhizosphere, and roots at two

time points: directly after pathogen inoculation and 9 days post-inoculation. The results revealed

strong evidence of increased α-diversity in the bacterial community 9 days after inoculation in both

healthy and inoculated niches, suggesting that bacterial communities are less influenced by *Pilidium*

lythri inoculation and more responsive to environmental factors. To test this, we analyzed the

mechanisms involved in bacterial microbiome assembly by calculating the βNRI index, treating the first

sampling as the source of microorganisms and the second sampling as the sinks. This index measures

how the differences in species composition between the two samplings differ from what would be

expected by the null model of random assembly. The results indicated that composition of bacterial

communities in healthy and diseased rhizosphere and roots samples was shaped by stochastic

processes (i.e. random colonization, dispersal, and ecological drift). Interestingly, we observed a shift

from deterministic processes in healthy bulk soil, to stochastic processes in *Pilidium lythri*-infected

samples, indicating a disruption in microbial turnover. Infected soil showed random processes that

could have been triggered by disturbances in plant biology caused by the infection, described by the

Anna Karenina Principle (AKP)(Arnault et al., 2023). The scenario described here can be explained by

the loss of plant control over microbiome assembly, resulting in a temporal AKP effect (C. Cheng et al.,

2013; Y. T. Cheng et al., 2019). Over time, this could lead to the recovery of the plant's control over

microbial assembly or the introduction of new deterministic processes to control bacterial assembly

(Gao et al., 2021). Analysis of microbial migration suggests increased migration of bacteria from the

soil in diseased samples, as well as increased migration of bacteria from the rhizosphere to the roots

in infected plant samples. This inconsistency between migration (suggesting plant filtering) and

microbial assembly (suggesting random processes) could be caused by the smaller scale of

deterministic processes being masked by larger-scale microbial community assembly. Furthermore,

these two mechanisms represent different time scales, where βNRI focuses on broader, longer-term

assembly processes (Arnault et al., 2023). Importantly, migration of microorganisms can be caused by

random dispersal (stochastic processes) but may appear deterministic as migration increases. Thus,

these two processes are not necessarily contradictory.

In the fungal microbiome, we observed a localized increase in α-diversity in the infected rhizosphere at the second sampling, which contrasts with recent results from strawberry fields that showed no differences in microbiome composition based on health status (Siegieda et al., 2023). We also observed increased migration of fungi from bulk soil to the rhizosphere, highlighting the importance of this niche in fungal microbiome dynamics and the role of plant in microbial recruitment under pathogen stress (Bais et al., 2006; Berendsen et al., 2012). To further asses the influence of the pathogen in microbial metacommunity, we analyzed the fungal networks in the rhizosphere. Comparison of healthy and infected samples revealed the important role *Pilidium lythri* plays in rhizosphere networks and overall microbiome assembly. The fungal networks in infected and control samples were similar, yet infected rhizospheres exhibited lower percentages of positive edges and longer average path lengths. This suggests that the presence of the pathogen does not significantly disrupt fungal networks but does decrease network connectivity, indicating a shift toward increased competition and antagonism within the fungal microbiome. This could destabilize the fungal meta-community, making it more vulnerable to further infections and environmental stressors (Hernandez et al., 2021). Furthermore, we noted the central role of *Pilidium lythri*, as it exhibited higher betweenness and closeness centrality in the infected samples. This suggests that the pathogen plays a pivotal role in reshaping the fungal community balance. Increased betweenness centrality indicates that the pathogen acts as a key connector in the infected rhizosphere, potentially by disrupting connections between beneficial fungal microorganisms or recruiting microbial partners to facilitate its distribution within and between the soil and plant niches.

Concluding, in this study, we isolated and molecularly identified pure isolates of spreading fungal phytopathogen from symptomatic strawberry plants - *Pilidium lythri*. We observed pathogen's low sensitivity against various chemical substances in 30°C compared with 24°C, highlighting dangers of its spread due to climate change. We revealed that only Niaproof is a promising substance that could be used to control its spread, but further investigations on its safety in the field conditions and mechanisms of actions are necessary. We used *P. lythri* as a model phytopathogenic microorganism to evaluate ecological impacts of spreading fungal phytopathogens on native microorganism metacommunities in bulk soil, rhizosphere and roots of strawberry with high-throughput marker sequencing. We revealed that *Pilidium lythri* infection had a major effect on both the structure and dynamics of the plant associated microbiome. Its ability to shift bacterial microbial assembly mechanism from deterministic to stochastic in the bulk soil, underscores its capacity to destabilize plant-associated microbiomes. Observed mechanism aligned with Anna Karenina Principle, suggesting loss of control over microbiota assembly in infected soil, which is particularly important for perennial crops, such as strawberry. Additionally, we observed that *P. lythri* plays an important role in fungal network in the strawberry rhizosphere, where it shifted the fungal microbiome dynamics towards

competition and antagonism, amplifying vulnerability of strawberry rhizosphere to secondary infections and other (a)biotic stresses.

These findings underline the critical role of emerging fungal pathogens like *Pilidium lythri* in destabilizing native plant-microbiome interactions. This is significantly important in the context of global warming, which accelerates spread and impact of such pathogens.

References

Aguín, O., Ferreiroa, V., González-Jartín, J. M., Alfonso, A., Botana, L. M., Mansilla, J. P., & Sainz, M. J. (2018). *Pilidium lythri* is associated with bunch rot of grapevine (Vitis vinifera). *American Journal of Enology and Viticulture*, *69*(4), 410–416. https://doi.org/10.5344/ajev.2018.17093

Arnault, G., Mony, C., & Vandenkoornhuyse, P. (2023). Plant microbiota dysbiosis and the Anna Karenina Principle. *Trends in Plant Science*, *28*(1), 18–30. https://doi.org/10.1016/j.tplants.2022.08.012

Arzanlou, M., Torbati, M., & Jafary, H. (2013). Fruit rot on olive caused by *Pilidium concavum* in Iran. *Australasian Plant Disease Notes*, *8*(1), 117–121. https://doi.org/10.1007/s13314-013-0111-0

Ayoubi, N., Soleimani, M. J., & Zare, R. (2016). Pilidium concavum, causing tan-brown rot on strawberry in Iran. *Journal of Plant Pathology*, *98*(3), 667–669. https://doi.org/10.4454/JPP.V98I3.063

Bais, H. P., Weir, T. L., Perry, L. G., Gilroy, S., & Vivanco, J. M. (2006). The role of root exudates in rhizosphere interactions with plants and other organisms. *Annual Review of Plant Biology*, *57*(1), 233–266. https://doi.org/10.1146/annurev.arplant.57.032905.105159

Bakker, P. A. H. M., Pieterse, C. M. J., de Jonge, R., & Berendsen, R. L. (2018). The Soil-Borne Legacy. *Cell*, *172*(6), 1178–1180. https://doi.org/10.1016/j.cell.2018.02.024

Baldrian, P. (2016). Forest microbiome: diversity, complexity and dynamics. *FEMS Microbiology Reviews*, fuw040. https://doi.org/10.1093/femsre/fuw040

Bebber, D. P., Ramotowski, M. A. T., & Gurr, S. J. (2013). Crop pests and pathogens move polewards in a warming world. *Nature Climate Change*, *3*(11), 985–988. https://doi.org/10.1038/nclimate1990

Berendsen, R. L., Pieterse, C. M. J., & Bakker, P. A. H. M. (2012). The rhizosphere microbiome and plant health. *Trends in Plant Science*, *17*(8), 478–486. https://doi.org/10.1016/j.tplants.2012.04.001

Berg, G. (2009). Plant–microbe interactions promoting plant growth and health: perspectives for controlled use of microorganisms in agriculture. *Applied Microbiology and Biotechnology*, *84*(1), 11–18. https://doi.org/10.1007/s00253-009-2092-7

Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., Alexander, H.,
Alm, E. J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J. E., Bittinger, K., Brejnrod, A., Brislawn, C.
J., Brown, C. T., Callahan, B. J., Caraballo-Rodríguez, A. M., Chase, J., … Caporaso, J. G. (2019).
Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nature
Biotechnology*, *37*(8), 852–857. https://doi.org/10.1038/s41587-019-0209-9

Bruckart, W. L., Eskandari, F. M., Coombs, E. M., Rossman, A. Y., & Palm, M. E. (2013). First report of
Pilidium concavum causing leaf necrosis on Fallopia japonica in the United States. *Plant Disease*,
97(1), 146–146. https://doi.org/10.1094/PDIS-07-12-0688-PDN

Caesar, A. J., Lartey, R. T., & Caesar-TonThat, T. (2012). First report of *Pilidium concavum* causing root
lesions of meadow hawkweed in France. *Plant Disease*, *96*(12), 1830–1830.
https://doi.org/10.1094/PDIS-05-12-0509-PDN

Cameron, E. S., Schmidt, P. J., Tremblay, B. J. M., Emelko, M. B., & Müller, K. M. (2021). Enhancing
diversity analysis by repeatedly rarefying next generation sequencing data describing microbial
communities. *Scientific Reports*, *11*(1), 1–13. https://doi.org/10.1038/s41598-021-01636-1

Cao, Y., Dong, Q., Wang, D., Zhang, P., Liu, Y., & Niu, C. (2022). microbiomeMarker: an R/Bioconductor
package for microbiome marker identification and visualization. *Bioinformatics*, *38*(16), 4027–
4029. https://doi.org/10.1093/bioinformatics/btac438

Cardin, L., Vincenot, L., & Balesdent, M. H. (2009). First report of Pilidium concavum on Bergenia
crassifolia in France. *Plant Disease*, *93*(5), 548–548. https://doi.org/10.1094/PDIS-93-5-0548B

Chaloner, T. M., Gurr, S. J., & Bebber, D. P. (2021). Plant pathogen infection risk tracks global crop
yields under climate change. *Nature Climate Change*, *11*(8), 710–715.
https://doi.org/10.1038/s41558-021-01104-8

Cheng, C., Gao, X., Feng, B., Sheen, J., Shan, L., & He, P. (2013). Plant immune response to pathogens
differs with changing temperatures. *Nature Communications*, *4*(1), 2530.
https://doi.org/10.1038/ncomms3530

Cheng, Y. T., Zhang, L., & He, S. Y. (2019). Plant-Microbe Interactions Facing Environmental Challenge.
Cell Host & Microbe, *26*(2), 183–192. https://doi.org/10.1016/j.chom.2019.07.009

Debode, J., Van Hemelrijck, W., Heungens, K., Maes, M., & Creemers, P. (2011). First Report of Pilidium
concavum Causing Tan-Brown Rot on Strawberry Fruit in Belgium. *Plant Disease*, *95*(8), 1029–
1029. https://doi.org/10.1094/PDIS-10-10-0752

Delgado-Baquerizo, M., Maestre, F. T., Reich, P. B., Jeffries, T. C., Gaitan, J. J., Encinar, D., Berdugo, M.,
Campbell, C. D., & Singh, B. K. (2016). Microbial diversity drives multifunctionality in terrestrial
ecosystems. *Nature Communications*, *7*, 1–8. https://doi.org/10.1038/ncomms10541

Duan, Y. B., Kang, Y. B., & Yu, Z. Z. (2010). First report of *Pilidium concavum* on *Paeonia suffruticosa* in
China. *Plant Disease*, *94*(2), 271–271. https://doi.org/10.1094/PDIS-94-2-0271B

Fernández-Ortuño, D., Bryson, P. K., & Schnabel, G. (2014). First Report of Pilidium concavum Causing
Tan-brown Rot on Strawberry Nursery Stock in South Carolina. *Plant Disease*, *98*(7), 1010–1010.
https://doi.org/10.1094/PDIS-01-14-0048-PDN

Frąc, M., Hannula, E. S., Bełka, M., Salles, J. F., & Jedryczka, M. (2022). Soil mycobiome in sustainable
agriculture.   *Frontiers   in   Microbiology*,   *13*(November),   1–9.
https://doi.org/10.3389/fmicb.2022.1033824

Frąc, M., Jezierska-Tys, S., & Yaguchi, T. (2015). Occurrence, detection, and molecular and metabolic
characterization of heat-resistant fungi in soils and plants and their risk to human health.
Advances in Agronomy, *132*, 161–204.

Gao, M., Xiong, C., Gao, C., Tsui, C. K. M., Wang, M. M., Zhou, X., Zhang, A. M., & Cai, L. (2021). Disease-
induced changes in plant microbiome assembly and functional adaptation. *Microbiome*, *9*(1), 1–
18. https://doi.org/10.1186/s40168-021-01138-2

Garfinkel, A. R., & Chastagner, G. A. (2019). Survey reveals a broad range of fungal pathogens and an
oomycete on peonies in the United States. *Plant Health Progress*, *20*(4), 250–254.
https://doi.org/10.1094/PHP-09-19-0065-S

Geng, W. L., Hu, P., Ma, Z., Zhao, X. Y., & Wei, Y. M. (2012). First Report of Pilidium concavum Causing
Tan-Brown Leaf Spot on Strawberry in China. *Plant Disease*, *96*(9), 1377–1377.
https://doi.org/10.1094/PDIS-03-12-0310-PDN

Gołębniak, B., & Jarosz, A. (2004). First report of tan-brown rot (*Hainesia lythri*) on strawberry fruits in
Poland. *Phytopathogia Polonica*, *31*, 57–60.

Gupta, R., Anand, G., Gaur, R., & Yadav, D. (2021). Plant–microbiome interactions for sustainable
agriculture: a review. *Physiology and Molecular Biology of Plants*, *27*(1), 165–179.
https://doi.org/10.1007/s12298-021-00927-1

Helfer, S. (2014). Rust fungi and global change. *New Phytologist*, *201*(3), 770–780.
https://doi.org/10.1111/nph.12570

Hernandez, D. J., David, A. S., Menges, E. S., Searcy, C. A., & Afkhami, M. E. (2021). Environmental stress destabilizes microbial networks. *ISME Journal*, *15*(6), 1722–1734. https://doi.org/10.1038/s41396-020-00882-x

Jost, L. (2006). Entropy and diversity. *Oikos*, *113*(2), 363–375. https://doi.org/10.1111/j.2006.0030-1299.14714.x

Karimi, K., Arzanlou, M., Babai-ahari, A., & Pertot, I. (2016). Biological and molecular characterisation of *Pilidium lythri*, an emerging strawberry pathogen in Iran. *Phytopathologia Mediterranea*, *55*(3), 366–379. https://doi.org/10.14601/Phytopathol_Mediterr-18391

Karimi, K., Arzanlou, M., & Pertot, I. (2019). Development of novel species-specific primers for the specific identification of *Colletotrichum nymphaeae* based on conventional PCR and LAMP techniques. *European Journal of Plant Pathology*. https://doi.org/10.1007/s10658-019-01895-9

Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., & Glöckner, F. O. (2013). Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Research*, *41*(1), e1–e1. https://doi.org/10.1093/nar/gks808

Knights, D., Kuczynski, J., Charlson, E. S., Zaneveld, J., Mozer, M. C., Collman, R. G., Bushman, F. D., Knight, R., & Kelley, S. T. (2011). Bayesian community-wide culture-independent microbial source tracking. *Nature Methods*, *8*(9), 761–765. https://doi.org/10.1038/nmeth.1650

Lefcheck, J. S., Byrnes, J. E. K., Isbell, F., Gamfeldt, L., Griffin, J. N., Eisenhauer, N., Hensel, M. J. S., Hector, A., Cardinale, B. J., & Duffy, J. E. (2015). Biodiversity enhances ecosystem multifunctionality across trophic levels and habitats. *Nature Communications*, *6*. https://doi.org/10.1038/ncomms7936

Liu, C., Cui, Y., Li, X., & Yao, M. (2021). microeco: an R package for data mining in microbial community ecology. *FEMS Microbiology Ecology*, *97*(2). https://doi.org/10.1093/femsec/fiaa255

Liu, C., Yao, M., Stegen, J. C., Rui, J., Li, J., & Li, X. (2017). Long-term nitrogen addition affects the phylogenetic turnover of soil microbial community responding to moisture pulse. *Scientific Reports*, *7*(1), 1–11. https://doi.org/10.1038/s41598-017-17736-w

Liu, X., Ma, Z., Cadotte, M. W., Chen, F., He, J.-S., & Zhou, S. (2019). Warming affects foliar fungal diseases more than precipitation in a Tibetan alpine meadow. *New Phytologist*, *221*(3), 1574–1584. https://doi.org/10.1111/nph.15460

Lopes, U. P., Zambolim, L., Lopes, U. N., Pereira, O. L., & Costa, H. (2010). First report of *Pilidium*
concavum causing tan-brown rot in strawberry fruits in Brazil. *Plant Pathology*, *59*, 1171.
https://doi.org/10.1111/j.1365-3059.2010.02336.x

Lundberg, D. S., Lebeis, S. L., Paredes, S. H., Yourstone, S., Gehring, J., Malfatti, S., Tremblay, J.,
Engelbrektson, A., Kunin, V., Rio, T. G. Del, Edgar, R. C., Eickhorst, T., Ley, R. E., Hugenholtz, P.,
Tringe, S. G., & Dangl, J. L. (2012). Defining the core *Arabidopsis thaliana* root microbiome.
Nature, *488*(7409), 86–90. https://doi.org/10.1038/nature11237

Malarczyk, D. G., Panek, J., & Frąc, M. (2020). Triplex Real-Time PCR Approach for the Detection of
Crucial Fungal Berry Pathogens—*Botrytis* spp., *Colletotrichum* spp. and *Verticillium* spp.
International Journal of Molecular Sciences, *21*(22), 8469. https://doi.org/10.3390/ijms21228469

Marco, S., Loredana, M., Riccardo, V., Raffaella, B., Walter, C., & Luca, N. (2022). Microbe-assisted crop
improvement: a sustainable weapon to restore holobiont functionality and resilience.
Horticulture Research, *9*(October). https://doi.org/10.1093/hr/uhac160

Marin-Felix, Y., Groenewald, J. Z., Cai, L., Chen, Q., Marincowitz, S., Barnes, I., Bensch, K., Braun, U.,
Camporesi, E., Damm, U., de Beer, Z. W., Dissanayake, A., Edwards, J., Giraldo, A., Hernández-
Restrepo, M., Hyde, K. D., Jayawardena, R. S., Lombard, L., Luangsa-ard, J., … Crous, P. W. (2017).
Genera of phytopathogenic fungi: GOPHY 1. *Studies in Mycology*, *86*, 99–216.
https://doi.org/10.1016/j.simyco.2017.04.002

Martin, M. (2011). Cutadapt removes adapter sequences from hight-throughput sequencing reads.
EMBnet.Journal, *17*(1), 10–12. https://doi.org/https://doi.org/10.14806/ej.17.1.200

Maússe-Sitoe, S. N., Chen, S., Wingfield, M. J., & Roux, J. (2016). Diseases of eucalypts in the central
and northern provinces of Mozambique. *Southern Forests: A Journal of Forest Science*, *78*(3),
169–183. https://doi.org/10.2989/20702620.2015.1126780

McMurdie, P. J., & Holmes, S. (2013). Phyloseq: An R Package for Reproducible Interactive Analysis and
Graphics of Microbiome Census Data. *PLoS ONE*, *8*(4).
https://doi.org/10.1371/journal.pone.0061217

McMurdie, P. J., & Holmes, S. (2014). Waste Not, Want Not: Why Rarefying Microbiome Data Is
Inadmissible. *PLoS Computational Biology*, *10*(4), e1003531.
https://doi.org/10.1371/journal.pcbi.1003531

Michl, K., Berg, G., & Cernava, T. (2023). The microbiome of cereal plants: The current state of
knowledge and the potential for future applications. *Environmental Microbiome*, *18*(1), 28.
https://doi.org/10.1186/s40793-023-00484-y

Milicević, T., Kaliterna, J., Ivić, D., & Milović, M. (2014). Records of phytopathogenic fungal species on
native plants new to Croatia. *Natura Croatica*, *23*(1), 179–187.

Nilsson, R. H., Larsson, K. H., Taylor, A. F. S., Bengtsson-Palme, J., Jeppesen, T. S., Schigel, D., Kennedy,
P., Picard, K., Glöckner, F. O., Tedersoo, L., Saar, I., Kõljalg, U., & Abarenkov, K. (2019). The UNITE
database for molecular identification of fungi: Handling dark taxa and parallel taxonomic
classifications. *Nucleic Acids Research*, *47*(D1), D259–D264.
https://doi.org/10.1093/nar/gky1022

Palm, M. E. (1991). Taxonomy and Morphology of the Synanamorphs *Pilidium concavum* and *Hainesia*
lythri (Coelomycetes). *Mycologia*, *83*(6), 787. https://doi.org/10.2307/3760436

Panek, J., Frąc, M., & Bilińska-Wielgus, N. (2016). Comparison of chemical sensitivity of fresh and long-
stored heat resistant *Neosartorya fischeri* environmental isolates using Biolog Phenotype
MicroArray system. *PLoS ONE*, *11*(1), 4–8. https://doi.org/10.1371/journal.pone.0147605

Peschel, S., Müller, C. L., Von Mutius, E., Boulesteix, A. L., & Depner, M. (2021). NetCoMi: Network
construction and comparison for microbiome data in R. *Briefings in Bioinformatics*, *22*(4), 1–18.
https://doi.org/10.1093/bib/bbaa290

Phatak, H. C., & Payak, M. M. (1965). *Hainesia lythri* on strawberry - a new record for India. *Indian*
Phytopathology, *18*, 237–239.

Pruesse, E., Quast, C., Knittel, K., Fuchs, B. M., Ludwig, W., Peplies, J., & Glöckner, F. O. (2007). SILVA:
A comprehensive online resource for quality checked and aligned ribosomal RNA sequence data
compatible with ARB. *Nucleic Acids Research*, *35*(21), 7188–7196.
https://doi.org/10.1093/nar/gkm864

Rivers, A. R., Weber, K. C., Gardner, T. G., Liu, S., & Armstrong, S. D. (2018). ITSxpress: Software to
rapidly trim internally transcribed spacer sequences with quality scores for marker gene analysis.
F1000Research, *7*(0), 1418. https://doi.org/10.12688/f1000research.15704.1

Sayami, M., Babaeizad, V., & Fathi, J. (2013). First report of *Pilidium concavum* as the causative agent
of tan-brown rot of *Prunus domestica* fruits. *Journal of Plant Pathology*, *95*(2), 451.

Schoch, C. L., Seifert, K. A., Huhndorf, S., Robert, V., Spouge, J. L., Levesque, C. A., Chen, W., Bolchacova,
E., Voigt, K., Crous, P. W., Miller, A. N., Wingfield, M. J., Aime, M. C., An, K. D., Bai, F. Y., Barreto,

R. W., Begerow, D., Bergeron, M. J., Blackwell, M., … Schindel, D. (2012). Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proceedings of the National Academy of Sciences of the United States of America*, *109*(16), 6241–6246. https://doi.org/10.1073/pnas.1117018109

Siegieda, D., Panek, J., & Frąc, M. (2023). Plant and soil health in organic strawberry farms – Greater importance of fungal trophic modes and networks than α -diversity of the mycobiome. *Applied Soil Ecology*, *188*(April), 1–15. https://doi.org/10.1016/j.apsoil.2023.104925

Siegieda, D., Panek, J., & Frąc, M. (2024). Ecological processes of bacterial microbiome assembly in healthy and dysbiotic strawberry farms. *BMC Plant Biology*, *24*(1). https://doi.org/10.1186/s12870-024-05415-8

Stegen, J. C., Lin, X., Fredrickson, J. K., Chen, X., Kennedy, D. W., Murray, C. J., Rockhold, M. L., & Konopka, A. (2013). Quantifying community assembly processes and identifying features that impose them. *ISME Journal*, *7*(11), 2069–2079. https://doi.org/10.1038/ismej.2013.93

Tamura, K., & Nei, M. (1993). *Estimation of the Number of Nucleotide Substitutions in the Control Region of Mitochondrial DNA in Humans and*. *10*(3).

Torbati, M., Arzanlou, M., Abed-Ashtiani, F., & Golmohammadi, H. (2019). Occurrence of fruit rot on cornelian cherry caused by *Pilidium lythri* in Iran. *Crop Protection*, *125*, 104884. https://doi.org/10.1016/j.cropro.2019.104884

Trivedi, P., Leach, J. E., Tringe, S. G., Sa, T., & Singh, B. K. (2020). Plant–microbiome interactions: from community assembly to plant health. *Nature Reviews Microbiology*, *18*(11), 607–621. https://doi.org/10.1038/s41579-020-0412-1

Vishwakarma, K., Kumar, N., Shandilya, C., Mohapatra, S., Bhayana, S., & Varma, A. (2020). Revisiting Plant–Microbe Interactions and Microbial Consortia Application for Enhancing Sustainable Agriculture: A Review. *Frontiers in Microbiology*, *11*(December), 1–21. https://doi.org/10.3389/fmicb.2020.560406

Ward, J. R., Kim, K., & Harvell, C. D. (2007). Temperature affects coral disease resistance and pathogen growth. *Marine Ecology Progress Series*, *329*, 115–121. https://doi.org/10.3354/meps329115

White, T. J., Bruns, T., Lee, S., & Taylor, J. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR Protocols: A Guide to Methods and Applications*, 315–322.

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G., Hayes, A.,
Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K., Ooms, J., Robinson,
D., Seidel, D., Spinu, V., … Yutani, H. (2019). Welcome to the Tidyverse. *Journal of Open Source*
Software, *4*(43), 1686. https://doi.org/10.21105/joss.01686

Zhang, M., Li, H. L., Zhao, A. L., & Zhang, J. X. (2008). First Report of *Hainesia lythri* Causing Leaf Spots
of *Paeonia suffruticosa* in China. *Plant Disease*, *92*(3), 486–486. https://doi.org/10.1094/PDIS-92-
3-0486A

Acknowledgments

This research was funded in whole or in part by National Science Centre, Poland, contract number: 2022/45/N/NZ9/02089. For the purpose of Open Access, the author has applied a CC-BY public copyright license to any Author Accepted Manuscript (AAM) version arising from this submission. The DS's 3-month visit at the Institute of Environmental Sciences, Leiden University, Leiden, Netherlands was financed by the Polish National Agency for Academic Exchange, contract number: BPN/BEK/2023/1/00125/U/00001.

Competing interests

The authors have declared that no competing interests exist.

CREdiT (https://credit.niso.org/)

Conceptualization: DS, MF, JP, Data curation: DS, JP, Formal analysis: DS, Funding acquisition: DS, Investigation: DS, JP, Methodology: DS, JP, MF, EH, Project administration: DS, Resources: MF, DS, EH, Software: DS, JP, EH, Supervision: DS, Validation: DS, Visualization: DS, Writing – original draft: DS, Writing – review & editing: DS, EH, JP, MF

A) bulk soil

Pilidium
Wardomyces
Ilyonectria
Unclassified Chytridiomycota
Rhizophydiales_gen_Incertae_sedis
Unclassified Lecanoromycetes
Onygenales_gen_Incertae_sedis
GS16_gen_Incertae_sedis
Lobulomycetales_gen_Incertae_sedis
Pyronemataceae_gen_Incertae_sedis
Entoloma
Cladochytriales_gen_Incertae_sedis
Preussia
Ascomycota_gen_Incertae_sedis
Pleotrichocladium

Importance score

B) rhizosphere

Pilidium
Pleotrichocladium
Eocronartium
Sistotrema
Trichoderma
Paraphysoderma
Solicoccozyma
Unclassified Mortierellaceae
Ramicandelaber
Unclassified Lecanoromycetes
Niesslia
Verrucariales_gen_Incertae_sedis
Ophiosphaerella
Hypocreales_gen_Incertae_sedis
Aphelidiomycetes_gen_Incertae_sedis

Importance score

C) roots

Peziza
Pseudeurotium
Unclassified Fungi
Unclassified Pleosporales
Leucoagaricus
Mucor
Parascedosporium
Cladosporium
Mortierella
Sporobolomyces
Kernia
Fungi_gen_Incertae_sedis
Ascomycota_gen_Incertae_sedis
Phialophora
Metarhizium

Importance score