Customizing the Structure of a Minimal TIM Barrel to Craft a De Novo Enzyme

Julian Beck,[1,†] Benjamin J. Smith,[2,3,†] Niayesh Zarifi,[2,3] Emily Freund,[1] Roberto A. Chica,[2,3,*] Birte Höcker[1,*]

[1] Department of Biochemistry, University of Bayreuth, 95447 Bayreuth, Germany.

[2] Department of Chemistry and Biomolecular Sciences, University of Ottawa, Ottawa, Ontario, Canada, K1N 6N5

[3] Center for Catalysis Research and Innovation, University of Ottawa, Ottawa, Ontario, Canada, K1N 6N5

[†] These authors contributed equally.

* Corresponding authors

Birte Höcker, E-mail: birte.hoecker@uni-bayreuth.de

Roberto A. Chica, E-mail: rchica@uottawa.ca

Keywords

De novo enzyme design, computational protein design, enzymes, TIM barrel, Kemp elimination

Abstract

The TIM barrel is the most prevalent fold in natural enzymes, supporting efficient catalysis of

diverse chemical reactions. While *de novo* TIM barrels have been successfully designed, their

minimalistic architectures lack structural elements essential for substrate binding and catalysis.

Here, we present CANVAS, a computational workflow that introduces a structural lid into a

minimal *de novo* TIM barrel to anchor catalytic residues and form an active-site pocket for

enzymatic function. Starting from two *de novo* TIM barrels, we designed nine variants with distinct

lids to form active sites for the Kemp elimination. Experimental testing identified one active variant

with catalytic efficiency comparable to previously reported Kemp eliminases, and mutational

analyses validated the essential role of the designed catalytic residues. Sequence optimization of

this variant improved solubility and stability, enabling X-ray structure determination, which

confirmed the designed lid structure. This study reports the first enzymatically active *de novo* TIM

barrel and establishes a platform for designing enzymes from minimal protein scaffolds.

The TIM barrel is one of the most versatile enzyme folds. It is present in six of the seven

enzyme classes and supports catalysis at diffusion-limited rates[1-3]. While idealized *de novo* TIM

barrels have been successfully designed[4-7], none have been converted into functional enzymes

due to their minimalistic architecture, which lacks key structural elements required for catalysis,

such as cavities, pockets, and extended loops. Efforts to address these deficiencies have led to

the introduction of secondary structural elements to create rudimentary pockets[8-10]. However,

these pockets are too solvent exposed and undefined to support the microenvironments required

for catalysis. Therefore, novel strategies are needed to craft custom active sites and endow *de*

novo TIM barrels with enzymatic function.

A computational approach for active-site scaffolding

Here, we introduce CANVAS (Customizing Amino-acid Networks for Virtual Active-site

Scaffolding), a computational workflow for transforming minimal proteins such as *de novo* TIM

barrels into functional enzymes (Figure 1a, Supplementary Figure 1). The process begins with a

minimal TIM barrel template and a theozyme, which is a computational model of an idealized

enzyme active site with catalytic groups arranged to stabilize a reaction's transition state[11]. Using

the protein design software Triad[12], a catalytic amino acid from the theozyme is placed onto the

TIM barrel template, and the transition state is built from its side chain to maintain the catalytic

geometry[13]. A second catalytic residue is then built from the transition state, ensuring proper

geometry for catalytic interaction. The alpha carbon of this second residue is thus positioned in

the empty space above the TIM barrel catalytic face. This allows it to serve as an anchor for

constructing a custom lid to form the active-site pocket, which is created using the generative AI

tool RFdiffusion[14] followed by sequence design with ProteinMPNN[15] and evaluation with

AlphaFold2[16]. The active site is further optimized with Triad to build a pocket complementary to

the transition state[17]. Designs are then filtered using key enzyme design criteria[18], including

solvent-accessible surface area of the transition state, active-site preorganization, energy, and

catalytic contact geometry.

Applying CANVAS to minimal TIM-barrel proteins

We applied CANVAS to the Kemp elimination (Figure 1b), a model organic transformation

commonly used as a benchmark for *de novo* enzyme design[17-19]. The reaction's theozyme

features an Asp as catalytic base and an Asn or Gln as H-bond donor to stabilize negative charge

buildup on the oxygen at the transition state. We used the crystal structures of two *de novo* TIM

barrels (PDB IDs: 7MCD[20] and 7OSV[5]) as templates. Since 7OSV has its N- and C-termini on the

catalytic face of the TIM barrel, we applied an *in silico* circular permutation based on the inpainting

method[21] to generate two modified structures termed NT6-CP1 and NT6-CP2 that contain termini

on the stability face (Supplementary Figure 2). Starting from these structures, we used CANVAS

to generate a distinct lid for each template tailored to the Kemp elimination (Figure 1c). The lid

diffused onto 7MCD consists of a single 26-residue fragment introduced in βα loop 7

(Supplementary Figure 3). By contrast, the designed lids in NT6-CP1 and NT6-CP2 comprise a

major fragment of 33 or 39 residues introduced into βα loops 2 or 6, respectively, to scaffold the

catalytic H-bond donor, as well as three or four elongated loops to enhance interactions with the

primary fragment. The resulting designs, termed KempTIMs, include variants 1–3 obtained from

NT6-CP2, variants 4–5 from NT6-CP1 and variants 6–9 from 7MCD, each with a unique active-

site configuration (Supplementary Figure 4) and sequence (Supplementary Table 1), which

yielded high AlphaFold2 pLDDT scores (Supplementary Figure 5).

Kinetic assays reveal an active Kemp eliminase

Of the nine designs, seven expressed in soluble form in *E. coli* but only five could be

purified to homogeneity (Supplementary Table 2). Notably, the NT6-CP1 template could not be

expressed, in contrast to NT6-CP2 and 7MCD. Enzymatic assays revealed a single active design,

KempTIM4, which followed Michaelis-Menten kinetics with substrate saturation (Figure 2a),

yielding k_{cat}, K_M, and k_{cat}/K_M values of 0.0066 ± 0.0006 s^{-1}, 0.6 ± 0.1 mM, and 11 ± 3 M^{-1} s^{-1},

respectively. This catalytic efficiency is comparable to those of other *de novo* Kemp

eliminases[17,19,22], such as KE07 (k_{cat} = 0.018 s^{-1}, K_M = 1.4 mM, and k_{cat}/K_M = 12.2 M^{-1} s^{-1})[19].

KempTIM4 expressed predominantly as a monomeric protein (Figure 3a,b, Supplementary Table

3) and displayed a mixed αβ secondary structure characteristic of TIM barrels (Figure 3c).

Compared to the minimal TIM barrel NT6-CP2, KempTIM4 exhibited increased random coil

content, indicated by a stronger circular dichroism (CD) signal at 208 nm. This structural feature

aligns with the design of its lid, which includes long loops (Figure 3a, Supplementary Figure 3).

KempTIM4 showed a strongly reduced melting temperature relative to NT6-CP2 (Figure 3d),

reflecting the destabilizing effects of incorporating a lid and active site onto the minimal TIM barrel

scaffold. Mutation of the designed catalytic base and H-bond donor to Ala abolished or

substantially reduced activity (Figure 2b) despite size-exclusion chromatograms and CD spectra

comparable to KempTIM4 (Supplementary Figure 6, Supplementary Table 3). These results

confirm that the loss of activity in these knockout variants is due to mutation, not structural

disruption. Moreover, the absence of enzymatic activity in NT6-CP2 demonstrates that catalysis

in KempTIM4 is driven by the designed lid and active site.

Redesign yields a more stable enzyme

Although active, KempTIM4 was prone to aggregation at high concentrations. This led us

to redesign its sequence for enhanced expression and solubility using ProteinMPNN, which has

been shown to produce stable and soluble proteins[23]. In this procedure, identities of active-site

residues designed with Triad were preserved to maintain catalytic function, while the rest of the

protein was redesigned. The redesigned protein, KempTIM4-2, demonstrated approximately 10-

fold higher expression yields compared to KempTIM4 (Supplementary Table 2) and existed

exclusively as a monomer with no tendency to form dimers (Figure 3b, Supplementary Table 3).

CD spectroscopy showed increased α-helicity compared to KempTIM4 and higher stability,

comparable to NT6-CP2 (Figure 3c,d). Despite these improvements, KempTIM4-2 exhibited lower

activity than KempTIM4 (k_{cat}/K_M = 7.0 ± 0.1 M^{-1} s^{-1}) and could not be saturated within the

substrate's solubility limit (Figure 2a), even though active-site residues remained unchanged. As

with KempTIM4, mutation of the catalytic base in KempTIM4-2 abolished activity (Figure 2b)

without disrupting structure (Supplementary Figure 6, Supplementary Table 3). However,

mutation of the H-bond donor caused only a small reduction in catalytic activity compared to

KempTIM4, suggesting that this designed Asn residue adopts a suboptimal orientation hindering

its catalytic function. Nevertheless, inhibition assays using the 6-nitrobenzotriazole (6NBT)

transition-state analogue demonstrated competitive binding in the active site (Figure 2c),

confirming the integrity of the designed active site pocket.

The crystal structure confirms the designed lid and provides active site details

Crystals of KempTIM4-2 were obtained in the presence of 6NBT, and the structure was

solved at 2.3 Å resolution (Supplementary Table 4). The structure confirmed the correct folding

of the designed lid, with the primary helix-turn-helix fragment and four elongated loops closely

matching the design model (Figure 4a). However, the lid was slightly displaced relative to the TIM

barrel core, resulting in a 1.8-Å shift of the catalytic H-bond donor N46 Cα atom (Figure 4b), which

likely reduces its contribution to catalysis (Figure 2b). While crystal contacts may contribute to this

displacement, size-exclusion chromatography coupled with small-angle X-ray scattering (SEC-

SAXS) confirmed that the designed lid adopts a similar conformation in solution (Supplementary

Figure 7).

Despite crystallization with 6NBT, no electron density corresponding to the transition-state

analogue was observed in the active site; instead, the density was best interpreted as a bound

glycerol molecule from the cryoprotectant. This suggests weak binding of the transition-state

analogue under crystallization conditions (Methods). Analysis of the active site revealed that the

catalytic base D153 adopts a similar rotameric configuration in both the crystal structure and

design model, whereas N46 adopts a rotameric state incompatible with transition-state

stabilization (Figure 4b). This aligns with the observation that mutation of N46 to Ala in KempTIM4-

2 causes only a small reduction in activity, while mutation of D153 abolishes activity (Figure 2b).

Additionally, the active site is poorly preorganized, with several amino acid side chains deviating

from the conformations designed to stabilize the transition state. These structural discrepancies

likely contribute to the modest catalytic efficiency of KempTIM4-2 and provide starting points for

future optimization.

Structural Dynamics and Active-Site Accessibility Underpin Catalytic Activity

To explore why only KempTIM4 and KempTIM4-2 were catalytically active, microsecond-

timescale molecular dynamics simulations were performed on these designs and the inactive

KempTIM9, both in the presence and absence of the transition-state analogue. KempTIM9 was

selected as a comparison due to its good expression (Supplementary Table 2), well-folded

structure and high melting temperature (67°C, Supplementary Figure 8), which rule out misfolding

as the cause of inactivity. The simulations revealed that the helical component of the lids, which

anchor the catalytic Asn, were rigid (Figure 5a). This rigidity stabilizes the transition-state

analogue within the binding pocket, maintaining catalytic interactions with the catalytic base for

much of the simulation, with active variants outperforming the inactive design KempTIM9 (Figure

5b). However, only KempTIM4 consistently maintained catalytic interactions with the H-bond

donor. Both KempTIM4-2 and KempTIM9 showed weaker contacts with this Asn residue, with

KempTIM4-2 failing to form this interaction throughout the simulation. This result aligns with the

lid displacement and unproductive rotameric state of N46 observed in the crystal structure, as

well as the small impact on activity when N46 was mutated to Ala in KempTIM4-2 (Figure 2d).

A key difference between the active designs and the inactive KempTIM9 was the

significantly narrower entrance to the active site in KempTIM9 (Figure 5b). Previous studies have

demonstrated that widening the active-site entrance enhances Kemp eliminase activity by

lowering the energy barriers associated with substrate entry and product release[24]. Thus, we

postulate that the inactivity of KempTIM9 is due to an insufficiently open active site, which likely

impedes substrate binding. By contrast, the lids of KempTIM4 and KempTIM4-2, which

incorporate five fragments rather than the single fragment found in KempTIM9 (Supplementary

Figure 3), may facilitate substrate entry. The additional inserted loops in these designs likely

contribute to active-site accessibility by providing greater flexibility to the lid in the unbound state

(Figure 5a), allowing the active site to open more readily. While other factors, such as incorrect

positioning of catalytic residues and lid misorientation, cannot be excluded, these findings suggest

that reducing barriers to substrate entry during lid generation could improve catalytic performance

in future designs.

Opportunities for CANVAS design

Natural TIM barrels use lids to bind substrates and form an active site conducive to

catalysis[25]. By contrast, minimal *de novo* TIM barrels lack such lids and are inherently

functionless. Using our CANVAS approach, we successfully designed a custom lid to form an

active site for a target reaction, resulting in the first enzyme constructed from a *de novo* TIM

barrel. While KempTIM4 exhibits modest activity, it matches the performance of many first-round

designs generated by traditional methods[17,19], despite the added complexity of engineering a

tailored lid comprising multiple loops to create an active site cavity. Further activity improvements

could be achieved by using the crystal structure reported here as a template for ensemble-based

design[24], a method that leverages crystallographic ensembles to design highly preorganized

active sites, leading to catalytic efficiency enhancements of several orders of magnitude[18]. Unlike

traditional enzyme design methods that rely on pre-existing scaffolds, CANVAS offers the

flexibility to customize lids for specific reaction requirements. It can also be readily adapted to

scaffold more than two catalytic residues, a task that becomes increasingly challenging with pre-

existing scaffolds as the number of catalytic groups grows. With CANVAS, we anticipate that any

minimal TIM barrel, including AI-generated virtual ones, could serve as a blank canvas for the

creation of a large diversity of *de novo* enzymes, unlocking the full potential of this versatile fold

for catalysis of a wide range of reactions.

Methods

Reagents and solutions. All experiments were conducted using analytical-grade chemicals, with

solutions prepared using double-distilled water.

Circular permutation of DeNovoTIM6-SB with inpainting. Circular permutation of

DeNovoTIM6-SB (PDB ID: 7OSV) was performed using the inpainting method described by Wang

et al.[21] The first residue of either the first or third β-strand was selected as the new N-terminus,

with the corresponding βα-loop deleted (Supplementary Figure 2). A new connection for the

corresponding βα-loop was modeled by inpainting 5 to 10 residues. Unresolved residues in the

crystal structure were remodeled, keeping intact the positions and identities of the remaining

residues. For each new connection, 100 designs were modeled, followed by relaxation and

scoring using the Rosetta protein design software[26]. Structures of each design were predicted

using AlphaFold2[16] with model 4 weights. Designs with a pLDDT value above 90 and a Rosetta

score below 3 Rosetta energy units per residue were kept. One design from each connection was

selected for experimental characterization (Supplementary Table 1). The proteins were named

NT6-CP1 (N-terminus on the first β-strand) and NT6-CP2 (N-terminus on the third β-strand).

Computational Enzyme Design. All calculations were performed with the Triad protein design

software (Protabit, Pasadena, CA, USA). Rotamer optimization was performed using a Monte

Carlo with simulated annealing search algorithm. Input structures were prepared for Triad

calculations via the *addH.py* application within Triad. The Kemp elimination transition state (TS)

structure was built using parameters from Privett *et al.*[17]. To provide sidechain conformations, a

backbone-independent conformer library (bda-bbind_1.0)[13] was used for theozyme placement,

and a backbone-independent rotamer library (bbind02.May.e2)[27] with expansions of ±1 standard

deviation around χ_1 and χ_2 was used for active-site repacking. Energies were calculated using a

modified version of the Phoenix energy function[12] consisting of a Lennard-Jones 12–6 van der

Waals term from the Dreiding II force field[28] with atomic radii scaled by 0.9, a direction-dependent

hydrogen bond term with a well depth of 8.0 kcal mol^{-1} and an equilibrium donor–acceptor

distance of 2.8 Å,[29] and an electrostatic energy term modeled using Coulomb's law with a

distance-dependent dielectric of 10. An energy benefit of −100 kcal mol^{-1} was applied when TS-

side-chain interactions satisfied catalytic contact geometries (Supplementary Table 5), as

described by Lassila *et al.*[13]

Theozyme placement. AlphaFold2 predictions of NT6-CP1 and NT6-CP2, along with the 7MCD

crystal structure, were used as backbone templates for theozyme placement. Amino-acid

residues with a C_α–C_β vector oriented towards the β-barrel interior and located at the C-terminus

of each β-strand were selected as positions for introduction of the catalytic base (Asp).

Neighbouring residues were mutated to alanine to avoid steric clashes with the TS

(Supplementary Table 6). TS poses were built using the contact geometries listed in

Supplementary Table 7, and those where the isoxazolic oxygen pointed toward the catalytic face

of the TIM barrel and remained coplanar with the catalytic base were selected (Supplementary

Figure 1). In parallel, the catalytic H-bond donor (Asn or Gln) was placed on a Gly-Asn/Gln-Gly

tripeptide scaffold, and TS poses were built using H-bond donor side-chain contacts specified in

Supplementary Table 7. The resulting TS was superimposed with the TS from the catalytic base

calculation to create the full theozyme, positioning the catalytic base within the minimal TIM barrel

and the H-bond donor in the empty space above the catalytic face. The Gly residues from the

tripeptide were then deleted, preserving only the Asn/Gln residue, whose alpha carbon atom

served as an anchor for the designed lid.

Lid design. For all theozyme placements, peptide connections between the catalytic H-bond

donor residue and the minimal TIM barrel structures were designed using RFdiffusion[14]. For each

position of the H-bond donor alpha carbon, various βα-loops were selected as insertion points for

a primary peptide fragment, and multiple fragment lengths were sampled to connect to the

catalytic residue while preserving the TIM-barrel topology. Additional βα-loops were sometimes

elongated to improve interactions with the primary fragment (Supplementary Figure 3). For each

inserted primary fragment, 100–200 structures were sampled. Primary fragments lacking

secondary structure, causing steric clashes with the TS, or displacing the H-bond donor from its

theozyme position, were discarded (Supplementary Figure 1). The filtered fragments were

sequence-optimized using ProteinMPNN[15] using a temperature factor of 0.1, ensuring the minimal

TIM barrel template sequence remained unchanged. Cysteine and methionine residues were

excluded during sequence design. Structures of designed sequences were predicted using

AlphaFold2[16] or ColabFold v1.3.0[30] with all five model weights, and those displaying an average

pLDDT value >90 and backbone RMSD < 3 Å relative to the RFdiffusion model were selected for

active-site repacking.

Active-site repacking. AlphaFold2 models generated above were used for a theozyme

placement step, as described previously, to verify that contacts between catalytic residues and

the TS could be formed on these scaffolds. Active-site repacking calculations were then

performed on the structures with this new theozyme placement. In these calculations, the TS was

translated by ±0.4 Å along each Cartesian coordinate in 0.2-Å steps and rotated about all three

axes (origin at the TS geometric center) in 5° increments over a 10° range (clockwise and

counterclockwise). This resulted in a total combinatorial search space of 15,625 possible poses.

Residues near the catalytic amino acids and TS were designated as design positions. During

repacking, these positions were allowed to sample rotamers of various amino acids, favoring

hydrophobic and aromatic residues, while the identities of the catalytic residues remained fixed

(Supplementary Table 8).

Computational library design. After repacking, the CLEARSS computational library design

algorithm[31] was used to generate a combinatorial library comprising the most favorable amino

acids predicted by Triad at each designed active-site position. In this method, libraries of a specific

size configuration are generated from a pre-scored list of sequences using the highest probability

set of amino acids at each position based on the sum of their Boltzmann weights. Using the list

of energy-ranked sequences from active-site repacking as input, libraries of 192 sequences were

generated. Rotamer configurations for each sequence in the library were optimized using the

cleanSequences.py application within Triad to find the lowest-energy conformation of each

sequence on its respective backbone, generating "cleaned" structures. To compare energies, the

energy difference between each "cleaned" structure and a corresponding all-Gly structure in the

absence of the TS was calculated. These energies are reported in Supplementary Table 9.

Structures were cleaned with and without the TS to evaluate preorganization, as described

previously[18].

Design filtering. Energies of the TS-bound and unbound "cleaned" structures were calculated

using Triad. Dihedral angles of catalytic residues relative to the TS, as well as the solvent-

accessible surface area (SASA) of the TS, were analyzed using PyMOL (v2.3.0, Schrödinger,

LLC). Bottleneck radii of active-site entrances were evaluated using CAVER v3.0[32], and the

number of residues mutated to glycine or methionine was counted. Designs were filtered based

on the following criteria (Supplementary Table 9): energy difference between bound and unbound

structures below 0 kcal mol^{-1}, catalytic residue dihedral angles outside 50° to 130° and –50° to –

130°, SASA of TS between 80 and 200 Å², < 6 Met and < 4 Gly, bottleneck radius of active-site

entrance > 0.9 Å, and < 7 non-preorganized residues. Designs passing all criteria were used for

structure prediction with AlphaFold2 or ColabFold (v1.3.0), where designs with a backbone RMSD

< 3 Å relative to the structure cleaned by Triad, overall pLDDT > 90, and lid pLDDT > 85 were

chosen for experimental characterization (Supplementary Table 1).

KempTIM4 redesign. Sequence optimization of KempTIM4 was performed using ProteinMPNN

with LigandMPNN weights[33]. Amino-acid identities of all residues designed during active-site

repacking of KempTIM4 were retained (Supplementary Table 8). Four sequences were

generated, omitting cysteine. The generated sequences were used for structure prediction with

ColabFold v1.3.0[30] using all five model weights. Predictions were filtered based on their all-atom

RMSD to the input KempTIM4 AlphaFold2 model. Theozyme placement and active-site repacking

for each sequence were performed as described earlier using the backbones of AlphaFold2

models that passed filtering. Following energy calculations with Triad, a single design,

KempTIM4-2, was selected for experimental characterization (Supplementary Table 1).

Cloning and generation of constructs. Genes for all proteins were ordered as codon-optimized

fragments flanked by restriction sites for *Nde*I and *Xho*I from Twist Bioscience (South San

Francisco, CA, USA). After digestion with *Nde*I and *Xho*I, the gene fragment was ligated into

pET21b(+) for KempTIM1–5, or pET29b(+) for KempTIM6–9. Catalytic knockout mutations were

introduced by a modified QuikChange PCR protocol utilizing KAPA Polymerase followed by *Dpn*I

digestion. *E. coli* Top10 or BL21 (DE3) cells were transformed with ligated vector or reaction

mixture and plated on Lysogeny Broth (LB) agar plates containing 100 μg mL^{-1} ampicillin or 50

μg mL^{-1} kanamycin as selection markers. Single colonies were used to inoculate 5 mL LB

supplemented with 100 μg mL^{-1} ampicillin or 50 μg mL^{-1} kanamycin. After overnight growth

(37 °C, 250 rpm), cells were harvested by centrifugation and DNA was isolated using NucleoSpin

Plasmid EasyPure-Kit (Machery & Nagel) according to the manufacturer's protocol. Vector

assembly and introduction of mutations were validated by sequencing (Eurofins Genomics) using

standard T7 primers.

Protein expression and purification. *E.coli* BL21 (DE3) cells (Novagen) were transformed with

plasmid, plated on agar plates containing antibiotic, and incubated overnight at 37 °C. Single

colonies were picked to inoculate precultures and incubated at 30–37 °C overnight. On the next

day, 1 L LB supplemented with antibiotic was inoculated with 10 mL of the preculture and

incubated at 37 °C until OD600 reached a value between 0.6 and 0.8. Overexpression was

induced by adding isopropyl-β-thiogalactoside (IPTG) to a final concentration of 0.1 mM for

KempTIM1–5 or 1 mM for KempTIM6–9 followed by overnight incubation at 16–20 °C with

shaking. Cells were harvested by centrifugation and pellets were either frozen at –20 °C or used

directly for purification.

Two different purification schemes were employed. In the first scheme, cell pellets were

resuspended in 35 mL buffer A (50 mM sodium phosphate pH 7.0, 100 mM NaCl, 10 mM

imidazole). Resuspended cells were lysed by sonication (Branson Ultrasonic Sonifier 250, output

4, duty cycle 40%, 3 × 3 min) and centrifuged. The supernatant was loaded onto a HisTrapHP

column (5 mL, Cytiva Life Science) equilibrated with buffer A and coupled to an ÄKTApure system

(Cytiva Life Science). After washing with 10 column volumes (CV) of buffer A, the protein was

eluted with a linear gradient over 20 CV to 60 % buffer B (50 mM sodium phosphate pH 7.0, 100

mM NaCl, 500 mM imidazole). For KempTIM4-2L, which contains an additional GSG tripeptide

linker between its His-tag and first TIM barrel residue (Supplementary Table 1), an additional

TEV-cleavage over night was performed during a dialysis against buffer C (50 mM of sodium

phosphate pH 7.0, 100 mM NaCl). The dialyzed protein was loaded onto a HisTrapHP column

(5 mL, Cytiva Life Science) equilibrated with buffer C and coupled to an ÄKTApure system (Cytiva

Life Science) and the flowthrough was collected. Fractions containing the protein were pooled,

concentrated with a centrifugal concentrator, and loaded onto a HiLoad 26/600 Superdex 75

preparative grade column (Cytiva Life Sciences) preequilibrated in buffer C and eluted with 1 CV

buffer C.

In the second scheme, cell pellets were resuspended in 8 mL buffer D (5 mM imidazole in

100 mM potassium phosphate buffer at pH 8.0) supplemented with 1 mg mL^{-1} lyophilized

lysozyme (MP Biomedicals) and 1 U mL^{-1} benzonase nuclease (Merck Millipore). Cells were

homogenized using an Avestin EmulsiFlex-B15 cell disruptor. Proteins were purified from the

lysate by immobilized metal affinity chromatography using Ni-NTA agarose (Qiagen) pre-

equilibrated with buffer D in individual Econo-Pac gravity-flow columns (Bio-Rad). Contaminants

were washed away using buffer E (10 mM imidazole in 100 mM potassium phosphate buffer pH

8.0) and then 20 mM imidazole in the same buffer. Proteins were eluted with 5 mL of buffer F

(250 mM imidazole in 100 mM potassium phosphate buffer pH 8.0). Proteins were further

subjected to gel filtration in buffer G (25 mM HEPES pH 7.25 and 100 mM sodium chloride) using

an ENrich SEC 650 size-exclusion chromatography column (Bio-Rad). Purification was checked

by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and fractions

containing the protein of interest were pooled. Protein concentration was determined

spectrophotometrically using the absorption at 280 nm and applying Beer-Lambert's law using

calculated extinction coefficients obtained from the ExPASy ProtParam tool

(https://web.expasy.org/protparam/).

Far-UV circular dichroism (CD) spectroscopy. CD spectra were collected with a Jasco J-710

or J-815 spectrometer. Spectra were recorded using a 1-mm quartz cuvette at 20 °C with

scanning speed 10–100 nm min^{-1}, bandwidth 1 nm, response time 1 s. Experiments were

performed in 20 mM sodium phosphate buffer pH 7.0 or 50 mM sodium phosphate buffer pH 8.0

supplemented with 100 mM sodium chloride, using a protein concentration of approximately

0.2 mg mL^{-1}. For each protein, 3 to 10 spectra were accumulated and averaged. For data

normalization, a buffer spectrum was subtracted and the signal was converted to mean residue

molar ellipticity using $[\theta_{MRE}] = \theta/(n \times d \times c)$, where n is the number of residues in the protein, θ the

collected ellipticity in mdeg, d the path length in mm, and c the protein concentration in M. Thermal

denaturation assays were performed by heating the samples from 20 to 95 °C at a rate of 1 °C

per minute and ellipticity at 222 nm was measured every 1 °C. Melting temperature (T_m) was

determined following the protocol by Greenfield[34].

Size-exclusion chromatography-multi angle light scattering. SEC-MALS measurements

were performed as described in Beck *et al*.[8] Each protein was measured at 2 mg mL^{-1} except

KempTIM4 N46A (1 mg mL^{-1}). For all measurements, buffer C and an injection volume of 50 µL

were used.

Steady-state kinetics. All assays were performed in buffer C or G at 27 °C in a Spark (Tecan)

or Synergy H1 (Biotek) plate reader. Enzyme concentrations varied from 5 to 80 µM. Reactions

(200 µL final volume) were initiated by addition of varying concentrations (0.03–2 mM) of

5-nitrobenzisoxazole (abcr or AAblocks) dissolved in methanol (final methanol concentration

10 %). Product formation was monitored at 380 nm (ε = 15,800 M^{-1} cm^{-1})[17] in individual wells of a

96-well plate (Nunc or Greiner Bio-One). Path lengths for each well were calculated ratiometrically

using the difference in absorbance at 900 and 975 nm. Control measurements without protein

were conducted for each substrate concentration and subtracted from the enzyme

measurements. Linear phases of the kinetic traces were used to measure initial reaction rates. If

the enzyme showed saturation within the substrate concentration used, the data were fitted to the

Michealis-Menten model using ($v_0 = (v_{max} \times [S]) / (K_M + [S])$ where v_0 is the initial velocity rate, v_{max}

is the maximal velocity rate, [S] is the substrate concentration, K_M is the Michaelis constant. The

catalytic constant k_{cat} was calculated with $k_{cat} = v_{max} / [E]$ where [E] is the enzyme concentration.

If saturation was not achieved within the substrate's solubility limit, a linear equation ($v_0 = k_{cat}/K_M$

× [S]) was used.

Inhibition assays. Inhibition assays of KempTIM4-2 were performed in 25 mM HEPES buffer

(pH 7.25) with 100 mM NaCl and 1 % DMSO at 27 °C using a Biotek Synergy H1 multimode plate

reader. Reactions (200 µL final volume) were initiated by addition varying concentrations (0.25–

2 mM) of 5-nitrobenzisoxazole (AAblocks) dissolved in methanol (10 % final concentration) to the

enzyme mixed with varying concentrations (0.5–1 mM) of 6-nitro-1H-benzotriazole (Aaron

Chemicals) in wells of a 96-well plate (Greiner Bio-One). The formation of the product was

monitored spectrophotometrically at 380 nm (ε = 15,800 M^{-1} cm^{-1}). Path lengths for each well

were calculated ratiometrically using the difference in absorbance at 900 and 975 nm. Linear

phases of kinetic traces were used to measure initial reaction rates. Michaelis-Menten and

Lineweaver-Burk plots were created from the resulting data to determine the type of inhibition.

Crystallization and structure determination. KempTIM4-2L (Supplementary Table 1) was

purified using the first purification scheme, with the final size-exclusion chromatography step

performed in 20 mM HEPES buffer pH 8.0 supplemented with 20 mM NaCl. Crystallization

screens were set up using the sitting-drop vapor diffusion method with a Phoenix pipetting robot

(Art Robbins Instruments) and commercially available sparse-matrix screens (NeXtal) in 96-well

sitting-drop plates (3-drop Intelli-Plates, Art Robbins Instruments). Protein and reservoir solutions

were mixed in ratios of 1:1, 2:1, and 1:2. The protein concentration was 12 mg mL^{-1},

supplemented with 5 mM 6NBT (5% DMSO). Crystals were grown at 293 K, and diffracting

crystals were obtained after 40 days in 4 M sodium formate. Cryoprotection was achieved by

adding glycerol to a final concentration of 25%, and crystals were mounted using cryo-loops on

SPINE standard bases and flash-cooled in liquid nitrogen. Diffraction data were collected at 100

K at beamline BL 14.1 with a Pilatus3 S 6M detector at the BESSY II synchrotron (Helmholtz-

Zentrum Berlin). Data processing was performed using X-ray Detector Software APP3

(XDSAPP3)[35] with XDS[36], and data quality was assessed using phenix.xtriage[37]. Phases were

solved by molecular replacement using the AlphaFold2 prediction as search model with Phaser[38].

The resulting model was manually rebuilt using Coot[39] and refined with phenix.refine[37] in an

iterative process. Refinement statistics and crystallographic data are shown in Supplementary

Table 4. Coordinates and structure factors were validated and deposited in the PDB under

accession code 9QKX.

Size-Exclusion Chromatography Small-angle X-ray Scattering (SEC-SAXS). SEC-SAXS

measurements were conducted at the BioSAXS beamline BM29 of the ESRF in Grenoble,

France. A Superdex 75 Increase 10/300 GL column (Cytiva Life Sciences) was used with a flow

rate of 0.8 mL min^{-1} in buffer C. KempTIM4-2L (Supplementary Table 1) was measured at a

concentration of 5 mg mL^{-1}, with an injection volume of 100 μL. Data processing and analysis

were performed using ATSAS 3.2.1[40,41]. For analysis, AlphaFold2 models of KempTIM4-2 and

NT6-CP2, as well as an AlphaFlow[42] model of KempTIM4-2 with a displaced lid obtained using

the base molecular dynamics weights were used. For each of these models a theoretical

scattering curve were generated and fitted to the experimental data using CRYSOL[43].

Molecular dynamics (MD) simulations. The Amber 2020 software (http://ambermd.org/) with

the ff19SB protein force field[44], gaff2 ligand force field[45], and OPC water force field[46] was used for

all simulations. A cutoff of 10 Å was applied to electrostatics modeled using the particle mesh

Ewald method[47]. All MD trajectories were run with a time step of 2 fs. Models of each design

generated using Triad as described above were prepared for MD. MD trajectories were run for

each enzyme in the presence and absence of bound transition-state analogue 6-

nitrobenzotriazole (6NBT). Parameters for 6NBT were generated using the Antechamber

package[48]. Hydrogen atoms were added using Reduce[49], and the prepared structures were

solvated with OPC water in a truncated octahedral box with periodic boundary conditions where

the distance between the protein surface and the box edges was set to 10 Å. The addions2

algorithm in Amber was used to place counterions to neutralize the system. The system was first

minimized through the method of steepest descent using a force constant of 500 kcal mol^{-1} Å^{-2}.

The system was then heated to a temperature of 300 K over 240 ps while restraints were gradually

removed, followed by equilibration under an NPT ensemble at 300 K and 1 bar for 10 ns and

subsequent equilibration under an NVT ensemble at 300 K for another 10 ns. Constant

temperature and pressure were achieved using the Langevin thermostat[50] and Berendsen

barostat[51], respectively. Following equilibration, triplicate 1-μs production simulations were run.

For analyzing bottleneck radii, 1000 snapshots separated by 1 ns each were extracted from the

production trajectory. A minimum cutoff of 0.9 Å was used to identify active-site entrance

bottlenecks in the unbound MD trajectories using CAVER 3.0[32]. To analyze catalytic interactions

and RMSF, 5×10^4 snapshots separated by 20 ps each were extracted from the production

trajectory. Catalytic interactions and RMSF values were extracted using CPPTRAJ and PYTRAJ,

respectively[52].

Figures

464
465
Figure 1. *De novo* enzyme design using CANVAS. (a) CANVAS follows three general steps: (1) placement of the
theozyme (blue) onto the minimal TIM-barrel scaffold using a single catalytic residue as an anchor; (2) construction of
a custom lid (magenta) to form the active-site pocket and anchor the second catalytic residue; and (3) active-site
repacking to further stabilize the transition state. (b) Kemp elimination reaction. (c) Structure of three distinct lids
(magenta) tailored to the Kemp elimination, designed using CANVAS. Transition state and designed catalytic residues
are shown as sticks.

472
473
Figure 2. Kinetic characterization of KempTIMs. (a) Michaelis–Menten plots of normalized initial rates as a function
of 5-nitrobenzisoxazole (5-NBZ) concentrations. Data represent the average of 18 or nine replicate measurements from
five or four independent protein batches for KempTIM4 and KempTIM4-2, respectively (mean ± SEM). (b) Reaction
progress curves ([5NBZ] = 2 mM) demonstrate the contributions of designed catalytic residues D153 and N46 to
catalysis. (c) Michaelis-Menten (left) and Lineweaver-Burk (right) plots reveal that the transition-state analogue 6NBT
acts as a competitive inhibitor of KempTIM4-2. Data represent the average of four replicate measurements from a
single protein batch (mean ± SEM).
481

482
483
484 **Figure 3. Structural characterization.** (a) Computational models of minimal TIM-barrel NT6-CP2 and active
485 KempTIMs with the designed lid shown in magenta and the theozyme as sticks. (b) SEC-MALS indicates that the
486 proteins are predominantly monomeric, with molecular weights close to their expected values. (c) CD spectra reveal a
487 mixed αβ signal characteristic of TIM barrels. MRE: mean residue ellipticity. (d) Melting curves demonstrate that
488 KempTIM4 is substantially destabilized compared to NT6-CP2. Redesign to yield KempTIM4-2 restores stability.
489

490
491
Figure 4. Crystal structure of KempTIM4-2. (a) The crystal structure (blue) aligns closely with the design model
(minimal TIM barrel and lid colored white and magenta, respectively). The backbone RMSD is 0.86 Å for the full
structure and 1.34 Å for the lid. (b) Active-site residues in the crystal structure (blue) and design model (white and
magenta) show greater divergence, possibly due to the absence of a bound transition-state analogue in the crystal
structure. While the catalytic base D153 adopts a similar rotamer in both structures (side chain RMSD = 0.76 Å), the
catalytic H-bond donor N46 adopts a rotameric configuration incompatible with transition-state (TS) stabilization (side
chain RMSD = 3.09 Å).
499

500
501
Figure 5. Molecular dynamics in the presence and absence of transition-state analogue 6NBT. (a) Root-mean-
square fluctuations (RMSF) of the protein backbone, represented as putty plots, reveal that the lid α-helix anchoring
the catalytic H-bond donor (boxed) is rigid, whereas loops exhibit greater flexibility. 6NBT and catalytic residues are
depicted as sticks. (b) Distributions of catalytic contact distances between 6NBT and catalytic base or H-bond donor
are shown alongside the active-site entrance bottleneck radius, presented as violin plots, with the median value
indicated by a black line.
508

Acknowledgments

J.B. and B.H. acknowledge support from the Elite Network of Bavaria and its study program

"Biological Physics". B.H. acknowledges support from European Union's Horizon 2020 research

and innovation program under grant agreement No 951375 (ArtMotor). R.A.C. acknowledges

grants from the Natural Sciences and Engineering Research Council of Canada (RGPIN-2021-

03484 and RGPAS-2021-00017) and the Canada Foundation for Innovation (26503). B.J.S. was

supported by an NSERC CREATE scholarship. This research was enabled in part by support

provided by Compute Ontario (www.computeontario.ca) and the Digital Research Alliance of

Canada (alliancecan.ca). We acknowledge financial support and allocation of synchrotron

beamtime by HZB and thank the beamline staff at BESSY for support. We thank Sabrina Wischt

for technical support, Janosch Hennig for help with SAXS data analysis and the University of

Bayreuth Centre of International Excellence "Alexander von Humboldt" for facilitating this

collaboration through a short-term grant to R.A.C.

Competing interests

The authors declare no competing financial interest.

ORCID identifiers

Julian Beck: 0009-0007-3555-9890

Roberto A. Chica: 0000-0003-3789-9841

Emily Freund: 0009-0006-2876-3319

Birte Höcker: 0000-0002-8250-9462

Benjamin J. Smith: 0009-0003-5953-4998

Niayesh Zarifi: 0000-0003-4748-7082

References

Nagano, N., Orengo, C. A. & Thornton, J. M. One Fold with Many Functions: The Evolutionary Relationships between TIM Barrel Families Based on their Sequences, Structures and Functions. *Journal of Molecular Biology* **321**, 741-765, doi:https://doi.org/10.1016/S0022-2836(02)00649-6 (2002).

Höcker, B., Jürgens, C., Wilmanns, M. & Sterner, R. Stability, catalytic versatility and evolution of the (βα)8-barrel fold. *Current Opinion in Biotechnology* **12**, 376-381, doi:https://doi.org/10.1016/S0958-1669(00)00230-5 (2001).

Wierenga, R. K., Kapetaniou, E. G. & Venkatesan, R. Triosephosphate isomerase: a highly evolved biocatalyst. *Cellular and Molecular Life Sciences* **67**, 3961-3982, doi:10.1007/s00018-010-0473-9 (2010).

Huang, P.-S. *et al.* De novo design of a four-fold symmetric TIM-barrel protein with atomic-level accuracy. *Nature Chemical Biology* **12**, 29-34, doi:10.1038/nchembio.1966 (2016).

Kordes, S., Romero-Romero, S., Lutz, L. & Höcker, B. A newly introduced salt bridge cluster improves structural and biophysical properties of de novo TIM barrels. *Protein Science* **31**, 513-527, doi:https://doi.org/10.1002/pro.4249 (2022).

Romero-Romero, S. *et al.* The Stability Landscape of de novo TIM Barrels Explored by a Modular Design Approach. *Journal of Molecular Biology* **433**, 167153, doi:https://doi.org/10.1016/j.jmb.2021.167153 (2021).

Chu, A. E., Fernandez, D., Liu, J., Eguchi, R. R. & Huang, P.-S. De Novo Design of a Highly Stable Ovoid TIM Barrel: Unlocking Pocket Shape towards Functional Design. *BioDesign Research* **2022**, 9842315, doi:https://doi.org/10.34133/2022/9842315 (2022).

Beck, J., Shanmugaratnam, S. & Höcker, B. Diversifying de novo TIM barrels by hallucination. *Protein Science* **33**, e5001, doi:https://doi.org/10.1002/pro.5001 (2024).

Kordes, S., Beck, J., Shanmugaratnam, S., Flecks, M. & Höcker, B. Physics-based approach to extend a de novo TIM barrel with rationally designed helix-loop-helix motifs. *Protein Engineering, Design and Selection* **36**, gzad012, doi:10.1093/protein/gzad012 (2023).

Wiese, J. G., Shanmugaratnam, S. & Höcker, B. Extension of a de novo TIM barrel with a rationally designed secondary structure element. *Protein Science* **30**, 982-989, doi:https://doi.org/10.1002/pro.4064 (2021).

Tantillo, D. J., Jiangang, C. & Houk, K. N. Theozymes and compuzymes: theoretical models for biological catalysis. *Current Opinion in Chemical Biology* **2**, 743-750, doi:https://doi.org/10.1016/S1367-5931(98)80112-9 (1998).

Lee, F. S., Anderson, A. G. & Olafson, B. D. Benchmarking TriadAb using targets from the second antibody modeling assessment. *Protein Engineering, Design and Selection* **36**, gzad013, doi:10.1093/protein/gzad013 (2023).

Lassila, J. K., Privett, H. K., Allen, B. D. & Mayo, S. L. Combinatorial methods for small-molecule placement in computational enzyme design. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 16710-16715, doi:10.1073/pnas.0607691103 (2006).

Watson, J. L. *et al.* De novo design of protein structure and function with RFdiffusion. *Nature* **620**, 1089-1100, doi:10.1038/s41586-023-06415-8 (2023).

Dauparas, J. *et al.* Robust deep learning–based protein sequence design using ProteinMPNN. *Science* **378**, 49-56, doi:10.1126/science.add2187 (2022).

Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-589, doi:10.1038/s41586-021-03819-2 (2021).

17 Privett, H. K. *et al.* Iterative approach to computational enzyme design. *Proceedings of the National Academy of Sciences of the United States of America* **109**, 3790-3795, doi:10.1073/pnas.1118082108 (2012).

18 Rakotoharisoa, R. V. *et al.* Design of Efficient Artificial Enzymes Using Crystallographically Enhanced Conformational Sampling. *J Am Chem Soc* **146**, 10001-10013, doi:10.1021/jacs.4c00677 (2024).

19 Rothlisberger, D. *et al.* Kemp elimination catalysts by computational enzyme design. *Nature* **453**, 190-195 (2008).

20 Anand, N. *et al.* Protein sequence design with a learned potential. *Nature Communications* **13**, 746, doi:10.1038/s41467-022-28313-9 (2022).

21 Wang, J. *et al.* Scaffolding protein functional sites using deep learning. *Science* **377**, 387-394, doi:10.1126/science.abn2100 (2022).

22 Zhang, S., Zhang, J., Luo, W., Wang, P. & Zhu, Y. A preorganization oriented computational method for de novo design of Kemp elimination enzymes. *Enzyme and Microbial Technology* **160**, 110093, doi:https://doi.org/10.1016/j.enzmictec.2022.110093 (2022).

23 Sumida, K. H. *et al.* Improving Protein Expression, Stability, and Function with ProteinMPNN. *Journal of the American Chemical Society* **146**, 2054-2061, doi:10.1021/jacs.3c10941 (2024).

24 Broom, A. *et al.* Ensemble-based enzyme design can recapitulate the effects of laboratory directed evolution in silico. *Nature Communications* **11**, 4808, doi:10.1038/s41467-020-18619-x (2020).

25 Sterner, R. & Höcker, B. Catalytic Versatility, Stability, and Evolution of the (βα)8-Barrel Enzyme Fold. *Chemical Reviews* **105**, 4038-4055, doi:10.1021/cr030191z (2005).

26 Leaver-Fay, A. *et al.* ROSETTA3: an object-oriented software suite for the simulation and design of macromolecules. *Methods Enzymol* **487**, 545-574, doi:10.1016/B978-0-12-381270-4.00019-6 (2011).

27 Dunbrack, R. L. Rotamer Libraries in the 21st Century. *Current Opinion in Structural Biology* **12**, 431-440, doi:https://doi.org/10.1016/S0959-440X(02)00344-5 (2002).

28 Mayo, S. L., Olafson, B. D. & Goddard, W. A. Dreiding - a Generic Force-Field for Molecular Simulations. *Journal of Physical Chemistry* **94**, 8897-8909, doi:Doi 10.1021/J100389a010 (1990).

29 Dahiyat, B. I. & Mayo, S. L. Probing the role of packing specificity in protein design. *Proceedings of the National Academy of Sciences of the United States of America* **94**, 10172-10177 (1997).

30 Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nature Methods* **19**, 679-682, doi:10.1038/s41592-022-01488-1 (2022).

31 Allen, B. D., Nisthal, A. & Mayo, S. L. Experimental library screening demonstrates the successful application of computational protein design to large structural ensembles. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 19838-19843, doi:10.1073/pnas.1012985107 (2010).

32 Chovancova, E. *et al.* CAVER 3.0: A Tool for the Analysis of Transport Pathways in Dynamic Protein Structures. *PLOS Computational Biology* **8**, e1002708, doi:10.1371/journal.pcbi.1002708 (2012).

33 Dauparas, J. *et al.* Atomic context-conditioned protein sequence design using LigandMPNN. *bioRxiv*, 2023.2012.2022.573103, doi:10.1101/2023.12.22.573103 (2023).

34 Greenfield, N. J. Determination of the folding of proteins as a function of denaturants, osmolytes or ligands using circular dichroism. *Nature Protocols* **1**, 2733-2741, doi:10.1038/nprot.2006.229 (2006).

35 Sparta, K. M., Krug, M., Heinemann, U., Mueller, U. & Weiss, M. S. XDSAPP2.0. *Journal*
of applied crystallography **49**, 1085-1092, doi:doi:10.1107/S1600576716004416 (2016).
36 Kabsch, W. XDS. *Acta Crystallographica Section D* **66**, 125-132,
doi:doi:10.1107/S0907444909047337 (2010).
37 Liebschner, D. *et al.* Macromolecular structure determination using X-rays, neutrons and
electrons: recent developments in Phenix. *Acta crystallographica. Section D, Structural*
biology **75**, 861-877, doi:10.1107/S2059798319011471 (2019).
38 McCoy, A. J. *et al.* Phaser crystallographic software. *Journal of applied crystallography*
40, 658-674, doi:10.1107/S0021889807021206 (2007).
39 Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot.
Acta crystallographica. Section D, Biological crystallography **66**, 486-501,
doi:10.1107/S0907444910007493 (2010).
40 Hopkins, J. B., Gillilan, R. E. & Skou, S. BioXTAS RAW: improvements to a free open-
source program for small-angle X-ray scattering data reduction and analysis. *Journal of*
applied crystallography **50**, 1545-1553, doi:doi:10.1107/S1600576717011438 (2017).
41 Manalastas-Cantos, K. *et al.* ATSAS 3.0: expanded functionality and new tools for small-
angle scattering data analysis. *Journal of applied crystallography* **54**, 343-355,
doi:10.1107/S1600576720013412 (2021).
42 Jing, B., Berger, B., Jaakkola, T. AlphaFold Meets Flow Matching for Generating Protein
Ensembles. *arXiv* **2402**, 04845 (2024).
43 Franke, D. *et al.* ATSAS 2.8: a comprehensive data analysis suite for small-angle
scattering from macromolecular solutions. *Journal of applied crystallography* **50**, 1212-
1225, doi:doi:10.1107/S1600576717007786 (2017).
44 Lindorff-Larsen, K. *et al.* Improved side-chain torsion potentials for the Amber ff99SB
protein force field. *Proteins: Structure, Function, and Bioinformatics* **78**, 1950-1958,
doi:https://doi.org/10.1002/prot.22711 (2010).
45 He, X., Man, V. H., Yang, W., Lee, T.-S. & Wang, J. A fast and high-quality charge model
for the next generation general AMBER force field. *The Journal of Chemical Physics* **153**,
114502, doi:10.1063/5.0019056 (2020).
46 Izadi, S., Anandakrishnan, R. & Onufriev, A. V. Building Water Models: A Different
Approach. *The Journal of Physical Chemistry Letters* **5**, 3863-3871,
doi:10.1021/jz501780a (2014).
47 Darden, T., York, D. & Pedersen, L. Particle mesh Ewald: An N·log(N) method for Ewald
sums in large systems. *The Journal of Chemical Physics* **98**, 10089-10092,
doi:10.1063/1.464397 (1993).
48 Wang, J., Wang, W., Kollman, P. A. & Case, D. A. Automatic atom type and bond type
perception in molecular mechanical calculations. *Journal of Molecular Graphics and*
Modelling **25**, 247-260, doi:https://doi.org/10.1016/j.jmgm.2005.12.005 (2006).
49 Word, J. M., Lovell, S. C., Richardson, J. S. & Richardson, D. C. Asparagine and
glutamine: using hydrogen atom contacts in the choice of side-chain amide orientation. *J*
Mol Biol **285**, 1735-1747, doi:10.1006/jmbi.1998.2401 (1999).
50 Izaguirre, J. A., Catarello, D. P., Wozniak, J. M. & Skeel, R. D. Langevin stabilization of
molecular dynamics. *The Journal of Chemical Physics* **114**, 2090-2098,
doi:10.1063/1.1332996 (2001).
51 Berendsen, H. J. C., Postma, J. P. M., van Gunsteren, W. F., DiNola, A. & Haak, J. R.
Molecular dynamics with coupling to an external bath. *The Journal of Chemical Physics*
81, 3684-3690, doi:10.1063/1.448118 (1984).
52 Roe, D. R. & Cheatham, T. E., III. PTRAJ and CPPTRAJ: Software for Processing and
Analysis of Molecular Dynamics Trajectory Data. *Journal of Chemical Theory and*
Computation **9**, 3084-3095, doi:10.1021/ct400341p (2013).