1   **Ancient biases in phenotype production drove the functional evolution of a protein family**

2   Santiago Herrera-Álvarez[1]†, Jaeda E. J. Patton[2]†, Joseph W. Thornton[1,3]*

3   [1]Department of Ecology and Evolution; [2]Committee on Genetics, Genomics, and Systems
4   Biology; [3]Department of Human Genetics, University of Chicago; Chicago, IL, USA.

5   †These authors contributed equally to this work; the order of their names was determined by
6   coin flip.

7   *Corresponding author. Email: joet1@uchicago.edu

8   ABSTRACT

Biological systems may be biased in the phenotypes they can access by mutation[1–7], but the
extent of these biases and their causal role in the evolution of extant phenotypic diversity
remains unclear. There are three major challenges: it is difficult to isolate the effect of bias in the
genotype-phenotype (GP) map from that of natural selection in producing natural diversity[6,8–11],
the universe of possible genotypes and phenotypes is so vast and complex that a direct
characterization has been impossible, and most extant phenotypes evolved long ago in species
whose GP maps cannot be recovered. Here we develop exhaustive multi-phenotype deep
mutational scanning to experimentally characterize the complete GP maps of two reconstructed
ancestral steroid receptor proteins, which existed during an ancient phylogenetic interval when a
new phenotype—specific binding of a new DNA response element—evolved[12]. We measured all
possible DNA specificity phenotypes encoded by all possible amino acid combinations at sites in
the protein's DNA binding interface. We found that the ancestral GP maps are structured by
strong global bias—unequal propensity to encode the various phenotypes—and extreme
heterogeneity in the phenotypes accessible around each genotype, which strongly affect
evolution on both long and short timescales. Distinct biases in the two ancestral maps steered
evolution toward the lineage-specific functional phenotypes that evolved during history. Our
findings establish that ancient biases in the GP relationship were causal factors in the
evolutionary process that produced the present-day patterns of phenotypic conservation and
diversity in this protein family.

MAIN TEXT

Countless conceivable lifeforms have evolved rarely or never, and those that exist are mostly restricted to specific lineages[13–16]. No flying vertebrates have two pairs of wings, for example, and no turtles or frogs fly. What explains the biased distribution of phenotypes in nature? Classical explanations focus on the influence of selection[17,18], but it is possible that the propensities of biological systems to produce phenotypic variation could also shape evolutionary outcomes. A phenotype can become fixed in an evolving population only if it is first generated by mutation. If biological systems are more likely to produce some phenotypes than others[1–7], and if these propensities change over time as lineages diverge[19,20], then some phenotypes will be more likely to evolve in some taxa than in others.

Whether phenotype production has been an important cause of evolutionary outcomes is unclear, because most patterns of phenotypic variation observed in nature could arise from production biases, natural selection, or both, and disentangling their past influences is extremely challenging [6,8–11]. Ideally, we would isolate the phenotype production process by directly characterizing the complete genotype-phenotype (GP) map, which maps all possible combinations of mutations to the phenotypes they encode. This would allow us to precisely quantify the ability of a system to produce phenotypic variation, both on a global scale and by mutation from each particular genotype. The total space of genotypes and phenotypes is vast, but we reasoned that by combining three recent technical advances, this goal could become tractable for proteins and their biochemical phenotypes. The first technique is deep mutational scanning (DMS), which allows huge libraries of protein variants to be characterized experimentally[21]. The scope of genetic variation to be measured in a DMS study can be defined as all combinations of all 20 possible amino acid states at the sequence sites that determine the protein's phenotype of interest, thus encompassing all potential genetic variation at those sites[22–26]. Complete combinatorial DMS studies to date, however, have assayed only one or a few phenotypes that exist in extant proteins. Although this approach can reveal all genetic variants that encode these phenotypes, it cannot address why those particular phenotypes evolved in the first place; to understand why evolution turned out as it did, we must characterize the propensity of mutations to produce not only the phenotypes that evolved historically but also all the phenotypes that did not. The second technique—comprehensive multi-phenotype profiling—addresses this limitation by quantifying all possible phenotypes that a single protein can perform, such as binding of all possible substrates or DNA elements in a defined class[27–31]. We reasoned that by combining DMS with comprehensive multi-phenotype assays, we could map all possible phenotypes onto all possible genotypes within a defined scope. This would allow us to characterize the total capacity of a protein system to produce and access phenotypic variation by genetic change.

The phenotypes of extant lineages evolved long ago, so understanding the causal role of phenotype production in historical evolution requires GP maps to be characterized as they existed in the deep past. The third technique— ancestral protein reconstruction[32]— can address this problem by providing the protein backgrounds on which a comprehensive combinatorial multi-phenotype DMS study is performed. Moreover, characterizing such GP maps across a phylogenetic time series of reconstructed ancestral proteins[24,33] would reveal how biases in phenotype production may have changed over time and whether these biases are congruent with the trajectories of phenotypic evolution that actually unfolded during history.

Here we apply this approach to assess how phenotype production shaped the functional diversification of the steroid hormone receptor protein family. We use comprehensive multi-phenotype DMS to experimentally characterize GP maps of the binding interface of two reconstructed ancestral steroid hormone receptor DNA binding domains (SR DBDs) and their ability to encode specific recognition of all possible DNA response elements. We then analyze

these maps to understand 1) how they could shape potential phenotypic outcomes of evolution on
short and long timescales, 2) characterize the mechanisms that changed key features of the maps
across evolutionary time, and 3) assess the impact of the maps on the historical evolutionary
processes that yielded the lineage-specific patterns of DNA specificity in extant steroid hormone
receptors.

Two complete ancestral GP maps
SRs are a family of transcription factors that regulate physiological and reproductive biology in
bilaterian animals. Most bilaterian taxa have a single SR, which specifically binds to inverted
palindromes of the motif AGGTCA, called the estrogen response element (ERE; Fig. 1a). In
chordates, a gene duplication of the ancestral SR (AncSR1) produced two major SR classes,
which have different DNA specificity phenotypes: chordate estrogen receptors (ERs) retain the
ancestral ERE specificity, but a novel specificity for a palindrome of AGAACA, called the
steroid response element (SRE), evolved in the lineage leading to AncSR2, the common ancestor
of the chordate ketosteroid receptors (kSRs; Fig. 1a, b)[12]. Specificity for DNA is determined
primarily by the amino acid sequence of a recognition helix (RH) that binds in the DNA major
groove[34,35]. AncSR1 and AncSR2 DBDs differ by 34 amino acid replacements, but experiments
on the reconstructed proteins established that three amino acid changes in the RH were the
primary cause of the evolution of SRE specificity[12].
To understand how phenotype production may have shaped the evolution of SR-DBD
specificity, we characterized combinatorially complete GP maps of the DBD-response element
(RE) interface at the key ancestral timepoints AncSR1 and AncSR2. The scope of genotypes is
all possible $20^4 = 160,000$ amino acid variants at four variable sites in the recognition helix—the
three that changed between AncSR1 and AncSR2, plus one other that varies in the broader
nuclear receptor family (Fig. 1c). The scope of specificity phenotypes consists of all $4^2 = 16$
possible RE sequences that can be produced by all combinations of nucleotides at the two base
positions that vary between ERE and SRE. These two maps of the recognition helix-RE interface
can be thought of as submaps within the much larger GP map of the entire DBD, which are
connected by the 31 other "background" substitutions that occurred between the AncSR1 and
AncSR2 proteins (Fig. 1b).
We engineered two protein libraries, each containing all 160,000 variants of the
recognition helix in the background of either the AncSR1 or AncSR2 DBD, along with16 yeast
strains, each containing a GFP reporter driven by one of the REs (Fig. 1c, Extended Data Fig.
1a–e). We transformed each RE strain separately with the two protein libraries, with barcodes to
mark the strain and the ancestral background, for a total of 5.12 million protein-DNA complexes.
, we used an initial round of fluorescence-activated cell sorting to enrich the yeast libraries for
GFP-positive cells, pooled the enriched libraries, sorted cells in three replicates by their
fluorescence, and sequenced the sorted bins (Fig. 1d, Extended Data Fig. 1f, g). Using this
strategy, we obtained empirical fluorescence estimates for the majority of complexes with good
replicability ($r^2 = 0.92$ across replicates, excluding complexes at the lower bound of
fluorescence; Extended Data Fig. 2). Fluorescence of the remaining complexes was predicted
using a generalized linear model trained on the experimental data (Methods, Extended Data Fig.
3a–d)[36,37].
Each protein variant was assigned a DNA specificity phenotype based on these
experiments. A protein variant is classified as specific if it is functional in complex with only one
RE, promiscuous if it is functional on multiple REs, or nonfunctional if it is not functional on
any RE. We defined functional complexes as those having fluorescence at least as great as the

wild-type complex in each background (*i.e.* EGKA:ERE for the AncSR1 library and GSKV:SRE
for AncSR2) (Methods, Extended Data Fig. 3e–g).

Fig. 1 | Characterizing ancestral GP maps using multi-phenotype DMS. a, Amino acid
sequence of the recognition helix (RH) in extant and ancestral steroid receptor (SR) proteins and
the sequence of the RE they bind to. Colored residues are responsible for differences in protein-
RE specificity. **b**, Phylogeny of SRs. Each clade of proteins is colored by the RE sequence it
recognizes. In chordates, a historical transition from ERE to SRE specificity occurred along the
branch between AncSR1 (the common ancestor of all chordate SRs) and AncSR2 (the common
ancestor of vertebrate kSRs). The number of historical sequence changes along the AncSR1-
AncSR2 branch is shown; three of these in the recognition helix (RH) caused the specificity
switch[12]. **c, d**, DMS experiment to assay effects of RH genotype on binding to variable REs. **c**,
We built combinatorial libraries of all combinations of 20 amino acid states at four variable sites
in the RH (pink Xs), using the rest of the AncSR1 and AncSR2 DBDs as backgrounds (top left).
These were transformed into 16 *S. cerevisiae* strains, each containing one of the 16 possible RE
motifs (pink Ns, bottom left) genomically integrated upstream of a GFP reporter gene (right). **d**,
We assayed binding of DBD-RE complexes using FACS coupled with deep sequencing. For
each library, we performed an initial enrichment sort to select for GFP+ cells. We then grew up
the selected cells, pooled them across the 32 libraries, and resorted them into four fluorescence
bins in triplicate (binned sort). Sorted cells were deep sequenced to estimate the mean $\log_{10}$GFP
(*F*) of each combination of protein and RE genotypes.

Global bias in the AncSR1 GP map

The probability that a phenotype will evolve equals the probability that it will be produced by
mutation times the probability that, once produced, it will be fixed. The GP map
would have no effect on evolutionary outcomes if and only if it had two properties: isotropy—

148 encoding all phenotypes with equal probability—and homogeneity—producing the same
149 distribution of phenotypes from all starting genotypes in the map[38–41]. If the map is anisotropic,
150

Fig. 2 | Global and local bias in the AncSR1 GP map. a, Global production distribution in the
AncSR1 GP map. Bars represent the number of protein variants that bind specifically to each
RE. The dashed line shows the expected frequencies if the distribution were unbiased. *B,*
phenotype bias, calculated as one minus the entropy (base 16) of the distribution. **b**, Sequence
space network of the AncSR1 GP map. Nodes represent functional protein variants, colored by
their RE specificity; white nodes, promiscuous genotypes. Edges connect protein variants that
can be interconverted by a single nucleotide change. Genotype clusters (1–6, ordered by
decreasing size) identified by a community structure detection algorithm are shown in gray.
Sequence logos show amino acid frequencies at the variable RH sites in each cluster. **c**, Bottom:
Frequencies of specificity phenotypes within each genotype cluster; the global production
distribution is shown for comparison. Asterisks, phenotypes significantly enriched within a
cluster relative to the global production distribution (Fisher's exact test, $p < 0.05$ after Bonferroni
correction). Top: strength of phenotype bias (*B*) in each cluster. Red line, *B* of global production
distribution. **d**, Distribution of phenotype bias (*B*) of the 1-mutant neighborhood of every RE-
specific protein variant in the main network component. Dot shows the mean. Dashed red line,
global phenotype bias. **e**, Proportion of neutral neighbors per RE-specific protein variant in the
main component of the AncSR1 map. Dot shows the mean. *P-value,* probability that the mean
would be at least as great as observed if phenotypes were randomly reassigned in the main
component ($n = 91$). **f**, Distribution of the number of new phenotypes accessible within one
mutation, across all RE-specific variants in the AncSR1 main component. **g**, Mean distance
between pairs of phenotypes in the AncSR1 main component. The color of each cell shows the
mean of the length of the most direct path from every genotype encoding one phenotype to every
genotype encoding the other. Bonferroni corrected *p*-values for a two-sided permutation test
where phenotype associations were shuffled within the main component: * $p < 0.001$.

then phenotypes more likely to be produced would be more likely to evolve; if the map is
heterogeneous, then the probability that each phenotype will be produced—and hence evolve—
would change as lineages diverge from each other across the map.
We assessed the isotropy of the AncSR1 GP map by characterizing the frequency
distribution of DNA specificity phenotypes encoded by all functional protein variants. Only 107
out of 160,000 total genotypes in the library were functional (0.07%). Of these, the majority (91)
were specific for a single RE. We calculated the bias (B) of this global phenotype distribution,
defined as 1 minus the Shannon entropy (base 16); B can range from 0 when specificity for all 16
REs is encoded with equal frequency to 1 when only a single phenotype is encoded. We found
that the distribution is strongly anisotropic ($B = 0.42$). Two specificity phenotypes—ERE and
SRE—together account for >60% of all specific genotypes, and only five others can be produced
at all; nine phenotypes are not encoded by any protein variant (Fig. 2a).
We refer to this anisotropy as global bias in the GP map[40]. Global bias in the AncSR1
map imposes hard limits on phenotypic evolution—the majority of conceivable phenotypes
could never evolve in this map, even if they conferred strong fitness advantages. The global bias
is also congruent with evolutionary history—the phenotypes that evolved historically in the two
lineages descending from AncSR1 are also the most frequently encoded.

Local bias in the AncSR1 GP map
We next assessed the homogeneity of the AncSR1 GP map using Maynard-Smith's classic
network model of sequence space[42]. Each functional protein variant is a node with its
experimentally defined phenotype. Nodes are connected by edges if their amino acid sequences
can be interconverted by a single nucleotide change given the standard genetic code.
Nonfunctional variants are excluded from the network, based on the assumption that they will be
removed quickly from evolving populations by purifying selection.
We found that the distribution of phenotypes in AncSR1 sequence space is strongly
heterogeneous. Although the majority of functional genotypes (91%) and phenotypes (6 of 7) are
mutually connected in a single main network component, each phenotype tends to be sequestered
in a local region (Fig. 2b). Using a community structure detection algorithm[43], we found that the
main network component can be partitioned into six clusters of genotypes that have dense
connectivity within clusters and weak connectivity between (Fig. 2b). The phenotype bias B
within every single cluster is higher than the global bias of the map, and 5 of 6 clusters are
significantly enriched for a single specificity phenotype, which differ among all 5 clusters (Fig.
2c). The clumpy distribution of phenotypes in sequence space arises from the simple fact that
similar genotypes, which are connected to each other in sequence space, are likely to encode
similar phenotypes (Fig. 2b, logos).
This heterogeneity creates local bias[40]: the propensity to produce phenotypes depends
strongly on the particular genotype occupied at the time. The one-mutant neighborhood around
every genotype has extremely high bias (mean $B = 0.91$; Fig. 2d), indicating that individual
genotypes can access much less phenotypic variation than is encoded across genotype space as a
whole. Most mutations are phenotypically neutral (79% of edges; Fig. 2e), and most genotypes
can directly access at most one new phenotype (Fig. 2f). The historical starting genotype
(EGKA), for example, has access to only one functional neighbor, which also has ERE
specificity. Another consequence of heterogeneity is that phenotypes, aggregated over the
genotypes that encode them, are differentially accessible to each other, with substantial variation
in the number of mutations required to transform each phenotype into the others (Fig. 2g). For
example, SRE-specific protein genotypes are directly accessible from nodes encoding specificity

for AT and GA, but they are multiple substitutions away from all ERE-specific genotypes (Fig.
2b).
The GP map of AncSR1 is therefore both anisotropic and heterogeneous, and these
properties impose global and local biases on the production of phenotypes. Global bias favors
production of the historical phenotypes ERE and SRE and entirely prevents the production of
most conceivable phenotypes. Local bias further restricts the number of accessible phenotypes
from each particular genotype, favoring conservation over the evolution of new phenotypes,
including from the historical genotype EGKA.

Biases in the GP map affect phenotypic outcomes of evolution
To characterize the potential influence of the AncSR1 GP map on the outcomes of evolution, we
modeled evolution on the network of functional amino acid genotypes as a discrete-time Markov
chain from every possible starting genotype given a variable trajectory length. Each time-step in
a trajectory is an amino acid substitution, the probability of which is weighted by the number of
single-nucleotide mutations that can mediate it; the relative probability of evolving a given
phenotype at the end of the trajectory is the sum of the probabilities of evolving all genotypes
that encode it. This model, which corresponds to neutral molecular evolution in which all
functional genotypes have equal fitness[42,44], represents a null scenario: the fixation process
imposes no biases on evolutionary outcomes except to prevent the loss of function via purifying
selection, thus allowing us to isolate the influence of biases imposed by the GP map on
evolutionary outcomes.
We first computed the equilibrium distribution of phenotypic outcomes after an infinite
number of substitutions. This represents the limiting case at which the distribution of outcomes is
insensitive to the starting genotype and does not change with additional substitutions. The
equilibrium outcome distribution is well correlated with the global production distribution (Fig.
3a, $r^2 = 0.82$), reflecting the constraints imposed by the global production bias. However, there
are differences: the equilibrium distribution is more biased ($B = 0.46$), and whereas ERE and SRE
specificity are the two most frequently encoded phenotypes, ERE and AT specificity are the
most likely equilibrium outcomes (Fig. 3a). This difference arises because most AT-specific
genotypes are located centrally within the network, while SRE-specific genotypes are in a more
peripheral cluster (Fig. 2b) and are therefore less likely to be occupied. The heterogeneous
connectivity of the GP network and global production bias therefore affect evolutionary
outcomes, even over infinitely long timescales.
On finite timescales, local bias strongly affects evolutionary outcomes. After 3
substitutions, for example—the shortest path between the historical ancestral and derived
genotypes—the outcome distributions are very strongly biased (mean $B = 0.8$ across starting
genotypes, Fig. 3b), because most genotypes can reach only a few new specificity phenotypes by
a path of this length (Extended Data Fig. 4). The bias in outcomes gradually decays as
trajectories get longer, but it takes 42 substitutions (10.5 per site) for the mean bias to decrease to
within 0.05 units of the equilibrium (Fig. 3b, vertical dashed line). By comparison, the maximum
root-to-tip branch length in the steroid receptor DBD phylogeny (Fig. 1b), which spans over 500
million years of evolution, is just 2.2 substitutions per site. The phenotypes likely to evolve on
phylogenetically relevant timescales are therefore strongly affected by local bias in the GP map.
Another consequence of local bias is that outcomes are strongly contingent on the genetic
starting point. Consider a trajectory length of 8 substitutions—long enough for new phenotypes
to become accessible from most starting points, but not so long that the influence of local bias is
lost. At this timescale, genotypes differ dramatically in the distribution of phenotypes that evolve

from them (Fig. 3c). Much of this variation is explained by the genotype cluster to which the
starting node belongs (Fig. 3c), because evolutionary trajectories rarely jump between weakly
271

Fig. 3 | The AncSR1 GP map biases evolutionary outcomes towards phenotype
conservation. a, Comparison between the global production distribution and the long-term
equilibrium distribution of phenotypic outcomes in the AncSR1 main network. Each dot shows
the frequency of one specificity phenotype in the two distributions. Black dot at the origin
represents nine phenotypes not encoded in the map. Dashed gray line, $y = x$. Squared Pearson's
correlation coefficient is shown. **b**, Strength of bias (B) in evolutionary outcomes as a function of
the length of evolutionary trajectories. Each gray dot shows the B of the outcome distribution for
trajectories of a given number of substitutions starting from one node on the main network
component. Solid blue and black lines show the mean across all starting genotypes and from
EGKA, respectively. Dashed horizontal red and cyan lines show B of the global production
distribution and the equilibrium distribution, respectively. Vertical dashed line shows the number
of substitutions required for mean B to reach within 0.05 units of the equilibrium value. The
secondary x-axis (above) shows the trajectory length as substitutions per site. **c**, Distribution of
evolutionary outcomes after 8 substitution steps from every starting genotype in the AncSR1
main network component, organized by the cluster of the starting genotype (top). Bottom bar
shows the phenotype of each starting genotype. Bars at right show the average outcome
distribution for all starting genotypes. **d**, Distribution of the probability of phenotype
conservation after 8 substitution steps across all specific starting genotypes in the AncSR1 main
network component. Dot shows the mean. **e**, Evolutionary outcomes become less similar as
starting genotypes diverge from each other. Each dot shows the similarity of the distributions of
phenotypic outcomes (Pearson's r^2) of 8-step trajectories starting from a pair of genotypes,
versus the number of network edges between the pair. Blue line, mean similarity across all pairs
of starting genotypes. **f**, Probability of evolving each specificity phenotype starting from EGKA
as a function of the number of substitutions.

connected clusters and clusters are strongly enriched for individual phenotypes. Even at this

timescale, the direction of phenotypic evolution on average favors conservation of the starting
phenotype (Fig. 3d), but when new phenotypes evolve, these too differ strongly among starting
genotype (Fig. 3c).
A final consequence of local bias is that as lineages diverge from each other across the
map, the distributions of phenotypic outcomes likely to evolve from them become increasingly
dissimilar. The correlation between the distributions of phenotypic outcomes after eight-step
evolutionary trajectories from pairs of starting genotypes depends strongly on the distance
between those genotypes in the network. For pairs of genotypes that are one substitution apart,
the average r^2 is 0.88, but this correlation drops to 0.50 when the genotypes are three steps apart
and is entirely lost at 11 steps ($r^2 = 0.02$, the maximum distance on the network) (Fig. 3e). Biases
in the outcomes of phenotypic evolution therefore become distinct among lineages as they
traverse the GP map.

The AncSR1 GP map favored historical conservation of ERE specificity
Local and global bias have a particularly strong and long-lasting impact on the outcomes of
evolutionary trajectories that begin from the historical genotype of the recognition helix in
AncSR1 (EGKA). It takes 80 substitutions for the bias in phenotypic outcomes from this starting
point to decay to within 0.05 units of equilibrium, almost double the average across genotypes
(Fig. 3b, blue vs. black solid lines). It takes at least 5 substitutions for any new specificity
phenotype to be accessed, and even after 8 substitutions the probability of conserving ERE
specificity is still 0.90 (Fig. 3f). The AncSR1 GP map heavily favors phenotypic conservation
from the historical starting genotype across phylogenetically relevant timescales. Bias imposed
by the GP map is therefore congruent with the long-term historical conservation of ERE
specificity in the lineages that descend from AncSR1 and lead to modern-day estrogen receptors.
The historical outcome that evolved in AncSR1's other descendant linage—acquisition of
SRE specificity in the kSR clade—was very unlikely on phylogenetic timescales. SRE-specific
genotypes are distant from EGKA (Fig. 2b), so the probability of evolving SRE specificity after
eight substitutions is only 0.0008 (Fig. 3f), despite the fact that this is the second-most frequently
encoded specificity phenotype in the network overall. Strong local bias around EGKA therefore
overrides the global bias towards SRE specificity, making the historical outcome in the kSR
clade extremely unlikely.

Evolution of a different GP map in AncSR2
Given that local bias made SRE specificity unlikely to evolve from the ancestral genotype in the
AncSR1 map, how could this phenotype have historically evolved in the kSRs? We reasoned that
the GP map must have changed along the branch leading to AncSR2 when SRE specificity was
acquired. Previous experiments showed that the background substitutions that occurred outside
the recognition helix during this interval had a nonspecific permissive effect on both ERE and
SRE activation, allowing the protein to tolerate the historical substitutions and other mutations in
the RH (Fig. 1b)[12,24]. We predicted that the background substitutions had a similarly permissive
effect across all REs, increasing the number of functional genotypes in the map and the number
of phenotypes they encode, including SRE specificity and others.
To assess this hypothesis, we characterized the GP map of the RH sites in AncSR2 and
compared it to the map in the AncSR1 background. As predicted, the number of functional
genotypes and phenotypes both massively increased (Fig. 4a, b). There are 2,407 functional
protein genotypes in the AncSR2 map, an increase of >20-fold over the AncSR1 background.
Fourteen of the 16 possible specificity phenotypes are now encoded in the map, twice as many as

in AncSR1 (Fig. 2a, 4a). The background substitutions therefore dramatically expanded the functional genetic and phenotypic variation that can be produced within the recognition helix.

Fig. 4 | Global and local bias and connectivity changed in the AncSR2 GP map. a, Global production distribution and global B of the AncSR2 GP map. **b**, Sequence space network of the AncSR2 GP map. **c**, Number of one-step neighbors per protein variant in each network. Dots show the mean of each distribution. **d**, Bottom: Frequencies of specificity phenotypes within each genotype cluster (1–14, ordered by decreasing size); the global production distribution is shown for comparison. Only the 14 largest clusters, which contain >90% of genotypes, are shown. Asterisks, phenotypes significantly enriched within a cluster relative to the global production distribution (Fisher's exact test, $p < 0.05$ after Bonferroni correction). Top: strength of phenotype bias (B) in each cluster. Red line, B of global production distribution. **e**, Distribution of the number of new phenotypes accessible within one mutation, across all RE-specific protein variants in the AncSR2 main component. **f**, Proportion of neutral neighbors per RE-specific variant in the main network component of the AncSR1 and AncSR2 maps. Dots show the mean. *p-value,* probability that the mean would be at least as great as observed if phenotypes were randomly reassigned in the main component of each map (AncSR1 $n = 91$, AncSR2 $n = 2,402$).

Connectivity between genotypes in the map increased, reducing local bias and facilitating access to new phenotypes. In the AncSR2 network, all but five of the 2,407 functional nodes are connected in a single main component (Fig. 4b), and the mean number of edges per node is 10.7, a three-fold increase compared to the AncSR1 network (Fig. 4c). Genotype clusters are still present, but bias within clusters is weaker than in the AncSR1 map (Fig. 2c, 4d). As a

consequence, genotypes have more access to new phenotypes: >50% of genotypes in the
AncSR2 map can access between 1 and 4 new phenotypes within a single mutation (Fig. 4e,
compare to Fig. 2e), because genotypes are typically connected to far more non-neutral
neighbors (Fig. 4f).
Finally, the global production distribution of phenotypes also changed across this
interval. In the AncSR2 map, SRE became the most frequently encoded phenotype (39% of
specific variants), and ERE's rank declined from first to seventh (encoding just 3% of specific
variants) (Fig. 2a, 4a). The background substitutions therefore realigned the global phenotype
bias from favoring the ancestral specificity to producing the derived specificity.

The AncSR2 GP map favored evolution of SRE specificity
These changes in the AncSR2 GP map dramatically altered the likely phenotypic outcomes of
evolution. At long-term equilibrium using our Markov model and the AncGR2 map, the most
likely evolutionary outcome is now SRE specificity, with a probability close to 40% (Fig. 5a,
compared to <20% in the AncSR1 map). At moderate timescales as well, SRE specificity is the
most likely outcome across the majority of starting genotypes (Fig. 5b). The probability of
evolving new phenotypes overall is considerably higher in the AncSR2 network compared to
AncSR1 (mean probability of conservation after 8 steps 0.47 in AncSR2 but 0.61 in AncSR1,
Fig. 3d, 5c).
387

Fig. 5 | The AncSR2 GP map biases evolutionary outcomes towards SRE specificity. a,
Comparison between the global production distribution and the long-term equilibrium
distribution of phenotypic outcomes in the AncSR2 main network. Dashed gray line, $y = x$. **b**,
Distribution of evolutionary outcomes after 8 substitution steps from every starting genotype in
the AncSR2 main network component, organized by the cluster of the starting genotype (top).
Bottom bar shows the phenotype of each starting genotype. Bars at right show the average
outcome distribution for all starting genotypes and all non-SRE-specific starting genotypes,
respectively. **c**, Distribution of the probability of phenotype conservation after 8 substitution
steps across all specific starting genotypes in the AncSR2 main network component. Dot shows

the mean. **d**, Number of genotypes (top) and phenotypes (bottom) accessible as a function of the
length of evolutionary trajectories. Lines show the mean across all starting genotypes in each
network. Gold, AncSR1 network; teal, AncSR2 network. **e**, Strength of bias (B) in evolutionary
outcomes as a function of the length of evolutionary trajectories. Lines and colors are the same
as in Fig. 3b. **f**, Distribution of the similarity in outcome distributions (Pearson's r^2) for 8-step
trajectories starting from all pairs of genotypes in the AncSR1 and AncSR2 main networks. Dots
show means. **g**, Probability of evolving each specificity phenotype starting from EGKA as a
function of the number of substitutions.
406
These changes in evolutionary outcomes are attributable to the increased connectivity of
the AncSR2 network and the shift in the global production distribution. From any starting point,
the increase in functional nodes and connectivity allows access to far more genotypes and new
phenotypes (Fig. 5d). As a result, the influence of local bias is lost faster, and trajectories more
rapidly converge on the equilibrium distribution (Fig. 5e), which more closely resembles the
production distribution than in the AncSR1 background (Fig. 5a). Evolutionary outcomes are
also more similar across pairs of starting points than they were in the AncSR1 map (Fig. 5f).
Combined with the shift in the global production distribution, this causes SRE specificity—
which was already the second-most likely outcome in the AncSR1 map—to become the most
likely outcome from a majority of starting points in the AncSR2 background.
From the historical RH genotype EGKA (the AncSR2 protein with the RH states reverted
to their ancestral states), the likely outcomes of phenotypic evolution are dramatically different
than in the AncSR1 map. EGKA is much less mutationally isolated in the AncSR2 network, so
the probability of conserving ERE specificity after 8 substitutions drops from 0.9 in the AncSR1
map (Fig. 3f) to 0.07 in the AncSR2 map (Fig. 5g). The probability of evolving new specificity
phenotypes on moderate timescales increases accordingly: after just three steps, two new
phenotypes—including SRE specificity—are more likely than conservation of ERE. By six
steps, SRE specificity becomes the most likely of all phenotypic outcomes.
The background substitutions that occurred along the branch to AncSR2 therefore
changed the GP map of the RH in a way that dramatically changed the probable phenotypic
outcomes of evolution. This map strongly favors phenotypic diversification, and it makes the
particular phenotype that historically evolved in the kSR lineage the most likely of all possible
outcomes.

Simple biophysical mechanisms changed the GP map

Finally, we sought insight into the biophysical mechanisms that changed the GP map of the
recognition helix between AncSR1 and AncSR2. Although our experiments provide a functional
rather than biophysical readout, different biophysical mechanisms predict different patterns of
functional change between the AncSR1 and AncSR2 maps. We therefore analyzed the change in
fluorescence of each protein-DNA complex variant between the two backgrounds to identify
potential biophysical mechanisms and considered them in light of existing crystal structures. We
found evidence for two major mechanisms.
First, the background substitutions between AncSR1 and AncSR2 appear to have caused
a universal increase in affinity across all protein-DNA complexes. Previous experiments and
crystal structures showed that the background substitutions improve nonspecific DNA contacts
and binding cooperativity to both ERE and SRE[12,24]; we therefore hypothesized that affinity
increased universally for all amino acid variants across all 16 REs. To test this hypothesis, we fit
a simple model in which fluorescence in each ancestral background is a function of a complex's

Fig. 6 | Nonspecific effects of background substitutions on DBD-RE affinity. a, Fluorescence of each complex in the AncSR1 vs. AncSR2 background, scaled between the upper and lower bounds for each background. Curve shows best-fit model assuming that the affinity of every complex in the AncSR2 background is related to its affinity in the AncSR1 background by the same scaling factor. Shaded region around the curve (barely visible) shows bootstrapped 95% confidence interval (CI). The Pearson's r^2 between the data and model predictions is shown ($n = 2,627$). Histograms show distribution of F in each background. Dashed lines show the fluorescence of the wild type complex in each background (AncSR1-EGKA:ERE or AncSR2-GSKV:SRE). Colors indicate the backgrounds in which each genotype is functional. **b**, Amino acid frequencies at the variable RH sites across all functional protein variants in the AncSR1 and AncSR2 maps. **c**, Distribution of the number of neighbors gained in the AncSR2 background across all functional protein variants in the AncSR1 background that remain functional in the AncSR2 background. Dashed line, mean. **d**, Correlation between the number of protein variants bound per RE in each background. Black line, linear fit to all REs except ERE; shaded region, 95% CI. **e**, Same as **a**, but fitting a model in which the background substitutions affect affinity of all variants for ERE by one scaling factor and for all other REs by a different scaling factor. Purple, observed fluorescence and best-fit model predictions for ERE complexes; gray, for non-ERE complexes. **f**, Crystal structure of the AncSR1-EGKA protein in complex with ERE (PDB 4OLN). The RH backbone is shown as a ribbon, with key side chains shown as sticks. The gray surface shows ERE, with variable bases and backbone as sticks. In this complex, glutamine (q) at site 36 forms a hydrogen bond (yellow dashed line) with the DNA backbone, and lysine (k) at site 32 forms two hydrogen bonds to the ERE-specific bases G and T. **g**, Same as **f**, but with the AncSR2-EGKA crystal structure (PDB 4OND). Substitution to glutamic acid (E) at site 36 abolishes the ancestral hydrogen bond to the DNA backbone and results instead in electrostatic repulsion from the backbone. This deforms the recognition helix, abolishing the hydrogen bonds between K32 and the G and T bases. In **F** and **G**, lowercase letters represent ancestral amino acid states, and uppercase derived.

affinity, and affinity is scaled by a constant factor in AncSR2 relative to AncSR1. The model fits the data very well ($r^2 = 0.95$; Fig. 6a), with an estimated 70-fold universal improvement in affinity in the AncSR2 background. This apparent increase in affinity explains the vast increase

in functional genotypes and specificity phenotypes between AncSR1 and AncSR2, because many
protein-DNA complexes that had weak affinity in the AncSR1 background—and were therefore
nonfunctional—bind strongly enough in AncSR2 to produce functional levels of fluorescence.
The number of promiscuous protein variants also increases, because many variants cross the
threshold for functionality on multiple REs (Extended Data Fig. 5a). A universal improvement in
affinity explains not only the increased size but also the greater connectivity of the AncSR2
network: the background substitutions do not qualitatively change the amino acid determinants
of binding but instead make them less stringent (Fig. 6b), so many of the newly functional nodes
in AncSR2 are close neighbors of those that were already functional in AncSR1, with an average
gain of 11 new neighbors per node (Fig. 6c).
The second apparent mechanism is that the background substitutions negatively affect
specific binding to ERE, shifting the global production bias away from ERE and leaving SRE as
the most-encoded phenotype in the AncSR2 background. A universal affinity increase predicts
that the number of variants with every specificity phenotype should increase proportionally
across the AncSR1-AncSR2 interval; this pattern holds, but ERE is an outlier, with far fewer
variants than would be expected given the pattern for other phenotypes (Fig. 6d). Moreover, ERE
complexes exhibit notably lower fluorescence in the AncSR2 background than predicted by a
universal increase in affinity (Extended Data Fig. 5b). We estimated the effect of the background
substitutions on ERE affinity by incorporating a background-by-ERE interaction term into our
affinity-fluorescence model; adding this parameter improves the fit to the data ($r^2 = 0.97$), with
the background substitutions improving ERE affinity by an estimated 2.3-fold, compared to 99-
fold for all other REs (Fig. 6e). The extent of the relative reduction in fluorescence differs among
protein variants, however, suggesting additional specific interactions between background
substitutions and amino acids in the recognition helix (Extended Data Fig. 5c). Crystal structures
of the EGKA:ERE complex[12] suggest a possible structural basis for the global reduction in ERE
affinity: one of the background substitutions (q36E) deforms the protein backbone of the
recognition helix, abolishing two hydrogen bonds that are formed between a conserved residue
and bases in the ERE (Fig. 6f, g). Corroborating this mechanism, the background substitutions
also shift the global bias away from AT specificity (Fig. 6d), and this is the only other RE that
can form these hydrogen bonds.
The structure of the GP map therefore changed between AncSR1 and AncSR2 via two
simple biophysical mechanisms. By increasing all proteins' affinity for all REs, while also
impairing their affinity for ERE, the background substitutions reduced local bias and changed the
direction of global bias, facilitating the evolution of many new genotypes and phenotypes and
shifting the protein's global propensity away from conserving ERE specificity to evolving the
new specificity for SRE.

Robustness to assumptions
To assess whether our conclusions are sensitive to assumptions that we made in our analysis, we
reanalyzed our experimental data under different models and assumptions. First, we applied
different thresholds to classify genotypes as functional or nonfunctional, included promiscuous
genotypes when characterizing global production distributions, and characterized these
distributions using only genotypes with experimentally measured phenotypes. In every case, we
observed similar forms of bias in both the AncSR1 and AncSR2 GP maps to those reported
above (Extended Data Fig. 6).
Second, instead of treating the protein as an evolutionary unit independent of the RE, we
repeated our analyses using an alternative sequence space network in which the protein and RE

coevolve as a complex. In this model, evolution may occur via single-step amino acid mutations
in the protein or nucleotide mutations in the RE. Our main conclusions again hold: global and
local biases impact phenotypic evolution over long and short timescales, favoring ERE
conservation in the AncSR1 map and evolution of SRE specificity in AncSR2 (Extended Data
Fig. 7).
Finally, we addressed uncertainty about the ancestral sequences. AncSR1 and AncSR2
DBD reconstructions have very high confidence, containing just five and zero ambiguously
reconstructed sites, respectively[33]. Experimental data from a prior single-mutant DMS study
show that the effects of mutations in the RH are virtually identical when they are introduced into
the AncSR1 background or into an alternative reconstruction of AncSR1 that incorporates all
plausible alternative amino acids at the ambiguously reconstructed sites ($r^2 > 0.99$; Extended
Data Fig. 8)[33]. The very limited uncertainty about the AncSR1 ancestral sequence is therefore
likely to have little or no effect on our conclusions.

The GP map was a cause of historical phenotypic evolution
Our data establish that global and local biases in the two ancestral GP maps we studied were
causal factors in the historical lineage-specific evolution of DNA specificity. Establishing
causality in a multifactorial framework requires 1) evidence that a putative cause increases the
probability of the outcome(s) of interest, and 2) evidence for a specific mechanism by which the
cause affects the outcome's probability[45]. Concerning the first requirement, our experiments
show that biases in the AncSR1 map increased the probability that ERE specificity would be
evolutionarily conserved, and biases in the AncSR2 map increased the probability that SRE
specificity would be acquired. The second requirement is satisfied by a simple axiom of
population genetics: the probability that a phenotype will evolve is the product of its probability
of production and its probability of fixation under the influence of selection and drift. If biases in
the GP map increase the production probability, then evolutionary outcomes will in turn be
biased.
A cause must precede its effect. The biases that favored the conservation of ERE
specificity in the AncSR1 map are ancestral to the ER lineage in which that outcome occurred
(Fig. 1b). This map persisted unchanged for hundreds of millions of years of phenotype
conservation, because zero amino acid changes anywhere in the DBD occurred along the
descendant branches leading from AncSR1 to ERα in the ancestor of all bony vertebrates. Even
most present-day ERa DBDs contain zero or at most a single substitution relative to AncSR1
(Extended Data Fig. 9). As for the acquisition of SRE specificity in the AncSR2 lineage, the
global bias that favors production of SRE specificity as the second-most encoded phenotype was
already present in the AncSR1 map. Further, the massive increase in connectivity of the AncSR2
map, which dramatically increased the propensity for new phenotypes to evolve, must have been
acquired before SRE specificity actually evolved, because the recognition helix substitutions that
conferred SRE specificity during history cannot be tolerated unless the background substitutions
that nonspecifically increased DNA affinity occurred first[12]. Our experiments do not resolve
whether the third major property of the AncSR2 map—a shift in global bias away from ERE
specificity that further enhanced the propensity to encode SRE specificity—occurred before or
after this phenotype was historically acquired.
We do not argue that selection played no historical role in the evolution of specificity. It
seems likely that purifying selection would have favored conservation of ERE specificity in the
chordate ERs, and positive selection could have contributed to fixation of SRE specificity in the

AncSR2 lineage. If so, however, selection would have further increased the probability of
outcomes that were already favored by the biases imposed by the GP map.
Our data show that the GP map's influence is strong enough to override the influence of
selection in many cases. For example, some global biases we observed are absolute. There are 9
specificity phenotypes that cannot be encoded at all in the AncSR1 GP map, and two cannot be
encoded in AncSR2; these could phenotypes never evolve, no matter how large a fitness benefit
they might confer. Local bias is also absolute in many cases: from every starting point, the vast
majority of phenotypes are impossible to produce directly by mutation, and most require many
substitutions before they become accessible. Selection would therefore be powerless to fix these
phenotypes over short or medium timescales. The GP map limited evolution to a small subset of
possible phenotypes; history, further influenced by selection and chance, played out within this
set.
There is evidence that features similar to those we observed in the steroid receptor GP
map affect biological systems and their evolution across levels of organization. Global bias is
apparent in other molecular[46,47] and developmental systems[48–51], and the resulting biases are
often congruent with natural patterns of diversity[52–55]. Local bias also appears to be widespread,
because most random mutations are phenotypically neutral if they are tolerated[51,56–59], and long-
term phenotype conservation is widespread in the fossil record[60]. When new phenotypes are
acquired, identical perturbations often yield different phenotypes in different lineages[61–64], and
convergent evolution becomes less likely among distantly related lineages[65]. As lineages evolve
across their GP maps, their biology inevitably changes, imposing new biases on the production
and future evolution of genotypes and phenotypes. It therefore seems likely that anisotropy and
heterogeneity are near-universal characteristics of GP maps[2,39,41], and that the biases these
properties create have shaped large-scale patterns of phenotype conservation and lineage-specific
evolutionary change across the tree of life.
Our study differs in kind from previous combinatorial DMS studies, which have
addressed the distribution in sequence space of just one or a few phenotypes that are encoded by
extant proteins, rather than the space of all possible phenotypes [22,24,25,63,66–68]. These studies have
shown that sequence landscapes are rugged, so the probability of reaching particular genotypes
encoding those phenotype may depend on the starting point and intermediate mutational steps.
Because those studies take the phenotypic "destination" for granted, they cannot address why
those phenotypes, rather than all the other conceivable outcomes, exist at all.
Our work shows that as a protein or other biological system moves through sequence
space, the set of phenotypes that it can produce changes at every step. Life is astonishing in its
diversity, but an even deeper puzzle lies in the fact that only a tiny fraction of conceivable
phenotypes have ever evolved, and those which have evolved are mostly limited to particular
taxa[14–16,69]. Chance and selection are likely important factors in explaining the patchy
distribution of phenotypes on Earth. But the very particular biology we observe today must also
reflect the constantly changing potential of biological systems, as they vary and diverge, to
generate new forms of life at every moment in time.

Methods

RE reporter strains

To measure binding of SR DBD to the 16 RE variants, we adapted a yeast GFP reporter system previously developed to measure binding to ERE and SRE, where GFP expression is well correlated with DNA affinity over a range of at least 2 M^{-2} ($r^2 = 0.74$)[33]. We engineered 16 yeast strains, each of which reports on binding of the DBD to one RE. We modified the yeast strain CM997 (YPS1000 MATa ho::KMX)[70] to replace the *KMX* gene at the *HO* locus with a construct containing yeast-enhanced GFP downstream of a minimal *CYC1* promoter with an array of four palindromic RE sites (tcaAG<u>NN</u>CAcagTG<u>NN</u>CTtga), each separated by a 19-nt sequence, along with a *HygR* gene. To ensure a consistent dynamic range of fluorescence across strains, we made changes to two RE strains in the nucleotide sequences flanking the palindromes at sites that do not affect specificity[34,35] (see Supplementary Methods for details). These constructs were transformed into yeast using the lithium acetate method[71] and selected for resistance to hygromycin and susceptibility to G418; integration was confirmed by Sanger sequencing.

To validate this reporter system, we measured fluorescence of each strain in the presence and absence of a DBD variant with universally high affinity to all REs (AncSR1+11P+GGKA)[12,29]. We used a low-copy yeast vector (pDBD) to express this DBD variant as a C-terminal fusion with an SV40 nuclear localization signal and a *S. cerevisiae* Gal4 activation domain (Gal4AD) under control of a pGAL1 promoter. We transformed this construct into each yeast strain using the lithium acetate method followed by G418 selection (50 μg/mL). Single colonies were inoculated in YPD+G418 and transferred to YPGal+G418 media for 6 hours to induce DBD expression. GFP fluorescence was measured on a BD LSRFortessa flow cytometer using a 488 nm laser with 505 nm long pass and 525/50 nm band pass filter. We used as the metric of fluorescence $\log_{10}(GFP/FSC-A^{1.5})$, which normalizes fluorescence to cell volume. All 16 strains showed DBD-dependent fluorescence across a similar dynamic range (Extended Data Fig. 1a–c).

AncSR1 and AncSR2 combinatorial library construction

We used as the wild-type protein sequences the maximum *a posteriori* AncSR1 and AncSR2 DBD sequences inferred from a maximum likelihood phylogeny of nuclear receptors[33].

We optimized codon usage for yeast and cloned the ancestral DBDs into the pDBD2.1 expression vector, which is modified from the pDBD vector[24,33] to express GFP at a level within the dynamic range of fluorescence for the wild type AncSR1:ERE and AncSR2:SRE complexes. A bidirectional pGAL1/GAL10 promoter simultaneously drives DBD and mCherry expression, which allowed us to monitor plasmid retention in yeast (Extended Data Fig. 1d).

Combinatorial mutant libraries were created by synthesizing oligos (IDT) with degenerate NNS codons to encode all 20 amino acids and a stop codon at four recognition helix sites of each ancestral protein (Extended Data Fig. 1e). To distinguish sequencing reads coming from different RE strains, 16 synonymously barcoded versions of the library were designed for each background (Extended Data Fig. 1e, Supplementary Table 1). Each barcode (REBC) differed by at least three nucleotides to ensure accurate read assignment despite sequencing errors. The oligos were cloned into the pDBD2.1 vector using the BsaI-HF Golden Gate Assembly kit (NEB), transformed into Invitrogen ElectroMAX DH5α-E E. coli, and maxiprepped (Supplementary Methods). Transformation yields exceeded 1.08×10^7 cfu per barcoded library, providing 56-fold coverage of the amino acid library size (Supplementary Table 2). Assemblies

were validated by Sanger sequencing of independent transformants and PCR of the plasmid
libraries to confirm the correct insert size.

Maxiprepped libraries (GenElute HP, Sigma-Aldrich) were transformed into the yeast reporter
strains using an optimized yeast electroporation protocol (Supplementary Methods).
Transformation yields exceeded 10^7 cfu per library (50-fold coverage), estimated by dilution
plating (Supplementary Table 2). Yeast libraries were flash-frozen in liquid N_2 in 200 OD_{600}-mL
aliquots with 25% glycerol and stored at –80°C. Multiple transformant rates estimated from
Sanger sequencing of individual colonies[72] were estimated to result in 0.03% or fewer cells with
multiple plasmid copies at time of sorting.

Cell sorting

We used fluorescence-activated cell sorting (FACS) to separate cells based on their GFP
expression. We performed two rounds of sorting: an initial "enrichment sort" to enrich for GFP+
variants in the full libraries, and a second, higher resolution "binned sort" on the enriched
libraries to generate quantitative fluorescence estimates for each variant. Enrichment sorting was
performed in batches of 8 libraries. Two glycerol stocks per library were thawed on ice, after
which cells were recovered for 2 hours in 400 mL YPD+chloramphenicol (chlor) per library at
30°C and 225 rpm. After recovery, G418 was added to the culture and a sample of cells was
taken for dilution plating. We recovered a minimum of 1.6×10^7 cfu per library (82-fold
coverage). After 15 hours of overnight growth, libraries were washed once in PBS, resuspended
to OD_{600} 0.25 in 50 mL YPGal+G418, and grown for 6 hours to induce DBD expression. Cells
were then spun down, washed once in PBS, resuspended in 5 mL PBS, and kept on ice for
sorting.

Sorting was performed at the University of Chicago Cytometry and Antibody Technology
Facility on a BD FACSAria Fusion machine. We used a 488 nm laser with 495 nm long pass
filter and 515/20 nm band pass filter for GFP detection, and a 561 nm laser with 595 nm long
pass filter and 610/20 nm band pass filter for mCherry detection. After gating on homogeneous
single cells and mCherry expression, we sorted cells into GFP– and GFP+ populations (Extended
Data Fig. 1f). To normalize fluorescence to cell volume, GFP gates were drawn to have a slope
of 1.5 on a log(FSC-A)-log(GFP) plot. We sorted 2.5×10^7 cells per library in the enrichment
stage (129-fold coverage, Supplementary Table 2).

Enriched cells from different libraries were pooled by GFP bin and grown in either 700 mL
(GFP+) or 2 L (GFP–) of YPD+G418+chlor. Cultures were grown overnight at 225 rpm and 22–
30°C, depending on the ratio of cells to media, until they were at least OD_{600} 3 but not yet
saturated. 200 OD_{600}-mL 25% glycerol stocks were then made for both the GFP+ and GFP–
cultures. 10 OD_{600}-mL of the GFP– culture was used for plasmid extraction using a previously
described protocol[21].

The binned sort was performed to yield three replicates per library. For each replicate, two 200
OD_{600}-mL glycerol stocks of GFP+ cells per enrichment sort batch were thawed on ice,
recovered in 400 mL YPD+chlor for 2 hours, and sampled for dilution plating. After adding
G418, cultures were grown overnight, achieving a recovery rate at least 4X the number of GFP+
cells collected during the enrichment sort (Supplementary Table 3). Overnight cultures were
pooled proportionally to the GFP+ cell counts from the enrichment sort, yielding a total of 100
OD_{600}-mL. The pooled cells were washed with PBS, induced for DBD expression in 400 mL
YPGal+G418 for 6 hours, washed again, resuspended in 40 mL PBS, and kept on ice for sorting.
Binned sorting followed the enrichment sort protocol but used four GFP bins instead of two

(Extended Data Fig. 1g), with ~1.6×10^8 cells collected per replicate. The number of sorted cells
and recovered reads was consistent across libraries and replicates (Supplementary Table 4).

Deep sequencing

After sorting, cells were grown in 100 mL YPD+G418+chlor per 10^7 sorted cells, or at least 100
mL per bin. Cultures were grown overnight to at least OD_{600} 3.0 but not yet saturated, and 50
OD_{600}-mL was collected per 10^7 sorted cells for plasmid extraction.

Sequencing libraries were constructed from plasmids extracted from the enrichment sort GFP–
population and the four binned sort populations using two rounds of amplification. In the first
round, the RH scanning and REBC regions of the DBD were amplified with primers that added a
6-nt barcode for bin and replicate identification (BRBC)[73]. For every 10 OD_{600}-mL of yeast used
for plasmid extraction, 3 µL of plasmid template was used in a 10 µL Q5 PCR reaction (NEB).
AncSR1- and AncSR2-specific primers were mixed proportionally to background-specific cell
counts (estimated from flow cytometry) to minimize amplification bias. To introduce nucleotide
diversity for improved cluster identification during Illumina sequencing, eight unique forward
and reverse primer pairs were used per bin and background to encode frameshift diversity and
attach read 1 primer sequences in both directions. PCR conditions included 52°C annealing for
13 cycles. Reactions were then pooled by bin/replicate and purified using the Zymo DNA Clean
& Concentrator Kit. In the second round, half of the first-round product was amplified with
primers to add Illumina P5 and P7 adapter sequences. PCR was performed in 50 µL Q5 reactions
(NEB) per 10 µL round 1 product reaction at 68.4°C annealing for 12 cycles. The final product
was size-selected on a 2% agarose gel, excised, purified using the Qiagen Gel Extraction Kit, and
re-purified with the Zymo DNA Clean & Concentrator Kit.

Final sequencing library concentrations were quantified by Qubit. Libraries were pooled
according to the number of cells sorted per bin/replicate, and 1.8 pM dilutions were prepared
according to Illumina's standard protocol. Replicate 1 of the binned sort libraries was sequenced
on a NextSeq High Output run. The remaining replicates were sequenced on a NovaSeq S1 run
at the University of Chicago Genomics Facility. We used standard read primers and 86 cycles for
read 1 and 80 cycles for read 2. This enabled us to bidirectionally sequence the region containing
the variable RH codons and REBC.

Mean fluorescence estimation, data cleaning and validation

Sequencing reads were processed using a custom pipeline. We used *sickle* v1.33[74] to filter reads
based on their quality: we kept reads with a Phred score $\geq$ 30 and a minimum length of 79
nucleotides. We then used *PEAR* v0.9.6[75] to merge the trimmed paired-end reads (minimum
assembly length 100 nucleotides). Finally, we used Biopython toolkit v1.79[76] to demultiplex the
assembled reads by DBD background, REBC, and BRBC. We only considered reads that
mapped exactly to the DBD background and allowed reads with at most one mismatch in the
REBC and one in the BRBC.

The mean fluorescence for protein:RE complexes observed in the binned sort data was estimated
as previously described[33]. We first estimated the proportion of cells of each complex g in each
bin b ($c_{g,b}$) from the proportion of reads in b that mapped to g. The mean fluorescence estimate
F_g for each complex was then estimated by taking the weighted mean fluorescence across bins
(mean fluorescence of each bin was measured during sorting), with weights $c_{g,b} / \sum_b c_{g,b}$.

We applied several filtering and correction steps to reduce global measurement error and
normalize fluorescence estimates between replicates. First, complexes with fewer than 27 reads
per replicate were removed to ensure >95% had a standard error (SE) of ≤ 0.1 (5% of the assay
range; Extended Data Fig. 2a). Second, complexes observed in only one replicate were excluded.
Third, batch effects were corrected by fitting I-splines to normalize fluorescence between
replicates (Extended Data Fig. 2b). Finally, SE was recalculated and complexes with SE > 0.1
were removed (Extended Data Fig. 2c). The final dataset had a mean pairwise Pearson's $r^2 =$
0.55 across replicates. The poor correlation arises primarily because the vast majority of
complexes are at the lower fluorescence bound, so r^2 is dominated by measurement noise; for
variants with fluorescence above the lower bound (roughly $F \geq -4.0$), r^2 improved to 0.92.
Altogether, we obtained fluorescence estimates for 628,732 AncSR1 and 658,475 AncSR2
variants, covering 24.6% and 25.7% of possible variants, respectively (excluding nonsense
variants).

Many variants were observed at high read depth in the GFP– bin of the enrichment sort but not in
the binned sort. We assigned these a null phenotype (lower-bound fluorescence) using a
statistical procedure based on read depth (see Supplementary Methods), resulting in 859,171
AncSR1 and 638,762 AncSR2 protein:RE null complexes (FDR = 0.1; Extended Data Fig. 2d).
This increased the total phenotyped variants to 1,487,903 in AncSR1 and 1,297,237 in AncSR2,
covering 58% and 51% of all possible variants, respectively.

To evaluate the accuracy of the sort-seq fluorescence values, we measured the fluorescence of 5
isogenic variants by flow cytometry, which were also spiked into the DMS libraries prior to the
binned sort. We found a high correlation between the fluorescence estimates from flow
cytometry and sorting (Pearson's $r^2 = 0.87$, Extended Data Fig. 2e). We additionally compared
the fluorescence estimates of the same variants that were contained in the DMS libraries and
again observed a strong correlation with flow cytometry measurements (Pearson's $r^2 = 0.97$,
Extended Data Fig. 2e).

To evaluate whether the REBC mutations affected fluorescence, we constructed AncSR1 and
AncSR2 "mini-libraries" consisting of each of the 16 REBCs engineered into the respective
wild-type protein variant. These were transformed via electroporation into the ERE or SRE
reporter strain, respectively, at 1:16 the scale of the full libraries, and spiked into the full-scale
libraries before sorting. The fluorescence of the mini-library variants did not differ significantly
by REBC ($p = 0.98$ AncSR1, $p = 0.99$ AncSR2, one-way ANOVA), indicating that fluorescence
estimates are directly comparable between libraries with different REBC mutations.

Fluorescence inference for missing complexes

To predict the fluorescence of the remaining complexes for which we did not obtain
experimental estimates, we fit a generalized linear model based on reference-free analysis
(RFA)[36,37] to the experimental data. The model estimates a sigmoid function to capture the
measurement bounds of the assay, plus additive and interaction effects (specific epistasis) for all
amino acid states at the four variable sites in the DBD and all nucleotide states at the two
variable sites in the RE. All possible intramolecular interactions up to third order amino acid
interactions in the DBD and second order nucleotide interactions in the RE, and intermolecular
interactions up to third order amino acid-by-second order nucleotide interactions were included.
L2 regularization with 10-fold cross validation was used to reduce overfitting (Extended Data
Fig. 3a; Supplementary Methods). We fit separate RFA models for each ancestral background

using the *glmnet* v4.1-6 R package[77]. Model fits to the observed data were $R^2 = 0.96$ for AncSR1
active complexes (0.31 all complexes) and $R^2 = 0.99$ for AncSR2 active complexes (0.88 all
complexes) (Extended Data Fig. 3b). These models were used to predict fluorescence values for
unobserved protein-RE complexes. We also used the fitted models to correct the predictions for
complexes in one of the modified strains that had systematically lower fluorescence
(Supplementary Methods; Extended Data Fig. 3c, d).

Classification of functional complexes

We classified complexes as functional if their fluorescence was not significantly lower than the
wild type complex, *i.e.* EGKA:ERE in the AncSR1 background and GSKV:SRE in the AncSR2
background. Complexes inferred as null from the enrichment sort were classified as
nonfunctional. For complexes observed in the binned sort, we used a *t*-test to account for
measurement error. For complexes with predicted fluorescence from the RFA models, we
performed a nonparametric bootstrap test using the distribution of model residuals concatenated
over the ten cross-validation fits to account for model prediction error (Supplementary Methods;
Extended Data Fig. 3e). For both tests, we used a Benjamini-Hochberg FDR threshold of 0.25 to
classify variants as nonfunctional if they were significantly less fluorescent than the wild type
complex (Extended Data Fig. 3f). The low stringency of the FDR threshold was chosen to reduce
the false positive rate for calling variants functional. The majority of complexes classified as
functional in both backgrounds had fluorescence estimates obtained from the binned sort
experiment (59.3% AncSR1, 75.4% AncSR2; Extended Data Fig. 3g).

Protein genotype networks

Following Maynard Smith's sequence space formalism[42], we built genotype networks consisting
of all functional RH variants in each DBD background. RH genotypes are connected by an edge
if they differ by a single amino acid mutation that can be produced via a single nucleotide
mutation given the standard genetic code. Genotype networks for joint protein-DNA models
follow a similar logic (Supplementary Methods). We used the R package *igraph* v1.5.1[78] to build
and analyze the genotype networks, and the software *gephi* v0.10.1[79] for network visualization.
To identify clusters of densely connected genotypes within the networks, we used the
cluster_edge_betweenness function from the R *igraph* package.

Model of evolution on GP maps

We modeled evolution on the genotype networks as an origin-fixation process under a strong
selection-weak mutation regime[80,81] To isolate the effect of the GP map's structure on evolution,
we considered a scenario in which all functional genotypes have equal fitness, so the fixation
probability is affected only by drift, and nonfunctional variants are removed by purifying
selection. The relative probability $P(i,j)$ of substitution from protein genotype i to genotype j is
therefore equal to the amino acid mutation rate μ_{ij}, normalized over all single-step neighbors of i
in the network. We assumed that there are no biases in the nucleotide mutation process (*e.g.*
transition vs. transversion rate), so μ_{ij} is affected only by unequal mutational access between
amino acids imposed by the genetic code. To incorporate this effect, we scaled μ_{ij} by the number
of possible nucleotide mutations that can change any nucleotide sequence that encodes i to any
nucleotide sequence that encodes j:

$$\mu_{ij} = \eta_{ij}^{o*} \times \prod_{o \neq o*} c_o \qquad (1)$$

where o indexes the amino acid position, $o*$ is the position at which the amino acid change
occurs, η_{ij}^{o*} is the number of possible single nucleotide changes that can produce the state in j
from the state in i at site $o*$, and c_o is the number of possible codons for the invariant amino acid
state at site o.

We used these transition probabilities to specify a discrete time Markov model for each ancestral
genotype network, where each step is a single amino acid substitution. Genotypes that are more
than one nucleotide change apart cannot access each other in a single time step, and the
probability of staying in the same genotype across a single step in the Markov chain is also zero.
We only considered functional genotypes within the main component of each network (the
largest connected component). With this model, we computed the probability distribution $\pi_{(k)}$ of
evolving all possible genotypes after k substitution steps given any specified set of starting
genotypes:

$$\pi_{(k)} = \pi_{(0)} \times P^k \qquad (2)$$

where P is the transition matrix with entries $P(i, j)$, $k > 0$, and $\pi_{(0)}$ is the vector of the probability
distribution of genotypes at time step $k = 0$. Setting a single element i of $\pi_{(0)}$ to 1 and all others to
zero corresponds to evolution from a single starting genotype; setting all elements of $\pi_{(0)}$ to $1/n$,
where n is the number of functional genotypes in the network, averages over all possible starting
genotypes. We calculated the relative probability of evolving a given specificity phenotype at
time step k by summing over all elements of $\pi_{(k)}$ that encode that specificity and normalizing by
the total probability across all specific protein genotypes.

Effects of background substitutions

To estimate the effect of the background substitutions between AncSR1 and AncSR2 on binding
affinity, we first considered a model where the background substitutions have a universal
nonspecific effect on affinity across all RH and RE genotypes. We assumed that fluorescence is
proportional to the fraction of protein bound to DNA. If a complex g has dissociation constant
$K_d(g)$ in the AncSR1 background, then its AncSR1 fluorescence (normalized to scale between 0
and 1) is:

$$F(g)_{AncSR1} = \frac{1}{1 + \dfrac{K_d(g)}{[RE]}} \qquad (3)$$

where $[RE]$ is the concentration of RE. If the background substitutions scale $K_d(g)$ by a factor α,
then fluorescence in the AncSR2 background is

$$F(g)_{AncSR2} = \frac{1}{1 + \alpha\left(\dfrac{K_d(g)}{[RE]}\right)} \qquad (4)$$

Rearranging these equations gives an expression for fluorescence in the AncSR2 background as
a function of fluorescence in the AncSR1 background and α:

$$F(g)_{AncSR2} = \cfrac{1}{1 + \alpha\left(\cfrac{1 - F(g)_{AncSR1}}{F(g)_{AncSR1}}\right)}$$ (5)

We fit this model to the AncSR1 and AncSR2 fluorescence data using orthogonal regression,
which accounts for measurement error in both backgrounds. We used only complexes that had
fluorescence measurements from the binned sort in both backgrounds, and whose fluorescence
was significantly greater than that of nonsense variants in either background ($n = 2{,}627$).
Fluorescence was normalized in each background to scale between the upper and lower bounds
inferred from the RFA models. Confidence intervals (CI) were constructed by bootstrapping the
data and refitting the model. The effect of the background substitutions was estimated to be $\alpha =$
0.014 (95% CI: 0.010–0.014), corresponding to a 70-fold increase in affinity (95% CI: 70–99).

We next considered a model where the background substitutions have a different effect on ERE
affinity than they do on other REs. We modified the model such that α_1 represents the ERE-
specific effect of the background substitutions and α_2 the effect on the other 15 REs. We fit this
model as before and obtained parameter estimates of $\alpha_1 = 0.43$ (95% CI: 0.19–0.76) and $\alpha_2 =$
0.010 (95% CI: 0.0028–0.010), corresponding to fold-increases in affinity of 2.3 (95% CI: 1.3–
5.2) on ERE and 99 (95% CI: 99–361) on other REs.

Code availability

Scripts for analysis are available at www.github.com/JoeThorntonLab/RH-RE_scanning.

References

1. Wagner, G. P. & Altenberg, L. Perspective : Complex Adaptations and the Evolution of Evolvability. *Evolution* **50**, 967–976 (1996).
2. Stoltzfus, A. & Yampolsky, L. Y. Climbing Mount Probable: Mutation as a Cause of Nonrandomness in Evolution. *J. Hered.* **100**, 637–647 (2009).
3. Hodgins-Davis, A., Duveau, F., Walker, E. A. & Wittkopp, P. J. Empirical measures of mutational effects define neutral models of regulatory evolution in Saccharomyces cerevisiae. *Proc. Natl. Acad. Sci.* **116**, 21085–21093 (2019).
4. Gould, S. J. & Lewontin, R. C. The Spandrels of San Marco and the Panglossian Paradigm: A Critique of the Adaptationist Programme. *Proc. R. Soc. B Biol. Sci.* **205**, 581–598 (1979).
5. Schluter, D. Adaptive radiation along genetic lines of least resistance. *Evolution* **50**, 1766–1774 (1996).
6. Arthur, W. The interaction between developmental bias and natural selection: From centipede segments to a general hypothesis. *Heredity* **89**, 239–246 (2002).
7. Maynard-Smith, J. *et al.* Developmental constraints and evolution. *Q. Rev. Biol.* **60**, 265–287 (1985).
8. Wake, D. B. & Larson, A. Multidimensional analysis of an evolving lineage. *Science* **238**, 42–48 (1987).
9. Fay, J. C. & Wittkopp, P. J. Evaluating the role of natural selection in the evolution of gene regulation. *Heredity* **100**, 191–199 (2008).
10. Dugand, R. J., Aguirre, J. D., Hine, E., Blows, M. W. & McGuigan, K. The contribution of mutation and selection to multivariate quantitative genetic variance in an outbred population of *Drosophila serrata*. *Proc. Natl. Acad. Sci.* **118**, e2026217118 (2021).
11. McGlothlin, J. W. *et al.* Adaptive radiation along a deeply conserved genetic line of least resistance in *Anolis* lizards. *Evol. Lett.* 310–322 (2018) doi:10.1002/evl3.72.
12. McKeown, A. N. *et al.* Evolution of DNA specificity in a transcription factor family produced a new gene regulatory module. *Cell* **159**, 58–68 (2014).
13. Raup, D. M. Geometric analysis of shell coiling: General Problems. *J. Paleontol.* **40**, 1178–1190 (1966).
14. Vermeij, G. J. Forbidden phenotypes and the limits of evolution. *Interface Focus* **5**, 20150028 (2015).
15. Deline, B. *et al.* Evolution of metazoan morphological disparity. *Proc Nat Acad Sci* E8909–E8918 (2018) doi:10.1073/pnas.1810575115.
16. Clark, J. W. *et al.* Evolution of phenotypic disparity in the plant kingdom. *Nat. Plants* (2023) doi:10.1038/s41477-023-01513-x.
17. Dawkins, R. *Climbing Mount Improbable*. (WW Norton & Company., 1996).
18. Grant, P. R. & Grant, B. R. *40 Years of Evolution: Darwin's Finches on Daphne Major Island*. (Princeton university press, Princeton, New Jersey, 2014). doi:10.5860/choice.52-0821.
19. Jablonski, D. Developmental bias, macroevolution, and the fossil record. *Evol. Dev.* 103–125 (2019) doi:10.1111/ede.12313.
20. Steppan, S. J., Phillips, P. C. & Houle, D. Comparative quantitative genetics: evolution of the Gmatrix. *Trends Ecol Evol* **17**, 320–327 (2002).
21. Fowler, D. M., Stephany, J. J. & Fields, S. Measuring the activity of protein variants on a large scale using deep mutational scanning. *Nat. Protoc.* **9**, 2267–2284 (2014).
22. Podgornaia, A. I. & Laub, M. T. Pervasive degeneracy and epistasis in a protein-protein interface. *Science* **347**, 673–677 (2015).
23. Wu, N. C., Dai, L., Olson, C. A., Lloyd-Smith, J. O. & Sun, R. Adaptation in protein fitness landscapes is facilitated by indirect paths. *eLife* **5**, 1–21 (2016).
24. Starr, T. N., Picton, L. K. & Thornton, J. W. Alternative evolutionary histories in the sequence space of an ancient protein. *Nature* **549**, 409–413 (2017).
25. Lite, T.-L. V. *et al.* Uncovering the basis of protein-protein interaction specificity with a combinatorially complete library. *eLife* **9**, e60924 (2020).

26. Kemble, H., Nghe, P. & Tenaillon, O. Recent insights into the genotype–phenotype relationship from
massively parallel genetic assays. *Evol. Appl.* **12**, 1721–1742 (2019).

27. Bulyk, M. L., Huang, X., Choo, Y. & Church, G. M. Exploring the DNA-binding specificities of zinc
fingers with DNA microarrays. *Proc. Natl. Acad. Sci.* **98**, 7158–7163 (2001).

28. Newburger, D. E. & Bulyk, M. L. UniPROBE: an online database of protein binding microarray data
on protein-DNA interactions. *Nucleic Acids Res.* **37**, D77–D82 (2009).

29. Anderson, D. W., McKeown, A. N. & Thornton, J. W. Intermolecular epistasis shaped the function
and evolution of an ancient transcription factor and its DNA binding sites. *eLife* **4**, e07864–e07864
(2015).

30. Wheeler, L. C. & Harms, M. J. Were Ancestral Proteins Less Specific? *Mol. Biol. Evol.* **38**, 2227–
2239 (2021).

31. Patwardhan, R. P. *et al.* High-resolution analysis of DNA regulatory elements by synthetic saturation
mutagenesis. *Nat. Biotechnol.* **27**, 1173–1175 (2009).

32. Hochberg, G. K. A. & Thornton, J. W. Reconstructing Ancient Proteins to Understand the Causes of
Structure and Function. *Annu. Rev. Biophys.* **46**, 247–269 (2017).

33. Park, Y., Metzger, B. P. H. & Thornton, J. W. Epistatic drift causes gradual decay of predictability in
protein evolution. *Science* **376**, 823–830 (2022).

34. Carroll, J. S. *et al.* Genome-wide analysis of estrogen receptor binding sites. *Nat. Genet.* **38**, 1289–
1297 (2006).

35. Watson, L. C. *et al.* The glucocorticoid receptor dimer interface allosterically transmits sequence-
specific DNA signals. *Nat. Struct. Mol. Biol.* **20**, 876–883 (2013).

36. Park, Y., Metzger, B. P. H. & Thornton, J. W. The simplicity of protein sequence-function
relationships. *Nat. Commun.* **15**, 7953 (2024).

37. Metzger, B. P. H., Park, Y., Starr, T. N. & Thornton, J. W. Epistasis facilitates functional evolution in
an ancient transcription factor. *eLife* **12**, (2024).

38. Gerber, S. Not all roads can be taken: Development induces anisotropic accessibility in morphospace.
Evol. Dev. **16**, 373–381 (2014).

39. Stadler, B. M. R., Stadler, P. F., Wagner, G. P. & Fontana, W. The Topology of the Possible: Formal
Spaces Underlying Patterns of Evolutionary Change. *J. Theor. Biol.* **213**, 241–274 (2001).

40. Psujek, S. & Beer, R. D. Developmental bias in evolution: evolutionary accessibility of phenotypes in
a model evo-devo system. *Evol. Dev.* **10**, 375–390 (2008).

41. Salazar-Ciudad, I. Why call it developmental bias when it is just development? *Biol. Direct* **16**, 1–13
(2021).

42. Maynard-Smith, J. Natural Selection and the Concept of a Protein Space. *Nature* **225**, 726–734
(1970).

43. Newman, M. E. J. & Girvan, M. Finding and evaluating community structure in networks. *Phys. Rev.*
E **69**, 026113 (2004).

44. Kimura, M. *The Neutral Theory of Molecular Evolution*. (Cambridge University Press, 1983).
doi:10.1016/B978-1-55938-802-3.50013-4.

45. Russo, F. & Williamson, J. Interpreting Causality in the Health Sciences. *Int. Stud. Philos. Sci.* **21**,
157–170 (2007).

46. Schuster, P., Fontana, W., Stadler, P. F. & Hofacker, I. L. From sequences to shapes and back: A case
study in RNA secondary structures. *Proc R Soc Lond B* **255**, 279–284 (1994).

47. Fontana, W. & Schuster, P. Shaping space: The possible and the attainable in RNA genotype-
phenotype mapping. *J Theor Biol* **194**, 491–515 (1998).

48. Alberch, P. Ontogenesis and Morphological Diversification. *Am. Zool.* **20**, 653–667 (1980).

49. Chipman, A. D., Arthur, W. & Akam, M. A Double Segment Periodicity Underlies Segment
Generation in Centipede Development. *Curr. Biol.* **14**, 1250–1255 (2004).

50. Salazar-Ciudad, I. & Jernvall, J. A computational model of teeth and the developmental origins of
morphological variation. *Nature* **464**, 583–586 (2010).

51. Fuqua, T. *et al.* Dense and pleiotropic regulatory information in a developmental enhancer. *Nature*
587, 235–239 (2020).

52. Arthur, W. & Farrow, M. The Pattern of Variation in Centipede Segment Number as an Example of
Developmental Constraint in Evolution. *J. Theor. Biol.* **200**, 183–191 (1999).

53. Harjunmaa, E. *et al.* Replaying evolutionary transitions from the dental fossil record. *Nature* **512**, 44–
48 (2014).

54. Dingle, K., Ghaddar, F., Šulc, P. & Louis, A. A. Phenotype Bias Determines How Natural RNA
Structures Occupy the Morphospace of All Possible Shapes. *Mol. Biol. Evol.* **39**, 1–11 (2022).

55. Rohner, P. T. & Berger, D. Developmental bias predicts 60 million years of wing shape evolution.
Proc. Natl. Acad. Sci. **120**, e2211210120 (2023).

56. Galupa, R. *et al.* Enhancer architecture and chromatin accessibility constrain phenotypic space during
Drosophila development. *Dev. Cell* **58**, 51-62.e4 (2023).

57. Ferrada, E. & Wagner, A. Evolutionary innovations and the organization of protein functions in
genotype space. *PLoS ONE* **5**, (2010).

58. Ciliberti, S., Martin, O. C. & Wagner, A. Innovation and robustness in complex regulatory gene
networks. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 13591–13596 (2007).

59. Matias Rodrigues, J. F. & Wagner, A. Evolutionary Plasticity and Innovations in Complex Metabolic
Reaction Networks. *PLoS Comput. Biol.* **5**, e1000613 (2009).

60. Gould, S. J. & Eldredge, N. Punctuated Equilibria : The Tempo and Mode of Evolution
Reconsidered. *Paleobiology* **3**, 115–151 (1977).

61. Alberch, P. & Gale, E. A. A developmental analysis of an evolutionary trend: digital reduction in
amphibians. *Evolution* **39**, 8–23 (1985).

62. Braendle, C., Baer, C. F. & Félix, M.-A. Bias and Evolution of the Mutationally Accessible
Phenotypic Space in a Developmental System. *PLoS Genet.* **6**, e1000877 (2010).

63. Phillips, A. M. *et al.* Binding affinity landscapes constrain the evolution of broadly neutralizing anti-
influenza antibodies. *eLife* **10**, 1–40 (2021).

64. Starr, T. N. *et al.* ACE2 binding is an ancestral and evolvable trait of sarbecoviruses. *Nature* **603**,
913–918 (2022).

65. Ord, T. J. & Summers, T. C. Repeated evolution and the impact of evolutionary history on
adaptation. *BMC Evol. Biol.* **15**, 137 (2015).

66. Aakre, C. D. *et al.* Evolving New Protein-Protein Interaction Specificity through Promiscuous
Intermediates. *Cell* **163**, 594–606 (2015).

67. Poelwijk, F. J., Kiviet, D. J., Weinreich, D. M. & Tans, S. J. Empirical fitness landscapes reveal
accessible evolutionary paths. *Nature* **445**, 383–386 (2007).

68. Anderson, D. W., Baier, F., Yang, G. & Tokuriki, N. The adaptive landscape of a metallo-enzyme is
shaped by environment-dependent epistasis. *Nat. Commun.* **12**, 3867 (2021).

69. Lewontin, R. C. Four complications in understanding the evolutionary process. in *SFI Bulletin* vol. 18
(2003).

70. Maclean, C. J. *et al.* Deciphering the Genic Basis of Yeast Fitness Variation by Simultaneous
Forward and Reverse Genetics. *Mol. Biol. Evol.* **34**, 2486–2502 (2017).

71. R.D. Gietz & R.A. Woods. Yeast Transformation by the LiAc/SS Carrier DNA/PEG Method. in
Yeast Protocol, W. Xiao, Ed. 107–120 (Humana Press, Totowa, NJ, 2006).

72. Scanlon, T. C., Gray, E. C. & Griswold, K. E. Quantifying and resolving multiple vector
transformants in S. cerevisiae plasmid libraries. *BMC Biotechnol.* **9**, 95 (2009).

73. Mir, K., Neuhaus, K., Bossert, M. & Schober, S. Short Barcodes for Next Generation Sequencing.
PLOS ONE **8**, e82933 (2013).

74. N.A. Joshi & J.N. Fass. Sickle: A sliding-window, adaptive, quality-based trimming tool for FastQ
files. (2011).

75. J. Zhang, K. Kobert, T. Flouri, & A. Stamatakis. PEAR: A fast and accurate Illimuna Paired-End
reAd mergeR. (2015).

76. Cock, P. J. A. *et al.* Biopython: Freely available Python tools for computational molecular biology
and bioinformatics. *Bioinformatics* **25**, 1422–1423 (2009).

77. Jerome Friedman *et al.* glmnet: Lasso and Elastic-Net Regularized Generalized Linear Models.

78. Csardi, G. & Nepusz, T. The igraph software package for complex network research. *InterJournal*
Complex Systems, 1695 (2006).
79. Bastian, M., Heymann, S., & Jacomy, M. Gephi: an open source software for exploring and
manipulating networks. *Proc. Int. AAAI Conf. Web Soc. Media* **3**, 361–362 (2009).
80. Gillespie, J. Molecular Evolution Over the Mutational Landscape. *Evolution* **38**, 1116–1129 (1984).
81. Mccandlish, D. M. & Stoltzfus, A. Modeling Evolution Using the Probability of Fixation: History
and Implications. *Q. Rev. Biol.* **89**, 225–252 (2014).

1034

Acknowledgements: We thank Yeonwoo Park and Brian Metzger for advice throughout the
project and all Thornton Lab members for comments on the manuscript. This work was
supported by the University of Chicago's Research Computing Center, Cytometry Core, and
Genomics Core. Funding was provided by the National Institutes of Health grants
R01GM131128 (J.W.T.), R01GM121931 (J.W.T.), R35GM14533601 (J.W.T.), an NSF
Graduate Research Fellowship (J.E.J.P.), and a Rosemary Grant Award from the Society for the
Study of Evolution (S.H.A.).

Author contributions: All authors conceived the project; S.H.A. and J.E.J.P. performed
experiments and analyzed data; all authors interpreted results and wrote the manuscript.

Competing interests: The authors declare that they have no competing interests.

Additional information: Supplementary information is available for this paper. Correspondence
and material requests should be addressed to J.W.T (joet1@uchicago.edu).

1047

Extended Data Figure 1 | DBD library construction and sorting. a, Design of the DBD
expression vector used for DMS. The SR DBD is fused to an N-terminal *S. cerevisiae* Gal4
Activation Domain. Its expression is under control of a bidirectional pGAL1/10 promoter, which
simultaneously drives mCherry expression to select cells that maintain the plasmid during
sorting. **b**, Design of DBD library oligos. NNS codons (pink) were used to generate all possible
combinations of amino acid mutations at the four RH scanning sites (marked as X in the amino
acid sequence). For each background (AncSR1, left; AncSR2, right), we synthesized 16 libraries,
each with a unique set of synonymous barcode mutations at five codons (purple, Supplementary
Table 1), which allows each to be associated with one RE strain. BsaI sites (orange) were used
for Golden Gate assembly into the pDBD2.1 backbone. **c–e**, Validation of the RE reporter
strains. GFP fluorescence was measured by flow cytometry in each strain in the presence
(+DBD) or absence (–DBD) of a universally high-affinity DBD variant (AncSR1+GGKA+11P,
[29]). In each row, the left peak corresponds to autofluorescence from cells that do not express

GFP, either due to lack of DBD binding or loss of the DBD expression plasmid; the right peak
corresponds to cells that are expressing GFP in response to DBD-RE binding. "FS mut" denotes
strains with mutations in the flank/spacer regions of the RE that correct anomalous expression
patterns shown in red (see Supplementary Methods). Red strains were not used in the final DMS
experiment. Experiments were conducted on the same day within each panel. **c**, Fluorescence in
the presence of high-affinity DBD. **d**, Fluorescence in the absence of DBD expression plasmid.
e, Fluorescence in the CC and GA FS mut strains, with the ERE strain included as a negative
control. **f–g**, Sorting gates used for DMS. **f**, Enrichment sort gates. Homogeneous single cells
were first selected by gating on FSC-A vs. SSC-A and FSC-A vs. FSC-H (top). Plasmid
retention was then selected for by gating on mCherry expression (PE-Texas Red-A, bottom left).
Finally, cells were sorted into GFP+ (P4) and GFP– (P5) populations (bottom right). The
boundary between the GFP+ and GFP– gates was drawn to have a slope of 1.5 on a log-FSC-A
vs. log-GFP (FITC) scale so that populations were sorted by GFP expression relative to cell
volume. **g**, Binned sort gates. Gates P1–P3 were drawn as in **C**. Cells were then sorted into four
GFP bins, which were drawn to have roughly equal heights (P5–P7). The boundaries between
GFP gates were again drawn to have a log-log slope of 1.5.
1077

1078

Extended Data Figure 2 | DMS data cleaning. a, Curves show characteristics of the binned sort dataset as a function of the read count threshold used to retain protein-RE complexes for further analysis (*x*-axis). Black, standard error of F (SE, left axis); red, complexes retained, expressed as a fraction of the number of complexes in the binned sort (right axis); gold, fraction of complexes retained that have SE ≤ 0.1 (right axis). We used a read count threshold of 27 (vertical dashed line), at which ≥95% of complexes have SE ≤ 0.1 (horizontal dashed line). **b**, Correcting and filtering estimates of F from the binned sort. Left, correlation in F between replicates before correction. Pearson's r^2 is shown for all complexes, and for the subset of complexes with $F > -4$ in both replicates, which roughly corresponds to the boundary between active and inactive complexes (gray dotted lines). Red curves, I-splines fit using complexes with SE of $F < 0.1$. Center, correlation in F between replicates after correcting using the I-spline transforms. Right, correlation in F between replicates after filtering corrected variants for SE ≤ 0.1. **c**, Distribution of SE across all complexes in the binned sort after the I-spline correction. Complexes with SE > 0.1 were discarded. **d**, Read count distribution for complexes sequenced in the enrichment sort GFP– bin. Complexes were inferred to be inactive (gray) if they were not observed in the binned sort, but had high enough inferred cell count in the enrichment sort to have been detectable in the binned sort had they been at least minimally fluorescent (see Supplementary Methods). **e**, Correlations between estimates of F from flow cytometry (*x*-axis) and DMS (*y*-axes). Left *y*-axis (black points) shows estimates from isogenic strains that were spiked into the DMS libraries prior to the binned sort. Right *y*-axis (red points) shows estimates from complexes that were encoded in the DMS libraries.

1100

1101

Extended Data Figure 3 | Fluorescence imputation, GA fluorescence correction, and functional genotype classification. a, **b**, A generalized linear model that predicts the fluorescence of each protein-RE complex from its sequence was fit to the data for each background, using L2 regularization to address overfitting. **a**, Ten-fold cross-validation (CV) was used to identify the optimal L2 penalty parameter (λ). Red and black, mean and SE of the out-of-sample mean squared error (MSE) across the 10 folds. Initial range finding was performed using two folds (pink and cyan). Vertical line, λ that minimizes mean MSE. **b**, Genetic score versus observed F for the regularized RFA models. Red line, best-fit nonspecific epistasis function. For display, the distribution was discretized; colors show the number of variants in the interval defined by each square. Coefficient of determination (R^2) is reported for all complexes and for the subset of active complexes (above the gray line). **c**, **d**, Fluorescence correction for the GA strain. **c**, Affinity (K_A) versus F for a panel of DBD variants measured on ERE, SRE, and GA. Affinities, measured by fluorescence anisotropy on the three REs, all with the original flank/spacer sequence, were previously reported[12,29]. F was measured by flow cytometry in the

RE strains that were used for DMS, of which the ERE and SRE strains had the original
flank/spacer sequence, and the GA strain had a mutated flank/spacer sequence (see
Supplementary Methods, Extended Data Fig. 1c–e). Curves, best-fit sigmoidal function. The
same midpoint parameter was used for ERE and SRE (black); that for GA was independently
estimated (gray). Dashed lines, sigmoidal functions using 95% confidence intervals on the
midpoints. **d**, GA fluorescence correction based on the affinity effect estimated in **c**. Plots show
F before and after the correction. Dashed gray lines, mean boundary between active and null
variants. Red line, $y = x$. **e**, Bootstrap sampling strategy for classifying functional complexes with
model-inferred fluorescence. Plots show concatenated out-of-sample predictions versus observed
F across all 10 CV models. Bootstrap-sampled residuals from the interval within ± 0.1 units of a
complex's predicted F were used to test whether a variant with model-inferred F was not
significantly worse than the wild-type complex (dashed gray lines). An example for a complex
with inferred $F = -3.75$ (solid black line) is shown, with the bootstrap interval shown as a shaded
rectangle. Solid red line, $y = x$. **f**, Distribution of F across all 2,560,000 complexes in each DBD
background. Solid vertical lines, upper and lower bounds of fluorescence inferred from the RFA
models; dashed vertical lines, fluorescence of wild type complex (EGKA: ERE for AncSR1 and
GSKV:SRE for AncSR2). Colors indicate the source from which F was estimated. Darker colors
show functional variants, lighter colors nonfunctional. All "enrichment sort" complexes were
assigned to the lower bound of fluorescence, except for GA RE variants whose fluorescence was
corrected upward (**d**). Some model-predicted variants in the AncSR1 background have predicted
F below the reference but are classified as functional, because the bootstrap test accounts for the
AncSR1 RFA model's tendency to under-predict fluorescence (**e**, left). **g**, Bars show the number
of functional RH variants per RE per DBD background, colored by source of F estimate as in **f**.
1139

1140

Extended Data Figure 4 | Accessible new phenotypes after 3 substitution steps in the AncSR1 network. Bars show the distribution over every starting genotype in the AncSR1 main component. Dashed line, mean.

1141
1142
1143
1144

1145

Extended Data Figure 5 | Additional analyses for effects of background substitutions on DBD-RE affinity. a, Changes in phenotype across the AncSR1-to-AncSR2 transition. Bars represent the set of protein variants in AncSR2 that have different classes of phenotypes: specificity phenotypes that were encoded in the AncSR1 map (old specificity), specificity phenotypes not encoded in the AncSR1 map (new specificity), or promiscuous in AncSR2. Colored sections show the fraction of variants in each class whose functional category in the AncSR1 background was specific, promiscuous, or nonfunctional. **b**, Plots are the same as in Fig. 6A, but split into panels by RE. Blue points, protein-DNA complexes with significantly lower fluorescence in the AncSR2 background than predicted by the model; red, all other variants. Numbers at the bottom-right of each panel show the fraction of plotted variants with significantly lower than expected AncSR2 fluorescence. **c**, Amino acid frequencies at the RH variable sites among all complexes that are significantly more (left) or less (right) fluorescent in the AncSR2 background than predicted by the ERE-specific model in Fig. 6e. To test for significance in **b** and **c**, we tested whether their Bonferroni-corrected 95% CI of fluorescence was outside of the 95% CI of the model in both the AncSR1 and AncSR2 backgrounds.

1161

Extended Data Figure 6 | Robustness to alternative phenotype assignment methods. a, Global production distribution in the AncSR1 background, counting variants that bind specifically (colored bars) and promiscuously (white bars) to each RE. Dashed line shows the expected frequencies if the production distribution were isotropic. The bias, *B*, of the distribution and r^2 to the production distribution for specific variants (Fig. 2a) are reported. **b**, Same as in **a**, with phenotypes calculated using data from variants with fluorescence significantly higher than that of nonsense variants (active variants). **c**, Sequence space network for AncSR1 active variants. **d**, Bottom: Frequencies of specificity phenotypes within each genotype cluster in the AncSR1 active variant networks; the global production distribution is shown for comparison. Top: strength of phenotype bias (*B*) in each cluster. Red line, *B* of global production distribution. **e–g**, Same as in **b–d**, but with phenotypes calculated using only data from the binned sort experiment; protein-DNA complexes without experimental fluorescence measurements were

assumed to have null fluorescence. **h–o**, Same as in **a–g**, but for the AncSR2 background. Note that the active variant datasets are likely to be enriched for false positives due to misclassification of variants whose fluorescence is by chance slightly higher than the nonsense variant distribution. This may explain the high frequency of variants that do not share any mutational connections to other active variants. It may also explain the high frequency of CG-specific variants compared to the original classification scheme, since the CG yeast strain has a slightly higher null fluorescence level than most other strains (Extended Data Fig. 1c, d) and most CG-specific variants are unconnected in the active variant genotype networks. **p**, Number of protein variants in each network under different methods of phenotype assignment. "Functional" indicates the original method used in the main text; note that this yields the same number of protein variants as the "specific + promiscuous" method. **q**, Number of edges per node in each network, with the original phenotype classification method (functional) shown for comparison.

1189

Extended Data Figure 7 | Robustness to model of evolution using joint protein-DNA networks. a, AncSR1 protein-DNA coevolution network. Nodes represent functional protein-RE complexes, colored by the RE specificity of the protein genotype; colors are as in Fig. 2b and 4b. Promiscuous protein genotypes are represented by multiple nodes, one for each RE it binds. Edges connect complexes that can be interconverted by a single nucleotide change in the RE or the coding sequence of the protein. **b**, AncSR2 protein-DNA coevolution network. **c**, Bottom: Frequencies of specificity phenotypes within each genotype cluster in the AncSR1 (left) and AncSR2 (right) coevolution networks; the global production distribution (right-most column) is shown for comparison. Top: strength of phenotype bias (*B*) in each cluster. Red line, *B* of global production distribution. **d**, Bias (*B*) in evolutionary outcomes as a function of the length of evolutionary trajectories. Solid curves, mean *B* across starting genotypes in the protein (cyan) or coevolution (orange) networks. Dashed horizontal lines, *B* of the equilibrium distribution in each

network; dashed horizontal red line, global bias. Vertical dashed lines show the number of substitutions required for mean B to reach within 0.05 units of the equilibrium value within each type of network. The equilibrium distributions are more biased in the coevolution networks, and require more amino acid substitutions to be reached, because changes in protein genotype must occur between variants that can bind to the same RE sequence. **e**, Comparison between equilibrium outcome distributions of the protein-only evolution and protein-DNA coevolution networks in each AncSR1 (left) and AncSR2 (right) backgrounds. Pearson's r^2 between the two distributions are shown. Dashed line, $y = x$. **f**, Probability of conservation of each phenotype after 8 amino acid substitution steps in the protein vs. coevolution networks. **g**, Mean shortest amino acid distance between all possible pairs of phenotypes in the coevolution vs. protein networks, calculated as in Fig. 2g. Circles, AncSR1 networks, triangles, AncSR2 networks. Dashed line, y = x. **h**, Probability of evolving each specificity phenotype as a function of the number of amino acid substitutions away from EGKA:ERE in the AncSR1 (left) and AncSR2 (right) coevolution networks. In both backgrounds, conservation is more likely at short trajectory lengths than in the corresponding protein networks (Fig. 3g, 5f), but the relative likelihood of achieving each phenotypic outcome is similar. **i**, Distribution of the number of neighbors per genotype with distinct RH sequences in each type of network. Dots, means of distributions. **j**, Distribution of the fraction of neutral neighbors per node with distinct RH genotypes in each network. Dots, means of distributions.

1222

Extended Data Figure 8 | Robustness of RH mutation effects to uncertainty in ancestral reconstruction. Effects on ERE binding of all possible single amino acid mutations at the four variable RH sites in the background of the maximum *a posteriori* (MAP) wild type AncSR1 protein (*x*-axis), and in the background of the AltAll wild type AncSR1 protein, which has the second-most likely amino acid state at all sites at which the posterior probability of the MAP state is less than 0.8 (*y*-axis)[33]. Pearson's r^2 is shown.

a

```
                 6              16              26              36              46              56              66              76
A. AncSR1        CAVCSDYASGYHYGVWSCEGCKAFFKRSIQGHNDYMCPATNQCTIDKNRRKSCQACRLRKCYEVGMMKGGIRKDRR
B. AncChordate_ER  ·····························································································
C. AncVertebrate_ER ·····························································································
D. AncGnathostome_ER ·····························································································
E. AncGnathostome_ERa ·····························································································
F. AncEuteleostomi_ERa ·····························································································
G. AncActinopterygii_ERa ·····························································································
H. E_burgeri_ER    ··························T···············R···············V·-·····K
I. P_marinus_ER    ··························T···············R···············V·-·V···K
J. H_sapiens_ERa   ····N·················································
K. D_rerio_ERa     ························V···············R···············G
L. L_oculatus_ERa  ···········································V·····
M. C_milii_ERa     ·····························································································
N. H_sapiens_ERb   ·························I························V·C·S·RE·C
O. X_laevis_ERb    ·S·······················I····················F·····C·T·RE·C
P. D_rerio_ERb     ·························I····················C·L·R··S
Q. L_oculatus_ERb  ·························R··················C·T·RE·C
R. R_typus_ERb     ····N·F··············A·I··················SDT·R··S
S. C_milii_ERb     ····N·F··············A·I··················SDT·R··C
T. AncSR2          ·LI·G·E···C····LT·GS··V····AVE·QHN·L·AGR·D·I····I···N·P·········QA··VL·A-··LKK
```

b

Extended Data Figure 9 | Amino acid changes along the SR phylogeny. a, Amino acid alignment of extant vertebrate ERs and the MAP protein sequences for key ancestral nodes in the SR phylogeny[33]. The AncSR1 sequence is used as the reference to indicate amino acid changes; dots, same amino acid state as that in AncSR1; dashes, gaps; red circles, variable sites in DMS experiment; yellow circle, historical substitution (q36E) that likely contributed to the shift in the direction of the global bias away from ERE. **b**, Cladogram of SRs showing the number of substitutions that occurred along each branch. Letters, nodes shown in alignment in **a**; black nodes, AncSR1 and AncSR2; orange nodes, ancestral ER sequences identical to AncSR1. Branches and clades are colored according to their DNA specificity phenotype: purple, ERE-specificity; green, SRE-specificity.

13