Eye movement patterns under exposure to spatial disorientation illusions during simulated flight

Maya Harel (M.Sc.)[1, 2], Idan Nakdimon (M.Sc.)[1], Oded Ben-Ari (M.D, M.H.A)[1,3,4], Tom Schonberg (Prof)*[2,5]

[1]The Israeli Air Force Aeromedical Center, Tel-Hashomer, Ramat Gan, Israel

[2]Sagol School of Neuroscience, Tel Aviv University, Tel Aviv, Israel

[3]Department of Military Medicine, Faculty of Medicine, The Hebrew University of Jerusalem, Israel

[4]The Adelson School of Medicine, Ariel University, Ariel, Israel

[5]School of Neurobiology, Biochemistry and Biophysics, The George S. Wise Faculty of Life Sciences and Sagol School of Neuroscience, Tel Aviv University, Israel

*Corresponding author contact information – schonberg@tauex.tau.ac.il

Running Head – Eye movements during Spatial Disorientation

Manuscript type – Research Article

Word count – 3,700

Abstract

Objective

To identify eye movement patterns which are correlated with spatial disorientation (SD) events during flights in a flight simulator that induces SD.

Background

Spatial disorientation is one of the main causes of aviation mishaps. It results from illusions caused by misinterpreted vestibular or visual sensory cues, leading to an incorrect perception of an aircraft's position, attitude, or motion. SD prevention is crucial, as there is no objective tool to detect its occurrence.

Method

Eye movements of 45 participants (30 aircrew members, 15 cadets) were recorded using Tobii Pro Glasses 2 in a Gyro-IPT SD flight simulator. Illusions were either vestibular or visual. Gaze metrics—such as fixations, saccades (rapid gaze transfers), and visits—were compared between participants who experienced SD and those who did not. Statistical analyses were conducted to identify significant differences.

Results

Among 284 flight profiles, 136 SD occurrences were recorded (48%). During visual illusions, participants who more frequently checked the instrument panel were more likely to avoid SD. In contrast, during vestibular illusions, participants who examined the head-up display (HUD) more frequently had a lower probability of experiencing SD.

Conclusion

Efficient SD mitigation requires task-specific eye-movement strategies: mitigating visual illusions requires increased focus on the instrument panel, while mitigating vestibular illusions involves greater engagement with the HUD, challenging current standard instructions.

Application

These findings may inform training programs aimed at improving performance in high-risk SD profiles. Additionally, the results support the development of real-time SD alert systems to help mitigate or prevent SD-related incidents.

Keywords

1. Pilot, crew behavior
2. Situation awareness
3. Simulation and training
4. Eye movements, tracking
5. Flight displays

Precis

This study examined eye-movement patterns during flight in a flight simulator to identify mechanisms underlying spatial disorientation (SD) and strategies for SD avoidance. Findings reveal that to avoid visual illusions greater instrument focus is required,

56   whereas to avoid vestibular illusions, HUD engagement is beneficial. Results could be

57   applied to training programs to enhance flight safety and reduce SD incidents.

Background

Spatial orientation refers to an individual's ability to correctly perceiving their position and motion relative to the earth. This perception is achieved through visual and vestibular cues or intellectually constructed from focal visual, verbal, or other symbolic data (Gillingham, 1992). In aviation, spatial orientation pertains to an aircrew member's ability to recognize the aircraft's velocity, altitude, heading direction, rotation, and pitch angle. **Spatial Disorientation (SD)** occurs when these elements are misperceived, leading to errors in interpreting motion, position, or attitude due to absent or misleading cues, primarily from the visual or vestibular systems (Cheung & Hofer, 2003).

SD in flight is a leading cause of aircraft mishaps attributable to operator error (Gillingham, 1992). It is responsible for 5 to 10% of all mishaps, 90% of which are fatal (Antunano, 2005). A U.S. Air Force report found SD contributed to 30% of mishaps from 1990 to 1999 and 26% from 1999 to 2009 (Gibb et al., 2011). Similarly, SD was a factor in 20% of fatal mishaps in the U.S. Navy between 1990 and 2000 and 38% between 2000 and 2017 (Meeks et al., 2024). SD-related mishaps are frequently caused by undetected illusions, with 80% of incidents occurring without aircrew awareness (Gregg, 1955)

Several visual and vestibular illusions that can occur during flight pose a risk for SD in aircrew members. The main illusions are described as follows:

Visual illusions	
Black Hole Approach (BHA)	When approaching a runway without ambient visual cues, estimating the distance from the runway is difficult. The absence of global and local objects impairs depth perception, leading the aircrew member to overestimate the glide path and initiate an excessively steep descent. The miscalculation can result is a shallow approach, increasing the risk of a mishap (Gibb, 2007).
Ground-Light and Stars (GLS)	On a dark night, the ground lights and the stars may appear similar, creating confusion about the horizon line. Attempting to rely on external visual cues under this situation can distort altitude and pitch perception, leading to SD (Sánchez￼Tena et al., 2018).
False Horizon (FH)	A misleading external horizon can occur when the tilted ground line (e.g., a mountain slope) or a tilted cloud deck is mistakenly perceived as the true horizon. This illusion may cause an aircrew member to align the aircraft is based on the incorrect reference, leading to hazardous flight adjustments (Zhdan'ko et al., 2012).
Vestibular illusion	
Somatogravic Illusion (Dark Takeoff, DT)	A linear acceleration is misinterpreted as a pitch up movement (Davis et al., 2008). During take-off in reduced visibility, inertia creates the sensation of excessive pitch, leading to the aircrew member to unnecessarily lower the pitch angle, potentially resulting in a crush.
Somatogyral Illusion (SGI)	After a prolonged rotation, an attempt to straighten the aircraft can trigger a false sensation of a rotation opposite to the initial rotation. If an aircrew member reacts to this illusion, they may inadvertently re-enter the original rotation rather than correcting it (Boril et al., 2020).
Graveyard	Similar to the Somatogyral Illusion, after a prolonged spin, a corrective

Spin (GS)	straightening motion may stimulate the semicircular canals in a way that creates a sensation of rotation in the opposite direction. An attempt to correct this sensation can result in re-entering the initial spin, further exacerbating the disorientation (Guzy et al., 2008).
Coriolis	Head movements during rotation activate multiple semicircular canals, leading to disorientation and nausea (Houben et al., 2022).

High-fidelity simulators replicating SD-inducing conditions are recommended for SD

training (Cheung & Wong, 1998). An SD simulator is a flight simulator specifically

designed to induce and study spatial disorientation by integrating both visual and

vestibular stimuli: (1) Visual stimuli – The simulator includes screens that replicate the

aircraft's external environment, simulating real-world flight conditions. Additionally,

instrument panels are displayed, providing an objective representation of the aircraft's

state for the aircrew member. (2) Vestibular stimuli – The simulator incorporates motion

capabilities across the three principal axes of movement: pitch, roll, and yaw. The

aircrew member actively controls these movements using the throttle lever and flight

control stick, creating a dynamic training experience. An SD simulator integrates visual

and vestibular stimuli to replicate flight conditions, enhancing spatial disorientation

training. These simulators allow researchers to examine physiological and perceptual

changes that occur during SD events.

Studies have demonstrated that vestibular stimulation can influence heart rate and

respiration (Ilbasmis & Yildiz, 2017). Research has also shown that nighttime climbing

perception is affected by optokinetic and cervico-ocular reflexes (Tamura et al., 2016;

Maurice & Gioanni, 2004). Additionally, participants consistently perceive roll tilt during a

turn, yet their pitch perception during takeoff is slower and less consistent (Tribukait et

al., 2013). Furthermore, when no visual cues were available, attempts to straighten after

a prolonged turn often led to an involuntary return to the original turn direction (Nooij &

Groen, 2011).

Eye-tracking technology provides valuable insights into pilots' visual scanning patterns,

revealing their attention to critical instruments and external cues. *Fixations*, indicate

sustained attention (Valliappan et al., 2020; Unsworth et al., 2020) and may reflect

thorough cognitive processing of critical information (Just & Carpenter, 1976), or,

conversely, suggest cognitive overload (Hauland, 2019). Research has shown that

prolonged fixations on critical instruments were associated with better situational

awareness, as pilots dedicated more time to interpreting vital information needed for

safe flight operations (Goldberg & Kotval, 1999). Meanwhile, the *Visit* measure provides

a broader perspective on how often a pilot re-engages with complex visual stimuli,

suggesting the re-visited areas contain necessary information for decision making

(McLaughlin et al. 2017).

Saccades, or rapid eye movements between fixations, are crucial for active scanning

(Barfield & Furness, 1995). Frequent and large saccades indicate active situational

awareness, whereas reduced saccadic activity may signal a narrowed focus or cognitive

tunneling, where a pilot becomes overly reliant on a single instrument and may neglect

other critical information (Di Stasi et al., 2013). Additionally, mental effort (Joshi et al.,

2016), and unexpected sensory outcomes (Yokoi & Weiler, 2021).

Thus, eye-tracking can help us understand how aircrew members respond to situational

changes, such as SD illusions. Eye-tracking serves as both an objective physiological

response indicator—providing information about saccades, and visual movement

patterns—as well as a subjective indicator of experience, as fixation areas reveal which

areas of interest received attention. When examining visual movement patterns in pilots

using imagery that simulated the external visual environment of the cockpit, research

found that efficient movement across multiple areas of interest (AOIs), supported by

well-timed saccades, was crucial for maintaining situational awareness and preventing

SD (Thomas & Wickens, 2016). Similarly, prolonged fixations and visits on critical

instruments were associated with better situational awareness, as pilots spent more

time interpreting vital information needed to maintain safe flight operations (Goldberg &

Kotval, 1999).

Research conducted in the Korean Air Force examined differences between movement

patterns across flight instruments between a group where a verbal report of their flight

data was required and one that wasn't (Kang et al., 2021). That study found that the

verbal report group focused more on specific areas of interest (AOI) and had a higher

fixation frequency in the altitude measure in the Head-Up Display (HUD) in both types of

SD: Vestibular (Coriolis, GS) and Visual (GS). A Japanese Air Force study examined

eye movements during DT illusions found an automatic downward motion during the

pitch-up motion, a reflexive response meant to stabilize the visual input (Tamura et al.,

2016). When comparing experienced pilots and non-pilot participants in Vestibular

illusions (FH, SGI, and Coriolis), research found that experienced pilots were as

susceptible to SD as non-pilots (Bałaj et al., 2018).

Previous studies focused on a limited range of AOIs and illusions although a broader

range of scenarios is required to gain a more comprehensive understanding of SD

mechanisms. As Cheung (2013) claims, there is a need for renewed training that will

improve pilots' physical and mental performance during sensory conflict (Bob Cheung,

2013). Previous studies have also focused only on one aircrew member population.

Comparing experienced pilots to navigators and cadets can help us create better

learning paradigms to help prevent SD events. A broader model including different

populations can add a data-driven angle to identify underlying mechanisms.

Therefore, based on existing literature, we aimed to use eye-tracking to study the

mechanisms leading to SD events. Specifically, we aimed to compare eye-movement

patterns of aircrew members during both visual and vestibular illusions that have the

potential to create an SD event. To simplify the multiple variants of analysis we focused

on several factors: role (pilot and navigator), experience (cadets vs. experiences pilot)

and two main AOIs (HUD vs. instruments). This aimed to separate aircrew members

who experienced the SD effect from those who maintained correct situational

orientation. The overreaching goal was to lay the foundations to create more targeted

training to mitigate causes of SD events, using the objective tool of eye-tracking.

Methods

Study Design

We collected participants as part of their pre-existing training program. The participants were from two groups: flying cadets and experienced pilots. The study was approved by the Ethics Committee of the Israeli Defense Forces based on the ICH-GCP principles. This is an exploratory study and was not pre-registered, yet the procedure and sample size were predetermined by the ethics protocol.

Apparatus and measurement

Eye-tracking glasses: The Tobii Pro Glasses 2 Live View Wireless 100 were used to collect data, the parameters extracted from the glasses are elaborated in *Table 1* in the variables section.

SD trainer: We used The Gyro-IPT simulator (Environmental Tectonics Corporation, Inc., Southampton, US), located at the Israeli Air Force Aeromedical Center. This dynamic motion-based simulator has 3 degrees of freedom (roll ±25°, pitch ±25°, yaw 360°, and Heave =/- 25cm) and has an out-the-window visual display (120cm * 40cm 3 windows wide field of view). The flight instruments panel displayed in the cabin represents typical indicators that are applied in the pilots' aircraft (Military Institute of Aviation Medicine, Warsaw, Poland et al., 2023).

175 *Figure 1 - SD simulator*

177 (1*) Left: the SD simulator from the outside, which has a motion ability in 3 axes. (2)*
178 *Right: internal view of the SD simulator containing a visual external display and the*
179 *cockpit instruments panel.*

180 Participants

Aircrew members (n = 30): operational experienced high-performance aircrew
members: pilots and navigators, 30-50 years old (average 37.74±5.46 years, 1 SD),
with 12-22 years of experience.

Flying cadets (n = 15): pilot cadets and navigator cadets, in their second year of the
flying academy (average age 19.84±0.01 years, 1 SD). The training profile for cadets
included only 4 out of the 7 illusions, 2 visual (GLS, BHA) and 2 vestibular (SGI, and
Coriolis) in the aircrew member training profile, as elaborated **Error! Reference source**
not found..

Variables

The demographic parameters: Aircraft type (F - 15, F - 16, or T - 6), Aircrew member

role (navigator or pilot), and age were collected and are reported below.

Eye-tracking parameters were extracted from the eye-tracking glasses using the Tobii

Pro lab program. Flight parameters were extracted from the Gyro IPT program per flight

profile. The data consisted of the aeromodel parameters (*Table 1*).

Table 1 – Eye-tracking and SD trainer parameters

Apparatus	Key indicator	Unit
Eye-tracker	The total duration of fixation and visits in AOIs	ms
Eye-tracker	Number of fixations and visits in AOIs	
Eye-tracker	Time to first visit in AOI	ms
Eye-tracker	Number of saccades in AOIs and total.	
Eye-tracker	Velocity of saccades	(°)/s
SD trainer	Altitude	m
SD trainer	Velocity – ground speed, airspeed, North velocity, east velocity	m/s
SD trainer	Heading Position	(°)
SD trainer	Pitch, Roll, and Yaw positions	(°)
SD trainer	Pitch, Roll, and Yaw velocity	(°)/s

Procedure

Forty-five participants were recruited, thirty active aircrew members and fifteen flying

cadets. They were approached when they arrived at the SD simulator according to the

Israeli Air Force training regulations. Prior to the beginning of the training session, they

were presented with the study's aims and the procedure. They were explained the study

would not change the course of the mandatory training they arrived for and would only

add the eye-tracking glasses. Those who agreed to participate signed the informed

consent form. Then, a briefing session was provided for each participant about the

simulator and the training procedure. The participants were equipped with the Tobii

glasses, and a calibration process was conducted using the internal calibration

algorithm of the Tobii Glasses 2. During the calibration process, the participants had to

wear the Eye-tracking glasses while focusing on the center of the calibration target (

Figure 2 - calibration target) located about one meter from the participant

Figure 2 - calibration target

Then the participants entered the SD trainer and underwent the training procedure,

which included up to 8 flight profiles, elaborated in the supplementary material. All

subjects underwent the same illusions ordered as presented in the table. Each profile

lasted 1 to 3 minutes, between profiles the subjects remained inside the SD trainer

while the instructor debriefed about the SD illusion they experienced and for subjects

who showed SD response explained what they should have done differently. The

training procedure is summarized in *Figure 3*.

For each illusion the training profile, the SD window, criteria for identifying SD response,

and a visual example of the profile timeline are presented in the supplementary

material.

Figure 3 - training procedure

AOIs

An AOI was marked for each instrument, HUD element, and external aircraft display

(*Figure 4*):

Figure 4 – AOI list and mark over aircraft's display.

(A) *The list of AOIs divided by location (B) The T-6 aircraft display of instruments, it*

does not have an HUD display (C) The F-15 aircraft display of the instruments panel,

and the HUD. (D) The F-16 aircraft display of the instruments panel, and the HUD.

Statistical analyses

The initial analysis of the eye-tracking data was performed using Tobii Pro Lab (Tobii

PRO LAB), which produced the eye-tracking parameters outlined in *Table 1* for each

Time of Interest (TOI) and Area of Interest (AOI) (*Figure 5*). These parameters were

then entered into Python 3.11, where logistic regressions predicting SD response were

conducted. Given the multiple comparisons between SD and non-SD groups across

various illusions and AOIs, statistical significance was set at 0.05, and the Holm-

Bonferroni multiple comparison correction was applied.

The data analysis code and the row data are attached in the github repository:

https://github.com/mayahar/Harel-et-al

Figure 5 - Tobii PRO Lab

The Tobii PRO lab statistical analysis program: (A): Video recording from the eye-

tracker. (B): A reference snapshot for the compatible aircraft display. (C): Recording

timeline with the following marks (up to down) - relevant event marks, interval, SD time,

and TOI marks, gaze mapping from video to snapshot affinity.

Results

Demographics

Data from active aircrew members and flying cadets were divided into pilots and navigators. (*Table 2*).

Table 2 - participants demographics

Position	Navigator cadet	Pilot cadet	Navigator	Pilot	Navigator	Pilot
Aircraft	T-6		F-15		F-16	
n	5	10	3	4	10	13
Percentages out of entire sample	11%	22%	7%	9%	22%	29%
Average age	19.5	19.5	38.2	40.4	36.0	38.1
Number of flight profiles	20	42	20	28	81	92

Within the navigator cadets: all 5 participants performed the 4 profiles (summing to 20). Within the pilot cadets all 10 participants performed the 4 profiles and two performed BHA twice, (summing to 42 sorties). Within the F-15 navigators two of the participants performed all 7 profiles, one performed BHA twice, one performed DT twice, and one performed only 4 profiles (DT, SGI, GLS, and BHA) due to an eye-tracker malfunction

(summing to 20 sorties). Within F-15 pilots all four participants performed all 7 profiles

(summing to 28 sorties). Within F-16 navigators all ten participants performed the 7

profiles, seven performed BHA twice, and four performed DT twice (summing to 81

sorties). Within F-16 pilots All thirteen participants performed the 7 profiles, and one

performed DT twice (summing to 92).

Active aircrew members (n=30, average age = 37.74±5.46) performed all 7 profiles.

Some participants performed BHA and DT twice, and one did not complete the entire

procedure (as elaborated above in *Table 2*), summing to 221 sorties. Flying cadets

(n=15) performed only 4 of those profiles (two of each type: the visual illusions BHA and

GLS, and the vestibular illusions GLS and Coriolis) and two pilot cadets performed the

BHA profile twice, summing to 62 profiles. Overall, 283 profiles were used in analyses.

The flight experiment of the aircrew members was deduced from their age, as they

started pilot training at 18, the flight experience of the flying cadets when undergoing the

SD training during the course is 1.5 years.

Out of the 283 profiles, 136 profiles contained an SD response, i.e. it managed to

induce SD. One flying cadet did not experience SD in any sorties, 2 experienced SD in

1 sortie, the rest of the cadets experienced SD either 2 or 3 of the 4 sorties. All aircrew

members experienced SD during between 1 to 5 of the 7 sorties, with the majority

experiencing SD in 3 sorties. The percentages of flight profiles containing an SD

response was calculated per group (*Figure 6*).

Figure 6 - percentages for SD responses by aircraft and position.

		Active aircrew members				Cadets	
		Pilot F-15	Navigator F-15	Pilot F-16	Navigator F-16	Pilot T-6	Navigator T-6
DT		0% N=4	25% N=4	0% N=14	42.9% N=14		
SGI		0% N=4	33.3% N=3	7.7% N=13	20% N=10	70% N=10	60% N=5
BHA		75% N=4	100% N=4	64.3% N=14	88.2% N=17	58.3% N=12	60% N=5
GLS		50% N=4	66.7% N=3	69.2% N=13	80% N=10	0% N=10	0% N=5
Coriolis		100% N=4	50% N=2	46.2% N=14	60% N=17	80% N=12	80% N=5
GS		0%	0%	7.7%	0%		

	N=4	N=2	N=14	N=10		
FH	100% N=14	100% N=4	64.3% N=10	60% N=2		
Mean Sum	46.4% N=28	55% N=20	37.2% N=94	53.1% N=81	52.4% N=42	50% N=20

The percentage of SD responses and the total number of flight profiles for each

category is shown at the table below the graph.

We used logistic regression to examine the effects of experience and role within the

aircraft on the probability of experiencing SD, while accounting for participant-level

random effects. When tested separately, experience showed no significant effects—

neither when modeled as a continuous variable (years of experience) nor as a binary

variable (cadets vs. active aircrew members). A marginal effect was found for role, with

navigators experiencing more SD responses (p = 0.06). No significant interaction

between experience and role was observed.

Figure 7 – Effect of role on SD events

Eye-movement patterns

The eye gaze throughout the flight was analyzed to create heatmaps (see *Figure 8*).

Figure 8 - gaze heat map

Heat map of each cockpit display as calculated in the Tobii pro lab program using the

Tobii I-VT (Fixation) gaze filter, from left to right: T-6, F-15, F-16. On the left there is a

colorbar presenting a linear interpolation of number of fixations in each area.

The heat map showed that T-6 pilots and navigators, visit the instruments more than the

external view, particularly on the ball instrument that indicates the roll and pitch degree,

while F-15 and F-16 pilots visited the HUD more. The percentage of the intervals spent

fixating on the different AOI categories is brought in *Table 3*:

Table 3 - fixation time to AOIs

	Instruments	HUD	Outside view
Navigator	3.92%	28.26%	36.77%
Pilot	11.63%	25.20%	34.53%
Navigator Cadet	38.87%	-	9.24%
Pilot Cadet	40.14%	-	10.40%

SD window analyses

To simplify and focus our analyses we categorized the four illusions experienced by all participants into visual illusions (BHA and GLS) and vestibular illusions (SGI and DT). Additionally, we grouped the Areas of Interest (AOIs) into instrument AOIs and HUD AOIs and selected five key eye movement parameters for each AOI: number of visits, number of fixations, number of saccades, total duration of visits, and total duration of fixations.

Using these data, we conducted logistic regression analyses separately for vestibular and visual illusion sorties, and separately for eye movement parameters related to instrument and HUD AOIs. Each model predicted the likelihood of an SD response based on the eye movement parameter, the role of the participant (navigator vs. pilot), and their interaction, while accounting for participant-level random effects.

After correcting for multiple comparisons, two results remained significant: within the visual illusions, longer total durations of fixations and visits on the instrument AOIs were associated with fewer SD responses ($p < 0.001$ for both; Figure 9).

Figure 9 - Total duration of eye tracking parameter during visual illusiond in instruments AOIs

To further explore the interaction between illusion type (visual vs. vestibular), display type (HUD vs. instruments), and

*the eye tracking parameters, we ran a model testing a three-way interaction. The following effects were found (*A

main effect for total duration of fixations and total duration of visits, with longer durations

predicting increased SD responses (p < 0.001 for both).

A two-way interaction between eye tracking parameter and display type for total

duration of visits, total duration of fixations, number of visits, and number of fixations:

greater engagement with instrument AOIs was linked to higher SD occurrences, while

greater engagement with the HUD was linked to lower SD occurrences (p < 0.001 for

all).

A two-way interaction between eye tracking parameter and the illusion type for total

duration of visits, total duration of fixations, and number of visits: greater engagement

with the AOIs during vestibular was linked to higher SD occurrences, while in visual

illusions it was linked to lower SD occurrences (p < 0.001 for all).

A two-way interaction between display type and illusion type for total duration of visits,

number of visits, and number of fixations: increased HUD engagement was associated

with more SD events in visual illusions but fewer in vestibular illusions, whereas

instrument engagement showed the opposite pattern ($p < 0.001$ for all).

A triple interaction between the three arguments for total duration of visits, total duration

of fixations, number of visits, and number of fixations: combining all the effects showed

above ($p < 0.001$ for all).

Figure):

A main effect for total duration of fixations and total duration of visits, with longer

durations predicting increased SD responses ($p < 0.001$ for both).

A two-way interaction between eye tracking parameter and display type for total

duration of visits, total duration of fixations, number of visits, and number of fixations:

greater engagement with instrument AOIs was linked to higher SD occurrences, while

greater engagement with the HUD was linked to lower SD occurrences ($p < 0.001$ for

all).

A two-way interaction between eye tracking parameter and the illusion type for total

duration of visits, total duration of fixations, and number of visits: greater engagement

with the AOIs during vestibular was linked to higher SD occurrences, while in visual

illusions it was linked to lower SD occurrences ($p < 0.001$ for all).

A two-way interaction between display type and illusion type for total duration of visits,

number of visits, and number of fixations: increased HUD engagement was associated

with more SD events in visual illusions but fewer in vestibular illusions, whereas

instrument engagement showed the opposite pattern ($p < 0.001$ for all).

A triple interaction between the three arguments for total duration of visits, total duration

of fixations, number of visits, and number of fixations: combining all the effects showed

above ($p < 0.001$ for all).

Figure 10 - combined effect in total duration of visits

Discussion

In this study, we investigated eye-tracking patterns during exposure to the common

illusions that cause spatial disorientation (SD) during flight both visual and vestibular.

We aimed to identify which eye-tracking patterns could be related to SD avoidance.

Among 284 flight profiles performed by 30 active aircrew members and 15 flying cadets,

136 SD responses were identified across 7 flight profiles divided into **visual illusions**:

Black-Hole Approach (BHA), Ground-lights and Stars (GS), and False Horizon (FH);

and **vestibular illusions**: Dark Takeoff (DT), Somatogyral illusion (SGI), Coriolis, and

Graveyard Spin (GS). SGI illusion was found mainly in navigators, and DT only in

navigators. FH had a higher effect on F-15 aircrew than F-16, as it affected all F-15

aircrew members and only partially on F-16 aircrew members. GLS affected only active

aircrew members as opposed to cadets.

Using eye-tracking we found that visual illusions had a higher probability of causing SD

responses in participants who examined the instruments' panel measurements (altitude,

pitch, and ball) in greater frequencies than the HUD. In contrast, vestibular illusions

showed the opposite effect of having a higher probability of causing SD responses in

participants who examined the HUD in greater frequencies than the instruments' panel

measurements (altitude, pitch, and ball).

The study shows that no population avoided SD responses completely, as they affected

cadets (in 50% to 52.4% of the sorties) and active aircrew members (in 37.2% to 55% of

sorties), and GLS and BHA illusions even affected active pilots to a greater degree than

cadets. This result is similar to a previous study that showed that SD cues affected

pilots and nonpilots similarly ability to navigate the aircraft as per instructions (Bałaj et

al., 2018).

In a previous study performed on transportation pilots, HUD usage increased in critical

flight stages, particularly in difficult weather conditions when there was an increase in

the amount of time the pilot spends looking out of the front window (Dill & Young, 2015).

While the transportation pilots in that study used the HUD only 4% of the overall time,

combat pilots in the current research engaged heavily on the HUD, around 90% of the

overall time. This might be due to the need for high surrounding awareness as they

must constantly assess external threats thus requiring a dynamic use of external visual

information (Zhang et al., 2014 , Martins, 2016). This effect might have occurred

because the HUD allowed them to maintain visual contact with the external environment

while monitoring critical flight data (Cummings et al., 2016, Johnson et al., 2006), thus a

HUD usage is beneficial to reduce SD responds and is generally more relayed upon.

Our results across the two types of illusions presented to the aircrew members – visual

vs. vestibular, suggest that the HUD is irrelevant in visual illusions and those who visited

it numerous times for longer durations succeeded less in the event of instruments shut

down (GLS).

In the vestibular illusions SGI, an increased time spent examining the HUD and a

decreased time examining the instruments panel was related to better SD avoidance.

This finding is supported by a former comparison of how experts' and novices'

movement patterns change between easy and difficult routes during flight (Yang et al.,

2013). For the easy routes, experts spent less time scanning out the window (OTW) and

had shorter OTW dwellings. In contrast, for the difficult routes, experts appeared to slow

down their scan by spending as much time scanning out the window as the novices

while also having fewer Map fixations and shorter OTW dwellings. The HUD allows

aircrew members to scan OTW view while examining instruments, thus allowing them to

relay on the HUD in challenging flight paths.

The focus specifically directed to the HUD rather than generally examining the OTW is

supported by another study that found that experienced pilots had better situational

awareness performance and paid more attention to the HUD, but focused less outside

the cockpit when compare with novice pilots (Yu et al., 2016). Another supporting finding

to the increased need for HUD scanning to avoid SD during vestibular illusions comes

from research where Coriolis was found to elevate the use of the HUD compared to

other illusions and yielded a higher attentional load score (Kim et al., 2021).

422   Applications

Repeated spatial training using a simulated environment has been shown to be

beneficial to reduce SD, as was shown by a study conducted on neurologic patients

(Kober et al., 2013), as well as in pilots (Landman et al., 2018, Qiu et al., 2023). It was

found that training eye movements based on knowledge acquired about which eye

movement type was the most efficient for a specific task could improve performance in

said task (Shapiro & Raymond, 1989). Thus, implementing the knowledge and training

based on the current results can potentially improve performance in flight profiles with

high SD risk.

Another study (Peysakhovich et al., 2018) highlighted the need for real-time alert

systems based on differences in eye-movement patterns associated with spatial

disorientation (SD). Their study demonstrated that such systems, when installed in

aircraft during real flights, can help mitigate SD-related incidents by providing timely

alerts and reducing the likelihood of fatal responses.

Limitations

Although the procedure and sample size were pre-determined in the ethics protocols, the study was exploratory and not pre-registered, leading to multiple comparisons, which hinders our knowledge on replicability of our results. Using the knowledge gained in this study, future research could create a more accurate procedure to induce stronger effects and means to address them.

Conclusions

This study highlighted the role of targeted eye-movement behaviors underlying SD. Our results suggest that dynamic, broad scanning across instruments could reduce SD experienced in various flight illusions. Specifically, we found that flight profiles involving limited visual cues and thus a potential for visual illusions require higher attention to the HUD to avoid visually induced SD. This is opposed to prolonged turns, that induce a higher potential for vestibular illusions, required greater attention to the instruments panel to avoid SD. We also found that SD risk remains throughout all experience levels from flight cadets to experienced pilots. Training that involves lessons from this study regarding the importance of different instruments in different flight situations could improve SD resilience. Further research should broaden the findings beyond the scope of combat pilots only. In addition, examining whether training using the knowledge accumulated in the study could potentially improve SD resilience.

Key points

- **SD Responses**: 136 SD responses across 7 flight profiles from 45 subjects; specific effects varied by role and aircraft type.
- **Illusions and Eye-Movement**: Increased instruments focus helped mitigate SD risk in visual illusions (DT, GLS, FH). Increased HUD focus improved performance vestibular illusions (SGI, Coriolis):
- **HUD Usage**: Combat pilots rely heavily on HUD (90% usage). HUD is effective to mitigate SD during vestibular illusions but less so during visual illusions.
- **Experience and SD**: Experience does not prevent SD; active pilots can be as affected as cadets.
- **Applications**: Simulated training and efficient eye-movement strategies could reduce SD. Real-time alert systems may help mitigate SD during flights.
- **Conclusion**: Training focused on tailored eye-movement and instrument use could enhance SD resilience.

References

Antunano, M. J. (2005). Spatial Disorientation. https://rosap.ntl.bts.gov/view/dot/39656

Bałaj, B., Lewkowicz, R., Francuz, P., Augustynowicz, P., Fudali-Czyż, A., Stróżak, P.,

& Truszczyński, O. (2018). Spatial disorientation cue effects on gaze behaviour in

pilots and non-pilots. Cognition, Technology & Work, 1–14.

https://doi.org/10.1007/s10111-018-0534-7

Barfield, W., & Furness, T. A. (Eds.). (1995). Virtual environments and advanced

interface design. Oxford University Press.

https://doi.org/10.1093/oso/9780195075557.001.0001

Boril, J., Smrz, V., Blasch, E., & Lone, M. (2020). Spatial disorientation impact on the

precise approach in simulated flight. Aerospace Medicine and Human Performance,

91(10), 767–775. https://doi.org/10.3357/AMHP.5591.2020

Cheung, B, & Hofer, K. (2003). Eye tracking, point of gaze, and performance

degradation during disorientation. Aviat Space Environ Med, 74:11–20.

Cheung, B, & Wong, W. T. (1998). Recommendation to Implement GYRO-IPT for

Disorientation Training at CFSAT.

Cheung, Bob. (2013). Spatial disorientation: more than just illusion. Aviation, Space,

and Environmental Medicine, 84(11), 1211–1214.

https://doi.org/10.3357/asem.3657.2013

Cummings, M. L., Stimpson, A., & Clamann, M. (2016, January 4). Functional

requirements for onboard intelligent automation in single pilot operations. AIAA

Infotech @ Aerospace. AIAA Infotech @ Aerospace, Reston, Virginia.

https://doi.org/10.2514/6.2016-1652

Davis, J. R., Johnson, R., & Stepanek, J. (2008). Fundamentals of aerospace medicine.

Lippincott Williams & Wilkins.

Di Stasi, L. L., Antolí, A., & Cañas, J. J. (2013). Evaluating mental workload while

interacting with computer-generated artificial environments. Entertainment

Computing, 4(1), 63–69. https://doi.org/10.1016/j.entcom.2011.03.005

Dill, E. T., & Young, S. D. (2015, June 22). Analysis of Eye-Tracking Data with Regards

to the Complexity of Flight Deck Information Automation and Management -

Inattentional Blindness, System State Awareness, and EFB Usage. 15th AIAA

Aviation Technology, Integration, and Operations Conference. 15th AIAA Aviation

Technology, Integration, and Operations Conference, Reston, Virginia.

https://doi.org/10.2514/6.2015-2901

Gibb, R., Ercoline, B., & Scharff, L. (2011). Spatial disorientation: decades of pilot

fatalities. Aviation, Space, and Environmental Medicine, 82(7), 717–724.

https://doi.org/10.3357/asem.3048.2011

Gibb, R. W. (2007). Visual Spatial Disorientation: Revisiting the Black Hole Illusion.

Aviation, Space, and Environmental Medicine, 78, 801–808.

Gillingham, K. K. (1992). The spatial disorientation problem in the united states air

force. Journal of Vestibular Research, 2(4), 297–306. https://doi.org/10.3233/VES-

1992-2404

Goldberg, J. H., & Kotval, X. P. (1999). Computer interface evaluation using eye

movements: methods and constructs. International Journal of Industrial Ergonomics,

24(6), 631–645. https://doi.org/10.1016/S0169-8141(98)00068-7

Gregg, L. A. (1955). Fundamentals of Internal Medicine, 4th edition. Academic

Medicine, 30(1), 63–65. https://doi.org/10.1097/00001888-195501000-00017

Guzy, L. T., Albery, W. B., & Goodyear, C. (2008). Vibrotactile stimulators and virtual 3-

D audio countermeasures, training and motion sickness symptoms with a simulated

graveyard spin illusion. Journal of Vestibular Research, 18(5–6), 287–294.

https://doi.org/10.3233/VES-2008-185-605

Hauland, G. (2019). Measuring Team Situation Awareness by means of Eye Movement

Data.

Houben, M. M. J., Meskers, A. J. H., Bos, J. E., & Groen, E. L. (2022). The perception

threshold of the vestibular Coriolis illusion. Journal of Vestibular Research, 32(4),

317–324. https://doi.org/10.3233/VES-210073

Ilbasmis, S., & Yildiz, S. (2017). Respiratory and Pulse Changes Due to Vestibular

Stimulations in a Motion-Based Simulator. Aerospace Medicine and Human

Performance, 88(1), 48–51. https://doi.org/10.3357/AMHP.4534.2017

Johnson, N., Wiegmann, D., & Wickens, C. (2006). Effects of Advanced Cockpit

Displays on General Aviation Pilots' Decisions to Continue Visual Flight Rules Flight

into Instrument Meteorological Conditions. Proceedings of the Human Factors and

Ergonomics Society Annual Meeting, 50(1), 30–34.

https://doi.org/10.1177/154193120605000107

Joshi, S., Li, Y., Kalwani, R. M., & Gold, J. I. (2016). Relationships between Pupil

Diameter and Neuronal Activity in the Locus Coeruleus, Colliculi, and Cingulate

Cortex. Neuron, 89(1), 221–234. https://doi.org/10.1016/j.neuron.2015.11.028

Just, M. A., & Carpenter, P. A. (1976). Eye fixations and cognitive processes. Cognitive

Psychology, 8(4), 441–480. https://doi.org/10.1016/0010-0285(76)90015-3

Kang, Y., Lazaro, M. J., & Kim, S. (2021). Crosschecking through verbal reports under

spatial disorientation scenarios: Evidence from eye tracking metrics. International

Journal of Industrial Ergonomics, 86, 103202.

https://doi.org/10.1016/j.ergon.2021.103202

Kim, S., Kang, Y., Lazaro, M. J., & Yun, M. H. (2021). The Effects of Verbal Reports and

Spatial Disorientation on Attention: an Eye-Tracking Study.

Kober, S. E., Wood, G., Hofer, D., Kreuzig, W., Kiefer, M., & Neuper, C. (2013). Virtual

reality in neurologic rehabilitation of spatial disorientation. Journal of

Neuroengineering and Rehabilitation, 10, 17. https://doi.org/10.1186/1743-0003-10-

17

Landman, A., van Oorschot, P., van Paassen, M. M. R., Groen, E. L., Bronkhorst, A. W.,

& Mulder, M. (2018). Training pilots for unexpected events: A simulator study on the

advantage of unpredictable and variable scenarios. Human Factors, 60(6), 793–805.

https://doi.org/10.1177/0018720818779928

Lounis, C., Peysakhovich, V., & Causse, M. (2021). Visual scanning strategies in the

cockpit are modulated by pilots' expertise: A flight simulator study. Plos One, 16(2),

e0247061. https://doi.org/10.1371/journal.pone.0247061

Martins, A. P. G. (2016). A review of important cognitive concepts in aviation. Aviation,

20(2), 65–84. https://doi.org/10.3846/16487788.2016.1196559

Maurice, M., & Gioanni, H. (2004). Role of the cervico-ocular reflex in the "flying"

pigeon: interactions with the optokinetic reflex. Visual Neuroscience, 21(2), 167–180.

https://doi.org/10.1017/s0952523804042087

McLaughlin, L., Bond, R., Hughes, C., McConnell, J., & McFadden, S. (2017).

Computing eye gaze metrics for the automatic assessment of radiographer

performance during X-ray image interpretation. *International journal of medical*

informatics, 105, 11-21 . https://doi.org/10.1016/j.ijmedinf.2017.03.001.

Meeks, R. K., Anderson, J., & Bell, P. M. (2024). Physiology of spatial orientation. In

StatPearls. StatPearls Publishing.

Military Institute of Aviation Medicine, Warsaw, Poland, Lewkowicz, R., Bałaj, B.,

Nicolaus Copernicus University, Torun, Poland, Francuz, P., & John Paul II Catholic

University of Lublin, Lublin, Poland. (2023). The incidence of simulator sickness in

pilots and non-pilots exposed to spatial disorientation events in the gyro-ipt simulator.

The Polish Journal of Aviation Medicine, Bioengineering and Psychology, 26(4), 13–

25. https://doi.org/10.13174/pjambp.17.05.2023.02

Nooij, S. A. E., & Groen, E. L. (2011). Rolling into spatial disorientation: simulator

demonstration of the post-roll (Gillingham) illusion. Aviation, Space, and

Environmental Medicine, 82(5), 505–512. https://doi.org/10.3357/asem.2946.2011

Peysakhovich, V., Lefrançois, O., Dehais, F., & Causse, M. (2018). The

Neuroergonomics of Aircraft Cockpits: The Four Stages of Eye-Tracking Integration to

Enhance Flight Safety. Safety, 4(1), 8. https://doi.org/10.3390/safety4010008

Qiu, R., Gong, Y., Cao, Y., You, X., & Zhu, X. (2023). Influence of self-efficacy on male

military pilots' capability to handle special situations: a moderated mediation model.

Scientific Reports, 13(1), 10794. https://doi.org/10.1038/s41598-023-38009-9

Sánchez-Tena, M. Á., Alvarez-Peregrina, C., Valbuena-Iglesias, M. C., & Palomera, P.

R. (2018). Optical illusions and spatial disorientation in aviation pilots. Journal of

Medical Systems, 42(5), 79. https://doi.org/10.1007/s10916-018-0935-4

Shapiro, K. L., & Raymond, J. E. (1989). Training of efficient oculomotor strategies

enhances skill acquisition. Acta Psychologica, 71(1–3), 217–242.

https://doi.org/10.1016/0001-6918(89)90010-3

Tamura, A., Wada, Y., Shimizu, N., Inui, T., & Shiotani, A. (2016). Correlation of

climbing perception and eye movements during daytime and nighttime takeoffs using

a flight simulator. Acta Oto-Laryngologica, 136(5), 433–438.

https://doi.org/10.3109/00016489.2015.1132844

Thomas, L. C., & Wickens, C. D. (2016). Eye-tracking and Individual Differences in off-

Normal Event Detection when Flying with a Synthetic Vision System Display. SAGE

Publications. https://doi.org/10.1177/154193120404800148

Tribukait, A., Eiken, O., Lemming, D., & Levin, B. (2013). Use of an adjustable hand

plate in studying the perceived horizontal plane during simulated flight. Aviation,

Space, and Environmental Medicine, 84(7), 739–745.

https://doi.org/10.3357/asem.3470.2013

Unsworth, N., Miller, A. L., & Robison, M. K. (2020). Individual differences in lapses of

sustained attention: Ocolumetric indicators of intrinsic alertness. Journal of

Experimental Psychology. Human Perception and Performance, 46(6), 569–592.

https://doi.org/10.1037/xhp0000734

Valliappan, N., Dai, N., Steinberg, E., He, J., Rogers, K., Ramachandran, V., Xu, P.,

Shojaeizadeh, M., Guo, L., Kohlhoff, K., & Navalpakkam, V. (2020). Accelerating eye

movement research via accurate and affordable smartphone eye tracking. Nature

Communications, 11(1), 4553. https://doi.org/10.1038/s41467-020-18360-5

Yang, J. H., Kennedy, Q., Sullivan, J., & Fricker, R. D. (2013). Pilot performance:

assessing how scan patterns & navigational assessments vary by flight expertise.

Aviation, Space, and Environmental Medicine, 84(2), 116–124.

https://doi.org/10.3357/ASEM.3372.2013

Yokoi, A., & Weiler, J. (2021). Pupil-linked arousal modulates trial-by-trial motor

adaptation in humans. BioRxiv. https://doi.org/10.1101/2021.04.03.438075

Yu, C.-S., Wang, E. M.-Y., Li, W.-C., Braithwaite, G., & Greaves, M. (2016). Pilots'

visual scan patterns and attention distribution during the pursuit of a dynamic target.

Aerospace Medicine and Human Performance, 87(1), 40–47.

https://doi.org/10.3357/AMHP.4209.2016

Zhang, Y., Sun, Y., & Chen, Y. (2014). A framework for ergonomics design of transport

category airplane cockpit. Procedia Engineering, 80, 573–580.

https://doi.org/10.1016/j.proeng.2014.09.113

Zhdan'ko, I. M., Chulaevskiĭ, A. O., & Kovalenko, P. A. (2012). [Visual illusions and

moving horizon]. Voenno-Meditsinskii Zhurnal, 333(9), 52–62.

Biographies

Maya Harel (M.Sc.)[1, 2], Idan Nakdimon (M.Sc.)[1], Oded Ben-Ari (M.D, M.H.A)[1,3,4], Tom Schonberg (Prof)*[2,5]

1. Maya Harel
   a. Current affiliation
      i. The Israeli Air Force Aeromedical Center, Tel-Hashomer, Ramat Gan, Israel
      ii. Segol School of Neuroscience, Tel Aviv University, Tel Aviv, Israel
   b. Degree – M.Sc. in Neuroscience, obtained in 2025 from Tel Aviv University.
2. Idan Nakdimon
   a. Current affiliation
      i. The Israeli Air Force Aeromedical Center, Tel-Hashomer, Ramat Gan, Israel
   b. Degree – M.Sc. in Medicine Sciences, obtained in 2019 from Tel Aviv University.
3. Oded Ben-Ari
   a. Current affiliation
      i. Department of Military Medicine, Faculty of Medicine, The Hebrew University of Jerusalem, Israel
      ii. The Adelson School of Medicine, Ariel University, Ariel, Israel
   b. Degree – MD obtained in 2009 from Tel Aviv University

645   4. Tom Schonberg

646   a. Current affiliation

647   i. School of Neurobiology, Biochemistry and Biophysics, The George

648   S. Wise Faculty of Life Sciences and Sagol School of

649   Neuroscience, Tel Aviv University, Israel

650   b. Degree – PhD (Department of Neurobiology and Sagol School of

651   Neuroscience), obtained in 2010 from Tel Aviv University.