The *Pseudomonas aeruginosa* T3SS can contribute to traversal of an *in situ* epithelial

multilayer independently of the T3SS needle

Eric Jedel [a, b], Daniel Schator [a], Naren G. Kumar [a], Aaron B. Sullivan [a], Arne Rietsch [c], David J.

Evans [a, d], Suzanne M. J. Fleiszig [a, b, e #]

a. Herbert Wertheim School of Optometry & Vision Science, University of California,

Berkeley, CA USA

b. Graduate Program in Infectious Diseases and Immunity, University of California,

Berkeley, CA USA

c. Department of Molecular Biology and Microbiology, Case Western Reserve University,

Cleveland, OH USA

d. College of Pharmacy, Touro University California, Vallejo, CA USA

e. Graduate Groups in Vision Science and Microbiology, University of California,

Berkeley, CA USA

Running Title: T3SS needle-independent epithelial traversal by *P. aeruginosa*

[#] Corresponding author: Herbert Wertheim School of Optometry & Vision Science, University of

California, Berkeley, CA 94720, USA. Tel: +1 (510) 643-0990. Email: fleiszig@berkeley.edu

ABSTRACT

Multilayered epithelia lining our tissue surfaces normally resist traversal by opportunistic

bacteria. Previously, we developed a strategy to experimentally perturbate this resistance *in situ*

in the corneas of mouse eyes and used it to show that traversal of a multilayered epithelium by

Pseudomonas aeruginosa requires ExsA, the transcriptional activator of its type 3 secretion

system (T3SS). Here, we developed a novel strategy for quantitively localizing individual

traversing bacteria within the *in situ* multilayered corneal epithelium and explored contributions

of T3SS components. The results showed that T3SS translocon and T3SS effector mutants had

reduced epithelial traversal efficiency. Surprisingly, a Δ*pscC* mutant unable to assemble the

T3SS needle traversed as efficiently as wild-type *P. aeruginosa*, while a Δ*exsD* mutant

'constitutively on' for T3SS expression was traversal defective. Dispensability of the T3SS

needle for effector-mediated traversal was confirmed using a mutant lacking the T3SS operon

except the effector genes (Δ*pscU-L* mutant). That mutant reacquired the ability to traverse if

complemented with rhamnose-inducible *exsA*, but not if the effector genes were also deleted

(Δ*pscU-L*Δ*exoSTY*). Western immunoblot confirmed ExoS in culture supernatants of rhamnose-

induced *exsA*-complemented Δ*pscU-L* mutants lacking all T3SS needle protein genes. Together,

these results show that epithelial traversal by *P. aeruginosa* can involve T3SS effectors and

translocon proteins independently of the T3SS needle previously thought essential for T3SS

function. This advances our understanding of *P. aeruginosa* pathogenesis and has relevance to

development of therapeutics targeting the T3SS system.

IMPORTANCE

While the capacity to cross an epithelial barrier can be a critical step in bacterial pathogenesis, our understanding of mechanisms involved is derived largely from cell culture experimentation. The latter is due to practical limitations of *in vivo*/*in situ* models and challenge of visualizing individual bacteria in the context of host tissue. Here, factors used by *P. aeruginosa* to traverse an epithelial multilayer *in situ* were studied by: 1) leveraging the transparent properties and superficial location of the cornea, 2) using our established method for enabling bacterial traversal susceptibility, and 3) developing a novel strategy for accurate and quantitative localization of individual traversing bacteria *in situ*. Outcomes showed that T3SS translocon and T3SS effector proteins synergistically contribute to epithelial traversal efficiency independently of the T3SS needle. These findings challenge the assumption that the T3SS needle is essential for T3SS effectors or translocon proteins to contribute to bacterial pathogenesis.

INTRODUCTION

Pseudomonas aeruginosa is a common cause of opportunistic infections, able to cause disease in many body sites including the airways, GI tract, skin, urinary tract, eye, brain, heart and blood (1, 2). Susceptibility can be associated with wounds, burns, immunocompromise, chemotherapy, surgery, predisposing diseases (e.g. cystic fibrosis), and use of indwelling medical devices (3–5). In the eye, *P. aeruginosa* is a leading cause of corneal infection (microbial keratitis), estimated to cause vision loss in approximately 2 million people worldwide each year (6–8). A major risk factor for keratitis is contact lens wear (8–10) and *P. aeruginosa* is the most frequently isolated pathogen from lens-associated infections (6, 10). Recently, an outbreak of multidrug resistant *P. aeruginosa* infections was reportedly transmitted *via* simple artificial tears eyedrops, causing infections of the eye and beyond. These infections resulted in multiple instances of vision loss and several deaths, in some instances without contact lens wear or prior eye disease (11–13).

Type three secretion systems (T3SSs) are among various tools Gram-negative bacteria use to export factors across their otherwise impermeable cell membranes. A T3SS is additionally able to inject effector toxins across host cell membranes into the cytoplasm of a target host cell to alter its biology. *P. aeruginosa* encodes a single T3SS that has been shown to make significant contributions to virulence during acute infections of the cornea (14–17) and other body sites (18–22). The *P. aeruginosa* T3SS is encoded by 42 known genes encoding proteins of a multimeric needle core, a translocon pore, and one or more effector toxins (23). ExsA is the only known transcription factor for these genes. Effectors of the *P. aeruginosa* T3SS, like those of other Gram-negative bacteria, can have a multitude of effects on host cells, including modifying their biology to support an intracellular lifestyle (24–27).

All environment-exposed body surfaces are covered by multilayers of epithelial cells, that

when healthy resist traversal by *P. aeruginosa* and other opportunistic bacteria. This includes the

cornea of the eye, supremely capable of preventing bacterial colonization through a repertoire of

intrinsic defenses, some inherent within the corneal epithelium, others conferred by other factors

present at the ocular surface (28–34). The cornea's unique efficacy in this regard makes the

corneal surface our only environmentally exposed body surface devoid of a viable bacterial

microbiome (35, 36).

If *P. aeruginosa* does adhere to the cornea, which can occur after superficial injury,

additional barriers prevent it from traversing the multilayered epithelium to reach the underlying

stroma, access to which is required for the initiation of infectious pathology (i.e. keratitis) (28,

37–41). Thus, infection susceptibility requires alterations to the epithelial barrier beyond

superficial injury. While that can be accomplished by full thickness injury (42, 43), *P.*

aeruginosa corneal infection is most commonly associated with contact lens wear, which

predisposes the epithelium to bacterial traversal more subtly by mechanisms not yet well

understood (44, 45).

A plethora of studies have been done to explore how *P. aeruginosa* crosses a cultured

epithelial cell layer *in vitro*, using a variety of epithelial cell types. For example, airway

epithelial cells or MDCK cells were used to show roles for elastase, exotoxin A, type 4 pili,

flagella, and the T3SS (46–50). Our own *in vitro* studies using cultured corneal epithelial cells

revealed roles for proteases (41), type 4 pilus-associated twitching motility (51), and the T3SS

effector ExoU; the latter encoded by only a subset of *P. aeruginosa* strains (cytotoxic strains)

and which enables traversal by killing epithelial cells (52). Recently, a study using *in vitro* grown

organoids of human airway epithelial cells showed that both the T3SS and type VI secretion

system (T6SS) can contribute to *P. aeruginosa* "translocation" (traversal) of an epithelial barrier

via goblet cell invasion (53).

Thus, our current understanding of how *P. aeruginosa* (and other bacteria) traverse

susceptible epithelial layers is based largely on *in vitro* cell culture study outcomes wherein

bacteria and host cells are studied in isolation from other factors normally found in their

environment *in vivo*. Yet our prior studies have shown that *in vivo* factors modify how bacteria

interact with cells, including basement membranes, mucosal fluids, other cell types, nerves,

soluble factors, and environmental conditions specific to the tissue site (28, 32, 37, 39, 41, 54,

55). Other limitations of the *in vitro* literature on this topic are that most studies used epithelial

cell monolayers that differ from *in vivo* multilayers that contain layers with cells in multiple

states of differentiation. Further, transformed cells used by many studies are generally locked

into one state of differentiation irrespective of whether they can polarize correctly *in vitro* (e.g.

MDCK cells, HeLa cells).

Factors that have hindered development of *in vivo/in situ* models for studying epithelial

traversal by *P. aeruginosa* and other bacteria include deliberately bypassing epithelial and other

tissue barriers to enable infection (e.g. by wounding or bacterial injection through them).

Moreover, introducing bacteria *in vivo/in situ* can trigger inflammation that can break down

subsequent barriers to bacterial dissemination, reducing or even eliminating the need for bacteria

to contribute. Another obstacle to an *in vivo/in situ* study if individual bacteria need to be

localized is the challenge of visualizing them in the context of host tissue.

Our published studies have shown that pretreatment of *in vitro* grown corneal epithelial cells

with mucosal fluid (tear fluid) increased their resistance to *P. aeruginosa* adhesion, invasion, and

cytotoxicity, and increased their traversal resistance when grown as multilayers (39, 56, 57). This

was accompanied by profound changes to the epithelial cell's transcriptome (57). Tear fluid

exposure also changes gene expression in *P. aeruginosa*, including multiple genes involved in

virulence and survival, reducing its ability to traverse cultured corneal epithelial cells - without

directly impacting bacterial viability (39, 57, 58). Other *in vivo* factors that are not present in

epithelial cell cultures can modify epithelial-microbe interactions. In the corneal epithelium they

include nerves and immune cells, both of which contribute to maintaining epithelial cell

homeostasis while also directly recognizing and responding to microbes (28, 31, 32, 59–61).

Thus, while cell culture studies have provided a good starting point for understanding epithelial

barrier function and mechanisms by which *P. aeruginosa* traverses epithelial cells, *in vivo/in situ*

studies are warranted.

Some of our published studies have attempted to address this knowledge gap. In one study,

we enabled traversal susceptibility *in vivo* in mice by scratching through the corneal epithelium,

then waiting for the time point at which the cell multilayer was reestablished but still remained

permissive to bacterial traversal, which occurred 6 hours after scratching (38). Bacterial location

and inoculation was quantified in fixed and stained tissue sections. The results showed that

mutants lacking ExsA, the transcriptional activator for the T3SS, were unable to traverse the

epithelium (38). Since wound healing might have played into the outcome, we later performed a

second study enabling susceptibility using more gentle superficial injury, by gently blotting the

surface with a KimWipe™, then EGTA-treatment prior to inoculation (37). Using this strategy,

we found that bacteria traversing the epithelium did not penetrate past the basal lamina, which

when intact functions as a non-specific size exclusion filter (37, 41). This avoids bacterial entry

into the stroma and therefore subsequent pathology/inflammation that could complicate studies

of how bacteria overcoming epithelial barrier function. In a later study, the blot/EGTA model

was used to quantify *P. aeruginosa* traversal of murine corneal epithelium by combining 3D

confocal imaging to identify fluorescent bacteria with confocal reflectance microscopy to image

the non-fluorescent cornea (62). Results again showed that ExsA was required for *P. aeruginosa*

to traverse the corneal epithelium when the mouse was immunocompetent (62).

A limitation of both of our prior studies was that neither method accounted for the complex

shape of the cornea, which is curved rather than flat. The problem this presents is that at any

plane of imaging the back and front surfaces of the corneal epithelium are not in line. This made

it difficult to accurately determine depth of penetration across the population and ascertain

whether T3SS mutants were penetration defective, or simply less efficient at traversing the

epithelium.

Here, we report development of an advanced analytical imaging approach allowing for

precise metrics of location to be determined for all individual bacteria within the curved

multilayered epithelium *in situ*. Use of these methods showed that the T3SS impacts

penetration/traversal efficiency of *P. aeruginosa* rather than being absolutely required, with roles

played by both the T3SS translocon and T3SS effector proteins. Unexpectedly, results showed

that the T3SS needle was dispensable for this phenotype, and actually interfered when

constitutively expressed.

RESULTS

Transcription of the T3SS promotes *P. aeruginosa* traversal efficiency in the multilayered

corneal epithelium *in situ*. Our prior study showed that the transcriptional activator of the T3SS

(ExsA) contributed to *P. aeruginosa* traversal of the mouse eye's corneal epithelium (62). Here,

we leveraged recent advances in imaging and image analysis to obtain more detail about how

ExsA impacts bacterial location, taking into account the complexities of the cornea's shape while

more accurately assessing location of individual members of the traversing population.

As in our previous study, we used confocal reflectance microscopy (CRM) to image the

corneal epithelium and GFP-expression to detect bacteria. Raw confocal images were then

imported into Imaris software v9.9 for processing, the CRM signal pseudo-colored red (Fig. 1A).

As detailed in the methods section, the reflectance signal from the stroma was manually

excluded, and a "Surface" representing only signal from the epithelium was generated (Fig. 1B).

Apical and basal boundaries of the epithelium were then identified (Fig. 1C). Individual bacteria

were identified as a "Spot", creating objects in the image that each represented one population

member (Fig. 1D). The distance (in microns) from the apical and basal boundary of the

epithelium was measured for each bacterial "Spot", and the depth of penetration of each

bacterium was calculated as a percentage to provide its relative position in between the upper

and lower epithelial boundaries, with 0% being surface adherent and 100% meaning that the

bacterium reached the underlying basal lamina (Fig. 1E). Another metric quantified was the

proportion of the population able to penetrate beyond the 50% (midpoint) normalized to the

thickness of the epithelium in that region of the tissue (as in Fig. 2B).

The first experiment was conducted using *P. aeruginosa* strain PAO1F wild-type compared

to an isogenic Δ*exsA* mutant: corneas were incubated with 600 μl of ~10^{11} CFU/ml bacteria for 6

h (see Methods). The goal was to compare the outcome to our prior study that used more

rudimentary methods. For wild-type, the median traversal depth of individual bacteria was

47.29% of the epithelial thickness, the upper quartile reaching at least 76.02% depth (Fig. 2A).

Fig. 2B shows that 47.27% of the bacterial population penetrated beyond 50% thickness of the

epithelium. In contrast, median penetration depth for ∆*exsA* mutant was only 7.31% depth, the

upper quartile being at only 12.86% depth (Fig. 2A). Only 2.67% penetrated beyond the midway

point (50% depth) (Fig. 2B). While these findings confirmed the importance of the T3SS in

corneal epithelial traversal by *P. aeruginosa* shown in our earlier publications, they provided

more granularity on how it impacts bacterial location/distribution. The pattern that emerged

showed that the *exsA* mutants lacking the entire T3SS were not completely defective in ability to

penetrate beyond the epithelial surface, with significant variability among the population. While

wild-type penetrated more deeply than T3SS mutants over the 6 h time span of these

experiments, there was again significant variability with only a fraction fully traversing to the

level of the underlying basal lamina. For this reason, the capacity to penetrate the corneal

epithelium is referred to as <u>traversal efficiency</u> for the remainder of this manuscript.

Efficient traversal involves T3SS effectors and the T3SS translocon but does not require

the T3SS needle apparatus. To explore which ExsA-regulated T3SS components are required

for traversal efficiency, we compared wild-type *P. aeruginosa* to mutants lacking the T3SS

needle apparatus (∆*pscC*), the T3SS effectors (∆*exoSTY*), or the T3SS translocon proteins

(∆*popBD*).

As shown in Fig. 2A, both the T3SS effector mutants (∆*exoSTY*) and the T3SS translocon

mutants (∆*popBD*) were defective in traversal efficiency compared to wild-type reaching median

traversal depths of 11.16% and 10.91% respectively, comparable to the ∆*exsA* mutant.

Compared to the ∆*exsA* mutant, a greater proportion of the ∆*popBD* or ∆*exoSTY* mutant

populations traversed to over 50% depth; 30.18% and 19.27% respectively (Fig. 2B).

Surprisingly, the ∆*pscC* mutant lacking the T3SS needle, was the most traversal efficient of all of

the mutants examined, with median traversal at 45.24% depth, similar to wild-type.

These outcomes were surprising given that the T3SS needle is considered important for T3SS

functions. Thus, we additionally tested a Δ*exsD* mutant that is constitutively active for the T3SS

and therefore consistently expresses T3SS components (including the needle). This differs from

wild-type which instead requires induction by host cell contact or low calcium to express the

T3SS, and it is effectively the opposite of a Δ*exsA* mutant which consistently lacks T3SS

expression. Results with the Δ*exsD* mutant showed that constitutive expression of the T3SS

actually <u>interferes</u> with traversal efficiency, showing a median traversal depth of only 12.66%

(Fig. 2A), with only 21.42% of the population penetrating beyond 50% depth (Fig. 2B).

Representative examples of epithelium traversal by wild-type bacteria and T3SS mutants are

shown in Supplemental Fig. S1.

Since traversal efficiency data within these different populations was not normally

distributed, the Kruskal-Wallis test with Dunn's multiple comparisons was used to compare each

mutant to the Δ*exsA* mutant. Due to the very large sample sizes (10,000-60,000 datapoints each

group), all comparisons were found to be highly statistically significant ($P < 0.0001$), even when

median values and distributions appeared visually comparable (Fig. 2A). This was also true

when comparing mutants to wild-type. For this reason, the magnitude and the direction of the

differences are important to consider in evaluating the biological significance of the outcomes.

Taking those parameters into consideration, the data showed that traversal efficiency was

supported by combined efforts of the T3SS effectors and the T3SS translocon. They further

suggested that the T3SS needle was not required, and that constitutively expressing it and other

T3SS factors in the entire population actually <u>detracted</u> from traversal efficiency. Since this

result was unexpected, we used Sanger sequencing to re-confirm that the Δ*pscC* mutant used

contained a clean deletion of *pscC* (data not shown). In other controls, the T3SS-GFP reporter

pJNE05 (P*exoS* expression) was used to confirm that P*exoS* was activated upon EGTA-exposure

similarly to wild-type in each of the mutants shown to be defective in traversal in Fig. 2:

Δ*popBD*, Δ*exoSTY* and Δ*exsD* (Supplemental Fig. S2). Since P*exoS* expression is a reliable

surrogate for *exsA* expression, this outcome confirmed that the mutants were not defective for

T3SS induction, at least under the conditions used.

Complementation of ExsA in the Δ*pscU-L* T3SS operon mutant restored T3SS effector-

dependent traversal efficiency. The above results using T3SS needle mutants and constitutive

expression of the T3SS showed that the capacity to make the T3SS needle did not correlate with

efficient traversal despite involvement of the T3SS effectors. Here, we directly tested the

hypothesis that the effectors can function without the needle.

First, we generated a mutant lacking the 36 genes in the main T3SS operons, from Δ*pscU* to

Δ*pscL* (designated Δ*pscU-L*). This mutant still encodes the known T3SS effectors in PAO1

(ExoS, ExoT and ExoY) and their chaperones under endogenous promoters. Results showed the

Δ*pscU-L* mutant had low traversal efficiency (Fig. 3A, B) thereby phenocopying the Δ*exsA*

mutant. This was to be expected since the Δ*pscU-L* mutant lacks the gene encoding ExsA in

addition to genes encoding all of the T3SS machinery proteins, and while it does encode the

T3SS effectors, these are not transcribed, translated or released without ExsA.

Next, we tested if expressing the T3SS effectors in this mutant lacking T3SS machinery

related proteins could rescue traversal efficiency. This was done by complementing the mutant

with ExsA using a chromosome-integrating vector to induce expression of *exsA* under a

rhamnose-sensitive promoter. This vector was first tested in the background of a Δ*exsA* mutant

using a GFP reporter pJNE05 for P_{exoS} expression as a surrogate for *exsA* induction (as above,

Supplemental Fig. S2). Rhamnose addition induced GFP expression and it significantly

increased bacterial traversal compared to the no rhamnose control for the Δ*exsA*:P$_{rha}$*exsA* mutant

(Supplemental Fig. S3A, B). We next showed that Rhamnose-induction of *exsA* expression in the

Δ*pscU-L* mutant background increased median traversal to 11.09% depth versus the rhamnose-

induced vector control at 7.34% depth (Fig. 3A), and it also increased the percentage of the

population that traversed to at least 50% depth to 18.35% vs. 0.00 % (Fig. 3B). Showing that

partial restoration of traversal efficiency by Δ*pscU-L*:P$_{rha}$*exsA* depended on the T3SS effectors, it

was abrogated when the three exotoxin genes were also deleted (Median: 5.76% depth,

population over halfway 0.00%) (Fig. 3A, B). An *in vitro* control with pJNE05 confirmed

activation of P$_{exoS}$ in the Δ*pscU-L* mutant after *exsA* complementation with rhamnose addition

relative to a vector-complemented control: Mean +/- SD P$_{exoS}$ expression (FITC/OD$_{600}$) for

Δ*pscU-L*:P$_{rha}$*exsA* + rhamnose was 3397 ± 1033 relative fluorescence units versus 1567 ± 1235

for Δ*pscU-L*:P$_{rha}$Vector + rhamnose (P = 0.0061, Student's t-Test).

In Fig. 3A all comparisons between groups were significant (P < 0.0001 for most

comparisons, and P = 0.025 comparing Δ*pscU-L* [no rhamnose] *vs.* Δ*pscU-L*Δ*exoSTY*:P$_{rha}$*exsA*

[+ rhamnose], Kruskal-Wallis test with Dunn's multiple comparisons). The latter comparison

compared two mutants both unable to express T3SS effectors or any of the T3SS machinery-

related proteins, only one able to express the transcriptional activator ExsA (Fig. 3A, B). Thus,

ExsA expression in the absence of those other components had the least impact on traversal

efficiency. Taken together, these results show that ExsA-driven expression of *exoSTY* can

increase traversal efficiency of a mutant lacking all the T3SS machinery and thus one or more

effectors are necessary for traversal. However, it remains possible that ExsA regulates the

expression of a factor(s) outside of the main T3SS operon that is/are needed for effector-

mediated traversal.

The T3SS translocon contributes to traversal efficiency beyond roles of the T3SS effectors,

also independently of the T3SS needle. To explore if the T3SS translocon contributes to

traversal efficiency beyond the impact of the T3SS effectors, we studied deletion of *popB* in the

background of a Δ*exoSTY* mutant (already reduced in traversal efficiency compared to wild-

type). This led to a small but statistically significant further reduction in median traversal depth:

Δ*exoSTY* median at 10.91% depth vs. Δ*popB*Δ*exoSTY* median at 8.10% (Fig. 4A, Δ*exoSTY*

data reproduced from Fig. 2) (P < 0.0001, Kruskal-Wallis test with Dunn's multiple

comparisons). A much bigger reduction was seen in the percent of population traversing beyond

the 50% point: Δ*exoSTY* 19.27% vs. Δ*popB*Δ*exoSTY* 1.62%.

We next tried the opposite experiment, deleting a T3SS effector gene (*exoS*) in a T3SS

translocon mutant (*popB*). This also showed separable contributions for the T3SS effectors and

the T3SS translocon, the median depth for a Δ*popB* mutant at 12.30% depth vs. median for

Δ*popB*Δ*exoS* mutants at 9.42% depth (Fig. 4A) (P < 0.0001, Kruskal-Wallis test with Dunn's

multiple comparisons). Again, the difference became more obvious when assessing the percent

of the population penetrating beyond 50% depth; Δ*popB* 17.76% vs. Δ*popB*Δ*exoS* 5.86% (Fig.

4B). Taken together, these results showed that T3SS translocon proteins and T3SS effector

proteins can have additive impacts on traversal efficiency and specifically implicate ExoS and

PopB. However, difference in traversal beyond 50% depth shown between Δ*popB*Δ*exoS* 5.86%

and Δ*popB*Δ*exoSTY* 1.62% implicate a role(s) for ExoT and/or ExoY.

Having already shown that the T3SS needle was dispensable for effector mediated traversal,

we next asked if it was needed for the translocon's contribution. Thus, we mutated the translocon

pore proteins PopB and PopD ($\Delta popBD$ mutant) in the background of the T3SS needle mutant

($\Delta pscC$) already shown to be traversal competent (Fig. 2). This reduced traversal efficiency of

the needle mutant: $\Delta pscC$ mutant median 45.24% depth, population over halfway 43.84% vs.

$\Delta pscC\Delta popBD$ mutant median at 36.11% depth, population over halfway at 35.00% (Fig. 4A, B,

$\Delta pscC$ data reproduced from Fig. 2) (P < 0.0001, Kruskal-Wallis test with Dunn's multiple

comparisons). This showed that the T3SS translocon can contribute to traversal efficiency in the

absence of the T3SS needle.

All traversal results and strain characteristics are summarized in Table 1. Together, these

outcomes implicate two separate contributors to T3SS-mediated traversal efficiency: the T3SS

effectors (including ExoS), and the T3SS translocon pore (including PopB), each functioning

independently of the T3SS needle which can instead play an inhibitory role. Interestingly, the

<u>magnitude</u> of their individual contributions appear similar in magnitude, the $\Delta exoSTY$ and $\Delta popB$

mutants showing no significant difference in impact (Fig. 4).

Exotoxin S protein can be released by a T3SS operon mutant. Results in Figs. 2-4 show that

the T3SS effectors can contribute to traversal efficiency without the T3SS needle genes. Our

current understanding of T3SS transcription and expression is that in the presence of Ca^{2+} or

absence of host cell contact, low levels of T3SS expression occur due to repression of ExsA by

ExsD (63). Upon host cell contact or Ca^{2+} deprivation, the T3SS regulator ExsE is secreted,

leading to ExsA de-repression and T3SS transcription. *In vitro*, $\Delta pscC$ needle mutants do not

secrete ExsE and do not activate high levels of T3SS expression (64). To test if an alternate

mechanism of T3SS effector toxin expression might be operating in our *ex vivo* traversal model

or T3SS mutants, the Δ*exsA* or Δ*pscU-L* mutant carrying rhamnose-inducible *exsA* were tested

for ExoS protein expression *in vitro* using LB broth without EGTA calcium chelation to

specifically test rhamnose induction compared to controls (vector only or no rhamnose). After

growth in broth culture to 'mid-log' phase, bacterial pellet and supernatant samples were

collected and probed for ExoS *via* affinity-purified polyclonal antibody using SDS-PAGE and

Western immunoblot as described previously (64). ExoS protein was detected in the pellet and

supernatant of Δ*exsA*:P$_{rha}$*exsA* with rhamnose induction (as expected) and in Δ*pscU-L*:P$_{rha}$*exsA*

with inclusion of rhamnose but not in its vector control (Fig. 5). These data confirmed that ExsA-

driven ExoS expression and release can occur for mutants lacking needle protein genes.

DISCUSSION

Previously, we showed that the transcriptional activator of the T3SS (ExsA) is required for *P.*

aeruginosa to traverse susceptible multilayered epithelium in the context of live tissue, shown

using the eyes of immunocompetent mice (38, 62). Here, we explored the role of T3SS

components.

The cornea has several advantages over other tissues for this type of investigation being

readily accessible for manipulation or imaging, optically clear when healthy allowing imaging

without 2-photon microscopy or tissue clearing protocols and is relatively separated from other

body sites reducing complexity associated with cross-talk that can occur during infection. Its

epithelial surface also lacks a microbiome (35, 36), allowing a pathogen to be studied alone or in

combination with other bacteria in a controlled fashion. Use of subtle epithelial injury to enable

susceptibility to bacterial traversal, as done in this study using tissue paper blotting then EGTA

treatment, preserves the underlying basal lamina barrier, prohibiting bacterial access to the

vulnerable stroma and allowing study of epithelial traversal by bacteria without complexities

introduced by later steps of infection development. Thus, this mouse eye model for studying

epithelial traversal shares some advantages of *in vitro* cell/tissue culture models in reducing

complexity but benefits from *in situ* context.

A challenge of *in situ* tissue imaging for quantitively localizing bacteria is variations in tissue

shape that can make it difficult to ascertain position relative to upper and lower boundaries

across an entire tissue sample. Here, we developed methods that account for the cornea's curved

surface, variations in thickness and topography in different regions and resulting from cell

exfoliation, and any changes to these parameters after exposure to bacteria. This allowed for

more precise quantitation of bacterial location between the epithelial surface and underlying

basal lamina, separately accounting for all individual bacteria present in the epithelial layer.

To calibrate outcomes to our previously published methods, we first compared wild-type *P.*

aeruginosa to mutants lacking the entire T3SS *(exsA* mutants) which confirmed the critical role

for the T3SS in corneal epithelium traversal. However, the more accurate localization method

showed traversal was not an absolute quality, with both wild-type *P. aeruginosa* and *exsA*

mutants showing a distribution/spread for individual bacteria in their penetration rates. Thus, we

referred to "traversal efficiency" rather than "traversal" to describe subsequent outcomes.

When roles of specific T3SS components were examined, some results aligned with *in vitro*

cell culture study findings (65). For example, mutants lacking genes for known T3SS exotoxins

had reduced traversal efficacy compared to wild-type. However, our study showed that mutants

in T3SS translocon pore proteins also showed reduced traversal efficiency, with mutants lacking

both translocon and exotoxin proteins showing an even more profound defect similar to *exsA*

mutants unable to express any T3SS components. This implicated both the T3SS effectors and

the T3SS translocon proteins, their roles being additive.

Not all outcomes aligned with *in vitro* study findings. Mutants lacking the T3SS needle were

just as traversal efficient as wild-type *in situ.* This was a surprising and confusing result because

the T3SS needle is generally thought critical for T3SS function, including for secretion or

expression of the very same T3SS components we found required for traversal efficiency. Thus,

we used a mutant lacking the entire T3SS operon except the genes encoding the effector

exotoxins (Δ*pscU-L* mutant) (66) and induced effector expression by complemented it with

rhamnose inducible *exsA* (the transcriptional activator of all T3SS-related genes). This promoted

traversal efficiency in the mutant despite it lacking T3SS needle and other machinery related

proteins. Confirming it was T3SS effector gene expression driving the phenotype in the T3SS

operon mutant not than other genes outside the T3SS operon, traversal was no longer promoted

by *exsA* complementation when the effector genes were also mutated (i.e. Δ*pscU-L exoSTY*

mutant). Since Δ*pscU-L* mutants lack T3SS translocon genes additionally needed for full

traversal efficiency, it was not surprising that *exsA* complementation did not fully restore the

operon mutant to wild-type traversal efficiency. Indeed, efficiency approximated translocon

(*popBD*) mutants that similarly encode T3SS effectors but not translocon proteins, but that differ

in encoding T3SS needle proteins. Together these results confirmed that the T3SS needle protein

genes are not needed for T3SS effector-mediated traversal efficiency in this *in situ* model.

Since T3SS needle proteins are thought necessary for exporting T3SS effectors out of

bacteria and for injecting them into host cells this data raised two related questions. Can T3SS

effectors become extracellular without the needle, and if so, how do they impact traversal

without being injected into host cells?

Western immunoblotting confirmed the presence of ExoS in both supernatant and pellet of *in*

vitro grown Δ*pscU-L*:P$_{rha}$-*exsA* after rhamnose induction, showing that ExoS can become

extracellular without needle protein genes. While this might be an unusual feature of this mutant,

that would not explain results with other mutants that also show the needle is dispensable for

T3SS mediated traversal efficiency. A potential explanation is that bacterial lytic cell death

releases effectors, e.g. explosive cell lysis in *P. aeruginosa* (67), and a feature of other toxin

release by other bacteria (68, 69) and quite possible given the dense bacterial inoculum and

routine 'natural' bacterial cell death as a result of nutrient deprivation or from formation of

biofilms (67). In the context of our *in situ* assay, the presence of host-derived antimicrobial

peptides (29, 33, 70) could further contribute to bacterial cell death. Alternatively, T3SS proteins

are packaged into OMVs (Outer Membrane Vesicles) that are released into the extracellular

environment, as shown for *Salmonella enterica* T3SS-1 effectors and translocation proteins (71).

Indeed, some T3SS proteins were efficiently secreted *via* OMVs by mutants lacking a functional

T3SS needle, and effector toxins were shown delivered *via* OMVs into the cytoplasm of

epithelial cells where they enhanced pathogenesis of T3SS-deficient mutants (72). Pathogenic *E.*

coli (O157:H7) can also efficiently package T3SS translocation and effector proteins into OMVs

in the absence of essential T3SS needle proteins (73). Our own work showed that exposing *P.*

aeruginosa to ocular surface tear fluid or to purified lysozyme (a tear fluid ingredient) generates

OMVs that are cytotoxic to the corneal epithelium of mouse eyes *in vivo*. We also showed that

priming the corneal surface with these OMVs reduces defense against bacterial adhesion, and

that tear fluid/lysozyme triggered OMVs contain a ~48 kDa protein similar in size to ExoS (~49

kDa) and ExoT (~53 kDa) (74). Another potential pathway for delivering T3SS effectors into a

host cell without the T3SS needle could be from an intracellular location. In this regard, it is

interesting that the results specifically implicated PopB and ExoS in traversal efficiency as our

work has shown critical roles for both in the intracellular lifestyle of *P. aeruginosa* (24–27, 75).

Notwithstanding, even T3SS mutants are internalized by epithelial cells and can persist inside

intracellular vacuoles that can subsequently fail to contain them (24–26, 75–78). In other words,

T3SS factors can facilitate but are not essential for bacterial entry into a host cell's cytoplasm.

Inside epithelial or other cells, bacteria (including *P. aeruginosa*) can use factors beyond T3SS

factors to exert impacts on the host cell, and the host cell can detect intracellular bacteria and in

response alter its biology and/or induce programmed cell death. Others have shown that

extracellular ExoS itself can be internalized by eukaryotic cells, and activates host cell TNF-α

responses by triggering surface TLRs (79). While mechanisms might overlap with some of the

above, ExoS has also been shown to affect host cell function from an extracellular location (80).

Given these many possibilities, a separate study will be needed to determine how T3SS

effector mediated traversal occurs without the T3SS needle. Importantly, much of our

understanding of gene expression in *P. aeruginosa* has been done using *in vitro* methods, often

without host cells present. Given its many environmental sensors and regulators of gene

expression, regulation of the *P. aeruginosa* T3SS *in situ/in vivo* could differ vastly from what has

been shown *in vitro*. It is also likely to be context dependent/complex due to microenvironments,

and lack of homogeneity and synchrony. Indeed, the result showing that *exsD* mutants (which

differ from wild-type in constitutively expressing the T3SS) are equally as traversal defective as

mutants lacking the entire T3SS are particularly poignant and highlight the importance of

regulation. Further research on this topic would benefit from more *in situ* experimentation to set

the stage for well-designed *in vitro* studies that can then tease apart mechanisms for the

phenomena shown relevant to pathogenesis *in situ/in vivo*.

The T3SS translocon proteins (PopB/PopD) were found to play roles additional to the

contribution of the effectors. This could relate to their ability to form pores in host cell

membranes, which can damage a cell or otherwise alter host cell function. Indeed, PopB-PopD

complexes in isolation have been shown to function as a pore-forming toxin by allowing K^+

efflux, resulting in histone H3 modification and host cell subversion (81). A potential mechanism

for the additive impacts of translocon and effector proteins could be pore formation providing an

entry mechanism for effector delivery into the host cell. Whether the T3SS needle is also

dispensable for translocon-mediated traversal efficiency remains an open question that will

require further investigation, although it is suggested by the fact that needle mutants remain fully

efficient.

Another factor to consider in interpreting the data is that T3SS needle proteins of *P.*

aeruginosa and other Gram-negative bacteria are recognized by mouse (and human) cells

causing inflammasome activation (82, 83), which can help the cornea and other tissues

recognize, respond and clear *P. aeruginosa* during experimental challenge *in situ* (35, 84). In this

way, expressing T3SS needle proteins could be detrimental to bacteria *in situ* (even if not *in*

vitro) and using needle-independent mechanism for traversal could be advantageous. Other *in*

situ factors (absent *in vitro*) may also impact bacterial viability or growth during traversal and

thus data interpretation.

Providing additional insights into pathogenesis, traversing bacteria tended to be bimodally

distributed across the epithelium. Typically, there was a cluster in the apical region from 0-15%

traversed, few across the midway point, then a group from 50-80% traversed. This distribution

may reflect bistability of T3SS expression in *P. aeruginosa* populations (26, 85), and/or a host

defense barrier, possibly related to specialized junctions in the suprabasal region of the corneal

epithelium that have been shown to be a barrier to leukocytes (86). Also worth noting, the very

large sample sizes (10,000-60,000 datapoints for each group) provided such a powerful analysis,

that most comparisons yielded statistically significant differences even when they were small.

Thus, we considered the magnitude of the differences in central tendency (median) and data

spread (upper and lower quartiles) to draw conclusions about biological significance.

In conclusion, this study used novel and powerful image analysis tools and an *in situ* model

with pared-down complexity to focus on an early step in pathogenesis, traversal of a superficial

multilayered epithelium. The goal was to determine which T3SS components contribute. Results

showed roles for T3SS effectors, T3SS translocon proteins, and regulation of T3SS expression.

Differing from *in vitro* culture studies and challenging our general understanding about how the

P. aeruginosa T3SS supports infection pathogenesis, the T3SS needle was dispensable. Follow

up experiments confirmed that the T3SS effector ExoS could become extracellular in the absence

of the T3SS needle. How T3SS effectors and translocon proteins influence host cells to promote

traversal without the T3SS needle will require further work. Published knowledge about *P.*

aeruginosa and other Gram-negative bacteria suggest multiple possibilities (67, 72, 73, 87).

Outcomes of this study may or may not be applicable to how *P. aeruginosa* traverses

epithelial barriers other than the mouse corneal epithelium, but if so they might generally inform

efforts to develop therapies targeting the T3SS (88, 89). Importantly, the results highlight the

need to move beyond *in vitro* and cell culture studies when studying pathogenesis. At the same

time, they support complementary *in vitro* and cell culture experimentation that can further

reduce complexity when teasing apart details, and to study regulation in a more controlled

system.

The model and imaging methods we have developed enable a single early stage of infection

pathogenesis to be studied *in situ* in the absence of various confounders generally present in

infection models (e.g. inflammation), and they enable individual bacteria to be visualized across

the entire tissue with their location quantified. These methods could prove useful for studying

pathogenic strategies of other microbes also able to traverse epithelial layers during infection,

including those not considered to be eye pathogens.

MATERIALS AND METHODS

Bacteria. *Pseudomonas aeruginosa* strain PAO1F was used for all experiments. Mutants in

T3SS components were either obtained from in-house strain collections or generated using two-

step allelic exchange (90). For this, pEXG2 vectors containing overlapping upstream and

downstream regions of the genes of interest (excluding the ORF) were cloned and transformed to

E. coli SM10 (λ*pir*), which were then used as donor strains to deliver the vector to recipient *P.*

aeruginosa strains. Successful merodiploid colonies underwent sucrose counter-selection and

PCR verification of mutagenesis. For integration of rhamnose-inducible ExsA, Dr. Arne Rietsch

provided strains generated by Tn7 vector-mediated insertion and Flp-mediated excision of

antibiotic resistance markers (91). All bacteria, mutants and plasmids used are shown in Table 2.

In control experiments not shown, T3SS mutants did not show any growth defects relative to

PAO1 in trypticase soy broth culture at 37 $^{\circ}$C *in vitro*.

For all experiments, bacteria were streaked from glycerol stocks stored at -80 ºC, plated onto

trypticase soy agar (TSA) plates with appropriate antibiotics (see Table 2) and grown at 37 ºC

overnight. For the rhamnose-induction traversal experiments, bacteria were grown on TSA also

supplemented with rhamnose (2 %). For T3SS-induction growth curves with plasmid pJNE05,

bacteria were grown in trypticase soy broth (TSB) supplemented with gentamicin (200 μg/ml),

monosodium glutamate (100 mM), glycerol (1%) and EGTA (2 mM) or alternatively with

rhamnose (2 %) instead of EGTA as specified.

For the *ex vivo* murine model of traversal, one bacterial "lawn-covered" TSA plate was used

per bacterial strain per condition. After overnight growth at 37 ℃, bacteria were collected with a

sterile loop and carefully suspended into serum-free DMEM (Dulbecco's Modified Eagle

medium; Gibco). DMEM was supplemented with 2% rhamnose when indicated. OD_{600} was

measured, with A_{600} 1.0 ~ 4 x 10^8 CFU/ml used for calculating density. Suspensions of 1.0 x 10^{11}

CFU/ml were used for *ex vivo* incubations, and concentrations confirmed by plating serial

dilutions for viable counts.

***Ex vivo* murine model of traversal.** All procedures were approved by the Animal Care and Use

Committee of the University of California, Berkeley, an AAALAC accredited institution. For all

experiments, male or female C57BL6/J or mT/mG mice with tdTomato-labeled cell membranes

were used, 6 to 12 weeks old, and 3-6 eyes per experimental condition. Contralateral eyes from

the same mouse were used as controls when possible. All experiments were performed *ex vivo*,

as mice were euthanized by isoflurane inhalation prior to any further manipulations. After

euthanasia, each eye was lightly blotted with a Kimwipe (Kimtech), enucleated, then placed in

0.2μm filter-sterilized 1X Phosphate-Buffered Saline (PBS) in a 48-well dish for subsequent

steps. Eyes were rinsed three times in PBS, then incubated in a 0.1M solution of EGTA at room

temperature for 1 h. Eyes were rinsed three more times in PBS, then completely submerged in a

600 μl suspension of bacteria, prepared as described above, and incubated at 37 ℃, 5 % CO_2 for

6 h, consistent with our previous studies (62, 92). Following this incubation, eyes were rinsed

with PBS three times to remove excess non-adherent bacteria before fixation in 2 %

paraformaldehyde (PFA) overnight at 4 ℃. Fixed eyes were then whole mounted on a glass

cover slip with Loctite superglue with the central cornea facing up before submersion in PBS and

imaging.

Imaging. Whole-mounted fixed corneas were imaged using an Olympus Fluoview FV1000

upright laser scanning confocal microscope with a water immersion 20X objective (NA = 1.0).

For each eye, three to five non-intersecting fields of view on the central cornea were selected for

imaging with 3x optical zoom (60x total magnification) at 1024 x 1024 resolution and 0.8µm

step size. Fluorescent bacteria were imaged using FITC (488/509) and/or RFP (555/580)

channels, and corneal cells were imaged using an 640nm excitation channel for confocal

reflectance microscopy (CRM) (92, 93).

Image analysis. Images of corneas were processed on Imaris Software (v 9.9) as follows. First, a

"surface object" was manually drawn using the reflectance channel to select the total signal

underneath the corneal basement membrane. This object represented the corneal stroma and was

used to mask the remaining reflectance signal. This created a new channel with signal from only

the epithelium, which was used to automatically generate a new surface object.

From the created Epithelium surface object, a distance transformation was performed, and

the basement membrane and apical surface were identified as two new surface objects ~1µm

away from the Epithelium. Variability in the created objects due to differing corneal reflectance

intensity was minimized through use of automatic thresholds, and manual surveillance of the

entire process ensured consistency in appearances. Next, bacteria were identified through the

'Spots' tool, using an XY size of 1.25 µm in the appropriate channel. Parameters for Spots

creation were assessed using an automatic threshold, adjusted manually to reduce false positive

results (e.g., Spots where there were no bacteria).

For each field of view, the following were measured: total volume and maximum thickness

of epithelium, total number of bacteria, and distance of each bacterium from apical surface and

basement membrane. Percent depth of traversal for each bacterium was calculated using the

formula: % Depth = Distance to Apical Surface / (Distance to Apical + Distance to Basement

Membrane) * 100 %. An inverted cumulative histogram was also generated from the % depth for

all bacteria in a condition with 5µm bins. The percent of the total population remaining deeper

was graphed, so 100 % of a population remained at 0 % traversal depth and 0 % remained at 100

% depth.

Western immunoblot. Bacteria were inoculated in liquid cultures of LB media (containing 200

mM NaCl, supplemented with 10 mM $MgCl_2$) and grown with shaking overnight at 37 ℃.

Overnight cultures were diluted 1:300 into fresh media, grown for 2 h, then 1 ml of culture was

added to 1 ml pre-warmed media with 2 % rhamnose and 1 ml of culture was added to 1 ml pre-

warmed media as a control. The bacteria were grown for another hour before the OD_{600} was

measured for each tube, which were then placed on ice. 1 ml of each sample was pelleted, and

the entire supernatant was added to a fresh tube containing 10 % trichloroacetic acid for

precipitation. After 10 min on ice, two washes with acetone were conducted before drying out

the precipitate at room temperature. The cell pellet and supernatant precipitate were both

resuspended in the same volume of 1x SDS Buffer (Laemmli Solution [Bio-Rad], 1 mM DTT) to

create a concentration of 4 x 10^9 CFU/ml, or an OD_{600} of 10 – typically 60-80 µl. Samples were

boiled at 90 ℃ for 10 min to denature proteins before brief centrifugation.

For SDS-PAGE, 50 µl of each sample and Precision Plus Protein Standards (Bio-Rad) were

then applied to a 4-20 % Protean TGX Stain Free Gel (Bio-Rad) before separation at 200V for

35 min. Gels were imaged on the Bio-Rad Gel Dock, then transferred to nitrocellulose

membranes using the Mixed MW program on the TransBlot Turbo Transfer System. Membranes

were blocked with EveryBlot buffer (Bio-Rad) before incubation overnight at 4 ℃ with an

affinity-purified Rabbit polyclonal anti-ExoS primary antibody (from Dr. Arne Rietsch, 1:5000).

After washing 3x with TBS-T (0.1 % Tween-20 in 1x TBS Buffer) for 5 min each, membranes

were then incubated for 1 h at RT with Goat anti-Rabbit HRP-conjugated secondary antibody

(ThermoFisher A16096, 1:1000). Membranes were again washed with TBS-T, then exposed to

Clarity Western ECL substrate (Bio-Rad) for 5 min before chemiluminescence and colorimetric

imaging.

Statistical analysis. Data on bacterial depth from Imaris were exported into Microsoft Excel and

GraphPad Prism 9 or 10 for visualization and analysis. The percent depth for each individual

bacterium was considered a unique data point and were aggregated across all fields of view for

each eye as one biological replicate. Data from 3-6 biological replicates for each condition were

combined into a single column for graphing and shown as median ± interquartile range. For all

ex vivo experiments, data did not pass normal distribution tests. The Kruskal-Wallis test with

Dunn's multiple comparisons or Kolmogorov-Smirnov test were used to compare groups. For

growth curve comparisons, One-way ANOVA with Dunnett's multiple comparisons or a paired

Student's t-Test were used. $P < 0.05$ was considered significant.

Funding Information. This work was supported by the National Institutes of Health; R01

EY011221 (SMJF) and R21 AI180421 (AR). The funding agency had no role in study design,

data collection and interpretation, or decision to publish.

Acknowledgements. Thanks to Dr. Alain Filloux (Imperial College London, UK) for originally

providing *P. aeruginosa* wild-type strain PAO1F.

Contributions. EJ, DS, NGK, AR, DE and SF designed the experiments; EJ, DS, NGK and AR

performed the experiments; EJ, AR, AS, DS, NGK, DE and SF analyzed and interpreted the

data; EJ, DE, AR and SF wrote the manuscript; DE and SF supervised the study.

REFERENCES

1. Morin CD, Déziel E, Gauthier J, Levesque RC, Lau GW. 2021. An organ system-based
synopsis of *Pseudomonas aeruginosa* virulence. *Virulence* 12:1469-1507

2. Juan C, Peña C, Oliver A. 2017. Host and pathogen biomarkers for severe *Pseudomonas*
aeruginosa infections. *J Infect Dis* 215:S44–S51.

3. Vincent JL, Sakr Y, Singer M, Martin-Loeches I, MacHado FR, Marshall JC, Finfer S,
Pelosi P, Brazzi L, Aditianingsih D, Timsit JF, Du B, Wittebole X, MácA J, Kannan S,
Gorordo-Delsol LA, De Waele JJ, Mehta Y, Bonten MJM, Khanna AK, Kollef M, Human
M, Angus DC. 2020. Prevalence and outcomes of infection among patients in intensive
care units in 2017. *JAMA* 323:1478-1487.

4. Magill SS, Edwards JR, Bamberg W, Beldavs ZG, Dumyati G, Kainer MA, Lynfield R,
Maloney M, McAllister-Hollod L, Nadle J, Ray SM, Thompson DL, Wilson LE, Fridkin
SK. 2014. Multistate point-prevalence survey of health care–associated infections. *N Engl*
J Med 370:1198-1208.

5. Reynolds D, Kollef M. 2021. The epidemiology and pathogenesis and treatment of
Pseudomonas aeruginosa infections: An update. *Drugs* 81:2117-2131

6. Cabrera-Aguas M, Khoo P, Watson SL. 2022. Infectious keratitis: A review. *Clin Exp*
Ophthalmol 50:543-562

7. Ung L, Bispo PJM, Shanbhag SS, Gilmore MS, Chodosh J. 2019. The persistent dilemma
of microbial keratitis: Global burden, diagnosis, and antimicrobial resistance. *Surv*
Ophthalmol 64:255-271.

8. Collier SA, Gronostaj MP, MacGurn AK, Cope JR, Awsumb KL, Yoder JS, Beach MJ.

2014. Estimated burden of keratitis--United States, 2010. *MMWR* 63:1027–1030.

9.   Stapleton F, Keay L, Edwards K, Naduvilath T, Dart JKG, Brian G, Holden BA. 2008.

The incidence of contact lens-related microbial keratitis in Australia. *Ophthalmology*

115:1655–1662.

10.   Durand ML, Barshak MB, Chodosh J. 2021. Infectious keratitis in 2021. *JAMA* 326:1319-

1320

11.   Morelli MK, Kloosterboer A, Fulton SA, Furin J, Newman N, Omar AF, Rojas LJ,

Marshall SH, Yasmin M, Bonomo RA. 2023. Investigating and treating a corneal ulcer

due to extensively drug-resistant *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother*

67:e0027723.

12.   Velcani F, Kuo IC, Shanks RMQ, Chodosh J, Garg P, Amescua G, Zegans ME. 2023.

Association of artificial tears with ocular and systemic infection: Carbapenem-resistant

Pseudomonas aeruginosa (VIM-GES-CRPA) Outbreak. *Ophthalmology* 130:1118-1120

13.   Tribin FE, Lieux C, Maestre-Mesa J, Durkee H, Krishna K, Chou B, Neag E, Tothova JD,

Martinez JD, Flynn Jr HW, Parel JM, Miller D, Amescua G. 2024. Clinical features and

treatment outcomes of carbapenem-resistant *Pseudomonas aeruginosa* keratitis. *JAMA*

Ophthalmol 142:407–415.

14.   Lee EJ, Cowell BA, Evans DJ, Fleiszig SMJ. 2003. Contribution of ExsA-regulated

factors to corneal infection by cytotoxic and invasive *Pseudomonas aeruginosa* in a

murine scarification model. *Invest Ophthalmol Vis Sci* 44:3892–3898.

15.   Tam C, Lewis SE, Li WY, Lee E, Evans DJ, Fleiszig SMJ. 2007. Mutation of the

phospholipase catalytic domain of the *Pseudomonas aeruginosa* cytotoxin ExoU abolishes

colonization promoting activity and reduces corneal disease severity. *Exp Eye Res*

85:799–805.

16.   Zolfaghar I, Evans DJ, Ronaghi R, Fleiszig SMJ. 2006. Type III secretion-dependent

modulation of innate immunity as one of multiple factors regulated by *Pseudomonas*

aeruginosa RetS. *Infect Immun* 74:3880–3889.

17.   Zolfaghar I, Angus AA, Kang PJ, To A, Evans DJ, Fleiszig SMJ. 2005. Mutation of *retS*,

encoding a putative hybrid two-component regulatory protein in *Pseudomonas*

aeruginosa, attenuates multiple virulence mechanisms. *Microbes Infect* 7:1305-1316

18.   Vance RE, Rietsch A, Mekalanos JJ. 2005. Role of the type III secreted exoenzymes S, T,

and Y in systemic spread of *Pseudomonas aeruginosa* PAO1 *in vivo*. *Infect Immun*

73:1706–1713.

19.   Shaver CM, Hauser AR. 2004. Relative contributions of *Pseudomonas aeruginosa* ExoU,

ExoS, and ExoT to virulence in the lung. *Infect Immun* 72:6969–6977.

20.   Goodman AL, Kulasekara B, Rietsch A, Boyd D, Smith RS, Lory S. 2004. A signaling

network reciprocally regulates genes associated with acute infection and chronic

persistence in *Pseudomonas aeruginosa*. *Dev Cell* 7:745–754.

21.   Mekonnen SA, Husseini N El, Turdiev A, Carter JA, Belew AT, El-Sayed NM, Lee VT.

2022. Catheter-associated urinary tract infection by *Pseudomonas aeruginosa* progresses

through acute and chronic phases of infection. *Proc Natl Acad Sci U S A* 119:

e2209383119.

22.   Rangel SM, Diaz MH, Knoten CA, Zhang A, Hauser AR. 2015. The role of ExoS in

dissemination of *Pseudomonas aeruginosa* during pneumonia. *PLoS Pathog* 11:e1004945.

23.   Hauser AR. 2009. The type III secretion system of *Pseudomonas aeruginosa*: infection by

injection. *Nat Rev Microbiol* 7:654-665.

24. Angus AA, Lee AA, Augustin DK, Lee EJ, Evans DJ, Fleiszig SMJ. 2008. *Pseudomonas*

aeruginosa induces membrane blebs in epithelial cells, which are utilized as a niche for

intracellular replication and motility. *Infect Immun* 76: 1992-2001.

25. Angus AA, Evans DJ, Barbieri JT, Fleiszig SMJ. 2010. The ADP-ribosylation domain of

Pseudomonas aeruginosa ExoS is required for membrane bleb niche formation and

bacterial survival within epithelial cells. *Infect Immun* 78:4500-4510.

26. Kroken AR, Chen CK, Evans DJ, Yahr TL, Fleiszig SMJJ. 2018. The impact of ExoS on

Pseudomonas aeruginosa internalization by epithelial cells is independent of *fleQ* and

correlates with bistability of type three secretion system gene expression. *MBio* 9:e00668-

18.

27. Kroken AR, Gajenthra Kumar N, Yahr TL, Smith BE, Nieto V, Horneman H, Evans DJ,

Fleiszig SMJ. 2022. Exotoxin S secreted by internalized *Pseudomonas aeruginosa* delays

lytic host cell death. *PLoS Pathog* 18:e1010306.

28. Metruccio MME, Tam C, Evans DJ, Xie AL, Stern ME, Fleiszig SMJ. 2017.

Contributions of MyD88-dependent receptors and CD11c-positive cells to corneal

epithelial barrier function against *Pseudomonas aeruginosa*. *Sci Rep* 7:13829.

29. Augustin DK, Heimer SR, Tam C, Li WY, Le Due JM, Evans DJ, Fleiszig SMJ. 2011.

Role of defensins in corneal epithelial barrier function against *Pseudomonas aeruginosa*

traversal. *Infect Immun* 79:595-605.

30. Fleiszig SMJ, Kroken AR, Nieto V, Grosser MR, Wan SJ, Metruccio MME, Evans DJ.

2019. Contact lens-related corneal infection: Intrinsic resistance and its compromise. *Prog*

Retin Eye Res 76: 100804.

31. Reins RY, Courson J, Lema C, Redfern RL. 2017. MyD88 contribution to ocular surface

homeostasis. *PLoS One* 12:0182153.

32.   Wan S, Datta A, Flandrin O, Metruccio MME, Ma S, Nieto V, Kroken AR, Hill RZ,

Bautista DM, Evans DJ, Fleiszig SMJ. 2021. Nerve-associated transient receptor potential

ion channels can contribute to intrinsic resistance to bacterial adhesion in vivo. *FASEB J*

35:e21899.

33.   Tam C, Mun JJ, Evans DJ, Fleiszig SMJ. 2012. Cytokeratins mediate epithelial innate

defense through their antimicrobial properties. *J Clin Invest* 122:3665-3677.

34.   Mantelli F, Mauris J, Argüeso P. 2013. The ocular surface epithelial barrier and other

mechanisms of mucosal protection: from allergy to infectious diseases. *Curr Opin Allergy*

Clin Immunol 13:563-8.

35.   Wan SJ, Sullivan AB, Shieh P, Metruccio MME, Evans DJ, Bertozzi CR, Fleiszig SMJ.

2018. IL-1R and MyD88 contribute to the absence of a bacterial microbiome on the

healthy murine cornea. *Front Microbiol* 9:1117.

36.   Wan SJ, Ma S, Evans DJ, Fleiszig SMJ. 2020. Resistance of the murine cornea to bacterial

colonization during experimental dry eye. *PLoS One* 15:e0234013.

37.   Alarcon I, Tam C, Mun JJJ, LeDue J, Evans DJ, Fleiszig SMJ. 2011. Factors impacting

corneal epithelial barrier function against *Pseudomonas aeruginosa* traversal. *Invest*

Ophthalmol Vis Sci 52:1368-1377.

38.   Lee EJ, Evans DJ, Fleiszig SMJ. 2003. Role of *Pseudomonas aeruginosa* ExsA in

penetration through corneal epithelium in a novel *in vivo* model. *Investig Ophthalmol Vis*

Sci 44:5220-5227.

39.   Kwong MSF, Evans DJ, Ni M, Cowell BA, Fleiszig SMJ. 2007. Human tear fluid protects

against *Pseudomonas aeruginosa* keratitis in a murine experimental model. *Infect Immun*

75:2325-2332.

40.   Fleiszig SMJ, Evans DJ, Do N, Vallas V, Shin S, Mostov KE. 1997. Epithelial cell

polarity affects susceptibility to *Pseudomonas aeruginosa* invasion and cytotoxicity. *Infect*

Immun 65: 2861-2867.

41.   Alarcon I, Kwan L, Yu C, Evans DJ, Fleiszig SMJ. 2009. Role of the corneal epithelial

basement membrane in ocular defense against *Pseudomonas aeruginosa. Infect Immun*

77:3264-3271.

42.   Marquart ME. 2011. Animal models of bacterial Keratitis. *J Biomed Biotechnol*

2011:680642.

43.   Hazlett LD. 2004. Corneal response to *Pseudomonas aeruginosa* infection. *Prog Retin*

Eye Res 23:1-30.

44.   Tam C, Mun JJ, Evans DJ, Fleiszig SMJ. 2010. The impact of inoculation parameters on

the pathogenesis of contact lens-related infectious keratitis. *Invest Ophthalmol Vis Sci*

51:3100-3106.

45.   Metruccio MME, Wan SJ, Horneman H, Kroken AR, Sullivan AB, Truong TN, Mun JJ,

Tam CKP, Frith R, Welsh L, George MD, Morris CA, Evans DJ, Fleiszig SMJ. 2019. A

novel murine model for contact lens wear reveals clandestine IL-1R dependent corneal

parainflammation and susceptibility to microbial keratitis upon inoculation with

Pseudomonas aeruginosa. Ocul Surf 17:119-133.

46.   Azghani AO, Gray LD, Johnson AR. 1993. A bacterial protease perturbs the paracellular

barrier function of transporting epithelial monolayers in culture. *Infect Immun* 61:2681-

2686.

47.   Azghani AO. 1996. *Pseudomonas aeruginosa* and epithelial permeability: role of

virulence factors elastase and exotoxin A. *Am J Respir Cell Mol Biol* 15:132-140.

48.   Heiniger RW, Winther-Larsen HC, Pickles RJ, Koomey M, Wolfgang MC. 2010.

Infection of human mucosal tissue by *Pseudomonas aeruginosa* requires sequential and

mutually dependent virulence factors and a novel pilus-associated adhesin. *Cell Microbiol*

12:1158-73.

49.   Golovkine G, Faudry E, Bouillot S, Elsen S, Attrée I, Huber P. 2016. *Pseudomonas*

aeruginosa transmigrates at epithelial cell-cell junctions, exploiting sites of cell division

and senescent cell extrusion. *PLoS Pathog* 12:e1005377.

50.   Nomura K, Obata K, Keira T, Miyata R, Hirakawa S, Takano K ichi, Kohno T, Sawada N,

Himi T, Kojima T. 2014. *Pseudomonas aeruginosa* elastase causes transient disruption of

tight junctions and downregulation of PAR-2 in human nasal epithelial cells. *Respir Res*

15:21.

51.   Alarcon I, Evans DJ, Fleiszig SMJ. 2009. The role of twitching motility in *Pseudomonas*

aeruginosa exit from and translocation of corneal epithelial cells. *Invest Ophthalmol Vis*

Sci 50:2237-2244.

52.   Ramirez JC, Fleiszig SMJ, Sullivan AB, Tam C, Borazjani R, Evans DJ. 2012. Traversal

of multilayered corneal epithelia by cytotoxic *Pseudomonas aeruginosa* requires the

phospholipase domain of *exoU*. *Invest Ophthalmol Vis Sci* 53:448-453.

53.   Leoni Swart A, Laventie B-J, Sütterlin R, Junne T, Lauer L, Manfredi P, Jakonia S, Yu X,

Karagkiozi E, Okujava R, Jenal U. 2024. *Pseudomonas aeruginosa* breaches respiratory

epithelia through goblet cell invasion in a microtissue model. *Nat Microbiol* 9:1725-1737.

54.   Li J, Metruccio MME, Evans DJ, Fleiszig SMJ. 2017. Mucosal fluid glycoprotein

DMBT1 suppresses twitching motility and virulence of the opportunistic pathogen

Pseudomonas aeruginosa. PLoS Pathog 13:e1006392

55. Fleiszig SM, Zaidi TS, Ramphal R, Pier GB. 1994. Modulation of *Pseudomonas*

aeruginosa adherence to the corneal surface by mucus. *Infect Immun* 62:1799-1804.

56. Fleiszig SMJ, Kwong MSF, Evans DJ. 2003. Modification of *Pseudomonas aeruginosa*

interactions with corneal epithelial cells by human tear fluid. *Infect Immun* 71:3866-3874.

57. Mun JJ, Tam C, Evans DJ, Fleiszig SMJ. 2011. Modulation of epithelial immunity by

mucosal fluid. *Sci Rep* 1:8.

58. Tabor LM, Grosser MR, Metruccio MMME, Kumar NG, Wu YT, Nieto V, Evans DJ,

Fleiszig SMJ. 2021. Human tear fluid modulates the *Pseudomonas aeruginosa*

transcriptome to alter antibiotic susceptibility. *Ocul Surf* 22: 94-102.

59. Gao N, Lee P, Yu FS. 2016. Intraepithelial dendritic cells and sensory nerves are

structurally associated and functional interdependent in the cornea. *Sci Rep* 6: 36414

60. Jiao H, Ivanusic JJ, McMenamin PG, Chinnery HR. 2021. Distribution of corneal TRPV1

and its association with immune cells during homeostasis and injury. *Investig*

Opthalmology Vis Sci 62:6.

61. Downie LE, Zhang X, Wu M, Karunaratne S, Loi JK, Senthil K, Arshad S, Bertram K,

Cunningham AL, Carnt N, Mueller SN, Chinnery HR. 2023. Redefining the human

corneal immune compartment using dynamic intravital imaging. *Proc Natl Acad Sci U S A*

120: e2217795120.

62. Sullivan AB, Tam KPC, Metruccio MME, Evans DJ, Fleiszig SMJ. 2015. The importance

of the *Pseudomonas aeruginosa* type III secretion system in epithelium traversal depends

upon conditions of host susceptibility. *Infect Immun* 83:1629-1640

63. McCaw ML, Lykken GL, Singh PK, Yahr TL. 2002. ExsD is a negative regulator of the

Pseudomonas aeruginosa type III secretion regulon. *Mol Microbiol* 46:1123-1133.

64.   Rietsch A, Vallet-Gely I, Dove SL, Mekalanos JJ. 2005. ExsE, a secreted regulator of type

III secretion genes in *Pseudomonas aeruginosa*. *Proc Natl Acad Sci U S A* 102:8006-

8011.

65.   Soong G, Parker D, Magargee M, Prince AS. 2008. The type III toxins of *Pseudomonas*

aeruginosa disrupt epithelial barrier function. *J Bacteriol* 190:2814-2821.

66.   Toska J, Sun Y, Carbonell DA, Foster ANS, Jacobs MR, Pearlman E, Rietsch A. 2014.

Diversity of virulence phenotypes among type III secretion negative *Pseudomonas*

aeruginosa clinical isolates. *PLoS One* 9:e86829.

67.   Turnbull L, Toyofuku M, Hynen AL, Kurosawa M, Pessi G, Petty NK, Osvath SR,

Cárcamo-Oyarce G, Gloag ES, Shimoni R, Omasits U, Ito S, Yap X, Monahan LG,

Cavaliere R, Ahrens CH, Charles IG, Nomura N, Eberl L, Whitchurch CB. 2016.

Explosive cell lysis as a mechanism for the biogenesis of bacterial membrane vesicles and

biofilms. *Nat Commun* 7:11220.

68.   Gonzalez D, Mavridou DAI. 2019. Making the best of aggression: The many dimensions

of bacterial toxin regulation. *Trends Microbiol* 27:897-905.

69.   Wydau-Dematteis S, El Meouche I, Courtin P, Hamiot A, Lai-Kuen R, Saubaméa B,

Fenaille F, Butel MJ, Pons JL, Dupuy B, Chapot-Chartier MP, Peltier J. 2018. Cwp19 is a

novel lytic transglycosylase involved in stationary-phase autolysis resulting in toxin

release in clostridium difficile. *MBio* 9:e00648.

70.   Redfern RL, Reins RY, McDermott AM. 2011. Toll-like receptor activation modulates

antimicrobial peptide expression by ocular surface cells. *Exp Eye Res* 92:209-220.

71.   Bai J, Kim SI, Ryu S, Yoon H. 2014. Identification and characterization of outer

membrane vesicle-associated proteins in *Salmonella enterica* serovar Typhimurium. *Infect*

Immun 82:4001-4010.

72. Kim SI, Kim S, Kim E, Hwang SY, Yoon H. 2018. Secretion of *Salmonella* pathogenicity

I island 1-encoded type III secretion system effectors by outer membrane vesicles in

Salmonella enterica Serovar Typhimurium. *Front Microbiol* 9:2810.

73. Sirisaengtaksin N, O'Donoghue EJ, Jabbari S, Roe AJ, Krachler AM. 2023. Bacterial

outer membrane vesicles provide an alternative pathway for trafficking of *Escherichia coli*

O157 type III secreted effectors to epithelial cells. *mSphere* 8:e0052023

74. Metruccio MME, Evans DJ, Gabriel MM, Kadurugamuwa JL, Fleiszig SMJ. 2016.

Pseudomonas aeruginosa outer membrane vesicles triggered by human mucosal fluid and

lysozyme can prime host tissue surfaces for bacterial adhesion. *Front Microbiol* 7:1-19.

75. Hritonenko V, Evans DJ, Fleiszig SMJ. 2012. Translocon-independent intracellular

replication by *Pseudomonas aeruginosa* requires the ADP-ribosylation domain of ExoS.

Microbes Infect 14:1366-1373.

76. Heimer SR, Evans DJ, Stern ME, Barbieri JT, Yahr T, Fleiszig SMJ. 2013. *Pseudomonas*

aeruginosa utilizes the type III secreted toxin ExoS to avoid acidified compartments

within epithelial cells. *PLoS One* 8:e73111.

77. Kumar NG, Nieto V, Kroken AR, Jedel E, Grosser MR, Hallsten ME, Mettrucio MME,

Yahr TL, Evans DJ, Fleiszig SMJ. 2022. *Pseudomonas aeruginosa* can diversify after host

cell invasion to establish multiple intracellular niches. *mBio* 13:e0274222

78. Kroken AR, Klein KA, Mitchell PS, Nieto V, Jedel EJ, Evans DJ, Fleiszig SMJ. 2023.

Intracellular replication of *Pseudomonas aeruginosa* in epithelial cells requires

suppression of the caspase-4 inflammasome. mSphere 8:e00351.

79.   Epelman S, Stack D, Bell C, Wong E, Neely GG, Krutzik S, Miyake K, Kubes P,

Zbytnuik LD, Ma LL, Xie X, Woods DE, Mody CH. 2004. Different domains of

Pseudomonas aeruginosa exoenzyme S activate distinct TLRs . *J Immunol* 173:2031-

2040.

80.   Epelman S, Neely GG, Ma LL, Gjomarkaj M, Pace E, Melis M, Woods DE, Mody CH.

2002. Distinct fates of monocytes and T cells directly activated by *Pseudomonas*

aeruginosa exoenzyme S . *J Leukoc Biol* 71:458-68

81.   Dortet L, Lombardi C, Cretin F, Dessen A, Filloux A. 2018. Pore-forming activity of the

Pseudomonas aeruginosa type III secretion system translocon alters the host epigenome.

Nat Microbiol 3:378-386.

82.   Miao EA, Mao DP, Yudkovsky N, Bonneau R, Lorang CG, Warren SE, Leaf IA, Aderem

A. 2010. Innate immune detection of the type III secretion apparatus through the NLRC4

inflammasome. *Proc Natl Acad Sci U S A* 107:3076-3080.

83.   Rayamajhi M, Zak DE, Chavarria-Smith J, Vance RE, Miao EA. 2013. Cutting edge:

Mouse NAIP1 detects the type III secretion system needle protein. *J Immunol* 191: 3986-

3989

84.   Mun JJ, Tam C, Kowbel D, Hawgood S, Barnett MJ, Evans DJ, Fleiszig SMJ. 2009.

Clearance of *Pseudomonas aeruginosa* from a healthy ocular surface involves surfactant

protein D and is compromised by bacterial elastase in a murine null-infection model.

Infect Immun 77:2392-2398.

85.   Rietsch A, Mekalanos JJ. 2006. Metabolic regulation of type III secretion gene expression

in *Pseudomonas aeruginosa*. *Mol Microbiol* 59:807-820.

86.   Sosnová-Netuková M, Kuchynka P, Forrester J V. 2007. The suprabasal layer of corneal

epithelial cells represents the major barrier site to the passive movement of small

molecules and trafficking leukocytes. *Br J Ophthalmol* 91:372-378.

87. MacDonald IA, Kuehna MJ. 2013. Stress-induced outer membrane vesicle production by

Pseudomonas aeruginosa. *J Bacteriol* 195:2971-2981.

88. Simonis A, Kreer C, Albus A, Rox K, Yuan B, Holzmann D, Wilms JA, Zuber S, Kottege

L, Winter S, Meyer M, Schmitt K, Gruell H, Theobald SJ, Hellmann AM, Meyer C,

Ercanoglu MS, Cramer N, Munder A, Hallek M, Fätkenheuer G, Koch M, Seifert H,

Rietschel E, Marlovits TC, van Koningsbruggen-Rietschel S, Klein F, Rybniker J. 2023.

Discovery of highly neutralizing human antibodies targeting *Pseudomonas aeruginosa*.

Cell 186:5098-5113

89. Goldberg JB, Crisan C V., Luu JM. 2022. *Pseudomonas aeruginosa* antivirulence

strategies: Targeting the type III secretion system. *Adv Exp Med Biol* 1386:257-280

90. Hmelo LR, Borlee BR, Almblad H, Love ME, Randall TE, Tseng BS, Lin C, Irie Y,

Storek KM, Yang JJ, Siehnel RJ, Howell PL, Singh PK, Tolker-Nielsen T, Parsek MR,

Schweizer HP, Harrison JJ. 2015. Precision-engineering the *Pseudomonas aeruginosa*

genome with two-step allelic exchange. *Nat Protoc* 10:1820-1841.

91. Choi KH, Schweizer HP. 2006. Mini-Tn7 insertion in bacteria with single attTn7 sites:

Example *Pseudomonas aeruginosa*. *Nat Protoc* 1:153-161.

92. Tam C, LeDue J, Mun JJ, Herzmark P, Robey EA, Evans DJ, Fleiszig SMJ. 2011. 3D

quantitative imaging of unprocessed live tissue reveals epithelial defense against bacterial

adhesion and subsequent traversal requires MyD88. *PLoS One* 6:e24008.

93. Guthoff RF, Zhivov A, Stachs O. 2009. In vivo confocal microscopy, an inner vision of

the cornea - A major review. *Clin Exp Ophthalmol* 37:100-117.

94.   Bleves S, Soscia C, Nogueira-Orlandi P, Lazdunski A, Filloux A. 2005. Quorum sensing

909   negatively controls type III secretion regulon expression in *Pseudomonas aeruginosa* PA

910   95.   Cisz M, Lee PC, Rietsch A. 2008. ExoS controls the cell contact-mediated switch to

911   effector secretion in *Pseudomonas aeruginosa*. *J Bacteriol* 190:2726-38

912   96.   Bloemberg G V, O'Toole G a, Lugtenberg BJ, Kolter R. 1997. Green fluorescent protein

913   as a marker for *Pseudomonas* spp. *Appl Environ Microbiol* 63:4543-4551.

914   97.   Brutinel ED, Vakulskas CA, Brady KM, Yahr TL. 2008. Characterization of ExsA and of

915   ExsA-dependent promoters required for expression of the *Pseudomonas aeruginosa* type

916   III secretion system. *Mol Microbiol* 68:657-671.

918   Table 1. Properties and epithelial traversal results for *P. aeruginosa PAO1F and T3SS mutants*

Strain	Exotoxins	Translocon	Needle	Epithelial Traversal
PAO1F	+	+	+	++
Δ*exsD*	++	++	++	+
Δ*pscC*	+	+	-	++
Δ*popBD*	+	-	+	+
Δ*exoSTY*	-	+	+	+
Δ*pscC*Δ*popBD*	+	-	-	++
Δ*popB*Δ*exoSTY*	-	-	+	-
Δ*pscU-L:P$_{rha}$exsA*	+	-	-	+
Δ*pscU-L*Δ*exoSTY:P$_{rha}$exsA*	-	-	-	-

934   Table 2. Strains, mutants, and plasmids used in this study

Strain or mutant	Description	Source
PAO1F	Wild-type *P. aeruginosa*; All mutants made within this background	Dr. Alain Filloux *via* Dr. Arne Rietsch (94)
Δ*exsA*	*exsA* mutant; Constitutively T3SS^{Off}	Dr. Arne Rietsch
Δ*exsD*	*exsD* mutant; Constitutively T3SS^{On}	Dr. Timothy Yahr (63)
Δ*pscC*	*pscC* mutant; Lacking needle	Dr. Arne Rietsch (95)
Δ*popBD*	*popBD* double mutant; Lacking translocon pore	Dr. Arne Rietsch (95)
Δ*exoSTY*	*exoSTY* triple mutant; Lacking known exotoxins	Dr. Arne Rietsch (95)
Δ*pscU-L*	Total T3SS-operon mutant (knockout of all genes *pscU* to *pscL*); called *ΔU-L*	Dr. Arne Rietsch (66)
Δ*exsA:Prha-A*	*exsA* mutant with chromosome-integrated rhamnose-inducible *exsA*	Dr. Abby Kroken (27)
Δ*pscU-L:Prha-A*	*ΔU-L* with rhamnose-inducible *exsA*	This study
Δ*pscU-L:Prha-V*	*ΔU-L* with rhamnose-inducible vector control	This study
Δ*pscU-L* Δ*exoSTY:Prha-A*	*ΔU-L* lacking all known exotoxins, with rhamnose-inducible *exsA*	This study
Δ*popB*Δ*exoSTY*	*popB* mutant lacking all known exotoxins	Dr. Victoria Hritonenko (75)
Δ*pscC*Δ*popBD*	*popBD* mutant lacking T3SS needle	This study
Δ*popB*Δ*exoS*	*popB* mutant lacking *exoS*	This study
E. coli SM10(λ*pir*)	Mutagenesis vector donor strain	New England Biolabs
Plasmid	**Description (Resistance conferred, µg/ml)**	**Source**
pSMC2	Constitutive GFP (Carbenicillin, 400)	(96)
pJNE05	T3SS-GFP (P*exoS*). Plasmid encoding *gfp* fused to the ExsA-dependent promoter of *exoS*. (Gentamicin, 200)	(97)
pEXG2-Δ*pscU-L*	Integrating suicide plasmid for mutagenesis	Dr. Arne Rietsch
pEXG2-Δ*exoS*	Integrating suicide plasmid for mutagenesis	Dr. Arne Rietsch
pUC18-mini-Tn7T-PrhaB-*exsA*	Delivery vector for integrating *exsA* under Rhamnose-sensitive promoter to *att*Tn7 site (Carbenicillin, 100)	This study
pUC18-mini-Tn7T-PrhaB-Vector	Delivery vector for integrating empty expression vector under rhamnose-sensitive promoter to *att*Tn7 site (Carbenicillin, 100)	This study
pTNS1	Helper plasmid for conjugation of Tn7 integration plasmid (Ampicillin, 100)	(91)
pRK2013	Helper plasmid for conjugation of Tn7 integration plasmid (Kanamycin, 35)	(91)
pFLP2	Sucrose-sensitive vector to excise antibiotic resistance marker from *att*Tn7 site	(91)

  FIGURE LEGENDS

Figure 1. Method for quantitative imaging and analysis of bacterial traversal of the corneal

epithelium. Whole murine eyeballs *ex vivo* were blotted with tissue paper, incubated in 0.1M

EGTA solution (1 h), then infected with ~1 x 10^{11} CFU/ml *P. aeruginosa* expressing GFP. Eyes

were fixed, whole mounted *en face*, and imaged by confocal microscopy. (A) Raw image

imported into Imaris 9.9 for 3D analysis, confocal reflectance signal shown in red and bacterial

fluorescence in green. (B) Boundary between stroma and epithelium manually delineated in

sequential XZ slices to exclude signal from stroma and create epithelium object. (C) Objects

representing apical and basal boundaries of epithelium. (D) Bacteria automatically identified by

fluorescent signal quality. (E) Bacteria colored by their depth of traversal, measured as a

percentage of its position from the apical to basal boundaries.

Figure 2. Epithelial traversal by wild-type or T3SS mutants. (A) Traversal depth after *ex*

vivo infection by PAO1F and T3SS mutants in genes encoding the transcriptional activator *exsA*,

transcriptional repressor *exsD*, known exotoxins *exoSTY*, needle component *pscC*, or translocon

pore *popBD*. Each data point represents the relative position (% depth) of one bacterium between

the apical and basal boundaries of the 3D corneal epithelium. All data points pooled from ≥ 4

eyes per group. Error bars show the median +/- interquartile range. Note: All comparisons

between the T3SS mutants with the Δ*exsA* mutant were significant as were all comparisons with

wild-type PAO1F (P < 0.0001, Kruskal-Wallis test with Dunn's multiple comparisons). (B)

Cumulative histogram graphing traversal data (from Fig. 2A) of wild-type and T3SS mutants as

a percent of the total bacterial population found deeper in the epithelium. The vertical line

indicates halfway through the epithelium, highlighting the proportion of each strain that

traversed more than 50% of the epithelial layer.

Figure 3. ExsA complementation in a T3SS operon-deficient background. (A) Traversal

depth after *ex vivo* infection by a Δ*pscU-L* mutant which lacks all T3SS genes except *exoSTY*

(and their respective chaperones) and a Δ*pscU-L*Δ*exoSTY* mutant which also lacks effector

toxins. Controls include expression of an empty vector or *exsA* induced by overnight growth on

and inclusion of 2% rhamnose during infection. Data were pooled from ≥ 3 eyes per strain. Error

bars show median +/- interquartile range. All comparisons between groups were significant at P

< 0.0001 except for Δ*pscU-L* vs. Δ*pscU-L*Δ*exoSTY*:P_{rha}-*exsA* + rhamnose, which had P = 0.025

(Kruskal-Wallis test with Dunn's multiple comparisons). (B) Cumulative histogram graphing

traversal data (from Fig. 3A) of T3SS operon-deficient mutants with ExsA complementation or

controls as a percent of the total bacterial population found deeper in the epithelium. The vertical

line indicates halfway through the epithelium, highlighting the proportion of each strain that

traversed more than 50% of the epithelial layer.

Figure 4. Traversal by translocon and needle- or exotoxin-deficient mutants. (A) Traversal

depth after *ex vivo* infection by T3SS translocon mutants in a background of Δ*pscC*, Δ*exoS*, or

Δ*exoSTY*. Data from Fig. 2A was used for Δ*pscC* and Δ*exoSTY* mutants. Data pooled from ≥3

eyes per strain. Error bars show median +/- interquartile range. All comparisons between groups

were significant (P < 0.0001, Kruskal-Wallis test with Dunn's multiple comparisons) except for

Δ*exoSTY* vs. Δ*popB*. (B) Cumulative histogram graphing traversal data (from Fig. 4A) of the

T3SS mutants as a percent of the total bacterial population found deeper in the epithelium. The

vertical line indicates halfway through the epithelium, highlighting the proportion of each strain

that traversed more than 50% of the epithelial layer.

Figure 5. Rhamnose-induction of ExsA-driven ExoS expression in both bacterial cells and

supernatant of a needle-deficient mutant Δ*pscU-L*. Western immunoblot showing bacterial

cell pellets and supernatant fractions collected from Δ*exsA*:P$_{rha}$*exsA*, Δ*pscU-L*:P$_{rha}$Vector,

Δ*pscU-L*:P$_{rha}$*exsA* after growth to 'mid-log' phase in LB media, with induction of *exsA* by 2%

rhamnose inclusion for the conditions indicated. The location of bands corresponding with ExoS

(~49 kDa) detected by affinity-purified antibody and chemiluminescence are evident below the

protein standard annotated at 50 kDa.

Supplemental Material

Supplemental Figure S1. Comparison of corneal epithelium traversal by *P. aeruginosa* and

its T3SS mutants. Representative images of corneal epithelium traversal by wild-type PAO1F

compared to the Δ*pscC*, Δ*popBD* and Δ*exoSTY* mutants color-coded for traversal depth.

Supplemental Figure S2. EGTA induction of *P. aeruginosa* T3SS gene expression was

similar between PAO1 and T3SS mutants. OD$_{600}$-normalized GFP signal from pJNE05

(P*exoS*) carried by PAO1F or its T3SS mutants in the translocon pore Δ*popBD*, known exotoxins

Δ*exoSTY*, or the transcriptional repressor Δ*exsD* after 24 h growth in T3SS-induction media

[TSB plus gentamicin (200 μg/ml), monosodium glutamate (100 mM), glycerol (1 %) and EGTA

(2 mM)]. Similar P*exoS* induction, i.e. T3SS induction, was observed across wild-type and

mutants. N = 3 separate growth curves per group. Mean +/- standard deviation, ns = not

significant (One-way ANOVA with Dunnett's Multiple Comparisons).

1006

Supplemental Figure S3. Rhamnose-induction of ExsA rescues *ex vivo* traversal. (A) OD_{600}-

normalized GFP signal from pJNE05 (P*exoS*) carried by a Δ*exsA* mutant complemented with

rhamnose-inducible *exsA* with or without inclusion of rhamnose (Rha) (2 %) in the growth

medium [TSB with gentamicin (200 μg/ml), monosodium glutamate (100 mM) and glycerol

(1%)] after 24 h of growth. N = 3 separate growth curves per group. Mean +/- standard

deviation, ** $P < 0.01$ (Paired Student's t-Test). (B) Traversal depth after *ex vivo* infection of

Δ*exsA* with expression of *exsA* remaining off or induced by overnight growth with rhamnose (2

%). Data pooled from 3 eyes per strain. Error bars show the median with interquartile range.

**** $P < 0.0001$ (Kolmogorov-Smirnov test).

A)

Wild-Type

Median % Depth: 47.29　7.31　12.66　45.24　11.16　10.91

Apical

% of Epithelium traversed

Basal

Strain Expresses:	PAO1F	ΔexsA	ΔexsD	ΔpscC	ΔpopBD	ΔexoSTY
ExsA	+	–	+	+	+	+
Needle	+	–	+	–	+	+
Translocon	+	–	+	+	–	+
Effectors	+	–	+	+	+	–

B)

Pop. Over Halfway

- PAO1F　47.27%
- ΔexsA　2.67%
- ΔexsD　21.42%
- ΔpscC　43.84%
- ΔpopBD　30.18%
- ΔexoSTY　19.27%

% of Total Population Remaining Deeper

0　25　50　75　100

Apical　　　　　　　　　　Basal

% Depth of Epithelium

A)

Median % Depth:

| 5.19 | 7.34 | 11.09 | 5.76 |

Strain Expresses:	ΔpscU-L	ΔpscU-L:P$_{rha}$Vector +Rha	ΔpscU-L:P$_{rha}$exsA +Rha	ΔpscU-LΔexoSTY:P$_{rha}$exsA +Rha
ExsA	–	–	+	+
Needle	–	–	–	–
Translocon	–	–	–	–
Effectors	–	–	+	–

B)

	Pop. Over Halfway
ΔpscU-L	0.15%
ΔpscU-L:P$_{rha}$Vector +Rha	0.00%
ΔpscU-L:P$_{rha}$exsA +Rha	18.35%
ΔpscU-LΔexoSTY:P$_{rha}$exsA +Rha	0.00%

A)

Median % Depth:

| | 10.91 | 12.30 | 9.42 | 8.10 | | 45.24 | 36.11 |

Strain Expresses:	ΔexoSTY	ΔpopB	ΔpopBΔexoS	ΔpopBΔexoSTY	ΔpscC	ΔpscCΔpopBD
ExsA	+	+	+	+	+	+
Needle	+	+	+	+	–	–
Translocon	+	–	–	–	+	–
Effectors	–	+	only *exoTY*	–	+	+

B)

Pop. Over Halfway

ΔpscC		43.84%
ΔpscCΔpopBD		35.00%
ΔexoSTY		19.27%
ΔpopB		17.76%
ΔpopBΔexoS		5.86%
ΔpopBΔexoSTY		1.62%

% of Total Population Remaining Deeper

% Depth of Epithelium

Apical — Basal