Divergent ecological selection causes partial reproductive isolation and vast genomic

instability in yeast

Running head: Ecological isolation barriers in yeast

Devin P. Bendixsen[1,2], Ciaran Gilchrist[1], Chloé Haberkorn[1], Karl Persson[3], Cecilia Geijer[3], Jonas Warringer[4],

Rike Stelkens[1]*

[1] Department of Zoology, Stockholm University, Svante Arrheniusväg 18 B, 106 91 Stockholm, Sweden

[2] MRC Human Genetics Unit, Institute of Genetics and Cancer, University of Edinburgh, Edinburgh, UK

[3] Division of Industrial Biotechnology, Department of Life Sciences, Chalmers University of Technology,

Kemigården 1, 412 96 Göteborg, Sweden

[4] Department of Chemistry and Molecular Biology, University of Gothenburg, Medicinaregatan 9C,

41390 Göteborg, Sweden

* Corresponding author: Rike Stelkens, Department of Zoology Stockholm University, telephone: +46 (0)

816 4223; email: rike.stelkens@zoologi.su.se

Devin Bendixsen: 0000-0003-0831-7646; devin.bendixsen@gmail.com

Ciaran Gilchrist: 0000-0002-7639-6131; ciaran.gilchrist@umu.se

Chloé Haberkorn: 0000-0002-7371-9177; chloe.haberkorn@zoologi.su.se

Karl Persson: 0000-0002-2173-8165; karperss@chalmers.se

Cecilia Geijer: 0000-0002-4158-2938; cecilia.geijer@chalmers.se

Jonas Warringer: 0000-0001-6144-2740; jonas.warringer@cmb.gu.se

Rike Stelkens: 0000-0002-8530-0656; rike.stelkens@zoologi.su.se

Abstract

Populations evolving independently in divergent environments accumulate genetic differences and potentially evolve reproductive isolation as a by-product of divergence. The speed and mechanisms underlying this process are difficult to investigate because we rarely get the opportunity to witness them in natural settings, and histories of selection and gene flow between populations are often unknown. Here, we experimentally evolved yeast for 1000 generations of evolution in both divergent and parallel environments. At regular time points during experimental evolution, we made crosses between parallel- and divergent-evolving populations to measure postzygotic reproductive isolation (gamete viability). We used whole genome population sequencing to determine the mutational load, the number and types of structural variation, and other genomic features of the parent, F1, and F2 intraspecific hybrids. We found evidence for large scale phenotypic and genome-wide differentiation in response to divergent laboratory selection. Divergent-selected populations produced hybrids with reduced gamete viability - a classic signature of postzygotic reproductive isolation in the form of hybrid breakdown. Parallel-selected populations on the other hand remained reproductively compatible. We found that F2 hybrid genomes contained vast genomic instability, i.e., new structural variants (especially insertions, deletions, and interchromosomal translocations) that were not observed in parent and F1 genomes, which is likely a result of chromosome missegregation and recombination errors in hybrid meiosis. Our results provide phenotypic and genomic evidence that partial reproductive isolation evolved due to adaptation to divergent environments, consistent with predictions of ecological speciation theory.

Keywords: experimental evolution, hybridization, mutational load, reproductive isolation, *Saccharomyces,* structural variation

Introduction

Ecological speciation implies a process by which adaptation to different ecological niches, in allopatry or sympatry, leads to the evolution of reproductive isolation between populations (Mayr 1947; Schluter 2001; McKinnon et al. 2004; Nosil 2012). Long-term monitoring of natural populations has been valuable in linking ecological traits with the emergence of reproductive barriers (e.g. (Filchak et al. 2000; Rundle et al. 2000; Grant and Grant 2003; Funk et al. 2006; Sawada et al. 2022)), providing convincing arguments for ecological factors to play a role in driving population divergence. However, ecological speciation remains difficult to demonstrate because it usually proceeds over extended evolutionary times (Hendry et al. 2007), trait divergence is often cryptic (Struck et al. 2018; Sparks et al. 2022), and the evolutionary histories of natural populations are challenging to reconstruct. It is usually unknown whether divergent selection was present, how long it lasted, and what the timing and amount of gene flow was between populations. Thus, it often remains unclear, especially between allopatric species pairs occupying different niches, whether selection was the driving force underlying their divergence, eventually leading to speciation (Anderson and Weir 2022; Irestedt et al.; Müller et al.). Due to these constraints, studies have often relied on combinatorial approaches, using i) data on niche divergence in traits related to habitat use, such as resource utilization or foraging behavior (e.g. (Hatfield and Schluter 1999; Nosil and Crespi 2006; Munar-Delgado et al. 2024), ii) data on reproductive barriers, such as differences in mating behaviors or flowering times reducing gene flow between populations adapted to different ecological niches (e.g. (Dieckmann et al. 2004; Villa et al. 2019)), and iii) genetic data, detecting signatures of selection and divergence in genetic regions associated with ecological adaptation and reproductive isolation (e.g. (Nosil and Schluter 2011; Härer et al. 2021; Louder et al. 2024)).

Microbial experimental evolution provides a powerful tool for investigating questions in evolutionary biology. It has been used to describe the dynamics and genetic mechanisms underlying adaptation to stable (Lenski et al. 1991; Good et al. 2017; Johnson and Desai 2022), heterogenous

(Rainey and Travisano 1998), or changing environments (Bell and Gonzalez 2011; Gorter et al. 2016;

Gorter et al. 2017) and to test theoretical predictions about adaptation rates in sexual versus asexual

populations (McDonald et al. 2016; Kosheleva and Desai 2018; Leu et al. 2020). Experimental evolution

with microbes can also provide insights on fundamental questions in speciation, especially about the

early stages of divergence, i.e. why, how and when lineages split (Stelkens 2024). Microbial genomes are

often small and generation times are short, allowing for evolve-and-resequencing approaches to capture

speciation and its underlying genetic basis in action. For instance, experimental evolution with a virus

(lambda bacteriophage) and its bacterial host (*Escherichia coli*) has demonstrated that divergent

ecological selection caused the phages to split into two genetically distinct lineages with different host

preferences, and that barriers to gene flow formed as a consequence of ecological specialization, which

resulted in reproductive isolation (Meyer et al. 2016). Experiments with yeast and other fungi have

shown that divergent selection can lead to phenotypic divergence and rapid ecological differentiation

(Dettman et al. 2007; Dettman et al. 2008; Ament-Velásquez et al. 2022), even under high rates of gene

flow (Tusso et al. 2021). But generally, there is a lack of studies that combine both organismal and

genetic approaches to measure reproductive isolation (Westram et al. 2022). Especially work

investigating the emergence and dynamics of reproductive isolation using repeated measures of hybrid

viability across generational time scales, from the beginning to later stages of parental divergence under

controlled environmental conditions, could significantly advance our understanding of what drives the

evolution of reproductive isolation in the early stages of speciation.

Here, we used experimental evolution with the budding yeast *Saccharomyces cerevisiae* and

propagated replicate, asexually reproducing populations in both different and parallel (same)

environments for 1000 mitotic generations (**Figure 1**). At regular time points of population divergence,

we generated sexual F1 and F2 (hybrid) crosses between and within environments, to test whether

reproductive isolation evolved and increased with time. We profiled parental phenotypes in 90 different

environments to test for phenotypic divergence over time, and used whole genome sequencing on

parental, F1, and F2 hybrid populations to measure their mutational load and other measures of

genomic stability. By the endpoint of experimental evolution, we found that divergent-selected

populations were up to 50% reproductively isolated due reduced hybrid gamete viability, while crosses

made between parallel-selected populations remained genetically compatible. We detected vast

structural variation in the F2 hybrid genomes, including up to three times more deletions, insertions,

inversions, tandem duplications, and interchromosomal translocations than in the diploid parental and

F1 hybrid genomes. Together, our results suggest that structural genetic differences accumulating in

populations under divergent ecological selection led to severe genomic instability in hybrid genomes

due to recombination errors and chromosome missegregation which caused reproductive isolation,

consistent with predictions of ecological speciation theory.

Materials and Methods

Founder populations

We generated large diploid, and genetically diverse, founder populations of yeast through mating and

mass sporulation of two *S. cerevisiae* strains (SK1 and Y55; Figure 1) as previously described (Ament-

Velásquez et al. 2022). SK1 and Y55 are relatively closely related and high quality reference genomes

are available for both (Peter et al. 2018; Devin P Bendixsen et al. 2021). Briefly, SK1 and Y55 were

genetically modified to introduce auxotrophies, which could be used as selective markers to exclude

non-mated haploids (non-hybrids). The founders of populations (called "N founder" below) evolving in

sodium chloride (NaCl) are the diploid recombined offspring of a cross between SK1 (MATα ho:KanMX

ura3::ClonNat) and Y55 (MATa ho leu2::HygMX ura3::ClonNat). All other founders (called "LE founder"

below) of the populations evolving in ethanol and two different concentrations of lithium acetate, are

the diploid recombined offspring of a cross between SK1 (MATα ho:KanMX lys2::ClonNat) and Y55

(MATa ho leu2::HygMX lys2::ClonNat).

Experimental evolution of parental populations

From these founders, replicate populations (from here on 'parental´ populations) were experimentally

evolved in four different (divergent) and same (parallel) environments, in four independent replicates

each, for up to 1000 asexual generations (**Figure 1**). The number of generations was estimated by

counting cells after 48h growth in selective media in the highest stress environment (0.75M NaCl, see

Ament-Velásquez et al. 2022). This approach is conservative because it likely underestimates the

number of generations in the other, less stressful environments. All evolution environments contained

synthetic complete (SC) medium (2% glucose, 0.67% bacto-yeast nitrogen base without amino acids,

0.00079% Formedium CSM powder (product code DCS0019), made up to 1L with sterilised deionised

water) plus one of the following substances to induce stress: 0.75M sodium chloride (from here on

NaCl), 8% ethanol (C_2H_6O, from here on EtOH), and two different concentrations of lithium acetate

($C_2H_3LiO_2$, from here on LiAc0.01M and LiAc0.02M). The correct volume of EtOH was added to the test

tube just before inoculation to limit evaporation. We have previously demonstrated that these four

media significantly differ in how much they reduce the fitness of populations, and we have provided

phenotypic and genetic evidence that populations become differentially adapted to these environments

(Ament-Velásquez et al. 2022).

Three to four replicate populations were evolved in separate test tubes containing 5mL of SC

media with one of the four substances. Then, we serially transferred 10μL of yeast culture every 48

hours to fresh media (approximately seven generations, estimated by counting cells after 48 hours).

Every five transfers (10 days, i.e. ~35 generations), a 900μL sample of each population was frozen in 44%

glycerol at -70°C for later use. The experiment was run for a total of ~300 days. After some cross-

replicate contamination was detected using sequencing data (Ament-Velásquez et al. 2022), we

excluded two NaCl-evolved replicate populations (R1 and R4) at 1000 generations, all LiAc0.01M and

LiAc0.02M-evolved populations at 1000 generations, and one LiAc0.01M-evolved population (R3) at all

generations. No crosses were generated from these populations. Retained populations showed no signs

of cross-contamination in terms of shared single nucleotide variants.

148
Figure 1: Experimental Design. A founder population was generated by the mating of two *S. cerevisiae*
strains (SK1 and Y55), followed by mass sporulation (meiosis). This genetically diverse, diploid founder
was split into four different environments, containing growth medium with either 0.75M NaCl, 8% EtOH,
0.01M lithium acetate, or 0.02M lithium acetate, and evolved for up to 1000 asexual generations by
serial transfer, in three or four independent biological replicates per environment (Ament-Velásquez et
al. 2022). At seven time points of divergence between these 'parent' populations, sexual hybrid crosses
were made between NaCl-populations and populations from all other environments (with replication).
Control crosses were made between populations evolving in parallel, in the same environment. All
crosses were then sporulated and the viability of the haploid F2 offspring was assessed after tetrad
dissection. 2N indicates diploid, 1N indicates haploid genotypes.
159

Hybridization at increasing timepoints of parental divergence

We generated hybrid crosses at 100, 300, 500, 700, and 1000 generations of parental divergence,

between all four replicate NaCl-populations and up to four replicate populations from each of the three

other environments (LiAc 0.01M (3 replicates), LiAc 0.02M (4 replicates) and EtOH (4 replicates); **Figure**

1). Due to the selectable markers present in the parents, NaCl- populations could be mass-crossed with

the other three populations while safely removing all non-hybrid parental genotypes using a selective

medium (described below). First, we mass-sporulated the diploid parental populations. For this, 500μL

of saturated culture were added to 50mL of sporulation medium (1% potassium acetate, 0.05% glucose,

0.1% yeast extract, and H_2O to 1L) and left for 5 days with gentle shaking at room temperature. We then

mixed 500μL of each of the two parent spore cultures together in a 1.5mL Eppendorf tube, and

centrifuged the culture down to a pellet to induce mating. To select for F1 hybrids, we inoculated 25μL

of the pellet in 5mL of -lys SD medium with G418 (0.67% bacto-yeast nitrogen base without amino acids,

2% glucose, dropout powder (-lys), G418 400mgL^{-1} and H_2O to 1L). This selects for F1 hybrid genotypes

in bulk, which are heterozygous for both selective markers (-lys and G418). Note that this mass mating

and sporulation protocol permitted F1 hybrid populations to contain genetic diversity, depending on

how many different parental genotypes participated in the mass mating. To generate F2 hybrid

populations, we put F1 hybrid populations through one round of meiosis by mass sporulating them in

50mL of sporulation medium for 5 days with gentle shaking at room temperature. F1 AND F2 hybrid

populations were stored at -70°C in 22% glycerol solution for whole genome sequencing.

Assessing parental phenotypic divergence

We phenotypically profiled all parent populations (i.e. populations evolved in NaCl, LiAc 0.01M, LiAc

0.02M, and EtOH) in 90 different environments, at increasing times of divergence, using the Scan-o-

Matic system for high-throughput microbial phenotyping (Zackrisson et al. 2016; Li et al. 2019; Alalam et

al. 2020; Mukherjee et al. 2021; Persson et al. 2022; Stenberg et al. 2022) version 2.2

(https://github.com/Scan-o-Matic/scanomatic.git). Parental populations were precultured in SC media

for 48h in 96-well plates at 30°C without shaking and frozen. Then, cultures were pinned directly from

thawed cryo-stocks in 96-format for preculturing on agar plates (for more detailed methods see

(Persson et al. 2024)). Samples were split into duplicates (two 96-well plates). A third plate was used for

propagation of a spatial control, containing a Y55 diploid strain. After preculture, populations were

pinned onto 1536 colony arrays on agar plates robotically (Singer ROTOR HDA), allowing for accurate

measuring of colony growth using the Scan-o-matic software. The spatial control was placed in every

fourth position (in total 384 colonies per plate) to remove any spatial bias, irregularities in the plates, or

other bias from the local area. In total, 52 distinct stressors were used in different concentrations,

resulting in 90 distinct environments, including superoxidative stress (e.g. paraquat), heavy metals (e.g.

$CuSO_4$), different carbon sources (e.g. maltose), antifungal drugs (e.g. Clotrimazole). Phenotypic data, i.e.

growth measurements, are reported in Table S1, with the full list of environments in Table S2.

Cultivation plates were maintained undisturbed and without lids for the entire duration of the assay

(72h) in high-resolution desktop scanners (Epson Perfection V800 PHOTO scanners, Epson Corporation,

UK) at 30°C, in moisture-controlled thermostatic cabinets with air circulation. We imaged plates at 20-

minute intervals using transmissive scanning at 600 dpi and extracted intensities for pixels included in,

and outside, each colony. We estimated the sum pixel intensity for each colony, subtracted the median

pixel intensity of the local background, and transformed the remaining cell-associated pixel intensity to

cell counts by using a pre-established calibration function, established using both spectrometry and flow

cytometry (Zackrisson et al. 2016). We smoothed and quality-controlled growth curves, rejecting

approximately 0.3% of growth curves as erroneous, blind with respect to sample identities. Based on the

number of cells after 72h, we used the yield normalized against the spatial control (Y55 diploid).

A custom 'ParentalDivergence' function was generated in R, using normalized yield, the number

of generations in experimental evolution, and the type of population comparison. Mean and double

standard deviations of normalized yields were first computed for each of the 90 environments for each

of the replicate parent 1 (P1) and 2 (P2) populations. This was done for each type of population

comparison, i.e. divergent-environment comparisons (NaCl x EtOH, NaCl x LiAc0.01, NaCl x LiAc0.02) and

parallel-environment comparisons (NaCl x NaCl, EtOH x EtOH, LiAc0.01 x LiAc0.01, LiAc0.02 x LiAc0.02).

We considered fitness to be higher if the normalized mean yield of P2 was larger than the normalized

mean yield of P1 plus P1's double standard deviation (i.e. mean P2 > mean P1 + 2σ). We considered

fitness to be lower if the normalized mean yield of P2 was lower than the normalized mean yield of P1

minus P1's double standard deviation, following previous protocols (i.e. mean P2 < mean P1 - 2σ; see

(deVicente and Tanksley 1993; Brice et al. 2021). Parental phenotypic divergence is thus defined as the

proportion of the 90 environments in which there was a large difference in yield between the two

populations compared.

Assessing reproductive isolation

To assess the presence and degree of reproductive isolation between the parental yeast populations

evolving in divergent environments, we mated individual parent genotypes after 100, 300, 500, 700 and

1000 generations of experimental evolution. The crosses were plated onto solid YPD agar medium (1%

yeast extract, 2% peptone, 2% glucose, 2% agar made up to 1L with sterilized deionized H_2O) and grown

for 2 days, followed by replica plating onto solid KAc medium (1% potassium acetate, 0.05% glucose,

0.1% yeast extract and H_2O to 1L) for sporulation. Populations were left for 2-7 days, until tetrads were

identified under the microscope. Yeast tetrads usually contain four spores (gametes), which are the

recombined haploid products of meiosis. Sporulated cultures with tetrads were incubated for one hour

at room temperature in 10μL of filter sterilized zymolyase solution (1mg of zymolyase per 1mL of 1M

sorbitol solution) to kill vegetative (unsporulated) cells and digest the outer membrane of the tetrads,

surrounding the ascospores. Cell lysis was terminated by adding 500µL of sterilized H_2O. To assess the

fertility of F1 hybrid crosses at increasing timepoints of parental divergence, we dissected 10-20

individual F1 tetrads, and measured the survival rates of their haploid F2 offspring. Tetrads were

dissected on YPD plates using a Singer MSM 400 robot. Individual spores were then left to germinate

and grow for 72h at 30°C, at which point spore viability was assessed. Spores were counted as viable

based on the presence of colonies visible to the eye. Thus gamete viability was assessed as the capacity

to germinate and show sufficient growth to form macroscopic colonies. The total proportion of unviable

spores per cross was calculated from 20 dissected tetrads (80 spores) at 100, 700, and 1000 generations,

and 10 tetrads (40 spores) at 300 and 500 generations of parental divergence.

We also measured the F1 fertility of the control crosses, i.e. crosses between replicate parental

populations that evolved in parallel environments. We used the same mating, sporulation and dissection

protocols as described above. The total proportion of unviable spores (gametes) per cross was

calculated from 10 dissected tetrads (40 spores) at 100, 500, 700 and 1000 generations of parental

divergence.

DNA extraction and whole-genome sequencing

DNA was extracted as described in (Ament-Velásquez et al. 2022). We extracted DNA from whole

parental, F1, and F2 hybrid populations, respectively, at seven time points each (0, 100, 200, 300, 400,

500, 700, and 1000 generations). Extraction was carried out on overnight culture, with ~1mL of culture

used. Library preparation and sequencing were performed at the Science for Life Laboratories

(Stockholm, Sweden), using the Illumina Nextera Flex protocol. Sequencing was done using either

NovaSeq S4-300 (2x150bp) or NovaSeq S-Prime (2x150bp) to an estimated median depth of coverage

per sample of around 30x to 780x. The raw sequences were deposited in the European Nucleotide

Archive, parental populations are available at accession number PRJEB46680, while hybrids are available

at accession number PRJEB83568.

Sequencing read quality and genome alignment

FastQC v.0.11.9 (Andrews 2010) was used to assess the quality of raw sequencing reads and reads were

trimmed using Trimmomatic v0.36 (Bolger et al. 2014). Sequencing reads were aligned to a S288C

reference genome (R64; GenBank accession GCA_000146045.2) using BWA mem v0.7.17-r1188 (Li and

Durbin 2009) and SAMtools v1.11 (Danecek et al. 2021) was used to sort and index the resulting

mapping file. Duplicated reads were flagged and removed using Picard v3.1.0

(http://broadinstitute.github.io/picard/), followed by quality assessment using Qualimap v2.3

(Okonechnikov et al. 2016).

Assessing the genomic mutational landscape

To assess the mutational landscape of each experimentally evolved, parental yeast population and the

F1 and F2 crosses made between them, we measured the amount of variants detected across

mutational scales, ranging from single nucleotide variants to entire chromosome aneuploidies. To

identify small mutational variants (SNV and indels), we used freebayes v1.3.6 (Garrison and Marth 2012)

with default parameters. Identified variants were filtered using VCFtools v0.1.16 (Danecek et al. 2011) to

include only variants with minimum quality (minQ) >30 and sequencing read depth (minDP) >5. SnpEff

v5.2 (Cingolani et al. 2012) was used to assess and annotate the effect of identified mutational variants

on genes using the database of the reference genome R64-1-1.105 with canonical transcripts. To

identify structural variants including insertions, deletions, inversions, tandem duplications and

interchromosomal translocations we used Manta v1.6.0 (Chen et al. 2016) with default parameters.

Structural variant detection is limited by the read pair fragment size. To more comprehensively assess

the genome-wide copy number variation (amplifications and deletions), we used Control-FREEC v11.6

(Boeva et al. 2011; Boeva et al. 2012) with baseline ploidy = 2, a break-point threshold = 0.8 and

coefficient of variation = 0.5 (Scopel et al. 2021). We further optimized the CNV detection by limiting the

expected GC content to between 0.35 and 0.55. Variants were further filtered to only include copy

number calls that were statistically significant (p<0.05) using Wilcoxon Rank Sum Test and a Kolmogorov

Smirnov Test. Two metrics of CNVs were assessed for differences across time and between populations:

CNV load and the fraction of the genome altered. CNV mutational load is the number of CNV events

(amplification or deletion) detected within a population. The fraction of genome altered incorporates

the size of each CNV event to determine the amount of the genome that is altered by CNVs. We also

differentiated between small (<10kb) and large (>10kb) CNVs. Chromosomes were further considered

aneuploid if >60% of the chromosome was identified as amplified or lost.

Calculating population differentiation (F_{st}) and genome similarity (Jaccard index)

We calculated pairwise F_{st} between all divergent- and parallel-evolving parental populations at each of

the sampled timepoints, based on single nucleotide polymorphisms (SNPs). We first combined the vcf

files generated from freebayes using bcftools v1.19 (Danecek et al. 2011), with parallelization of file

compression by gnuparallel v20230422 (Tange 2011), using the default parameters of bcftools merge

and the R package poolfstats v2.2.0 (Gautier et al. 2022) on R v3.4 (R Core Team 2023). First the vcf files

were converted to pooldata objects, using vcf2pooldata with the default parameters. F_{st} was then

calculated using pairwiseFST with the nsnp.per.bjack.block = 150 to calculate the Block-Jackknife

estimation of standard error, using an ANOVA framework (Hivert et al. 2018). Similarities between copy

number variation calls in different populations and through time were assessed based on the Jaccard

similarity coefficient using CNVmetrics R package v3.20 (Belleau et al. 2021; Deschênes et al. 2022).

Results

We experimentally evolved yeast populations for 1000 generations in divergent and parallel environments. The four selection environments were media containing either salt, ethanol, or two different concentrations of lithium acetate. These media significantly differ in how much they reduced the fitness of the founder populations, and we have previously provided phenotypic and genetic evidence that these populations become differentially adapted to these environments (Ament-Velásquez et al. 2022). At regular time points during experimental evolution, we made crosses between the parallel- and divergent-evolving populations (the 'parents') to measure postzygotic reproductive isolation (F1 spore viability) and used whole population genome sequencing to assess their mutational load, i.e. the number and types of structural variation, and other genomic features of the parent, F1, and F2 hybrids.

Parental phenotypic divergence

We phenotypically profiled the evolving parent populations by measuring their growth capacity (yield) in 90 different environments, at increasing times of divergence, using the Scan-o-Matic system for high-throughput microbial phenotyping. These 90 non-selection environments contained the same medium (SC) as the selective environments we used for experimental evolution, but they also each contained an additional growth limiting substance (Table S2). Any phenotypic divergence detected between populations is likely a conservative estimate, as multivariate phenotypes were measured in non-selection environments, and no other traits than growth yield were considered. Also note that we only assessed the phenotypic divergence of evolving parent populations here (not the F1 or F2 crosses), to test for and illustrate their adaptation to the four selection environments due to divergent ecological selection.

As expected, with increasing generations of experiment evolution in different environments, divergent-selected parent populations significantly diverged in phenotype, i.e. their yield differed

327   significantly (regression across all divergent-evolved populations: R = 0.14, p = 0.007; **Figure 2A**). In

328   contrast, parent populations evolving in parallel in the same environment did not diverge phenotypically

329   (R = 9.34e-04, p = 0.81; **Figure 2B**). This suggests that the phenotypic divergence of populations evolving

330   under different selection pressures was caused by ecological differences between the four selection

331   environments.

333
Figure 2: **Phenotypic divergence of evolving parental populations.** Regression of parental phenotypic
divergence (the proportion of 90 environments, in which populations significantly differed in yield)
against the increasing time (in number of generations) that parental population evolved in the
experiment. **A)** Comparisons between divergent-evolved populations in different environments. **B)**
Comparisons between parallel-evolved populations in the same environment.

Genetic population differentiation

When comparing parental populations, genetic population differentiation (genome-wide F_{st}) was

generally larger (mean = 0.25) in pairwise comparisons between divergent-selected (grey boxes in **Figure**

3A) than between parallel-selected populations (mean = 0.19, colored boxes). The only exception were

comparisons between parallel-selected LiAc0.01 populations (purple boxes), which produced slightly

higher F_{st}-values but were not significantly larger than the divergent-selected populations. F_{st} -values

increased with the number of generations of evolution in divergent environments (**Figure 3A**, Figure S1),

except for comparisons between divergent-selected NaCl and LiAc0.02 populations, which tended to

decrease in F_{st} over time. Consistent with their lower F_{st} -values overall, the genomes of parallel-selected

populations remained significantly more similar (using the Jaccard similarity index) than the genomes of

divergent-selected populations (gray boxes in **Figure 3B**). Together, this suggests that the larger genetic

divergence of populations evolving under different selection pressures was caused by ecological

differences between the four selection environments.

354
Figure 3: Parental genomic population differentiation. **A)** Genetic population divergence (F_{st}) against
the number of parental generations in experimental evolution. Boxplots indicate the first and third
quartiles with whiskers extending to the furthest point not exceeding 1.5 x the interquartile range and
outliers beyond this are shown as dots. The solid line indicates the median value. **B)** Genomic similarity
(Jaccard similarity index) against the number of generations. Box plots show means (thick horizontal
line) across all generations of each population type. Circles indicate the mean measurement and the
vertical line indicates the standard error of the mean. 'Divergent' indicates comparisons between
divergent-selected populations. Asterisks indicate statistical significance using Kruskal-Wallis and
pairwise Wilcoxon post hoc tests with Bonferroni corrections (adjusted p-values < 0.001 = ***, < 0.01 =
**, < 0.05 = *). Dashed horizontal lines indicate the measured differentiation between the founder
populations at the beginning of evolution.
366

F2 hybrid viability

To measure reproductive isolation, we mated pairs of strains evolving in different environments

(divergent selection) or the same environments (parallel selection) to generate F1 hybrids. We then

sporulated these to produce F1 spores (gametes), which are their recombined, haploid F2 offspring. We

then measured their viability as the fraction of gametes capable of germination and cell colony

formation as a measure of reproductive isolation between populations. The viability of F1 spores made

between divergent-selected populations was significantly lower than the viability of crosses made

between parallel-selected populations (Wilcoxon rank sum test with continuity correction across all

generations; W = 1395.5, p = 0.046). Analyzing cross types separately, the viability of offspring from

parallel-selected populations was significantly higher than the viability of hybrids made from divergent-

selected parents in two cross types (Kruskal-Wallis: NaCl x EtOH: X^2 = 7.6216, p = 0.022; NaCl x LiAc0.02:

X^2 = 14.209, p = 0.001). In the NaCl x LiAc0.01 cross type, the viability of crosses made from parallel-

selected populations was similar to that of crosses from divergent-selected populations (X^2 = 5.834, p =

0.054).

Following the breakdown of hybrid viability over time, we found that viability between

divergent-selected parents was still relatively high at early time points of divergence (on average 89.1%

± 0.02 at 100 generations) but then decreased substantially between 250 and 500 generations to on

average 66% ± 0.08 (**Figure 4**, top left panel). On the contrary, the viability of crosses between parallel-

selected populations stayed high throughout the duration of experimental evolution. One exception

were the crosses made from parallel-selected populations in LiAc0.02, which dropped in viability after

500 generations, to a similar extent as the divergent-selected NaCl x LiAc0.02 hybrids. We discuss

potential explanations for this below.

390
Figure 4: Analysis of reproductive isolation. The viability of F1 spores (i.e. the recombined haploid F2 offspring), which is an inverse measure of reproductive isolation, against the number of parental generations in experimental evolution. Circles indicate the mean measurement and the vertical line indicates the standard error of the mean. Crosses made between parallel-selected populations evolving in the same environment are coloured (or white in the TOTAL plot), crosses made between divergent-selected populations, evolving in different environments, are grey. Box plots show means (thick horizontal line) across all generations of each cross type. 'Divergent' are the crosses between divergent-selected populations (NaCl vs LiAc0.01, NaCl vs LiAc0.02, NaCl vs EtOH). Asterisks indicate statistical significance using Kruskal-Wallis and pairwise Wilcoxon post hoc tests with Bonferroni corrections (adjusted p-values < 0.001 = ***, < 0.01 = **, < 0.05 = *).

The genome-wide mutational landscape of adaptation and hybridization

In an effort to find the genetic mechanisms potentially underlying postzygotic reproductive isolation, we

assessed the genome-wide mutational landscape of the divergent-evolved parental populations, and the

F1 and F2 hybrid crosses made between them, through all sampled timepoints. We focused on three

major mutational classes: single nucleotide variants (SNVs), large structural variants (SVs), and copy

number variants (CNVs).

We found that all populations showed reduced SNV mutational diversity compared to the

founder population within the first 100 generations of evolution (**Figure 5A**). As expected, most

identified mutational variants were single nucleotide variants, fewer were insertions and deletions

(indels) (Figure S2). The mutational SNV load in LiAc0.02 populations was consistently lower than in

populations evolving in NaCl and in their F1 and F2 hybrids, and remained low through time (Kruskal-

Wallis: X^2 = 44.95, p <0.001, **Figure 5A**). This pattern remained after filtering for mutational variants

estimated by SnpEff to have high impact (Kruskal-Wallis: X^2 = 13.07, p=0.004, **Figure 5C**, Figure S3).

LiAc0.02 populations also showed a lower SNV mutational load than the populations evolving in the

weaker concentration of lithium acetate (LiAc0.01). It is possible that this is the result of stronger

purifying selection in LiAc0.02 populations from the start. But since populations evolving in the stronger

concentration of lithium acetate have likely also undergone fewer mitotic generations, this may also

reflect mutations accumulating primarily as a function of reproductive time. In the other three parental

population types (NaCl, LiAc0.01 and EtOH), SNV mutational load decreased over time (**Figure 5B**) and

the pattern stayed consistent when filtering for only mutations with high impact on fitness (Figure S3A).

Conversely, in the F1 (R = 0.48, p < 0.01, in grey) and F2 hybrid populations (R = 0.57, p < 0.01, in yellow),

we found a total increase in mutational load over time (**Figure 5B**). Thus, the more the parental

populations diverged phenotypically and genetically over time, the more single nucleotide mutational

variants were detected in their resulting F1 and F2 hybrid populations.

Figure 5: The single nucleotide variant mutational landscape of hybridization. A) The SNV mutational
load consists of the number of mutational variants and indels (insertions and deletions) within a
population. The median SNV mutational load of populations is indicated by the colored solid line. Gray
plots show mutational load averaged across all sampled timepoints (TOTAL) for the three types of hybrid
crosses. Other plots show data from crosses made at 7 timepoints of increasing parental divergence
(number of generations). The mean mutational load of the founder population is indicated by the
dashed black line. Boxplots indicate the first and third quartiles with whiskers extending to the furthest
point not exceeding 1.5 x the interquartile range and outliers beyond this are shown as dots. The solid
line indicates the median value. Letters indicate statistical significance using Kruskal-Wallis and pairwise
Wilcoxon post hoc tests with Bonferroni corrections. **B)** Linear regression (Pearson R) between SNV
mutational load and increasing timepoints of parental divergence (number of generations). Only
significant linear relationships are shown. The gray plot includes F1 and F2 hybrids across all crosses. **C)**
SNV mutational load after filtering for high impact mutations on gene function using SnpEff v5.2
(Cingolani et al. 2012). The mutational load is averaged across all timepoints (TOTAL) for each hybrid
cross. The mean mutational load of the founder population is indicated by the dashed black line.

Different letters (A, B and C) indicate statistically significant differences using Kruskal-Wallis and pairwise
Wilcoxon post hoc tests with Bonferroni corrections.

Analysis of structural variants (i.e. insertions, deletions, duplications, inversions and

interchromosomal translocations that are > 8bp) showed that the total number of SVs found in the

evolving parental populations remained comparable to the levels in the founder population (**Figure 6A**).

Unlike the pattern we saw in SNV mutational load, the number of SVs harbored in parental populations

increased over time (**Figure 6C**). This positive linear relationship was consistent for populations evolving

in NaCl (R = 0.72, p < 0.01), LiAc0.01 (R = 0.6, p < 0.01) and LiAc0.02 (R = 0.61, p < 0.01), but not for EtOH

populations, which already had elevated levels of SVs (p < 0.001) early on but then decreased over time

(R = -0.7, p < 0.01; **Figure 6A**).

Strikingly, we found very high levels of SVs in all types of F2 hybrid crosses, right from the

beginning of parental population divergence (Kruskal-Wallis: NaCl x LiAc0.01: X^2 = 22.87, p <0.001; NaCl

x LiAc0.02: X^2 = 30.28, p <0.001; NaCl x EtOH: X^2 = 34.45, p <0.001). In most cases, F2 hybrid populations

harboured almost twice as many structural variants compared to the corresponding parental

populations and the F1 hybrids. This pattern was seen across SV types collectively, but was particularly

prominent for insertions, deletions and interchromosomal translocations (breakend mutations, **Figure**

6B, Figure S4-S8). Duplications were rare compared to deletions and insertions and are therefore not

driving the elevated levels of SVs observed in F2 hybrid genomes (**Figure 6B)**. Because F2 hybrids are

distinguished from F1 hybrids and the parent populations only by having passed through one round of

meiosis, the additional load of structural variants we found in F2 hybrids likely arose due to meiotic

errors.

466
467
Figure 6: The structural variant mutational landscape of hybridization. A) Structural variant (SV)
mutational load consists of the number of deletions, insertions, inversions, tandem duplications and
interchromosomal translocations (> 8bp) detected in the population. Gray plots show SV mutational
load averaged across all sampled timepoints (TOTAL) for the three types of hybrid crosses. Other plots
show data from crosses made at 7 timepoints of increasing parental divergence (number of
generations). The mean mutational load of the founder population is indicated by the dashed black line.
Boxplots indicate the first and third quartiles with whiskers extending to the furthest point not
exceeding 1.5 x the interquartile range and outliers beyond this are shown as dots. The solid line
indicates the median value. Different letters (A, B and C) indicate statistically significant differences
using Kruskal-Wallis and pairwise Wilcoxon post hoc tests with Bonferroni corrections. **B)** Mean
structural variant count of each type of structural variant (deletion, insertion, breakend, inversion and
duplication) for each environment and cross. Node color and size indicates the mean count of each
variant type across all timepoints. **C)** Linear regression (Pearson R) between SV mutational load and
increasing timepoints of parental divergence (number of generations). Only significant linear
relationships are shown. The gray plot includes F1 and F2 hybrids from all crosses.

In addition to structural variants, we also more comprehensively assessed copy number

variations (CNV) across the genome, which are a subtype of structural variation. In the structural

variation analysis, detection of variants was limited by the read pair fragment size. However, copy

number variants can range from small focal amplifications or deletions to chromosome arm, or whole

chromosome aneuploidies. Therefore, we used an approach that employs sequencing read depth and

allelic imbalances to estimate segmented copy number uniformly throughout the genome. On average,

parental populations, F1 and F2 hybrids had a lower CNV mutational load compared to the founder

population (dashed black lines in **Figure 7A**). This pattern remained consistent when including the size of

each copy number variant to estimate the fraction of altered genome content (**Figure 7B**). Most

populations showed moderate levels (< 10%) of total alteration (amplification + deletion) of the genome

by CNV (**Figure 7B**). However, NaCl populations had a larger fraction of their genomes altered in copy

number than parental populations evolved in LiAc and EtOH (pairwise Wilcoxon test: vs LiAc0.01: p

<0.001; vs LiAc0.02: p <0.001; vs EtOH: p =0.012), mainly due to a large fraction of the genome

alterations (> 20%) in three parallel-evolved NaCl populations at three different time points.

Strikingly, the F2 hybrid genomes of all three crosses contained significantly fewer CNVs than

parental populations and most F1 hybrids (**Figure 7A**; Kruskal-Wallis: NaCl x LiAc0.01: X^2 = 30.1, p <

0.001; NaCl x LiAc0.02: X^2 = 24.4, p < 0.001; NaCl x EtOH: X^2 = 37.7, p < 0.001) and F2 hybrid genomes of

all crosses were significantly less altered by CNVs (in % genome) than the parental populations (**Figure

7B**). Overall, we found that amplifications were more common than deletions across populations and

through time (Figure S9). Similar to SNVs and SVs, we found consistent patterns of linear increase or

decrease in CNVs with the number of parental generations of evolution in divergent environments

(**Figure 7C**). In the parental NaCl (R = -0.59, p < 0.01) and LiAc0.02 (R = -0.56, p < 0.01) populations, the

CNV mutational load decreased with the number of generations. In the LiAc0.01 parents, the same

pattern was seen but was not significant (R = -0.37, p = 0.08). Interestingly, CNV mutational load was

often relatively stable until approximately 500 generations, followed by a steep decrease (**Figure 7A,**

7C). In the EtOH parents, we found the opposite pattern, with CNV load increasing with the number of

generations (R=0.72, p <0.01), affecting the F1 hybrid genomes between NaCl and EtOH in the same way

(R = 0.73, p = 0.01).

Lastly, we sought to understand if these patterns of CNV load were driven by small focal variants

or larger multi-kilobase alterations, including whole chromosomal aneuploidies. We found that, as

expected, the majority of variants were small (< 10kb) and contributed significantly to the global pattern

found in CNV load (compare upper row of plots in **Figure 7D** to **Figure 7A**). A closer examination of the

number of larger alterations (> 10kb; middle row of plots in **Figure 7D)** revealed that the F2 hybrid

genomes of two hybrid crosses (NaCl x LiAc0.01, NaCl x LiAc0.02), showed significantly fewer large

alterations (on average 12.1) than their respective parents (on average 20.8) and F1 hybrids (on average

19.4). This pattern is further emphasized when examining exclusively the number of aneuploid

chromosomes (>60% amplified or deleted) found in each sample (lower row of plots in **Figure 7D**). On

average, F2 hybrid genomes carried fewer aneuploidies (mean=0.24 chromosomes) compared to parent

(mean=0.67) and F1 hybrid genomes (mean=0.64).

Figure 7: The copy number variant mutational landscape of hybridization. A) Copy number variant (CNV) mutational load consisting of amplifications, deletions and aneuploidies. Gray plots show CNV mutational load averaged across all sampled timepoints (TOTAL) for the three types of hybrid crosses. Other plots show data from crosses made at 7 timepoints of increasing parental divergence (number of generations). The mean mutational load of the founder population is indicated by the dashed black line. Boxplots indicate the first and third quartiles with whiskers extending to the furthest point not exceeding 1.5 x the interquartile range and outliers beyond this are shown as dots. The solid line indicates the median value. Different letters (A, B and C) indicate statistically significant differences using Kruskal-Wallis and pairwise Wilcoxon post hoc tests with Bonferroni corrections. **B)** The fraction of the genome altered (%) across all sampled timepoints (TOTAL) for the three types of hybrid crosses. Colors, boxplots and statistical significance are consistent with panel A. **C)** Linear regression (Pearson R) between CNV mutational load and increasing timepoints of parental divergence (number of generations). Only significant linear relationships are shown. The gray plot includes F1 and F2 hybrids from all crosses. **D)** The distribution of large (>10kb) CNVs, small (<10kb) CNVs and chromosome aneuploidy (>60% amplified or deleted) across all sampled timepoints for the three types of hybrid crosses. Colors, boxplots and statistical significance are consistent with panel A.

The distribution of mutational events across the genome

After assessing the genome-wide mutational landscape in terms of SNVs, SVs, and CNV mutational load,

we determined the distribution of these mutational events across the genome. We found that the rate

of SNVs was not uniformly distributed, but instead showed chromosomal biases (Kruskal-Wallis: X^2 =

147.28, p <0.001, **Figure 8A**). Chromosomes VI and VIII had significantly higher frequencies of SNVs (16.7

and 14.1 mutations/kb, respectively). Chromosome I and IX also showed slightly elevated SNV

frequencies (10.8 and 9.35 mutations/kb, respectively). Chromosome VII on the other hand showed a

significantly lower frequency of SNVs (2.29 mutations/kb). The remaining chromosomes had

approximately equal frequencies of SNVs (6.39-7.34 mutations/kb). These chromosomal biases in SNV

distribution already existed in the founder population and remained consistent across all parental

lineages and all F1 and F2 hybrids (**Figure 8A**).

The distribution of structural variant (SV) events were also not uniformly distributed throughout

the genome (Kruskal-Wallis: X^2 = 60.06, p <0.001, **Figure 8B**). Similar to the SNV distribution, we found

highly elevated rates of SVs on chromosome VI (0.03 mutations/kb). We also found peaks in the rate of

SVs in chromosomes I, III, and IX in most parental and hybrid populations.

The proportion of whole chromosomes altered by copy number variants was not consistent

across environments and hybrid crosses (**Figure 8C**). Interestingly, F2 hybrids of all crosses showed the

lowest levels of copy number alterations, and aneuploidy (defined as > 60% of chromosome

amplification) was consistently rare across all F2 hybrid chromosomes. We discuss the implications of

the near absence of aneuploidies and large genome alterations in F2 hybrid genomes below. Our

analysis shows that the two founding populations each carried aneuploid chromosomes. The NaCl

founder population (N founder, **Figure 8C**) had an extra chromosome X, while the LE founder population

had three aneuploid chromosomes (I, III and VI). Parental NaCl populations subsequently lost the extra

chromosome X over the course of experimental evolution, but evolved an aneuploid chromosome I and

underwent alterations of large portions of chromosome III and VI. Chromosome I was also heavily

altered in parent populations evolving in LiAc 0.01 (40%), LiAc 0.02 (37%), and EtOH (25%). Similarly, F1

hybrid crosses between NaCl and the two LiAc populations (to a lesser extent the NaCl x EtOH F1

hybrids) also showed large copy number alterations in chromosomes I, III, and VI (**Figure 8C**). Confirming

our previous result (Ament-Velásquez et al. 2022), we also recovered a chromosome X aneuploidy in

two parental EtOH populations and the chromosome XI aneuploidy in one parental EtOH population

here.

The 16 chromosomes of *S. cerevisiae* range widely in size from 270kb (chrVI) to 1,531kb (chrIV;

Figure 8D). As might be expected, larger chromosomes are larger targets for SNVs, SVs and CNVs and

therefore have higher overall mutational load for each mutational class. Even after correcting

mutational load for chromosome size, either by reporting SNVs and SVs as mutations per kilobase or

reporting CNVs as the percentage of chromosome altered, we still found a strong negative relationship

between chromosome size and SNV, SV and CNV mutational rate (**Figure 8D**). Smaller chromosomes had

elevated rates of mutation while larger chromosomes were rarely affected. This pattern is most striking

in CNVs where, as previously mentioned, chromosomes I, III and VI were disproportionately altered. This

is interesting because chromosomes I, III, and VI are the smallest chromosomes in the yeast genome and

have been frequently reported to carry aneuploidies in other yeast studies as well (reviewed in Gilchrist

and Stelkens 2019), likely because aneuploidies of smaller chromosomes cause fewer deleterious gene

dosage effects and can be compensated for at the proteome level (Muenzner et al. 2024). This pattern

of chromosome size bias for CNVs extends beyond intraspecific crosses and has been found across

interspecific yeast hybrids with diverse ancestries (Bendixsen et al. 2021).

Figure 8: The distribution of mutational events across the genome. A) The mean number of events
(mutations/kb) for each chromosome across all available timepoints. The top graph shows the
distribution of mean SNV counts for each chromosome, while the right graph shows the distribution for
each environment and hybrid cross. Boxplots indicate the first and third quartiles with whiskers
extending to the furthest point not exceeding 1.5 x the interquartile range and outliers beyond this are
shown as dots. The solid line indicates the median value. Different letters (A, B, C, D) indicate statistically
significant differences using Kruskal-Wallis and pairwise Wilcoxon post hoc tests with Bonferroni
corrections. **B)** The mean number of structural variant (SV) events (mutations/kb). **C)** The mean amount
of genome altered (%) by copy number variants (losses and gains), including whole chromosome
aneuploidies. **D)** The relationship between chromosome size and SNVs, SVs or CNVs. The barplot
indicates the size of each of the 16 chromosomes in *Saccharomyces cerevisiae* reported in kilobases. The
linear correlations show the mean SNV (muts/kb), SV (muts/kb) or CNV (% genome altered) for each
population and for each chromosome as shown in the top graphs of panels A, B and C. The colors of
each node indicate the populations (NaCl, LiAc0.01, LiAc0.02, EtOH, Hybrid F1, Hybrid F2).

Discussion

We used experimental evolution with budding yeast to test for the evolution of reproductive isolation as

a result of ecologically divergent selection, a process called ecological speciation (Mayr 1947; Schluter

2001; McKinnon et al. 2004; Nosil 2012; Jarrett et al. 2024). Previous studies with yeast have tested for

reproductive isolation between species that have diverged millions of years ago, which translates into

billions of generations (Greig et al. 2002; Shapira et al. 2014; Stelkens et al. 2014; Bernardes et al. 2017;

Eberlein et al. 2019; Gallone et al. 2019; Zhang et al. 2020; Bautista et al. 2021; Brice et al. 2021;

Bautista et al. 2023). Here, instead of starting with already divergent species, separated by billions of

generations, each with a complex evolutionary history of selection, genetic drift, recombination, and

gene flow, we generated phenotypic and genetic divergence over only a few hundred generations. We

evolved replicate populations in allopatry, in four different environments, under both ecologically

divergent and parallel selection. Hence, we controlled the type and degree of selection, minimized

genetic drift and recombination, and excluded gene flow. We systematically produced hybrid crosses

between these populations at seven time points of experimental evolution, to test whether

reproductive isolation evolved between the divergent-selected populations, as predicted by ecological

speciation theory.

First, we confirmed that populations when evolved in different environments, increasingly

diverged in phenotypic traits that were not directly under selection. The parallel-selected (control)

populations on the other hand remained phenotypically similar (**Figure 2**). This provides evidence for

large scale phenotypic differentiation under divergent selection. Second, the genomes of divergent-

selected populations became more genetically differentiated over time, showing higher F_{st} - values and

lower similarity (Jaccard index) than those of the parallel-selected populations (**Figure 3**). Third, we

found that divergent-selected populations produced hybrid gametes with reduced viability, which is a

classic signature of postzygotic reproductive isolation in the form of hybrid breakdown (**Figure 4**).

Zygotes generated from parallel-selected populations on the other hand remained reproductively

compatible (**Figure 4**). Together, our results suggest that partial reproductive isolation evolved due to

divergent ecological selection in our experiment. Alternative processes, such as mutation-order effects

(Schluter 2009; Ono et al. 2017), may have contributed to the build up of reproductive barriers, but

since most parallel-selected populations remained compatible, neither the timing nor the order in which

mutations occurred likely played a large role.

Early experiments, e.g. in fruit flies with similar set-ups of divergent- and parallel-selected lines

on different media (Dodd 1989) and a large body of theory (reviewed in Kirkpatrick and Ravigné 2002)

have demonstrated the evolution of reproductive isolation as a consequence of adaptive divergence

(Nosil 2012). Here, we have taken additional steps to understand the genetic basis of this hybrid

breakdown. Our population genomic analyses show that, as parental populations diverged

phenotypically and genetically over time, the mutational load of single nucleotide variants (SNVs)

steadily increased in F1 and F2 hybrids (**Figure 5**), with a bias for chromosomes I, VI, and VIII (**Figure 8A**).

This may point to increasingly defective DNA repair pathways in hybrid genomes. But even more

strikingly, F2 hybrid populations harboured more than twice as many structural variants (SVs) than their

respective parental populations and the F1 hybrids. This pattern was seen across all SV types (**Figure 6**),

but was particularly strong for insertions, deletions, and interchromosomal translocations (breakend

mutations, Figure S4-S8) with strong chromosomal biases for chromosomes I, III, VI, and IX (**Figure 8B**).

Because F2 hybrids differ from F1 hybrids only by one meiotic event (and a few mitotic generations

before and after sporulation that are negligible), the increase in structural variants seen in F2 hybrids

can only reasonably be explained by errors during meiosis. This suggests that ecological divergence has

led to hybrid offspring with higher genomic instability due to meiotic recombination errors (e.g. unequal

or incorrect crossing overs), resulting in frequent chromosomal rearrangements (SVs).

Interestingly, F2 hybrids showed the lowest levels of copy number alterations, and aneuploidy

rates were consistently low across all chromosomes in F2 hybrid genomes (**Figure 7, Figure 8C**). This is

surprising, because in contrast to the high SV load found in hybrid genomes (**Figure 6**), errors during

meiosis seem to not have led to a detectable increase in chromosome non-disjunction and aneuploidy.

Meanwhile, all parent and F1 hybrid populations contained various aneuploidies, especially in

chromosomes I, III, and VI, which are the smallest chromosomes in the yeast genome and frequently

reported to carry aneuploidies (reviewed in (Gilchrist and Stelkens 2019). A plausible explanation for the

surprising absence of aneuploidies in F2 hybrids is that, by default, we only sequenced viable genotypes

that had successfully passed through meiosis and germinated. Average mortality due to lethal

aneuploidies is likely high in the recombined haploid gametes, where copy number imbalances imposed

by aneuploidies are more pronounced, and lethality has additional opportunities to manifest when

strains are pushed through germination. As a result, aneuploidies may have been efficiently purged from

F2 hybrid genomes before the sequencing stage.

Experimental and genetic work in interspecific yeast crosses, made from species separated by

billions of generations and high nucleotide divergence (>10%), has shown that the main reason for

hybrid breakdown in yeast is antirecombination (Hunter et al. 1996; Rogers et al. 2018; Bozdag et al.

2021). Antirecombination suppresses crossovers during meiosis when nucleotide divergence along

chromosomes is too high. The ensuing mispairing of homologous chromosomes causes missegregation

into the hybrid gametes, leading to lethal aneuploidies and typically > 90% spore mortality in

interspecific yeast crosses. This process has also been shown to cause hybrid sterility in mice, where it is

controlled by the Prdm9 gene (Forejt and Jansa 2023). However, antirecombination due to genome-

wide nucleotide divergence is unlikely to have played a role in the hybrid breakdown we observed here,

given the short evolutionary time span of our experiment. Instead, we suggest that the following

process underlies the build up of reproductive barriers here: First, divergent ecological selection in the

parent populations has led to SVs with positive (adaptive) fitness effects and additional neutral SVs

hitchhiking along with them. These parental SVs then interfered with hybrid meiosis, causing additional

SVs to emerge in the hybrid genomes, due to unequal or incorrect crossing over. Cumulatively, the high

SV load of the hybrid genomes then negatively affected gamete viability, germination, and colony

formation, disrupting cell function and homeostasis (Chang et al. 2013; Liu et al. 2022). In short, we

suggest that the most likely genetic mechanism underlying hybrid breakdown in our experiment is the

high structural variant load of the hybrid genomes, which was jump-started by the increasing number of

SVs accumulating in the divergent-selected parent populations (**Figure 6**).

We saw one interesting exception in crosses between parallel-selected LiAc0.02 populations

where, against our expectations, gamete viability decreased over time (**Figure 4**). We previously found

that three of four replicate populations evolving in LiAc0.02 had turned (functionally) haploid early on in

experimental evolution (Ament-Velásquez et al. 2022). In haploid populations, beneficial mutations can

sweep to fixation faster and deleterious mutations are more quickly purged by purifying selection, as

they are not masked by heterozygosity (Gerstein et al. 2011; Cervantes et al. 2023). These processes are

particularly efficient in large populations exposed to stressful environments with strong selection, which

the LiAc0.02 environment represents (Ament-Velásquez et al. 2022). Propelled by clonal interference,

this may have affected the adaptive dynamics (the rate and direction of allele frequency changes) in the

LiAc0.02 populations in idiosyncratic ways, leading to substantial genetic differentiation and

incompatibilities between them, despite parallel selection pressures.

But what is the genetic architecture of the partial reproductive isolation in our experiment? The

pattern of decreasing genomic similarity and increasing F_{st} between divergent-selected populations

(while parallel-selected populations remained similar), and the vast genome-wide instability found in

hybrid genomes, does suggest that reproductive isolation is based on multiple genomic regions and that

'continents' rather than 'islands' of differentiated loci are operating together (Michel et al. 2010).

However, due to the absence of sexual recombination during the evolution of the parent populations in

our experiment, other evolutionary mechanisms may have also caused large genomic regions to become

differentiated. Effects of clonal interference and linked selection for example, i.e. the competition

between multiple beneficial mutations arising independently in the same population leading to only one

lineage sweeping through the population, may have caused genetic differentiation between populations

that is not due to ecological selection alone. Although our previous work has identified the number and

types of adaptive mutations segregating and fixing in the evolving parental populations (Ament-

Velásquez et al. 2022), our pool-seq approach does not provide haplotype information. Thus, we cannot

infer which and how many genes are causing reproductive isolation, how large their effect sizes are, and

whether they are genomically clustered (Nosil and Feder 2012). Directed crosses between parents with

known genotypes, and dissecting and genotyping a large number of recombined F2 gametes, could

identify incompatible chromosomes, chromosomal regions, or individual genes causing reproductive

isolation in this system.

Conclusions

Selection against hybrids due to ecological factors, when hybrid phenotypes are ill-adapted to existing

ecological niches, has traditionally been considered as being different from selection arising from

epistatic (Dobzhansky-Muller) incompatibilities (Thompson et al. 2023). There is a recent push to close

this gap between evolutionary ecology and speciation genetics, and to focus more on the role of the

environment and ecological adaptation in driving what we traditionally think of as 'intrinsic' genetic

isolation barriers (Anderson et al. 2023; Thompson et al. 2023). Our study contributes to bridging this

gap by synthesizing phenotypic and whole genome data, and demonstrating that divergent ecological

adaptation can indeed promote intrinsic reproductive isolation, consistent with predictions of ecological

speciation theory. Our population genomics results show that divergent ecological selection, even for

relatively few mitotic generations, can lead to vast genomic instability in F2 hybrids, including deletions,

inversions, and interchromosomal translocations. Thus, our multifaceted approach of combining

experimental evolution with sexual microbes, hybridization at increasing time points of divergence,

high-throughput phenotyping, and comparative genomics, confirms the role of ecology in driving

reproductive isolation and speciation. Our approach can be further developed to provide a

comprehensive empirical framework to test different concepts in speciation research. This includes, for

instance, i) testing if the speciation continuum is a 'continuum of reproductive isolation' (Stankowski

and Ravinet 2021), by measuring whether the accumulation of reproductive isolating barriers is gradual

over extended evolutionary time frames, ii) testing if the 'speciation hypercube' captures the complexity

of species divergence (Dieckmann et al. 2004; Bolnick et al. 2023), by including more phenotypic and

genetic variables, and iii) to better understand the evolution of hybrid fitness during speciation by

including additional data sets (e.g. gene interaction networks (Dagilis et al. 2019), which could reconcile

the contrasting outcomes of epistatic genetic interactions leading to hybrids with both increased

(heterosis) and decreased fitness (hybrid inviability) during different stages of speciation.

Acknowledgements

We thank Maria de la Paz Celorio Mancera, Viktoria Köppä, and Zebin Zhang for helping with serial

transfers and media preparation, and Lorena Ament-Velásquez for bioinformatics advice. Computation

was performed on resources provided by SNIC through Uppsala Multidisciplinary Center for Advanced

Computational Science (UPPMAX) under Projects NAISS2023/23-519 (Small Storage) and NAISS2023/22-

62 and NAISS2023/22-924 (Small Compute). We acknowledge support from Science for Life Laboratory,

the National Genomics Infrastructure, NGI, and Uppmax for providing assistance in massive parallel

sequencing and computational infrastructure. This work was supported by Swedish Research Council

Project Grants (2022-03427 to R.S. and 2022-03024 to J.W.), a Knut and Alice Wallenberg Foundation

Grant (2017.0163 to R.S.), and the Wenner-Gren Foundations (UPD2018-0196, UPD2019-0110 to D.B.).

References

Alalam H, Graf FE, Palm M, Abadikhah M, Zackrisson M, Boström J, Fransson A, Hadjineophytou C,
Persson L, Stenberg S, et al. 2020. A High-Throughput Method for Screening for Genes
Controlling Bacterial Conjugation of Antibiotic Resistance. *mSystems* 5:e01226-20.
Ament-Velásquez SL, Gilchrist C, Rêgo A, Bendixsen DP, Brice C, Grosse-Sommer JM, Rafati N, Stelkens R.
2022. The Dynamics of Adaptation to Stress from Standing Genetic Variation and de novo
Mutations. *Molecular Biology and Evolution* 39:msac242.
Anderson SAS, López-Fernández H, Weir JT. 2023. Ecology and the Origin of Nonephemeral Species. *The*
American Naturalist 201:619–638.
Anderson SAS, Weir JT. 2022. The role of divergent ecological adaptation during allopatric speciation in
vertebrates. *Science* 378:1214–1218.
Andrews S. 2010. FastQC: A Quality Control tool for High Throughput Sequence Data. Available from:
https://www.bioinformatics.babraham.ac.uk/projects/fastqc/
Bautista C, Gagnon-Arsenault I, Utrobina M, Fijarczyk A, Bendixsen DP, Stelkens R, Landry C. 2023.
Hybrid adaptation is hampered by Haldane's sieve. *bioRxiv*:2023–12.
Bautista C, Marsit S, Landry CR. 2021. Interspecific hybrids show a reduced adaptive potential under
DNA damaging conditions. *Evolutionary Applications* 14:758–769.
Bell G, Gonzalez A. 2011. Adaptation and Evolutionary Rescue in Metapopulations Experiencing
Environmental Deterioration. *Science* 332:1327–1330.
Belleau P, Deschênes A, Beyaz S, Tuveson DA, Krasnitz A. 2021. CNVMetrics package: Quantifying
similarity between copy number profiles. *F1000Research* [Internet] 10. Available from:
https://f1000research.com/slides/10-737
Bendixsen Devin P, Gettle N, Gilchrist C, Zhang Z, Stelkens R. 2021. Genomic Evidence of an Ancient East
Asian Divergence Event in Wild Saccharomyces cerevisiae. *Genome Biology and Evolution*
13:evab001.
Bendixsen, Peris D, Stelkens R. 2021. Patterns of Genomic Instability in Interspecific Yeast Hybrids With
Diverse Ancestries. *Frontiers in Fungal Biology* [Internet] 2. Available from:
https://www.frontiersin.org/article/10.3389/ffunb.2021.742894
Bernardes JP, Stelkens RB, Greig D. 2017. Heterosis in hybrids within and between yeast species. *J Evol*
Biol 30:538–548.
Boeva V, Popova T, Bleakley K, Chiche P, Cappo J, Schleiermacher G, Janoueix-Lerosey I, Delattre O,
Barillot E. 2012. Control-FREEC: a tool for assessing copy number and allelic content using next-
generation sequencing data. *Bioinformatics* 28:423–425.
Boeva V, Zinovyev A, Bleakley K, Vert J-P, Janoueix-Lerosey I, Delattre O, Barillot E. 2011. Control-free
calling of copy number alterations in deep-sequencing data using GC-content normalization.
Bioinformatics 27:268–269.
Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics 30:2114–2120.
Bolnick DI, Hund AK, Nosil P, Peng F, Ravinet M, Stankowski S, Subramanian S, Wolf JBW, Yukilevich R.
2023. A multivariate view of the speciation continuum. *Evolution* 77:318–328.
Bozdag GO, Ono J, Denton JA, Karakoc E, Hunter N, Leu J-Y, Greig D. 2021. Breaking a species barrier by

788   enabling hybrid recombination. *Current Biology* 31:R180–R181.

789 Brice C, Zhang Z, Bendixsen D, Stelkens R. 2021. Hybridization Outcomes Have Strong Genomic and
790   Environmental Contingencies. *The American Naturalist* 198:E53–E67.

791 Cervantes S, Kesälahti R, Kumpula TA, Mattila TM, Helanterä H, Pyhäjärvi T. 2023. Strong Purifying
792   Selection in Haploid Tissue–Specific Genes of Scots Pine Supports the Masking Theory.
793   *Molecular Biology and Evolution* 40:msad183.

794 Chang S-L, Lai H-Y, Tung S-Y, Leu J-Y. 2013. Dynamic Large-Scale Chromosomal Rearrangements Fuel
795   Rapid Adaptation in Yeast Populations. *PLOS Genetics* 9:e1003232.

796 Chen X, Schulz-Trieglaff O, Shaw R, Barnes B, Schlesinger F, Källberg M, Cox AJ, Kruglyak S, Saunders CT.
797   2016. Manta: rapid detection of structural variants and indels for germline and cancer
798   sequencing applications. *Bioinformatics* 32:1220–1222.

799 Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM. 2012. A program for
800   annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the
801   genome of Drosophila melanogaster strain w1118; iso-2; iso-3. *Fly (Austin)* 6:80–92.

802 Dagilis AJ, Kirkpatrick M, Bolnick DI. 2019. The evolution of hybrid fitness during speciation. *PLOS*
803   *Genetics* 15:e1008125.

804 Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth GT,
805   Sherry ST, et al. 2011. The variant call format and VCFtools. *Bioinformatics* 27:2156–2158.

806 Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T, McCarthy SA,
807   Davies RM, et al. 2021. Twelve years of SAMtools and BCFtools. *GigaScience* 10:giab008.

808 Deschênes A, Belleau P, Tuveson DA, Krasnitz A. 2022. Quantifying similarity between copy number
809   profiles with CNVMetrics package. *F1000Research* [Internet] 11. Available from:
810   https://f1000research.com/posters/11-816

811 Dettman JR, Anderson JB, Kohn LM. 2008. Divergent adaptation promotes reproductive isolation among
812   experimental populations of the filamentous fungus Neurospora. *BMC Evol Biol* 8:35.

813 Dettman JR, Sirjusingh C, Kohn LM, Anderson JB. 2007. Incipient speciation by divergent adaptation and
814   antagonistic epistasis in yeast. *Nature* 447:585–588.

815 deVicente MC, Tanksley SD. 1993. Qtl Analysis of Transgressive Segregation in an Interspecific Tomato
816   Cross. *Genetics* 134:585–596.

817 Dieckmann U, Doebeli M, Metz JAJ, Tautz D eds. 2004. Adaptive Speciation. Cambridge: Cambridge
818   University Press Available from: https://www.cambridge.org/core/books/adaptive-
819   speciation/A2413E0B1E5DDAB5C5848F0666468A35

820 Dodd DMB. 1989. Reproductive Isolation as a Consequence of Adaptive Divergence in Drosophila
821   pseudoobscura. *Evolution* 43:1308–1311.

822 Eberlein C, Hénault M, Fijarczyk A, Charron G, Bouvier M, Kohn LM, Anderson JB, Landry CR. 2019.
823   Hybridization is a recurrent evolutionary stimulus in wild yeast speciation. *Nat Commun* 10:923.

824 Filchak KE, Roethele JB, Feder JL. 2000. Natural selection and sympatric divergence in the apple maggot
825   Rhagoletis pomonella. *Nature* 407:739–742.

826 Forejt J, Jansa P. 2023. Meiotic Recognition of Evolutionarily Diverged Homologs: Chromosomal Hybrid
827   Sterility Revisited. *Molecular Biology and Evolution* 40:msad083.

828 Funk DJ, Nosil P, Etges WJ. 2006. Ecological divergence exhibits consistently positive associations with
829   reproductive isolation across disparate taxa. *Proceedings of the National Academy of Sciences*
830   103:3209–3213.

831 Gallone B, Steensels J, Mertens S, Dzialo MC, Gordon JL, Wauters R, Theßeling FA, Bellinazzo F, Saels V,
832   Herrera-Malaver B, et al. 2019. Interspecific hybridization facilitates niche adaptation in beer
833   yeast. *Nat Ecol Evol* 3:1562–1575.

Garrison E, Marth G. 2012. Haplotype-based variant detection from short-read sequencing. *arXiv* 1207.

Gautier M, Vitalis R, Flori L, Estoup A. 2022. f-Statistics estimation and admixture graph construction

with Pool-Seq or allele count data using the R package poolfstat. *Mol Ecol Resour* 22:1394–1416.

Gerstein AC, Cleathero LA, Mandegar MA, Otto SP. 2011. Haploids adapt faster than diploids across a
range of environments. *Journal of Evolutionary Biology* 24:531–540.

Gilchrist C, Stelkens R. 2019. Aneuploidy in yeast: Segregation error or adaptation mechanism? *Yeast*
36:525–539.

Good BH, McDonald MJ, Barrick JE, Lenski RE, Desai MM. 2017. The dynamics of molecular evolution
over 60,000 generations. *Nature* 551:45–50.

Gorter FA, Derks MF, van den Heuvel J, Aarts MG, Zwaan BJ, de Ridder D, de Visser JAG. 2017. Genomics
of adaptation depends on the rate of environmental change in experimental yeast populations.
Molecular Biology and Evolution 34:2613–2626.

Gorter FA, Scanlan PD, Buckling A. 2016. Adaptation to abiotic conditions drives local adaptation in
bacteria and viruses coevolving in heterogeneous environments. *Biol. Lett.* 12:20150879.

Grant RB, Grant PR. 2003. What Darwin's Finches Can Teach Us about the Evolutionary Origin and
Regulation of Biodiversity. *BioScience* 53:965–975.

Greig D, Louis EJ, Borts RH, Travisano M. 2002. Hybrid Speciation in Experimental Populations of Yeast.
Science 298:1773–1775.

Härer A, Bolnick DI, Rennison DJ. 2021. The genomic signature of ecological divergence along the
benthic-limnetic axis in allopatric and sympatric threespine stickleback. *Molecular Ecology*
30:451–463.

Hatfield T, Schluter D. 1999. ECOLOGICAL SPECIATION IN STICKLEBACKS: ENVIRONMENT-DEPENDENT
HYBRID FITNESS. *Evolution* 53:866–873.

Hendry AP, Nosil P, Rieseberg LH. 2007. The speed of ecological speciation. *Functional Ecology* 21:455–
464.

Hivert V, Leblois R, Petit EJ, Gautier M, Vitalis R. 2018. Measuring Genetic Differentiation from Pool-seq
Data. *Genetics* 210:315–330.

Hunter N, Chambers SR, Louis EJ, Borts RH. 1996. The mismatch repair system contributes to meiotic
sterility in an interspecific yeast hybrid. *EMBO J* 15:1726–1733.

Irestedt M, Müller I, Thörn F, Joseph L, Nylander J, Guinet B, Valk TV der, Jønsson K. Reticulate and
hybrid speciation is promoted by environmental instability in an Indo-Pacific species complex of
whistlers (Aves: Pachycephala). Available from:
https://www.authorea.com/doi/full/10.22541/au.173053090.08198521?commit=f442f52dffaef
d5b78602e94558c1591a8dff4e5; accepted for publication in Molecular Ecology, Special Issue on
Genomics of Speciation.

Jarrett BJM, Downing PA, Svensson EI. 2024. Divergent selection and plasticity shape reproductive
isolation at the onset of speciation. :2024.09.23.614431. Available from:
https://www.biorxiv.org/content/10.1101/2024.09.23.614431v1

Johnson MS, Desai MM. 2022. Mutational robustness changes during long-term adaptation in laboratory
budding yeast populations.Cooper VS, Walczak AM, Miller C, editors. *eLife* 11:e76491.

Kirkpatrick M, Ravigné V. 2002. Speciation by natural and sexual selection: models and experiments. *Am
Nat* 159 Suppl 3:S22-35.

Kosheleva K, Desai MM. 2018. Recombination Alters the Dynamics of Adaptation on Standing Variation
in Laboratory Yeast Populations. *Mol Biol Evol* 35:180–201.

Lenski RE, Rose MR, Simpson SC, Tadler SC. 1991. Long-Term Experimental Evolution in Escherichia coli.
I. Adaptation and Divergence During 2,000 Generations. *The American Naturalist* 138:1315–
1341.

Leu J-Y, Chang S-L, Chao J-C, Woods LC, McDonald MJ. 2020. Sex alters molecular evolution in diploid
experimental populations of S. cerevisiae. *Nat Ecol Evol* 4:453–460.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler transform.

884    *Bioinformatics* 25:1754–1760.

885 Li J, Vázquez-García I, Persson K, González A, Yue J-X, Barré B, Hall MN, Long A, Warringer J, Mustonen V,
886    et al. 2019. Shared Molecular Targets Confer Resistance over Short and Long Evolutionary
887    Timescales. *Molecular Biology and Evolution* 36:691–708.

888 Liu Y, Ma G, Gao Z, Li J, Wang J, Zhu X, Ma R, Yang J, Zhou Y, Hu K, et al. 2022. Global chromosome
889    rearrangement induced by CRISPR-Cas9 reshapes the genome and transcriptome of human cells.
890    *Nucleic Acids Research* 50:3456–3474.

891 Louder MIM, Justen H, Kimmitt AA, Lawley KS, Turner LM, Dickman JD, Delmore KE. 2024. Gene
892    regulation and speciation in a migratory divide between songbirds. *Nat Commun* 15:98.

893 Mayr E. 1947. Ecological Factors in Speciation. *Evolution* 1:263–288.

894 McDonald MJ, Rice DP, Desai MM. 2016. Sex speeds adaptation by altering the dynamics of molecular
895    evolution. *Nature* 531:233–236.

896 McKinnon JS, Mori S, Blackman BK, David L, Kingsley DM, Jamieson L, Chou J, Schluter D. 2004. Evidence
897    for ecology's role in speciation. *Nature* 429:294–298.

898 Meyer JR, Dobias DT, Medina SJ, Servilio L, Gupta A, Lenski RE. 2016. Ecological speciation of
899    bacteriophage lambda in allopatry and sympatry. *Science* 354:1301–1304.

900 Michel AP, Sim S, Powell THQ, Taylor MS, Nosil P, Feder JL. 2010. Widespread genomic divergence
901    during sympatric speciation. *Proceedings of the National Academy of Sciences* 107:9724–9729.

902 Muenzner J, Trébulle P, Agostini F, Zauber H, Messner CB, Steger M, Kilian C, Lau K, Barthel N, Lehmann
903    A, et al. 2024. Natural proteome diversity links aneuploidy tolerance to protein turnover. *Nature*
904    630:149–157.

905 Mukherjee V, Lind U, St. Onge RP, Blomberg A, Nygård Y. 2021. A CRISPR Interference Screen of Essential
906    Genes Reveals that Proteasome Regulation Dictates Acetic Acid Tolerance in Saccharomyces
907    cerevisiae. *mSystems* 6:10.1128/msystems.00418-21.

908 Müller I, Thörn F, Rajan S, Olsen R-A, Ericson P, Peona V, Smith B, Maiah G, Koane B, Iova B, et al.
909    Ephemeral speciation in a New Guinean honeyeater complex (Aves: Melidectes). Available from:
910    https://www.authorea.com/doi/full/10.22541/au.173639442.28305532?commit=c6fffa9cc0ca7
911    1d3e44f2de72f5001a07ea2ab20; accepted for publication in Molecular Ecology, Special Issue on
912    Genomics of Speciation.

913 Munar-Delgado G, Pulido F, Edelaar P. 2024. Performance-based habitat choice can drive rapid adaptive
914    divergence and reproductive isolation. *Current Biology* 34:5564-5569.e4.

915 Nosil P. 2012. Ecological Speciation. Oxford University Press Available from:
916    https://academic.oup.com/book/5499

917 Nosil P, Crespi BJ. 2006. Experimental evidence that predation promotes divergence in adaptive
918    radiation. *Proceedings of the National Academy of Sciences* 103:9090–9095.

919 Nosil P, Feder JL. 2012. Genomic divergence during speciation: causes and consequences. *Philosophical*
920    *Transactions of the Royal Society B: Biological Sciences* 367:332–342.

921 Nosil P, Schluter D. 2011. The genes underlying the process of speciation. *Trends Ecol Evol* 26:160–167.

922 Okonechnikov K, Conesa A, García-Alcalde F. 2016. Qualimap 2: advanced multi-sample quality control
923    for high-throughput sequencing data. *Bioinformatics* 32:292–294.

924 Ono J, Gerstein AC, Otto SP. 2017. Widespread Genetic Incompatibilities between First-Step Mutations
925    during Parallel Adaptation of Saccharomyces cerevisiae to a Common Environment. *PLOS*
926    *Biology* 15:e1002591.

927 Persson K, Onyema VO, Nwafor IP, Peri KVR, Otti C, Nnaemeka P, Onyishi C, Okoye S, Moneke A, Amadi
928    O, et al. 2024. Lactose-assimilating yeasts with high fatty acid accumulation uncovered by
929    untargeted bioprospecting. *Applied and Environmental Microbiology* 0:e01615-24.

930 Persson K, Stenberg S, Tamás MJ, Warringer J. 2022. Adaptation of the yeast gene knockout collection is
931    near-perfectly predicted by fitness and diminishing return epistasis. *G3*

Genes|Genomes|Genetics 12:jkac240.

Peter J, De Chiara M, Friedrich A, Yue J-X, Pflieger D, Bergström A, Sigwalt A, Barre B, Freel K, Llored A, et
al. 2018. Genome evolution across 1,011 Saccharomyces cerevisiae isolates. *Nature* 556:339–
344.

R Core Team. 2023. R: A Language and Environment for Statistical Computing. Available from:
https://www.R-project.org/

Rainey PB, Travisano M. 1998. Adaptive radiation in a heterogeneous environment. *Nature* 394:69–72.

Rogers DW, McConnell E, Ono J, Greig D. 2018. Spore-autonomous fluorescent protein expression
identifies meiotic chromosome mis-segregation as the principal cause of hybrid sterility in yeast.
PLOS Biology 16:e2005066.

Rundle HD, Nagel L, Wenrick Boughman J, Schluter D. 2000. Natural selection and parallel speciation in
sympatric sticklebacks. *Science* 287:306–308.

Sawada A, Iwasaki T, Akatani K, Takagi M. 2022. Mate choice for body size leads to size assortative
mating in the Ryukyu Scops Owl Otus elegans. *Ecology and Evolution* 12:e9578.

Schluter D. 2001. Ecology and the origin of species. *Trends Ecol Evol* 16:372–380.

Schluter D. 2009. Evidence for Ecological Speciation and Its Alternative. *Science* 323:737–741.

Scopel EFC, Hose J, Bensasson D, Gasch AP. 2021. Genetic variation in aneuploidy prevalence and
tolerance across Saccharomyces cerevisiae lineages. *Genetics* 217:iyab015.

Shapira R, Levy T, Shaked S, Fridman E, David L. 2014. Extensive heterosis in growth of yeast hybrids is
explained by a combination of genetic models. *Heredity (Edinb)* 113:316–326.

Sparks MM, Kraft JC, Blackstone KMS, McNickle GG, Christie MR. 2022. Large genetic divergence
underpins cryptic local adaptation across ecological and evolutionary gradients. *Proceedings of
the Royal Society B: Biological Sciences* 289:20221472.

Stankowski S, Ravinet M. 2021. Defining the speciation continuum. *Evolution* 75:1256–1273.

Stelkens R. 2024. A microbial perspective on speciation. *Evolutionary Journal of the Linnean Society*
3:kzae023.

Stelkens RB, Brockhurst MA, Hurst GDD, Miller EL, Greig D. 2014. The effect of hybrid transgression on
environmental tolerance in experimental yeast crosses. *J. Evol. Biol.* 27:2507–2519.

Stenberg S, Li J, Gjuvsland AB, Persson K, Demitz-Helin E, González Peña C, Yue J-X, Gilchrist C, Ärengård
T, Ghiaci P, et al. 2022. Genetically controlled mtDNA deletions prevent ROS damage by
arresting oxidative phosphorylation.Gruber J, Tyler JK, Toledano MB, editors. *eLife* 11:e76095.

Struck TH, Feder JL, Bendiksby M, Birkeland S, Cerca J, Gusarov VI, Kistenich S, Larsson K-H, Liow LH,
Nowak MD, et al. 2018. Finding Evolutionary Processes Hidden in Cryptic Species. *Trends in
Ecology & Evolution* 33:153–163.

Tange O. 2011. Gnu parallel-the command-line power tool. *Usenix Mag* 36:42.

Thompson KA, Brandvain Y, Coughlan JM, Delmore KE, Justen H, Linnen CR, Ortiz-Barrientos D,
Rushworth CA, Schneemann H, Schumer M, et al. 2023. The Ecology of Hybrid Incompatibilities.
Cold Spring Harb Perspect Biol:a041440.

Tusso S, Nieuwenhuis BPS, Weissensteiner B, Immler S, Wolf JBW. 2021. Experimental evolution of
adaptive divergence under varying degrees of gene flow. *Nat Ecol Evol* 5:338–349.

Villa SM, Altuna JC, Ruff JS, Beach AB, Mulvey LI, Poole EJ, Campbell HE, Johnson KP, Shapiro MD, Bush
SE, et al. 2019. Rapid experimental evolution of reproductive isolation from a single natural
population. *Proceedings of the National Academy of Sciences* 116:13440–13445.

Westram AM, Stankowski S, Surendranadh P, Barton N. 2022. What is reproductive isolation? *Journal of
Evolutionary Biology* 35:1143–1164.

Zackrisson M, Hallin J, Ottosson L-G, Dahl P, Fernandez-Parada E, Ländström E, Fernandez-Ricaud L,
Kaferle P, Skyman A, Stenberg S, et al. 2016. Scan-o-matic: High-Resolution Microbial Phenomics
at a Massive Scale. *G3 Genes|Genomes|Genetics* 6:3003–3014.

Zhang Z, Bendixsen DP, Janzen T, Nolte AW, Greig D, Stelkens R. 2020. Recombining Your Way Out of
Trouble: The Genetic Architecture of Hybrid Fitness under Environmental Stress. *Molecular*
Biology and Evolution 37:167–182.
983

Data Accessibility and Benefit-Sharing Section

Phenotypic data are available in Table S1. The raw sequences were deposited in the European

Nucleotide Archive, parental populations are available at accession number PRJEB46680, while hybrids

are available at accession number PRJEB83568. The code used for analyses and plotting are available at

https://github.com/devinbendixsen/EvolvingRI. Benefits from this research accrue from the sharing of

our data and results on public databases as described above.

Author Contributions

Conceived research: Rike Stelkens; Designed research: Rike Stelkens, Ciaran Gilchrist; Performed

research: Devin P. Bendixsen, Ciaran Gilchrist, Chloé Haberkorn, Karl Persson; Contributed new reagents

or analytical tools: Karl Persson, Jonas Warringer; Data analysis: Devin P. Bendixsen, Chloé Haberkorn,

Karl Persson; Writing: Rike Stelkens, Devin P. Bendixsen, Chloé Haberkorn, Ciaran Gilchrist; Funding

acquisition: Rike Stelkens, Devin P. Bendixsen, Cecilia Geijer, Jonas Warringer.