Uniform bacterial genetic diversity along the gut

Michael Wasney[1]*, Leah Briscoe[2]*, Richard Wolff[3], Hans Ghezzi[4], Carolina Tropini[5,6,7],

Nandita Garud[1,2,3]

[1]University of California, Los Angeles, Human Genetics, Los Angeles, CA, [2]University of

California, Los Angeles, Interdepartmental Program in Bioinformatics, Los Angeles, CA,

[3]University of California, Los Angeles, Ecology and Evolutionary Biology, Los Angeles, CA, [4]

University of British Columbia, Department of Bioinformatics, Vancouver, Canada, [5]University

of British Columbia, Department of Microbiology and Immunology, Vancouver, Canada,

[6]University of British Columbia, School of Biomedical Engineering, Vancouver, Canada,

[7]Canadian Institute for Advanced Research, Humans and the Microbiome Program, Toronto,

Canada

*These authors contributed equally to this work.

Abstract

Environmental gradients throughout the digestive tract shape spatial variation in the composition

and abundance of bacterial species along the gut. However, much less is known about how

genetic diversity *within* bacterial species is distributed along the gut. Understanding this

distribution is important because bacterial genetic variants confer traits that influence both

microbiome function and host physiology, including local inflammation and nutrient

metabolism. Thus, to understand how the microbiome functions at a mechanistic level, it is

essential to understand how genetic diversity is organized along the gut. In this study, we

profiled genetic diversity of approximately 30 common gut commensal bacteria in five regions

along the gut lumen in germ-free mice colonized with the same healthy human stool sample.

Although species composition varied considerably along the gut, genetic diversity within species

was substantially more uniform. Driving this uniformity were similar strain frequencies along the

gut, implying that multiple, genetically divergent strains of the same species can coexist within a

host without spatially segregating. Additionally, the approximately 60 unique evolutionary

adaptations arising within mice tended to sweep throughout the gut, showing little gut region

specificity. We then analyzed metagenomic samples collected along the guts of conventional

mice and healthy humans and found similar dynamics with their natural microbiomes, suggesting

that uniform bacterial genetic diversity may be common to multiple host species. Together, our

findings demonstrate that uniform spatial distribution of genetic diversity along the

gastrointestinal tract is a robust feature of mammalian gut ecosystems.

Introduction

The spatial organization of microbiota along the gut plays a critical role in microbiome function[1].

Previous research in humans and mice has established that different regions of the gut, such as

the small and large intestines, harbor distinct assemblages of bacterial species[2,3], and that

disruption of this organization is associated with disease[4]. Spatial organization in the microbiome

is thought to arise from distinct niches imposed by environmental and physiological gradients

along the gut[3,5,6]. However, it remains unknown whether such gradients drive similar spatial

organization of within-species bacterial genetic diversity, such as by determining the occupancy

of genetically distinct strains or individual genetic variants along the gut. Understanding spatial

organization at this level is crucial, as bacterial genetic diversity drives traits that support proper

microbiome function and are associated with multiple host-relevant phenotypes[7], including

ability to digest food[8], local inflammation in specific regions of the gut[9], antibiotic resistance[10],

metabolic capabilities[8,11–14], and pathogen resistance[15].

There are several ecological and evolutionary mechanisms that could result in either i) spatially

segregated or ii) well-mixed genetic variation along the gut (**Figure 1A**). Spatial segregation

could arise because environmental gradients select for different strains and mutations along the

gut, co-colonizing strains occupy distinct physical niches to reduce competition, or slow

migration rates limiting the spread of genetic variation. Contrastingly, spatial uniformity could

arise because adaptive variants are globally adaptive along the gut, co-colonizing strains occupy

distinct non-physical (e.g., metabolic) niches, or rapid migration rates.

Previous studies have sought to examine the spatial distribution of strains and individual

nucleotide variants. One study looking at a small number of commensal bacterial species found

that different strains can coexist along the human gut[16], but did not determine whether strain

frequencies of these co-colonizing strains were variable in different gut regions or whether

coexistence was a general phenomenon. Other recent work found that single nucleotide variants

(SNVs) can arise locally along the gut for a few bacterial species[16,17], though these studies did

not demonstrate that spatially specific evolutionary changes were adaptive or common across

bacterial species or host. In another case, a uropathogenic strain of *Enterococcus gallinarum*

evolved adaptive mutations specific to the mucosa, which caused inflammation[9]. However, it is

unclear how commonly adaptive mutations spatially segregate along the longitudinal axis of the

gastrointestinal (GI) tract (i.e., from small to large intestine), particularly among commensal

microbes not facing any strong selective perturbations such as antibiotics. Here we will test the

hypothesis that strains and their adaptations may also display spatial segregation, similar to the

spatial heterogeneity observed at the species level.

To date, characterizing the spatial structure of genetic diversity along the mammalian gut has

been challenging because of the invasive nature of directly sampling the GI tract, the limited

bacterial biomass obtained from such attempts, and because of the difficulties in culturing a

broad diversity of taxa found natively in the human gut microbiome. However, these challenges

can be circumvented using a combination of experimental and statistical approaches. First, the

use of a gnotobiotic mouse model for studying the gut microbiome enables the study of human

gut-specific species of bacteria *in vivo* in a controlled manner as a single inoculum can be

administered to multiple mice, allowing us to measure variability across replicates[18–21]. Second,

the use of mice overcomes the hurdle of invasive sampling in humans as the full gut contents can

be extracted after sacrifice and then processed for sequencing. Third, the use of shotgun

metagenomic sequencing enables the characterization of genetic diversity across a broad panel of

bacterial taxa, including those undetectable using culture-based methods[22]. Finally, recent

statistical innovations in phasing genotypes of individual strains from shotgun metagenomic data

enables the ability to quantify strain frequencies and their evolutionary modifications[23–25].

In this study, we capitalize on these advances to assess the distribution of strains and

evolutionary modifications along the gut. Specifically, we analyzed the frequencies of bacterial

strains and the evolutionary modifications they carry within luminal samples obtained from

along the GI tracts of germ-free mice eight weeks after they were inoculated with the

microbiome of a single healthy human (**Figure 1B**). We previously analyzed this same

humanized mouse dataset with 16S rRNA sequencing of luminal contents and found significant

shifts in abundance of taxonomic family members along the gut[3]. By using metagenomic

sequencing to analyze the same samples at a deeper level, we were able to estimate strain

frequencies and identify evolutionary modifications. We report that the frequency of strains and

evolutionary modifications found in most bacterial species was unexpectedly uniform across the

gut. Confirming that this uniformity extends to other natural microbiomes, we applied the same

analyses to (i) a cohort of conventional mice with their own naturally acquired, host-adapted

microbiomes and (ii) a cohort of healthy humans whose luminal contents from multiple gut

regions were sampled using ingestible capsule devices swallowed over several days. In all

systems, we observed spatial uniformity of genetic diversity along the GI tract, suggesting that

this pattern may be a common feature of multiple mammalian gut microbiomes. These results

further imply that species abundance differences along the gut may play a greater role than

genetic differences in responding to environmental gradients along the gut, and that this could be

a robust feature of gut microbial ecology across both mouse and human hosts.

Figure 1. Hypotheses and experimental design. (A) Possible organization of genetic diversity

along the gut. Strains and their evolutionary changes may be either spatially segregated, or

evenly mixed. **(B)** Experimental design. Eight germ-free Swiss Webster mice were inoculated

with a single healthy human-derived stool sample[3]. Subsequently, the microbiota was allowed to

equilibrate over eight weeks before shotgun sequencing was performed on samples collected

from five regions along the gut (small intestine: duodenum, jejunum, and ileum; large intestine:

107   cecum and colon). The duodenum, jejunum, and ileum are considered regions of the small

108   intestine, and cecum and colon are considered regions of the large intestine. To quantify species

109   abundances and SNV frequencies, metagenomic reads were mapped against a reference database

110   of genomes from almost 6,000 bacterial species.

Results

To investigate the effect of gut region on microbial strain diversity, we performed shotgun

sequencing on microbiomes from eight mice that were orally gavaged germ-free Swiss Webster

mice with a single stool sample (hereafter referred to as the inoculum) from a healthy human

donor in a previous study[3] (**Figure 1B**). Luminal contents were collected from five intestinal

regions in each mouse: the duodenum, jejunum, ileum, cecum, and colon. DNA was extracted

and metagenomic sequencing was performed on each of the five mouse intestinal samples as

well as the original human inoculum (**Methods**).

To infer species abundances and SNV frequencies, we mapped shotgun reads to a database of

reference genomes[26] for the most prevalent and abundant bacterial species present in these mice

(**Figure 1B**) (see **Methods** for more details on the bioinformatic pipeline used). Summaries of

species and genetic-level diversity have been previously quantified from stool with similar

approaches (e.g., Nayfach *et al.*, 2016)[26], but here we investigate this diversity along the gut.

Bacterial species and functional diversity differ along the gut

First, we examined whether bacterial species diversity, measured using alpha diversity (the

Shannon index), varied along the tract of the gut. We hypothesized the large intestine would

have higher alpha diversity compared to other regions of the gut, as previous studies in humans[27]

and mice[19,20] have found that species diversity is highest in the colon due to lower flow rates and

the relatively high abundance of host- and food-derived complex polysaccharides[5,6,28]. As

expected, gut region was a major driver of alpha diversity, with the large intestine having

significantly higher diversity than all regions in the small intestine (average fold change per

mouse = 1.22; Wilcoxon signed-rank test, p-value = 0.0156) (**Figure 2A**).

Next, we hypothesized that the relative abundance of individual bacterial community members

differ along the gut, as factors such as oxygen availability, antimicrobial peptides, and carbon-

source favor different dominant taxa in the mammalian small and large intestines,

respectively[1,2,29,30]. To quantify shifts in community composition along the gut, we examined the

relative abundance of bacterial families, expecting to see saccharolytic, obligate anaerobic taxa

dominate in the large intestine and rapidly dividing facultative anaerobes dominate in the small

intestine[2,6,26,28,29]. Supporting our prediction, several families in the order Bacteroidales—an

order of anaerobic bacteria comprising many species which are involved in breaking down

complex sugars in the colon[31]—were enriched in the large intestine (**Figure 2B, Supplementary**

Figure 1), including the families Bacteroidaceae (median log2 fold change per mouse = 0.338;

Wilcoxon signed-rank test, p-value = 0.0143), Porphyromonadaceae (median log2 fold change

per mouse = 8.58×10^{-2}; Wilcoxon signed-rank test, p-value = 0.0143), and Rikenellaceae

(median log2 fold change per mouse = 6.23×10^{-3}; Wilcoxon signed-rank test, p-value =

0.0209). Meanwhile, the family Lactobacillaceae, which is known to colonize the human small

intestine with the aid of unique mucus-binding proteins (MUBs) which adhere to the small

intestinal mucosal layer[29], was enriched in the small intestine relative to the large intestine

(median log2 fold change per mouse = -0.121; Wilcoxon signed-rank test, p-value = 0.0225)

(**Supplementary Figure 1**). These enrichments in the small intestine confirm that despite human

stool more closely resembling the community composition of the human large intestine[30],

inoculation with a human fecal sample is sufficient to support colonization by small intestine-

associated taxa. Together, these results are in line with previous work showing that the

composition of microbial taxa differs along the gut[3,28].

Lastly, to investigate community-wide functional diversity along the gut, using HUMAnN3[32] we

mapped metagenomic reads to gene families and MetaCyc pathways. Across all samples,

HUMAnN3 identified 808,204 gene families corresponding to 470 MetaCyc pathways, which

were subsequently aggregated at the community level by summing read counts across taxa.

These pathway profiles were then analyzed using MaAsLin3[33] to test for differences between the

small intestine and large intestine. The model incorporated gut location, cage, and read count as

fixed effects and used subject-level stratification for paired comparisons—revealing 80 pathways

significantly associated with gut location (q < 0.05). Among these, 33 pathways showed highly

significant associations (q < 0.001), and their coefficients are shown in **Supplementary Figure 2**

(Supplementary Table 2). The same families we found to be enriched in the large intestine were

also the source of many of the pathways most significantly associated with the large intestine,

namely Bacteroidaceae (25 pathways) and Rikenellaceae (14 pathways). Meanwhile, a family

enriched in the small intestine, Lactobacillaceae, was the source of 7 pathways significantly

associated with the small intestine.

Figure 2. Increase in taxonomic diversity and change in community membership along the length of the gut. (A) Alpha diversity (Shannon Index) estimates for bacteria in different regions of the gut. Two-sided Wilcoxon rank sum tests were performed between all possible pairs of small intestinal versus large intestinal samples (* indicates p ≤ 0.05; ** indicates p ≤ 0.01). **(B)** Relative abundance of bacterial families in the five gut regions and inoculum. Bacterial families are grouped by order in the legend.

Bacterial genetic diversity along the gut

Having confirmed that taxonomic diversity varies spatially along the gut, we next investigated whether the same was true of within-species genetic diversity. First, we quantified the distribution of nucleotide diversity (π) for the 30 most abundant species detected in the inoculum and mice (**Methods**) (**Supplementary Figure 3**, **Supplementary Table 3**). Across these species, nucleotide diversity measured in the inoculum was highly variable, ranging from 2.2×10^{-4} in *Odoribacter splanchnicus* to 1.2×10^{-2} in *Bacteroides vulgatus,* spanning two orders of magnitude (**Figure 3A**). High values of π ($> 10^{-3}$) are generally inconsistent with the colonization and subsequent diversification of a single strain during a host's lifetime, and instead support the presence of two or more strains that diverged long before colonization[23]. Meanwhile, low values of π are more consistent with colonization by a single strain[23]. Thus, the large range of π values observed across species indicates that in some instances, there are multiple co-colonizing strains of the same species, and in other instances there is only one strain.

Average nucleotide diversity values in the mice were positively correlated with nucleotide diversity in the inoculum (Spearman $\rho = 0.845$, p-value $= 7.55 \times 10^{-7}$) (**Figure 3A, Supplementary Figure 4**). However, despite this overall similarity, a range of nucleotide diversity values was observed within individual mice, with some mice having levels comparable to those seen in the inoculum and others having substantially lower values, suggesting variation in lineage structure across mice. Specifically, for the species with the highest values of nucleotide diversity in the inoculum ($\pi > 1 \times 10^{-3}$), values across mice were often bimodally distributed. For example, π values for *Bacteroides vulgatus* were either all clustered around 10^{-2} in some mice or around 10^{-3} and lower for other mice (**Figure 3A**). This variation suggests that in some mice, multiple strains colonized, and in other mice, a single strain colonized even when

multiple were present in the inoculum. However, π never spanned an order of magnitude within

a single mouse, potentially implying more similar strain colonization dynamics along the gut

within mice than across mice.

To understand if nucleotide diversity differed between regions of the gut, we measured F_{ST}, or

fixation index, between the small and large intestine for each species. F_{ST} is a distance metric

where high values indicate large genetic differentiation between two samples[39,40]. We

hypothesized that F_{ST} would be elevated between gut regions within hosts, either due to

evolutionary changes accruing or variable strain frequencies. Contrary to this expectation, we

observed that F_{ST} between the small and large intestine of the same host was on average low for

the 30 species examined (0.119), especially relative to the average F_{ST} between mice at the same

gut region (0.225) (**Figure 3B**). This difference was more pronounced in species with high

nucleotide diversity in the inoculum ($\pi > 10^{-3}$) (Wilcoxon rank sum test, p-value = 2.3×10^{-7})

compared to species with low diversity ($\pi \leq 10^{-3}$) (Wilcoxon rank sum test, p-value=0.043).

Species with high nucleotide diversity in the inoculum may have larger differences in F_{ST} within

versus between mice than species with low nucleotide diversity in the inoculum because the

former exhibits more variable strain colonization dynamics. Specifically, when multiple strains

are present in the inoculum, there is a potential for different combinations of strains to colonize

different mice, leading to higher F_{ST} and differences in π between mice. Within mice, F_{ST} tends

to be low and π tends to be uniform for these species, suggesting that similar combinations of

strains are found all along the gut tract of the same host. Meanwhile, when there is only one

strain present in the inoculum, the same strain should colonize all mice and gut regions. This is

consistent with the observed low values of F_{ST} and more uniform values of π both within and

between mice for these latter species. In the next section, we further investigate whether similar

combinations of strains are found along the gut tract when nucleotide diversity is high.

Figure 3: Genetic diversity measured in 30 gut commensal species. (A) Nucleotide diversity
(π) estimates for the 30 most abundant and prevalent species (**Methods**). Each dot represents
nucleotide diversity (π) in a specific gut region of an individual mouse. Dot color corresponds to
the mouse, while dot shape indicates the gut region. Red asterisks denote nucleotide diversity
measured in the inoculum for that species. Species names in light red correspond to those species
with $\pi > 1 \times 10^{-3}$ in the inoculum. **(B)** Fixation index (F_{ST}) for between all pairs of mouse

samples for the 30 species analyzed in panel **(A)**. Each data point represents a single species for a

given pair of samples, with points colored according to whether the sample comparison is within

host (grey) or between hosts (black). Data points are further separated based on nucleotide

diversity of the species in the inoculum. Two separate Wilcoxon rank sum tests were performed

comparing the distributions of F_{ST} within versus between hosts for species with $\pi > 10^{-3}$ and

species with $\pi \leq 10^{-3}$, respectively (* indicates p $\leq$ 0.05; **** indicates p $\leq$ 0.0001).

Strain frequencies are uniform along the gut within hosts

We next asked whether the identities and frequencies of strains colonizing different parts of the

gut of a given mouse are the same or different when multiple strains are present in the inoculum.

Variation in occupancy or frequency along the gut would suggest that physical niche partitioning

is important in maintaining multiple conspecific strains in the same host that might otherwise

compete in the same physical niche, or that distinct environmental pressures along the gut select

for different strains.

To understand whether frequencies of co-colonizing strains of the same species vary along the

gut, we inferred strain frequencies for the 14 species with levels of nucleotide diversity $> 10^{-3}$

in the inoculum, as these samples had sufficiently high enough diversity to potentially harbor at

least two strains (**Figure 3**). To infer strain frequencies, we used an algorithm previously applied

to metagenomic time series data[24,25] (**Methods**). For the species *Alistipes shahii*, *Bacteroides*

ovatus, *Coprococcus comes*, and *Faecalibacterium prausnitzii*, only one strain was inferred to be

present within the mouse samples. For the rest of the species, we inferred that two strains were

present (**Figure 3**; **Supplementary Figure 5**). We next asked whether the frequencies of strains

for species with two strains were uniform or variable along the gut.

Application of the strain phasing algorithm revealed that frequencies of co-colonizing strains

were relatively uniform across gut regions within hosts for most bacterial species and mice,

contrary to the expectation that their frequencies may differ due to niche partitioning or

environmental selection. For example, for the species *Blautia wexlerae*, the two strains were able

to coexist at roughly 40:60% frequency along all five regions of the gut in mouse 8, and for the

species *Adlercreutzia equilofaciens*, the two strains were able to coexist at roughly 20:80%

frequency in all parts of the gut in mice 4 and 5 (**Figure 4A**). To understand whether co-

colonizing strains generally displayed uniform frequencies along the gut, we calculated change

in major strain frequency (Δf) between mouse samples for 7 species that were present in

multiple mice, cages, and gut regions (**methods**). Across the seven species, median change in

frequency between gut regions within a host was only 9.36×10^{-3}, much smaller than the

median change in frequency we observed across mice within the same cage (3.49×10^{-2}) or

between mice in different cage (3.09×10^{-1}) (**Figure 4B**).

To confirm that strain frequencies were generally more homogeneous within versus between

hosts for all species in our dataset, we performed a series of ANOVA tests to estimate the

proportion of variance in major strain frequencies attributable to gut region ("gut region"),

mouse identity ("mouse"), and cage identity ("cage") (**Supplementary Figure 6**). Consistent

with the observations of uniformity in genetic variation along the gut, gut region explained the

lowest variance for all seven bacterial species (average variance explained = 2.70%), suggesting

that gut region is less important in explaining strain frequencies than mouse identity or cage

identity. By contrast, mouse and cage explained substantially greater variation (average

combined variance explained = 90.0%). These observations support the idea that strain

frequencies along the gut were generally uniform across species and reveal that multiple strains

were able to coexist in the same region of the gut. Furthermore, they imply some ecological

mechanism other than spatial segregation must be responsible for permitting strain coexistence

within the same host.

Social transmission of bacteria, through behaviors like coprophagy[14,41], may also facilitate

between-host spread of strains, resulting in similar strain frequencies between mice in the same

cage. Supporting this notion, strain frequencies were more similar among cohoused mice

compared to mice housed in different cages, indicating that migration of strains between mice

plays a role in homogenizing frequencies of strains. For example, for the species *A.*

equilofaciens, which was present in multiple mice and cages, the within-cage variance of strain

frequencies measured in the jejunum ranged from 1.65×10^{-6} in cage 3 to 1.29×10^{-3} in

cage 2, whereas the between-cage variance was at least two orders of magnitude greater,

1.18×10^{-1} (**Supplementary Figure 7, Supplementary Table 4**). These results align with

previous studies that have found coprophagic behavior in mice can facilitate the spread of strains

and adaptive variants in *Bacteroides thetaiotaomicron*[42] and *Escherichia coli*[14,41] between co-

housed mice, enforcing similar strain frequencies between mice[14]. However, not all species have

similar within-cage strain frequencies (e.g., *B. uniformis*, **Figure 4**), suggesting barriers to

between-host migration in these cases. This species-specific migration effect is reflected in the

fact that "cage" explains most of the variance in strain frequency for four species, but very little

variance in the remaining three species (**Supplementary Figure 6**).

Together, these results indicate that co-colonizing strains do not need to occupy physically

distinct niches to colonize the same host. To the contrary, strain frequencies tended to be

relatively uniform along the gut. This uniformity may be driven by rapid migration rates between

gut regions, which also likely results in greater similarity in strain frequencies among mice of the

same cage versus different cages due to social transmission.

Figure 4. Strain frequencies are similar along the gut but may differ between mice housed in different cages. (A) Strain frequencies for four bacterial species detected in the guts of eight mice inoculated with the same human stool sample. A total of 14 bacterial species had high levels of inoculum nucleotide diversity ($\pi \geq 1 \times 10^{-3}$). See **Supplementary Figure 5** for the strain frequencies of the ten species not shown here). "Strain 1" and "strain 2" represent the same strain across all samples for a given species. Error bars represent bootstrapped 95% confidence intervals. Mice 1-3 were co-housed in cage 1, mice 4 and 5 in cage 2, and mice 6-8 in cage 3, with the housing scheme indicated by the alternating light and dark purple boxes. Absence of a strain frequency bar in a sample indicates insufficient coverage or species absence (**Methods**). **(B)** Change in strain frequency Δf was calculated for seven species present in multiple mice, cages, and gut regions. (**Methods**). Each dot represents the change in major strain frequency between a pair of samples from within the same host (red), between hosts in the same cage (light purple), and between hosts in the different cages (dark purple).

Evolutionary changes are not localized and spread throughout the gut

Even if bacterial strain frequencies are relatively homogenous along the gut, it is possible that individual genetic variants within species are locally restricted along the gut, potentially due to local adaptation. To assess the possibility of local adaptation, we asked if any SNVs have extreme frequency differences between gut regions, or if they too are homogeneously distributed.

SNV frequency changes between gut regions could arise within a host as a consequence of (i) evolutionary changes (e.g., adaptation), (ii) fluctuations in frequencies of genetically distinct strains, or (iii) sampling error. Here, we were interested in distinguishing SNV frequency changes arising as a consequence of evolution from those arising from strain fluctuations or sampling error. To distinguish SNV frequency changes arising due to evolution as opposed to strain fluctuations, we inferred the genotype of one of the strains present in the sample, or 'quasi-phased' the genotype[23] and then tracked SNV frequency changes on the background of that genotype between pairs of samples. Doing so allowed us to detect new mutations modifying a resident lineage as opposed to alleles rising to high frequency due to strain fluctuations. To distinguish SNV frequency changes due to evolution (e.g., adaptation) versus sampling error, we considered only large allele frequency changes ($f \leq 0.2$ in one sample to $f \geq 0.8$ in another

sample) between samples with a median read coverage of 20 or more and at sites with a

minimum read coverage of 20 or more in both samples, which are unlikely to arise due to

sampling error or drift[23,43]. Previously, we showed that the genotype of the dominant lineage in

samples with a single, high-frequency strain can be confidently quasi-phased[23], and that

evolutionary changes with large allele frequency changes can subsequently be detected

(**Methods**).

Using this method, we identified 366 quasi-phaseable genomes from 62 bacterial species present

in the intestinal contents of the eight mice and the inoculum (**Supplementary Figure 3**). Next,

we detected SNV frequency changes between pairs of quasi-phaseable genomes. These pairs

were between the inoculum and any mouse (10 bacterial species), different gut regions *within*

mice (spanning 33 bacterial species), and gut regions *between* mice (28 bacterial species). To

avoid comparing distantly diverged lineages that were not descended from the same inoculum

strain, we only considered pairs with ≤20 SNV frequency changes. We observed a total of 65

unique SNV frequency changes across all possible pairs of quasi-phaseable genomes in our

dataset (**Figure 5A**, **Supplementary Figure 3, Supplementary Figure 8**, **Supplementary**

Table 5).

By having an inoculum to compare to, we can ascertain if a lack of SNV differences along the

gut is due to the no evolutionary changes accruing, or the global spread of an evolutionary

change throughout the gut. We observed 11 SNV frequency changes rising from low frequency

in the inoculum to high frequency in mice in the 10 species with at least one inoculum-mouse

sample quasi-phaseable genome pair (**Figure 5A**, **Supplementary Figure 8**, **Supplementary**

Table 5). The rapid timescales over which these SNVs rose to high frequency are consistent with

these SNV frequency changes potentially representing adaptations occurring during the host

colonization process. Only one of these SNVs, detected in *Coprococcus* sp., showed spatial

differentiation in a single host (SNV frequency of 0 in jejunum versus 0.935 in cecum of mouse

1) (**Figure 5A**, **Supplementary Table 5**), indicating that advantageous SNVs tend to spread

along the gut. Since not all SNVs observed within mice had sufficient coverage to be analyzed in

the inoculum, we next examined SNV frequency changes arising between gut regions.

Supporting the observation that SNVs spread along the gut, we observed only 4 unique *within-*

host SNV frequency changes in 3 out of the 33 species, all between the small intestine and large

intestine, including the SNV in *Coprococcus* sp. (**Supplementary Table 5**). By contrast, we

observed 57 unique *between-host* SNV frequency changes in 15 of the 28 species (**Figure 5A**,

Supplementary Figure 8, **Supplementary Table 5**). The rate of SNV frequency changes

observed between hosts was significantly higher than that within hosts (bootstrapped 95%

confidence interval for between-host SNV changes is 1.54×10^{-7} differences/bp -

4.42×10^{-7} differences/bp; bootstrapped 95% confidence interval for within-host SNV changes

is 0 differences/bp - 6.62×10^{-8} differences/bp) (**Figure 5B**). The low rate of SNV frequency

changes within mice relative to between-host and inoculum-mouse SNV frequency changes rates

suggests that potentially advantageous SNVs overwhelmingly transmit along the gut within

hosts.

To understand whether SNVs spreading along the gut showed additional signals of being

adaptive beyond extreme allele frequency changes on short time scales, we asked if the same

SNV changes arise in multiple mice, as parallelism is a hallmark of adaptation[44–48]. Adaptive

variation can arise in multiple mice via independent recurrent *de novo* mutation or the parallel

rise in frequency of standing genetic variation (SGV) already present in the inoculum[48]. Of the

21 unique SNVs that were observed to undergo an extreme allele frequency change between any

pair of samples and had sufficient coverage in the inoculum (minimum of 20 reads), 15 had non-

zero frequency in the inoculum, indicating that SGV is an important source of genetic variation

in these mice (**Figure 5A**, **Supplementary Figure 8**, **Supplementary Table 5**). Of these, five

nonsynonymous and one synonymous SNVs were observed to undergo parallel sweeps from low

frequency in the inoculum to high frequency in multiple independent cages. One of these parallel

SNVs was in *P. distasonis* in a gene encoding a TonB-dependent transporter, which rose from

low frequency in the inoculum ($f = 0.057$) to high frequency in multiple mice in all three

cages (ranging from an average $f = 0.319$ in mouse 4 to an average $f = 1.00$ in mouse 6),

consistent with parallelism (**Figure 5A**, **Supplementary Table 5**). TonB-dependent transporters

are ubiquitous outer membrane-associated proteins involved in nutrient transport[49] with

structural variation having been found to be associated with niche partitioning by different

Bacteroides taxa[50]. Moreover, TonB-dependent transporters have been found to be a common

target of adaptation in the human gut microbiome[47,51,52]. Notably, all the SNVs observed to

undergo parallelism were not found to be differentiated along the gut within hosts. This fact

would seem to suggest that even when SNVs are adaptive, they are either broadly adaptive to the

entire gut, or alternatively have high migration rates that outweighs local selection pressures,

perhaps due to mixing and peristalsis along the gut[53].

Figure 5: Evolutionary changes are uniformly found along the gut. (A) SNVs exhibiting
extreme allele frequency changes (allele frequency $f \leq 0.2$ in one sample to $f \geq 0.8$ in another
sample) between any pair of samples sharing the same quasi-phaseable strain genome (< 20 SNV
frequency changes) are shown for *Coprococcus* sp., *P. distasonis*, *R. intestinalis*, and
Ruminococcus sp. Asterisks represent the frequency of a given SNV, with each SNV within a
species having its own unique color. Samples lack asterisks for a particular SNV when those loci
do not meet the minimum coverage requirement of 20 reads to infer frequency. When two strains
of the same species are present, strain abundances of co-colonizing strains are represented by
two different shades of brown (see **Figure 4** for legend). The remaining 11 species in which
SNV frequency differences were detected are shown in **Supplementary Figure 8**. **(B)**
Comparison of the rate of SNV frequency changes aggregated across species within mice,
between mice, and between inoculum and mice. Colored in red are the observed SNV frequency
change rates aggregated across species. Bars represent bootstrapped 95% Cis (**Methods**).

Genetic homogeneity is generalizable to the gut of conventional mice and healthy humans

Having established that uniform strain frequencies and spread of evolutionary adaptations along
the gut is a common feature in humanized mice, we next investigated whether these same
patterns also arise in hosts with established native microbiota, specifically conventional mice and
healthy humans.

We first analyzed a cohort of nine conventional mice that were cohoused until weaning at three
weeks of age before being housed into three cages of three mice each. At six weeks of age, we
sampled and sequenced luminal contents from the same five gut regions that were previously
examined in humanized mouse guts (**Methods**). Large intestinal samples harbored higher alpha
diversity and a different family-level composition than small intestinal samples (**Methods**)
(**Supplementary Figure 9**), confirming that the expected spatial organization does exist at
higher taxonomic levels along the guts of the conventional mice.

Further replicating our findings from humanized mice, we found that in conventional mice, co-
colonizing strains of the same species exhibited largely uniform frequencies across gut regions.
This pattern contrasts with the substantial variation in strain frequencies between mice, even if
co-housed (**Figure 6A**, **Supplementary Figure 10**). Among the nine bacterial species with co-

colonizing strains that were present in multiple mice, cages, and gut regions, median change in frequency Δf between gut segments within hosts was only 2.21×10^{-2}, an order of magnitude smaller than the median frequency changes observed between mice in the same cage (2.16×10^{-1}) and in different cages (1.42×10^{-1}) (**Figure 6B**). Corroborating these results, gut region explained the least variation (average variance explained = 0.75%), whereas mouse identity explained the most variation (average variance explained = 78.39%) (**Supplementary Figure 11**). Cage explained a moderate amount of variation (average variance explained = 19.97%). These results indicate that uniform frequencies of strains along the gut is not unique to humanized mice and may be a common feature of mouse microbiota.

Additionally, we did not observe evolutionary changes accruing between gut regions within hosts even while changes arose between hosts at a rate of 1.03×10^{-6} changes/bp (**Figure 6C**). These results indicate that any evolutionary changes arising within hosts sweep globally throughout the gut.

Next, to determine whether spatially uniform patterns of genetic diversity are a feature of human guts as well, we analyzed previously published metagenomic data[54] collected using ingestible sampling capsules designed to sample luminal contents from four successive regions of the gastrointestinal tract: pyloric sphincter to the upper small intestine, upper to mid-small intestine, mid- to lower small intestine and the lower small intestine into the ascending colon. In this dataset, 15 healthy adult subjects ingested four capsule types twice daily for as long as a week.

We hypothesized that strain frequencies in humans might vary more along the gut than in mice, given the greater length and compartmentalization of the human intestine, as well as the potential for host-specific niche adaptation over long residence times[25,27]. Contrary to this expectation, strain frequencies were generally homogeneous not only along the gut but also across timepoints. For example, the frequency of the major strain of *Ruminococcus obeum* in Subject 11 fluctuated around a mean of 72% with a variance of 0.4% across gut locations and timepoints (**Figure 6D**). More generally, across the 25 species x host pairs with co-colonizing strains in multiple timepoints and device types, the median change in major strain frequency Δf between gut regions within the same timepoint was only 3.79×10^{-2}. This was the same order of magnitude as the median change observed between consecutive timepoints (2.92×10^{-2}), or even between

the first and last observed timepoint for a species x host pair (3.55×10^{-2}) (**Figure 6E**). These

results indicate that strain frequency tends to be uniform along the human gut as well as across

hourly and daily timescales.

While the majority of species x host pairs never showed a temporal or spatial strain fluctuation in

strain frequency exceeding 25%, there were several outlier species x host that harbored strains

undergoing large frequency temporal shifts, and to a lesser degree, spatial shifts. However, these

fluctuations tended to be transient. For example, the frequency of the major strain of

Burkholderiales bacterium in Subject 6 fluctuated between a minimum of ~10% and maximum

of ~95% within and across timepoints, but displayed no consistency in the direction of change

across either space or time (**Supplementary Figure 12**). We observed this in other species x

host pairs as well, illustrating that strain fluctuations that do occur along the gut and over time

are short-lived, consistent with previous work showing that stably co-colonizing strains can

fluctuate in frequency around a carrying capacity over longer timescales[25].

We next asked whether extreme SNV frequency changes arising as a consequence of within-host

evolution were spatially segregated. To do so, we identified SNV frequency changes between

distinct regions of the gut for each of the 56 host x species pairs (representing 35 species

detected in 11 subjects) with enough data in at least two capsules for a given time point. Out of

the 56 pairs, only 7 of these (12.5% of species x host pairs) had detectable SNV frequency

changes between gut regions within a single timepoint (**Supplementary Table 6**), indicating that

spatial segregation of SNVs is a feature in only a minority of samples.

Finally, we investigated whether the SNV changes arising along the gut are observable over

multiple time points, or whether they are transient. Persistence would suggest that environmental

gradients maintain spatial segregation of genetic variation along the gut over time. Of the seven

host x species pairs with spatial differences, four had sufficient temporal and spatial data (at least

two timepoints with at least two device types per timepoint). In three of four host x species pairs,

spatial differences did not persist over multiple time points, with allele frequencies often

reverting to their original state (**Supplementary Figure 13**, **Supplementary Table 6**). These

results imply that sustained extreme allele frequency changes along the gut are rare.

Figure 6. Uniform genetic variation along the gut in conventional mice and healthy humans. (A) Strain frequencies for two co-colonizing strains of an *Oscillospiriceae* species long the guts of nine conventional mice housed across three cages. Error bars represent bootstrapped 95% confidence intervals. Mice 1-3 were co-housed in cage 1, mice 4-6 in cage 2, and mice 7-9 in cage 3, as indicated by the light and dark purple boxes. Absence of a strain frequency bar indicates insufficient coverage or species absence in a given sample. For additional strain frequency plots corresponding to species analyzed in **(B)**, see **Supplementary Figure 10**. **(B)** Change in strain frequency Δf was calculated for seven species present in multiple mice, cages, and gut regions. (**Methods**). Each dot represents the change in major strain frequency between a pair of samples from within the same host (red), between hosts in the same cage (light purple), and between hosts in the different cages (dark purple). **(C)** Comparison of the rate of SNV frequency changes within and between mice. Colored in red are the observed SNV frequency change rates aggregated across species. Bars represent bootstrapped 95% CIs. **(D)** Strain frequencies for two co-colonizing strains of *Ruminococcus obeum* were inferred from capsule devices that sampled luminal contents along the gut of a healthy human subject (subject 11) across six timepoints. Device type 1 targeted the pyloric sphincter to the upper small intestine, device type 2 targeted the upper to mid-small intestine, device type 3 targeted the mid- to lower small intestine, and device type 4 targeted the lower small intestine into the ascending colon[54]. Error bars represent bootstrapped 95% confidence intervals. Timepoints represent the time at which capsules were swallowed, with days being relative to the first timepoint plotted and times of day being coarsened into "morning" (before 12 pm PST), "afternoon" (after 12pm and before 8 pm PST), and "evening" (after 8 pm PST) bins. For additional strain frequency plots corresponding to species analyzed in **(E)**, see **Supplementary Figure 12**. **(E)** Change in strain frequency Δf was calculated between samples belonging to 25 species x host pairs with co-colonizing strains present in multiple timepoints and device types (**Methods**). Subject is indicated by the parenthetical. Each dot represents the change in major strain frequency between samples from different device types swallowed during the same timepoint, (red), collected during consecutive timepoints (light purple), and collected during the first and last timepoint (dark purple). Strain frequency change over the course of the entire sampling period rarely exceeded changes occurring between consecutive timepoints, suggesting that hourly or daily shifts in strain frequency are transient.

Discussion

In this study, we analyzed the frequencies of bacterial strains and evolutionary changes along the guts of eight mice inoculated with the same human-stool-derived inoculum. In sharp contrast to the species-level heterogeneity reported here and in previous studies[3,6,28,29,55–63], we found substantial *genetic* homogeneity along the gut (**Figure 4, Figure 5**, **Supplementary Figure 5**, **Supplementary Figure 8)**. Spatial uniformity was a feature of not only the bacterial species analyzed in humanized mice, but also those in conventional mice and healthy humans, indicating that uniformity is a robust feature of mammalian gut ecosystems. Together, these results indicate that spatial segregation of strains is not a requisite for their coexistence and that environmental gradients do not necessarily generate large strain fluctuations or allele frequency differences along the gut lumen.

Uniformity likely arises due to rapid migration rates of bacteria between gut regions and, in the case of the mice studies, due to rapid migration between mice (**Figure 4-6**, **Supplementary Figure 5-7**, **Supplementary Figure 11**). This latter conclusion is supported by the observation that strain frequencies are more similar between co-housed mice than between mice housed in different cages, consistent with recent work demonstrating extensive social transmission of bacteria among co-housed mice[14,41]. Additionally, this conclusion is supported by the observation that evolutionary changes at the SNV level are usually found at similar frequencies along the gut, suggesting migration of variants arising from a single origin as opposed to the same evolutionary change arising in different locations along the gut independently of each other. The possibility of rapid migration facilitating uniformity is consistent with recent work showing that there is significant mixing of luminal contents along the gut[53].

Uniform strain frequencies are not only found along the length of the gut but also over time, reflecting stable coexistence of strains. Stable coexistence is a common feature of healthy human gut microbiomes[25,27] and in experimental evolution studies[64,65]. Our own analysis of the healthy human cohort sampled over multiple time points show that while strains tended to fluctuate in frequency more across time than they did across space (i.e., between capsule) within any one timepoint, these temporal strain frequency fluctuations were short-lived and tended to revert to the overall trend at subsequent sampling points (**Figure 6E**, **Supplementary Figure 12**). Stable

coexistence of conspecific strains is thought to be facilitated at least in part by metabolic niche

partitioning[66]. Our findings support the role of metabolic niche partitioning in stably maintaining

strain diversity, as spatial segregation does not seem required to reduce competition among

strains.

Future work employing higher-resolution sampling strategy across both time and space could

reveal bacterial genetic heterogeneity that was not observed in this study. For example, sampling

the gut prior to eight weeks post-inoculation might have captured distinct spatial distribution of

alleles before they migrated to all gut regions. Additionally, in this study we focused on extreme

allele frequency changes, as these are unlikely to be confounded with sampling error in shotgun

data and instead are likely to have arisen only due to adaptation. However, more subtle allele

frequency differences may manifest along the gut, especially among neutral genetic variants,

consistent with recent human microbiome studies analyzing metagenome-assembled genomes

reporting locally restricted, potentially neutral evolutionary changes in a limited number of

species[16,17].

In addition, while we focused on luminal contents from five gut regions and observed uniform

genetic variation, other physiological structures found in the mammalian gut could exhibit

greater spatial heterogeneity. For example, different crypts in the intestinal epithelium could

potentially be occupied by distinct strains, similar to that of skin pores[67]. Crypts are known to

have communities distinct from the lumen[55,68], and may perform ecological filtering at the strain

and nucleotide-level as well. Similarly, the mucosa displays different biogeography from the

lumen[56], and mucosal surfaces are known to generate population structure in culture[69], promote

site-specific binding in certain bacterial species that have co-evolved with their hosts[70], and drive

adaptive differentiation relative to populations in the gut lumen[9]. However, the technical

challenges in obtaining sufficient bacterial biomass for metagenomics sequencing from crypts or

mucosa is significant[71]. The continued improvement in host-depletion and fine-scale sampling

strategies (e.g., Wu-Woods *et al.*, 2023)[17] will enable investigation of bacterial population

structure at unprecedented spatial resolution.

Our findings in healthy mammalian hosts raise the question of whether spatial genetic structure

may emerge under disease conditions or altered environments. Local inflammation or

environmental changes could generate distinct niches that foster local bacterial adaptation or

filter for specific, disease-adapted strains. In future work, it will be interesting to investigate if

spatial structure at the genetic level arises in diseased hosts. Intestinal disorders such as

inflammatory bowel disease (IBD) are known to generate local inflammation phenotypes and

disrupt gut microbiome composition[72,73] in specific gut regions[4,74,75]. For example, in a recent

mouse study, adaptive SNVs in a species of pathogenic bacteria were found to appear locally in

different colonic tissues, enabling translocation of a pathogenic bacteria from the colonic lumen

to the mucosa, and subsequently to the liver[9]. Thus, in pathogenic bacteria and disease scenarios,

we might observe specific strains or local adaptations associated with these acute conditions

along the gut, supporting our original hypothesis that environmental heterogeneity might induce

genetic diversity along the gut.

In conclusion, our findings demonstrate that bacterial strains and their evolutionary changes are

uniformly distributed along the gut lumen of across different mammalian hosts, in contrast to the

spatial heterogeneity observed at higher bacterial taxonomic levels. These findings indicate that

spatial segregation of strains is not a requisite for their co-existence, and that environmental

gradients do not result in local adaptation among alleles experiencing the largest allele frequency

changes. Future studies aimed at understanding the dynamics of migration facilitating this

uniformity and the metabolic strategies employed by strains to co-exist will be crucial for

elucidating the dynamics of strain-level diversity and its functional consequences within the gut

ecosystem.

Methods

Humanized mouse experimental design

For the humanized mouse study, we performed shotgun sequencing on the luminal contents of

mice we previously analyzed with 16S rRNA sequencing[3]. All animal experiments were

conducted in accordance with the ethical guidelines of the University of British Columbia's

(UBC) animal care procedures, following protocol numbers A19-0078 and A19-0122 approved

by the Animal Care Committee. Swiss Webster male and female mice at nine weeks of age were

used and were provided with an autoclaved standard diet (Purina LabDiet 5K67). Briefly, eight

germk-free Swiss Webster mice were orally gavaged with the same human stool sample (the

inoculum) from a healthy adult human previously labeled TL1[3]. This set up is akin to previous

studies conducting controlled mouse studies with human inoculums[18–21]. The eight mice were

housed in three different cages in the following configurations: cage 1 had mice 1-3 (sex male),

cage 2 had mice 4 and 5 (sex male), and cage 3 had mice 6-8 (sex female). Given that human

microbiota can rapidly colonize the mouse in a matter of days to weeks, we allowed for a total of

eight weeks to colonize and equilibrate in the mice[18–21], allowing for stabilization of community

structure and recovery following oral gavage. This period included six weeks on a standard

rodent diet (LabDiet 5k67), followed by two weeks in which mice in cages 2 and 3 were exposed

to a fiber-rich diet of 30% guar gum (TestDiet 5BSE) while those in cage 1 remained on the

same chow. With this experimental set up, we were able to ascertain replicability and

generalizability of the genetic homogeneity observed along the gut in three independent cages

with varying diets and sexes, as well as contrast within-host, across-host, and across-cage

dynamics (**Figure 1**).

Eight weeks after inoculation, mice were sacrificed using carbon dioxide with secondary cervical

dislocation, and luminal contents were collected from five intestinal regions: the duodenum,

jejunum, ileum, cecum, and colon. The intestines were carefully sectioned using a razor blade

into the corresponding regions and the contents were squeezed into 1.5 mL-microcentrifuge

tubes. DNA extraction was performed on luminal content samples and the original inoculum

(TL1) from 96-well plates using the DNeasy PowerSoil Pro Kit (Catalog number: 47016). For

more information on the approach used, see the methods section of Ng *et al.*, 2023[3].

Conventional mice experimental design

Conventional mice microbiomes were obtained from nine male Swiss Webster mice from the

same litter and with their own microbiomes at six weeks of age. All animal experiments were

conducted in accordance with the ethical guidelines of the University of British Columbia's

(UBC) animal care procedures, following protocol numbers A19-0078 and A19-0122 approved

by the Animal Care Committee. Mice were cohoused until three weeks of age, at which point

they were separated into three cages (three mice per cage). They were provided with an

autoclaved standard diet (Purina LabDiet 5K67) for the duration of the six weeks. At six weeks,

mice were sacrificed and luminal contents were collected from the same five intestinal regions as

with the humanized mice (duodenum, jejunum, ileum, cecum, and colon), and DNA extraction

was performed using an identical approach.

Metagenomic Sequencing

Metagenomic sequencing was performed on luminal content samples from humanized and

conventional mouse cohorts and the inoculum using a NovaSeq with 150 bp paired end reads.

Humanized mouse samples were sequenced to a depth ranging from 23M to 136M reads (median

= 33M reads), and the inoculum was sequenced to a depth of 119M reads. Samples from mice 7,

8 and the inoculum were sequenced in a separate batch than from samples from mice 1-6 at the

highest end of the range of sequencing depths. Conventional mouse samples were sequenced to a

depth ranging from 39M to 75M reads (median = 45M reads) (**Table S1**).

Estimation of species abundances, SNVs, and CNVs

For the humanized mice and data from Shalon *et al.*, 2023[54], we used the Metagenomic Intra-

species Diversity Analysis System 1 (MIDAS 1; version 1.2, downloaded November 24, 2021)[26]

to calculate relative species abundances, gene copy number variants (CNVs), and single

nucleotide variants (SNVs). MIDAS 1 calculated these values by mapping reads against a

database of genomes representing 5,952 bacterial species[26,76] found in the human gut

microbiome.

The conventional mice microbiomes harbored bacteria native to mice and thus was not well-

suited to the MIDAS 1 database used to process the humanized mouse and human microbiomes,

as the reference genomes in the database corresponded only to human-associated bacterial

species resulting in fewer reads being mapped. We therefore used MIDAS 3 (version 3.0.1;

downloaded August 5, 2024) to construct a mouse-specific database using isolate and

metagenome assembled genome (MAG) reference genomes from the Mouse Gastrointestinal

Bacteria Catalogue (MGBC)[77] using a standard approach[78]. MIDAS 3 was subsequently used to

map reads to this database and calculate relative species abundances, CNVs, and SNVs.

While the outputs from both pipelines were generated with permissive parameters, as described

below, we subsequently applied several stringent post-processing filters to rule out metagenomic

artifacts and to accurately estimate species, CNVs, and SNVs. The post processing pipeline we

use is also briefly described below and reflects the pipeline developed in Garud *et al.*, 2019[23].

Estimating species relative abundance

For humanized mouse and Shalon *et al.*, 2023 data, MIDAS 1 was used to estimate species

presence and abundance by mapping reads to 15 single-copy marker genes from 5,952

representative species[26,76]. Reads were mapped with default MIDAS 1 settings. Species with a

median single copy marker gene coverage ≥3 reads were marked as present within a sample.

These read coverages were then used by MIDAS 1 to compute species relative abundance. For

conventional mouse data, MIDAS 3 was used estimated species presence and abundance in a

similar manner, this time to 15 single-copy marker genes from 1,093 representative species[77] in

the custom-built mouse-specific database. Species were marked as present if they had a median

marker coverage ≥ 2 reads, per the recommendation of MIDAS 3 developers[78,79].

Quantifying pathway relative abundance

To quantify changes in pathway abundances along the gut, we used HUMAnN 3 (Beghini et al.,

2021) version 3.9. HUMAnN3 was downloaded on August 11, 2025 and run with default

settings using the CHOCOPhlAn pangenome database version v201901_v31 and the UniRef90-

DIAMOND 2019-01 protein database obtained through the HUMAnN3 download utility both on

August 11, 2025. UniRef gene families were mapped to KEGG Ortholog (KO) groups using the

HUMAnN3-provided mapping file map_ko_uniref90.txt.gz (downloaded December 5, 2025).

Gene family and pathway read counts were summed across all taxa to generate community-wide

functional profiles. Pathway abundance was automatically normalized in the MaAsLin3 analysis

described below via the "total sum scaling" setting.

To identify pathways differing between the upper and lower gut, we analyzed the raw pathway

count matrix using the MaAsLin3[33] R package version 1.2.0. MaAsLin3 was downloaded on

December 7, 2025 and run with the default parameters on the raw pathway count matrix

(minimum feature abundance of 0.0001, minimum prevalence of 0.1, and default FDR

correction). We specified the model formula: ~ coarse_position + cage + read_counts +

strata(subject_id) with normalization = TSS, transformation = LOG. Gut location, read count and

cage were modeled as fixed-effect covariates and stratification by subject ID. P-values were

adjusted for multiple comparisons using the Benjamini–Hochberg false discovery rate (FDR)

procedure, as implemented in MaAsLin3, and pathways with q < 0.05 were considered

significant.

Quantifying SNV frequencies

We estimated SNV frequencies for each species in each sample. Downstream, these frequencies

were used to estimate nucleotide diversity, identify quasi-phaseable samples, and quantify SNV

frequency differences between pairs of quasi-phaseable samples. To estimate SNV frequencies in

humanized mouse and Shalon et al., 2023 samples, reads were mapped to representative

reference genomes using default MIDAS 1 mapping thresholds and arguments[23]. SNV

frequencies were estimated in conventional mouse data using MIDAS 3, with mapping

thresholds and arguments set to match MIDAS 1 defaults.

To obtain accurate estimates of SNV frequencies, we imposed coverage requirements. MIDAS 1

and 3 report the read coverage D for each site in the reference genome for each sample. We

excluded any sample with a mean $D < 5X$ across all protein coding sites, as was done in Garud *et*

al., 2019. We further excluded from analysis species non-zero coverage at $\geq 40\%$ of reference

sites within a sample. Additionally, to minimize the possibility of any mapping errors, we

implemented two controls. First, as described below and as was done in Garud *et al.*, 2019[23], we

exclude genes with copy number greater than 3 in at least one sample in our study, as they were

potential candidates for read stealing. Second, we exclude individual sites with a coverage value

$< 0.3\overline{D}$ or $> 3\overline{D}$ where $\overline{D}$ is mean genome-wide coverage, as these sites had abnormal coverages

outside the range of expectations for single copy genes.

Finally, SNV frequency changes were identified only if median coverage values between two

samples from the same mouse were within a factor of three of each other, as we expect that

coverage should be relatively constant within a mouse. Below we describe additional coverage

filters that were imposed for various calculations, including SNV frequency changes.

Annotating SNVs as synonymous versus nonsynonymous

MIDAS 1 annotated SNVs as synonymous and nonsynonymous based on the reading frames of genes as annotated in the PATRIC database[80], and MIDAS 3 annotated SNVs as synonymous and nonsynonymous based on annotations from Prokka[81].

Quantifying gene copy number

We estimate copy number variation (CNV) to exclude any genes from further analysis that may be potential candidates for read stealing or donating. To estimate CNVs in humanized mice and Shalon et al., 2023 data, reads were mapped to pangenomes using default MIDAS 1 mapping thresholds and arguments[23]. CNVs were estimated in conventional mouse data using MIDAS 3 with mapping thresholds and arguments set to match MIDAS 1 defaults.

To minimize read mapping errors, we identified genes with abnormally high gene copy number values, which may be potentially indicative of erroneous read recruitment[82]. MIDAS 1 and 3 estimates CNVs by mapping reads to a *pangenome* consisting of several sequenced isolates belonging to the same species. Average gene coverage was calculated as the number of reads mapping to that gene normalized by gene length. Both versions of MIDAS compute copy number, c, as the ratio between a gene's average coverage and the median coverage of a set of 15 single-copy marker genes belong to that species. Genes with a copy number $c > 3$ in at least one sample in our study were excluded during post-processing, as they were potential candidates for stealing. To exclude any additional genes that may suffer from erroneous read recruitment, we further filtered out any genes with a copy number $c > 3$ in the human samples analyzed in Garud *et al.*, 2019[23]. Finally, any genes sharing $\geq 95\%$ average nucleotide identity with any other gene across species boundaries as identified in a set of "blacklisted" genes in Garud *et al.*, 2019 (section A.iv)[23] were excluded to further prevent erroneous read mapping.

Calculating alpha and beta diversity of gut samples

Alpha diversity was computed for humanized and conventional mouse data using the Shannon diversity index as implemented by the Vegan package in R[83] (**Figure 2A**, **Supplementary Figure 9A**). To assess if Shannon diversity between gut regions was significantly different, we used a paired Wilcoxon signed-rank test comparing pairs of samples from one region versus the

other from the same mouse. To calculate fold change in alpha diversity between the small and

large intestines, alpha diversity was averaged across small intestinal regions (duodenum,

jejunum, and ileum) and large intestinal regions (cecum and colon), and the per mouse fold

change was calculated as the ratio of the averaged large intestinal and small intestinal alpha

diversity values. The fold change values presented in the main text represent the average of these

per-mouse estimates.

To produce the stacked bar plots in **Figure 2B** and **Supplementary Figure 9B**, species

abundances within taxonomic families were summed. Species with less than 0.1% abundance

were excluded. To test for differential abundance of specific families between the small and

large intestines of humanized mice, relative abundances for each family were averaged across the

small intestine (duodenum, jejunum, and ileum) and large intestine (cecum and colon) in each

host, the wilcox.test() in R was used to apply a paired Wilcoxon signed-rank test to compare

average large and small intestinal relative abundances (paired by mouse) for each species,

generating per-species estimates for the median log2 fold change in mean relative abundance

between large and small intestinal samples in the same mouse, as well as p-values and 95%

confidence intervals to assess significance (**Supplementary Figure 1**).

Calculating nucleotide diversity

For every humanized mouse and sample and the inoculum, we estimated π, a population-level

metric of nucleotide diversity that represents the probability of two randomly sampled genomes

from the population have different alleles at a base pair. To do so, we used the formula for

nucleotide diversity applied in Schloissnig *et al.*, 2013[84], which accounts for total read counts:

$$\pi(S,G) = \frac{1}{|G|} \sum_{i=1}^{|G|} \sum_{B_1 \in \{ATGC\}} \sum_{B_2 \in \{ATGC\} \setminus B_1} \frac{x_{i,B_1}}{D_i} \frac{x_{i,B_2}}{D_i - 1}$$

Here, S is the sample; G is the genome of the focal species, with $|G|$ being its size; i is the locus

in the genome; B_1 and B_2 are the reference and alternative alleles, respectively; x_{i,B_j} is the

number of reads with allele B_j at locus i; and D_i is the total read depth at locus i. This formula is

an extension of a previously proposed π estimator on next generation sequencing data devised by

Begun *et al.*, 2007[85]. Sites with read coverage less than four were excluded from π calculations,

such that $|G|$ is equivalent to the total number of sites with adequate coverage rather than the

total size of the genome.

Samples from mice 7 and 8 were sequenced to a substantially higher depth compared to samples

from mice 1 through 6 (**Supplementary Table 1**), resulting in elevated levels of coverage per bp

for bacterial species in mice 7 and 8. As a consequence, there was an increased opportunity to

detect higher levels of nucleotide diversity in mice 7 and 8 merely due to systematic differences

in coverage. To be able to compare nucleotide diversity levels between mice, we applied a

random down sampling per nucleotide site such that the depth of mice 7 and 8 matched the

median coverage of mice 1 through 6.

Distributions of π in **Figure 3A** were plotted for the 30 species which met the following criteria:

minimum coverage of four reads in the inoculum and at least three mouse samples representing

at least two different mouse hosts. Species were only considered in samples in which at least

500,000 nucleotides met the minimum coverage threshold.

Calculating F_{ST}

π estimates were used to compute genome-wide F_{ST} as a quotient of the mean genome-wide π

subtracted from 1:

782
$$F_{ST}(S_1, S_2, G) = 1 - \frac{\pi_{within}}{\pi_{between}} = 1 - \frac{\big(\pi(S_1, G) + \pi(S_2, G)\big)/2}{\pi(S_1, S_2, G)}$$

783 Where π_{within} is given by the formula for $\pi(S, G)$ provided above, and pairwise $\pi_{between}$ is

784 given by the following formula also derived in Schloissnig *et al.*, 2013:

785
$$\pi(S_1, S_2, G) = \frac{1}{|G|} \sum_{i=1}^{|G|} \sum_{B_1 \in \{ATGC\}} \sum_{B_2 \in \{ATGC\}\setminus B_1} \frac{x_{i,B_1,S_1}}{D_{i,S_1}} \frac{x_{i,B_2,S_2}}{D_{i,S_2}}$$

786 Where k in S_k designates sample 1 or 2, respectively. We used ileum and colon to represent the

787 small and large intestine, respectively because those regions had the highest mean read coverage

788   across all samples in each gut region. In order to accurately estimate allele frequencies, each site

789   was only included in the F_{ST} calculation if a minimum of 4 reads was observed at that site for

790   both samples being compared.

Inferring strain frequencies

Previous work has shown that hosts tend to be colonized by multiple genetically distinct strains

(typically 1-4) of the same species[23,86]. When levels of recombination between strains within a

host is sufficiently low, these strains may co-exist as genetically distinct subpopulations[24].

Typically, two randomly drawn strains from different hosts harbor on order 10^3 - 10^4 SNV

changes[23,47,84,87], and two strains residing within a host are expected to be similarly diverged[23].

We leverage this expected divergence between strains to distinguish between the strain

genotypes and infer their overall frequencies.

Inferring strain genotypes and frequencies from short read sequencing data is a difficult problem

because linkage between co-segregating SNVs residing on the same strain backbone are

destroyed by the fact that they are unlikely to reside on the same short read. Even perfectly

linked SNVs may be difficult to assign to the correct strain given the potential for sampling

noise. However, recent work has shown that strain inference is feasible when there are many

samples collected from the same hosts, as alleles belonging to the same strain will display highly

correlated allele frequencies across samples[24,25,65,88–90].

We leverage recent work by Roodgar *et al.*, 2021[24] and extended by Wolff *et al.*, 2023[25] to infer

strain identity and frequency in samples from all three datasets analyzed in this study. The papers

cited above inferred strain genotypes and frequencies from *temporal* data. Here, we use *spatial*

data both within mice and across mice, and a mixture of *spatial* and *temporal* data within human

hosts sampled in Shalon *et al.*, 2023 to infer strain genotypes and frequencies within samples.

Because humanized mice were inoculated with the same human sample and conventional mice

shared homogenized microbiomes due to co-housing, we inferred strain frequency jointly across

all mouse samples belonging to the same dataset, reasoning that mice from the same dataset all

shared the same subset of strains. For Shalon *et al.*, 2023 data, we inferred strain frequency

jointly across all samples belonging to the same host, likewise reasoning that these samples

carried the same subset of strains.

Briefly, the approach clusters SNVs segregating on the same strain background by detecting

correlations between SNVs with highly similar allele frequencies across samples. The degree of

correlation between a pair of SNVs i and j is measured by distance metric d between allele

frequency trajectories $\hat{f}_i$ and $\hat{f}_j$:

821
$$d(\hat{f}_i, \hat{f}_j) = \frac{1}{S}\sum_{s=1}^{S} \frac{2(D_{is} + D_{js})(\hat{f}_{is} - \hat{f}_{js})^2}{(\hat{f}_{is} + \hat{f}_{js})(1 - \hat{f}_{is} + 1 - \hat{f}_{js})},$$

where S is the number of samples, $\hat{f}_{is}$ and $\hat{f}_{js}$ are the frequency of the alleles in a sample s, and

D_{is} and D_{js} represent the read depth of the alleles in sample s. SNVs that are in perfect linkage

with one another are expected to have a $d = 0$.

To ensure that high-quality and informative sites were included in our stain inference procedure,

we only considered loci that had a read coverage of $D \geq 10$ and were polymorphic in at least

20% of samples in which the species was detected. Filtering out non-polymorphic loci prevented

the algorithm from generating clusters at allele frequencies of 1 or 0 that reflect low frequency

variation within a strain population as opposed to actual genetically distinct strains.

Detecting correlated clusters of SNVs belonging to the same strain background can be

confounded without knowing if the alternative or reference allele belongs to one strain versus

another. To address this issue, as done in Roodgar *et al.*, 2021[24], we set d between a pair of loci

to the minimum of $d(\hat{f}_i, \hat{f}_j)$ and $d(\hat{f}_i, 1 - \hat{f}_j)$ to ensure that polarization differences (i.e., calling

an allele frequency 0.2 instead of 0.8, based on the reference allele used in the reference genome)

between loci did not prevent them from being clustered together.

To identify SNVs linked on the same strain background, we applied a greedy, network-based

algorithm to extract large cluster SNVs with highly correlated allele frequencies across samples.

The algorithm begins by forming a network in which nodes represent SNVs, with pairs SNVs

connected by an edge if they have a distance $d < 3.5$, which represents a maximum distance

threshold below which pairs of SNVs are likely to be linked on the same strain background, as

demonstrated in Roodgar *et al.*, 2021[24]. Next, the algorithm identifies a focal SNV as the SNV

with the maximum number of connections (i.e., edges) and extracts all polymorphic SNVs

connected to it. This cluster of SNVs was designated as a strain if it included at least 10^3 SNVs,

as this represents the lower bound for the typical number of alleles that are expected to segregate

between genetically distinct strains[23,47,84,87]. Note that SNVs connected to the focal SNV were

only extracted if they were also connected to at least 25% of other SNVs within a cluster. SNVs

included in the initial cluster were removed from the overall set of variable sites, and the

clustering process was repeated until no additional clusters of at least 10^3 SNVs were identified.

Strain frequencies were then inferred as the mean SNV allele frequencies in each cluster

(**Supplementary Figure 14**). However, despite requiring strain clusters to have a minimum of

10^3 SNVs, not all SNVs met our coverage requirement of $D \geq 10$ across all samples.

Therefore, we only report strain frequencies for samples in which every cluster has at least 100

high coverage SNVs ($D \geq 10$) to ensure that the inferred strain frequencies were well-

supported.

95% confidence intervals for strain frequencies in each sample were generated using a stationary

bootstrapping approach[91]. Specifically, 1,000 bootstrap samples each consisting of 100 SNVs

were drawn with replacement from each strain cluster within each sample, and subsequently the

mean of the subsampled SNV frequencies in each bootstrap sample was computed as strain

frequency. The upper and lower bounds of the 95% confidence interval was calculated as the 2.5

and 97.5 percentile of the bootstrapped strain frequency distribution.

In addition to the filters used by Wolff *et al.*, 2023, we required minor alleles to have a read

support of at least four reads to prevent rare mutations and sequencing errors from erroneously

giving rise to spurious low-frequency clusters inferred from conventional and humanize mouse

data. For strain frequency profiles in Shalon *et al.*, 2023, we filtered out spurious low-frequency

clusters and lowly abundant strains using a different approach by requiring all strains rise above

a frequency of 5% in at least one sample to be included in downstream analyses.

We visually inspected all strain frequency profiles and either applied manual modifications to the

algorithm or excluded from downstream analysis all profiles which violated one of the following

expectations. In a scenario when only one strain is present, we expect *zero* clusters to be

detected. In a scenario in which two strains are present, we expect only *one* cluster, representing

the alleles segregating between the two strains. The relative frequencies of the two strains can be

inferred as $\bar{f}$ and $1 - \bar{f}$, where $\bar{f}$ is the mean frequency of the clustered SNVs. In a scenario in

which three strains are present, we expect *three* clusters the frequencies of which roughly sum to

1 in each sample, representing the SNV segregating between each of the three pairs of strains.

This second scenario did not occur in our humanized or conventional mouse data. In the Shalon

et al., 2023 data, the strain frequency inference pipeline did at times yield a maximum of three

strain clusters. However, manual inspection revealed that these clusters tended to sum to an

overall frequency significantly greater or less than 1 in some samples, suggesting that the strain

frequency pipeline was failing to accurately capture all strains or spuriously inferring too many.

As a result, we excluded species x host pairs with three strain clusters and limited our analysis to

species x host pairs that had two strains. At times, the strain inference algorithm outputs two

clusters, which is inconsistent with a scenario of either 1, 2 or 3 strains colonizing and instead

may be arising from high variance in allele frequency. This happened once in the humanized

mouse data with *B. uniformis*, the strain inference algorithm inferred *two* clusters

(**Supplementary Figure 15A**). It was visually apparent in this case that the two clusters likely

represented the same group of SNVs segregating between two strains' genetic backgrounds

because of their highly correlated trajectories, but were separated into two distinct clusters due to

there being a large variance in individual SNV frequency. In this singular instance, we

reclustered by using the mean SNV frequency of all SNVs in both clusters as a focal SNV

(**Supplementary Figure 15B**) and extracting all other SNVs connected to it by an edge in the

network (i.e., with $d < 3.5$). This method produced a single cluster. The algorithm was able to

infer strain frequencies for all other species analyzed in this study from the humanized and

conventional mouse datasets without the need for *ad hoc* modifications to the pipeline. For

several species x host pairs in the Shalon *et al.*, 2023 data, the strain inference algorithm yielded

two clusters. These species x host pairs were excluded from downstream analysis of strain

frequency in favor of species x host pairs with more unambiguous strain phasing results, all of

which had two strains (*i.e.*, a single cluster).

When applying this algorithm to the humanized mouse data, it in some instances inferred only

single strain for a bacterial species across all samples despite the inoculum displaying high

nucleotide diversity, suggestive of multiple strains being present in the inoculum. This

phenomenon may occur when applying the algorithm to a collection of samples in which a strain

is present in only one sample (*e.g.,* the inoculum) but absent in all others (*e.g.*, the mouse

samples). As a result, the algorithm would be unable to detect correlations between the SNVs

private to a single sample and therefore fail to detect that strain. We identified when this might

be happening in the humanized mouse data by looking for species in which only a single strain

was inferred (i.e., the algorithm produced no clusters), but which had at least 1,000 SNVs in the

inoculum that were polymorphic and had a read support of four reads for the minor allele. When

this scenario occurred, we assumed two strains were present in the inoculum and inferred the

strain frequency $f_{strain\ 1,inoculum}$ of the first strain as the mean of the allele frequencies of these

polymorphic SNVs in the inoculum, and the strain frequency $f_{strain\ 2,inoculum}$ of the second

strain as $1 - f_{strain\ 1,inoculum}$. We generated 95% confidence intervals by applying the same

stationary bootstrapping approach to the polymorphic SNVs identified in the inoculum.

Calculating change in major strain frequency Δf between samples

The change in major strain frequency Δf was calculated as the magnitude of the difference in

frequencies of one of the strains between two samples. Because all samples with co-colonizing

strain analyzed in this study harbored two strains, Δf was the same regardless of whether strain 1

or 2 was used to calculate the frequency change. For the humanized and conventional mouse

data, we calculated Δf between samples for co-colonizing species present in at least at least two

mice, at least two cages, and at least two gut regions. When visualizing the distribution of Δf

values, we categorized pairwise comparisons as being "within host", "between host, within

cage", and "between host, between cage". For each species, we selected a single "within host"

pair per mouse, a single "between host, within cage" pair per cage, and a single "between host,

between cage" pair for the three possible cage-wise comparisons. For "within host" comparison

within each host, we selected the pair of samples that maximized the distance between the two

sampling sites, breaking ties by selecting the pair of samples with the sample that is furthest

down the gut (i.e., closest to the colon). For between host comparisons (both within and between

cage), we selected the pair of samples that represented the same gut region or two gut regions

proximal to one another along the gut, breaking ties by selecting pairs of samples with the

sample that is furthest down the gut (i.e., closest to the colon). For "between host, between cage"

comparisons, we ensured that no mouse was involved in more than one cross-cage comparison to

eliminate redundancy. For Shalon et al., 2023 data, we calculated Δf between samples belonging

to co-colonized species x host pairs if the species were present in at least two timepoints and at

least two device types. When visualizing the distribution of Δf values, we categorized pairwise

comparisons as being "within timepoint, between gut region", "between consecutive timepoints",

and "between host first and last timepoint". For each species x host pair, we selected a single

"within timepoint" pair per timepoint, a single "between consecutive timepoints" pair for each

consecutive pair of timepoints, and a single "between host first and last timepoint" pair. For

"within timepoint" comparisons within each timepoint, we selected the sample pair that

maximized the distance between device types (i.e., biasing towards device type 1 versus device

type 4 comparisons, if available), breaking ties by selecting the sample pair with the maximum

device type number (i.e., biasing towards pairs with device type 4). For between timepoint

comparisons (both consecutive and first and last timepoints), the sample pair that minimized the

distance between device types and was furthest along the gut was selected.

Performing ANOVA of major strain frequencies

To quantify the variance in major strain frequency across samples, we applied an ANOVA

approach. In the humanized mouse dataset, the major strain was designated as the strain at

frequency $f > 0.5$ in the inoculum. In conventional mouse and Shalon *et al.*, 2023 data, the

major strain was designated as the strain at frequency $f > 0.5$ in a majority of samples. The

centered log ratio (CLR) transformation was applied to major strain frequency in all samples in

which that species was detected, a commonly used approach to remove compositional artifacts

when applying non-compositional approaches such as ANOVA[92,93]. For each species, we built an

ANOVA model using the Stats package in R. In the humanized and conventional mouse data, the

model was specified with CLR-transformed major strain frequency as the dependent variable and

cage, mouse, and gut region as the independent categorical variables, and was applied jointly to

all samples belonging to the same study. Variance explained was calculated as the sum of

squares between groups corresponding to each variable divided by the sum of squares total. We

applied several prevalence requirements to ensure that the amount of variance explained by each

dependent variable was properly quantified. In humanized and conventional mouse data, we only

performed ANOVA on species detected across at least two cages (with the species detectable in

at least two mice in those cages), at least two mice (with the species detectable in at least two gut

regions in those mice), and two gut regions (with the species detectable in these same gut regions

in at least two mice).

Identifying quasi-phaseable lineages

SNV frequency differences between samples can arise due to evolution, or due to shifts in the

frequency of co-colonizing strains, which are likely to harbor thousands of differing nucleotides

between them. To distinguish between SNV frequency differences that arise due to evolution

versus those arising from strain frequency shifts, we identified quasi-phaseable lineages using an

approach developed in Garud *et al.*, 2019[23]. By inferring quasi-phaseable lineages, we then were

able to confidently infer evolutionary changes.

Briefly, the approach works by identifying samples in which a particular species has a single

dominant strain at $f \geq 0.8$, hereafter referred to as a quasi-phaseable sample. In such samples,

the allele corresponding to the dominant strain can be inferred with statistical confidence.

Specifically, the alleles with frequency $f \geq 0.8$ at genomic loci with coverage $D \geq 20$ are

assigned to the quasi-phaseable sample. Only samples with a median coverage of 20X for the

species of interest are processed through the quasi-phaseable pipeline. Employing such stringent

coverage and allele frequency thresholds ensures that the incorrect allele is not inferred, as it is

statistically improbable that sampling error would give rise to the incorrect allele[23].

We made an important alteration to the quasi-phaseable pipeline in its application to the

humanized and conventional mouse data relative to its application in Garud *et al.*, 2019[23].

Normally, the quasi-phaseable pipeline designates a sample as having a quasi-phaseable strain if

the number of intermediate frequency SNVs (i.e., SNVs with frequency $0.2 < f < 0.8$) which a

species harbors within the sample are 10% or less than the number of SNV differences that are

typically observed for that species between independent hosts. This heuristic arises from the

assumption that independent hosts should harbor genetically distinct strains that have diverged

over long evolutionary timescales, and is born out by empirical evidence[23]. The 10% threshold is

calibrated independently for each species, as genetically distinct strains of one species are not

necessarily expected to show the same level of nucleotide differentiation as genetically distinct

strains of a different species. In previous applications of the quasi-phaseable pipeline, the

number of SNV differences between independent hosts has been assessed from the dataset used

in those studies. For example, Garud *et al.*, 2019[23] studied evolutionary dynamics within 693

unique individuals sampled as a part of the human microbiome project[94,95], the TwinsUK

registry[96], a cohort of Chinese subjects[97], and four additional young twins[98]. Because most hosts

included in this compiled dataset were unrelated (and in some cases, on different continents), the

assumption that their resident strains were distantly diverged was a reasonable one. By contrast,

the humanized and conventional mice are expected to have low between-host strain diversity,

either because they were inoculated with the stool from a single human host or because their

microbiomes were homogenized early on in development through co-housing. As a result, we

could not accurately quantify the number of differences that are expected between two unrelated

hosts, making it impossible to detect quasi-phaseable strains using this dataset alone. We took

two different approaches to overcome this issue in humanized and conventional mice,

respectively. In humanized mice, we calculated the expected number of between-host SNV

differences for each focal species based on the aforementioned data used in Garud *et al.*, 2019[23],

which we had previously processed through MIDAS 1. Because this resource was not available

for the conventional mouse data processed through MIDAS 3, we designated samples as quasi-

phaseable if intermediate frequency sites (i.e., SNVs with frequency $0.2 < f < 0.8$) occurred at

a rate of 1×10^{-4} per bp or less, consistent with rates observed in quasi-phaseable samples from

humanized mice. Results published in this study (**Figure 3A**) and elsewhere[23] suggest that this is

a highly conservative threshold for the maximum amount of intermediate frequency diversity we

would expect to see if only a single strain were present within a host.

Inferring SNV frequency changes

To detect SNV frequency changes arising as a result of evolution, we follow the approach in

Garud *et al.*, 2019. A SNV frequency change is defined as an allele with low frequency in one

quasi-phaseable sample ($f \leq 0.2$) and high frequency in another quasi-phaseable sample ($f \geq$

0.8). SNV differences were only considered at loci that had a coverage $D \geq 20$ in both samples

being compared. An extreme allele frequency change of this magnitude and this minimum depth

is statistically unlikely to arise between quasi-phaseable strains due to sampling error or drift

given large population sizes and short timescales considered in this study[23,43]. Pairs of samples

harboring ≤ 20 SNV differences were considered to have the same inoculum strain evolving over

short timescales (e.g., the eight-week timescale considered in this study), as larger amounts of

SNV differences, such as $\mathcal{O}(10^3)$ or greater are inconsistent with within-host rates of

diversification[47,82,84,87]. Consequently, pairs of quasi-phaseable strains with > 20 SNV differences

were excluded from this evolutionary analysis. For more information, see the Supplementary

Information for Garud *et al.*, 2019[23].

Bootstrapping confidence intervals for SNV difference rates

To measure whether there were significant differences in the number of SNV frequency changes

occurring between samples within hosts, between hosts, and between the inoculum and mouse

hosts in the humanized mouse data, we employed a bootstrap approach to compute confidence

intervals for the rate of SNV difference between quasi-phaseable pairs of samples with the same

dominant strain. In 1,000 bootstrap iterations, we sampled 100 quasi-phaseable sample pairs (not

necessarily of the same species) with replacement from each comparison category: within host,

between host, and between inoculum and host. For the within-cage comparison category, we

only considered quasi-phaseable pairs from different hosts. We computed the actual rate of SNV

frequency changes for a particular comparison category as the total number of SNVs displaying

extreme allele frequency changes across all sampled quasi-phaseable pairs divided by the total

number of high coverage loci across all sampled quasi-phaseable pairs. We calculated 95%

confidence intervals using the cumulative distribution function (CDF) of the overall SNV change

difference rate in each comparison category generated from the 1,000 bootstrap iterations, taking

the 2.5 and 97.5 percentile as the upper and lower bound, respectively (**Figure 5B**). We applied

an identical approach for assessing whether SNV frequency change rates were significantly

different between samples from the same host or different hosts in the conventional mouse data

(**Figure 6C**).

Data Availability Statement

Raw metagenomics sequencing reads are available on NCBI under BioProject [project accession

number will be made available upon publication]. All necessary metadata, as well as the source

code for the sequencing pipeline, downstream analyses, and figure generation, are available at
GitHub (https://github.com/garudlab/Wasney-Briscoe/).

Acknowledgments

The authors thank Kristin Harper, PhD, of Harper Health & Science Communications, LLC, for providing editorial support in accordance with Good Publication Practice guidelines, and Alison Feder, PhD, for her helpful edits. The authors also thank the Garud and the Tropini labs for their valuable inputs. In particular, the authors thank Peter Laurin for his statistical advice and feedback on the manuscript. Finally, the authors thank Dr. Sidhartha Sinha and members of the Sinha Lab at Stanford University for helpful discussions. This work was funded by NIGMS NIH award R35GM151023 (to NRG), NSF CAREER award (no. 2240098) (to NRG), a Paul Allen Distinguished Investigator Award (to CT and NRG), a Canadian Institutes of Health Research Team Grant: Canadian Microbiome Initiative 2 (to CT), Crohn's and Colitis Canada, Canadian Institute for Advanced Research (to CT), and the Michael Smith Foundation for Health Research Scholar Award (to CT).

References

1. McCallum, G. & Tropini, C. The gut microbiota and its biogeography. *Nat. Rev. Microbiol.* **22**, 105–118 (2024).
2. Gu, S. *et al.* Bacterial Community Mapping of the Mouse Gastrointestinal Tract. *PLOS ONE* **8**, e74957 (2013).
3. Ng, K. M. *et al.* Single-strain behavior predicts responses to environmental pH and osmolality in the gut microbiota. *mBio* **14**, e00753-23 (2023).
4. D'Adamo, G. L. *et al.* Bacterial clade-specific analysis identifies distinct epithelial responses in inflammatory bowel disease. *Cell Rep. Med.* **4**, (2023).
5. Morowitz, M. J., Carlisle, E. & Alverdy, J. C. Contributions of Intestinal Bacteria to Nutrition and Metabolism in the Critically Ill. *Surg. Clin. North Am.* **91**, 771–785 (2011).
6. Miller, B. M., Liou, M. J., Lee, J.-Y. & Bäumler, A. J. The longitudinal and cross-sectional heterogeneity of the intestinal microbiota. *Curr. Opin. Microbiol.* **63**, 221–230 (2021).
7. Dapa, T. *et al.* Within-host evolution of the gut microbiome. *Curr. Opin. Microbiol.* **71**, 102258 (2023).
8. Dapa, T., Ramiro, R. S., Pedro, M. F., Gordo, I. & Xavier, K. B. Diet leaves a genetic signature in a keystone member of the gut microbiota. *Cell Host Microbe* **30**, 183-199.e10 (2022).
9. Yang, Y. *et al.* Within-host evolution of a gut pathobiont facilitates liver translocation. *Nature* **607**, 563–570 (2022).
10. Collins, J. *et al.* Dietary trehalose enhances virulence of epidemic Clostridium difficile. *Nature* **553**, 291–294 (2018).

11.   Paepe, M. D. *et al.* Trade-Off between Bile Resistance and Nutritional Competence Drives
Escherichia coli Diversification in the Mouse Gut. *PLOS Genet.* **7**, e1002107 (2011).
12.   Conway, T. & Cohen, P. S. Commensal and Pathogenic Escherichia coli Metabolism in the Gut.
in *Metabolism and Bacterial Pathogenesis* 343–362 (John Wiley & Sons, Ltd, 2015).
doi:10.1128/9781555818883.ch16.
13.   Barroso-Batista, J. *et al.* Specific Eco-evolutionary Contexts in the Mouse Gut Reveal
Escherichia coli Metabolic Versatility. *Curr. Biol. CB* **30**, 1049-1062.e7 (2020).
14.   Vasquez, K. S. *et al.* Quantifying rapid bacterial evolution and transmission within the mouse
intestine. *Cell Host Microbe* **29**, 1454-1468.e4 (2021).
15.   Kim, S. G. *et al.* Microbiota-derived lantibiotic restores resistance against vancomycin-resistant
Enterococcus. *Nature* **572**, 665–669 (2019).
16.   Dubinsky, V., Dotan, I. & Gophna, U. Strains Colonizing Different Intestinal Sites within an
Individual Are Derived from a Single Founder Population. *mBio* **0**, e03456-22 (2023).
17.   Wu-Woods, N. J. *et al.* Microbial-enrichment method enables high-throughput metagenomic
characterization from host-rich samples. *Nat. Methods* **20**, 1672–1682 (2023).
18.   Johansson, M. E. V. *et al.* Normalization of Host Intestinal Mucus Layers Requires Long-Term
Microbial Colonization. *Cell Host Microbe* **18**, 582–592 (2015).
19.   Seedorf, H. *et al.* Bacteria from diverse habitats colonize and compete in the mouse gut. *Cell* **159**,
253–266 (2014).
20.   Turnbaugh, P. J. *et al.* The Effect of Diet on the Human Gut Microbiome: A Metagenomic
Analysis in Humanized Gnotobiotic Mice. *Sci. Transl. Med.* **1**, 6ra14 (2009).
21.   Sonnenburg, E. D. *et al.* Diet-induced extinction in the gut microbiota compounds over
generations. *Nature* **529**, 212–215 (2016).
22.   Tikhonov, M., Leach, R. W. & Wingreen, N. S. Interpreting 16S metagenomic data without
clustering to achieve sub-OTU resolution. *ISME J.* **9**, 68–80 (2015).
23.   Garud, N. R., Good, B. H., Hallatschek, O. & Pollard, K. S. Evolutionary dynamics of bacteria in
the gut microbiome within and across hosts. *PLOS Biol.* **17**, e3000102 (2019).
24.   Roodgar, M. *et al.* Longitudinal linked-read sequencing reveals ecological and evolutionary
responses of a human gut microbiome during antibiotic treatment. *Genome Res.* **31**, 1433–1446 (2021).
25.   Wolff, R., Shoemaker, W. & Garud, N. Ecological Stability Emerges at the Level of Strains in the
Human Gut Microbiome. *mBio* **14**, e02502-22 (2023).
26.   Nayfach, S., Rodriguez-Mueller, B., Garud, N. & Pollard, K. S. An integrated metagenomics
pipeline for strain profiling reveals novel patterns of bacterial transmission and biogeography. *Genome
Res.* **26**, 1612–1625 (2016).
27.   Faith, J. J. *et al.* The Long-Term Stability of the Human Gut Microbiota. *Science* **341**, 1237439
(2013).
28.   Tropini, C., Earle, K. A., Huang, K. C. & Sonnenburg, J. L. The Gut Microbiome: Connecting
Spatial Organization to Function. *Cell Host Microbe* **21**, 433–442 (2017).
29.   Donaldson, G. P., Lee, S. M. & Mazmanian, S. K. Gut biogeography of the bacterial microbiota.
Nat. Rev. Microbiol. **14**, 20–32 (2016).
30.   Vaga, S. *et al.* Compositional and functional differences of the mucosal microbiota along the
intestine of healthy individuals. *Sci. Rep.* **10**, 14977 (2020).
31.   Zafar, H. & Saier, M. H. Gut Bacteroides species in health and disease. *Gut Microbes* **13**,
1848158.

32.	Beghini, F. *et al.* Integrating taxonomic, functional, and strain-level profiling of diverse microbial
communities with bioBakery 3. *eLife* **10**, e65088 (2021).

33.	Nickols, W. A. *et al.* MaAsLin 3: Refining and extending generalized multivariable linear models
for meta-omic association discovery. *BioRxiv Prepr. Serv. Biol.* 2024.12.13.628459 (2024)
doi:10.1101/2024.12.13.628459.

34.	Li, X. *et al.* Uncovering de novo polyamine biosynthesis in the gut microbiome and its alteration
in inflammatory bowel disease. *Gut Microbes* **17**, 2464225 (2025).

35.	Kennedy, M. S. *et al.* Dynamic genetic adaptation of *Bacteroides thetaiotaomicron* during murine
gut colonization. *Cell Rep.* **42**, 113009 (2023).

36.	Candeliere, F. *et al.* β-Glucuronidase Pattern Predicted From Gut Metagenomes Indicates
Potentially Diversified Pharmacomicrobiomics. *Front. Microbiol.* **13**, (2022).

37.	Walter, J. Ecological Role of Lactobacilli in the Gastrointestinal Tract: Implications for
Fundamental and Biomedical Research. *Appl. Environ. Microbiol.* **74**, 4985–4996 (2008).

38.	Mager, L. F. *et al.* Microbiome-derived inosine modulates response to checkpoint inhibitor
immunotherapy. *Science* **369**, 1481–1489 (2020).

39.	Wright, S. The Genetical Structure of Populations. *Ann. Eugen.* **15**, 323–354 (1949).

40.	Wright, S. The genetical structure of populations. *Ann. Eugen.* **15**, 323–354 (1951).

41.	Frazão, N. & Gordo, I. Shared Evolutionary Path in Social Microbiomes. *Mol. Biol. Evol.* **40**,
msad153 (2023).

42.	Vasquez, K. S. *et al.* High-resolution lineage tracking of within-host evolution and strain
transmission in a human gut symbiont across ecological scales. 2024.02.17.580834 Preprint at
https://doi.org/10.1101/2024.02.17.580834 (2024).

43.	Ghosh, O. M. & Good, B. H. Emergent evolutionary forces in spatial models of luminal growth
and their application to the human gut microbiota. *Proc. Natl. Acad. Sci.* **119**, e2114931119 (2022).

44.	Wichman, H. A., Badgett, M. R., Scott, L. A., Boulianne, C. M. & Bull, J. J. Different
Trajectories of Parallel Evolution During Viral Adaptation. *Science* **285**, 422–424 (1999).

45.	Woods, R., Schneider, D., Winkworth, C. L., Riley, M. A. & Lenski, R. E. Tests of parallel
molecular evolution in a long-term experiment with Escherichia coli. *Proc. Natl. Acad. Sci.* **103**, 9107–
9112 (2006).

46.	Lieberman, T. D. *et al.* Parallel bacterial evolution within multiple patients identifies candidate
pathogenicity genes. *Nat. Genet.* **43**, 1275–1280 (2011).

47.	Zhao, S. *et al.* Adaptive Evolution within Gut Microbiomes of Healthy People. *Cell Host Microbe*
25, 656-667.e8 (2019).

48.	Wong, D. P. G. H. & Good, B. H. Quantifying the adaptive landscape of commensal gut bacteria
using high-resolution lineage tracking. *Nat. Commun.* **15**, 1605 (2024).

49.	Gómez-Santos, N., Glatter, T., Koebnik, R., Świątek-Połatyńska, M. A. & Søgaard-Andersen, L.
A TonB-dependent transporter is required for secretion of protease PopC across the bacterial outer
membrane. *Nat. Commun.* **10**, 1–18 (2019).

50.	Moreno-Gámez, S. *et al.* Niche partitioning by resource size in the gut microbiome.
2025.11.13.688124 Preprint at https://doi.org/10.1101/2025.11.13.688124 (2025).

51.	Mills, R. H. *et al.* Multi-omics analyses of the ulcerative colitis gut microbiome link Bacteroides
vulgatus proteases with disease severity. *Nat. Microbiol.* **7**, 262–276 (2022).

52.	Chen, D. W. & Garud, N. R. Rapid evolution and strain turnover in the infant gut microbiome.
Genome Res. **32**, 1124–1136 (2022).

53.   Cremer, J., Arnoldini, M. & Hwa, T. Effect of water flow and chemical environment on
microbiota growth and composition in the human colon. *Proc. Natl. Acad. Sci.* **114**, 6438–6443 (2017).

54.   Shalon, D. *et al.* Profiling the human intestinal environment under physiological conditions.
Nature **617**, 581–591 (2023).

55.   Pédron, T. *et al.* A Crypt-Specific Core Microbiota Resides in the Mouse Colon. *mBio* **3**,
10.1128/mbio.00116-12 (2012).

56.   Li, H. *et al.* The outer mucus layer hosts a distinct intestinal microbial niche. *Nat. Commun.* **6**,
8292 (2015).

57.   Yasuda, K. *et al.* Biogeography of the Intestinal Mucosal and Lumenal Microbiome in the Rhesus
Macaque. *Cell Host Microbe* **17**, 385–391 (2015).

58.   Desai, M. S. *et al.* A Dietary Fiber-Deprived Gut Microbiota Degrades the Colonic Mucus Barrier
and Enhances Pathogen Susceptibility. *Cell* **167**, 1339-1353.e21 (2016).

59.   Chassaing, B. & Gewirtz, A. T. Identification of Inner Mucus-Associated Bacteria by Laser
Capture Microdissection. *Cell. Mol. Gastroenterol. Hepatol.* **7**, 157–160 (2019).

60.   Mas-Lloret, J. *et al.* Gut microbiome diversity detected by high-coverage 16S and shotgun
sequencing of paired stool and colon sample. *Sci. Data* **7**, 92 (2020).

61.   Duncan, K., Carey-Ewend, K. & Vaishnava, S. Spatial analysis of gut microbiome reveals a
distinct ecological niche associated with the mucus layer. *Gut Microbes* **13**, 1874815 (2021).

62.   Mondragón-Palomino, O. *et al.* Three-dimensional imaging for the quantification of spatial
patterns in microbiota of the intestinal mucosa. *Proc. Natl. Acad. Sci.* **119**, e2118483119 (2022).

63.   Guo, Y. *et al.* Oral pathobiont Klebsiella chaperon usher pili provide site-specific adaptation for
the inflamed gut mucosa. *Gut Microbes* **16**, 2333463 (2024).

64.   Lenski, R. E. Experimental evolution and the dynamics of adaptation and genome evolution in
microbial populations. *ISME J.* **11**, 2181–2194 (2017).

65.   Goyal, A., Bittleston, L. S., Leventhal, G. E., Lu, L. & Cordero, O. X. Interactions between
strains govern the eco-evolutionary dynamics of microbial communities. *eLife* **11**, e74987 (2022).

66.   Feng, L. *et al.* Identifying determinants of bacterial fitness in a model of human gut microbial
succession. *Proc. Natl. Acad. Sci.* **117**, 2622–2633 (2020).

67.   Conwill, A. *et al.* Anatomy promotes neutral coexistence of strains in the human skin
microbiome. *Cell Host Microbe* **30**, 171-182.e7 (2022).

68.   Lee, S. M. *et al.* Bacterial colonization factors control specificity and stability of the gut
microbiota. *Nature* **501**, 426–429 (2013).

69.   Jin, X. *et al.* Culturing of a complex gut microbial community in mucin-hydrogel carriers reveals
strain- and gene-associated spatial organization. *Nat. Commun.* **14**, 3510 (2023).

70.   Gutiérrez-García, K. *et al.* A conserved genetic basis for commensal-host specificity through live
imaging of colonization dynamics. 2024.04.19.590229 Preprint at
https://doi.org/10.1101/2024.04.19.590229 (2024).

71.   Tang, Q. *et al.* Current Sampling Methods for Gut Microbiota: A Call for More Precise Devices.
Front. Cell. Infect. Microbiol. **10**, (2020).

72.   Lloyd-Price, J. *et al.* Multi-omics of the gut microbial ecosystem in inflammatory bowel diseases.
Nature **569**, 655–662 (2019).

73.   Nishihara, Y. *et al.* Mucosa-associated gut microbiota reflects clinical course of ulcerative colitis.
Sci. Rep. **11**, 13743 (2021).

74.   Atreya, R. & Siegmund, B. Location is important: differentiation between ileal and colonic
Crohn's disease. *Nat. Rev. Gastroenterol. Hepatol.* **18**, 544–558 (2021).
75.   Kordahi, M. C. *et al.* Genomic and functional characterization of a mucosal symbiont involved in
early-stage colorectal cancer. *Cell Host Microbe* **29**, 1589-1598.e6 (2021).
76.   Wu, D., Jospin, G. & Eisen, J. A. Systematic identification of gene families for use as 'markers'
for phylogenetic and phylogeny-driven ecological studies of bacteria and archaea and their major
subgroups. *PloS One* **8**, e77033 (2013).
77.   Beresford-Jones, B. S. *et al.* The Mouse Gastrointestinal Bacteria Catalogue enables translation
between the mouse and human gut microbiotas via functional mapping. *Cell Host Microbe* **30**, 124-
138.e8 (2022).
78.   Smith, B. J. *et al.* Accurate estimation of intraspecific microbial gene content variation in
metagenomic data with MIDAS v3 and StrainPGC. 2024.04.10.588779 Preprint at
https://doi.org/10.1101/2024.04.10.588779 (2024).
79.   Zhao, C., Dimitrov, B., Goldman, M., Nayfach, S. & Pollard, K. S. MIDAS2: Metagenomic
Intra-species Diversity Analysis System. *Bioinformatics* **39**, btac713 (2023).
80.   Wattam, A. R. *et al.* Improvements to PATRIC, the all-bacterial Bioinformatics Database and
Analysis Resource Center. *Nucleic Acids Res.* **45**, D535–D542 (2017).
81.   Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068–2069
(2014).
82.   Garud, N. R., Good, B. H., Hallatschek, O. & Pollard, K. S. Evolutionary dynamics of bacteria in
the gut microbiome within and across hosts. *PLOS Biol.* **17**, e3000102 (2019).
83.   OKSANEN, J. Vegan : community ecology package. *Httpveganr-Forg.-Proj.*
https://cir.nii.ac.jp/crid/1570291225091856896 (2010).
84.   Schloissnig, S. *et al.* Genomic variation landscape of the human gut microbiome. *Nature* **493**, 45–
50 (2013).
85.   Begun, D. J. *et al.* Population Genomics: Whole-Genome Analysis of Polymorphism and
Divergence in Drosophila simulans. *PLOS Biol.* **5**, e310 (2007).
86.   Truong, D. T., Tett, A., Pasolli, E., Huttenhower, C. & Segata, N. Microbial strain-level
population structure and genetic diversity from metagenomes. *Genome Res.* **27**, 626–638 (2017).
87.   Van Rossum, T., Ferretti, P., Maistrenko, O. M. & Bork, P. Diversity within species: interpreting
strains in microbiomes. *Nat. Rev. Microbiol.* **18**, 491–506 (2020).
88.   Quince, C. *et al.* STRONG: metagenomics strain resolution on assembly graphs. *Genome Biol.*
22, 214 (2021).
89.   Kang, J. T. L. *et al.* Long-term ecological and evolutionary dynamics in the gut microbiomes of
carbapenemase-producing Enterobacteriaceae colonized subjects. *Nat. Microbiol.* **7**, 1516–1524 (2022).
90.   Zheng, W. *et al.* High-throughput, single-microbe genomics with strain resolution, applied to a
human gut microbiome. *Science* **376**, eabm1483 (2022).
91.   Politis, D. N. & Romano, J. P. The Stationary Bootstrap. *J. Am. Stat. Assoc.* **89**, 1303–1313
(1994).
92.   Tools for compositional data with a total - Vera Pawlowsky-Glahn, Juan José Egozcue, David
Lovell, 2015. https://journals.sagepub.com/doi/abs/10.1177/1471082X14535526.
93.   Greenacre, M. *Compositional Data Analysis in Practice.* (Chapman and Hall/CRC, New York,
2018). doi:10.1201/9780429455537.

94.	Human Microbiome Project Consortium. A framework for human microbiome research. *Nature*
486, 215–221 (2012).
95.	Lloyd-Price, J. *et al.* Strains, functions and dynamics in the expanded Human Microbiome
Project. *Nature* **550**, 61–66 (2017).
96.	Xie, H. *et al.* Shotgun Metagenomics of 250 Adult Twins Reveals Genetic and Environmental
Impacts on the Gut Microbiome. *Cell Syst.* **3**, 572-584.e3 (2016).
97.	Qin, J. *et al.* A metagenome-wide association study of gut microbiota in type 2 diabetes. *Nature*
490, 55–60 (2012).
98.	Korpela, K. *et al.* Selective maternal seeding and environment shape the human gut microbiome.
Genome Res. **28**, 561–568 (2018).
1263