Anterior cingulate cortex in complex associative learning: monitoring action state and
action content

Wenqiang Huang †[1], Arron F Hall †[1], Natalia Kawalec [1,2], Ashley N Opalka [1,3], Jun Liu [1], & Dong
V Wang *[1]

1 Department of Neurobiology & Anatomy, Drexel University College of Medicine, Philadelphia,
PA 19129.

2 School of Arts & Sciences, University of Pennsylvania, Philadelphia, PA 19104.

3 Current address: Department of Neuroscience, Chronobiology and Sleep Institute, Perelman
School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania 19104.

† Co-first authors

* Correspondence: dw657@drexel.edu

Abstract

Environmental changes necessitate adaptive responses, and thus the ability to monitor one's
actions and their connection to specific cues and outcomes is crucial for survival. The anterior
cingulate cortex (ACC) is implicated in these processes, yet its precise role in action monitoring
versus outcome tracking remains unclear. To investigate this, we developed a novel
discrimination–avoidance task for mice, designed with clear temporal separation between
actions and outcomes. Our findings show that ACC neurons primarily encode post-action
variables over extended periods, reflecting the animal's preceding actions rather than the
outcomes or values of those actions. Specifically, we identified two distinct subpopulations of
ACC neurons: one encoding the action state (whether an action was taken) and the other
encoding the action content (which action was taken). Importantly, increased post-action ACC
activity was associated with better performance in subsequent trials. These findings suggest
that the ACC supports complex associative learning through extended signaling of rich action-
relevant information, thereby bridging cue, action, and outcome associations.

Introduction

The ability to adapt behavior based on environmental cues and outcome-related feedback is essential for survival. Central to this process is the brain's capacity to integrate diverse information to form cue–action–outcome associations that guide future behaviors across varying conditions. However, the neural mechanisms underlying this cognitive flexibility remain poorly understood. The anterior cingulate cortex (ACC) has emerged as a critical node in mediating this process, with growing evidence underscoring its pivotal role in higher-order cognition and adaptive behavior [1-7]. Despite this, the precise role of the ACC in updating and modifying behavior is still under debate [1-7].

Historically, the ACC was thought to be essential for error detection or conflict monitoring [8-10]. However, accumulating evidence has challenged these views [11-17], leading to proposals that the ACC may support a diverse range of functions. These include value encoding, strategy updating, decision-making, action monitoring, and outcome tracking [18-29]. Some of these proposed functions remain controversial, such as the ACC's role in decision-making [18-20], while others are defined more broadly, such as the ACC's role in strategy updating [26, 28]. Nevertheless, there is broad agreement that ACC activity is closely linked to actions and/or action-related outcomes, particularly in tasks involving competing actions [18-29]. Supporting this view, lesions of the ACC impair the maintenance of newly acquired task performance and disrupt behavioral flexibility, including the ability to associate actions with their outcomes [27-30].

Despite evidence linking the ACC to action-related cognitive processes, past studies have primarily relied on simple motor tasks, such as saccades, licking, or joystick movements, leaving its role in more naturalistic behaviors largely unexplored. Moreover, the brief temporal separations between cues, actions, and outcomes in prior studies have made it challenging to determine whether ACC neuronal activity contributes to decision-making, action monitoring, or merely tracking the outcomes of those actions [17-31]. In this study, we aim to disentangle the ACC's role by employing a novel discrimination–avoidance task, designed to evoke robust, naturalistic action responses and establish a clear temporal separation between cues, actions, and outcomes. Our findings highlight a distinct role of the ACC in encoding post-action variables that capture detailed information about preceding actions, rather than tracking the outcomes of those actions or their associated values.

Results

A novel discrimination–avoidance task

We first developed a novel discrimination–avoidance task, tailored to investigate action-based
complex associative learning in mice. In this task, animals learn to discriminate between two
auditory cues that predict context-dependent footshocks. Specifically, sounds A and B signal
electric shocks in rooms A and B of a shuttle box at sound terminations, respectively (Fig. 1A;
Fig. S1). This design requires animals to either "stay" in the current room or "shuttle" to the
adjacent room during sound presentations to avoid shocks (Supplementary Videos 1–2). During
inter-trial intervals, animals are free to explore either room without consequence.

Notably, the auditory cues are not inherently associated with positive or negative valence;
instead, their meaning is dynamically determined by the animal's current location (room A or B)
at the time of cue presentation, signaling either safety or shock. Thus, this task, which requires
discrimination of sensory cues and environmental contexts, as well as the integration of cues,
actions, and outcomes, serves as an ideal tool to study complex associative learning. Our
results showed that mice gradually learned the task, achieving an average success rate of
76.7% in avoiding shocks by the 10th training session (Fig. 1B). An additional five training
sessions yielded a modest improvement, increasing the average success rate to 82.5% on the
15th session (Fig. S2).

Given that the task requires two distinct behavioral responses, we separated the trials into
shuttle and stay trials for further analysis. In shuttle trials, mice must move to the adjacent room
before the sound ends to avoid shocks. Across training sessions, correct shuttles increased
markedly, whereas incorrect shuttles decreased only modestly (Fig. 1C; Fig. S2). This pattern
suggests that animals choose to shuttle primarily when they are confident in the outcome (i.e.,
safety); otherwise, they remain in place. Consistent with this notion, in stay trials, where mice
must remain in their current room to avoid shocks, incorrect stays decreased markedly across
training sessions, mirroring the improvement in correct shuttle performance. By contrast, correct
stays increased only modestly (Fig. 1C; Fig. S2).

We also compared shuttle response latencies between correct and incorrect trials during the
late stages of training. On average, the shuttle response latency was 6.5 s during correct
shuttles, providing a 3.5-s temporal separation between actions (shuttles) and outcomes (safety
or shocks; Fig. 1D). This clear temporal distinction allows us to compare how information is
coded during the action *vs.* outcome periods. The longer shuttle latency during incorrect shuttles
suggests that last-second responses are more likely to be incorrect (Fig. 1D).

Figure 1. A novel discrimination–avoidance task. A, Top: Schematic of the task. Mice are trained to discriminate between two auditory cues (lasting 10 s) and shuttle between two distinct rooms of a shuttle box to avoid footshocks. Specifically, sounds A and B signal shocks in rooms A and B, respectively. Bottom: Two behavioral response scenarios. S1: If the mouse makes the correct response, either by staying in or shuttling to the correct room before the sound ends, no shock is administered. S2: If the mouse makes an incorrect response, either by staying in or shuttling to the incorrect room before the sound ends, up to 10 mild shocks (0.5 mA, 0.1 s; 2 s apart) are administered until the mouse shuttles to the correct room. Each training session comprises 50 trials, 60 s apart; sounds A and B are presented in a pseudorandom order. **B,** Learning curve showing the success rate in avoiding shocks across training sessions (mean ± s.e.m.; n = 11 mice). $F_{9, 90}$ = 14.78; P < 0.001; One-way ANOVA. **C,** Correct and incorrect trial counts for shuttle *vs.* stay trials across training sessions for the same mice shown in B. Correct shuttle: $F_{9, 90}$ = 14.62, P < 0.001; Incorrect shuttle: $F_{9, 90}$ = 1.94, P = 0.057; Correct stay: $F_{9, 90}$ = 1.55, P = 0.142; Incorrect stay: $F_{9, 90}$ = 18.73, P < 0.001; One-way ANOVA. **D,** The mean shuttle crossing latency, averaged over the last two sessions for individual mice, is significantly shorter in correct trials than that in incorrect trials (t_{10} = 2.62, P = 0.026; Effect Size: Cohen's d = 0.789; Power = 0.656; paired *t* test). The black line indicates the mean; grey lines indicate individual mice. Shuttle crossing is defined as the body center crossing the midline opening of the shuttle box.

Next, we conducted multi-channel *in vivo* electrophysiology recordings from the ACC (Fig. 2A;

Fig. S2) in mice performing the discrimination–avoidance task after they had completed 10

training sessions and achieved a success rate of 70% or above. We initially observed robust

changes in ACC activity following shuttle responses (Fig. 2 B&C). Most ACC neurons exhibited

prolonged activity changes lasting up to 30 s, either activation or suppression, primarily after

shuttle crossings (Fig. 2 C&E; Fig. S2). In contrast, the same ACC neurons showed no or limited

activity changes during stay trials in response to the auditory cues (Fig. 2 D&F).

Figure 2. ACC neurons primarily encode post-action variables. A, A representative brain section

showing electrode tracks (left) and the presumed implantation sites (red dashed lines; right), located

largely within the ACC. Scale bars, 0.5 mm. **B,** A representative shuttle response and corresponding Y-

position of the animal's body center. **C&D,** Peri-event rasters and histograms of a representative ACC

To assess ACC neuronal population activity dynamics, we performed Principal Component
Analysis (PCA) on simultaneously recorded ACC neurons across correct-shuttle and correct-
stay trials (Fig. 3A). The first three principal components (PCs) revealed pronounced changes in
population activity during shuttle trials, whereas activity remained relatively stable during stay
trials (Fig. 3B; Fig. S3). Quantifying the maximum trajectory distance within the 3D PC space
revealed consistent, prominent changes in population activity during shuttle responses across
animals in five representative sessions (Fig. 3C). The pronounced ACC activity following
shuttles, coupled with minimal activity in response to cues, suggests that ACC neurons primarily
encode post-action variables. Such sustained post-action activity may support complex
associative learning by linking temporally separated action- and outcome-relevant information.

Figure 3. ACC neurons primarily respond during "shuttle" but not "stay" trials. A, Heatmaps
showing the activity of simultaneously recorded ACC neurons (n = 29) during correct-shuttle (top) and
correct-stay trials (bottom) from the same recording session as shown in Fig. 2C. **B,** Principal component
analysis (PCA) of ACC neuronal population activity as shown in A. PC1, PC2, and PC3 are the first three
principal components; the numbers are the percentages of total variance explained by the corresponding
PCs. Each circle indicates a time lapse of 0.1 s. Note that there is a robust neural state change in the 3-D
PC space surrounding the shuttle response (top), but not the stay response (bottom). **C,** Maximum
trajectory distance between the center of the baseline period and any post–Time 0 data point, as shown
in B, across five animals (top), and the corresponding scree plots showing variance explained by the first
eight PCs during shuttle trials (bottom).

<u>ACC neurons exhibit limited pre-action firing changes</u>

To assess whether ACC activity has a role in encoding pre-action variables, we examined ACC
activity aligned to shuttle initiations, defined as locomotion velocity exceeding one standard
deviation (s.d.) above baseline (Fig. 4A). Our results revealed diverse response properties of
the ACC neuronal population (Fig. 4B). PCA classified these responses into three major
categories (Fig. 4 C&D). Pre-shuttle ramping activity was mainly observed in a small subset of
neurons, (Types 3 b&c, ~16% of the population), indicating a limited role of the ACC in pre-
action information coding, including decision-making and planning processes. The remaining
ACC neurons primarily exhibited post-shuttle responses, either activation (Types 1a, 1b, 3a) or
inhibition (Types 2a, 2b), highlighting the predominant role of the ACC in post-action information
processing.

Fig. 4. Characterizing ACC neuronal activity in relation to action initiations. A, Top, schematic
illustrating the definition of shuttle initiation, crossing, and termination (pause). Bottom, distributions of
half- (left) and full-shuttle durations (right). **B**, Z-scored activity of all ACC neurons (n = 376) during correct
shuttle trials. **C,** Principal-component analysis (PCA) classifies ACC neuronal activity (as shown in B) into
seven categories. PC1, PC2, and PC3 represent the first three principal components color coded from
low (dark) to high scores (white). **D**, Mean activity (± s.e.m.) of the seven categories of ACC neurons. **E**,
Fractions of individual categories of ACC neurons.

ACC neurons exhibit limited modulation by speed

Given that the post-action ACC activity is often prolonged and outlasts shuttle termination, we

hypothesized that this activity is distinct from locomotion encoding. To test this, we analyzed

ACC activity in relation to movement speed. Each task session included a 5-min free exploration

period before trials begin, allowing us to assess if ACC activity is associated with basic motor

functions. We found that only a small portion of neurons showed speed-correlated activity, with

either positive (7.7%) or negative correlation (6.9%), indicating that the majority of the neurons

are not tuned to movement speed (Fig. 5 A–C). This is consistent with our observation of

sustained post-shuttle ACC activity in the absence of movement, which is distinct from

locomotion encoding. Nevertheless, it remains unclear whether this small fraction of speed-

related neurons represents a distinct subpopulation within the ACC or reflects recordings from

nearby motor cortex. Postmortem examination of the recording sites suggests that most

neurons were recorded from the ACC, while a small subset located at the border between the

ACC and motor cortex (Fig. S2). Therefore, it is possible that the speed-related neurons

originated from the motor cortex.

Figure 5. ACC neurons exhibit limited modulation by speed. A, Speed-tuning curves of three
simultaneously recorded neurons during free exploration in shuttle boxes, showing positive (red),
negative (blue), or no correlation (gray) between firing rate and locomotion speed. "r" indicates Pearson's
correlation coefficient. **B,** Distributions of observed *vs.* shuffled correlation values across all recorded ACC
neurons (n = 376). Dashed lines mark the 99th percentile of shuffled distribution. Overall, 7.7% of
recorded neurons were positively modulated by speed, and 6.9% were negatively modulated. **C,**
Proportions of speed-modulation for each category of ACC neurons (see Fig. 4D).

    ACC neurons monitor actions independent of outcomes

To determine if the post-action ACC activity encodes information related to outcomes, we
analyzed ACC neuronal activity across three distinct conditions: correct shuttles, incorrect
shuttles, and post-shock shuttles (Fig. 6A; post-shock shuttles are defined as shuttles following
footshocks received during incorrect-shuttle and incorrect-stay trials). Our results revealed that
ACC neurons exhibited similar activity patterns across all three conditions, regardless of
outcomes (presumed safety *vs.* uncertainty *vs.* safety; Fig. 6B). Specifically, both activity
strength (Fig. 6C) and activity pattern (Fig. 6D) were significantly correlated across conditions,
indicating outcome-independent responses in ACC neurons. Consistently, we found that ACC
neurons showed limited responses to footshocks during incorrect trials (Fig. S4). Together,
these findings suggest that ACC neurons monitor actions independent of outcomes or values
associated with these actions.

Figure 6. ACC neurons monitor actions independent of outcomes. A, Peri-event rasters (trials) and
histograms of a representative ACC neuron during correct (left), incorrect (middle), and post-shock
shuttles (right) within a session. Red triangles indicate shock administrations. Note that incorrect shuttles
are followed by a second shuttle after animals receive footshocks, and approximately half of the post-
shock shuttles are preceded by incorrect shuttles. **B,** Heatmaps showing the activity of individual ACC
neurons (n = 348) during correct (left), incorrect (middle), and post-shock shuttles (right). Neurons are

ACC neurons monitor *action state* and *action content*

Since post-action ACC activity was outcome-independent, we next asked which other features

of the task ACC activity might differentiate. Leveraging the two distinct shuttle responses within

the task, we examined whether ACC neurons responded differently between rooms A→B

shuttles (in response to sound A) and rooms B→A shuttles (in response to sound B). Our

analyses identified two major groups of ACC neurons based on their responses to these

shuttles. The first group showed indiscriminate response, exhibiting either increased or

decreased activity without differentiating between A→B and B→A shuttles (Fig. 7: Neurons

1&2). We propose that these ACC neurons encode an *action state*, a variable representing the

change of behavioral state in response to the auditory cues.

In contrast, the second group of ACC neurons exhibited discriminate responses between rooms

A→B and B→A shuttles. These neurons either responded selectively in one set of shuttles or

show different levels of responses (activation or inhibition) between the two sets of shuttles (Fig.

7: Neurons 3&4; Fig. S5). We propose that these ACC neurons encode *action content*, a

variable representing distinct action information (i.e., shuttling from rooms A→B *vs.* B→A).

To quantify *action-state* and *action-content* neurons, we performed a generalized linear model

(GLM)-based analysis of ACC activity surrounding shuttle responses (see Methods). Based on

coefficient β or Δβ value differences, most ACC neurons were classified as either *action-content*

neurons (54.5%; Categories 1&2) or *action-state* neurons (19.1%; Categories 3&4; Fig. 7 C–E).

These findings suggest a prominent role for post-action ACC activity in encoding rich

information about preceding *action state* and *action content*.

Notably, ACC activity does not resemble place cell activity observed in the hippocampus [32], as

evidenced by our analysis of spike activity from the intertrial-interval periods (Fig. S6). This

aligns with prior findings that ACC neurons do not simply encode spatial information [33],

although *action-content* ACC neurons likely incorporate spatial variables, given their shuttle-

route response selectivity.

Figure 7. ACC neurons monitor *action state* and *action content*. A&B, Peri-event rasters (trials) & histograms of four simultaneously recorded ACC neurons during two sets of shuttles: rooms A→B shuttles (A) *vs.* rooms B→A shuttles (B). Notably, neurons 1&2 exhibit indiscriminate responses, either increasing or decreasing their activity after shuttles, thereby monitoring *action state* changes. In contrast, neurons 3&4 are selectively activated in one set of the shuttles, thereby monitoring *action content* (i.e., rooms A→B *vs.* B→A shuttles). **C,** Z-scored activity of all ACC neurons (n = 376) during A→B shuttles (left) and B→A shuttles (right). The color bar indicates z score. Neurons are categorized and sorted according to coefficient β or Δβ values (see Methods): Category 1 (n = 98), $\beta_1 > \beta_2$, sorted by β_1; Category 2 (n = 107), $\beta_1 < \beta_2$, sorted by β_2; Categories 3&4 (n = 29/43), both β_1 and β_2 are significantly positive or negative, sorted by the combined magnitude of β_1 and β_2. **D,** Coefficient values β_1, β_2, and Δβ ($\beta_1-\beta_2$) values are

To determine if the post-action ACC neuronal population activity can decode shuttle contents
(rooms A→B *vs.* B→A shuttles), we implemented a machine-learning approach. Specifically, we
trained binary support vector machine (SVM) classifiers and performed cross-validations (Fig.
8A). Our results revealed that the post-shuttle population ACC activity was highly effective in
decoding the animal's preceding actions of either A→B *vs.* B→A shuttles, reaching a decoding
accuracy close to 90% on average (Fig. 8B). Although the pre-shuttle ACC activity can also
decode the shuttle content, it did so with much lower accuracy (Fig. 8B). These findings provide
compelling evidence supporting post-action ACC activity in encoding detailed information about
the animal's preceding actions. Further analysis confirmed that model prediction accuracy was
derived primarily from *action-content* ACC neurons, as gradual removal of these neurons from
the analysis resulted in progressive decreases in prediction accuracy (Fig. 8C).

Figure 8. Post-shuttle ACC neuronal population activity decodes *action content*. C, Schematic
diagram of support vector machine (SVM) decoding. ACC neuronal population activity from pre-shuttle
period (-5–0), post-shuttle period (0–5), or shuffled spikes is used to train the decoder and subsequently
distinguish between action content (rooms A→B *vs.* B→A shuttles). **B,** Mean decoding accuracy (blue
line) and individual decoding accuracies for 15 sessions (grey lines). *Friedman test (*P < 0.001*) and post-
hoc Wilcoxon signed-rank test with Bonferroni correction* (***P < 0.001). **C**, SVM decoding accuracy
across all 15 sessions as a function of the fraction of neurons removed (20% per step), applied separately
to *action-content* neurons (blue) or the remaining neurons (grey) within each session (P < 0.001 for each
comparison between the two removals; *Wilcoxon signed-rank test*). Shaded areas denote ± s.e.m.

276 <u>Post-action ACC activity influences future performance</u>

277 We next investigated whether post-action ACC activity following a trial influenced performance
278 on the subsequent trial, which occurred ~1 min later. Specifically, we divided correct-shuttle
279 trials into two groups (Fig. 9A): those followed by correct trials (including both correct stays and

shuttles), and those followed by incorrect trials (including incorrect stays and shuttles). Our results revealed that post-shuttle ACC activity was notably higher when it preceded correct trials rather than incorrect ones (Fig. 9B). *The strength of post-shuttle ACC activity may reflect task engagement*, with greater engagement facilitating learning. Statistically, the top one third of the most responsive ACC neurons exhibit significantly higher activity that preceded correct trials than incorrect ones (Fig 9C). This correlation between higher ACC activity and future correct performance suggests that post-action ACC activity may contribute to trial-to-trial behavioral adjustment that underlie associative learning [34].

As a control, we also examined whether the status of the preceding trial influenced post-action ACC activity in the current trial. Specifically, we divided correct-shuttle trials into two groups: those preceded by correct trials, and those preceded by incorrect ones (Fig. 9D). Our results showed no difference in post-shuttle ACC activity between the two conditions (Fig. 9 E&F). This finding was expected, as both positive and negative reinforcement can similarly contribute to associative learning and neural plasticity.

Figure 9. Post-action ACC activity influences future performance within a task session. A, Schematic illustration. **B**, Mean activity (± s.e.m.) of post-shuttle activated neurons (orange lines; top 1/3), inhibited neurons (blue lines; bottom 1/3), and remaining ACC neurons (black lines; middle 1/3), which preceded either correct trials (solid lines) or incorrect trials (dashed lines). **C,** Further comparison of the activation strength for post-shuttle activated neurons (left) and inhibited neurons (right) between the two

Discussion

Our newly designed discrimination–avoidance task is unique in that it allows us to disentangle the roles of ACC neurons across several proposed functions, including value encoding, decision-making, action monitoring, and outcome tracking. First, this task requires animals to discriminate both sensory cues and environment contexts. Unlike established tasks that often assign fixed positive or negative values to cues, the cues in our task are not inherently associated with valence. Instead, their meaning is dynamically determined by the animal's location (context) at the time of cue presentation. By removing valence from the cues, this design helps disentangle the ACC's potential role in value encoding from other cognitive functions. Second, this task involves robust, ethologically relevant actions (i.e., shuttles), unlike many established paradigms that rely on less naturalistic behaviors such as saccades or lever presses. Finally, the clear temporal separation between actions and outcomes helps disentangle the ACC's roles in action monitoring *vs.* outcome tracking.

Utilizing this task, we find that the ACC primarily encodes post-action variables. Specifically, ACC neurons exhibit robust post-shuttle responses across various conditions, including correct, incorrect, and post-shock shuttles. Despite conditions signaling distinct outcomes (presumed safety *vs.* uncertainty *vs.* safety), the response properties of the ACC neurons remain consistent. Moreover, very few ACC neurons respond directly to positive outcomes (safety) or negative outcomes (shocks). Together, these findings indicate that post-shuttle ACC activity primarily monitors animal's most recent actions, rather than tracking the outcomes or values associated with those actions [21, 29, 30].

Our results further reveal two distinct groups of ACC neurons that encode different aspects of actions: *action state* and *action content*. Action state appears to represent the animal's preceding choice of whether an action was taken, updating changes in behavioral state within the ongoing task. In contrast, *action content* represents the specific actions taken (i.e., shuttles from rooms A→B *vs.* B→A), preserving detailed action information. The response selectivity of *action-content* ACC neurons reinforces the notion that ACC activity monitors preceding actions rather than action-associated outcomes or values, which are uniform across these actions. Notably, our findings do not support an alternative interpretation that *action content* reflects

correct *vs.* incorrect shuttles. Theoretically, categorizing actions solely as correct or incorrect
may not be necessary, as both positive and negative outcomes can guide future actions and
contribute to learning [30]. Potential neural networks mediating the sustained post-action ACC
activity include thalamic and retrosplenial inputs, given the well-established roles of
thalamocortical circuits in sustaining cortical activity [35, 36] and the retrosplenial cortex in
processing spatial and directional information [37-40].

Our study also reveals that ACC neurons play a limited role in encoding pre-action variables
associated with decision-making or planning, as evidenced by their minimal responses to
auditory cues and the modest activity changes prior to shuttle initiation. These findings align
with recent research, showing that ACC neurons are mainly involved in post-decisional
information processing rather than decision-making itself [18-20]. Nevertheless, substantial
evidence from other studies supports the ACC's involvement in value coding or value updating,
as reflected in its differential responses to discriminative sensory cues that precede decisions
[25, 31, 41]. One possible explanation for the discrepancy in our findings is that the sensory
cues used in our task are not intrinsically linked to specific values. Although these cues play a
crucial role in driving go (shuttle)/no-go (stay) decisions, their values are dynamic and vary
across trials. This lack of value association may explain why we observe minimal response of
ACC neurons to these cues. In contrast, the ACC activity reported in previous studies likely
reflects the stable value of the cues [31, 42, 43]. It remains to be determined if distinct ACC
subpopulations may be responsible for value encoding *vs.* action monitoring, or if the same ACC
neurons multiplex across these functions.

Our results suggest that ACC activity is not directly associated with locomotion. First, both
action-state and *action-content* neurons tend to show sustained activity even when the animals
remain immobile after completing shuttle behaviors, suggesting that their activity is not driven by
locomotion. Furthermore, *action-content* neurons are selectively engaged in only one of the two
shuttle categories, either rooms A→B or B→A shuttles. Therefore, differences in neuronal
activity are unlikely to reflect locomotor differences, given that both shuttle types involve similar
movement patterns. Finally, we show that only a small fraction of neurons (14.6%) exhibits
locomotion speed-correlated activity. Overall, these results suggest that post-action ACC activity
reflects information about preceding actions, independent of the animal's current movement.

One caveat of our study is that the discrimination–avoidance task requires weeks of training in
mice. By the time they master the task, ACC activity may reflect modified neural circuits.
Investigating ACC activity during early phase of learning, such as by introducing a new pair of

cues or contexts, could provide further insights into ACC's role in learning and cognitive
processes. Additionally, previous studies have highlighted ACC's key role in reversal learning
[44-47]. Future research examining how ACC neurons respond when task rules are reversed
could provide further insight into this function. Finally, a limitation of the current study is the lack
of evidence for the causal role of post-action ACC activity in complex associative learning.
Future investigations using closed-loop strategies to selectively disrupt ACC activity during the
post-action phase could help address this question.

374
Figure 10. Proposed model of cue–action–outcome associative learning. Top: Four distinct phases
of information coding: pre-shuttle, shuttle, post-shuttle, and outcome phases. **Bottom:** Integration of cue,
action, and outcome information occurs when all three components are concurrently available (indicated
by the oval). Notably, without the ACC, action-relevant information could be lost after action execution,
thereby disrupting cue–action–outcome associative learning.

Lastly, our findings suggest that post-action ACC activity plays a key role in shaping future
behavior. Specifically, we find that increased post-action ACC activity is linked to future
performance in subsequent trials, highlighting the ACC's role in facilitating learning and guiding
behavior. The level of post-shuttle ACC activity may reflect task engagement, with greater
engagement facilitating learning and improving future performance. This aligns with previous
research demonstrating the ACC's critical role in complex and flexible learning, including
discriminative avoidance learning [48], discriminative extinction learning [49], reversal learning
[44-47], task switching [25], trial-to-trial behavioral adaptation [34], and action–outcome
associative learning [28-30]. We speculate that both *action-state* and *action-content* ACC
neurons contribute to complex associative learning through extended signaling of action-

relevant information, thereby bridging cue, action, and outcome associations (Fig. 10).
Specifically, *action-state* ACC neurons signal whether an action was taken, while *action-content*
ACC neurons provide specific details about what the action was—the content. This action-
related information, eventually, is integrated with cue- and outcome-related variables to form
complex cue–action–outcome associations that guide future behavior (Fig. 10). Without the
ACC, action-relevant information could be lost after action execution, thereby disrupting cue–
action–outcome associative learning. Given our key finding that the ACC primarily encodes
action-related variables rather than cue- or outcome-related information, we speculate that the
integration of these complex associations takes place in other higher cognitive areas, such as
the prelimbic cortex and other medial prefrontal subregions [50-52].

Methods

Methods

Mice. Male C57BL/6 mice (Jackson Laboratory, stock #000664) were used in this study. The
mice were 8–10 weeks old at the start of discrimination–avoidance task training. All mice were
group-housed (2–4 mice per cage; 40 × 20 × 25 cm) with corn cob bedding and cotton nesting
material, except that after electrode implantation surgery, mice were singly housed. They were
maintained on a 12-h light/dark cycle with *ad libitum* access to food and water. All experimental
procedures were approved by the Institutional Animal Care and Use Committees at Drexel
University and adhered to the National Research Council's Guide for the Care and Use of
Laboratory Animals.

Sounds and shuttle box. Two 10-s auditory cues, 5-kHz tone at ~75 dB and white noise at ~65
dB, were chosen for sound discrimination. Both sounds included 50-ms shaped rise and fall
times to reduce abruptness and minimize potential startle responses. The shuttle box used in
the experiment was a square chamber measuring 25 × 25 × 32 cm, with a 36-bar shock grid
floor, illuminated by lights inside sound-attenuating cubicles (64 × 75 × 36 cm) equipped with
speakers (*Med Associates*). The shuttle box was divided at the midline by a plastic divider into
two rooms. These two rooms were slightly modified for discrimination purposes: one had two
black walls, one white wall, and one transparent wall, while the other room had two white walls,
one metal wall, and one transparent wall (Fig. S1). The divider had a 2-inch opening in the
center to allow the mice to move freely between the rooms. Animals' behaviors were recorded
using Video Freeze software (*Med Associates*) [53].

Discrimination–avoidance task. Prior to training, all mice underwent two daily handling
sessions (~10 min each). Once training began, the mice received one training session per day,

5 days per week. In the task, the mice were trained to discriminate between two distinct sounds
(A and B) and shuttle between two adjacent rooms (A and B) within a shuttle box to avoid
potential footshocks. Specifically, sounds A and B signaled electric shocks in rooms A and B,
respectively (Fig. 1A). During training, the mice were first allowed to freely explore the shuttle
box for 2 min before trials began, except on the first day of training, where the free exploration
period was extended to 10 min.

The training procedure consisted of two phases: pre-training and training. Pre-training phase:
mice underwent five daily sessions of sound alternation trials 5/5 (AAAAABBBBB …). Mice that
did not exhibit shuttling responses during the first session were excluded from further training.
Training phase: mice underwent 10 daily sessions (5 sessions per week) of alternation trials in a
pseudorandom order (ABBABAAB …).

Each training session comprised 50 trials, 60 s apart. On incorrect trials, up to 10 scrambled
electric shocks (0.5 mA; 0.1 s) were administered starting at sound terminations and continued
for an additional 18 s (1 shock every 2 s), except during the pre-training phase, where up to 20
scrambled electric shocks were administered. Shocks were terminated once the mice navigated
to the adjacent safe room. The mouse's success rate in avoiding shocks at sound terminations
was defined as: Success rate (%) = (Correct stays + Correct shuttles) / All trials.

Real-time location detection. We employed MATLAB functions to detect animal location and
subsequently control shock administration (Fig. S1). Specifically, *Med Associates* Video Freeze
software recorded real-time video footage of the animal's behaviors [53]. During each trial,
MATLAB captured screenshots of the ongoing video at sound terminations and every 2 s
thereafter during the shock period. MATLAB then performed background extraction on the
captured images to determine which room the mouse was located in. To deliver a shock,
MATLAB sent a "shock" signal to an *Arduino UNO* circuit board, which relayed that signal to
Med Associates, triggering shock administration.

Stereotaxic surgery. Mice that had completed 10 training sessions and surpassed a success
rate of 70% during discrimination–avoidance tasks were used for surgery. In brief, mice were
anesthetized with ketamine/xylazine mixture (~100/10 mg/kg, i.p.) and maintained on a heating
pad at 37°C. Following that, mice received an intra-ACC implantation of a custom-made
electrode array (8 tetrodes) [54, 55], and the implant was secured to the skull with stainless
screws and resin ionomer (*DenMat*). The coordinates used were AP 1.0 mm, ML 0.4 mm, and
DV 1.0 mm.

***In vivo* recording during discrimination–avoidance tasks.** We used tetrodes for recording
[54, 55]. Each tetrode consisted of four wires (90% platinum 10% iridium; ~18 μm diameter;
California Fine Wire). Neural signals were preamplified, digitized, and recorded using a
Blackrock Neurotech CerePlex, while the animals' behaviors were simultaneously recorded.
Spikes were digitized at 30 kHz and filtered between 600–6000 Hz. The recorded spikes were
manually sorted using *Plexon* Offline Sorter [56, 57], with key datasets verified by a second
experimenter. Tetrode arrays were gradually lowered through a microdrive [58, 59] to record at
multiple depths within the ACC, with 2–4 depths from each animal (DV ~1.0–1.3) used for
analysis.

Overall, ACC spikes from five mice across 15 sessions were analyzed, with neuron counts of
20/23/29/32 (mouse #1; 4 sessions), 17/20/22 (mouse #2; 3 sessions), 28/24 (mouse #3; 2
sessions), 33/25/21/27 (mouse #4; 4 sessions), and 22/25 (mouse #5; 2 sessions), respectively.
The total number of correct/incorrect shuttles used for analysis are 19/5, 19/4, 21/5, 20/4
(mouse #1); 20/7, 23/7, 20/7 (mouse #2); 19/4, 16/2 (mouse #3); 26/4, 23/4, 17/6, 25/5 (mouse
#4); 20/5, and 17/4 (mouse #5), respectively. For Figs. 2–5 & 7–8, all recorded neurons (n =
376) from the 15 sessions were included in the analyses. For Fig. 6, however, one or two
sessions were excluded because they contained too few trials of certain types (n ≤ 3).

The discrimination–avoidance task was similar to that during the training phase, except that five
direct-current electric shocks, which minimize electromagnetic artifacts, were administered
starting at sound terminations and continued for 8 s. This shorter shock period allowed the mice
more time to move freely within the shuttle box during the 42-s inter-trial intervals, when no
consequences were administered. Additionally, each task session included a 5-min free
exploration period before trials began. In some sessions, a small number of trials were excluded
from analysis, partly due to the electrode implant or recording cable occasionally interfering with
the animal's shuttle responses.

Shuttle behavior analysis. We used DeepLabCut [60] to analyze animals' locations during
discrimination–avoidance tasks, with all locations determined based on body center positions.
Shuttle crossing was defined as the body center crossing the midline opening of the shuttle box.
Shuttle initiations and terminations were defined as the time points when the animal's
movement velocity deviated 1 s.d. above mean (Fig. 4A).

Dimension reduction. We used Principal Component Analysis (PCA) to analyze the major
activity patterns of ACC neuronal populations. The first three principal components (PCs) were
used for 3-D visualizations. Specifically, the activity of each ACC neuron was first averaged

during shuttle responses (bin size, 0.1 s) and z scored. The z-scored activity of ACC neurons
was then processed with PCA (*pca*, MATLAB).

Speed tuning analysis and speed cell classification. Speed tuning was assessed using
neural and behavioral data collected during a ~5-min period of free movement in the shuttle box
prior to task onset. During this period, animals freely explored the environment and exhibited a
broad range of instantaneous running speeds. Running speed was extracted from behavioral
tracking data and sampled at 50 Hz. Spike trains from individual neurons were binned at 20 ms
resolution and converted to firing rates. Firing-rate time series were smoothed using a Gaussian
kernel with a 250 ms window. To reduce potential confounds associated with immobility-related
network states, time points with running speed below 2 cm/s were excluded from the analysis,
following established procedures [61]. For each neuron, a speed score was defined as the
Pearson correlation coefficient between the smoothed firing rate and the valid portion of the
instantaneous running speed.

Statistical significance was assessed using a circular time-shift shuffle procedure. For each
neuron, spike times were circularly shifted by a random offset greater than 30 s, preserving the
temporal structure of firing while disrupting its alignment with behavior. The firing rate–speed
correlation was recomputed for each shuffle, and this procedure was repeated 100 times to
generate a null distribution of speed scores. Neurons were classified as speed-modulated if
their observed speed score exceeded the 99th percentile or fell below the 1st percentile of the
shuffle distribution (two-sided criterion). All analysis procedures were adapted from previous
studies of speed-modulated neurons [61].

PCA-based classification of ACC neuronal types. We used PCA to classify major types of
ACC neurons based on their activity in reference to shuttle initiation. Specifically, the activity of
each ACC neuron was first averaged surrounding shuttle responses (-5–5 s; bin size, 25 ms)
and z-scored. The first three PCs were used in a hierarchical clustering algorithm (Linkage) to
find the similarity (Euclidean distance) between all pairs of activity patterns in PC space,
iteratively grouping the activity patterns into larger and larger clusters based on their similarity.
Lastly, we set a distance-criterion to extract major clusters from the hierarchical tree [62].

GLM-based classification of 'shuttle-content' and 'shuttle-state' neurons. For each neuron,
we computed peri-event firing rates using 25-ms bins and normalized activity relative to a
baseline window from −30 to −20 s preceding shuttle onset. Post-shuttle activity (0–5 s) was

analyzed using a generalized linear model (GLM) with separate regressors for each event.

Specifically, for Event1 (shuttle A→B) and Event2 (shuttle B→A) trials, we fit the model

$$r(t) = \beta1 \cdot Event1(t) + \beta2 \cdot Event2(t),$$

using a normal distribution with an identity link (*glmfit*, constant = 'off', MATLAB). We estimated

coefficients $\beta1$ and $\beta2$ (corresponding to shuttle A→B and shuttle B→A, respectively) and

assessed their significance using Wald tests against zero. To quantify event selectivity, we

tested the contrast: $\Delta\beta = \beta1–\beta2$. Resulting p-values were corrected for multiple comparisons

using Bonferroni correction across regressors and neurons ($\alpha\_bonf = 0.01/(3N)$, where N is the

number of neurons in a session). Neurons were classified as *action-content* neurons if the

corrected p-value for $\Delta\beta$ was significant and the absolute effect size exceeded a predefined

threshold ($|\Delta\beta| > 0.5$). Neurons were classified as *action-state* neurons if $\Delta\beta$ was not significant

but both $\beta1$ and $\beta2$ were individually significant after correction. All analysis procedures were

adapted from previous studies [63-65].

Machine learning decoding. We used Support Vector Machine (SVM) classifiers to train ACC

neuronal population activity to decode shuttle content (rooms A→B *vs.* B→A shuttles) and

subsequently performed 10-fold cross-validations. Specifically, the total number of spikes from

each ACC neuron, calculated during either the pre-shuttle (-5–0 s) or post-shuttle period (0–5 s),

were used for training and testing the SVM classifier. For each dataset, we performed SVM

classification training and cross-validation 100 times (*fitcsvm* and *crossval*, MATLAB), assigning

the mean correct classification rate as the decoding accuracy. Notably, using shorter or longer

time windows surrounding shuttle responses, such as 2.5 or 10 s, yielded similar conclusions

(Fig. S7). Additionally, ACC neuronal population activity moderately decoded Room-A *vs.* Room-

B stays (Fig. S7; SVM training and cross-validation repeated 500 times).

To assess the contribution of *action-content* neurons to decoding performance, neurons were

progressively removed in 20% increments (Fig. 8C). Within each session, *action-content*

neurons were ranked by $|\Delta\beta|$ in descending order, and the top fraction was removed at each

step prior to re-estimating decoding accuracy. As a control, the same number of neurons was

randomly removed from the $\Delta\beta$ non-significant pool. At each removal fraction, decoding

accuracies obtained after *action-content* neuron removal were compared to control accuracies

across sessions using a paired, two-sided, Wilcoxon signed-rank test.

Spatial information analysis (Fig. S6). To quantify spatial information coding, firing rate maps

for individual neurons were constructed using 1 × 1 cm spatial bins in NeuroExplorer and

smoothed with a Gaussian filter (filter width, 3 bins) before being exported to MATLAB for
further analyses. Spatial information for each neuron was calculated using the following formula:

$$Spatial\ Information\ (bits/spike) = \sum_i p_i \frac{\lambda_i}{\lambda} \log_2 \frac{\lambda_i}{\lambda},$$

where λ_i is the mean firing rate in the i-th bin, λ is the overall mean firing rate, and p_i is the
probability of the animal's being in the i-th bin, as described previously [55, 66].

Histology. To mark the final recording sites, we made electrical lesions by passing 10-s, 10-μA
currents through multiple tetrodes. Mice were deeply anesthetized and intracardially perfused
with ice-cold PBS or saline, followed by 10% formalin. The brains were removed and postfixed
in formalin for at least 24 hours. The brains were sliced into coronal sections of 50-μm thickness
using *Leica* vibratome. Brain sections were mounted with Mowiol mounting medium for
microscopic examination of electrode array placements.

Statistics. Sample sizes were based on previous similar studies [55, 62]. All statistics were
conducted in SPSS 30.0. Statistical analyses include repeated measures ANOVA, the
nonparametric Friedman test by post-hoc test (pair-wise Wilcoxon signed-rank test and
Bonferroni correction), the Wilcoxon signed-rank test (paired), and Student's *t* test (paired). All
statistical tests are two-sided; P-values of 0.05 or lower were considered significant.

Financial Disclosures

The authors declare no financial interests or potential conflicts of interest.

Acknowledgements

This work was supported by the National Institutes of Health grants R01MH119102 (D.V.W.),
R21MH134016 (D.V.W.), and F31MH134582 (A.F.H.).

References

1.   Heilbronner, S.R. and B.Y. Hayden, *Dorsal Anterior Cingulate Cortex: A Bottom-Up View.*
Annu Rev Neurosci, 2016. **39**: p. 149-70.
2.   Shenhav, A., J.D. Cohen, and M.M. Botvinick, *Dorsal anterior cingulate cortex and the*
value of control. Nat Neurosci, 2016. **19**(10): p. 1286-91.
3.   Shenhav, A., M.M. Botvinick, and J.D. Cohen, *The expected value of control: an*
integrative theory of anterior cingulate cortex function. Neuron, 2013. **79**(2): p. 217-40.
4.   Kolling, N., et al., *Multiple signals in anterior cingulate cortex.* Curr Opin Neurobiol, 2016.
37: p. 36-43.

5.   Clairis, N. and A. Lopez-Persem, *Debates on the dorsomedial prefrontal/dorsal anterior*
cingulate cortex: insights for future research. Brain, 2023. **146**(12): p. 4826-4844.

6.   Botvinick, M.M., *Conflict monitoring and decision making: reconciling two perspectives*
on anterior cingulate function. Cogn Affect Behav Neurosci, 2007. **7**(4): p. 356-66.

7.   Holroyd, C.B. and N. Yeung, *Motivation of extended behaviors by anterior cingulate*
cortex. Trends Cogn Sci, 2012. **16**(2): p. 122-8.

8.   Holroyd, C.B. and M.G.H. Coles, *The neural basis of human error processing:*
reinforcement learning, dopamine, and the error-related negativity. Psychol Rev, 2002.
109(4): p. 679-709.

9.   Botvinick, M.M., J.D. Cohen, and C.S. Carter, *Conflict monitoring and anterior cingulate*
cortex: an update. Trends Cogn Sci, 2004. **8**(12): p. 539-46.

10.   Botvinick, M., et al., *Conflict monitoring versus selection-for-action in anterior cingulate*
cortex. Nature, 1999. **402**(6758): p. 179-81.

11.   Brown, J.W. and T.S. Braver, *Learned predictions of error likelihood in the anterior*
cingulate cortex. Science, 2005. **307**(5712): p. 1118-21.

12.   Carter, C.S., et al., *Anterior cingulate cortex, error detection, and the online monitoring of*
performance. Science, 1998. **280**(5364): p. 747-9.

13.   Hayden, B.Y., et al., *Surprise signals in anterior cingulate cortex: neuronal encoding of*
unsigned reward prediction errors driving adjustment in behavior. J Neurosci, 2011.
31(11): p. 4178-87.

14.   Nakamura, K., M.R. Roesch, and C.R. Olson, *Neuronal activity in macaque SEF and*
ACC during performance of tasks involving conflict. J Neurophysiol, 2005. **93**(2): p. 884-
908.

15.   Amiez, C., J.P. Joseph, and E. Procyk, *Anterior cingulate error-related activity is*
modulated by predicted reward. Eur J Neurosci, 2005. **21**(12): p. 3447-52.

16.   Amiez, C., J.P. Joseph, and E. Procyk, *Reward encoding in the monkey anterior*
cingulate cortex. Cereb Cortex, 2006. **16**(7): p. 1040-55.

17.   Ito, S., et al., *Performance monitoring by the anterior cingulate cortex during saccade*
countermanding. Science, 2003. **302**(5642): p. 120-2.

18.   Blanchard, T.C. and B.Y. Hayden, *Neurons in dorsal anterior cingulate cortex signal*
postdecisional variables in a foraging task. J Neurosci, 2014. **34**(2): p. 646-55.

19.   Li, Y.S., et al., *Individual Neurons in the Cingulate Cortex Encode Action Monitoring, Not*
Selection, during Adaptive Decision-Making. J Neurosci, 2019. **39**(34): p. 6668-6683.

20.   Cai, X. and C. Padoa-Schioppa, *Neuronal activity in dorsal anterior cingulate cortex*
during economic choices under variable action costs. Elife, 2021. **10**.

21.   Quilodran, R., M. Rothe, and E. Procyk, *Behavioral shifts and action valuation in the*
anterior cingulate cortex. Neuron, 2008. **57**(2): p. 314-25.

22.   Azab, H. and B.Y. Hayden, *Correlates of decisional dynamics in the dorsal anterior*
cingulate cortex. PLoS Biol, 2017. **15**(11): p. e2003091.

23.   Rushworth, M.F., et al., *Frontal cortex and reward-guided learning and decision-making.*
Neuron, 2011. **70**(6): p. 1054-69.

24.   Hadland, K.A., et al., *The anterior cingulate and reward-guided selection of actions.* J
Neurophysiol, 2003. **89**(2): p. 1161-4.

25.   Cole, N., et al., *Prediction-error signals in anterior cingulate cortex drive task-switching.*
Nat Commun, 2024. **15**(1): p. 7088.

26.   Tervo, D.G.R., et al., *The anterior cingulate cortex directs exploration of alternative*
strategies. Neuron, 2021. **109**(11): p. 1876-1887 e6.

27.   Brockett, A.T., et al., *Anterior cingulate cortex is necessary for adaptation of action plans.*
Proc Natl Acad Sci U S A, 2020. **117**(11): p. 6196-6204.

28.   Akam, T., et al., *The Anterior Cingulate Cortex Predicts Future States to Mediate Model-*
Based Action Selection. Neuron, 2021. **109**(1): p. 149-163 e7.

29. Rudebeck, P.H., et al., *Frontal cortex subregions play distinct roles in choices between actions and stimuli.* J Neurosci, 2008. **28**(51): p. 13775-85.

30. Kennerley, S.W., et al., *Optimal decision making and the anterior cingulate cortex.* Nat Neurosci, 2006. **9**(7): p. 940-7.

31. Chen, W., et al., *Anterior cingulate cortex provides the neural substrates for feedback-driven iteration of decision and value representation.* Nat Commun, 2024. **15**(1): p. 6020.

32. Moser, E.I., E. Kropff, and M.B. Moser, *Place cells, grid cells, and the brain's spatial representation system.* Annu Rev Neurosci, 2008. **31**: p. 69-89.

33. Kennerley, S.W. and J.D. Wallis, *Encoding of reward and space during a working memory task in the orbitofrontal cortex and anterior cingulate sulcus.* J Neurophysiol, 2009. **102**(6): p. 3352-64.

34. Sheth, S.A., et al., *Human dorsal anterior cingulate cortex neurons mediate ongoing behavioural adaptation.* Nature, 2012. **488**(7410): p. 218-21.

35. Bolkan, S.S., et al., *Thalamic projections sustain prefrontal activity during working memory maintenance.* Nat Neurosci, 2017. **20**(7): p. 987-996.

36. Schmitt, L.I., et al., *Thalamic amplification of cortical connectivity sustains attentional control.* Nature, 2017. **545**(7653): p. 219-223.

37. Opalka, A.N. and D.V. Wang, *Hippocampal efferents to retrosplenial cortex and lateral septum are required for memory acquisition.* Learn Mem, 2020. **27**(8): p. 310-318.

38. Mao, D., et al., *Sparse orthogonal population representation of spatial context in the retrosplenial cortex.* Nat Commun, 2017. **8**(1): p. 243.

39. Alexander, A.S., et al., *Egocentric boundary vector tuning of the retrosplenial cortex.* Sci Adv, 2020. **6**(8): p. eaaz2322.

40. Alexander, A.S. and D.A. Nitz, *Retrosplenial cortex maps the conjunction of internal and external spaces.* Nat Neurosci, 2015. **18**(8): p. 1143-51.

41. Wu, K., et al., *Distinct circuits in anterior cingulate cortex encode safety assessment and mediate flexibility of fear reactions.* Neuron, 2023. **111**(22): p. 3650-3667 e6.

42. Kane, G.A., et al., *Rat Anterior Cingulate Cortex Continuously Signals Decision Variables in a Patch Foraging Task.* J Neurosci, 2022. **42**(29): p. 5730-5744.

43. Vertechi, P., et al., *Inference-Based Decisions in a Hidden State Foraging Task: Differential Contributions of Prefrontal Cortical Areas.* Neuron, 2020. **106**(1): p. 166-176 e6.

44. Rushworth, M.F., et al., *Role of the human medial frontal cortex in task switching: a combined fMRI and TMS study.* J Neurophysiol, 2002. **87**(5): p. 2577-92.

45. Johnston, K., et al., *Top-down control-signal dynamics in anterior cingulate and prefrontal cortex neurons following task switching.* Neuron, 2007. **53**(3): p. 453-62.

46. Meunier, M., R. Jaffard, and C. Destrade, *Differential involvement of anterior and posterior cingulate cortices in spatial discriminative learning in a T-maze in mice.* Behav Brain Res, 1991. **44**(2): p. 133-43.

47. Chudasama, Y., et al., *The role of the anterior cingulate cortex in choices based on reward value and reward contingency.* Cereb Cortex, 2013. **23**(12): p. 2884-98.

48. Gabriel, M., et al., *Effects of cingulate cortical lesions on avoidance learning and training-induced unit activity in rabbits.* Exp Brain Res, 1991. **86**(3): p. 585-600.

49. Bussey, T.J., et al., *Dissociable effects of anterior and posterior cingulate cortex lesions on the acquisition of a conditional visual discrimination: facilitation of early learning vs. impairment of late learning.* Behav Brain Res, 1996. **82**(1): p. 45-56.

50. Corbit, L.H. and B.W. Balleine, *The role of prelimbic cortex in instrumental conditioning.* Behav Brain Res, 2003. **146**(1-2): p. 145-57.

51. Klein-Flugge, M.C., A. Bongioanni, and M.F.S. Rushworth, *Medial and orbital frontal cortex in decision-making and flexible behavior.* Neuron, 2022. **110**(17): p. 2743-2770.

680   52.   Asaad, W.F., G. Rainer, and E.K. Miller, *Neural activity in the primate prefrontal cortex*
681   *during associative learning.* Neuron, 1998. **21**(6): p. 1399-407.
682   53.   Anagnostaras, S.G., et al., *Automated assessment of pavlovian conditioned freezing and*
683   *shock reactivity in mice using the video freeze system.* Front Behav Neurosci, 2010. **4**.
684   54.   Lin, L., et al., *Large-scale neural ensemble recording in the brains of freely behaving*
685   *mice.* J Neurosci Methods, 2006. **155**(1): p. 28-38.
686   55.   Liu, J., A.F. Hall, and D.V. Wang, *Emerging many-to-one weighted mapping in*
687   *hippocampus-amygdala network underlies memory formation.* Nat Commun, 2024.
688   **15**(1): p. 9248.
689   56.   Wang, D.V. and J.Z. Tsien, *Convergent processing of both positive and negative*
690   *motivational signals by the VTA dopamine neuronal populations.* PLoS One, 2011. **6**(2):
691   p. e17047.
692   57.   Wang, D.V., et al., *Neurons in the amygdala with response-selectivity for anxiety in two*
693   *ethologically based tests.* PLoS One, 2011. **6**(4): p. e18739.
694   58.   Wang, D.V., et al., *Mesopontine median raphe regulates hippocampal ripple oscillation*
695   *and memory consolidation.* Nat Neurosci, 2015. **18**(5): p. 728-35.
696   59.   Opalka, A.N., et al., *Hippocampal Ripple Coordinates Retrosplenial Inhibitory Neurons*
697   *during Slow-Wave Sleep.* Cell Rep, 2020. **30**(2): p. 432-441 e3.
698   60.   Mathis, A., et al., *DeepLabCut: markerless pose estimation of user-defined body parts*
699   *with deep learning.* Nat Neurosci, 2018. **21**(9): p. 1281-1289.
700   61.   Kropff, E., et al., *Speed cells in the medial entorhinal cortex.* Nature, 2015. **523**(7561): p.
701   419-24.
702   62.   Liu, J., L. Lin, and D.V. Wang, *Representation of Fear of Heights by Basolateral*
703   *Amygdala Neurons.* J Neurosci, 2021. **41**(5): p. 1080-1091.
704   63.   Truccolo, W., et al., *A point process framework for relating neural spiking activity to*
705   *spiking history, neural ensemble, and extrinsic covariate effects.* J Neurophysiol, 2005.
706   **93**(2): p. 1074-89.
707   64.   Vaccari, F.E., et al., *A Poisson generalized linear model application to disentangle the*
708   *effects of various parameters on neurophysiological discharges.* STAR Protoc, 2021.
709   **2**(2): p. 100413.
710   65.   Peer, N.D., H.G. Yamin, and D. Cohen, *Multidimensional encoding of movement and*
711   *contextual variables by rat globus pallidus neurons during a novel environment exposure*
712   *task.* iScience, 2022. **25**(9): p. 105024.
713   66.   Skaggs, W.E., et al., *Theta phase precession in hippocampal neuronal populations and*
714   *the compression of temporal sequences.* Hippocampus, 1996. **6**(2): p. 149-72.

Fig. S1. Experimental setup. A, The shuttle box used for behavioral training. Room A is configured with two black walls (left and right), one white wall (back), and a transparent front wall for video recording purposes. Room B is configured with two white walls (left and back), one metal wall (right), and a transparent front wall. **B,** Schematic diagram of the control setup utilizing MATLAB functions, with numbers indicating the sequence of control flow. **C,** A comprehensive flowchart illustrating the control setup as shown in B. **D,** Left, sounds A and B signal shocks in the bottom and top rooms of the shuttle box, respectively. Right, the Y position of a well-trained mouse in a ~40-min session.

Fig. S2. Extended training recording sites. A, Correct and incorrect trial counts for shuttle *vs.* stay trials
across 15 training sessions. In shuttle trials, mice must move to the adjacent room before the sound ends
to avoid shocks, whereas in stay trials, mice must remain in their current room to avoid shocks. **B,**
reconstructed recording sites for individual mice. **C,** Heatmaps showing the activity of all recorded ACC
neurons during correct-shuttle trials across individual mice.

Fig. S3. ACC neurons primarily respond during "shuttle" but not "stay" trials. A&B, Dimension
reduction analysis (i.e., PCA) indicates robust changes in ACC neuronal population activity during correct
shuttles (A) but not correct stays (B) from four representative recording sessions. Time "0" indicates
shuttle crossings (A) or sound onsets (B), respectively. For more details, see Fig. 3B.

735
736
Fig. S4. ACC shows limited response to cue, stay trials, and footshocks. Left, Heatmap showing the
activity of individual ACC neurons (n = 348) in relation to auditory cue onset. **Middle,** Heatmap showing
the activity of individual ACC neurons (n = 348) during stay trials. **C,** Heatmap showing the activity of
individual ACC neurons (n = 336) during footshock. Note, footshock trials with a shuttle response within 1
s shock onset were excluded to avoid shuttle response confound. As a result, some behavioral sessions
were excluded due to insufficient trial numbers.

Fig. S5. ACC pyramidal neurons and interneurons both monitor *action content*. A, Spike waveforms
(mean ± s.d.) of two representative ACC neurons: one putative pyramidal neuron and one interneuron. **B,**
Peri-event rasters (trials) & histograms of the same two ACC neurons surrounding shuttle responses.
Both neurons exhibit differential activity changes that discriminate between rooms A→B (top panels) *vs.*
B→A shuttles (bottom panels). **C,** Cross-correlation histograms between the putative interneuron (Neuron
#2; the same as shown in A) and three other pyramidal neurons (Neurons #1, #3, and #4). Neuron #4
appears to excite Neuron #2, which in turn inhibits Neuron #3, as indicated by short-latency (~2 ms)
excitatory or inhibitory interactions. The four ACC neurons were recorded simultaneously.

Fig. S6. ACC neurons do not display place cell characteristics. A, A representative navigation path during inter-trial intervals of a discrimination–avoidance task session. **B,** Place field activity of two representative ACC neurons. Both neurons show spiking activity across the chamber without place preference. The color bar indicates normalized firing rate. **C,** Spatial information coding distribution of all recorded ACC neurons (n = 376). **D–F,** Same as in A–C, but for neurons recorded from hippocampal dorsal CA1 (n = 161). Notably, more than half of CA1 neurons exhibit high information coding (>2 bits/spike; F), whereas very few ACC neurons show comparable levels of information coding (C).

Fig. S7. SVM decoding of multiple behaviors from ACC neuronal population activity. A, Mean (blue line) and individual session decoding accuracies (grey lines; 15 sessions) for decoding A→B *vs.* B→A shuttles using short (left, 2.5 s), medium (middle, 7.5 s), or long decoding windows (right, 10 s) surrounding shuttle crossings. **P < 0.01, ***P < 0.001, *Wilcoxon signed-rank test*. **B**, Mean (blue line) and individual session decoding accuracies (grey lines) for decoding correct *vs.* incorrect stays (left) or Room-A *vs.* Room-B stays (right), using a 10-s window after sound onset. Notably, in several sessions, lower-than-chance decoding accuracy for Corr/Incorr stays (colored dots; left) corresponded to high decoding accuracy for Room A/B stays (shown in the same colors; right), suggesting a potential confound between these variables.