Begging costs rather than food received cause brood size effect on growth in zebra finches

Running title: Begging costs drive brood size effect on zebra finch growth

Marianthi Tangili[1*], Michael Briga[2] & Simon Verhulst[1*]

[1] Groningen Institute for Evolutionary Life Sciences, University of Groningen, Groningen,
The Netherlands

[2] Department of Biology, University of Turku, Finland

Abstract

Altricial species rely on parental provisioning for early-life sustenance, and a larger brood
size leads to higher levels of competition between siblings for food provided by the
parents. Early-life stress can have severe and lifelong effects on Darwinian fitness. Indeed,
it is well established that being reared in a larger brood impairs growth and fitness
prospects of birds, but the mechanistic underpinnings of this effect are still largely
unknown. Specifically, it is not well known to what extent the reduced growth and fitness
prospects of nestlings reared in large broods is due to increased resource allocation to
competition versus a *per capita* reduction in parental provisioning rate, or a combination
of the two. We cross-fostered zebra finch (*Taeniopygia guttata*) chicks into small and large
broods, and recorded their growth as well as the behavior of parents and offspring
throughout the nestling period. As in previous experiments, growth rate was higher in
small broods. In large broods, chicks begged more and parents invested more time feeding
which more than compensated for the difference in brood size. We therefore conclude that
the lower growth rate for nestlings raised in large broods must be attributed to increased
energy expenditure on begging rather than a reduction in the amount of food received.
These results suggest significant energetic costs associated with begging and raise the
interesting possibility that brood size would not have negatively affected growth in large
broods if chicks had not increased their begging effort due to increased levels of
competition in nest.

Keywords: begging, parental provisioning, growth rate, cross-fostering, parent-offspring
interaction

Introduction

Early-life stress can have profound and long-lasting effects on the Darwinian fitness of individuals. Animals exposed to adversity during the developmental stages of their lives can exhibit reduced reproductive success (van de Pol et al., 2006) and lifespan (Briga et al., 2017; Tung et al., 2016), altered rates of senescence (Nussey et al., 2007) and differential social behavior (Gerritsma, 2025; Goldenberg & Wittemyer, 2018; Patterson et al., 2022). In birds, early-life stress has been found to affect foraging behavior and mass regulation (Andrews et al., 2015), attractiveness to mates (Holveck & Riebel, 2010; Mainwaring et al., 2012), telomere shortening rates (Boonekamp et al., 2014; Nettle et al., 2015) as well as inflammation (Nettle et al., 2017). Early-life challenges can have direct effects on Darwinian fitness by affecting growth (de Kogel, 1997; Nilsson & Svensson, 2001), recruitment (Schwagmeyer & Mock, 2008), nestling survival (Mock et al., 2009), foraging behavior (Andrews et al., 2015) and reproductive success (Spagopoulou et al., 2020).

In altricial avian species, for whom early-life sustenance is reliant on parental provisioning, intricate family dynamics emerge among individual chicks, their siblings, and their parents (e.g. Davies, 1976; Wright & Leonard, 2002). An increased brood size presents numerous challenges for parents and offspring. For parents, having to care for more offspring increases care demands, forcing parents to either reallocate resources that would otherwise be used for their own somatic maintenance (compensation hypothesis) or increase their own energy intake (increased-intake hypothesis) (Nilsson, 2002). For nestlings, being raised in larger broods results in higher rates of sibling competition for parental resources (Godfray, 1995; Godfray & Parker, 1992; Grodzinski & Johnstone, 2012) and the prioritization of investment in the development of structures that enhance their competitive ability, such as their gapes (Gil et al., 2008). Ultimately, being raised in a larger brood can not only lead to reduced growth rates and nestling mortality but also have long-term effects on fitness prospects in adulthood (Boonekamp et al., 2014, 2020; Briga et al., 2017; Dijkstra et al., 1990; Hõrak, 2003; Nicolaus et al., 2009).

Begging is the main behavior displayed by the chicks in the nest to attract the parents' attention and compete for parental resources with siblings (Godfray & Johnstone, 2000). Begging consists of intense and conspicuous vocalizations as well as head and body movements (e.g. wing flapping, gaping, Kilner & Johnstone, 1997; Zann, 1996). Theoretical work on begging behavior often relies on the assumption that begging entails a cost, either physiological as increased energy expenditure or by increasing predation risk. Experimental studies have confirmed that nestling begging calls can increase the risk of predation (Haskell, 1994, 1999; Leech & Leonard, 1997). The energetic costs of begging appear modest (Leech & Leonard, 1996; Mccarty, 1996), but even the observed energetic costs may constitute a substantial part of the energy that can be otherwise allocated to growth (Verhulst & Wiersma, 1997) as confirmed by experiments that manipulated begging effort independent from the amount of food received (Nettle et al., 2017; Rodriguez-Girones et al., 2001; Soler et al., 2014).

Given that begging is a possibly dangerous and energetically costly behavior, theory has been developed to understand how this behavior has evolved. Central in theoretical considerations is that there is an inherent evolutionary conflict between parents and

chicks: due to divergent genetic interests, the optimal level of parental care from the
perspective of the offspring is usually higher than the optimal level from the perspective
of the parents, hence there is a "parent-offspring conflict" (Trivers, 1974). In this context,
begging may serve as an honest signal of need, enabling parents to make informed
decisions on the allocation of resources to the brood and between nestlings (Godfray,
1991; Royle et al., 2002). Offspring need, and possibly thereby the optimal begging effort,
depends on brood size, because brood size manipulation typically depresses nestling
growth (Burness et al., 2000; de Kogel, 1997). At the same time, competition for food is
likely to increase with increasing brood size unless parents increase provisioning rate
sufficiently to fully match the demands of an increased brood size. Both processes are
likely to increase 'nestling need', and hence begging effort (Harper, 1986).

In zebra finches (*Taeniopygia guttata*), it is well established that being raised in larger
broods has multiple phenotypic effects that reduce fitness prospects, possibly through
reduced growth (Briga et al., 2017; de Kogel, 1997; Griffith & Buchanan, 2010). These
effects include reduced immunocompetence (Naguib et al., 2004), higher standard
metabolic rate (Verhulst et al., 2006, but see Briga & Verhulst, 2021), and altered mating
preferences (Holveck & Riebel, 2010). Our aim was to determine whether diminished
growth results from increased resource allocation to competition, a *per capita* reduction
in parental provisioning rate, or a combination of the two. We therefore cross-fostered
zebra finch chicks into broods of either two or six young and filmed the manipulated
broods to record the behavior of parents and offspring. We envisaged different scenarios
that could emerge, depending on responses of parents and offspring to the brood size
manipulation. If provisioning behavior and begging effort remain unchanged, the
resulting reduction in *per capita* provisioning would depress growth. If parents maintain
the same *per capita* provisioning rate, while the offspring increase their begging effort,
growth in large broods would be reduced due to increased resource allocation to begging.
Intermediate responses of both parents and offspring would lead to reduced growth due
to a combination of reduced *per capita* provisioning and increased resource allocation to
begging.

Methods

Cross fostering

Zebra finch pairs were randomly matched and placed in breeding cages (L x H x D: 80 x 40
x 40 cm) on a 14L:10D schedule at ∼ 25°C temperature and ∼60% humidity. In the cages,
there were nestboxes, nesting material (hay) as well as food and water *ad libitum*. Up until
the hatching of the first egg in the nest, egg food was also supplied to the parents.
Nestboxes were checked daily for the presence of eggs or chicks. Chicks were cross-
fostered until the maximum age of five days to broods of either two (n=21) or six (n=13)
chicks, which matches the brood size range observed in the wild (Zann, 1996). On the day
of cross-fostering, all chicks were weighed and marked by clipping one or more of the head
tufts for individual recognition. No siblings were placed in the same nests and no parents
were taking care of their own offspring. Chicks were ringed at the age of 12 days, and
weighed again at 15 and 35 days. At 35 days, having reached nutritional independence
(Zann, 1996), chicks were moved from the breeding cage into larger indoor aviaries with
two pairs of foster parents for sexual imprinting.

Extra-pair fertilizations are rare in natural zebra finch populations (Griffith et al., 2010;
Zann, 1996), and we assume therefore that as a rule the species has evolved to perceive
other brood members as siblings. This is relevant insofar that potential effects of variation
in relatedness within the families on interactions between brood members and parents
can be ignored in the context of this study.

Camera observations

Cameras were placed above the nest an hour before the start of the recording, to allow the
birds to adjust to them. Each observation consisted of an hour of filming per nest and each
nest was recorded multiple times at multiple ages of the chicks until they reached
nutritional independence. The video recordings were then uploaded and analyzed in
BORIS, an interactive software program for behavioral observations (Friard & Gamba,
2016). Frequency and duration of the following behaviors of chicks and parents was
recorded:

• *Resting*, the individual rests and does nothing else or the adult is just sitting in the nest;
• *Begging*, the chick makes mouth and body movements to extort parental resources;
• *Feeding*, the chick takes food from the parent or the parent gives food to the chick
(foraging time not included for the parents).
• *Cleaning*, the individual (either chick or parent) cleans its feathers;
• *Moving*, the individual (either chick or parent) moves within the nest or the parent
makes changes in the nest (e.g., nest-building);
• *Other,* the individual does not display any of the behaviors mentioned above, is outside
of the nest or is not visible.

We recorded 328 individual observations of chicks and 240 of adults in / rearing small
broods and 634 observations of chicks and 100 of adults in / rearing large broods. Each
individual appears multiple times in the dataset because each nest was recorded multiple
times throughout the nestling period.

Statistical analyses

155 All analyses were performed in R v.4.1.2 (R Core Team, 2023). All models were fitted using
156 package *lme4* v. 1.1.35.1 (Bates et al., 2015) and we tested the statistical significance of
157 variables and their interactions using the package *car* v. 3.1.2 (Fox & Weisberg, 2019).

Chick mass

159 We used linear mixed models to determine the effect of brood size on chick mass. To
160 account for variations in the age at cross-fostering, we included the age at which mass was
161 recorded as a covariate. Since not all chicks were cross-fostered and therefore weighed at
162 the same age, the average age at cross-fostering per brood served as the reference point
163 for the first timepoint, and day 15 served as the second timepoint for each observation.

Chick behavior

We used linear mixed models with the arcsine transformed proportion ($\theta = \arcsin \sqrt{p}$,
where p is the proportion) of time spent on each behavior per observation as the response
variable using brood size, age category of the recording (0-4, 5-9, 10-5 and 16+ days) and

average age at cross fostering as explanatory variables while brood and observer
identities and were included as random intercepts. We used a chi-square test to test the
significance of the interaction between brood size and age category.

Parent behavior

We used linear mixed models with the arcsine transformed proportion ($\theta = \arcsin \sqrt{p}$,
where p is the proportion) of time spent on feeding per parent pair as the response
variable using brood size as the explanatory variable and included brood identity as
random effect. Since parental behavior was not always recorded, parent provision rates
were calculated from chick provision rate data. Two chicks were observed in each video,
and we averaged the proportion of time the two chicks were observed being fed per
observation, then multiplied by two for small broods and by 6 for large broods. Lastly, we
divided the result by two to obtain the proportion of time spent feeding per parent
(because sex of the feeding parent was not always recorded).

Results

Chick mass

Brood size affected chick growth as evidenced by a significant interaction between age
and brood size, confirming diminished growth rate in chicks in large broods (Table 1).
Being raised in large broods resulted in gaining 0.04±0.2 g less per day (Fig.1a). Age at
cross-fostering did not significantly increase the explained variance ($p=0.80$) and was
therefore excluded from the final model. At age 15 days, chicks raised in large broods
weighed on average 0.5±2.33 g less than chick reared in small broods (Fig.1b).

Figure 1. Mass gain per day of chicks raised in small and large broods (a). Mass of chicks raised in
small and large broods on day 15 (b). Violin plot widths represent the data density, boxplots show
the median and quartiles while dashed lines show the average daily mass gain and average mass on
day 15 for chicks raised in small and large broods.

Table 1. Linear mixed effects model results of the effects of brood size and age on chick mass (gr).

	Chick mass (gr)			
	Estimate (± SE)	t-value	df	Pr(>|t|)
Fixed effects				
Intercept	0.34 (± 0.30)	1.15	77.92	0.25
large brood	0.13 (± 0.45)	0.3	63.32	0.77
age (days)	0.75(± 0.02)	37.09	98.14	< 0.001
large brood * age	-0.07 (± 0.03)	-2.29	98.68	0.02
Random effects				
bird id	0.12			
brood id	0.7			
Residual	1.33			

Begging behavior

Chicks in large broods begged on average 7% more than chicks from small broods. This
increase was not at the expense of resting time, which was not affected by brood size, but
rather of other behaviors (Fig.2). The proportion of time at which chicks displayed each
behavior per age category is summarized in Figure S1. Chicks in large broods consistently
begged more throughout the nestling period, independent of age ($\chi^2_{(3, N = 199)}$ = 0.98, *p*
>0.8). Begging intensity peaking between day five and fifteen (Table 2, Fig.3).

Figure 2. Proportion of time chicks in small and large broods spent on each behavior over all ages
pooled. Proportions of each behavior per observation were normalized and then square root
transformed. See Figure S1 for the results for the different age categories.

Figure 3. Proportion of time chicks in small and large broods spent begging over the nestling period. For graphical purposes proportions were square root transformed. Violin plot widths represent the data density, boxplots show the median and quartiles.

Table 2. Linear mixed effects model results brood size and age effects on the proportion of time (arcsine square root transformed) that nestlings spent begging (N=139).

	Proportion of time chicks spent begging			
	Estimate (± SE)	t-value	df	Pr(>|t|)
Fixed effects				
Intercept	0.0089 ± 0.0041	2.2	4.98	0.08
large brood	0.012 ± 0.0022	5.71	23.28	<0.001
age 5-9 days	0.017 ± 0.0027	6.3	189	<0.001
age 10-15 days	0.014 ± 0.0028	5.2	171.2	<0.001
age 16+ days	0.0028 ± 0.0031	0.89	191.7	0.39
Random effects				
brood id	1.59e-05			
observer	2.72e-05			
residual	1.08e-04			

Parent provisioning

Parents caring for large broods spent 4.6% and parents caring for small broods spent 0.9% of their time transferring food to the chicks (Fig.4, Table 3). The amount of time

parents spent feeding their brood did not significantly differ between the early days of the chicks' lives and the later age categories (Fig.4, Table 3).

Figure 4. Proportion of time each parent spent transferring food to the chicks in small and large broods over the nestling period. For graphical purposes proportions were square root transformed. Violin plot widths represent the data density, boxplots show the median and quartiles.

Table 3. Linear mixed effects model results of brood size effects on the proportion of time (arcsine square root transformed) parents spent feeding their chicks (N=73).

	Proportion of time spent feeding per parent			
	Estimate (± SE)	df	t-value	Pr(>|t|)
Fixed effects				
Intercept	0.0040 ± 0.0064	3.34	061	0.58
large brood	0.0099 ± 0.0022	32	4.43	<0.001
age 5-9 days	0.0030 ± 0.0036	62.19	0.83	0.41
age 10-15 days	0.00079 ± 0.0036	63.72	0.22	0.83
age 16+ days	-0.0022 ± 0.0040	58.77	-0.55	0.56
Random effects				
brood id	1.04e-05			
observer	9.01e-05			
residual	5.73e-05			

Per capita chick feeding

Chicks reared in large broods spent 0.16% more time being fed than chicks in small broods (Fig.5, Table 4). The interaction between brood size and age category did not explain a significant part of the variance when added to the model ($\chi^2_{(3, N = 199)}$ = 0.65, p >0.8). The proportion of time being fed peaked between five and fifteen days for chicks of both brood sizes (Fig.5b).

Figure 5. Proportion of time chicks in small and large broods spent being fed over the nestling period. For graphical purposes proportions were square root transformed. Violin plot widths represent the data density, boxplots show the median and quartiles.

Table 4. Linear mixed effects model results of brood size and age effects on proportion of time (arcsine square root transformed) individual chicks spent being fed (N=199).

	Proportion of time chicks spent being fed			
	Estimate (± SE)	df	t-value	Pr(>|t|)
Fixed effects				
Intercept	0.0037 ± 0.0039	2.33	0.96	0.42
large brood	0.0039 ±0.0013	29.95	2.88	<0.01
age 5-9 days	0.0029 ± 0.0012	184.37	2.48	0.014
age 10-15 days	0.0016 ± 0.0012	188.05	1.29	0.20
age 16+ days	-0.0029 ± 0.0014	180.41	-2.11	0.037
Random effects				
brood id	1.24e-05			
observer	3.89e-05			
residual	1.863e-05			

Discussion

It is well established that being raised in a larger brood adversely affects growth – even in captive zebra finches with an *ad libitum* food supply. To shed light on the mechanistic underpinnings of this effect we manipulated brood size and recorded parent and offspring behavior throughout the nestling period. We found that zebra finch nestlings raised in large broods are faced with a diminished growth rate, despite a slight increase in *per capita* feeding rate when compared to chicks raised in small broods. We hypothesize that

the diminished growth results from increased resource allocation to competition through begging, which is consistent with earlier studies across various bird species, which have shown that nestlings in larger broods beg more intensively (Leonard et al., 2000; Neuenschwander et al., 2003; Wright et al., 2002). Our findings suggest a complex trade-off between competitive behavior for survival and physical development in the nestling period of zebra finches, also supporting pervious claims about a significant energetic cost associated with begging.

When parents have to care for more chicks, they can adaptively adjust their provisioning rate to match the needs of a larger brood as displayed by the increased parental feeding for parents and *per capita* provisioning rates for chicks in large broods. Previous research has also demonstrated that larger broods are generally associated with increased parental provisioning rates in birds (Bowers et al., 2014; Lessells, 1993; Stoehr et al., 2001). Parents caring for large broods in our study spent more time feeding their chicks, which we can assume to be, at least in part, a response to the increase in begging, given that begging intensity as call playback experiments have previously successfully elevated parental provisioning (e.g. Burford et al., 1998; Ottosson et al., 1997; Price, 1998; Tarwater et al., 2009). In most studies, the effect of brood size manipulation on parental provisioning only partly compensated for the change in brood size, and, as a consequence, *per capita* provisioning rate is typically lower in enlarged broods (e.g. Dijkstra et al., 1990). Our study is unusual in this respect, in that the brood size effect on provisioning rate was such that there was an *increase* in *per capita* provisioning rate in large broods– although it should be kept in mind that time being fed is only a proxy of the actual amount of food the nestlings received. Begging is the foraging technique of altricial nestlings, and, as any foraging technique, the yield will depend on food availability in the environment. In our study, food availability was *ad libitum* from the perspective of the parents, which is likely to have shaped our results, in that the energy yield of begging was probably higher than it would have been in a more natural environment where parents usually have to make a greater effort to gather food.

With respect to the question how the brood size effect on nestling growth arises, we infer from the present findings that brood size increased *per capita* provisioning rate and that the brood size effect on growth must therefore be due to other factors than the amount of food received. Instead, we consider the increased energy allocation to begging the most likely cause of the brood size effect on nestling growth. Our findings thereby provide indirect evidence for a significant energetic cost associated with begging behavior, similar to experiments in which nestlings were forced to beg at a higher rate for the same amount of food, which also depressed growth (Nettle et al., 2017; Rodriguez-Girones et al., 2001; Soler et al., 2014). A new dimension revealed by the present study is that family dynamics can induce a qualitatively similar effect on growth in the absence of an experimentally set limit on how much food the nestlings consume. Interestingly, this suggests that the brood size effect on growth is in a way self-inflicted, because when parents had adjusted their provisioning rate spontaneously, i.e., without having been stimulated to do so by the nestling's begging intensity, nestlings in small and large broods could have allocated the same amount of energy to growth. Paying the begging costs, which affects inclusive fitness of all family members, may be the evolutionary stable solution that is the outcome of the parent-offspring conflict over the optimal level of parental care.

Acknowledgements

We thank the animal caretakers of the University of Groningen as well as numerous students including Gillian Bakx and Wietske van Buiten for assistance with data collection.

Funding

MT was supported by an Adaptive Life Scholarship awarded by the University of Groningen.

Conflict of interest

The authors declare no conflict of interest

Ethics approval

All methods and experiments for the zebra finches detailed in this manuscript were performed under approval of the Central Committee for Animal Experiments (Centrale Commissie Dierproeven) of the Netherlands, under license AVD1050020174344.

Data availability statement

The data and scripts can be made publicly available on Dryad upon acceptance.

References

Andrews, C., Viviani, J., Egan, E., Bedford, T., Brilot, B., Nettle, D., & Bateson, M. (2015). Early life adversity increases foraging and information gathering in European starlings , Sturnus vulgaris. *Animal Behaviour*, *109*(109), 123–132. https://doi.org/10.1016/j.anbehav.2015.08.009

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using {lme4}. *Journal of Statistical Software*, *67*(1), 1–48. https://doi.org/10.18637/jss.v067.i01

Boonekamp, J. J., Bauch, C., & Verhulst, S. (2020). Experimentally increased brood size accelerates actuarial senescence and increases subsequent reproductive effort in a wild bird population. *Journal of Animal Ecology*, *89*(6), 1395–1407. https://doi.org/10.1111/1365-2656.13186

Boonekamp, J. J., Mulder, G. A., Salomons, H. M., Dijkstra, C., & Verhulst, S. (2014). Nestling telomere shortening, but not telomere length, reflects developmental stress and predicts survival in wild birds. *Proceedings of the Royal Society B: Biological Sciences*, *281*(1785), 1–7. https://doi.org/10.1098/rspb.2013.3287

Bowers, E. K., Nietz, D., Charles, F., & Sakaluk, S. K. (2014). Parental provisioning in house wrens : effects of varying brood size and consequences for offspring. *Behavioral Ecology*, *25*(6), 1485–1493. https://doi.org/10.1093/beheco/aru153

Briga, M., Koetsier, E., Boonekamp, J. J., Jimeno, B., & Verhulst, S. (2017). Food availability affects adult survival trajectories depending on early developmental conditions. *Proceedings of the Royal Society B: Biological Sciences*, *284*(1846), 20162287. https://doi.org/10.1098/rspb.2016.2287

Briga, M., & Verhulst, S. (2021). Mosaic metabolic ageing: Basal and standard metabolic

rates age in opposite directions and independent of environmental quality, sex and life span in a passerine. *Functional Ecology*, *35*(5), 1055–1068. https://doi.org/10.1111/1365-2435.13785

Burford, J. E., Friedrich, T. J., & Yasukawa, K. (1998). Response to playback of nestling begging in the red-winged blackbird, Agelaius phoeniceus. *Animal Behaviour*, *56*(3), 555–561. https://doi.org/10.1006/anbe.1998.0830

Burness, G. P., McClelland, G. B., Wardrop, S. L., & Hochachka, P. W. (2000). Effect of brood size manipulation on offspring physiology: An experiment with passerine birds. *Journal of Experimental Biology*, *203*(22), 3513–3520. https://doi.org/10.1242/jeb.203.22.3513

Davies, N. B. (1976). Parental Care and the Transition To Independent Feeding in the Young Spotted Flycatcher (Muscicapa Striata). *Behavior*, 280–294.

de Kogel, C. H. (1997). Long-Term Effects of Brood Size Manipulation on Morphological Development and Sex-Specific Mortality of Offspring. *The Journal of Animal Ecology*, *66*(2), 167. https://doi.org/10.2307/6019

Dijkstra, C., Bult, A., S., B., Daan, S., Meijer, T., & ZIilster, M. (1990). Brood Size Manipulations in the Kestrel (Falco tinnunculus): Effects on Offspring and Parent Survival. *Journal of Animal Ecology*, *59*(1), 269–285.

Fox, J., & Weisberg, S. (2019). *An {R} Companion to Applied Regression* (Third). Sage. https://socialsciences.mcmaster.ca/jfox/Books/Companion/

Friard, O., & Gamba, M. (2016). BORIS : a free , versatile open-source event-logging software for video / audio coding and live observations. *Methods in Ecology and Evolution*, *7*, 1325–1330. https://doi.org/10.1111/2041-210X.12584

Gerritsma, Y. H. (2025). *From adversity to personality. The effect of early-life adversity on zebra finch behaviour.*

Gil, D., Bulmer, E., Celis, P., & López-Rull, I. (2008). Adaptive developmental plasticity in growing nestlings: Sibling competition induces differential gape growth. *Proceedings of the Royal Society B: Biological Sciences*, *275*(1634), 549–554. https://doi.org/10.1098/rspb.2007.1360

Godfray, H. C. J. (1991). Signaling of need by offspring to their parents. *Nature*, *352*, 328–330.

Godfray, H. C. J. (1995). Signaling of need between parents and young: parent-offspring conflict and sibling rivalry. *The American Naturalist*, *146*(1), 1–24.

Godfray, H. C. J., & Johnstone, R. (2000). Begging and bleating : the evolution of parent – offspring signalling Begging and bleating : the evolution of parent – offspring signalling. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *355*, 1581–1591. https://doi.org/10.1098/rstb.2000.0719

Godfray, H. C. J., & Parker, G. (1992). Sibling competition , parent-offspring conflict and clutch size. *Animal Behaviour*, *43*(1974), 473–490.

Goldenberg, S. Z., & Wittemyer, G. (2018). Orphaning and natal group dispersal are associated with social costs in female elephants. *Animal Behaviour*, *143*, 1–8.

374 https://doi.org/10.1016/j.anbehav.2018.07.002

Griffith, S. C., & Buchanan, K. L. (2010). Maternal effects in the Zebra Finch: A model mother reviewed. *Emu*, *110*(3), 251–267. https://doi.org/10.1071/MU10006

Griffith, S. C., Holleley, C. E., Mariette, M. M., Pryke, S. R., & Svedin, N. (2010). Low level of extrapair parentage in wild zebra finches. *Animal Behaviour*, *79*(2), 261–264. https://doi.org/10.1016/j.anbehav.2009.11.031

Grodzinski, U., & Johnstone, R. A. (2012). Parents and offspring in an evolutionary game : the effect of supply on demand when costs of care vary. *Proceedings of the Royal Society B: Biological Sciences*, *279*, 109–115. https://doi.org/10.1098/rspb.2011.0776

Harper, A. B. (1986). The evolution of begging: sibling competition and parent- offspring conflict. *American Naturalist*, *128*(1), 99–114. https://doi.org/10.1086/284542

Haskell, D. G. (1994). Experimental evidence that nestling begging behaviour incurs a cost due to nest predation. *Proceedings of the Royal Society B: Biological Sciences*, *257*(1349), 161–164. https://doi.org/10.1098/rspb.1994.0110

Haskell, D. G. (1999). The effect of predation on begging-call evolution in nestling wood warblers. *Animal Behaviour*, *57*(4), 893–901. https://doi.org/10.1006/anbe.1998.1053

Holveck, M. J., & Riebel, K. (2010). Low-quality females prefer low-quality males when choosing a mate. *Proceedings of the Royal Society B: Biological Sciences*, *277*(1678), 153–160. https://doi.org/10.1098/rspb.2009.1222

Hõrak, P. (2003). When to pay the cost of reproduction? A brood size manipulation experiment in great tits (Parus major). *Behavioral Ecology and Sociobiology*, *54*(2), 105–112. https://doi.org/10.1007/s00265-003-0608-1

Kilner, R. M., & Johnstone, R. A. (1997). Begging the question: are offspring solicitation behaviors signals of need? *Trends in Ecology & Evolution*, *12*(1), 11–15.

Leech, S. M., & Leonard, M. L. (1996). Is there an energetic cost to begging in nestling tree swallows (Tachycineta bicolor)? *Proceedings of the Royal Society B: Biological Sciences*, *263*(1373), 983–987. https://doi.org/10.1098/rspb.1996.0145

Leech, S. M., & Leonard, M. L. (1997). Begging and the risk of predation in nestling birds. *Behavioral Ecology*, *8*(6), 644–646. https://doi.org/10.1093/beheco/8.6.644

Leonard, M. L., Horn, A. G., Gozna, A., & Ramen, S. (2000). Brood size and begging intensity in nestling birds. *Behavioral Ecology*, *11*(2), 196–201.

Lessells, C. M. (1993). The cost of reproduction: do experimental manipulations measure the edge of the options set. *Etologia*, *3*, 95–111.

Mainwaring, M. C., Blount, J. D., & Hartley, I. A. N. R. (2012). Hatching asynchrony can have long-term consequences for offspring fitness in zebra finches under captive conditions. *Biological Journal of the Linnean Society*, *106*, 430–438.

Mccarty, J. P. (1996). The energetic cost of begging in nestling passerines. *The Auk*, *113*(1), 178–188.

Mock, D. W., Schwagmeyer, P. L., & Dugas, M. B. (2009). Parental provisioning and nestling mortality in house sparrows. *Animal Behaviour*, *78*(3), 677–684. https://doi.org/10.1016/j.anbehav.2009.05.032

Naguib, M., Riebel, K., Marzal, A., & Gil, D. (2004). Nestling immunocompetence and testosterone covary with brood size in a songbird. *Proceedings of the Royal Society B: Biological Sciences*, *271*(1541), 833–838. https://doi.org/10.1098/rspb.2003.2673

Nettle, D., Andrews, C., Reichert, S., Bed, T., Kolenda, C., Parker, C., Martin-ruiz, C., Monaghan, P., & Bateson, M. (2017). Early-life adversity accelerates cellular ageing and affects adult inflammation : Experimental evidence from the European starling. *Scientific Reports*, *7*(40794), 1–10. https://doi.org/10.1038/srep40794

Nettle, D., Monaghan, P., Gillespie, R., Brilot, B., Bedford, T., Bateson, M., & Nettle, D. (2015). An experimental demonstration that early-life competitive disadvantage accelerates telomere loss. *Proceedings of the Royal Society B: Biological Sciences*, *282*(20141610), 1–8.

Neuenschwander, S., Brinkhof, M. W. G., & Ko, M. (2003). *Brood size , sibling competition , and the cost of begging in great tits (Parus major).* *14*(4), 457–462.

Nicolaus, M., Michler, S. P. M., Ubels, R., Van Der Velde, M., Komdeur, J., Both, C., & Tinbergen, J. M. (2009). Sex-specific effects of altered competition on nestling growth and survival: An experimental manipulation of brood size and sex ratio. *Journal of Animal Ecology*, *78*(2), 414–426. https://doi.org/10.1111/j.1365-2656.2008.01505.x

Nilsson, J. Å. (2002). Metabolic consequences of hard work. *Proceedings of the Royal Society B: Biological Sciences*, *269*(1501), 1735–1739. https://doi.org/10.1098/rspb.2002.2071

Nilsson, J. Å., & Svensson, M. (2001). Sibling competition affects individual growth strategies in marsh tit , Parus palustris , nestlings. *Animal Behaviour*, *61*, 357–365. https://doi.org/10.1006/anbe.2000.1602

Nussey, D. H., Kruuk, L. E. B., Morris, A., & Clutton-brock, T. H. (2007). Environmental conditions in early life influence ageing rates in a wild population of red deer. *Current Biology*, *17*(23), 1000–1001.

Ottosson, U., Bäckman, J., & Smith, H. G. (1997). Begging affects parental effort in the pied flycatcher, Ficedula hypoleuca. *Behavioral Ecology and Sociobiology*, *41*(6), 381–384. https://doi.org/10.1007/s002650050399

Patterson, S. K., Strum, S. C., Silk, J. B., & Patterson, S. K. (2022). Early life adversity has long-term effects on sociality and interaction style in female baboons. *Proceedings of the Royal Society B: Biological Sciences*, *289*(20212244), 1–9.

Price, K. (1998). Benefits of begging for yellow-headed blackbird nestlings. *Animal Behaviour*, *56*(3), 571–577. https://doi.org/10.1006/anbe.1998.0832

R Core Team. (2023). *R: A language and environment for statistical computing.* (RStudio 2021.09.0+351 "Ghost Orchid" Release. https://www.r-project.org/

Rodriguez-Girones, M. A., Zuniga, J. M., & Redondo, T. (2001). Effects of begging on

growth rates of nestling chicks. *Behavioral Ecology*, *12*(3), 269–274.

Royle, N. J., Hartley, I. R., & Parker, G. A. (2002). Begging for control : when are offspring solicitation behaviours honest ? *Trends in Ecology & Evolution*, *17*(9), 434–440.

Schwagmeyer, P., & Mock, D. (2008). Parental provisioning and offspring fitness : size matters. *Animal Behaviour*, *75*, 291–298. https://doi.org/10.1016/j.anbehav.2007.05.023

Soler, M., Ruiz-Raya, F., Carra, L. G., Medina-Molina, E., Ibáñez-Alamo, J. D., & Martín-Gálvez, D. (2014). A long-term experimental study demonstrates the costs of begging that were not found over the short term. *PLoS ONE*, *9*(11). https://doi.org/10.1371/journal.pone.0111929

Spagopoulou, F., Teplitsky, C., Lind, M. I., Chantepie, S., Gustafsson, L., & Maklakov, A. A. (2020). Silver-spoon upbringing improves early-life fitness but promotes reproductive ageing in a wild bird. *Ecology Letters*, *23*(6), 994–1002. https://doi.org/10.1111/ele.13501

Stoehr, A. M., McGraw, K. J., Nolan, P. M., & Hill, G. E. (2001). Parental Care in Relation To Brood Size in the House Finch. *Journal of Field Ornithology*, *72*(3), 412–418. https://doi.org/10.1648/0273-8570-72.3.412

Tarwater, C. E., Kelley, J. P., & Brawn, J. D. (2009). Parental response to elevated begging in a high predation, tropical environment. *Animal Behaviour*, *78*(5), 1239–1245. https://doi.org/10.1016/j.anbehav.2009.07.040

Trivers, R. L. (1974). Parent-Offspring Conflict. *American Zoologist*, *264*, 249–264.

Tung, J., Archie, E. A., Altmann, J., & Alberts, S. C. (2016). Cumulative early life adversity predicts longevity in wild baboons. *Nature Communications*, *7*(11181), 1–7. https://doi.org/10.1038/ncomms11181

van de Pol, M., Bruinzeel, L. W., Heg, D., van der Jeugd, H., & Verhulst, S. (2006). A silver spoon for a golden future : long-term effects of natal origin on fitness prospects of oystercatchers (Haematopus ostralegus). *Journal*, *75*, 616–626. https://doi.org/10.1111/j.1365-2656.2006.01079.x

Verhulst, S., Holveck, M. J., & Riebel, K. (2006). Long-term effects of manipulated natal brood size on metabolic rate in zebra finches. *Biology Letters*, *2*(3), 478–480. https://doi.org/10.1098/rsbl.2006.0496

Verhulst, S., & Wiersma, P. (1997). Is begging cheap? *The Auk*, *114*(1), 134.

Wright, J., Hinde, C., Fazey, I., & Both, C. (2002). Begging signals more than just short-term need: Cryptic effects of brood size in the pied flycatcher (Ficedula hypoleuca). *Behavioral Ecology and Sociobiology*, *52*(1), 74–83. https://doi.org/10.1007/s00265-002-0478-y

Wright, J., & Leonard, M. L. (2002). *The Evolution of Begging*. Kluwer Academic Publishers.

Zann, R. A. (1996). *The zebra finch. A synthesis of field and laboratory studies.* Oxford University Press.

Supplementary information to

Begging costs rather than food received cause brood size effect on growth in zebra finches

Marianthi Tangili[1*], Michael Briga[2] & Simon Verhulst[1*]

[1] Groningen Institute for Evolutionary Life Sciences, University of Groningen, Groningen,
The Netherlands

[2] Department of Biology, University of Turku, Finland

Figure S1. Proportion of time spent on each behavior by chicks in small and large broods divided in
the age categories of the chicks. For graphical purposes proportions of each behavior per observation
were normalized and then square root transformed.