Belowground competition increases root allocation in agreement with game-theoretical predictions, but only when plants simultaneously compete aboveground

Emanuel B. Kopp[1, 2] [*], Niels P.R. Anten[3], Pascal A. Niklaus[2,4] and Samuel E. Wuest[1] [*]

1) Group Breeding Research, Agroscope, Mueller-Thurgau-Strasse 29, 8820 Waedenswil, Switzerland

2) Department of Evolutionary Biology and Environmental Studies, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland

3) Centre for Crop Systems Analysis, Wageningen University, Wageningen, The Netherlands

4) Zurich-Basel Plant Science Center

* correspondence to emanuel.kopp@proton.me or samuel.wuest@agroscope.admin.ch

Keywords: crop evolutionary ecology, allocation, soybean, plant competition, game theory, shade avoidance, crop breeding, aboveground–belowground interactions

Abstract

Competition among plants can lead to allocation strategies that favor individual competitiveness at the expense of group-level productivity. However, the role of root and shoot responses in driving these outcomes is intensely debated. Experimental approaches often have difficulty disentangling above- and belowground interactions due to the confounding effects of pot size and nutrient distribution. Here, we used physical dividers and varied inter-plant distances (3–24 cm) in soybean to isolate competitive interactions while controlling for these confounding effects. Simultaneous above- and belowground competition increased relative root allocation, but reduced total biomass. Aboveground competition alone had stronger effects, triggering shade avoidance and reducing both shoot and root biomass. This likely overrode belowground cues. These results underscore the significance of neighbor-induced responses, particularly under full competition, as pivotal drivers of allocation patterns and promising targets for breeding strategies that enhance collective crop performance.

Introduction

Evolutionary game theory posits that selection in competitive environments favors resource acquisition strategies that benefit the individual but are detrimental at the level of the population (Anten & Vermeulen 2016; Gersani *et al.* 2001; Zhang *et al.* 1999). Conversely, strategies optimal for population-level productivity typically are not evolutionary stable, because such stands can be invaded by individuals that optimize their own resource acquisition at the expense of the group (Aschehoug *et al.* 2016; Harper 1977). This phenomenon is known as "tragedy of the commons" and has profound implications in agriculture, where the selection of overly competitive individuals during breeding needs to be avoided (Denison 2012; Donald 1968; Jennings & Herrera 1968; Weiner 2019).

In crops, some important traits associated with competition are well-described. Examples are plant height and an open shoot architecture. In maize, rice, and wheat, breeding for short and compact shoots accordingly contributed to substantial yield increases (Duvick *et al.* 2003; Hedden 2003; Jennings & Herrera 1968). Another trait detrimental to yield is shade avoidance, where plants sense light quality changes caused by the presence of neighbors and respond by elongating shoots and increasing leaf area, especially at the top of the canopy, where they can shade neighbors (Weiner *et al.* 2010). In soybean (*Glycine max*), shade often increases petiole and internode lengths but reduces branching (Lyu *et al.* 2023), and often leads to higher disease susceptibility and lower yield per plant (Green-Tracewicz *et al.* 2011; Lyu *et al.* 2023). Shade avoidance responses typically are undesired because they divert resources away from seed production (Pantazopoulou *et al.* 2021; Weiner *et al.* 2010). At the same time, some degree of light sensing and shade avoidance-like growth responses are required for efficient light harvesting and for canopy organization (López Pereira *et al.* 2017; Zhou *et al.* 2024). This trade-off makes targeting shade avoidance in breeding difficult.

Plants also compete belowground, but the consequences for population-level growth

and yield are less well understood. Game-theoretical modelling suggests that the adaptation of

plants to root competition can lead to the diversion of resources from yield to root growth, i.e.

to a tragedy of the commons similar to the one observed under aboveground competition

(Gersani *et al.* 2001; Zhang *et al.* 1999). Indeed, experiments by Gersani *et al.* (2001) using

belowground root system separation showed that soybean plants competing belowground, i.e.

without separation, had higher root biomass and lower seed yield than plants whose root

systems were separated by a physical divider. However, one difficulty in interpreting this result

is that pot-bounding effects may have occurred because root system separation also halved the

total soil volume available to each plant (Hess & De Kroon 2007; McNickle 2020; Poorter &

Sack 2012; Semchenko *et al.* 2007). An ideal experiment would simultaneously account for the

effects of pot volume, total amount of nutrients per plant, and for soil nutrient concentrations,

which is difficult to achieve (Chen *et al.* 2015; Maina *et al.* 2002; McNickle 2020). Recently,

root-impermeable mesh dividers that allow nutrients to pass through have been used to keep

available root space constant (Chen *et al.* 2021; Zhu *et al.* 2019). However, the assumption

remains that root growth responses are caused by signals that pass through the mesh divider

and do not require direct root contact (Cabal 2022; Chen *et al.* 2021; McNickle 2020; Zhu *et al.*

2019). Collectively, these studies show that the presence of neighbor plants can increase root

biomass under belowground competition, although this effect is not always observed (Lepik *et*

al. 2021; Schmid *et al.* 2015). Indeed, a recent meta-analysis confirmed such conflicting root

responses in pea (Mobley *et al.* 2022).

Recently, Cabal *et al.* (2020) suggested that seemingly contradictory effects of

neighbors on root growth occur because they are distance dependent. Assuming that the cost of

growing and maintaining roots increases with distance from the stem, a game-theoretical model

predicted that competing plants would shift additional root production to the stem because the

cost of root competition is lower there. Thus, the phenomenon that emerges from such a model, which was termed "exploitative segregation of plant roots", is one where the same resource foraging strategy leads to different root allocation responses depending on the distance between competing plants (Cabal 2022; Cabal *et al.* 2020).

Here, we studied biomass allocation in soybean plants competing aboveground, belowground, both aboveground and belowground, or not at all. We assessed root and shoot biomass at flowering, as plants had reached a size at which they strongly competed but still were growing rapidly. This allowed us to determine the competitive strategy of the plant, with organ size differences both reflecting differences in resource allocation up to this point and determining its future growth (Sun *et al.* 2021). Although this would have allowed quantification of seed yield, we chose not to assess biomass at senescence because resources would have been remobilized from the vegetative biomass, obscuring the earlier effects of biomass allocation (Liu *et al.* 2019; Maillard *et al.* 2015).

Combining the traditionally used separation treatments with a range of distances between plants (Fig. 1) allowed us to modulate the strength of competition between neighbors while keeping the available soil volume constant. Accordingly, the primary metric of interest is the statistical interaction of the separation treatment with the distance to the neighbor. Specifically, we expected neighbor-induced allocation changes to be largely absent when plants were separated or at a large distance. Under belowground competition, we expect a preferential allocation to roots, i.e. a root-to-shoot ratio that increases as plant distance decreases (Fig. 1a). Conversely, under aboveground competition, we predicted a preferential allocation to shoots, i.e. a root-to-shoot ratio that decreases as plant get closer (Fig. 1b). When plants compete both aboveground and belowground, predictions are more difficult to make because aboveground and belowground responses are coupled. We therefore expected their effects to interact. For

105   example, the effects of belowground competition would depend on whether plants also compete

106   aboveground (Fig. 1c).

Material and Methods

In Summer 2023, an outdoor pot experiment was established in Wädenswil, Switzerland

(47.222°N, 8.669°E, 509 m a.s.l.) to study competition-effects on allocation in soybean. After

inoculating soybean seeds with commercial rhizobia (HiStick, BASF, Ludwigshafen, Germany;

containing *Bradyrhizobium japonicum*), plants were germinated and grown in propagator trays

in a greenhouse for 14 days, and seedlings then transplanted in pairs to 7.5 l rectangular pots

filled with lawn soil (Ökohum Art. 2633200, Herrenhof, Switzerland) and moved outside. There,

the plant pairs were separated aboveground, belowground, both aboveground and belowground,

or not at all. The aboveground separation consisted of a white polystyrene sheet (80 cm high ×

20 cm wide) that prevented shoot competition. The belowground separation consisted of a

water-impermeable plastic sheet dividing the pot into halves. Each divider combination was

combined with an inter-plant distance treatment, starting at a very close interplant distance

relative to plant size, that is at a distance where plants were strongly competing (3, 6, 9, 12, 15,

18, 21, and 24 cm). Each pot contained two plants of one of four genotypes (Weber - PI 548524,

Natsu Kurakake – PI 417187, Kanegawawase - PI 229330, or Kanagawa Wase - PI 506832;

USDA Soybean Germplasm Collection, Urbana, IL, USA). We previously established that these

four genotypes formed a tentative gradient in competitive strategies, with Weber being least

and Natsu Kurakake being most competitive. Each treatment combination was replicated three

times, resulting in a total of 384 pots (2 aboveground separation treatments × 2 belowground

separation treatments × 8 distances × 4 genotypes × 3 replicates). Pots were protected from hail

by a net and irrigated ad libitum (Figure S1 in Supporting Information).

The plants were grown for 55 days until the majority of them had started to flower.

Starting on day 20, we measured plant height once a week, for a total of six measurements. On

day 54, we quantified relative chlorophyll content (MultispeQ V 2.0;   PhotosynQ.com;

Kuhlgert *et al.* 2016). On day 55, we clipped shoots at ground level, washed roots, and dried

(60°C, 48 h) and weighed all biomass samples. Because the two plants in each pot were in

symmetrical positions, all biomass data were analyzed at the pot level.

We analyzed all data using linear models (aov function of R 4.3.1; http://www.r-

project.org). Biomass data was analyzed without allometric correction, because the biomass

data was independent of root-to-shoot ratio (R^2 = 0.001, p > 0.5). The linear models contained

row and column terms (the pots were arranged on a rectangular grid; Fig. S1) to account for

spatial effects within the hail protection tunnels, followed by the full-factorial combination of

aboveground and belowground dividers, the distance between the competing plants, and

genotype. Given our hypothesis, we were primarily interested in the dependence of the

separator effect on distance.

Results

Genotypes differed in all traits measured (P < 0.001 for main effect of genotype, Table 1, Fig.

2). This was expected, since genotypes were deliberately chosen to represent different growth

strategies in soybean. However, all other effects we were interested in were independent of

genotype, i.e. there was no statistical interaction of genotype with interplant distance and

aboveground or belowground separation.

Our primary interest was on effects of root competition. Some aspects of plant growth,

however, depended only on shoot separation. First, shoot competition caused shade avoidance

responses. This manifested in lower shoot-biomass-to-height ratios, i.e. plants were more

slender when close together (aboveground separation x distance $F_{1,347}$ = 4.03, P < 0.05; Table

1, Fig. 3a). Second, both root and total biomass decreased under shoot competition

(aboveground separation x distance: $F_{1,347} > 5$ and $P < 0.05$ for both; Table 1, Fig. 3b, c).

All other growth-related parameters also depended on root separation. Specifically, root

competition effects were contingent on shoot competition, manifesting as three-way

interactions between the two separation treatments and distance. In the absence of aboveground

separators, the effects of root competition were consistent with our hypothesis. Specifically,

root-to-shoot ratios increased under belowground competition, and this effect was strongest

when plants were close together. In contrast, no such increase was found when plants were

separated aboveground. These opposing patterns manifested as statistical three-way interaction

(aboveground separation x belowground separation x distance, $F_{1,347} = 4.16$, $P < 0.05$; Fig. 4a).

Conversely, shoot competition decreased root-to-shoot ratios, as expected, but only when plants

were separated belowground (Fig. 4b). Similar contingencies of root competition effects on

shoot competition and vice-versa were found for final plant height and relative chlorophyll

content (aboveground separation x belowground separation x distance, $F_{1,347} > 4$, P-values <

0.05; Table 1; Fig. S2). For example, plants competing above- and belowground were smaller

than plants competing belowground only, and plants had higher relative chlorophyll content

when competing belowground but not aboveground.

Shifts in root-to-shoot ratio may be caused by changes in root biomass, in shoot biomass,

or in both, and we therefore examined what caused the significant three-way interaction of

treatments and interplant distance on root-to-shoot ratios in our experiment. Belowground

competition had contrasting effects on shoot biomass, depending on whether plant competed

aboveground (aboveground separation x belowground separation x distance, $F_{1,347} = 5.4$, $P <$

0.05, Fig. 5). This increased root-to-shoot ratio when plants were not separated aboveground,

and reduced root-to-shoot ratios when they were. In contrast, root competition did not affect

root biomass. However, shoot competition decreased root biomass (Fig. 3b), a change which

lowers root-to-shoot ratio. Overall, our results indicate that the observed three-way interaction

in root-to-shoot ratio is predominantly driven by changes in shoot biomass.

Discussion

Concomitant aboveground and belowground competition shifts allocation to roots

In our experiment, we found increased relative allocation to roots and decreased total biomass

when plants were close to their neighbors, but only when plants were simultaneously competing

aboveground and belowground. In contrast, aboveground competition led to a reduction of total

biomass, irrespective of whether plants also competed belowground. Aboveground competition

also reduced final root biomass, which also was independent of belowground competition. Our

data thus indicates increased allocation to root growth under full competition, but nevertheless

a lower final root biomass. We think that occurs because increased allocation to root

competition reduces allocation to photosynthetically-active leaf area and thus slows the plants'

overall growth rates, which translates into a lower final root biomass because plants become

smaller. This pattern is in line with a tragedy of the commons in which competition with

neighbors reduces the combined performance of the two plants. Our results are in line with

Gersani *et al.* (2001), who manipulated belowground competition using physical dividers and

found evidence for such a tragedy of the commons for seed production (Hardin 1968). It is

interesting to note that Gersani *et al.* (2001) used a setup where competing plants were grown

relatively close to each other, without aboveground dividers and in a greenhouse over the winter

period requiring supporting artificial light – conditions that typically induce a shade avoidance

response. This growth condition was similar to our close-proximity and no-aboveground-

separator treatment, and the observed increased root allocation under such conditions is

consistent with our study. We therefore think that many discrepancies among studies could be

due to differences in experimental conditions such as light levels and light quality, interplant

distances, or pot sizes (Murphy & Dudley 2007). Particularly interplant distance, which in our

experiment interacted strongly with both above- and belowground competition, has received

surprisingly little attention in past work, despite its potential to modulate competition intensity

and neighbor detection (Cabal *et al.* 2020; Gottlieb & Gruntman 2022; McNickle 2020;

Semchenko *et al.* 2007).

208
Shade avoidance responses under aboveground competition led to a stronger decrease in

shoot biomass than in root biomass

Responses to aboveground competition were consistent with the typical shade avoidance

syndrome (Ballaré & Pierik 2017; Green-Tracewicz *et al.* 2011; Pantazopoulou *et al.* 2021), i.e.

plants were more slender and produced less root biomass. Plant height typically increases under

shade avoidance (Pierik & De Wit 2014). However, under shoot competition alone, final plant

height increased only slightly. On the contrary, simultaneous shoot and root competition had a

clear negative effect on final height. This reduction in height likely resulted from a reduction

in total plant biomass due to less-optimal allocation under shade avoidance. We think that this

is induced through light quality changes when neighbors are present, and that there was no

limitation of growth by available light (Fig. S1). Even though shade avoidance can increase the

fitness of individuals at high densities, our results and those of others suggest that the light-

foraging strategy invoked by the presence of neighbors is detrimental to the performance of the

overall stand (Pantazopoulou *et al.* 2021; Schmitt *et al.* 1995; Wille *et al.* 2017).

223
Aboveground competition has stronger effects than belowground competition

Competition between two plants ranges from fully symmetric to strongly asymmetric:

depending on the extent to which larger plants acquire amounts of resources that are

proportionately or disproportionately large relative to their size (Weiner 1990). Light

competition is usually asymmetric, because the taller plant can pre-empt light, as light is a

largely directional resource. Nutrient competition is more symmetric with nutrient acquisition

largely proportional to the size of the root system (Schwinning & Weiner 1998). Based on our

findings, we think that aboveground competition was stronger than belowground competition.

Indeed, the interaction between aboveground separation and neighbor distance was significant

for multiple measurements. However, the interaction between belowground separation and

neighbor distance was never significant in absence of an interaction with aboveground

separation. An integration of above- and belowground responses to competition on root growth

could occur through various processes. For example, plants may "sense" the strength of a

neighboring competitor aboveground and adjust their belowground foraging strategy (Gottlieb

& Gruntman 2022). Alternatively, as predicted by game theoretical models for belowground

interactions, aboveground interactions with neighbors may increase the costs of root production

and maintenance (Cabal *et al.* 2020).

241
The response to neighbors is a valuable breeding target

Here, we showed a novel and simple way to control for the confounding factors that have long

dominated the debate on how to interpret competition experiments involving pairs of plants.

We strongly suggest that future work investigating the effect of individual foraging strategies

on plant group performance should explicitly account for neighbor distance. It has long been

anticipated, most famously by Donald (1968), that reducing competitive traits in crop plants

may hold great potential for increasing yields (see also Duvick *et al.* 2003; Jennings & Herrera

1968). However, this has only recently become a renewed focus in the emerging research field

of "crop evolutionary ecology" (Anten & Vermeulen 2016; Denison 2012; Weiner 2019;

Weiner *et al.* 2017). Here, we found reduced biomass production under shoot competition,

despite similar resource availability. We also found a diversion of resources to root production

under combined root and shoot competition. We quantified plant biomass at flowering to

capture effects of allocation when plants were still rapidly growing, and found a tragedy of the

commons with respect to total plant biomass. It is plausible that a similar tragedy of the

commons would have been found for seed yield if the experiment had been run for longer (but

see Murphy & Dudley 2007). The fact that we found this effect across a range of genotypes that

varied substantially in allocation patterns suggest that this finding is general. Overall, this

suggests that responses to the presence of neighbors bear potential as breeding target.

Acknowledgements

262 This work was supported by the Swiss National Science Foundation, project 310030_192537.

263 We acknowledge Lukas Vonmetz and Jürgen Krauss (both Agroscope) for technical support.

Competing interests

265 The authors declare no competing interests.

Author contributions

267 E.B.K., S.E.W. and P.A.N conceptualized and designed the research (with input from N.P.R.A.).

268 E.B.K and S.E.W performed the experiments. E.B.K performed the analyses and wrote the

269 manuscript with input from S.E.W and P.A.N. All authors revised and approved the final version

270 of the manuscript.

Data accessibility statement

272 The data that support the findings of this study are openly available on Zenodo at

273 http://doi.org/10.5281/zenodo.15524543

Table 1: Statistical tests (F ratio and P-value) for effects of genotype (G), the distance between the pair of plants (D), the aboveground separation treatment (C_A), the belowground separation treatment (C_B), and including all interactions of the competition treatments with distance.

Response	Explanatory variable[1]	F (df_n, df_d)	p-value
	G	161.03 (3, 347)	**< 0.001 ***
	D	>0.001 (1, 347)	0.99
	D x C_A	6.845 (1, 347)	**0.009 ***
Total biomass	D x C_B	0.02 (1, 347)	0.9
	G x D x C_A	1.784 (3, 347)	0.15
	G x D x C_B	0.122 (3, 347)	0.95
	D x C_A x C_B	2.62 (1, 347)	0.106
	G	132.46 (3, 347)	**< 0.001 ***
	D	0.14 (1, 347)	0.71
	D x C_A	5.71 (1, 347)	**0.017 ***
Shoot biomass	D x C_B	0.15 (1, 347)	0.7
	G x D x C_A	1.04 (3, 347)	0.37
	G x D x C_B	0.2 (3, 347)	0.9
	D x C_A x C_B	5.4 (1, 347)	**0.021 ***
	G	187.11 (3, 347)	**< 0.001 ***
	D	2.66 (1, 347)	0.6
	D x C_A	5.49 (1, 347)	**0.02 ***
Root biomass	D x C_B	0.69 (1, 347)	0.79
	G x D x C_A	2.2 (3, 347)	0.088
	G x D x C_B	0.05 (3, 347)	0.99
	D x C_A x C_B	0.09 (1, 347)	0.76
	G	164.88 (3, 347)	**< 0.001 ***
	D	0.793 (1, 347)	0.37
	D x C_A	0.006 (1, 347)	0.94
Root-to-shoot ratio	D x C_B	0.58 (1, 347)	0.45
	G x D x C_A	0.786 (3, 347)	0.5
	G x D x C_B	0.172 (3, 347)	0.92
	D x C_A x C_B	4.163 (1, 347)	**0.042 ***
	G	136.85 (3, 347)	**< 0.001 ***
	D	>0.001 (1, 347)	0.98
	D x C_A	2.24 (1, 347)	0.135
Final height	D x C_B	0.611 (1, 347)	0.435
	G x D x C_A	0.113 (3, 347)	0.952
	G x D x C_B	0.3 (3, 347)	0.825
	D x C_A x C_B	4.774 (1, 347)	**0.03 ***
	G	67.55 (3, 347)	**< 0.001 ***
Shoot-biomass-to-height ratio	D	0.11 (1, 347)	0.74
	D x C_A	4.031 (1, 347)	**0.045 ***

	D x C_B	0.01 (1, 347)	0.908
	G x D x C_A	1.101 (3, 347)	0.349
	G x D x C_B	0.992 (3, 347)	0.397
	D x C_A x C_B	2.176 (1, 347)	0.141
	G	6.545 (3, 347)	**< 0.001 ***
	D	2.98 (1, 347)	0.085
	D x C_A	1.761 (1, 347)	0.185
Relative chlorophyll content	D x C_B	0.244 (1, 347)	0.621
	G x D x C_A	1.282 (3, 347)	0.28
	G x D x C_B	0.696 (3, 347)	0.555
	D x C_A x C_B	4.06 (1, 347)	**0.045 ***

276   [1]: To avoid confounding with pot volume, competition treatments can only be interpreted in interaction with inter-plant distance.

Figure 1: Expected ratio of root biomass to shoot biomass in dependence of root and shoot separation and inter-plant distance. Increasing inter-plant distance allows to decrease competition; hence, the effect of competition can be assessed as biomass at close distance relative to the situation when plants are far from each other. (a) When plants are always separated aboveground, we expect root competition to increase the root-to-shoot ratio. (b) When plants do not compete belowground, we expect aboveground competition to decrease the root-to-shoot ratio. (c) When plants simultaneously compete aboveground and belowground, we expect the change in root-to-shoot ratio to depend on whether the effect of competition is stronger aboveground or belowground.

Figure 2: The four genotypes strongly differed in all measured traits, as shown here for root

biomass. Error bars indicate +/– one standard error of the mean.

Figure 3: Shoot competition effects on (a) shoot biomass to height ratio, (b) root biomass, and (c) total biomass. The shoot separation effect is the difference between pots with and without shoot separation. We used the distance treatment to modulate the strength of competition. Hence, shoot competition manifests as difference between the separation effects when plants are close and when plants are distant. Lines show model predicted values.

Figure 4: (a) Root separation effects on root-to-shoot ratio, in dependence of inter-plant distance. When plants were not separated aboveground, root competition increased root-to-shoot ratio (green arrow). However, when plants were separated aboveground, root competition decreased root-to-shoot ratio (blue arrow). (b) Shoot separation effects on root-to-shoot ratio. When plants were not separated belowground, shoot competition increased root-to-shoot ratio (green arrow). However, when plants were separated belowground, shoot competition decreased root-to-shoot ratio (blue arrow). The dependency of root competition effects on shoot separation manifested as statistical interaction between the two separation treatments and distance ($F_{1,347} = 4.16$, $P < 0.05$, Table 1).

Figure 5: (a) Root separation effects on shoot biomass, in dependence of inter-plant distance. When plants were not separated aboveground, root competition decreased shoot biomass (green arrow). When plants were separated aboveground, root competition increased shoot biomass (blue arrow). (b) Shoot separation effects on shoot biomass, in dependence of inter-plant distance. When plants were not separated belowground, shoot competition decreased shoot biomass (green arrow). When plants were separated belowground, shoot competition slightly decreased shoot biomass (blue arrow). The dependency of root competition effects on shoot separation manifested as statistical interaction between the two separation treatments and distance ($F_{1,347} = 5.4$, $P < 0.05$, Table 1).

References

Anten, N.P.R. & Vermeulen, P.J. (2016). Tragedies and Crops: Understanding Natural
Selection To Improve Cropping Systems. *Trends Ecol. Evol.*, 31, 429–439.

Aschehoug, E.T., Brooker, R., Atwater, D.Z., Maron, J.L. & Callaway, R.M. (2016). The
Mechanisms and Consequences of Interspecific Competition Among Plants. *Annu.*
Rev. Ecol. Evol. Syst., 47, 263–281.

Ballaré, C.L. & Pierik, R. (2017). The shade-avoidance syndrome: multiple signals and
ecological consequences: Neighbour detection and shade avoidance. *Plant Cell*
Environ., 40, 2530–2543.

Cabal, C. (2022). Root tragedy of the commons: Revisiting the mechanisms of a
misunderstood theory. *Front. Plant Sci.*, 13, 960942.

Cabal, C., Martínez-García, R., De Castro Aguilar, A., Valladares, F. & Pacala, S.W. (2020).
The exploitative segregation of plant roots. *Science*, 370, 1197–1199.

Chen, B.J.W., During, H.J., Vermeulen, P.J., Kroon, H., Poorter, H. & Anten, N.P.R. (2015).
Corrections for rooting volume and plant size reveal negative effects of neighbour
presence on root allocation in pea. *Funct. Ecol.*, 29, 1383–1391.

Chen, B.J.W., Huang, L., During, H.J., Wang, X., Wei, J. & Anten, N.P.R. (2021). No
neighbour-induced increase in root growth of soybean and sunflower in mesh-divider
experiments after controlling for nutrient concentration and soil volume. *AoB*
PLANTS, 13, plab020.

Denison, R.F. (2012). *Darwinian agriculture: how understanding evolution can improve*
agriculture. Lightning Source UK Ltd., Princeton University Press, Milton Keynes
UK, Princeton, N.J.

Donald, C.M. (1968). The breeding of crop ideotypes. *Euphytica*, 17, 385–403.

Duvick, D.N., Smith, J.S.C. & Cooper, M. (2003). Long-Term Selection in a Commercial
Hybrid Maize Breeding Program. In: *Plant Breeding Reviews*. John Wiley & Sons,
Ltd, pp. 109–151.

Gersani, M., Brown, J.S., O'Brien, E.E., Maina, G.M. & Abramsky, Z. (2001). Tragedy of the
commons as a result of root competition: Game theory and root competition. *J. Ecol.*,
89, 660–669.

Gottlieb, R. & Gruntman, M. (2022). Can plants integrate information on above-ground
competition in their directional responses below ground? *Ann. Bot.*, 130, 763–771.

Green-Tracewicz, E., Page, E.R. & Swanton, C.J. (2011). Shade Avoidance in Soybean
Reduces Branching and Increases Plant-to-Plant Variability in Biomass and Yield Per
Plant. *Weed Sci.*, 59, 43–49.

Hardin, G. (1968). The Tragedy of the Commons. *Science*, 162, 1243–1248.

Harper, J. (1977). *Population biology of plants*. Academic Press, New York, NY.

Hedden, P. (2003). The genes of the Green Revolution. *Trends Genet.*, 19, 5–9.

Hess, L. & De Kroon, H. (2007). Effects of rooting volume and nutrient availability as an
alternative explanation for root self/non-self discrimination. *J. Ecol.*, 95, 241–251.

Jennings, P.R. & Herrera, R.M. (1968). Studies on Competition in Rice II. Competition in
Segregating Populations. *Evolution*, 22, 332–336.

Kuhlgert, S., Austic, G., Zegarac, R., Osei-Bonsu, I., Hoh, D., Chilvers, M.I., *et al.* (2016).
MultispeQ Beta: a tool for large-scale plant phenotyping connected to the open
PhotosynQ network. *R. Soc. Open Sci.*, 3, 160592.

Lepik, A., Abakumova, M., Davison, J., Zobel, K. & Semchenko, M. (2021). Spatial mapping
of root systems reveals diverse strategies of soil exploration and resource contest in
grassland plants. *J. Ecol.*, 109, 652–663.

Liu, Z., Marella, C.B.N., Hartmann, A., Hajirezaei, M.R. & Von Wirén, N. (2019). An Age-
Dependent Sequence of Physiological Processes Defines Developmental Root
Senescence. *Plant Physiol.*, 181, 993–1007.

López Pereira, M., Sadras, V.O., Batista, W., Casal, J.J. & Hall, A.J. (2017). Light-mediated
self-organization of sunflower stands increases oil yield in the field. *Proc. Natl. Acad.*
Sci., 114, 7975–7980.

Lyu, X., Mu, R. & Liu, B. (2023). Shade avoidance syndrome in soybean and ideotype toward
shade tolerance. *Mol. Breed.*, 43, 31.

Maillard, A., Diquélou, S., Billard, V., Laîné, P., Garnica, M., Prudent, M., *et al.*
(2015). Leaf mineral nutrient remobilization during leaf senescence and modulation
by nutrient deficiency. *Front. Plant Sci.*, 6.

Maina, G.G., Brown, J.S. & Gersani, M. (2002). Intra-plant versus Inter-plant Root
Competition in Beans: avoidance, resource matching or tragedy of the commons.
Plant Ecol., 160, 235–247.

McNickle, G.G. (2020). Interpreting plant root responses to nutrients, neighbours and pot
volume depends on researchers' assumptions. *Funct. Ecol.*, 34, 2199–2209.

Mobley, M.L., Kruse, A.S. & McNickle, G.G. (2022). *PISUM SATIVUM* has no competitive
responses to neighbors: A case study in (non)reproducible plant biology. *Plant Direct*,
6, e411.

Murphy, G.P. & Dudley, S.A. (2007). Above- and below-ground competition cues elicit
independent responses. *J. Ecol.*, 95, 261–272.

Pantazopoulou, C.K., Bongers, F.J. & Pierik, R. (2021). Reducing shade avoidance can
improve Arabidopsis canopy performance against competitors. *Plant Cell Environ.*,
44, 1130–1141.

Pierik, R. & De Wit, M. (2014). Shade avoidance: phytochrome signalling and other
aboveground neighbour detection cues. *J. Exp. Bot.*, 65, 2815–2824.

Poorter, H. & Sack, L. (2012). Pitfalls and Possibilities in the Analysis of Biomass Allocation
Patterns in Plants. *Front. Plant Sci.*, 3.

Schmid, C., Bauer, S. & Bartelheimer, M. (2015). Should I stay or should I go? Roots
segregate in response to competition intensity. *Plant Soil*, 391, 283–291.

Schmitt, J., McCormac, A.C. & Smith, H. (1995). A Test of the Adaptive Plasticity Hypothesis
Using Transgenic and Mutant Plants Disabled in Phytochrome-Mediated Elongation
Responses to Neighbors. *Am. Nat.*, 146, 937–953.

Schwinning, S. & Weiner, J. (1998). Mechanisms determining the degree of size asymmetry
in competition among plants. *Oecologia*, 113, 447–455.

Semchenko, M., Hutchings, M.J. & John, E.A. (2007). Challenging the tragedy of the
commons in root competition: confounding effects of neighbour presence and
substrate volume. *J. Ecol.*, 95, 252–260.

Sun, Y., Zang, H., Splettstößer, T., Kumar, A., Xu, X., Kuzyakov, Y., *et al.* (2021). Plant
intraspecific competition and growth stage alter carbon and nitrogen mineralization in
the rhizosphere. *Plant Cell Environ.*, 44, 1231–1242.

Weiner, J. (1990). Asymmetric competition in plant populations. *Trends Ecol. Evol.*, 5, 360–
364.

Weiner, J. (2019). Looking in the Wrong Direction for Higher-Yielding Crop Genotypes.
Trends Plant Sci., 24, 927–933.

Weiner, J., Andersen, S.B., Wille, W.K.-M., Griepentrog, H.W. & Olsen, J.M. (2010).
Evolutionary Agroecology: the potential for cooperative, high density, weed-
suppressing cereals. *Evol. Appl.*, 3, 473–479.

Weiner, J., Du, Y.-L., Zhang, C., Qin, X.-L. & Li, F.-M. (2017). Evolutionary agroecology:
individual fitness and population yield in wheat (*Triticum aestivum*). *Ecology*, 98,
2261–2266.
Wille, W., Pipper, C.B., Rosenqvist, E., Andersen, S.B. & Weiner, J. (2017). Reducing shade
avoidance responses in a cereal crop. *AoB PLANTS*, 9.
Zhang, D.-Y., Sun, G.-J. & Jiang, X.-H. (1999). Donald's ideotype and growth redundancy: a
game theoretical analysis. *Field Crops Res.*, 61, 179–187.
Zhou, Y., Kusmec, A. & Schnable, P.S. (2024). Genetic regulation of self-organizing
azimuthal canopy orientations and their impacts on light interception in maize. *Plant*
Cell, 36, 1600–1621.
Zhu, Y.-H., Weiner, J. & Li, F.-M. (2019). Root proliferation in response to neighbouring roots
in wheat (Triticum aestivum). *Basic Appl. Ecol.*, 39, 10–14.

Supporting Information

Figure S1: Experiment under hail protection net, at 41 days after seed germination. The conditions were chosen so that water or light limitations were excluded as much as possible.

Figure S2: Predicted marginal means for (a) final plant height, and (b) relative chlorophyll content depending on presence/absence of aboveground separation (panels) and on presence/absence of belowground separation (colors). The opposing patterns of the lines between the panels show the three-way interactions between the two separation treatments and the interplant distance.