Astrocytes expressing M$_4$ muscarinic acetylcholine receptor regulate

locomotion and survival in murine prion disease

Gonzalo S. Tejeda[1*], Colin Molloy[1], Carrie K. Jones[2], Craig W. Lindsley[2], Graeme

Milligan[1] and Andrew B. Tobin[1*]

[1] Centre for Translational Pharmacology, School of Molecular Biosciences, Advanced

Research Centre, 11 Chapel Lane, University of Glasgow, Glasgow G11 6EW, Scotland,

United Kingdom.

[2] Warren Centre for Neuroscience Drug Discovery, Department of Pharmacology,

Innovation Park, 393 Nichol Mills Road, Franklin, Tennessee, 37027, USA.

[*] Corresponding authors:

Gonzalo.Tejeda@glasgow.ac.uk

Andrew.Tobin@glasgow.ac.uk

Abstract

Whilst there has been much focus on neuronally expressed muscarinic acetylcholine

receptors (mAChRs), less attention has been paid to their expression and role in non-

neuronal brain cells. Using genetically engineered mice, we identify a previously

unappreciated subpopulation of astrocytes expressing the M4 mAChR subtype

dispersed across various regions that include the brainstem, hypothalamus and, most

abundantly, the cerebellum. Signalling and proteomic analysis of M4 mAChR positive

astrocytes from the cerebellum confirmed functional receptor expression and its role in

regulating protein expression. Genetic ablation of these astrocytes in mice revealed a

specific role in locomotion. Importantly, in the context of murine prion disease, we

observed a significant expansion of M4 mAChR positive astrocytes, and ablation

experiments established that this astrocyte subpopulation has a detrimental effect on

late-stage disease. Together we identify M4 mAChR expressing astrocyte subpopulation

that plays a specific role in normal neurophysiology and the progression of

inflammatory neurodegenerative disease.

Keywords: Astrocyte, M_4 muscarinic acetylcholine receptor, G protein-coupled

receptor, neuroinflammation, locomotion.

Main Text

Background

The muscarinic acetylcholine receptors (mAChRs) have been under intense

investigation due to the critical role of this family of G protein-coupled receptors

(GPCRs) in the regulation of various neuronal processes, including learning, memory,

reward and seek behaviour. Among the five subtypes (M_1-M_5 mAChRs), M_1 and M_4

have emerged as particularly important in brain function and they are considered

promising therapeutic targets for the treatment of several brain disorders (1, 2).

However, while most research on mAChRs in the central nervous system (CNS) has

focused on neuronal expression and function, much less is known about their roles in

non-neuronal cells.

Some of the strongest evidence for mAChRs function in glial cells comes from studies on M_1 mAChR in oligodendrocyte precursor cells (OPCs), where specific ablation of this receptor gene enhances remyelination in mouse models of autoimmune encephalomyelitis (3), hypoxia (4), aging (5) and Alzheimer's disease (6). Additional clues suggest that mAChRs may regulate neuroinflammation. For example, our studies in a prion model of neurodegeneration (7), and others in mouse models of Alzheimer's disease (8), have shown that the activation of the M_1 mAChR by selective positive allosteric modulators (PAMs) reduced markers of astrocyte and microglial activation. However, whilst transcripts for the mAChR family have been reported in astrocytes and microglia (9, 10), including the M_4 mAChR (11-13), little is known about the contribution that these receptors play in the regulation of glial cells.

Here, we used a transgenic mouse model expressing CRE recombinase under the control of the M_4 mAChR gene promoter (*Chrm4*) to discover a discrete population of M_4 mAChR-expressing astrocytes located in the brainstem, hypothalamus, and cerebellum. Signalling and proteomic studies confirmed the functional expression of this receptor subtype in astrocytes, particularly within the cerebellum. CRE-mediated deletion of M_4 mAChR expressing astrocytes exposed the role of this subpopulation of astrocytes in locomotion. Importantly, our study unveiled the expansion of the M_4 mAChR expressing astrocytes in parallel with the progression of prion neurodegenerative disease. Genetic ablation of these astrocytes significantly extended the lifespan of prion-diseased mice, suggesting a detrimental role in disease progression. Together our data point to a previously unappreciated subpopulation of M_4 mAChR expressing astrocytes that play a specific role in normal neurophysiology and inflammatory neurodegenerative disease.

Results

Generation of a genetic mouse model to identify M_4 mAChR expressing cells

To facilitate the study of the diversity and distribution of M_4 mAChR expressing cells (M_4^+ cells) in the brain, we generated a mouse line that expresses a CRE recombinase under the control of the *Chrm4*-promoter (Chrm4-CRE-IRES-mM4, hereinafter referred to as Chrm4^{CRE}) **(Fig. 1a)**. In these mice, the *Chrm4* locus was modified to contain an internal ribosomal entry site (IRES) cassette followed by the CRE recombinase coding sequence down-stream of the M_4 mAChR receptor exon 3 **(Fig. 1a)**. By crossing these mice with a CRE-dependent green fluorescent protein (GFP) reporter mouse strain (R26-CAG-GFP,-MetRS*^{L274G}, also called R26-MetRS*), we aimed to generate animals where the GFP acts as a marker of M_4 mAChR expressing cells (14) **(Fig. 1a)**. The resulting mouse strain (Chrm4^{CRE}::MetRS*), heterozygous for both alleles, showed similar levels of M_4 mAChR expression to that of wild-type mice as determined by radio-ligand binding experiments of striatal membranes **(Supplementary Fig. 1a)**. Examination of brain sections of Chrm4^{CRE}::MetRS* animals revealed GFP-positive neurons (stained with the neuronal marker MAP2) especially abundant in the striatum, cortex, and hippocampus **(Fig. 1b)**, areas where Western blots confirmed M_4 mAChR expression **(Supplementary Fig. 1b)** consistent with previous studies (15-18). These experiments, together with the lack of GFP signal detected in the brain of the parental lines **(Supplementary Fig. 1c)**, give confidence that the reporter-GFP system acts as a surrogate marker for M_4 mAChR expressing cells/neurons.

M_4 mAChRs expression in a discrete astrocytic subpopulation

In addition to M_4^+ neurons, we also detected dispersed populations of M_4^+ stellate cells

in the midbrain, hypothalamus, and the pons **(Fig. 1c)**. These cells did not co-localise

with markers for microglia, oligodendrocytes or neurons **(Supplementary Fig. 1d)** but

rather co-stained with antibody markers for glial fibrillary acidic protein (GFAP) and

$S100\beta$, thereby identifying them as M_4^+ astrocytes **(Fig. 1c, Supplementary Fig. 1e)**.

These experiments also revealed a surprisingly diverse array of M_4^+ cells in the

cerebellum **(Fig. 2a),** a region with low but detectable levels of M_4 mAChR transcript

(Supplementary Fig. 2a). In contrast, no significant GFP staining was observed in the

cerebellum of the parental mouse strains **(Supplementary Fig. 2b)**. Among these M_4^+

cells, we could identify a sparse population of M_4^+ Purkinje neurons, labelled with the

calcium-binding protein calbindin D-28k **(Fig. 2b)**. Most strikingly, however, was the

abundance of cerebellum M_4^+ cells that co-stained with astrocytic markers (GFAP,

NDRG2, $S100\beta$) **(Fig. 2c, Supplementary Fig. 2c)**. The location and morphology of these

cells indicated that they correspond to Bergmann glia in the Purkinje layer, some velate

astrocytes in the granule cell layer, dispersed fibrous astrocytes surrounding the white

matter and a large population of M_4^+ astrocytes within the cerebellar nuclei **(Fig. 2c).**

Signalling analysis of astrocyte M_4 mAChRs

To determine the functional expression of M_4 mAChRs in glial cells, primary astrocyte

cultures were prepared from the cerebrum and the cerebellum of Chrm4^{CRE}::MetRS*

mice. In line with the observations in intact brain, M_4^+ astrocytes represented only a

small proportion of the total astrocyte population prepared from the cerebrum (8.7 ±

2.4 %), whilst a higher proportion (30.4 ± 3.8 %) of M_4^+ astrocytes were evident in

cerebellar cultures **(Fig. 3a, b)**.

To establish if these M_4^+ astrocytes expressed functional M_4 mAChRs, we took

advantage of previous studies that showed that muscarinic agonists can induce

accumulation of the Immediate Early-Gene (IEG) c-fos in cultured astrocytes (19).

Therefore, we treated cerebellar astrocytes prepared from Chrm4[CRE]::MetRS* mice

with the muscarinic agonist oxotremorine (10 μM). This resulted in a significant up-

regulation of c-fos expression in M_4^+ astrocytes **(Fig. 3c, d)**. Furthermore, preincubation

with VU0467154 (VU154, 10 μM), a M_4 mAChR PAM (20), increased the number of

oxotremorine-induced c-fos$^+$ cells in the cerebellar astrocyte cultures, whilst

pretreatment with the highly selective M_4 mAChR antagonist VU6028418 (VU418, 5

μM) (21) prevented oxotremorine-induced stimulation of c-fos in these cells **(Fig. 3c,**

d). Additionally, we probed the phosphorylation of the extracellular signal-regulated

kinases 1/2 ($pERK_{1/2}$), a signalling pathway downstream of M_4 mAChR-Gi activation.

Treatment with oxotremorine increased $pERK_{1/2}$ levels in a concentration-dependent

manner (pEC_{50} = 5.57 ± 0.21, **Supplementary Fig. 3a**) whereas the M_4 mAChR

antagonist VU418 (1 μM) shifted the curve to higher agonist concentrations (pEC_{50} =

4.61 ± 0.22, p = 0.002) consistent with VU418 selective antagonism of the M_4 mAChR

$ERK_{1/2}$ response.

Proteomic evaluation of astrocyte M_4 mAChRs

To further characterise the functional M_4 mAChR response in cerebellar astrocytes, we

analysed changes in *de novo* synthesised proteins in response to receptor activation. In

these experiments we took advantage of the fact that, in addition to the expression of

GFP, the Chrm4[CRE]::MetRS* mice contain a mutated methionyl-tRNA synthetase (MetRS

L274G) expressed in a CRE-dependent manner. The enlarged binding pocket of this

mutant synthetase allows for the incorporation of the unnatural amino acid

azidonorleucine (ANL) into nascent proteins, which can later be tagged by click

chemistry for identification of labelled proteins or for biochemical purification (14).

Since our system results in CRE-expression only in M_4^+ cells (i.e. astrocytes) then the

mutated methionyl-tRNA synthetase will only be expressed in M_4^+ astrocytes. In this

way, although our cerebellar astrocyte cultures are mixed (~30% of the astrocytes are

M_4^+ **(Fig. 3b)**), we could conduct proteomic analysis on proteins synthesised exclusively

in M_4^+ astrocytes.

We first tested our system by exposing cerebellar astrocyte cultures, prepared from

Chrm4[CRE]::MetRS* mice, to ANL (4 mM) followed by fixation, permeabilization and

labelling ANL-incorporated proteins with the fluorescent Cy3-alkyne tag – process

referred to as fluorescent non-canonical amino acid tagging (FUNCAT). This established

a FUNCAT signal (Cy3) that was restricted to M_4^+ astrocytes (identified by GFP and

S100β) **(Supplementary Fig. 3b)**.

Next, we collected lysates from ANL treated cerebellar astrocytes prepared from

Chrm4[CRE]::MetRS* mice and using click chemistry incorporated a biotin-tag to ANL

incorporated proteins. Western blot analysis revealed an abundance of biotinylated

nascent proteins using this approach **(Supplementary Fig. 3c)**. These biotinylated

proteins were purified using avidin beads and proteins identified by mass spectrometry

(Supplementary Data 1). These experiments detected a total of 260 enriched

biotinylated proteins **(Fig. 3e)**. A gene ontology (GO) analysis revealed enrichment of

astrocytic markers (GFAP, vimentin, 3-phosphoglycerate dehydrogenase), confirming

the specific labelling. Importantly, terms associated with metabolic enzymes,

translation and cytoskeleton components were among the most significantly enriched

GO groups **(Fig. 3f)**. Furthermore, oxotremorine (10 μM) treatment induced the

synthesis of 67 unique proteins that were dependent on M_4 mAChR activation, as they

were not identified in the presence of the specific M_4 mAChR antagonist (VU418, 5

μM). A GO enrichment analysis showed that proteins up-regulated by M_4 mAChR

activation are involved in trafficking and protein translation **(Fig. 3g)**, including the

actin-binding profilin-2, the metallothionein subtypes II (MT2) and III (MT3), guanylate-

binding protein 2 (GBP2), and the MAP kinase ERK_2.

Diptheria toxin-mediated ablation of M_4^+ astrocytes effects locomotion

In order to decipher the role of M_4^+ astrocytes, we followed a genetic strategy of

targeted ablation. The mouse strain Aldh1l1-eGFP,-DTA (22), expresses eGFP in

astrocytes and a CRE-dependent diphteria toxin A (DTA). Crossing these animals with

the Chrm4^{CRE} strain results in a mouse (Chrm4^{CRE}:: Aldh1l1^{DTA}) where M_4^+ astrocytes are

selectively ablated **(Fig. 4a)**. As M_4 mAChR expression was observed in some

hypothalamic astrocytes **(Fig. 1b)**, we first evaluated the body weight of these animals

over time and found no differences compared to wild type littermates **(Fig. 4b)**. Also,

these animals showed similar food consumption and organ weights **(Supplementary**

Fig. 4a, b). Previous studies have indicated that alterations in cerebellar astrocytes,

specifically Bergmann glia, led to cerebellar atrophy and motor impairment (23).

However, we did not observe any abnormalities in the morphology or the overall glial

composition of the cerebellum of the Chrm4^{CRE}:: Aldh1l1^{DTA} animals when compared to

the cerebellum of the parental line **(Supplementary Fig. 4c)**. Furthermore, the

Chrm4^{CRE}:: Aldh1l1^{DTA} mice did not show ataxia or any deficits in motor coordination as

determined by the hindlimb clasping, ledge and rotarod tests **(Fig. 4c)**. In contrast,

exploratory behaviour evaluated in an open field test showed that Chrm4^{CRE}::

Aldh1l1^{DTA} mice exhibited significantly increased time in the centre of the arena

compared to littermate controls (74.1 ± 6.7 and 51.4 ± 2.6 s, respectively, $p < 0.01$, **Fig.**

4d), while the overall locomotion was reduced (24.1 ± 1.5 m travelled in the arena

versus 30.6 ± 1.7 m for control mice) and the time spent immobile (231.0 ± 13.7 s) was

greater than wild type littermates (164.7 ± 14.4 s) **(Fig. 4e, Supplementary Fig. 4d)**.

These results could be interpreted as deficits in anxiety-like behaviour, but a further

evaluation in an elevated plus maze test excluded this possibility, as Chrm4^{CRE}::

Aldh1l1^{DTA} mice performed similarly to wild type littermates in this paradigm **(Fig. 4e,**

Supplementary Fig. 4e).

These data suggest a potential role for M$_4^+$ astrocytes in locomotion which was tested

further by implant telemetry. Both Chrm4^{CRE}:: Aldh1l1^{DTA} and wild type littermate mice

had peaks of activity at similar times and of similar duration, excluding the possibility of

disturbed sleep **(Fig. 4g)**. Moreover, fluctuations in body temperature followed a

similar pattern, with no differences observed between the two groups of animals

(Supplementary Fig. 4f). Interestingly, when considering the overall activity in each

phase, it was evident that Chrm4^{CRE}:: Aldh1l1^{DTA} mice moved less intensely during the

dark phase (7.85 ± 0.61 counts) than their control littermates (9.93 ± 0.95 counts) **(Fig.**

4h). These findings are consistent with a deficit in the overall locomotor activity when

the M$_4^+$ astrocytes are ablated.

M$_4^+$ astrocytes do not regulate amphetamine hyperlocomotion

M_4 mAChRs are known to counteract hyperlocomotion induced by amphetamine or the

glutamate receptor agonist MK-801 (20, 24). This pivotal function in motor behaviours

has been attributed to neuronal (presynaptic) M_4 mAChR regulation of the dopamine

release and signalling in the basal ganglia (25-28). Here, we examined if M_4^+ astrocytes

might contribute to this by testing the efficacy of a M_4 mAChR selective PAM, VU154,

to reverse amphetamine-induced hyperlocomotion in Chrm4[CRE]:: Aldh1l1[DTA] and

littermate control mice. Administration of VU154 (10 mg/kg, i.p.) 20 mins before

amphetamine (2 mg/kg, i.p.) significantly decreased amphetamine-induced

hyperlocomotion in both groups of mice **(Fig. 4i, j)**. Likewise, both genotypes displayed

comparable measurements in mobility time **(Supplementary Fig. 4g)**. Together these

data indicate that the role of M_4^+ astrocytes in locomotion observed earlier is

independent from the dopamine-dependent motor function of M_4 mAChR in the basal

ganglia.

M_4^+ astrocytes during prion disease

There is currently intense interest in the role of astrocytes in regulating the onset and

progression of many neurodegenerative diseases and other brain disorders (29, 30). It

is clear, however, that the responses to pathological stimuli of reactive astrocytes are

very diverse, which could reflect the heterogeneous nature of astrocyte

subpopulations with established differences in origin and temporal function (31). Here

we probed the potential involvement of M_4^+ astrocytes in the pathology of murine

prion disease, a progressive terminal neurodegenerative disease model associated with

pronounced neuroinflammation and astrogliosis (32). Prion disease was induced in

Chrm4[CRE]::MetRS* mice by inoculation with Rocky Mountain Laboratory (RML) prion-

infected brain homogenate. At 20 weeks post-inoculation (w.p.i.), a sharp increase was

observed in the number of M_4^+ astrocytes **(Fig. 5a)**, as determined by co-staining with

the astrocytic marker GFAP **(Supplementary Fig. 5a)**. The lack of co-staining of these

M_4^+ cells with microglial marker Iba1 excluded the possibility that these cells were

infiltrating macrophages **(Supplementary Fig. 5b)**. In fact, the strong GFAP staining is

consistent with these M_4^+ cells showing characteristics of reactive astrocytes, which

was particularly evident in the region of the cerebellar nuclei **(Fig. 5b)**.

A   rostro-caudal   examination   of   serial   brain   sections   from   RML-inoculated

Chrm4^{CRE}::MetRS* mice demonstrated that M_4^+ astrocytes were profoundly up-

regulated in disease, even in regions such as the cortex, the hippocampus and the

thalamus where these cells were not evident under normal "non-disease" conditions

(Supplementary Fig. 5a, b, c). To interrogate regional differences, we estimated the

number of M_4^+ astrocytes in each area during prion disease and displayed the cellular

densities as heat maps overlaid onto the brain anatomy **(Fig. 5c)**. Our results indicated

that the hypothalamus hosts the largest number of M_4^+ astrocytes. As prion disease is

rapidly progressive, we further studied the evolution of these cellular subpopulations

by comparing three time points after RML-inoculation (15, 18 and 23 w.p.i., **Fig. 5c,**

Supplementary Fig. 5d). We found that the number of M_4^+ reactive astrocytes

gradually increased in almost all examined brain regions over time **(Fig. 5d)**.

Interestingly, this prion-induced expansion was not observed in one of the biggest

reservoirs of M_4^+ astrocytes, the cerebellar nuclei within the arbor vitae

(Supplementary Fig. 5e), although astrocytes in this region showed a prominent

reactivity during disease **(Fig. 5b)**. Critically, M_4^+ astrocytes showed expression of GBP2

during prion disease **(Fig. 5e)**, a marker associated with a neurotoxic or pro-

inflammatory phenotype (33). In addition to GBP2, these astrocytes also expressed

metallothionein III (MT3) **(Supplementary Fig. 5f)**, which was similarly identified in our

proteomic analysis as being upregulated upon activation of the M4 mAChR

(Supplementary Data 1). This spatio-temporal analysis of the M_4^+ astrocytes in prion

disease indicated that these cells become activated in a pathological context and the

number of reactive astrocytes that express the M_4 mAChR continuously increases

alongside the progression of the disease.

Increased lifespan of prion disease inoculated mice where M_4^+ astrocytes are ablated

To establish the potential role played by M_4^+ astrocytes in prion disease, we evaluated

the disease progression in mice where the M_4^+ astrocytes had been ablated.

Inoculation of both Chrm4^{CRE}:: Aldh1l1^{DTA} and wild type littermate mice with RML

resulted in a similar onset of prion disease and early/mid disease progression, as

determined by near equivalent times when at least two early "clinical" signs of prion

disease appeared **(Supplementary Fig. 6a)**. Similarly, there was equivalence in

progressive weight loss **(Supplementary Fig. 6b)**, decreased motor co-ordination and

decline in burrowing behaviour (a sign of hippocampal function) between the

genotypes **(Fig. 6a, b)**. Importantly, despite a similar progression of early and mid-

stages of disease, Chrm4^{CRE}:: Aldh1l1^{DTA} mice showed extended lifespan compared to

wild type littermates (mean survival 164.7 ± 0.9 and 158.6 ± 1.5 days, respectively, ****p

< 0.01) (Fig. 6c).** This survival effect was more prominent in females **(Fig. 6d)**, which

could be explained by their shorter incubation periods to Prp^{Sc} compared to males **(Fig.**

6e) when exposed to the same quantity of infectious material (34). Our result

emphasises the importance of sex as a biological variable in prion pathogenesis,

previously observed in mice depleted of NG2 glia (35) and in human sporadic

Creutzfeldt-Jakob disease (sCJD) (36).

We next assessed the accumulation of misfolded, insoluble scrapie prion protein

(Prp^{Sc}), which constitutes the major component of infectious prion disease. At the

terminal stage, a dramatic deposition of Prp^{Sc} was detected in the hippocampus,

cortex, striatum and cerebellum of prion-inoculated mice when compared to the non-

diseased controls **(Fig. 6f, g)**. Importantly, the accumulation of Prp^{Sc} was significantly

decreased in prion inoculated Chrm4^{CRE}:: Aldh1l1^{DTA} compared to wild type controls in

the cerebellum, but not in the rest of the brain regions **(Fig. 6g)**. However, this

reduction was not accompanied by changes in other indicators of neurodegeneration

(ApoE and serpinA3N) or markers of neuroinflammation (GFAP, Iba1 and clusterin)

(Supplementary Fig. 6c, d). Overall, these data indicate that the ablation of M$_4^+$

astrocytes does not affect the onset of the clinical symptoms or progression of prion

disease but extends the lifespan of mice, indicating a potential detrimental effect of

M$_4^+$ astrocytes in the later stages of disease.

Discussion

Despite the considerable progress in understanding the neuronal populations that

display the M$_4$ mAChR (37-39), the identity, location and function of non-neuronal CNS

cells, in particular glial cells, expressing this receptor subtype have remained largely

unexplored. Here, we reveal a discrete subpopulation of M$_4$ mAChR expressing

astrocytes dispersed across several areas of the mouse CNS. Genetic ablation of these

astrocytes demonstrated a role in neurological responses associated with locomotion,

while other M$_4$ mAChR-associated processes such as anxiety and the regulation of

responses to amphetamine remained unaffected. Importantly, there was a dramatic

expansion of this subpopulation of astrocytes during the progression of

neurodegeneration in a model of prion disease. Ablation studies indicated a

detrimental role for these M_4^+ astrocytes in later disease stages. In these ways our

study contributes to a growing awareness that astrocytes are not uniform but rather a

diverse set of cell types with varying cellular and functional characteristics.

Appreciating the nature and complexity of the various subtypes of astrocytes will be

essential if we are to understand their role in normal neurophysiology and if we are to

target these cells in brain diseases.

Under physiological conditions, we found M_4^+ astrocytes in the hypothalamus and

dorsal parts of the brainstem. These regions are critical regulators of energy balance

and appetite, and astrocytes in these areas have been reported to contribute to these

tasks by modulating the activity of adjacent neurons (40). Thus, manipulation of

hypothalamic and brainstem astrocytes has been reported to influence metabolic

processes including feeding and glucose metabolism (41-46), as well as circadian

rhythm, and sleep (47, 48). Interestingly, ablation of M_4^+ astrocytes in mice did not

result in alterations in the weight, body temperature or daily activity cycle. It was,

however, of relevance that M_4^+ astrocytes regulate locomotion, an activity consistent

with the expression of the receptor in a diverse array of the cerebellar astrocytes,

which are activated during voluntary and enforced locomotion (49, 50). It is important

to note that M_4^+ astrocytes are involved in specific aspects of motor control whilst

other behaviours such as balance and coordination, typically altered after a loss of

Bergmann glia support (23), remained normal in M_4^+ astrocyte-depleted mice.

Furthermore, the ablation of M_4^+ astrocytes had no impact on basal ganglia-mediated

hyperlocomotion responses, classically linked with neuronal M_4 mAChRs (25-28),

despite the fact that astrocytes in this region can modulate these psychomotor

behavioural effects (51). Hence, our studies have dissected a locomotion response that

may be under the regulation of M_4 mAChR expressed on a subpopulation of astrocytes.

While our study primarily identified the existence of a novel population of astrocytes,

the neuro-anatomical distribution of these cells, and the function that these astrocytes

might play under normal and neurodegenerative disease scenarios, it is a particular

challenge to relate this to the activity of M_4 mAChRs themselves. We started to

investigate this in primary cultures where we used a genetic approach that allows for

the signalling and proteomic interrogation of the M_4^+ cerebellar astrocytes amongst a

mixed population of astrocytes. This has provided an in-depth proteomic profile of M_4^+

astrocyte subpopulation and demonstrated that selective M_4 mAChR activation

resulted in the regulation of key cytoskeletal, metabolic and translational proteins that

can significantly impact astrocyte biology and activation status. For example, profilin-2

can modulate astrocytic morphology (52); MT3 plays a role in the clearance of the

prion-like amyloid β (53), and the low-density lipoprotein receptor-related protein-1

(LRP1) is important for the astrocyte-neuron communication (54, 55). Interestingly,

LRP1 is downregulated together with M_4 mAChR in the cerebellum of patients with

Rett syndrome (56), and mouse models show motor deficits when the *MeCP2* gene is

deleted from the cerebellum but not when is restricted to Purkinje or granular neurons

(57), highlighting the contribution of the cerebellar astrocyte pool.

The involvement of astrocytes, and indeed microglia, in protection against

neurodegenerative diseases is an extremely active area of research (58). Astrogliosis is

a widespread response in prion disease, although each brain region contains reactive

astrocytes with heterogenous phenotypes and functions determined by the

microenvironment (59). Our study showed a striking increase in M_4^+ astrocytes in the

thalamus and hypothalamus, brain regions identified as particularly susceptible to

prion deposition and neuroinflammatory cascades in RML and other prion strain

models (60, 61). Astrocytes are known to adopt either neuroprotective or neurotoxic

phenotypes in response to pathological stimuli. While their early activation may

support neuronal survival and clearance of misfolded proteins, functions that are

protective in diseases like Alzheimer's disease (62), the prolonged activation of

astrocytes is widely considered to be detrimental (63, 64). This certainly seems to be

the case for M_4^+ astrocytes which do not seem to make a significant impact on early-

and mid-stages of disease but rather are involved in acceleration of the later stages of

disease. The expression of GBP2 in M_4^+ astrocytes, a marker linked to neurotoxic

reactivity, further supports their detrimental role in late-stage disease. How the M_4

mAChR itself is involved in the regulation of these astrocytes at latter disease stages,

and whether agonists or antagonists might make an impact is at present uncertain.

What is clear, however, is that our findings support the selective targeting of neurotoxic

astrocytes while preserving glial subtypes with neuroprotective roles as a promising

therapeutic strategy for prion disease. A therapeutic option previously discarded after

observing that preventing the formation of the neurotoxic $C3^+$ astrocytes in mice

accelerated disease progression (33). This dichotomy highlights the delicate balance

between protective and harmful astrocyte states and the need for careful and

thorough characterisation of these astrocytic subpopulations.

In conclusion by identifying a previously unappreciated astrocyte subpopulation we not

only provide further evidence of the diversity of astrocytes but by ablation of M_4^+

astrocytes highlight the role of this astrocyte population in the regulation of

locomotion and neurodegenerative disease pathology.

Materials and Methods

Animals

Mice were maintained on a 12-h light–dark cycle with ad libitum standard mouse chow

food and water.

Generation of Chrm4^{CRE} mice

Transgenic Chrm4^{CRE} mice were generated by genOway (Lyon, France). Briefly, a CRE-

IRES cassette was inserted in frame of the ATG in exon 2 of the endogenous *Chrm4*

gene, by homologous recombination in embryonic stem (ES) cells. To this end a

targeting vector was created containing long and short homology arms to the *Chrm4*

genomic region. The targeted vector was completed with an FRT-flanked neomycin

resistance gene upstream the CRE sequence to select ES cells that had undergone

homologous recombination, and a diphtheria toxin cassette upstream the long

homology arm for negative selection. The linearised targeting construct was

transfected into C57BL/6N ES cells by electroporation. Positive clones were assessed by

PCR and whole locus sequence analysis to validate the presence of the correct

recombination event before being injected into blastocysts derived from C57BL/6J

female mice. The resulting highly chimeric males (chimerism rate above 50%) were

then crossed to a FLP recombinase-expressing deleter mouse strain to generate

heterozygous knock-in animals devoid of neomycin. Genotyping was initially confirmed

by PCR on genomic DNA obtained from ear biopsies using the primers 5'

CACTGCCCACCTGTCTCCAG and 5' GGTTGTGGGCTGTTGTGACCAG, which flank the

insertion site of the CRE-IRES cassette; and further confirmed with the pair of primers

5' GCTAGAGCCTGTTTTGCACG and 5' GCTAGAGCCTGTTTTGCACG, which amplify a

region of the CRE recombinase sequence. PCR-positive animals were further validated

by whole locus sequencing. The results were later corroborated by fluorescent

quantitative PCR (qPCR) using a commercial genotyping service (Transnetyx, Cordova,

TN). This service also analysed further samples to verify the genotype of the first litter

of every new breeder pair for the maintenance of the $Chrm4^{CRE}$ colony.

Homozygous $Chrm4^{CRE}$ were crossed with homozygous R26-CAG-GFP,-MetRS* (Jackson

Laboratory, strain: 028071) to generate $Chrm4^{CRE}$::MetRS*, and heterozygosity for both

alleles was confirmed by Transnetyx genotyping service. $Chrm4^{CRE}$ animals were mated

with hemizygous Aldh1l1-eGFP,-DTA mice (Jackson Laboratory, strain: 026033) to

produce $Chrm4^{CRE}$::Aldh1l1^{WT} and $Chrm4^{CRE}$:: Aldh1l1^{DTA}. All these animals were

genotyped with specific probes designed for CRE recombinase, eGFP and Chrm4

sequences at weaning by a commercial vendor (Transnetyx). $Chrm4^{CRE}$:: Aldh1l1^{DTA} and

wild type littermate animals were kept mixed in housing cages, except when measuring

the food and water intake and during telemetry monitoring experiments. The

generation of M_4 KO mice has been described elsewhere (65).

Measurement of organ weight, food and water consumption

$Chrm4^{CRE}$:: Aldh1l1^{DTA} and wild type littermate mice were housed separated by

genotype and food and water intakes were measured every 3 days for 1-2 weeks.

Average daily food and water consumption per animal were calculated and no

differences were observed between genotypes or sex.

The brain, liver, heart, spleen and kidneys from Chrm4^{CRE}:: Aldh1l1^{DTA} and wild type

littermate animals were collected (n=3-5, 4-5 months old) and the weight was recorded

and adjusted for body weight. Organ dimensions were also measured, and no

differences were observed (data not shown).

Rotarod

Motor performance was tested using the 5-lane rotarod (Cat. #47650, Ugo Basile,

Germany). Mice were weighed prior every testing, which happened weekly for both

prion infected and non-infected 10-week-old male mice for 15-20 weeks. Each day,

mice were placed on the 3 cm diameter rod for 2 habituation phases, consisting of a

constant rotation speed at 4 rpm for 90 s. For the testing phase, the animals were

placed on the rod that accelerated from 4 rpm to 40 rpm over a period of 5 min. The

mice were tested 2 times with a rest period of at least 15 min between each trial.

Animals that fell before the rod reached 15 rpm speed were placed back on the beam,

but if they fell more than once in that period, they were returned to the home cage. If

mice turned on the rod, the test was stopped, and the time recorded. The latency to

fall (time that the mouse remains on the rod) was registered automatically by a trip

switch under the floor of each rotating drum, and the result was used as an indicator of

motor coordination. The activity of Chrm4^{CRE}:: Aldh1l1^{WT} in this test was comparable to

wild type animals with a C57BL/6 background (data not shown).

Elevated Plus Maze (EPM)

The EPM test was used to assess anxiety-related behaviour in mice. 9-10 weeks old

male and female animals were placed in the centre of a cross shaped maze (arms of 50

x 10 cm), with 2 enclosed arms (black walls of 30 cm height) and 2 dimly illuminated

open arms opposite each other. The test was performed before the start of the light

cycle and lasted for 5 min. The time spent in the open arms versus the enclosed spaces

versus, the immobility time and the number of entries to each arm were recorded

using the ANY-maze video tracking software (Stoelting).

Open field

The open field activity of 8-12 weeks old male and female mice was monitored using a

clear Perspex square box (50 × 50 cm). Animals were placed in the centre of the

illuminated open field arena and allowed to freely explore for 10 minutes. The time

spent in the centre of the arena, the distance travelled and speed were assessed using

the ANY-maze tracking system (Stoelting).

Hindlimb clasping test

The degree of hindlimb clasping as an indicator of motor dysfunction was evaluated in

8-9 weeks old mice. Animals were suspended by their tails for 10-20 s and rated from 0

to 3 according to the following score system: 0 corresponded to hindlimbs

consistently splayed outward and away from the abdomen; 1 was assigned when

one hindlimb was retracted inwards towards the abdomen for more than 50% of the

time; a score of 2 meant that both hindlimbs were partially retracted inwards towards

the abdomen for at least 50% the time; and 3 was given when both hindlimbs were

entirely retracted inwards towards the abdomen for at least 50% of the observation

period (66). Two separate trials were performed per animal. All evaluated Chrm4^{CRE}::

Aldh1l1^{DTA} and Chrm4^{CRE}:: Aldh1l1^{WT} animals showed a consistent score of 0 in this test.

Ledge test

Mice aged 8-9 weeks old were placed on the cage ledge and recorded to measure

balance and coordination. Paw placement and forward movement as it walked along

the ledge were observed and scored from 0 to 3 as in (66): 0, for normal walking; 1, if

the mouse lost its footing; 2, if the hindlimbs were not used, and 3, if the animal fell off

the ledge. Data from three trials were averaged for each mouse and none of the

Chrm4^{CRE}:: Aldh1l1^{DTA} and Chrm4^{CRE}:: Aldh1l1^{WT} mice presented any deficits in balance

and coordination.

Telemetry

A TA-F10 temperature and activity probe (Data Sciences International) was implanted

in the abdominal cavity of 9-13 weeks old male and female Chrm4^{CRE}:: Aldh1l1^{DTA} and

wild type littermate mice (n=8, 19-30 g), using aseptic techniques under a surgical level

of isoflurane anaesthesia. Carprofen (10 mg/kg) and buprenorphine (0.1 mg/kg) were

s.c. administered 30 min before surgery for pain relief. After surgery, mice were housed

individually, and post-operative analgesia was given 6 h (buprenorphine) and 24 h later

(carprofen and buprenorphine). One week after surgery, when the animals completely

recovered, the cages were placed over the receiver panels and the TA-F10 transmitters

were activated. Activity and body temperature data were sampled every 1 min for 10 s

and continuously acquired during 4 consecutive days using the Ponemah software

(Data Sciences International). The measurements from the final 3 days were averaged

together for each mouse. No differences were observed between males and females.

Amphetamine-induced hyperlocomotion

Male and female Chrm4^{CRE}:: Aldh1l1^{DTA} and wild type littermate mice (n=6-7, 10-11

weeks old) were placed in an open field arena and the animals were tracked with the

ANY-maze software (Stoelting). Habituation lasted for 20 min before being i.p. injected

with vehicle (10% Tween 80 in sterile water) or VU0467154 (10 mg/kg). A dose of D-

amphetamine (2 mg/kg in saline, i.p.) was administered 20 min later, and locomotor

activity was recorded for an additional 90 min. The mobility and distance travelled over

time were expressed and averaged per 5 min bins. Total activity data were calculated

as the total distance travelled 60 min after the time of D-amphetamine administration.

Burrowing

The voluntary behaviour of removal material from an enclosed tube was assessed in

control and prion-infected Chrm4^{CRE}:: Aldh1l1^{DTA} and wild type littermate female mice

every week from 9 w.p.i. until survival time. Mice were placed into individual cages (22

× 36 cm) with an acrylic clear tube filled with food pellets for 2 h. The burrowing

activity was calculated by subtracting the weight of the remaining pellets at the end of

the experiment from the starting weight (140 g). Mice were acclimatised to the

burrowing cage the week before the start of the test.

Prion mouse model

Male and female Chrm4^{CRE}::MetRS*, Chrm4^{CRE}:: Aldh1l1^{DTA} and Chrm4^{CRE}:: Aldh1l1^{WT}

mice aged 3-4 weeks old were i.c. inoculated, under a surgical level of isoflurane

anaesthesia, into the right brain parietal lobe with 30 µl of either 1% NBH (Normal

Brain Homogenate) or 1% RML prion (Rocky Mountain Laboratories), as previously

described (67, 68).

Evaluation of prion disease progression and survival

Chrm4^{CRE}:: Aldh1l1^{DTA} and Chrm4^{CRE}:: Aldh1l1^{WT} control and prion-infected mice were

co-housed in mixed-genotype cages to minimise environmental and housing-related

bias. Clinical assessment of prion disease progression was performed daily by a trained

technician who was blinded to the genotype of the animals, and the scoring protocol

followed established criteria consistent with previously published methodologies (7,

68, 69). Disease progression was evaluated according to the presence of early indicator

and confirmatory signs of prion disease. Early indicator signs included piloerection,

mild loss of coordination, erect penis, clasping of hindlimbs when lifted by tail,

intermittent generalised tremors, sustained erect ears, rigid tail, discontinued hunched

posture, or subdued behaviour. Confirmatory signs of prion disease included righting

reflex impairment, sustained hunched posture, dragging of limbs, and/or significant

abnormal breathing. Animals were humanely killed when they showed two

confirmatory signs or two early indicators plus one confirmatory sign, and the days

after inoculation were considered as the survival time.

Preparation of protein extracts and immunoblot analysis

Mice were culled via cervical dislocation, and different regions of the brain were

dissected and snap frozen on dry ice. Samples were lysed in RIPA buffer (10 mM Tris-Cl,

1 mM EDTA, 1% Triton X-100, 0.1% sodium deoxycholate, 0.1% SDS, 140 mM NaCl, pH

7.5) supplemented with protease inhibitor tablets (Cat. #11697498001, Roche) and

incubated for 30 min at 4 °C. Lysates were then briefly sonicated (Cat. #12337338,

Fisherbrand) with 2 pulses of 15 s at an amplitude of 40% (~50 W) until fully dissolved.

Samples were then centrifuged at 10,000 × g for 10 min, supernatants collected, and

protein concentration calculated using a BCA assay kit (Cat. #23225, Pierce). Equal

concentrations of protein samples were resuspended in Laemmli loading buffer (10%

SDS, 300 mM Tris-HCl pH 7.2, 0.05% bromophenol blue, 10% β-mercaptoethanol) and

heated at 60°C for 10 min. For the assessment of PrP^{Sc} levels, 60 µg of each lysate were

treated with proteinase K (10 µg/ml, Cat. P2308, Sigma-Aldrich) for 10 min at 37°C. The

reaction was stopped by the addition of Laemmli sample buffer and boiling at 95°C for

5 min. 20-40 µg of each cell lysate were resolved on NuPAGE precast 4-12% gels (Cat.

#NP0321, Invitrogen) followed by transfer onto nitrocellulose membranes (Cat.

#88018, Thermo Scientific). Protein transfer was verified with Ponceau-S staining (0.2%

Ponceau-S red, 1% acetic acid), before membranes were blocked with 5% non-fat dry

milk in TBST (25 mM Tris-HCl; pH 7.6, 100 mM NaCl, 0.5% Tween 20) for 1 h at room

temperature. Membranes were incubated with the corresponding primary antibodies

in the blocking solution overnight at 4°C. After washing three times for 5 min each with

TBST, secondary antibodies were added to the membranes for 1 h at RT. Antibodies

used in this study are detailed in Tables 1 and 2. Immunoreactive bands were visualised

using the Licor Odyssey system and densitometry of fluorescence was measured using

Image Studio Lite (Licor Biosciences).

Astrocyte cultures

Murine astrocyte cultures were performed according to the AWESAM protocol (70)

with some modifications. Brains from Chrm4^{CRE}::MetRS* or WT pups (P0-P5) were

collected and immediately placed in ice cold HBSS (ThermoFisher Scientific). After

removing of meninges, the cerebellum or the cerebrum were dissected and incubated

with 0.05% trypsin-EDTA (Cat. #25300054, ThermoFisher Scientific) at 37°C for 10 min

followed by gentle mechanical dissociation. Cells were centrifuged at 1,000 x g for 3

min, resuspended in high glucose, Glutamax-supplemented DMEM (Cat. #61965026,

Gibco) with 10% Foetal Bovine Serum (Cat. #10500064, Gibco), 100 units/ml penicillin

and 100 μg/ml streptomycin (Cat. # 15070063, Gibco), and plated on flasks precoated

with poly-D-lysine (100 μg/ml, Cat. #P0899, Sigma-Aldrich). After 7 days in vitro (DIV),

the flasks were shaken at 180 rpm for 30 min at 37°C, media changed and further

shaken at 240 rpm for 6 h. Astrocytes were detached and seeded at a density of 0.2 x

10^6 cells/ml on the corresponding experimental poly-D-lysine plates. After a couple of

days, media was replaced with NB+H media, containing Neurobasal (Cat. # 21103049,

Gibco) supplemented with B-27 (Cat. #17504044, Gibco), 2 mM Glutamax (Cat. #

35050061, Gibco), 10 ng/mL heparin-binding EGF-like growth factor (Cat. #E4643,

Sigma-Aldrich), 100 units/ml penicillin and 100 μg/ml streptomycin for 5 days before

the start of the experiments.

ERK$_{1/2}$ phosphorylation assay

Cultured primary astrocytes from WT mice were seeded onto 96 well plates precoated

with poly-D-lysine at a density of 20,000 cells/well in NB+H media. One week later, the

media was replaced, and cells were incubated with 80 μl of Neurobasal without

additives for 4 h. Cells were treated for 10 min with vehicle or 10 μM of the M$_4$ mAChR

antagonist VU6028418 before stimulation with the indicated concentrations of

oxotremorine for 5 min at 37°C in a final volume of 100 μl. Maximum ERK$_{1/2}$

phosphorylation response was obtained by treating the cells with FBS. ERK

phosphorylation was evaluated using a Homogenous Time-Resolved FRET (HTRF)-

based phospho-ERK (Thr202/Tyr204) cellular kit (Cat. #64ERKPEH, Revvity). Briefly, lysis

buffer was added immediately after the end of the stimulation and incubated for 30

min. After homogenisation, 16 µl of the lysate were transferred to a 384-well white

ProxiPlate (PerkinElmer) and combined with 4 µl HTRF premixed antibodies. for 4 h at

room temperature. Fluorescence emission at 665 and 620 nm was measured after 4 h

with a PHERAstar plate reader (BMG Labtech) and results were normalised to the

response stimulated by FBS.

Immunocytochemistry analysis

Cerebral or cerebellar primary astrocytes from Chrm4[CRE]::MetRS* mice were grown

onto 24 well plates with 13 mm glass coverslips previously treated with poly-D-lysine as

indicated above. Cells were fixed with 4% formaldehyde for 1 h at room temperature.

Coverslips were then washed 3 times with PBS and blocked with PBS supplemented

with 1% BSA and 0.2% Triton X-100 for 1 h. Subsequently, astrocytes were incubated

with the indicated primary antibodies diluted in blocking solution overnight at 4°C in a

humidity chamber. Detection was achieved using appropriate secondary antibodies

conjugated to Alexa Fluor 488 or Alexa Fluor 594 (1:500, Life Technologies) and

coverslips were mounted on glass slides using Fluoromount-G with DAPI (ThermoFisher

Scientific). Confocal images were acquired with a Zeiss Pascal laser-scanning confocal

microscope (LSM) 880 using the LSM image acquisition software.

For analysis of c-fos expression, the NB+H media was replaced with non-supplemented

Neurobasal, and cells were incubated at 37°C and 5% CO_2 for 4 h, before being

stimulated with vehicle or 10 µM oxotremorine for 60 min. To evaluate M_4 mAChR

action, cultures were simultaneously treated with vehicle, 10 µM VU0467154 or 5 µM

VU6028418. Immunocytochemistry was performed as described above and, for each

replicate (n = 3), 8 confocal images (425 μm x 425 μm) per condition were capture,

randomly selected using DAPI staining from areas with equivalent number of cells. The

images were analysed using ImageJ Fiji (NIH Image). Automatic quantification of c-fos$^+$

cells was performed using the "Analyze particle" plugin. M_4 mAChR-expressing

astrocytes were manually counted and M_4^+/c-fos$^+$ double stained cells were noted.

Total number of cells were also counted automatically with the software tools. The

experimental conditions were blinded to the investigator performing the image

analysis. This methodology was also used to calculate the proportion of M_4^+ astrocytes

present in cerebral and cerebellar cultures.

For FUNCAT analysis, Chrm4^{CRE}::MetRS* astrocytes were incubated with 4 mM L-

Azidonorleucine hydrochloride (ANL, Tocris) for 24 h. Cells were fixed and blocked as

previously described, followed by click chemistry to visualize the ANL-labelled newly

synthesized proteins. The Cu(I)-catalysed azide/alkyne cycloaddition (CuAAC) reaction

comprised the addition of 0.45 mM $CuSO_4$, 0.9 mM Tris((1-hydroxy-propyl-1H-1,2,3-

triazol-4-yl) methyl) amine (THPTA, Cat. #CLK-1010, Jena Bioscience), 0.2 mM Cy3-

alkyne (Cat. #777331, Sigma-Aldrich) and 5 mM Sodium ascorbate to the cells for 24 h.

Coverslips were washed with PBS containing 0.5 mM EDTA for 10 min, followed by 3

washes with blocking solution and the remaining steps of the immunocytochemistry

proceeded as indicated above.

Immunohistochemistry

Mice were deeply anaesthetised with isofluorane and then they were transcardially

perfused with 20 ml saline solution followed by 20 ml 4% formaldehyde in PBS. Brains

were removed and incubated overnight in the same fixative. The whole brain was

coronally or sagittally sectioned (50 µm) using a VT1200S vibratome (Leica). After

blocking the free-floating sections with PBS supplemented with 0.5% Triton X-100 and

1% bovine serum albumin for 1 h at room temperature, they were then incubated with

the indicated primary antibodies diluted in blocking solution 24 to 48 h at 4°C.

Antibodies used in this study are listed in Tables 1 and 2. Brain sections were washed 3

times with PBS, followed by incubation with the corresponding secondary antibodies

diluted in blocking solution for 4 h at room temperature. Sections were placed onto 2%

gelatine-coated slides and mounted with Fluoromount-G with DAPI. Images were taken

with a Zeiss LSM880 confocal microscope using the LSM image acquisition software

Zen lite.

Cell counting

The right brain hemisphere of each mouse was sliced in 50 µm coronal sections that

were sequentially collected in series containing one every 8[th] slice. Randomly chosen

series of sections were used for immunohistochemistry against GFP and GFAP as

described above. Due to the low numbers of M_4^+ astrocytes in the brain, we counted all

visible cells positive for both GFP and GFAP within the indicated brain regions in every

section of the series under a Zeiss microscope. DAPI images of each section were taken

and the area of each region of interest was traced using the freehand drawing tool of

ImageJ Fiji. The determined area was then multiplied by the thickness of the section to

calculate the reference volume. The density of M_4^+ astrocytes (number of cells/mm^3)

was estimated by dividing the total number of positive cells by the reference volume of

the corresponding region.

The quantification of M_4^+ astrocytes in the arbor vitae of the cerebellum was estimated

by unbiased stereology using the physical dissector method adapted to confocal

microscopy, as previously described (71). One in eight coronal sections of the

cerebellum (50 μm) were immunostained for GFP and GFAP as before. 9-12 confocal

counting stacks, acquired with a LSM880 Zeiss microscope, were randomly selected

from the whole arbor vitae of the cerebellum. The counting frame size was 150 × 150

μm with a thickness of 10.5 μm and a Z-interval of 1.5 μm. The mean cell density of

M_4^+ astrocytes was estimated with the counts of the double-labelled cells and the size

of the dissector.

Saturation [^{3}H]-NMS binding assay

Striata from homozygous M_4 KO, R26-MetRS* (WT) or heterozygous Chrm4^{CRE}::MetRS*

were collected and homogenised in ice-cold HEPES buffer (10 mM HEPES, 1 mM EGTA,

1 mM dithiothreitol) containing 10% sucrose and protease inhibitors. The lysate was

first centrifuged at 1,000 x g for 10 min at 4°C, the pellet resuspended, re-homogenised

and further centrifuged at 20,000 x g for 1 h at 4°C. The resulting pellet containing

striatal membranes, was diluted and protein concentration calculated by the Bradford

method. 22.5 μg of the striatal membrane preparation were incubated in binding

buffer (50 mM HEPEs, 110 mM NaCl, 5.4 mM KCl, 1.8 mM CaCl$_2$, 1 mM MgSO$_4$, 25 mM

glucose, 58 mM sucrose, pH 7.4) with increasing concentrations (0.3-5 nM) of the

muscarinic receptor antagonist [^{3}H]-N-methyl scopolamine ([^{3}H]-NMS, Cat.

#NET636001MC, PerkinElmer) for 1 h at 37°C. Nonspecific binding was determined by

the addition of atropine (1 μM) during the incubation with [^{3}H]-NMS. Binding reactions

were terminated by rapid filtration over UniFilter GF/C plates (PerkinElmer) pre-soaked

in binding buffer, followed by three rapid washes with ice-cold 0.9% NaCl using a filter

plate system (Brandel). The plates were then air-dried and 50 µL Microscint-20 liquid

scintillation fluid (PerkinElmer) was added to each well. Membrane bound radioactivity

was then counted in a Topcount liquid scintillation counter (PerkinElmer). The specific

bound counts (dpm) were expressed as fmol/mg protein.

Quantitative RT-PCR

Striata or cerebella from homozygous M_4 KO and R26-MetRS* (WT) mice (10-12 weeks

old) were collected and total RNA was isolated using a RNeasy Plus Mini Kit (Cat.

#74134, Qiagen), following the manufacturer's instructions. 1 µg total RNA was treated

with DNase I (Cat. #79254, Qiagen) and used as template for cDNA synthesis with M-

MLV reverse transcriptase (Cat. #28025013, Invitrogen), according to the

manufacturer's recommendations. Real-time PCR was performed with probes for

Chrm4   (forward   5'-GGAGAAGAAGGCCAAGACTCTGG   and   reverse   5'-

GGCAGTCACACATTCACTGCCTG)   and   GAPDH   as   control   gene   (forward   5'-

CGGATTTGGCCGTATTGGG and reverse 5'-CTCGCTCCTGGAAGATGG), 2.4 µl of the cDNA

sample as a template and Fast SYBR Green Master Mix (Cat. #4385612, Applied

Biosystems) on a Quantstudio 5 Real Time PCR system (ThermoFisher Scientific). The

cycling conditions were 50°C for 2 min, 95°C for 10 min, followed by 45 cycles of 95°C

for 15 s and 60°C for 1 min. Melting curve analyses were performed for all real-time

PCR runs to confirm amplification of only one product and to eliminate false positive

results. Quantitative analysis was performed on Quantstudio software (ThermoFisher

Scientific), which calculated the threshold cycle (Ct) values. Expression of Chrm4 was

defined relative to GAPDH using the $2-\Delta\Delta Ct$ method.

687   BONCAT purification for LC-MS/MS analysis

Cerebellar primary astrocytes from Chrm4[CRE]::MetRS* mice grown onto 10 cm dishes

as before, were incubated with 4 mM ANL in combination with vehicle, oxotremorine

(10 μM) or oxotremorine in combination with VU6028418 (5 μM) for 24 h at 37°C. Cells

were lysed in PBS with 0.5% SDS, 0.2% Triton X-100 and proteinase inhibitors, briefly

sonicated with 2 pulses of 15 s at an amplitude of 40% (~50 W) and centrifuged at

10,000 x g for 10 min at 4°C. Supernatants were collected and protein concentration

calculated with a BCA assay kit. 500 μg of protein were heated at 65°C for 5 min

followed by incubation with 50 mM iodoacetamide for 2 h at room temperature in the

dark. Bio-orthogonal non-canonical amino acid tagging (BONCAT) was performed using

150 μM Biotin-PEG4-alkyne (Cat. #764213, Sigma-Aldrich), 0.5 mM $CuSO_4$, 1 mM

THPTA and 5 mM sodium ascorbate overnight at 4°C. Buffer exchange was done with

PD SpinTrap G-25 columns (Cat. #28-9180-04, Cytvia) and biotinylation of ANL-labelled

proteins was confirmed by Western Blot. Ultralink avidin beads (Cat. #53146, Pierce)

were used to purify clicked proteins. After overnight incubation at 4°C, beads were

loaded into micro-spin columns (Cat. #89879, Pierce) and washed three times with PBS

and three washes with ultrapure water. Proteins were eluted with 125 μl of 0.1%

Trifluoroacetic acid (TFA) followed by 125 μl of 0.1% TFA with 50% acetonitrile (ACN).

Eluted proteins were digested with 4 μg Trypsin/Lys-C (Cat. #V5073, Promega) in S-Trap

midi spin columns (Profiti) according to the manufacturer's instructions.

Dried peptides were resuspended in 5 % acetonitrile with 0.5 % formic acid for

nanoflow liquid chromatography–electrospray ionisation-tandem mass spectrometry

(nLC-ESI-MS/MS) analysis. The samples were loaded onto a C_{18} Acclaim PepMap

trapping column (ThermoFisher Scientific) using an UltiMate3000 RSLCnano System

(ThermoFisher Scientific). Peptide separation was performed on a Pepmap C_{18} reversed

phase column (2 μm, 75 μm × 50 cm, pore size 100 Å, Cat. #164540, ThermoFisher

Scientific) by a gradient formed by mobile phase A (0.1% formic acid) and B (80% ACN,

0.08% formic acid), with increasing concentrations of solvent B running from 4% for 10

min, 60% for 170 min to 99% for 15 min at a fixed solvent flow rate of 0.3 μl/min.

Eluted peptides were electrosprayed into an Orbitrap Elite MS (ThermoFisher Scientific)

using a NanoMate Triversa (Advion Biosciences) ion source at 1.7 kV. The mass

spectrometer was operated using full-scan MS spectra (m/z 375–1,500) at a resolution

of 120,000 RP, followed by top speed (3s) high energy collision dissociation (HCD)

fragmentation detection of the top precursor ions. Ion fragmentation was performed

by top speed (3s) high energy collision dissociation (HCD) and the top precursor ions

were detected at a resolution of 60,000 in MS2 spectra. HCD energy and isolation

width was adjusted depending on charge of the precursor ions (a width of 1.2 m/z with

40% HCD for 2^+ ions, 0.7 m/z with 38% for 3^+, and 0.5 m/z with 38% HCD for 4^+-6^+). The

resulting MS/MS spectra were analysed for protein identification using Proteome

Discoverer software 2.4 (ThermoFisher Scientific) and searched against the murine

UniProtKB/Swiss-Prot database using Mascot software (Matrix Science) with peptide

tolerance set to 10 ppm and fragment ion mass tolerance of 0.3 Da. The

carbamidomethylation of cysteine was programmed as fixed modification, while the

oxidation of methionine, the deamidation of asparagine and glutamine, the

replacement of methionine for ANL-PEG-biotin and the iodination of tyrosine were set

as variables. Proteins present in at least 2 independent experiments were considered

for GO enrichment analysis to identify overrepresented biological processes using

PANTHER (72) with a Fisher's exact test and a cutoff of FDR ≤0.05.

Statistical analysis

All data were expressed as mean ± SEM of at least 3 independent experiments.

Statistical significance was determined by two-tailed Student's t test, one-way ANOVA

with Tukey's post hoc correction or two-way ANOVA followed by a Tukey's or

Bonferroni's post hoc test, as indicated. For $ERK_{1/2}$ phosphorylation assays the

concentration–response data were plotted on a log axis, fitted to a three-parameter

sigmoidal curve and extra sum-of-squares F test applied to determine whether

differences in pEC_{50} existed. The Kaplan-Meier plot survival analyses were compared

using the Gehan-Breslow-Wilcoxon test. Significance was considered a p value smaller

than 0.05 (*p < 0.05, **p < 0.01, ***p < 0.001). Statistical analysis and graph plotting

was performed using GraphPad Prism software.

Table 1: Primary antibodies. ICC, immunocytochemistry; IHC, immunohistochemistry;

WB, western blot.

Antibody	Supplier	Cat. No	Host	Dilution	Use
S100β	Proteintech	15146-1-AP	Rabbit	1:500	ICC
c-fos	Abcam	ab190289	Rabbit	1:1,000	ICC
GFAP	Abcam	ab7260	Rabbit	1:500	ICC
GFP	Abcam	ab6673	Goat	1:500	ICC
S100β	Proteintech	15146-1-AP	Rabbit	1:1,000	IHC
Calbindin-D28k	Proteintech	14479-1-AP	Rabbit	1:1,000	IHC
GBP2	Proteintech	11854-1-AP	Rabbit	1:500	IHC
GFAP	Abcam	ab7260	Rabbit	1:1,000	IHC
GFAP	Sigma-Aldrich	G3893	Mouse	1:2,000	IHC
GFP	Abcam	ab6673	Goat	1:1,000	IHC
GFP	Abcam	Ab6556	Rabbit	1:1,000	IHC

Iba1	Abcam	ab178846	Rabbit	1:1,000	IHC
MAP2	Biolegend	822501	Chicken	1:1,000	IHC
MT3	Proteintech	12179-1-AP	Rabbit	1:500	IHC
Myelin basic protein (MBP)	Proteintech	10458-1-AP	Rabbit	1:2,000	IHC
NDRG2	Abcam	ab174850	Rabbit	1:1,000	IHC
Apolipoprotein E (APO-E)	Abcam	ab183596	Rabbit	1:1,000	WB
Chrm4	Aviva	OASG0431	Rabbit	1:500	WB
Clusterin	R&D Systems	AF2747	Goat	1:2,000	WB
GFAP	Sigma-Aldrich	G3893	Mouse	1:1,000	WB
GFP	Invitrogen	A-11122	Rabbit	1:1,000	WB
Iba1	Abcam	ab178846	Rabbit	1:2,000	WB
IRDye 800CW Streptavidin	Li-Cor Biosciences	926-32230	-	1:5,000	WB
Myelin basic protein (MBP)	Proteintech	10458-1-AP	Rabbit	1:4,000	WB
Prion protein (Prp)	Abcam	ab61409	Mouse	1:1,000	WB
SerpinA3N	R&D Systems	AF4709	Goat	1:1,000	WB
Vimentin	R&D Systems	MAB21052	Mouse	1:2,000	WB
α-tubulin	Abcam	ab7291	Mouse	1:10,000	WB

748 **Table 2: Secondary antibodies.** ICC, immunocytochemistry; IHC,

749 immunohistochemistry; WB, western blot.

Antibody	Supplier	Cat. No	Host	Dilution	Use
Anti-Rabbit Alexa Fluor Plus 647	Invitrogen	A32795	Donkey	1:500	ICC
Anti-Chicken Alexa Fluor 594	Invitrogen	A11042	Goat	1:1,000	IHC
Anti-Rabbit Alexa Fluor 488	Invitrogen	A32731	Goat	1:1,000	IHC
Anti-Goat Alexa Fluor Plus 488	Invitrogen	A32814	Donkey	1:1,000/ 1:500	IHC/ ICC
Anti-Mouse Alexa Fluor Plus 594	Invitrogen	A32744	Donkey	1:1,000/ 1:500	IHC/ ICC
Anti-Rabbit Alexa Fluor Plus 594	Invitrogen	A32754	Donkey	1:1,000/ 1:500	IHC/ ICC
IRDye 680RD Anti-Goat	Li-Cor Biosciences	925-68074	Donkey	1:5,000	WB

IRDye 680RD Anti-Mouse	Li-Cor Biosciences	925-68072	Donkey	1:5,000	WB/ IHC
IRDye 800CW Anti-Mouse	Li-Cor Biosciences	926-32212	Donkey	1:5,000	WB
IRDye 800CW Anti-Rabbit	Li-Cor Biosciences	925-32213	Donkey	1:5,000	WB

Abbreviations

CNS: central nervous system, $ERK_{1/2}$: extracellular signal-regulated kinases 1/2, GFAP:

glial fibrillary acidic protein, GFP: green fluorescent protein, eGFP: enhanced green

fluorescent protein, GPCR: G protein-coupled receptor, mAChR: muscarinic

acetylcholine receptor, NBH: normal brain homogenate, PAM: positive allosteric

modulator, pEC_{50}: half maximal response, PrP^{Sc}: scrapie prion protein, RML: rocky

mountain laboratory, w.p.i: weeks post-inoculation.

Declarations

Ethics approval

760 All animal experiments were conducted in accordance with the UK Home Office Animal

761 Procedures Act (1986) and all procedures were approved by the University of Glasgow

762 Animal Welfare and Ethical Review Board.

Consent for publication

764 Not applicable

Availability of data and materials

766 Source data are provided with this paper.

Competing Interests

The authors declare no competing interests.

Funding

This work was supported by grants from the Wellcome Trust (Collaborative Award 201529/Z/16/Z and Discovery Award 307319/Z/23/Z).

Author's Contributions

GST conducted the experiments, developed the study design and wrote the paper, ABT lead the study design and wrote the paper, GM contributed to the study design and writing of the paper, CM managed the mouse colonies and breeding and helped to conduct the animal experiments, CWL contributed with the muscarinic antagonist VU6028418 and the M_4 mAChR positive allosteric modulator VU0467154 and CKJ contributed with pharmacokinetic, pharmacology and metabolism data of these ligands.

Acknowledgments

The authors would like to thank the staff of the University of Glasgow Biomedical Services, genOway for generation of the muscarinic receptor genetically modified mice. The authors would also like to acknowledge the assistance of Dr. Karen Fairlie-Clarke to this work.

References

1.	Kaul I, Sawchak S, Correll CU, Kakar R, Breier A, Zhu H, et al. Efficacy and safety of the muscarinic receptor agonist KarXT (xanomeline-trospium) in schizophrenia

(EMERGENT-2) in the USA: results from a randomised, double-blind, placebo-
controlled, flexible-dose phase 3 trial. Lancet. 2024;403(10422):160-70.

2.   Kaul I, Sawchak S, Walling DP, Tamminga CA, Breier A, Zhu H, et al. Efficacy and Safety
of Xanomeline-Trospium Chloride in Schizophrenia: A Randomized Clinical Trial. JAMA
Psychiatry. 2024;81(8):749-56.

3.   Mei F, Lehmann-Horn K, Shen YA, Rankin KA, Stebbins KJ, Lorrain DS, et al. Accelerated
remyelination during inflammatory demyelination prevents axonal loss and improves
functional recovery. Elife. 2016;5.

4.   Wang F, Yang YJ, Yang N, Chen XJ, Huang NX, Zhang J, et al. Enhancing Oligodendrocyte
Myelination Rescues Synaptic Loss and Improves Functional Recovery after Chronic
Hypoxia. Neuron. 2018;99(4):689-701 e5.

5.   Wang F, Ren SY, Chen JF, Liu K, Li RX, Li ZF, et al. Myelin degeneration and diminished
myelin renewal contribute to age-related deficits in memory. Nat Neurosci.
2020;23(4):481-6.

6.   Chen JF, Liu K, Hu B, Li RR, Xin W, Chen H, et al. Enhancing myelin renewal reverses
cognitive dysfunction in a murine model of Alzheimer's disease. Neuron.
2021;109(14):2292-307 e5.

7.   Dwomoh L, Rossi M, Scarpa M, Khajehali E, Molloy C, Herzyk P, et al. M(1) muscarinic
receptor activation reduces the molecular pathology and slows the progression of
prion-mediated neurodegenerative disease. Sci Signal. 2022;15(760):eabm3720.

8.   Abd-Elrahman KS, Colson TL, Sarasija S, Ferguson SSG. A M1 muscarinic acetylcholine
receptor-specific positive allosteric modulator VU0486846 reduces neurogliosis in
female Alzheimer's mice. Biomed Pharmacother. 2024;173:116388.

9.   Pannell M, Meier MA, Szulzewsky F, Matyash V, Endres M, Kronenberg G, et al. The
subpopulation of microglia expressing functional muscarinic acetylcholine receptors
expands in stroke and Alzheimer's disease. Brain Struct Funct. 2016;221(2):1157-72.

10.   Andre C, Dos Santos G, Koulakoff A. Muscarinic receptor profiles of mouse brain
astrocytes in culture vary with their tissue of origin but differ from those of neurons.
Eur J Neurosci. 1994;6(11):1702-9.

11.   Endo F, Kasai A, Soto JS, Yu X, Qu Z, Hashimoto H, et al. Molecular basis of astrocyte
diversity and morphology across the CNS in health and disease. Science.
2022;378(6619):eadc9020.

12.   Zhang Y, Chen K, Sloan SA, Bennett ML, Scholze AR, O'Keeffe S, et al. An RNA-
sequencing transcriptome and splicing database of glia, neurons, and vascular cells of
the cerebral cortex. J Neurosci. 2014;34(36):11929-47.

13.   Yao Z, van Velthoven CTJ, Kunst M, Zhang M, McMillen D, Lee C, et al. A high-resolution
transcriptomic and spatial atlas of cell types in the whole mouse brain. Nature.
2023;624(7991):317-32.

14.   Alvarez-Castelao B, Schanzenbacher CT, Hanus C, Glock C, Tom Dieck S, Dorrbaum AR,
et al. Cell-type-specific metabolic labeling of nascent proteomes in vivo. Nat
Biotechnol. 2017;35(12):1196-201.

15.   Sjostedt E, Zhong W, Fagerberg L, Karlsson M, Mitsios N, Adori C, et al. An atlas of the
protein-coding genes in the human, pig, and mouse brain. Science. 2020;367(6482).

16.   Buckley NJ, Bonner TI, Brann MR. Localization of a family of muscarinic receptor
mRNAs in rat brain. J Neurosci. 1988;8(12):4646-52.

17.   Heintz N. Gene expression nervous system atlas (GENSAT). Nat Neurosci.
2004;7(5):483.

18.   Vilaro MT, Wiederhold KH, Palacios JM, Mengod G. Muscarinic cholinergic receptors in
the rat caudate-putamen and olfactory tubercle belong predominantly to the m4 class:
in situ hybridization and receptor autoradiography evidence. Neuroscience.
1991;40(1):159-67.

19. Arenander AT, de Vellis J, Herschman HR. Induction of c-fos and TIS genes in cultured
rat astrocytes by neurotransmitters. J Neurosci Res. 1989;24(1):107-14.
20. Bubser M, Bridges TM, Dencker D, Gould RW, Grannan M, Noetzel MJ, et al. Selective
activation of M4 muscarinic acetylcholine receptors reverses MK-801-induced
behavioral impairments and enhances associative learning in rodents. ACS Chem
Neurosci. 2014;5(10):920-42.
21. Spock M, Carter TR, Bollinger KA, Han C, Baker LA, Rodriguez AL, et al. Discovery of
VU6028418: A Highly Selective and Orally Bioavailable M(4) Muscarinic Acetylcholine
Receptor Antagonist. ACS Med Chem Lett. 2021;12(8):1342-9.
22. Tsai HH, Li H, Fuentealba LC, Molofsky AV, Taveira-Marques R, Zhuang H, et al. Regional
astrocyte allocation regulates CNS synaptogenesis and repair. Science.
2012;337(6092):358-62.
23. Cerrato V. Cerebellar Astrocytes: Much More Than Passive Bystanders In Ataxia
Pathophysiology. J Clin Med. 2020;9(3).
24. Gould RW, Grannan MD, Gunter BW, Ball J, Bubser M, Bridges TM, et al. Cognitive
enhancement and antipsychotic-like activity following repeated dosing with the
selective M(4) PAM VU0467154. Neuropharmacology. 2018;128:492-502.
25. Foster DJ, Wilson JM, Remke DH, Mahmood MS, Uddin MJ, Wess J, et al. Antipsychotic-
like Effects of M4 Positive Allosteric Modulators Are Mediated by CB2 Receptor-
Dependent Inhibition of Dopamine Release. Neuron. 2016;91(6):1244-52.
26. Pancani T, Bolarinwa C, Smith Y, Lindsley CW, Conn PJ, Xiang Z. M4 mAChR-mediated
modulation of glutamatergic transmission at corticostriatal synapses. ACS Chem
Neurosci. 2014;5(4):318-24.
27. Moehle MS, Pancani T, Byun N, Yohn SE, Wilson GH, 3rd, Dickerson JW, et al.
Cholinergic Projections to the Substantia Nigra Pars Reticulata Inhibit Dopamine
Modulation of Basal Ganglia through the M(4) Muscarinic Receptor. Neuron.
2017;96(6):1358-72 e4.
28. Jeon J, Dencker D, Wortwein G, Woldbye DP, Cui Y, Davis AA, et al. A subpopulation of
neuronal M4 muscarinic acetylcholine receptors plays a critical role in modulating
dopamine-dependent behaviors. J Neurosci. 2010;30(6):2396-405.
29. Zimmer TS, Orr AL, Orr AG. Astrocytes in selective vulnerability to neurodegenerative
disease. Trends Neurosci. 2024;47(4):289-302.
30. Endo F. Deciphering the spectrum of astrocyte diversity: Insights into molecular,
morphological, and functional dimensions in health and neurodegenerative diseases.
Neurosci Res. 2024.
31. Moulson AJ, Squair JW, Franklin RJM, Tetzlaff W, Assinck P. Diversity of Reactive
Astrogliosis in CNS Pathology: Heterogeneity or Plasticity? Front Cell Neurosci.
2021;15:703810.
32. Tahir W, Thapa S, Schatzl HM. Astrocyte in prion disease: a double-edged sword. Neural
Regen Res. 2022;17(8):1659-65.
33. Hartmann K, Sepulveda-Falla D, Rose IVL, Madore C, Muth C, Matschke J, et al.
Complement 3(+)-astrocytes are highly abundant in prion diseases, but their
abolishment led to an accelerated disease course and early dysregulation of microglia.
Acta Neuropathol Commun. 2019;7(1):83.
34. Akhtar S, Wenborn A, Brandner S, Collinge J, Lloyd SE. Sex effects in mouse prion
disease incubation time. PLoS One. 2011;6(12):e28741.
35. Liu Y, Guo J, Matoga M, Korotkova M, Jakobsson PJ, Aguzzi A. NG2 glia protect against
prion neurotoxicity by inhibiting microglia-to-neuron prostaglandin E2 signaling. Nat
Neurosci. 2024;27(8):1534-44.
36. Denouel A, Brandel JP, Seilhean D, Laplanche JL, Elbaz A, Haik S. The role of
environmental factors on sporadic Creutzfeldt-Jakob disease mortality: evidence from
an age-period-cohort analysis. Eur J Epidemiol. 2023;38(7):757-64.

37. Myslivecek J, Farar V, Valuskova P. M(4) muscarinic receptors and locomotor activity
regulation. Physiol Res. 2017;66(Suppl 4):S443-S55.

38. Chambers NE, Millett M, Moehle MS. The muscarinic M4 acetylcholine receptor
exacerbates symptoms of movement disorders. Biochem Soc Trans. 2023;51(2):691-
702.

39. Nunes EJ, Addy NA, Conn PJ, Foster DJ. Targeting the Actions of Muscarinic Receptors
on Dopamine Systems: New Strategies for Treating Neuropsychiatric Disorders. Annu
Rev Pharmacol Toxicol. 2024;64:277-89.

40. Camandola S. Astrocytes, emerging stars of energy homeostasis. Cell Stress.
2018;2(10):246-52.

41. Gonzalez-Garcia I, Garcia-Caceres C. Hypothalamic Astrocytes as a Specialized and
Responsive Cell Population in Obesity. Int J Mol Sci. 2021;22(12).

42. Yang L, Qi Y, Yang Y. Astrocytes control food intake by inhibiting AGRP neuron activity
via adenosine A1 receptors. Cell Rep. 2015;11(5):798-807.

43. Chen N, Sugihara H, Kim J, Fu Z, Barak B, Sur M, et al. Direct modulation of GFAP-
expressing glia in the arcuate nucleus bi-directionally regulates feeding. Elife. 2016;5.

44. Herrera Moro Chao D, Kirchner MK, Pham C, Foppen E, Denis RGP, Castel J, et al.
Hypothalamic astrocytes control systemic glucose metabolism and energy balance. Cell
Metab. 2022;34(10):1532-47 e6.

45. MacDonald AJ, Holmes FE, Beall C, Pickering AE, Ellacott KLJ. Regulation of food intake
by astrocytes in the brainstem dorsal vagal complex. Glia. 2020;68(6):1241-54.

46. Clyburn C, Carson KE, Smith CR, Travagli RA, Browning KN. Brainstem astrocytes control
homeostatic regulation of caloric intake. J Physiol. 2023;601(4):801-29.

47. Cai P, Huang SN, Lin ZH, Wang Z, Liu RF, Xiao WH, et al. Regulation of wakefulness by
astrocytes in the lateral hypothalamus. Neuropharmacology. 2022;221:109275.

48. Brancaccio M, Edwards MD, Patton AP, Smyllie NJ, Chesham JE, Maywood ES, et al. Cell-
autonomous clock of astrocytes drives circadian behavior in mammals. Science.
2019;363(6423):187-92.

49. Nimmerjahn A, Mukamel EA, Schnitzer MJ. Motor behavior activates Bergmann glial
networks. Neuron. 2009;62(3):400-12.

50. Paukert M, Agarwal A, Cha J, Doze VA, Kang JU, Bergles DE. Norepinephrine controls
astroglial responsiveness to local circuit activity. Neuron. 2014;82(6):1263-70.

51. Corkrum M, Covelo A, Lines J, Bellocchio L, Pisansky M, Loke K, et al. Dopamine-Evoked
Synaptic Regulation in the Nucleus Accumbens Requires Astrocyte Activity. Neuron.
2020;105(6):1036-47 e5.

52. Schweinhuber SK, Messerschmidt T, Hansch R, Korte M, Rothkegel M. Profilin isoforms
modulate astrocytic morphology and the motility of astrocytic processes. PLoS One.
2015;10(1):e0117244.

53. Lee SJ, Seo BR, Koh JY. Metallothionein-3 modulates the amyloid beta endocytosis of
astrocytes through its effects on actin polymerization. Mol Brain. 2015;8(1):84.

54. Zhou J, Zhang L, Peng J, Zhang X, Zhang F, Wu Y, et al. Astrocytic LRP1 enables
mitochondria transfer to neurons and mitigates brain ischemic stroke by suppressing
ARF1 lactylation. Cell Metab. 2024;36(9):2054-68 e14.

55. Romeo R, Glotzbach K, Scheller A, Faissner A. Deletion of LRP1 From Astrocytes
Modifies Neuronal Network Activity in an in vitro Model of the Tripartite Synapse.
Front Cell Neurosci. 2020;14:567253.

56. Gogliotti RG, Fisher NM, Stansley BJ, Jones CK, Lindsley CW, Conn PJ, et al. Total RNA
Sequencing of Rett Syndrome Autopsy Samples Identifies the M(4) Muscarinic
Receptor as a Novel Therapeutic Target. J Pharmacol Exp Ther. 2018;365(2):291-300.

57. Achilly NP, He LJ, Kim OA, Ohmae S, Wojaczynski GJ, Lin T, et al. Deleting Mecp2 from
the cerebellum rather than its neuronal subtypes causes a delay in motor learning in
mice. Elife. 2021;10.

58. Patani R, Hardingham GE, Liddelow SA. Functional roles of reactive astrocytes in
neuroinflammation and neurodegeneration. Nat Rev Neurol. 2023;19(7):395-409.

59. Makarava N, Chang JC, Kushwaha R, Baskakov IV. Region-Specific Response of
Astrocytes to Prion Infection. Front Neurosci. 2019;13:1048.

60. Makarava N, Chang JC, Molesworth K, Baskakov IV. Region-specific glial homeostatic
signature in prion diseases is replaced by a uniform neuroinflammation signature,
common for brain regions and prion strains with different cell tropism. Neurobiol Dis.
2020;137:104783.

61. Carroll JA, Striebel JF, Rangel A, Woods T, Phillips K, Peterson KE, et al. Prion Strain
Differences in Accumulation of PrPSc on Neurons and Glia Are Associated with Similar
Expression Profiles of Neuroinflammatory Genes: Comparison of Three Prion Strains.
PLoS Pathog. 2016;12(4):e1005551.

62. Munoz-Castro C, Serrano-Pozo A. Astrocyte-Neuron Interactions in Alzheimer's Disease.
Adv Neurobiol. 2024;39:345-82.

63. Carter SF, Herholz K, Rosa-Neto P, Pellerin L, Nordberg A, Zimmer ER. Astrocyte
Biomarkers in Alzheimer's Disease. Trends Mol Med. 2019;25(2):77-95.

64. Makarava N, Mychko O, Chang JC, Molesworth K, Baskakov IV. The degree of astrocyte
activation is predictive of the incubation time to prion disease. Acta Neuropathol
Commun. 2021;9(1):87.

65. Gomeza J, Zhang L, Kostenis E, Felder CC, Bymaster FP, Brodkin J, et al. Generation and
pharmacological analysis of M2 and M4 muscarinic receptor knockout mice. Life Sci.
2001;68(22-23):2457-66.

66. Guyenet SJ, Furrer SA, Damian VM, Baughan TD, La Spada AR, Garden GA. A simple
composite phenotype scoring system for evaluating mouse models of cerebellar ataxia.
J Vis Exp. 2010(39).

67. Mallucci G, Dickinson A, Linehan J, Klohn PC, Brandner S, Collinge J. Depleting neuronal
PrP in prion infection prevents disease and reverses spongiosis. Science.
2003;302(5646):871-4.

68. Scarpa M, Molloy C, Jenkins L, Strellis B, Budgett RF, Hesse S, et al. Biased M1
muscarinic receptor mutant mice show accelerated progression of prion
neurodegenerative disease. Proc Natl Acad Sci U S A. 2021;118(50).

69. Mirabile I, Jat PS, Brandner S, Collinge J. Identification of clinical target areas in the
brainstem of prion-infected mice. Neuropathol Appl Neurobiol. 2015;41(5):613-30.

70. Wolfes AC, Dean C. Culturing In Vivo-like Murine Astrocytes Using the Fast, Simple, and
Inexpensive AWESAM Protocol. J Vis Exp. 2018(131).

71. Llorens-Martin M, Torres-Aleman I, Trejo JL. Pronounced individual variation in the
response to the stimulatory action of exercise on immature hippocampal neurons.
Hippocampus. 2006;16(5):480-90.

72. Thomas PD, Ebert D, Muruganujan A, Mushayahama T, Albou LP, Mi H. PANTHER:
Making genome-scale phylogenetics accessible to all. Protein Sci. 2022;31(1):8-22.

Figure legends

Fig. 1: M$_4$ mAChR is expressed in neuronal and disperse astrocytic populations

throughout the brain. a, Schematic overview of the breeding between mice driving

CRE recombinase from the *Chrm4* promoter (Chrm4[CRE]) with mice carrying the

inducible transgene for expression of GFP and MetRS* L274G (R26-MetRS*). The

distribution of M_4 mAChR expressing cells can be detected in the resulting

Chrm4^{CRE}::MetRS* mice by GFP immunostaining (green) in a brain sagittal section. **b**,

Representative confocal images showing co-immunostaining of GFP (green) with the

neuronal marker MAP2 (magenta) in areas of the cortex, striatum and hippocampus.

The white square insets indicate the enlarged areas showed in the panels below by

each individual channel. **c**, Representative confocal images showing co-localisation of

GFP expressing cells with the astrocyte marker GFAP (magenta) in the hypothalamus

and the brainstem. Scale bar for panel **a** is 250 μm, and for **b** and **c**, 25 μm.

Fig. 2: The cerebellum contains a diverse array of cells expressing M_4 mAChR. a,

Distribution of M_4 mAChR expressing cells labelled with GFP staining in a sagittal

section of the cerebellum of Chrm4^{CRE}::MetRS* mice. **b**, Co-immunostaining of GFP-

labelled M_4^+ cells with Calbindin in the Purkinje layer of the cerebellum of

Chrm4^{CRE}::MetRS* mice. **c**, Immunohistochemistry analysis of different populations of

GFAP positive astrocytes (magenta) co-expressing M_4 mAChR (green) in the cerebellar

cortex (left panel), cerebellar nuclei (CN, middle panel) and white matter region (right

panel). Arrowheads indicate M_4^+ astrocytes. Scale bar for panels **a** and **b** is 250 μm, and

for **c**, 25 μm.

Fig. 3: Role of M_4 mAChR in cerebellar astrocytic cultures of Chrm4^{CRE}::MetRS*. a,

Immunostaining of cerebral or cerebellar astrocyte cultures from Chrm4^{CRE}::MetRS*

animals detecting M_4 expressing cells with GFP (green) and the astrocyte marker GFAP

(magenta). **b**, Quantification of the proportion of M_4^+ astrocytes in cerebral and

cerebellar cultures relative to the total astrocytes present in the preparation (n = 5 - 9).

c, The percentage of the population of M_4^+ astrocytes co-expressing c-fos was

estimated in astrocyte cerebellar cultures after stimulation with Oxotremorine (Oxo, 10

μM), a combination of Oxo and the M_4 mAChR PAM VU0467154 (VU154, 10 μM) or

Oxo plus the M_4 mAChR antagonist VU6028418 (VU418, 5 μM) for 60 min (n= 5). **d**,

Representative confocal images showing M_4 expressing astrocytes (green) co-stained

with c-fos (magenta) in cerebellar cultures after treatment with Vehicle (Control), Oxo

or a combination of Oxo and VU154 or Oxo with VU418, as indicated in **c**. **e**, Venn

diagram showing the newly synthesised proteins (NSPs) exclusively identified from M_4^+

cerebellar astrocytes after treatment with ANL (4 mM) in combination with Vehicle,

Oxo (10 μM) or Oxo and VU418 (5 μM). **f**, Analysis of enriched GO annotations (Fisher's

exact test performed with Panther v. 17.0) in M_4^+ cerebellar astrocytes after treatment

with ANL in basal conditions (n = 2). The size of each circle represents the number of

genes, and the colour intensity indicates the $-\log_{10} p$ value adjusted by Benjamini-

Hochberg method assigned to the corresponding GO term. **g**, Gene ontology biological

processes which are significantly enriched from the nascent proteins synthesized after

activation of M_4 mAChR in M_4^+ cerebellar astrocytes (NSPs exclusively found after Oxo

treatment and not identified when the M_4 antagonist is present). Graphs depict means

± SEM and statistical significance was calculated by two-tailed unpaired t test (**b**), or

one-way ANOVA followed by Tukey's pot hoc test (**c**) (*$p < 0.05$, **$p < 0.01$, ***$p <$

0.01,). Scale bars, 25 μm.

Fig. 4: Selective ablation of M_4^+ astrocytes leads to deficits in locomotor activity. a,

Depiction showing the breeding overview between the Chrm4[CRE] animals with the

hemizygous Aldh1l1-eGFP,-DTA mice, that express eGFP under the astrocyte promoter

Aldh1l1 and an inducible diphteria toxin A when Cre recombinase is present. This

mating results in Chrm4[CRE]:: Aldh1l1[DTA] mice (DTA[M4]*) with ablated astrocytes that

express the M_4 mAChR, and the control littermates Chrm4^{CRE}:: Aldh1l1^{WT} (WT^{M4}*), wild

type for the floxed gene. **b**, Average animal weight per group over time (n = 6-10). **c**,

Comparison of the performance on the accelerated rotarod test between DTA^{M4}* and

WT^{M4}* mice during 4 consecutive weeks (n = 10). **d**, Activity of mice in and open field

test measuring the time spent in the central area of the apparatus (n = 15-18). **e**, Total

distance travelled over the whole area of the open field arena. **f**, Analysis of the

anxiety-like behaviour of DTA^{M4}* and WT^{M4}* mice measuring the time spent in the

open arms of an elevated plus maze (n = 15-18). **g**, Locomotor activity in DTA^{M4}*

animals compared to WT^{M4}* littermates measured by telemetry. The graph depicts the

profile of activity counts averaged over 3 days of recording and plotted every five min.

Mice experience a 12/12 h light/dark cycle, with the dark cycle shown in the shaded

area (n = 8). **h**, Comparison of the average telemetry activity counts during the 12-h

light and dark phases between DTA^{M4}* and WT^{M4}* mice. **i**, Effect of VU0467154

(VU154, 10 mg/kg) or vehicle (Veh) to counteract the amphetamine-induced

hyperlocomotion (2 mg/kg) in DTA^{M4}* and WT^{M4}* mice. **j**, Total distance travelled

during the 60 min period following D-amphetamine administration. (n = 6-7). Data are

means ± SEM and statistical significance was calculated by two-tailed unpaired *t* test or

one-way ANOVA followed by Bonferroni's pot hoc test when analysis involved more

than two groups (*p < 0.05, **p < 0.01).

Fig 5. Regional and temporal expansion in the number of the M_4^+ astrocytes during

mouse prion disease. a, Representative confocal images of GFP-labelled M_4^+ cells in

coronal brain sections from Chrm4^{CRE}::MetRS* animals after 20 w.p.i. of RML prion

inoculation (right) or control animals treated with NBH (left). Panels below show a

detail of M_4^+ astrocytes (arrowheads) co-stained with GFP (green) and GFAP (magenta)

from the thalamic areas indicated with white square insets. **b**, Confocal images

showing M_4^+ astrocytes co-stained with GFP (green) and GFAP (magenta) in the

cerebellar nuclei of Chrm4^{CRE}::MetRS* mice after 20 w.p.i. of RML prion disease (right)

or NBH controls (left). **c**, Regional and temporal spreading pattern of the density of M_4^+

astrocytes at different stages of RML prion disease in Chrm4^{CRE}::MetRS* mice plotted

on brain anatomical maps. Darker red sections represent regions with higher densities

in M_4^+ astrocytes, while not quantified areas are depicted in grey. **d**, Quantification of

M_4^+ astrocytes density across brain regions in Chrm4^{CRE}::MetRS* mice treated with

RML prion after 15, 17 and 23 w.p.i. and compared to control animals inoculated with

NBH (n = 2-3). **e**, Representative confocal images showing co-localisation of GFP

(green), GFAP (magenta) and GBP2 in the hippocampus and thalamus of 20 w.p.i. RML-

inoculated Chrm4^{CRE}::MetRS* mice. Scale bar in **a** is 250 μm, and 25 μm in **b** and **e**.

Data shown are represented as mean ± SEM. Statistical significance was calculated by

two-way ANOVA followed by Bonferroni's post hoc test (***$p < 0.001$, **$p < 0.01$).

Fig. 6. Ablation of M_4^+ astrocytes prolongs survival and reduces Prp^{Sc} deposition in

the cerebellum of prion-infected mice. a, Evaluation of progressive motor deficits by

rotarod in prion-diseased DTA^{M4}* and WT^{M4}* mice from 12 w.p.i (n = 8-12). **b**,

Burrowing behaviour (percentage of food pellets displaced from the tube) of control or

prion-infected DTA^{M4}* and WT^{M4}* mice assessed from 12 w.p.i. (n = 5 for NBH, n = 10-

12 for RML). **c**, Kaplan-Meier plot showing a significant increase in survival probability

for RML-prion inoculated mice after ablation of M_4^+ astrocytes (DTA^{M4}*) compared to

wild-type littermates (WT^{M4}*) (n = 22 - 25). Sex-differences are observed when survival

curves are compared between females (**d**, n = 10-17) and males (**e**, n = 8-12). Curves

were analysed with a Gehan-Breslow-Wilcoxon test (**$p < 0.01$, ***$p < 0.001$). **f**,

Immunoblot analysis of PK-resistant Prp^{Sc} from the striatum, cortex, hippocampus and

cerebellum of DTA^{M4}* and WT^{M4}* mice inoculated with NBH or RML-prion. Evaluated

samples were collected at terminal stages of prion diseased mice with equivalent

survival times. **g**, Quantification of Prp^{Sc} shows accumulation of the PK-resistant protein

in all evaluated brain regions from diseased animals. Ablation of M$_4^+$ astrocytes

significantly reduces the deposition of Prp^{Sc} in the cerebellum. Graph represents means

± SEM of Prp^{Sc} levels relative to α-tubulin expression (n = 4-6) expressed as a

percentage of the values obtained for RML-prion diseased wild-type mice (WT^{M4}*).

Statistical significance was calculated by two-way ANOVA followed by Tukey's post-hoc

test (*$p < 0.05$, ***$p < 0.001$).

Fig. 1

a

Chrm4^CRE

R26-MetRS*

Chrm4^CRE::MetRS*

b

| Cortex | Striatum | Hippocampus |

GFP / MAP2

c

| Hypothalamus | Brainstem |

3V

GFP / GFAP

Fig. 2

a Cerebellar cortex / White matter / Cerebellar nuclei

DAPI / GFP

b DAPI / GFP / Calbindin

c Bergmann glia | CN astrocytes | Fibrous astrocytes

DAPI / GFP / GFAP

Fig. 3

Fig. 4

Fig. 5

a

NBH RML

GFP / GFAP

b

NBH

RML

DAPI / GFP / GFAP

c

M4+ astrocytes in prion disease

Control (NBH)   15 w.p.i.   17 w.p.i.   23 w.p.i.

Density (cells/mm³)

0   4,000

d

Density M4+ astrocytes (cells/mm³)

□ Control
□ RML 15 wpi
▨ RML 17 wpi
□ RML 23 wpi

Cortex Striatum Hippocampus Thalamus Hypothalamus Midbrain

e

RML (Thalamus)

25 um

DAPI / GFP / GBP2 / GFAP

Fig. 6