Identification of M$_4$ muscarinic acetylcholine receptor expressing astrocytes that regulate locomotion and survival in murine prion disease

Gonzalo S. Tejeda[*], Collin Molloy, Carrie K. Jones, Craig W. Lindsley, Graeme Milligan and Andrew B. Tobin[*]

Abstract

Whilst there has been much focus on the function of neuronally expressed members of the muscarinic acetylcholine receptor family (mAChR) less attention has been paid to the expression profile and role of the five members of this family (M$_1$-M$_5$ mAChRs) in non-neuronal cells of the brain. Using genetically engineered mice we identify a previously unappreciated sub-population of astrocytes expressing the M$_4$ mAChR subtype. These are located in various brain regions that include the brainstem, hypothalamus and, most abundantly, the cerebellum. Signalling and proteomic analysis of M$_4$ mAChR positive astrocytes from the cerebellum established the functional expression of this receptor subtype and its role in the regulation of protein expression. Genetic ablation of M$_4$ mAChR positive astrocytes in mice revealed a specific role in locomotion behaviour. Importantly, in the context of murine prion disease, a model of terminal neurodegeneration associated with profound neuroinflammation, we observed a significant expansion of M$_4$ mAChR positive astrocytes and report ablation experiments that established that this astrocyte sub-population has a detrimental effect on late-stage disease. Together we provide evidence of a sub-population of M$_4$ mAChR expressing astrocytes that play a specific role in normal neurophysiology and the progression of inflammatory neurodegenerative disease.

Keywords: Astrocyte, muscarinic acetylcholine receptor, M$_4$ mAChR, G protein-coupled receptor, neurodegeneration, neuroinflammation, locomotion.

Abbreviations: CNS: central nervous system, ERK1/2: extracellular signal-regulated kinases 1/2, GFAP: glial fibrillary acidic protein, GFP: green fluorescent protein, eGFP: enhanced green fluorescent protein, GPCR: G protein-coupled receptor, mAChR: muscarinic acetylcholine receptor, NBH: normal brain homogenate, PAM: positive allosteric modulator, pEC$_{50}$: half maximal response, Prp^{Sc}: scrapie prion protein, RML: rocky mountain laboratory, w.p.i: weeks post-inoculation.

Main

The muscarinic acetylcholine receptor (mAChR) family of G protein-coupled receptors (GPCRs), that consists of five subtypes (M_1-M_5 mAChRs), have been under intense investigation due to their role in the regulation of a range of neuronal processes from learning and memory to reward and seek behaviour. This is particularly true for the M_1 and M_4 mAChR subtypes that have emerged as important targets for the treatment of brain disorders, as exemplified by *Cobenfy*, otherwise known as KarXT, a M_1/M_4 mAChR agonist recently approved as a medicine for the treatment of schizophrenia [1, 2]. Studies on the mAChR effects on the central nervous system (CNS) have focused on the neurons on which the receptors are expressed, whilst much less attention has been given to the potential role of this receptor family on non-neuronal cells of the CNS.

Some of the best evidence that mAChRs play a role on glial cells has emerged from the study of the M_1 mAChR in oligodendrocyte precursor cells (OPCs), where specific ablation of this receptor gene has been shown to enhance remyelination in mouse disease models that include autoimmune encephalomyelitis [3], hypoxia [4], aging [5] and Alzheimer's disease [6]. Clues that the mAChRs might also regulate neuroinflammation have emerged from our studies in a prion model of neurodegeneration [7], and others in mouse models of Alzheimer's disease [8], where activation of the M_1 mAChR by selective positive allosteric modulators (PAMs) were shown to reduce markers of astrocyte and microglial activation. However, whilst transcripts for the mAChR family have been reported in astrocytes and microglia [9, 10], including the M_4 mAChR [11-13], little is known about the contribution that these receptors play in the regulation of glial cells.

Here we used a novel transgenic mouse model in which CRE recombinase is expressed under the control of the M_4 mAChR gene promoter (*Chrm4*) to discover the existence of a discrete population of M_4 mAChR-expressing astrocytes located in the brainstem, hypothalamus, and cerebellum. Signalling and proteomic studies established the functional expression of this receptor subtype in astrocytes, particularly of the cerebellum. CRE-mediated ablation of M_4 mAChR expressing astrocytes exposed the role of this sub-population of astrocytes in locomotion. Importantly, our study unveiled the expansion of the M_4 mAChR expressing astrocytes in parallel with the progression of prion neurodegenerative disease. The prospect that this expansion played a detrimental role was indicated by a significant extension of the lifespan of prion-diseased mice following genetic ablation of M_4 mAChR expressing astrocytes. Together our data point to a previously unappreciated sub-population of M_4 mAChR expressing astrocytes that play a specific role in normal neurophysiology and inflammatory neurodegenerative disease.

Results

Generation of a genetic mouse model to identify M_4 mAChR expressing cells

To facilitate the study of the diversity and distribution of M_4 mAChR expressing cells (M_4^+ cells) in the brain, we generated a mouse line that expresses a CRE recombinase under

the control of the *Chrm4*-promoter (Chrm4-CRE-IRES-mM4, hereinafter referred to as Chrm4^{CRE}) **(Fig. 1A)**. In these mice, the *Chrm4* locus was modified to contain an internal ribosomal entry site (IRES) cassette followed by the CRE recombinase coding sequence down-stream of the M$_4$ mAChR receptor exon 3 **(Fig. 1A)**. By crossing these mice with a CRE-dependent green fluorescent protein (GFP) reporter mouse strain (R26-CAG-GFP,-MetRS*^{L274G}, also called R26-MetRS*), we aimed to generate animals where the GFP acts as a marker of M$_4$ mAChR expressing cells [14] **(Fig. 1A)**. The resulting mouse strain (Chrm4^{CRE}::MetRS*), heterozygous for both alleles, showed similar levels of M$_4$ mAChR expression to that of wild-type mice as determined by radio-ligand binding experiments of striatal membranes **(Supplementary Fig. 1A)**. Examination of brain sections of Chrm4^{CRE}::MetRS* animals revealed GFP-positive neurons (stained with the neuronal marker MAP2) especially abundant in the striatum, cortex, and hippocampus **(Fig. 1B)**, areas where Western blots confirmed M$_4$ mAChR expression **(Supplementary Fig. 1B)** consistent with previous studies [15-18]. These experiments, together with the lack of GFP signal detected in the brain of the parental lines **(Supplementary Fig. 1C)**, give confidence that the reporter-GFP system acts as a surrogate marker for M$_4$ mAChR expressing cells/neurons.

M$_4$ mAChRs expression in a discrete astrocytic sub-population.

In addition to M$_4^+$neurons, we also detected dispersed populations of M$_4^+$ stellate cells in the midbrain, hypothalamus, and the pons **(Fig. 1C)**. These cells did not co-localise with markers for microglia, oligodendrocytes or neurons **(Supplementary Fig. 1D)** but rather co-stained with antibody markers for glial fibrillary acidic protein (GFAP) and S100β, thereby identifying them as M$_4^+$ astrocytes **(Fig.1C, Supplementary Fig. 1E)**.

These experiments also revealed a surprisingly diverse array of M$_4^+$ cells in the cerebellum **(Fig. 2A)**, a region with low but detectable levels of M$_4$ mAChR transcript **(Supplementary Fig. 2A)**. In contrast, no significant GFP staining was observed in the cerebellum of the parental mouse strains **(Supplementary Fig. 2B)**. Among these M$_4^+$ cells, we could identify a sparse population of M$_4^+$ Purkinje neurons, labelled with the calcium-binding protein calbindin D-28k **(Fig. 2B)**. Most strikingly, however, was the abundance of cerebellum M$_4^+$ cells that co-stained with astrocytic markers (GFAP, NDRG2, S100β) **(Fig. 2C, Supplementary Fig. 2C)**. The location and morphology of these cells indicated that they correspond to Bergmann glia in the Purkinje layer, some velate astrocytes in the granule cell layer, dispersed fibrous astrocytes surrounding the white matter and a large population of M$_4^+$ astrocytes within the cerebellar nuclei **(Fig. 2C)**.

Signalling analysis of astrocyte M$_4$ mAChRs

To determine the functional expression of M$_4$ mAChRs in glial cells, primary astrocyte cultures were prepared from the cerebrum and the cerebellum of Chrm4^{CRE}::MetRS* mice. In line with the observations in intact brain, M$_4^+$ astrocytes represented only a small proportion of the total astrocyte population prepared from the cerebrum (8.7 ± 2.4 %), whilst a higher proportion (30.4 ± 3.8 %) of M$_4^+$ astrocytes were evident in cerebellar cultures **(Fig. 3A, B)**.

To establish if these M_4^+ astrocytes expressed functional M_4 mAChRs, we took advantage of previous studies that showed that muscarinic agonists can induce accumulation of the Immediate Early-Gene (IEG) c-fos in cultured astrocytes [19]. Therefore, we treated cerebellar astrocytes prepared from Chrm4[CRE]::MetRS* mice with the muscarinic agonist oxotremorine (10 µM). This resulted in a significant up-regulation of c-fos expression in M_4^+ astrocytes **(Fig. 3C, D)**. Furthermore, preincubation with VU0467154 (VU154, 10 µM), a M_4 mAChR PAM [20], increased the number of oxotremorine-induced c-fos$^+$ cells in the cerebellar astrocyte cultures, whilst pretreatment with the highly selective M_4 mAChR antagonist VU6028418 (VU418, 5 µM) [21] prevented oxotremorine-induced stimulation of c-fos in these cells **(Fig. 3C, D)**. Additionally, we probed the phosphorylation of the extracellular signal-regulated kinases 1/2 (pERK1/2), a signalling pathway downstream of M_4 mAChR-Gi activation. Treatment with oxotremorine increased pERK1/2 levels in a concentration-dependent manner ($pEC_{50} = 5.57 \pm 0.21$, **Supplementary Fig. 3A**) whereas the M_4 mAChR antagonist VU418 (1 µM) shifted the curve to higher agonist concentrations ($pEC_{50} = 4.61 \pm 0.22$, $p = 0.002$) consistent with VU418 selective antagonism of the M_4 mAChR ERK1/2 response.

Proteomic evaluation of astrocyte M_4 mAChRs

To further characterise the functional M_4 mAChR response in cerebellar astrocytes, we analysed changes in *de novo* synthesised proteins in response to receptor activation. In these experiments we took advantage of the fact that, in addition to the expression of GFP, the Chrm4[CRE]::MetRS* mice contains a mutated methionyl-tRNA synthetase (MetRS L274G) expressed in a CRE-dependent manner. The enlarged binding pocket of this mutant synthetase allows for the incorporation of the unnatural amino acid azidonorleucine (ANL) into nascent proteins, which can later be tagged by click chemistry for identification of labelled proteins or for biochemical purification [14]. Since our system results in CRE-expression only in M_4^+ cells (i.e. astrocytes) then the mutated methionyl-tRNA synthetase will only be expressed in M_4^+ astrocytes. In this way, although our cerebellar astrocyte cultures are mixed (~30% of the astrocytes are M_4^+ **(Fig. 3B)**), we could conduct proteomic analysis on proteins synthesised exclusively in M_4^+ astrocytes.

We first tested our system by exposing cerebellar astrocyte cultures, prepared from Chrm4[CRE]::MetRS* mice, to ANL (4 mM) followed by fixation, permeabilization and labelling ANL-incorporated proteins with the fluorescent Cy3-alkyne tag – process referred to as fluorescent non-canonical amino acid tagging (FUNCAT). This established a FUNCAT signal (Cy3) that was restricted to M_4^+ astrocytes (identified by GFP and S100β) **(Supplementary Fig. 3B)**.

Next, we collected lysates from ANL treated cerebellar astrocytes prepared from Chrm4[CRE]::MetRS* mice and using click chemistry incorporated a biotin-tag to ANL incorporated proteins. Western blot analysis revealed an abundance of biotinylated nascent proteins using this approach **(Supplementary Fig. 3C)**. These biotinylated proteins were purified using avidin beads and proteins identified by mass spectrometry **(Supplementary Table 1)**. These experiments detected a total of 260 enriched biotinylated proteins **(Fig. 3E)**. A gene ontology (GO) analysis revealed enrichment of

astrocytic markers (GFAP, vimentin, 3-phosphoglycerate dehydrogenase), confirming the specific labelling. Importantly, terms associated with metabolic enzymes, translation and cytoskeleton components were among the most significantly enriched GO groups **(Fig. 3F)**. Furthermore, oxotremorine (10 μM) treatment induced the synthesis of 67 unique proteins that were dependent on M_4 mAChR activation, as they were not identified in the presence of the specific M_4 mAChR antagonist (VU418, 5 μM). A GO enrichment analysis showed that proteins up-regulated by M_4 mAChR activation are involved in trafficking and protein translation **(Fig. 3G)**, including the actin-binding profilin-2, the metallothionein subtypes II and III, and the MAP kinase ERK2.

Diptheria toxin-mediated ablation of M_4^+ astrocytes effects locomotion

In order to decipher the role of M_4^+ astrocytes, we followed a genetic strategy of targeted ablation. The mouse strain Aldh1l1-eGFP,-DTA [22], expresses eGFP in astrocytes and a CRE-dependent diphteria toxin A (DTA). Crossing these animals with the Chrm4^{CRE} strain results in a mouse (Chrm4^{CRE}:: Aldh1l1^{DTA}) where M_4^+ astrocytes are selectively ablated **(Fig. 4A)**. As M_4 mAChR expression was observed in some hypothalamic astrocytes **(Fig. 1B)**, we first evaluated the body weight of these animals over time and found no differences compared to wild type littermates **(Fig. 4B)**. Also, these animals showed similar food consumption and organ weights **(Supplementary Fig. 4A, B)**. Previous studies have indicated that alterations in cerebellar astrocytes, specifically Bergmann glia, led to cerebellar atrophy and motor impairment [23]. However, we did not observe any abnormalities in the morphology or the overall glial composition of the cerebellum of the Chrm4^{CRE}:: Aldh1l1^{DTA} animals when compared to the cerebellum of the parental line **(Supplementary Fig. 4C)**. Furthermore, the Chrm4^{CRE}:: Aldh1l1^{DTA} mice did not show ataxia or any deficits in motor coordination as determined by the hindlimb clasping, ledge and rotarod tests **(Fig. 4C)**. In contrast, exploratory behaviour evaluated in an open field test showed that Chrm4^{CRE}:: Aldh1l1^{DTA} mice exhibited significantly increased time in the centre of the arena compared to littermate controls (74.1 $\pm$ 6.7 and 51.4 $\pm$ 2.6 s, respectively, $p < 0.01$, **Fig. 4D**), while the overall locomotion was reduced (24.1 $\pm$ 1.5 m travelled in the arena versus 30.6 $\pm$ 1.7 m for control mice) and the time spent immobile (231.0 $\pm$ 13.7 s) was greater than wild type littermates (164.7 $\pm$ 14.4 s) **(Fig. 4E, Supplementary Fig. 4D)**. These results could be interpreted as deficits in anxiety-like behaviour, but a further evaluation in an elevated plus maze test excluded this possibility, as Chrm4^{CRE}:: Aldh1l1^{DTA} mice performed similarly to wild type littermates in this paradigm **(Fig. 4E, Supplementary Fig. 4E)**.

These data suggest a potential role for M_4^+ astrocytes in locomotion which was tested further by implant telemetry. Both Chrm4^{CRE}:: Aldh1l1^{DTA} and wild type littermate mice had peaks of activity at similar times and of similar duration, excluding the possibility of disturbed sleep **(Fig. 4G)**. Moreover, fluctuations in body temperature followed a similar pattern, with no differences observed between the two groups of animals **(Supplementary Fig. 4G)**. Interestingly, when considering the overall activity in each phase, it was evident that Chrm4^{CRE}:: Aldh1l1^{DTA} mice moved less intensely during the dark phase (7.85 $\pm$ 0.61 counts) than their control littermates (9.93 $\pm$ 0.95 counts) **(Fig.**

4H). These findings are consistent with a deficit in the overall locomotor activity when the M$_4$$^+$ astrocytes are ablated.

M$_4$$^+$ astrocytes do not regulate amphetamine hyperlocomotion

M$_4$ mAChRs are known to counteract hyperlocomotion induced by amphetamine or the glutamate receptor agonist MK-801 [20, 24]. This pivotal function in motor behaviours has been attributed to neuronal (presynaptic) M$_4$ mAChR regulation of the dopamine release and signalling in the basal ganglia [25-28]. Here, we examined if M$_4$$^+$ astrocytes might contribute to this by testing the efficacy of a M$_4$ mAChR selective PAM, VU154, to reverse amphetamine-induced hyperlocomotion in Chrm4^{CRE}:: Aldh1l1^{DTA} and littermate control mice. Administration of VU154 (10 mg/kg, i.p.) 20 mins before amphetamine (2 mg/kg, i.p.) significantly decreased amphetamine-induced hyperlocomotion in both groups of mice **(Fig. 4I, J)**. Likewise, both genotypes displayed comparable measurements in mobility time **(Supplementary Fig. 4H)**. Together these data indicate that the role of M$_4$$^+$ astrocytes in locomotion observed earlier is independent from the dopamine-dependent motor function of M$_4$ mAChR in the basal ganglia.

M$_4$$^+$ astrocytes during prion disease

There is currently intense interest in the role of astrocytes in regulating the onset and progression of many neurodegenerative diseases and other brain disorders [29, 30]. It is clear, however, that the responses to pathological stimuli of reactive astrocytes are very diverse, which could reflect the heterogeneous nature of astrocyte sub-populations with established differences in origin and temporal function [31]. Here we probed the potential involvement of M$_4$$^+$ astrocytes in the pathology of murine prion disease, a progressive terminal neurodegenerative disease model associated with pronounced neuroinflammation and astrogliosis [32]. Prion disease was induced in Chrm4^{CRE}::MetRS* mice by inoculation with Rocky Mountain Laboratory (RML) prion-infected brain homogenate. At 20 weeks post-inoculation (w.p.i.), a sharp increase was observed in the number of M$_4$$^+$ astrocytes **(Fig. 5A)**, as determined by co-staining with the astrocytic marker GFAP **(Supplementary Fig. 5A)**. The lack of co-staining of these M$_4$$^+$ cells with microglial marker Iba1 excluded the possibility that these cells were infiltrating macrophages **(Supplementary Fig. 5B)**. In fact, the strong GFAP staining is consistent with these M$_4$$^+$ cells showing characteristics of reactive astrocytes, which was particularly evident in the region of the cerebellar nuclei **(Fig. 5B)**.

A rostro-caudal examination of serial brain sections from RML-inoculated Chrm4^{CRE}::MetRS* mice demonstrated that M$_4$$^+$ astrocytes were profoundly up-regulated in disease, even in regions such as the cortex, the hippocampus and the thalamus where these cells were not evident under normal "non-disease" conditions **(Supplementary Fig. 5A, B, C)**. To interrogate regional differences, we estimated the number of M$_4$$^+$ astrocytes in each area during prion disease and displayed the cellular densities as heat maps overlaid onto the brain anatomy **(Fig. 5C)**. Our results indicated that the hypothalamus hosts the largest number of M$_4$$^+$ astrocytes. As prion disease is rapidly progressive, we further studied the evolution of these cellular sub-populations

by comparing three time points after RML-inoculation (15, 18 and 23 w.p.i., **Fig. 5C, Supplementary Fig. 5D)**. We found that the number of M_4^+ reactive astrocytes gradually increased in almost all examined brain regions over time **(Fig. 5D)**. Interestingly, this prion-induced expansion was not observed in one of the biggest reservoirs of M_4^+ astrocytes, the cerebellar nuclei within the arbor vitae **(Fig. 5E)**, although astrocytes in this region showed a prominent reactivity during disease **(Fig. 5C)**. This spatio-temporal analysis of the M_4^+ astrocytes in prion disease indicated that these cells become activated in a pathological context and the amount of reactive astrocytes that express the M_4 mAChR continuously increases alongside the progression of the disease.

Increased lifespan of prion disease inoculated mice where M_4^+ astrocytes are ablated

To establish the potential role played by M_4^+ astrocytes in prion disease, we evaluated the disease progression in mice where the M_4^+ astrocytes had been ablated. Inoculation of both Chrm4^{CRE}:: Aldh1l1^{DTA} and wild type littermate mice with RML resulted in a similar onset of prion disease and early/mid disease progression, as determined by near equivalent times when at least two early "clinical" signs of prion disease appeared **(Supplementary Fig. 6A)**. Similarly, there was equivalence in progressive weight loss **(Supplementary Fig. 6B)**, decreased motor co-ordination and decline in burrowing behaviour (a sign of hippocampal function) between the genotypes **(Fig. 6A, B)**. Despite a similar progression of early and mid-stages of disease, Chrm4^{CRE}:: Aldh1l1^{DTA} mice showed extended lifespan compared to wild type littermates (170 ± 3 days and 150 ± 2 days, respectively, *******p* < 0.01) **(Fig. 6C).** This survival effect was more prominent in females **(Supplementary Fig. 6C)**, which could be explained by their shorter incubation periods to Prp^{Sc} compared to males when exposed to the same quantity of infectious material [33]

We next assessed the accumulation of misfolded, insoluble scrapie prion protein (Prp^{Sc}), which constitutes the major component of infectious prion disease. At the terminal stage, a dramatic deposition of Prp^{Sc} was detected in the hippocampus, cortex, striatum and cerebellum of prion-inoculated mice when compared to the non-diseased controls **(Fig. 6D, E)**. Importantly, the accumulation of Prp^{Sc} was significantly decreased in prion inoculated Chrm4^{CRE}:: Aldh1l1^{DTA} compared to wild type controls in the cerebellum, but not in the rest of the brain regions **(Fig. 6E)**. However, this reduction was not accompanied by changes in other indicators of neurodegeneration (ApoE and serpinA3N) or markers of neuroinflammation (GFAP, Iba1 and clusterin) **(Supplementary Fig. 6D, E)**. Overall, these data indicate that the ablation of M_4^+ astrocytes does not affect the onset of the clinical symptoms or progression of prion disease but extends the lifespan of mice, indicating a potential detrimental effect of M_4^+ astrocytes in the later stages of disease.

Discussion

Despite the considerable progress in understanding the neuronal populations that display the M_4 mAChR [34-36], the identity, location and function of non-neuronal CNS cells, in particular glial cells, expressing this receptor subtype has remained largely

unexplored. Here, we reveal a discrete sub-population of M_4 mAChR expressing astrocytes dispersed across several areas of the mouse CNS. Genetic ablation of these astrocytes demonstrated a role in neurological responses associated with locomotion, whilst other M_4 mAChR-associated processes such as anxiety and the regulation of responses to amphetamine remained unaffected. Importantly, there was a dramatic expansion of this sub-population of astrocytes during the progression of neurodegeneration in a model of neuroinflammatory disease. Ablation studies indicated a detrimental role of these M_4^+ astrocytes in latter disease stages. In these ways our study contributes to a growing awareness that astrocytes are not uniform but rather a diverse set of cell types with varying cellular and functional characteristics. Appreciating the nature and complexity of the various subtypes of astrocytes will be essential if we are to understand their role in normal neurophysiology and if we are to target these cells in brain diseases.

Under physiological conditions, we found M_4^+ astrocytes in the hypothalamus and dorsal parts of the brainstem. These regions are critical regulators of energy balance and appetite, and astrocytes in these areas have been reported to contribute to these tasks by modulating the activity of adjacent neurons [37]. Thus, manipulation of hypothalamic and brainstem astrocytes has been reported to influence metabolic processes including feeding and glucose metabolism [38-43], as well as circadian rhythm, and sleep [44, 45]. Interestingly, ablation of M_4^+ astrocytes in mice did not result in alterations in the weight, body temperature or daily activity cycle. It was, however, of relevance that M_4^+ astrocytes regulate locomotion, an activity consistent with the expression of the receptor in a diverse array of the cerebellar astrocytes, which are activated during voluntary and enforced locomotion [46, 47]. It is important to note that M_4^+ astrocytes are involved in specific aspects of motor control whilst other behaviours such as balance and coordination, typically altered after a loss of Bergmann glia support [23], remained normal in M_4^+ astrocyte-depleted mice. Furthermore, the ablation of M_4^+ astrocytes had no impact on basal ganglia-mediated hyper-locomotion responses, classically linked with neuronal M_4 mAChRs [25-28], despite the fact that astrocytes in this region can modulate these psychomotor behavioural effects [48]. Hence, our studies have dissected a novel locomotion response that may be under the regulation of M_4 mAChR expressed on a sub-population of astrocytes.

Whilst our study primarily identified the existence of a novel population of astrocytes, the neuro-anatomical distribution of these cells, and the function that these astrocytes might play under normal and neurodegenerative disease scenarios, it is a particular challenge to relate this to the activity of M_4 mAChRs themselves. We started to investigate this in primary cultures where we used a genetic approach that allows for the signalling and proteomic interrogation of the M_4^+ cerebellar astrocytes amongst a mixed population of astrocytes. This has provided an in-depth proteomic profile of M_4^+ astrocyte sub-population and demonstrated that selective M_4 mAChR activation resulted in the regulation of key cytoskeletal, metabolic and translational proteins that can significantly impact astrocyte biology and activation status. For example, profilin-2 can modulate astrocytic morphology [49]; metallothionein III plays a role in the clearance of

the prion-like amyloid β [50], and the low-density lipoprotein receptor-related protein-1 (LRP1) is important for the astrocyte-neuron communication [51, 52]. Interestingly, LRP1 is downregulated together with M$_4$ mAChR in the cerebellum of patients with Rett syndrome [53], and mouse models show motor deficits when the *MeCP2* gene is deleted from the cerebellum but not when is restricted to Purkinje or granular neurons [54], highlighting the contribution of the cerebellar astrocyte pool.

The involvement of astrocytes, and indeed microglia, in protection against neurodegenerative disease is an extremely active area of research [55]. It is therefore of relevance that M$_4^+$ astrocytes expand following RML inoculation and that ablation of M$_4^+$ astrocytes increased the lifespan of prion diseased mice. There are numerous studies suggesting that the neurotrophic and neuroprotective properties of astrocytes as well as their role in clearing misfolded proteins is important in protecting against *"prion-like"* misfolded neurodegenerative diseases such as Alzheimer's disease [56]. Whilst this is true for early stages of disease under conditions of progressive neurodegeneration associated with chronic neuroinflammation, the prolonged activation of astrocytes is widely considered to be detrimental [57, 58]. This certainly seems to be the case for M$_4^+$ astrocytes which do not seem to make a significant impact on early- and mid-stages of disease but rather are involved in acceleration of the latter stages of disease. How the M$_4$ mAChR itself is involved in the regulation of M$_4^+$ astrocytes at these latter disease stages, and whether agonists or antagonists might make an impact is at present uncertain. What is clear, however, is that targeting M$_4^+$ astrocytes might be a selective therapeutic strategy that would directly target astrocytes with detrimental functions leaving intact other astrocyte sub-types that might be beneficial.

In conclusion by identifying a previously unappreciated astrocyte sub-population we not only provide further evidence of the diversity of astrocytes but by ablation of M$_4^+$ astrocytes highlight the role of this astrocyte population in the regulation of locomotion and neurodegenerative disease pathology.

Methods

Animals

All experiments were conducted in accordance with the UK Home Office Animal Procedures Act (1986) and all procedures were approved by the University of Glasgow Animal Welfare and Ethical Review Board. Mice were maintained on a 12-h light–dark cycle with ad libitum standard mouse chow food and water.

Generation of Chrm4^{CRE} mice

Transgenic Chrm4^{CRE} mice were generated by genOway (Lyon, France). Briefly, a CRE-IRES cassette was inserted in frame of the ATG in exon 2 of the endogenous *Chrm4* gene, by homologous recombination in embryonic stem (ES) cells. To this end a targeting vector was created containing long and short homology arms to the *Chrm4* genomic region. The long arm was obtained from a 4,656 bp size fragment containing most of intron 1. This

was generated by PCR from genomic DNA template using primers 5' ATTGAGAAGGCTAGGGGAGGAATAGTCC and 5' ACTGGAGACAGGTGGGCAGTGGC. The short arm was generated from a 3,278 bp fragment containing the 3' UTR part of exon 2 and downstream sequences, using primers 5' TTAGGAGGTGCTGGTGTCAAGAGTGTGTAC and 5' AAGCGCACTGGTAAAACCGAACTCC. The targeted vector was completed with an FRT-flanked neomycin resistance gene upstream the CRE sequence to select ES cells that had undergone homologous recombination, and a diphtheria toxin cassette upstream the long homology arm for negative selection. The linearised targeting construct was transfected into C57BL/6N ES cells by electroporation. Positive clones were assessed by PCR and whole locus sequence analysis to validate the presence of the correct recombination event before being injected into blastocysts derived from C57BL/6J female mice. The resulting highly chimeric males (chimerism rate above 50%) were then crossed to a FLP recombinase-expressing deleter mouse strain to generate heterozygous knock-in animals devoid of neomycin. Genotyping was initially confirmed by PCR on genomic DNA obtained from ear biopsies using the primers 5' CACTGCCCACCTGTCTCCAG and 5' GGTTGTGGGCTGTTGTGACCAG, which flank the insertion site of the CRE-IRES cassette; and further confirmed with the pair of primers 5' GCTAGAGCCTGTTTTGCACG and 5' GCTAGAGCCTGTTTTGCACG, which amplify a region of the CRE recombinase sequence. PCR-positive animals were further validated by whole locus sequencing. The results were later corroborated by fluorescent quantitative PCR (qPCR) using a commercial genotyping service (Transnetyx, Cordova, TN). This service also analysed further samples to verify the genotype of the first litter of every new breeder pair for the maintenance of the Chrm4^{CRE} colony.

Homozygous Chrm4^{CRE} were crossed with homozygous R26-CAG-GFP,-MetRS* (Jackson Laboratory, strain: 028071) to generate Chrm4^{CRE}::MetRS*, and heterozygosity for both alleles was confirmed by Transnetyx genotyping service. Chrm4^{CRE} animals were mated with hemizygous Aldh1l1-eGFP,-DTA mice (Jackson Laboratory, strain: 026033) to produce Chrm4^{CRE}::Aldh1l1^{WT} and Chrm4^{CRE}:: Aldh1l1^{DTA}. All these animals were genotyped with specific probes designed for CRE recombinase, eGFP and Chrm4 sequences at weaning by a commercial vendor (Transnetyx). Chrm4^{CRE}:: Aldh1l1^{DTA} and wild type littermate animals were kept mixed in housing cages, except when measuring the food and water intake and during telemetry monitoring experiments. The generation of M$_4$ KO mice has been described elsewhere [59].

Measurement of organ weight, food and water consumption

Chrm4^{CRE}:: Aldh1l1^{DTA} and wild type littermate mice were housed separated by genotype and food and water intakes were measured every 3 days for 1-2 weeks. Average daily food and water consumption per animal were calculated and no differences were observed between genotypes or sex.

The brain, liver, heart, spleen and kidneys from Chrm4^{CRE}:: Aldh1l1^{DTA} and wild type littermate animals were collected (n=3-5, 4-5 months old) and the weight was recorded and adjusted for body weight. Organ dimensions were also measured, and no differences were observed (data not shown).

Rotarod

Motor performance was tested using the 5-lane rotarod (Cat. #47650, Ugo Basile, Germany). Mice were weighed prior every testing, which happened weekly for both prion infected and non-infected 10-week-old male mice for 15-20 weeks. Each day, mice were placed on the 3 cm diameter rod for 2 habituation phases, consisting of a constant rotation speed at 4 rpm for 90 s. For the testing phase, the animals were placed on the rod that accelerated from 4 rpm to 40 rpm over a period of 5 min. The mice were tested 2 times with a rest period of at least 15 min between each trial. Animals that fell before the rod reached 15 rpm speed were placed back on the beam, but if they fell more than once in that period, they were returned to the home cage. If mice turned on the rod, the test was stopped, and the time recorded. The latency to fall (time that the mouse remains on the rod) was registered automatically by a trip switch under the floor of each rotating drum, and the result was used as an indicator of motor coordination. The activity of Chrm4^{CRE}:: Aldh1l1^{WT} in this test was comparable to wild type animals with a C57BL/6 background (data not shown).

Elevated Plus Maze (EPM)

The EPM test was used to assess anxiety-related behaviour in mice. 9-10 weeks old male and female animals were placed in the centre of a cross shaped maze (arms of 50 x 10 cm), with 2 enclosed arms (black walls of 30 cm height) and 2 dimly illuminated open arms opposite each other. The test was performed before the start of the light cycle and lasted for 5 min. The time spent in the open arms versus the enclosed spaces versus, the immobility time and the number of entries to each arm were recorded using the ANY-maze video tracking software (Stoelting).

Open field

The open field activity of 8-12 weeks old male and female mice was monitored using a clear Perspex square box (50 × 50 cm). Animals were placed in the centre of the illuminated open field arena and allowed to freely explore for 10 minutes. The time spent in the centre of the arena, the distance travelled and speed were assessed using the ANY-maze tracking system (Stoelting).

Hindlimb clasping test

The degree of hindlimb clasping as an indicator of motor dysfunction was evaluated in 8-9 weeks old mice. Animals were suspended by their tails for 10-20 s and rated from 0 to 3 according to the following score system: 0 corresponded to hindlimbs consistently

splayed outward and away from the abdomen; 1 was assigned when one hindlimb was retracted inwards towards the abdomen for more than 50% of the time; a score of 2 meant that both hindlimbs were partially retracted inwards towards the abdomen for at least 50% the time; and 3 was given when both hindlimbs were entirely retracted inwards towards the abdomen for at least 50% of the observation period [60]. Two separate trials were performed per animal. All evaluated Chrm4^{CRE}:: Aldh1l1^{DTA} and Chrm4^{CRE}:: Aldh1l1^{WT} animals showed a consistent score of 0 in this test.

Ledge test

Mice aged 8-9 weeks old were placed on the cage ledge and recorded to measure balance and coordination. Paw placement and forward movement as it walked along the ledge were observed and scored from 0 to 3 as in [60]: 0, for normal walking; 1, if the mouse lost its footing; 2, if the hindlimbs were not used, and 3, if the animal fell off the ledge. Data from three trials were averaged for each mouse and none of the Chrm4^{CRE}:: Aldh1l1^{DTA} and Chrm4^{CRE}:: Aldh1l1^{WT} mice presented any deficits in balance and coordination.

Telemetry

A TA-F10 temperature and activity probe (Data Sciences International) was implanted in the abdominal cavity of 9-13 weeks old male and female Chrm4^{CRE}:: Aldh1l1^{DTA} and wild type littermate mice (n=8, 19-30 g), using aseptic techniques under a surgical level of isoflurane anaesthesia. Carprofen (10 mg/kg) and buprenorphine (0.1 mg/kg) were s.c. administered 30 min before surgery for pain relief. After surgery, mice were housed individually, and post-operative analgesia was given 6 h (buprenorphine) and 24 h later (carprofen and buprenorphine). One week after surgery, when the animals completely recovered, the cages were placed over the receiver panels and the TA-F10 transmitters were activated. Activity and body temperature data were sampled every 1 min for 10 s and continuously acquired during 4 consecutive days using the Ponemah software (Data Sciences International). The measurements from the final 3 days were averaged together for each mouse. No differences were observed between males and females.

Amphetamine-induced hyperlocomotion

Male and female Chrm4^{CRE}:: Aldh1l1^{DTA} and wild type littermate mice (n=6-7, 10-11 weeks old) were placed in an open field arena and the animals were tracked with the ANY-maze software (Stoelting). Habituation lasted for 20 min before being i.p. injected with vehicle (10% Tween 80 in sterile water) or VU0467154 (10 mg/kg). A dose of D-amphetamine (2 mg/kg in saline, i.p.) was administered 20 min later, and locomotor activity was recorded for an additional 90 min. The mobility and distance travelled over time were expressed and averaged per 5 min bins. Total activity data were calculated as the total distance travelled 60 min after the time of D-amphetamine administration.

Burrowing

The voluntary behaviour of removal material from an enclosed tube was assessed in control and prion-infected Chrm4^{CRE}:: Aldh1l1^{DTA} and wild type littermate female mice every week from 9 w.p.i. until survival time. Mice were placed into individual cages (22 × 36 cm) with an acrylic clear tube filled with food pellets for 2 h. The burrowing activity was calculated by subtracting the weight of the remaining pellets at the end of the experiment from the starting weight (140 g). Mice were acclimatised to the burrowing cage the week before the start of the test.

Prion mouse model

Male and female Chrm4^{CRE}::MetRS*, Chrm4^{CRE}:: Aldh1l1^{DTA} and Chrm4^{CRE}:: Aldh1l1^{WT} mice aged 3-4 weeks old were i.c. inoculated, under a surgical level of isoflurane anaesthesia, into the right brain parietal lobe with 30 µl of either 1% NBH (Normal Brain Homogenate) or 1% RML prion (Rocky Mountain Laboratories), as previously described [61, 62].

Evaluation of prion disease progression and survival

Chrm4^{CRE}:: Aldh1l1^{DTA} and Chrm4^{CRE}:: Aldh1l1^{WT} control and prion-infected mice were mixed in housing cages. Progression of the disease was evaluated according to the presence of early indicator and confirmatory signs of prion disease and scored by an experimenter blind to the genotype. Early indicator signs included piloerection, mild loss of coordination, erect penis, clasping of hindlimbs when lifted by tail, sustained erect ears, rigid tail, discontinued hunched posture, or subdued behaviour. Confirmatory signs of prion disease included ataxia, righting reflex impairment, sustained hunched posture, dragging of limbs, and/or significant abnormal breathing. When the animals showed two confirmatory signs or two early indicators plus one confirmatory sign, they were humanely killed and the days after inoculation were considered as the survival time.

Preparation of protein extracts and immunoblot analysis

Mice were culled via cervical dislocation, and different regions of the brain were dissected and snap frozen on dry ice. Samples were lysed in RIPA buffer (10 mM Tris-Cl, 1 mM EDTA, 1% Triton X-100, 0.1% sodium deoxycholate, 0.1% SDS, 140 mM NaCl, pH 7.5) supplemented with protease inhibitor tablets (Cat. #11697498001, Roche) and incubated for 30 min at 4 °C. Lysates were then briefly sonicated (Cat. #12337338, Fisherbrand) with 2 pulses of 15 s at an amplitude of 40% (~50 W) until fully dissolved. Samples were then centrifuged at 10,000 × g for 10 min, supernatants collected, and protein concentration calculated using a BCA assay kit (Cat. #23225, Pierce). Equal concentrations of protein samples were resuspended in Laemmli loading buffer (10% SDS, 300 mM Tris-HCl pH 7.2, 0.05% bromophenol blue, 10% β-mercaptoethanol) and heated at 60°C for 10 min. For the assessment of PrP^{Sc} levels, 60 µg of each lysate were treated with proteinase K (10 µg/ml, Cat. P2308, Sigma-Aldrich) for 10 min at 37°C. The reaction was stopped by the addition of Laemmli sample buffer and boiling at 95°C for 5

min. 20-40 μg of each cell lysate were resolved on NuPAGE precast 4-12% gels (Cat. #NP0321, Invitrogen) followed by transfer onto nitrocellulose membranes (Cat. #88018, Thermo Scientific). Protein transfer was verified with Ponceau-S staining (0.2% Ponceau-S red, 1% acetic acid), before membranes were blocked with 5% non-fat dry milk in TBST (25 mM Tris-HCl; pH 7.6, 100 mM NaCl, 0.5% Tween 20) for 1 h at room temperature. Membranes were incubated with the corresponding primary antibodies in the blocking solution overnight at 4°C. After washing three times for 5 min each with TBST, secondary antibodies were added to the membranes for 1 h at RT. Antibodies used in this study are detailed in Supplementary Tables 2 and 3. Immunoreactive bands were visualised using the Licor Odyssey system and densitometry of fluorescence was measured using Image Studio Lite (Licor Biosciences).

Astrocyte cultures

Murine astrocyte cultures were performed according to the AWESAM protocol [63] with some modifications. Brains from Chrm4[CRE]::MetRS* or WT pups (P0-P5) were collected and immediately placed in ice cold HBSS (ThermoFisher Scientific). After removing of meninges, the cerebellum or the cerebrum were dissected and incubated with 0.05% trypsin-EDTA (Cat. #25300054, ThermoFisher Scientific) at 37°C for 10 min followed by gentle mechanical dissociation. Cells were centrifuged at 1,000 x g for 3 min, resuspended in high glucose, Glutamax-supplemented DMEM (Cat. #61965026, Gibco) with 10% Foetal Bovine Serum (Cat. #10500064, Gibco), 100 units/ml penicillin and 100 μg/ml streptomycin (Cat. # 15070063, Gibco), and plated on flasks precoated with poly-D-lysine (100 μg/ml, Cat. #P0899, Sigma-Aldrich). After 7 days in vitro (DIV), the flasks were shaken at 180 rpm for 30 min at 37°C, media changed and further shaken at 240 rpm for 6 h. Astrocytes were detached and seeded at a density of 0.2×10^6 cells/ml on the corresponding experimental poly-D-lysine plates. After a couple of days, media was replaced with NB+H media, containing Neurobasal (Cat. # 21103049, Gibco) supplemented with B-27 (Cat. #17504044, Gibco), 2 mM Glutamax (Cat. # 35050061, Gibco), 10 ng/mL heparin-binding EGF-like growth factor (Cat. #E4643, Sigma-Aldrich), 100 units/ml penicillin and 100 μg/ml streptomycin for 5 days before the start of the experiments.

ERK1/2 phosphorylation assay

Cultured primary astrocytes from WT mice were seeded onto 96 well plates precoated with poly-D-lysine at a density of 20,000 cells/well in NB+H media. One week later, the media was replaced, and cells were incubated with 80 μl of Neurobasal without additives for 4 h. Cells were treated for 10 min with vehicle or 10 μM of the M_4 mAChR antagonist VU6028418 before stimulation with the indicated concentrations of oxotremorine for 5 min at 37°C in a final volume of 100 μl. Maximum ERK1/2 phosphorylation response was obtained by treating the cells with FBS. ERK phosphorylation was evaluated using a Homogenous Time-Resolved FRET (HTRF)-based phospho-ERK (Thr202/Tyr204) cellular

kit (Cat. #64ERKPEH, Revvity). Briefly, lysis buffer was added immediately after the end of the stimulation and incubated for 30 min. After homogenisation, 16 μl of the lysate were transferred to a 384-well white ProxiPlate (PerkinElmer) and combined with 4 μl HTRF premixed antibodies. for 4 h at room temperature. Fluorescence emission at 665 and 620 nm was measured after 4 h with a PHERAstar plate reader (BMG Labtech) and results were normalised to the response stimulated by FBS.

Immunocytochemistry analysis

Cerebral or cerebellar primary astrocytes from Chrm4[CRE]::MetRS* mice were grown onto 24 well plates with 13 mm glass coverslips previously treated with poly-D-lysine as indicated above. Cells were fixed with 4% formaldehyde for 1 h at room temperature. Coverslips were then washed 3 times with PBS and blocked with PBS supplemented with 1% BSA and 0.2% Triton X-100 for 1 h. Subsequently, astrocytes were incubated with the indicated primary antibodies diluted in blocking solution overnight at 4°C in a humidity chamber. Detection was achieved using appropriate secondary antibodies conjugated to Alexa Fluor 488 or Alexa Fluor 594 (1:500, Life Technologies) and coverslips were mounted on glass slides using Fluoromount-G with DAPI (ThermoFisher Scientific). Confocal images were acquired with a Zeiss Pascal laser-scanning confocal microscope (LSM) 880 using the LSM image acquisition software.

For analysis of c-fos expression, the NB+H media was replaced with non-supplemented Neurobasal, and cells were incubated at 37°C and 5% CO_2 for 4 h, before being stimulated with vehicle or 10 μM oxotremorine for 60 min. To evaluate M_4 mAChR action, cultures were simultaneously treated with vehicle, 10 μM VU0467154 or 5 μM VU6028418. Immunocytochemistry was performed as described above and, for each replicate (n = 3), 8 confocal images (425 μm x 425 μm) per condition were capture, randomly selected using DAPI staining from areas with equivalent number of cells. The images were analysed using ImageJ Fiji (NIH Image). Automatic quantification of c-fos[+] cells was performed using the "Analyze particle" plugin. M_4 mAChR-expressing astrocytes were manually counted and M_4^+/c-fos[+] double stained cells were noted. Total number of cells were also counted automatically with the software tools. The experimental conditions were blinded to the investigator performing the image analysis. This methodology was also used to calculate the proportion of M_4^+ astrocytes present in cerebral and cerebellar cultures.

For FUNCAT analysis, Chrm4[CRE]::MetRS* astrocytes were incubated with 4 mM L-Azidonorleucine hydrochloride (ANL, Tocris) for 24 h. Cells were fixed and blocked as previously described, followed by click chemistry to visualize the ANL-labelled newly synthesized proteins. The Cu(I)-catalysed azide/alkyne cycloaddition (CuAAC) reaction comprised the addition of 0.45 mM $CuSO_4$, 0.9 mM Tris((1-hydroxy-propyl-1H-1,2,3-triazol-4-yl) methyl) amine (THPTA, Cat. #CLK-1010, Jena Bioscience), 0.2 mM Cy3-alkyne (Cat. #777331, Sigma-Aldrich) and 5 mM Sodium ascorbate to the cells for 24 h.

Coverslips were washed with PBS containing 0.5 mM EDTA for 10 min, followed by 3 washes with blocking solution and the remaining steps of the immunocytochemistry proceeded as indicated above.

Immunohistochemistry

Mice were deeply anaesthetised with isofluorane and then they were transcardially perfused with 20 ml saline solution followed by 20 ml 4% formaldehyde in PBS. Brains were removed and incubated overnight in the same fixative. The whole brain was coronally or sagittally sectioned (50 μm) using a VT1200S vibratome (Leica). After blocking the free-floating sections with PBS supplemented with 0.5% Triton X-100 and 1% bovine serum albumin for 1 h at room temperature, they were then incubated with the indicated primary antibodies diluted in blocking solution 24 to 48 h at 4°C. Antibodies used in this study are listed in Supplementary Tables 2 and 3. Brain sections were washed 3 times with PBS, followed by incubation with the corresponding secondary antibodies diluted in blocking solution for 4 h at room temperature. Sections were placed onto 2% gelatine-coated slides and mounted with Fluoromount-G with DAPI. Images were taken with a Zeiss LSM880 confocal microscope using the LSM image acquisition software Zen lite.

Cell counting

The right brain hemisphere of each mouse was sliced in 50 μm coronal sections that were sequentially collected in series containing one every 8th slice. Randomly chosen series of sections were used for immunohistochemistry against GFP and GFAP as described above. Due to the low numbers of M_4^+ astrocytes in the brain, we counted all visible cells positive for both GFP and GFAP within the indicated brain regions in every section of the series under a Zeiss microscope. DAPI images of each section were taken and the area of each region of interest was traced using the freehand drawing tool of ImageJ Fiji. The determined area was then multiplied by the thickness of the section to calculate the reference volume. The density of M_4^+ astrocytes (number of cells/mm^3) was estimated by dividing the total number of positive cells by the reference volume of the corresponding region.

The quantification of M_4^+ astrocytes in the arbor vitae of the cerebellum was estimated by unbiased stereology using the physical dissector method adapted to confocal microscopy, as previously described [64]. One in eight coronal sections of the cerebellum (50 μm) were immunostained for GFP and GFAP as before. 9-12 confocal counting stacks, acquired with a LSM880 Zeiss microscope, were randomly selected from the whole arbor vitae of the cerebellum. The counting frame size was 150 × 150 μm with a thickness of 10.5 μm and a Z-interval of 1.5 μm. The mean cell density of M_4^+ astrocytes was estimated with the counts of the double-labelled cells and the size of the dissector.

Saturation [^{3}H]-NMS binding assay

Striata from homozygous M_4 KO, R26-MetRS* (WT) or heterozygous Chrm4[CRE]::MetRS* were collected and homogenised in ice-cold HEPES buffer (10 mM HEPES, 1 mM EGTA, 1 mM dithiothreitol) containing 10% sucrose and protease inhibitors. The lysate was first centrifuged at 1,000 x g for 10 min at 4°C, the pellet resuspended, re-homogenised and further centrifuged at 20,000 x g for 1 h at 4°C. The resulting pellet containing striatal membranes, was diluted and protein concentration calculated by the Bradford method. 22.5 µg of the striatal membrane preparation were incubated in binding buffer (50 mM HEPEs, 110 mM NaCl, 5.4 mM KCl, 1.8 mM $CaCl_2$, 1 mM $MgSO_4$, 25 mM glucose, 58 mM sucrose, pH 7.4) with increasing concentrations (0.3-5 nM) of the muscarinic receptor antagonist [3H]-N-methyl scopolamine ([3H]-NMS, Cat. #NET636001MC, PerkinElmer) for 1 h at 37°C. Nonspecific binding was determined by the addition of atropine (1 µM) during the incubation with [3H]-NMS. Binding reactions were terminated by rapid filtration over UniFilter GF/C plates (PerkinElmer) pre-soaked in binding buffer, followed by three rapid washes with ice-cold 0.9% NaCl using a filter plate system (Brandel). The plates were then air-dried and 50 µL Microscint-20 liquid scintillation fluid (PerkinElmer) was added to each well. Membrane bound radioactivity was then counted in a Topcount liquid scintillation counter (PerkinElmer). The specific bound counts (dpm) were expressed as fmol/mg protein.

Quantitative RT-PCR

Striata or cerebella from homozygous M_4 KO and R26-MetRS* (WT) mice (10-12 weeks old) were collected and total RNA was isolated using a RNeasy Plus Mini Kit (Cat. #74134, Qiagen), following the manufacturer's instructions. 1 µg total RNA was treated with DNase I (Cat. #79254, Qiagen) and used as template for cDNA synthesis with M-MLV reverse transcriptase (Cat. #28025013, Invitrogen), according to the manufacturer's recommendations. Real-time PCR was performed with probes for Chrm4 (forward 5'-GGAGAAGAAGGCCAAGACTCTGG and reverse 5'- GGCAGTCACACATTCACTGCCTG) and GAPDH as control gene (forward 5'-CGGATTTGGCCGTATTGGG and reverse 5'-CTCGCTCCTGGAAGATGG), 2.4 µl of the cDNA sample as a template and Fast SYBR Green Master Mix (Cat. #4385612, Applied Biosystems) on a Quantstudio 5 Real Time PCR system (ThermoFisher Scientific). The cycling conditions were 50°C for 2 min, 95°C for 10 min, followed by 45 cycles of 95°C for 15 s and 60°C for 1 min. Melting curve analyses were performed for all real-time PCR runs to confirm amplification of only one product and to eliminate false positive results. Quantitative analysis was performed on Quantstudio software (ThermoFisher Scientific), which calculated the threshold cycle (Ct) values. Expression of Chrm4 was defined relative to GAPDH using the $2-\Delta\Delta Ct$ method.

BONCAT purification for LC-MS/MS analysis

Cerebellar primary astrocytes from Chrm4[CRE]::MetRS* mice grown onto 10 cm dishes as before, were incubated with 4 mM ANL in combination with vehicle, oxotremorine (10 µM) or oxotremorine in combination with VU6028418 (5 µM) for 24 h at 37°C. Cells were lysed in PBS with 0.5% SDS, 0.2% Triton X-100 and proteinase inhibitors, briefly sonicated

with 2 pulses of 15 s at an amplitude of 40% (~50 W) and centrifuged at 10,000 x g for 10 min at 4°C. Supernatants were collected and protein concentration calculated with a BCA assay kit. 500 µg of protein were heated at 65°C for 5 min followed by incubation with 50 mM iodoacetamide for 2 h at room temperature in the dark. Bio-orthogonal non-canonical amino acid tagging (BONCAT) was performed using 150 µM Biotin-PEG4-alkyne (Cat. #764213, Sigma-Aldrich), 0.5 mM CuSO$_4$, 1 mM THPTA and 5 mM sodium ascorbate overnight at 4°C. Buffer exchange was done with PD SpinTrap G-25 columns (Cat. #28-9180-04, Cytvia) and biotinylation of ANL-labelled proteins was confirmed by Western Blot. Ultralink avidin beads (Cat. #53146, Pierce) were used to purify clicked proteins. After overnight incubation at 4°C, beads were loaded into micro-spin columns (Cat. #89879, Pierce) and washed three times with PBS and three washes with ultrapure water. Proteins were eluted with 125 µl of 0.1% Trifluoroacetic acid (TFA) followed by 125 µl of 0.1% TFA with 50% acetonitrile (ACN). Eluted proteins were digested with 4 µg Trypsin/Lys-C (Cat. #V5073, Promega) in S-Trap midi spin columns (Profiti) according to the manufacturer's instructions.

Dried peptides were resuspended in 5 % acetonitrile with 0.5 % formic acid for nanoflow liquid chromatography–electrospray ionisation-tandem mass spectrometry (nLC-ESI-MS/MS) analysis. The samples were loaded onto a C$_{18}$ Acclaim PepMap trapping column (ThermoFisher Scientific) using an UltiMate3000 RSLCnano System (ThermoFisher Scientific). Peptide separation was performed on a Pepmap C$_{18}$ reversed phase column (2 µm, 75 µm × 50 cm, pore size 100 Å, Cat. #164540, ThermoFisher Scientific) by a gradient formed by mobile phase A (0.1% formic acid) and B (80% ACN, 0.08% formic acid), with increasing concentrations of solvent B running from 4% for 10 min, 60% for 170 min to 99% for 15 min at a fixed solvent flow rate of 0.3 µl/min. Eluted peptides were electrosprayed into an Orbitrap Elite MS (ThermoFisher Scientific) using a NanoMate Triversa (Advion Biosciences) ion source at 1.7 kV. The mass spectrometer was operated using full-scan MS spectra (m/z 375–1,500) at a resolution of 120,000 RP, followed by top speed (3s) high energy collision dissociation (HCD) fragmentation detection of the top precursor ions. Ion fragmentation was performed by top speed (3s) high energy collision dissociation (HCD) and the top precursor ions were detected at a resolution of 60,000 in MS2 spectra. HCD energy and isolation width was adjusted depending on charge of the precursor ions (a width of 1.2 m/z with 40% HCD for 2$^+$ ions, 0.7 m/z with 38% for 3$^+$, and 0.5 m/z with 38% HCD for 4$^+$-6$^+$). The resulting MS/MS spectra were analysed for protein identification using Proteome Discoverer software 2.4 (ThermoFisher Scientific) and searched against the murine UniProtKB/Swiss-Prot database using Mascot software (Matrix Science) with peptide tolerance set to 10 ppm and fragment ion mass tolerance of 0.3 Da. The carbamidomethylation of cysteine was programmed as fixed modification, while the oxidation of methionine, the deamidation of asparagine and glutamine, the replacement of methionine for ANL-PEG-biotin and the iodination of tyrosine were set as variables. Proteins present in at least 2 independent

experiments were considered for GO enrichment analysis to identify overrepresented biological processes using PANTHER [65] with a Fisher's exact test and a cutoff of FDR ≤0.05.

Statistical analysis

All data were expressed as mean ± SEM of at least 3 independent experiments. Statistical significance was determined by two-tailed Student's t test, one-way ANOVA with Tukey's post hoc correction or two-way ANOVA followed by a Tukey's or Bonferroni's post hoc test, as indicated. For ERK1/2 phosphorylation assays the concentration–response data were plotted on a log axis, fitted to a three-parameter sigmoidal curve and extra sum-of-squares F test applied to determine whether differences in pEC_{50} existed. The Kaplan-Meier plot survival analyses were compared using the Gehan-Breslow-Wilcoxon test. Significance was considered a p value smaller than 0.05 (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$). Statistical analysis and graph plotting was performed using GraphPad Prism software.

Supplementary Table 2: Primary antibodies. ICC, immunocytochemistry; IHC, immunohistochemistry; WB, western blot.

Antibody	Supplier	Cat. No	Host	Dilution	Use
S100β	Proteintech	15146-1-AP	Rabbit	1:500	ICC
c-fos	Abcam	ab190289	Rabbit	1:1,000	ICC
GFAP	Abcam	ab7260	Rabbit	1:500	ICC
GFP	Abcam	ab6673	Goat	1:500	ICC
S100β	Proteintech	15146-1-AP	Rabbit	1:1,000	IHC
Calbindin-D28k	Proteintech	14479-1-AP	Rabbit	1:1,000	IHC
GFAP	Abcam	ab7260	Rabbit	1:1,000	IHC
GFP	Abcam	ab6673	Goat	1:1,000	IHC
GFP	Abcam	Ab6556	Rabbit	1:1,000	IHC
Iba1	Abcam	ab178846	Rabbit	1:1,000	IHC
MAP2	Biolegend	822501	Chicken	1:1,000	IHC
Myelin basic protein (MBP)	Proteintech	10458-1-AP	Rabbit	1:2,000	IHC
NDRG2	Abcam	ab174850	Rabbit	1:1,000	IHC
Apolipoprotein E (APO-E)	Abcam	ab183596	Rabbit	1:1,000	WB
Chrm4	Aviva	OASG0431	Rabbit	1:500	WB
Clusterin	R&D Systems	AF2747	Goat	1:2,000	WB
GFAP	Sigma-Aldrich	G3893	Mouse	1:1,000	WB
GFP	Invitrogen	A-11122	Rabbit	1:1,000	WB
Iba1	Abcam	ab178846	Rabbit	1:2,000	WB
IRDye 800CW Streptavidin	Li-Cor Biosciences	926-32230	-	1:5,000	WB
Myelin basic protein (MBP)	Proteintech	10458-1-AP	Rabbit	1:4,000	WB

Prion protein (Prp)	Abcam	ab61409	Mouse	1:1,000	WB
SerpinA3N	R&D Systems	AF4709	Goat	1:1,000	WB
Vimentin	R&D Systems	MAB21052	Mouse	1:2,000	WB
α-tubulin	Abcam	ab7291	Mouse	1:10,000	WB

Supplementary Table 3: Secondary antibodies. ICC, immunocytochemistry; IHC, immunohistochemistry; WB, western blot.

Antibody	Supplier	Cat. No	Host	Dilution	Use
Anti-Rabbit Alexa Fluor Plus 647	Invitrogen	A32795	Donkey	1:500	ICC
Anti-Chicken Alexa Fluo 594	Invitrogen	A11042	Goat	1:1,000	IHC
Anti-Rabbit Alexa Fluor 488	Invitrogen	A32731	Goat	1:1,000	IHC
Anti-Goat Alexa Fluor Plus 488	Invitrogen	A32814	Donkey	1:1,000/ 1:500	IHC/ ICC
Anti-Mouse Alexa Fluor Plus 594	Invitrogen	A32744	Donkey	1:1,000/ 1:500	IHC/ ICC
Anti-Rabbit Alexa Fluor Plus 594	Invitrogen	A32754	Donkey	1:1,000/ 1:500	IHC/ ICC
IRDye 680RD Anti-Goat	Li-Cor Biosciences	925-68074	Donkey	1:5,000	WB
IRDye 680RD Anti-Mouse	Li-Cor Biosciences	925-68072	Donkey	1:5,000	WB
IRDye 800CW Anti-Mouse	Li-Cor Biosciences	926-32212	Donkey	1:5,000	WB
IRDye 800CW Anti-Rabbit	Li-Cor Biosciences	925-32213	Donkey	1:5,000	WB

References

1. Kaul, I.*, et al.* Efficacy and safety of the muscarinic receptor agonist KarXT (xanomeline-trospium) in schizophrenia (EMERGENT-2) in the USA: results from a randomised, double-blind, placebo-controlled, flexible-dose phase 3 trial. *Lancet* **403**, 160-170 (2024).

2. Kaul, I.*, et al.* Efficacy and Safety of Xanomeline-Trospium Chloride in Schizophrenia: A Randomized Clinical Trial. *JAMA Psychiatry* **81**, 749-756 (2024).

3. Mei, F.*, et al.* Accelerated remyelination during inflammatory demyelination prevents axonal loss and improves functional recovery. *Elife* **5** (2016).

4. Wang, F.*, et al.* Enhancing Oligodendrocyte Myelination Rescues Synaptic Loss and Improves Functional Recovery after Chronic Hypoxia. *Neuron* **99**, 689-701 e685 (2018).

5. Wang, F.*, et al.* Myelin degeneration and diminished myelin renewal contribute to age-related deficits in memory. *Nat Neurosci* **23**, 481-486 (2020).

6. Chen, J.F.*, et al.* Enhancing myelin renewal reverses cognitive dysfunction in a murine model of Alzheimer's disease. *Neuron* **109**, 2292-2307 e2295 (2021).

7. Dwomoh, L.*, et al.* M(1) muscarinic receptor activation reduces the molecular pathology and slows the progression of prion-mediated neurodegenerative disease. *Sci Signal* **15**, eabm3720 (2022).

8. Abd-Elrahman, K.S., Colson, T.L., Sarasija, S. & Ferguson, S.S.G. A M1 muscarinic acetylcholine receptor-specific positive allosteric modulator VU0486846 reduces neurogliosis in female Alzheimer's mice. *Biomed Pharmacother* **173**, 116388 (2024).

9. Pannell, M.*, et al.* The subpopulation of microglia expressing functional muscarinic acetylcholine receptors expands in stroke and Alzheimer's disease. *Brain Struct Funct* **221**, 1157-1172 (2016).

10. Andre, C., Dos Santos, G. & Koulakoff, A. Muscarinic receptor profiles of mouse brain astrocytes in culture vary with their tissue of origin but differ from those of neurons. *Eur J Neurosci* **6**, 1702-1709 (1994).

11. Endo, F.*, et al.* Molecular basis of astrocyte diversity and morphology across the CNS in health and disease. *Science* **378**, eadc9020 (2022).

12. Zhang, Y.*, et al.* An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex. *J Neurosci* **34**, 11929-11947 (2014).

13. Yao, Z.*, et al.* A high-resolution transcriptomic and spatial atlas of cell types in the whole mouse brain. *Nature* **624**, 317-332 (2023).

14. Alvarez-Castelao, B.*, et al.* Cell-type-specific metabolic labeling of nascent proteomes in vivo. *Nat Biotechnol* **35**, 1196-1201 (2017).

15. Sjostedt, E.*, et al.* An atlas of the protein-coding genes in the human, pig, and mouse brain. *Science* **367** (2020).

16. Buckley, N.J., Bonner, T.I. & Brann, M.R. Localization of a family of muscarinic receptor mRNAs in rat brain. *J Neurosci* **8**, 4646-4652 (1988).

17. Heintz, N. Gene expression nervous system atlas (GENSAT). *Nat Neurosci* **7**, 483 (2004).

18. Vilaro, M.T., Wiederhold, K.H., Palacios, J.M. & Mengod, G. Muscarinic cholinergic receptors in the rat caudate-putamen and olfactory tubercle belong predominantly to the M_4 class: in situ hybridization and receptor autoradiography evidence. *Neuroscience* **40**, 159-167 (1991).

19. Arenander, A.T., de Vellis, J. & Herschman, H.R. Induction of c-fos and TIS genes in cultured rat astrocytes by neurotransmitters. *J Neurosci Res* **24**, 107-114 (1989).

20. Bubser, M.*, et al.* Selective activation of M_4 muscarinic acetylcholine receptors reverses MK-801-induced behavioral impairments and enhances associative learning in rodents. *ACS Chem Neurosci* **5**, 920-942 (2014).

21. Spock, M.*, et al.* Discovery of VU6028418: A Highly Selective and Orally Bioavailable M(4) Muscarinic Acetylcholine Receptor Antagonist. *ACS Med Chem Lett* **12**, 1342-1349 (2021).

22. Tsai, H.H.*, et al.* Regional astrocyte allocation regulates CNS synaptogenesis and repair. *Science* **337**, 358-362 (2012).

23. Cerrato, V. Cerebellar Astrocytes: Much More Than Passive Bystanders In Ataxia Pathophysiology. *J Clin Med* **9** (2020).

24. Gould, R.W.*, et al.* Cognitive enhancement and antipsychotic-like activity following repeated dosing with the selective M(4) PAM VU0467154. *Neuropharmacology* **128**, 492-502 (2018).

25. Foster, D.J.*, et al.* Antipsychotic-like Effects of M_4 Positive Allosteric Modulators Are Mediated by CB2 Receptor-Dependent Inhibition of Dopamine Release. *Neuron* **91**, 1244-1252 (2016).

26. Pancani, T.*, et al.* M_4 mAChR-mediated modulation of glutamatergic transmission at corticostriatal synapses. *ACS Chem Neurosci* **5**, 318-324 (2014).

27.   Moehle, M.S.*, et al.* Cholinergic Projections to the Substantia Nigra Pars Reticulata Inhibit Dopamine Modulation of Basal Ganglia through the M(4) Muscarinic Receptor. *Neuron* **96**, 1358-1372 e1354 (2017).

28.   Jeon, J.*, et al.* A subpopulation of neuronal M_4 muscarinic acetylcholine receptors plays a critical role in modulating dopamine-dependent behaviors. *J Neurosci* **30**, 2396-2405 (2010).

29.   Zimmer, T.S., Orr, A.L. & Orr, A.G. Astrocytes in selective vulnerability to neurodegenerative disease. *Trends Neurosci* **47**, 289-302 (2024).

30.   Endo, F. Deciphering the spectrum of astrocyte diversity: Insights into molecular, morphological, and functional dimensions in health and neurodegenerative diseases. *Neurosci Res* (2024).

31.   Moulson, A.J., Squair, J.W., Franklin, R.J.M., Tetzlaff, W. & Assinck, P. Diversity of Reactive Astrogliosis in CNS Pathology: Heterogeneity or Plasticity? *Front Cell Neurosci* **15**, 703810 (2021).

32.   Tahir, W., Thapa, S. & Schatzl, H.M. Astrocyte in prion disease: a double-edged sword. *Neural Regen Res* **17**, 1659-1665 (2022).

33.   Akhtar, S., Wenborn, A., Brandner, S., Collinge, J. & Lloyd, S.E. Sex effects in mouse prion disease incubation time. *PLoS One* **6**, e28741 (2011).

34.   Myslivecek, J., Farar, V. & Valuskova, P. M(4) muscarinic receptors and locomotor activity regulation. *Physiol Res* **66**, S443-S455 (2017).

35.   Chambers, N.E., Millett, M. & Moehle, M.S. The muscarinic M_4 acetylcholine receptor exacerbates symptoms of movement disorders. *Biochem Soc Trans* **51**, 691-702 (2023).

36.   Nunes, E.J., Addy, N.A., Conn, P.J. & Foster, D.J. Targeting the Actions of Muscarinic Receptors on Dopamine Systems: New Strategies for Treating Neuropsychiatric Disorders. *Annu Rev Pharmacol Toxicol* **64**, 277-289 (2024).

37.   Camandola, S. Astrocytes, emerging stars of energy homeostasis. *Cell Stress* **2**, 246-252 (2018).

38.   Gonzalez-Garcia, I. & Garcia-Caceres, C. Hypothalamic Astrocytes as a Specialized and Responsive Cell Population in Obesity. *Int J Mol Sci* **22** (2021).

39.   Yang, L., Qi, Y. & Yang, Y. Astrocytes control food intake by inhibiting AGRP neuron activity via adenosine A1 receptors. *Cell Rep* **11**, 798-807 (2015).

40.   Chen, N.*, et al.* Direct modulation of GFAP-expressing glia in the arcuate nucleus bi-directionally regulates feeding. *Elife* **5** (2016).

41.   Herrera Moro Chao, D.*, et al.* Hypothalamic astrocytes control systemic glucose metabolism and energy balance. *Cell Metab* **34**, 1532-1547 e1536 (2022).

42.   MacDonald, A.J., Holmes, F.E., Beall, C., Pickering, A.E. & Ellacott, K.L.J. Regulation of food intake by astrocytes in the brainstem dorsal vagal complex. *Glia* **68**, 1241-1254 (2020).

43.   Clyburn, C., Carson, K.E., Smith, C.R., Travagli, R.A. & Browning, K.N. Brainstem astrocytes control homeostatic regulation of caloric intake. *J Physiol* **601**, 801-829 (2023).

44.   Cai, P.*, et al.* Regulation of wakefulness by astrocytes in the lateral hypothalamus. *Neuropharmacology* **221**, 109275 (2022).

45.   Brancaccio, M.*, et al.* Cell-autonomous clock of astrocytes drives circadian behavior in mammals. *Science* **363**, 187-192 (2019).

46.   Nimmerjahn, A., Mukamel, E.A. & Schnitzer, M.J. Motor behavior activates Bergmann glial networks. *Neuron* **62**, 400-412 (2009).

47.   Paukert, M.*, et al.* Norepinephrine controls astroglial responsiveness to local circuit activity. *Neuron* **82**, 1263-1270 (2014).

48.   Corkrum, M.*, et al.* Dopamine-Evoked Synaptic Regulation in the Nucleus Accumbens Requires Astrocyte Activity. *Neuron* **105**, 1036-1047 e1035 (2020).

49.   Schweinhuber, S.K., Messerschmidt, T., Hansch, R., Korte, M. & Rothkegel, M. Profilin isoforms modulate astrocytic morphology and the motility of astrocytic processes. *PLoS One* **10**, e0117244 (2015).

50.   Lee, S.J., Seo, B.R. & Koh, J.Y. Metallothionein-3 modulates the amyloid beta endocytosis of astrocytes through its effects on actin polymerization. *Mol Brain* **8**, 84 (2015).

51.   Zhou, J.*, et al.* Astrocytic LRP1 enables mitochondria transfer to neurons and mitigates brain ischemic stroke by suppressing ARF1 lactylation. *Cell Metab* **36**, 2054-2068 e2014 (2024).

52.   Romeo, R., Glotzbach, K., Scheller, A. & Faissner, A. Deletion of LRP1 From Astrocytes Modifies Neuronal Network Activity in an in vitro Model of the Tripartite Synapse. *Front Cell Neurosci* **14**, 567253 (2020).

53.   Gogliotti, R.G.*, et al.* Total RNA Sequencing of Rett Syndrome Autopsy Samples Identifies the M(4) Muscarinic Receptor as a Novel Therapeutic Target. *J Pharmacol Exp Ther* **365**, 291-300 (2018).

54.   Achilly, N.P.*, et al.* Deleting Mecp2 from the cerebellum rather than its neuronal subtypes causes a delay in motor learning in mice. *Elife* **10** (2021).

55.   Patani, R., Hardingham, G.E. & Liddelow, S.A. Functional roles of reactive astrocytes in neuroinflammation and neurodegeneration. *Nat Rev Neurol* **19**, 395-409 (2023).

56.   Munoz-Castro, C. & Serrano-Pozo, A. Astrocyte-Neuron Interactions in Alzheimer's Disease. *Adv Neurobiol* **39**, 345-382 (2024).

57.   Carter, S.F.*, et al.* Astrocyte Biomarkers in Alzheimer's Disease. *Trends Mol Med* **25**, 77-95 (2019).

58.   Makarava, N., Mychko, O., Chang, J.C., Molesworth, K. & Baskakov, I.V. The degree of astrocyte activation is predictive of the incubation time to prion disease. *Acta Neuropathol Commun* **9**, 87 (2021).

59.   Gomeza, J.*, et al.* Generation and pharmacological analysis of M2 and M$_4$ muscarinic receptor knockout mice. *Life Sci* **68**, 2457-2466 (2001).

60.   Guyenet, S.J.*, et al.* A simple composite phenotype scoring system for evaluating mouse models of cerebellar ataxia. *J Vis Exp* (2010).

61.   Mallucci, G.*, et al.* Depleting neuronal PrP in prion infection prevents disease and reverses spongiosis. *Science* **302**, 871-874 (2003).

62.   Scarpa, M.*, et al.* Biased M1 muscarinic receptor mutant mice show accelerated progression of prion neurodegenerative disease. *Proc Natl Acad Sci U S A* **118** (2021).

63.   Wolfes, A.C. & Dean, C. Culturing In Vivo-like Murine Astrocytes Using the Fast, Simple, and Inexpensive AWESAM Protocol. *J Vis Exp* (2018).

64.   Llorens-Martin, M., Torres-Aleman, I. & Trejo, J.L. Pronounced individual variation in the response to the stimulatory action of exercise on immature hippocampal neurons. *Hippocampus* **16**, 480-490 (2006).

65.   Thomas, P.D.*, et al.* PANTHER: Making genome-scale phylogenetics accessible to all. *Protein Sci* **31**, 8-22 (2022).

Acknowledgments

The authors would like to thank the staff of the University of Glasgow Biomedical Services, genOway for generation of the muscarinic receptor genetically modified mice and the grant support of the Wellcome Trust (Collaborative Award 201529/Z/16/Z and Discovery Award 307319/Z/23/Z). The authors would also like to acknowledge the assistance of Dr. Karen Fairlie-Clarke to this work.

Author information

Authors and Affiliations

Centre for Translational Pharmacology, School of Molecular Biosciences, Advanced Research Centre, 11 Chapel Lane, University of Glasgow, Glasgow G11 6EW, Scotland, United Kingdom.

Gonzalo S. Tejeda, Collin Molloy, Graeme Milligan and Andrew B. Tobin.

Warren Centre for Neuroscience Drug Discovery, Department of Pharmacology, Innovation Park, 393 Nichol Mills Road, Franklin, Tennessee, 37027, USA.

Carrie K. Jones and Craig W. Lindsley.

Corresponding authors

Gonzalo S. Tejeda: email Gonzalo.Tejeda@glasgow.ac.uk
Andrew B. Tobin: andrew.tobin@glasgow.ac.uk

Contributions

GST conducted the experiments, developed the study design and wrote the paper, ABT lead the study design and wrote the paper, GM contributed to the study design and writing of the paper, CM managed the mouse colonies and breeding and helped to conduct the animal experiments, CWL contributed with the muscarinic antagonist VU6028418 and the M_4 mAChR positive allosteric modulator VU0467154 and CKJ contributed with pharmacokinetic, pharmacology and metabolism data of these ligands.

Competing interests

The authors declare no competing interests.

Fig. 1

Fig. 1. M$_4$ mAChR is expressed in neuronal and disperse astrocytic populations throughout the brain. a, Schematic overview of the breeding between mice driving CRE recombinase from the *Chrm4* promoter (Chrm4^{CRE}) with mice carrying the inducible transgene for expression of GFP and MetRS* L274G (R26-MetRS*). The distribution of M$_4$ mAChR expressing cells can be detected in the resulting Chrm4^{CRE}::MetRS* mice by GFP immunostaining (green) in a brain sagittal section. **b,** Representative confocal images showing co-immunostaining of GFP (green) with the neuronal marker MAP2 (magenta) in areas of the cortex, striatum and hippocampus. The white square insets indicate the enlarged areas showed in the panels below by each individual channel. **c,** Representative confocal images showing co-localisation of GFP expressing cells with the astrocyte marker GFAP (magenta) in the hypothalamus and the brainstem. Scale bar for panel **a** is 250 μm, and for **b** and **c**, 25 μm.

Fig. 2

Fig.2. The cerebellum contains a diverse array of cells expressing M$_4$ mAChR. a, Distribution of M4 mAChR expressing cells labelled with GFP staining in a sagittal section of the cerebellum of Chrm4^{CRE}::MetRS* mice. **b,** Co-immunostaining of GFP-labelled M$_4^+$ cells with Calbindin in the Purkinje layer of the cerebellum of Chrm4^{CRE}::MetRS* mice. **c,** Immunohistochemistry analysis of different populations of GFAP positive astrocytes (magenta) co-expressing M$_4$ mAChR (green) in the cerebellar cortex (left panel), cerebellar nuclei (CN, middle panel) and white matter region (right panel). Arrowheads indicate M$_4^+$ astrocytes. Scale bar for panels **a** and **b** is 250 μm, and for **c**, 25 μm.

Fig. 3

Fig. 3. Role of M$_4$ mAChR in cerebellar astrocytic cultures of Chrm4^{CRE}::MetRS*. a, Immunostaining of cerebral or cerebellar astrocyte cultures from Chrm4^{CRE}::MetRS* animals detecting M$_4$ expressing cells with GFP (green) and the astrocyte marker GFAP (magenta). **b,** Quantification of the proportion of M$_4^+$ astrocytes in cerebral and cerebellar cultures relative to the total astrocytes present in the preparation (n = 5 -9). **c,** The percentage of the population of M$_4^+$ astrocytes co-expressing c-fos was estimated in astrocyte cerebellar cultures after

stimulation with Oxotremorine (Oxo, 10 μM), a combination of Oxo and the M_4 mAChR PAM VU0467154 (VU154, 10 μM) or Oxo plus the M_4 mAChR antagonist VU6028418 (VU418, 5 μM) for 60 min (n= 11- 25). **d,** Representative confocal images showing M_4 expressing astrocytes (green) co-stained with c-fos (magenta) in cerebellar cultures after treatment with Vehicle (Control), Oxo or a combination of Oxo and VU154 or Oxo with VU418, as indicated in **c**. **e,** Venn diagram showing the newly synthesised proteins (NSPs) exclusively identified from M_4^+ cerebellar astrocytes after treatment with ANL (4 mM) in combination with Vehicle, Oxo (10 μM) or Oxo and VU418 (5 μM). **f,** Analysis of enriched GO annotations (Fisher's exact test performed with Panther v. 17.0) in M_4^+ cerebellar astrocytes after treatment with ANL in basal conditions (n = 2). The size of each circle represents the number of genes, and the colour intensity indicates the $-\log_{10} p$ value adjusted by Benjamini-Hochberg method assigned to the corresponding GO term. **g,** Gene ontology biological processes which are significantly enriched from the nascent proteins synthesized after activation of M_4 mAChR in M_4^+ cerebellar astrocytes (NSPs exclusively found after Oxo treatment and not identified when the M_4 antagonist is present). Graphs depicts means ± SEM and statistical significance was calculated by two-tailed unpaired t test (**b**), or one-way ANOVA followed by Tukey's pot hoc test (**c**) (*$p <$ 0.05, **$p <$ 0.01, ***$p <$ 0.01,). Scale bars, 25 μm.

Fig. 4

Fig. 4. Selective ablation of M$_4$$^+$ astrocytes leads to deficits in locomotor activity. a, Depiction showing the breeding overview between the Chrm4^{CRE} animals with the hemizygous Aldh1l1-eGFP,-DTA mice, that express eGFP under the astrocyte promoter *Aldh1l1* and an inducible diphteria toxin A when Cre recombinase is present. This mating results in Chrm4^{CRE}:: Aldh1l1^{DTA} mice (DTA^{M4}*) with ablated astrocytes that express the M$_4$ mAChR, and the control littermates Chrm4^{CRE}:: Aldh1l1^{WT} (WT^{M4}*), wild type for the floxed gene. **b,** Average animal weight per group over time (n = 6-10). **c,** Comparison of the performance on the accelerated rotarod test between DTA^{M4}* and WT^{M4}* mice during 4 consecutive weeks (n = 10). **d,** Activity of mice in and open field test measuring the time spent in the central area of the apparatus (n = 15-18). **e,** Total distance travelled over the whole area of the open field arena. **f,** Analysis of the anxiety-like behaviour of DTA^{M4}* and WT^{M4}* mice measuring the time spent in the open arms of an elevated plus maze (n = 15-18). **g,** Locomotor activity in DTA^{M4}* animals compared to WT^{M4}* littermates measured by telemetry. The graph depicts the profile of activity counts averaged over 3 days of recording and plotted every five min. Mice experience a 12/12 h light/dark cycle, with the dark cycle shown in the shaded area (n = 8). **h,** Comparison of the average telemetry activity counts during the 12-h light and dark phases between DTA^{M4}* and WT^{M4}* mice. **i,** Effect of VU0467154 (VU154, 10 mg/kg) or vehicle (Veh) to counteract the amphetamine-induced hyperlocomotion (2 mg/kg) in DTA^{M4}* and WT^{M4}* mice. **j,** Total distance travelled during the 60 min period following D-amphetamine administration. (n = 6-7). Data are means ± SEM and statistical significance was calculated by two-tailed unpaired *t* test or one-way ANOVA followed by Bonferroni's pot hoc test when analysis involved more than two groups (*$p < 0.05$, **$p < 0.01$).

Fig. 5

a

NBH | RML

GFP / GFAP

b

NBH

RML

DAPI / GFP / GFAP

c

M₄⁺ astrocytes in prion disease

Control (NBH) | 15 w.p.i. | 17 w.p.i. | 23 w.p.i.

Density (cells/mm³)
0   4,000

d

Density M₄⁺ astrocytes (cells/mm³)

- Control
- RML 15 wpi
- RML 17 wpi
- RML 23 wpi

Cortex Striatum Hippocampus Thalamus Hypothalamus Midbrain

*** ** *** *** ***

e

Density M₄⁺ astrocytes (cells/mm³)

NBH | 18 wpi | 21 wpi

RML

Fig 5. Regional and temporal expansion in the number of the M_4^+ astrocytes during mouse prion disease. a, Representative confocal images of GFP-labelled M_4^+ cells in coronal brain sections from Chrm4[CRE]::MetRS* animals after 20 w.p.i. of RML prion inoculation (right) or control animals treated with NBH (left). Panels below show a detail of M_4^+ astrocytes (arrowheads) co-stained with GFP (green) and GFAP (magenta) from the thalamic areas indicated with white square insets. **b,** Confocal images showing M_4^+ astrocytes co-stained with GFP (green) and GFAP (magenta) in the cerebellar nuclei of Chrm4[CRE]::MetRS* mice after 20 w.p.i. of RML prion disease (right) or NBH controls (left). **c,** Regional and temporal spreading pattern of the density of M_4^+ astrocytes at different stages of RML prion disease in Chrm4[CRE]::MetRS* mice plotted on brain anatomical maps. Darker red sections represent regions with higher densities in M_4^+ astrocytes, while not quantified areas are depicted in grey. **d,** Quantification of M_4^+ astrocytes density across brain regions in Chrm4[CRE]::MetRS* mice treated with RML prion after 15, 17 and 23 w.p.i. and compared to control animals inoculated with NBH (n = 2-3). **e,** Stereological counting of M_4^+ astrocytes density in the cerebellar nuclei of mice inoculated with RML prion (18 and 21 w.p.i.) or NBH (n = 5-7). Scale bar in **a** is 250 μm, and 25 μm in **b**. Data shown are represented as mean ± SEM. Statistical significance was calculated by two-way ANOVA followed by Bonferroni's post hoc test (***$p < 0.001$, **$p < 0.01$).

Fig. 6

Fig. 6. Ablation of M$_4$⁺ astrocytes prolongs survival and reduces Prp^{Sc} deposition in the cerebellum of prion-infected mice. a, Evaluation of progressive motor deficits by rotarod in prion-diseased DTA^{M4}* and WT^{M4}* mice from 12 w.p.i (n = 8-12). **b,** Burrowing behaviour (percentage of food pellets displaced from the tube) of control or prion-infected DTA^{M4}* and WT^{M4}* mice assessed from 12 w.p.i. (n = 5 for NBH, n = 10-12 for RML). **c,** Kaplan-Meier plot showing a significant increase in survival probability for RML-prion inoculated mice after ablation of M$_4$⁺ astrocytes (DTA^{M4}*) compared to wild-type littermates (WT^{M4}*) (n = 22 - 25). Curves were analysed with a Gehan-Breslow-Wilcoxon test (**p < 0.01). **d,** Immunoblot analysis of PK-resistant Prp^{Sc} from the striatum, cortex, hippocampus and cerebellum of DTA^{M4}* and WT^{M4}* mice inoculated with NBH or RML-prion. Evaluated samples were collected at terminal stages of prion diseased mice with equivalent survival times. **e,** Quantification

of Prp^{Sc} shows accumulation of the PK-resistant protein in all evaluated brain regions from diseased animals. Ablation of M$_4^+$ astrocytes significantly reduces the deposition of Prp^{Sc} in the cerebellum. Graph represents means ± SEM of Prp^{Sc} levels relative to α-tubulin expression (n = 4-6) expressed as a percentage of the values obtained for RML-prion diseased wild-type mice (WT^{M4}*). Statistical significance was calculated by two-way ANOVA followed by Tukey's post-hoc test (*$p < 0.05$, ***$p < 0.001$).