**Spatial transcriptomic analysis of HIV and tuberculosis coinfection in a humanized mouse**

**model reveals specific transcription patterns, immune responses and early morphological**

**alteration signaling**

Sitaramaraju Adduri[1#], Jose Alejandro Bohorquez[1#], Omoyeni Adejare[1], Diego Rincon[2], Torry

Tucker[1], Nagarjun V Konduru[1], Guohua Yi[1*]

1.  Department of Cellular and Molecular Biology, School of Medicine, The University of

Texas Health Science Center at Tyler, 11937 US HWY 271, Tyler, TX 75708, USA

2.  Unaffiliated pathologist

*Corresponding author: Guohua.yi@uttyler.edu

#Authors contributed equally to this work

**Abstract**

*Mycobacterium tuberculosis* (*Mtb*) and human immunodeficiency virus (HIV) coinfection is one

of the biggest public health concerns worldwide. Both pathogens are adept at modulating immune

response and, in the case of *Mtb*, even inducing structural modification of the affected tissue. The

present study aimed at understanding the early phenotypical and functional changes in immune

cell infiltration in the affected organ, using a humanized mouse model. The humanized mice were

infected with either HIV or *Mtb* in single infection, or with both pathogens in coinfection. Three

weeks after the infection, lung samples were collected, and spatial transcriptomics analysis was

performed. This analysis revealed high infiltration of $CD4^+$ T cells in *Mtb* infection, but not in HIV

or coinfection. Coinfected mice also showed a minimal number of NK cells compared to the other

groups. In addition to infection status, histological features also influenced the immune cell

infiltration pattern in the lungs. Two distinct airway regions with distinct immune cell abundance

patterns were detected by spatial transcriptome profiling.  A lymphoid cell aggregate detected in

coinfection lung exhibited distinct transcript profile. The cellular architecture in the lymphoid cell

aggregate did not follow the spatial patterns seen in mature granulomas. However, lymphoid cell

aggregates exhibited granuloma gene expression signatures, and pathways associated with reactive

oxygen species production, oxidative phosphorylation, and TGFβ and interferon signaling similar

to granulomas. This study revealed specific transcription patterns, immune responses and

morphological alteration signaling in the early stage of HIV and *Mtb* infections.

**Introduction**

Tuberculosis (TB), caused by *Mycobacterium tuberculosis* (*Mtb*), continues to be a major public health concern. It has been estimated that as much as 25% of the world population is infected with *Mtb*, although most of these cases will develop the latent form of TB (LTBI)[1]. One of the main features of *Mtb* is its ability to modulate immune response to generate the latent disease form. This immune modulation takes place in circulating immune cells, but also extends to tissue resident cells, where *Mtb* is able to induce structural modification of the hosts tissue to create specialized histological structures called granulomas[2,3]. These structures are composed of a necrotic core, surrounded by different immune cell populations that contain the pathogen, preventing its spread throughout the host and the development of active TB. Consequently, increasing interest has been granted to the cellular infiltration in lung tissue during *Mtb* infection, in general, and in granulomas, in particular[4-6]. The modification of cellular subsets in the affected organ and the inter-cellular communication taking place provides an explanation for how they favor or deter pathogen dissemination. It has been posed that there are multiple microenvironments present within granuloma structures that induce an immunoregulatory environment with differential signaling, as determined by pathway activation and cytokine production, playing a crucial role in the maintenance of the structure[7].

Accordingly, local or generalized immune alterations can lead to changes in the granuloma structures that can facilitate bacterial growth[8,9]. The Human immunodeficiency virus (HIV) is one of the most important disruptors of the immune response and causes instability in the granuloma structures[9]. Coinfection with HIV in *Mtb*-infected individuals severely increases the probability of developing active TB[10], due to the generalized immunosuppression caused by the depletion of CD4+ T cells, the main target for HIV infection, as well as the specific depletion of this cell subset

in the granulomas. Reports show that CD4$^+$ populations within granulomas are particularly

susceptible to HIV infection, due to increased expression of the CCR5 coreceptor[8]. Additionally,

given the importance of CD4$^+$ cells for CD8$^+$ cell maturation, HIV-induced CD4$^+$ cell depletion

reduces the circulating effector and memory CD8$^+$ cells and promotes immunosenescence in these

cells, rendering them dysfunctional[11].

Studies dedicated to understanding the cellular composition and intercellular networks

present locally in lung during *Mtb* infection and HIV/*Mtb* coinfection have focused on infection

models or samples in which the infections and structural changes are already set in the host, leaving

a gap in the knowledge of the early events taking place after infection[5-8,12]. This is partially due to

the use of stored human samples or non-human primate (NHP) animal models for these

experiments, which, though highly reliable, limit the availability of samples from early infection

timepoints. Recent reports show the feasibility of humanized mice as a model for the study of *Mtb*

and HIV[13,14]. These models are able to reproduce immune changes caused by infection, including

formation of lung granulomas. Given this, the aim of the present study was to evaluate the changes

in the cellular repertoire in the lung induced by HIV and *Mtb* in single infections, as well as in

coinfection, at an early timepoint in which granulomas have not been fully established.

Furthermore, we analyzed the transcriptomic changes in these cells, using a spatial approach, to

determine alterations in cytokine production in the affected areas. This allowed us to assess the

molecular processes taking place during infection, including evidence of changes in early

granuloma formation.

**Materials and methods**

**Generation of humanized mice**

Mice from the NOD.Cg-Prkdcscid Il2rgtm1Wjl Tg(CMV-IL3,CSF2,KITLG)1Eav

Tg(IL15)1Sz/J (NSG-SGM3-IL15) strain (The Jackson laboratory, Bar Harbor, ME), maintained

at the animal housing facility in the University of Texas Health Science Center at Tyler (UTHSCT)

were used for this study. Four to five weeks old mice were irradiated at a dose of 100cgy/mouse

for myeloablation. This was followed 6 hours later by intravenous (IV) inoculation with $2 \times 10^5$

human CD34$^+$ hematopoietic stem cells (STEMCELL technologies, Vancouver, Canada) per

mouse for humanization.

At 16 weeks after inoculation with human CD34$^+$ cells, humanization was confirmed in

accordance with a previously stablished protocol. Briefly, the whole blood samples were collected

by puncture of the submandibular vein and density gradient was used to separate peripheral blood

mononuclear cells (PBMCs). Once PBMCs were obtained, flow cytometry for specific human and

mouse immune cell markers was performed and humanization was confirmed for 13 mice. The

criteria for humanization included a positive human/mouse leukocyte ratio, as determined by

human CD45$^+$ vs. mouse CD45$^+$ expression, as well as the presence of multiple human immune

cell subsets, determined by markers for T-cells (CD3$^+$), B-cells (CD20$^+$), monocytes (CD14$^+$) and

natural killer (NK) cells (CD56$^+$), in accordance with a previously stablished protocol in our lab[13].

**Animal infection and experimental design**

Once mouse humanization was confirmed, the animals were randomly divided into 4

experimental groups (Figure 1A): A) No infection (n=3), B) HIV single infection (n=3), C) *Mtb*

single infection (n=3) and D) HIV/*Mtb* coinfection (n=4). Animals in groups C and D were infected

with aerosolized *Mtb*, H37Rv strain, using a Madison chamber[13,15]. Three non-humanized mice

were included in the infection chamber to confirm the infection dose. These three mice were

euthanized the following day and lung samples were collected, macerated and plated in 7H10

plates, supplemented with OADC (BD biosciences, Franklin Lakes, NJ). These mice confirmed

that the infection was successful, and the dose was determined to be ~40 CFU/lung.

Two days after the *Mtb* infection was carried out, mice in groups B and D were injected

intraperitoneally (IP) with $10^7$ TCID of HIV-1 BaL strain (obtained from NIH AIDS Reagent

Program). Two weeks after inoculation, whole blood sample was obtained and RNA was extracted

from plasma sample of these mice, using the NucleoSpin RNA isolation kit (Macherey-Nagel,

Allentown, PA), according to manufacturer's instructions. The viral RNA load in each animal was

determined by RT-qPCR, using a control standard (obtained from NIH AIDS Reagent Program)

with known quantities of HIV-1 genome copies, as previously described[16]. All HIV-inoculated

mice were positive for HIV RT-qPCR and showed similar viral RNA loads.

Three weeks after infection, all animals were humanely euthanized, in accordance with the

NIH guidelines for euthanasia of rodents using carbon dioxide, followed by confirmation by neck

dislocation. During necropsy, lung samples from each mouse were collected in accordance with

previous protocols. Briefly, once euthanasia was confirmed, the thoracic cavity was opened, and

the lungs were perfused and inflated by injection of 10% neutral buffered formalin (Thermo

Scientific, Waltham, MA) into the right heart ventricle and trachea, respectively. Afterwards,

samples were immediately submerged in the same buffer and kept at 4 °C until paraffin

embedding[17,18]. All animal procedures were approved by the UTHSCT Institutional Animal Care

and Use Committee (IACUC) (Protocol #763).

**10x Visium spatial transcriptome assay:**

Formalin fixed and paraffin embedded tissue sections were used to prepare sample slides

as per the 10x Genomics protocol (CG000518 and CG000520). RNA was isolated from the FFPE

122 sections using RNeasy FFPE kit (Qiagen, # 73504) following the kit protocol. Samples with

123 DV200 value of > 30% were selected for the spatial transcriptomics assay. The qualified samples

124 after H&E staining and image were decrosslinked based the protocol (CG000520) and the

125 sequencing libraries were prepared as per the manufacturer's protocol (CG000495). The quality of

126 the final libraries was examined using Agilent High Sensitive DNA Kit (#5067-4626) using

127 Agilent Bioanalyzer 2100 (Agilent Technologies, Santa Clara, USA). The qualified libraries were

128 sequenced using the paired-end sequencing on an Illumina Novaseq System (Illumina, Inc., USA).

129 **Visium spatial sequencing data analysis:**

130  With demultiplexed sequence fastq files and CytAssist tissue image as input, Space ranger

131 was used to map the reads to the latest human genome (hg38) and quantified read counts per spot.

132 The counts were normalized using 'sctranform' method in Seurat package[19]. For comparison

133 between different samples, the data was further normalized using minimum median counts

134 function. The samples were clustered and projected on UMAP. Find markers function was used

135 for identifying cluster specific markers and differentially expressed genes. Genes exhibiting a

136 minimum |log2FC| value of 2 at p.adj value of <0.05 were considered differentially expressed

137 genes. Pathway analysis was performed using GprofileR[20]. Loupe browser was used to examine

138 the spot-wise expression level of individual genes. Deconvolution of spatial gene expression

139 profiles were performed using RCTD package[21]. Single cell RNA sequencing (GSE192483) of

140 lung tissues with 18F-FDG avidity and nearby uninvolved tissues from six tuberculosis patients[22]

141 were used as reference for deconvolution analysis. Azimuth[23] was used for annotating the cells in

142 the reference dataset. Visualizations were created using ggplot2 package in R environment (4.1.1).

143 **Results**

**Immune cell infiltration patterns in the lungs are different for each infection type:**

We first investigated the effect of HIV and *Mtb* coinfection on immune responses in the lung. In our earlier study on humanized mice model of coinfection, we observed that the CD4$^+$ T cells depletion occurred rapidly after day 15 post infection and plateaued after day 28. Therefore, to investigate the status of immune responses during rapid declining of CD4$^+$ cells, we euthanized mice 21 days post infection and performed spatial transcriptome analysis of lung specimens from humanized mice with HIV infection, *Mtb* infection and coinfection using 10x Genomics Visium platform. The "No infection" group was used as control (Figure 1A). Excluding the poor-quality regions and empty spots, our spatial transcriptomics analysis detected expression of 18,000 genes across 3633, 2698, 2778, and 1397 spots in no infection, HIV, *Mtb*, and coinfection lungs, respectively.

Since TB is predominantly a lung infectious disease, we assume that examining the frequency of infiltration of different immune cells into lung will help us understand the immune response against the pathogen. CD4$^+$ cells play crucial role in mitigating *Mtb* infection by releasing signals that stimulate B cells to produce antibodies, cytotoxic T cells to kill infected cells, and macrophages to engulf and destroy pathogen. Therefore, we were also interested in understanding the effect of declining CD4$^+$ T cells titer on immune cell infiltration in lung tissues during coinfection. To compare the abundance of immune cell markers across all 4 infections groups, we normalized the spatial gene expression data from all samples using sctranform method followed by recorrecting the SCT counts minimum median counts. This data is suitable for comparing the gene expression between samples. As shown in Figure 1B, we observed that expression of CD4 is minimal in HIV single infection and in coinfection compared to control and *Mtb* single infection. This is expected as HIV infection and coinfection depletes CD4$^+$ T cell population. These results

suggest the possibility of existence of infection specific immune cell infiltration landscape in the

lungs.

To further investigate infiltration frequencies of immune cells into lungs across different

infection groups and control, we performed RCTD based cell type deconvolution analysis by

integrating the previously published single cell RNA sequencing data of human lung granuloma

(as detected by PET-CT) and their matched normal tissues of TB patients with our spatial

transcriptome data of all 4 infection groups. We first annotated the cell type labels of all cells in

scRNAseq data using Azimuth annotation database at the finest annotation level and transferred

these labels to our spatial transcriptome data during deconvolution analysis in 'full mode'. Since

Visium spots may span up to 6 individual cells, we opted to run deconvolution in 'full mode' as it

allows identification of any number of cells existing in each pixel. Corroborating with CD4 gene

expression in the tissues, cell type weights for CD4$^+$ T cells also showed the highest abundance of

CD4$^+$ T cells in *Mtb* infection while showing less abundance in HIV and coinfection mice (Figure

1C). Supporting previous reports, CD4$^+$ cells were less coinfection mice compared to HIV single

infection[24] (Figure 1C).  Surprisingly, the abundance of nonclassical monocytes reduced in all

infections compared to control (Figure 1C). NK cells abundance was less in the coinfection mice

compared to the rest of the three groups. NK cell abundance was the highest in *Mtb* single infection

lung (Figure 1C). The lower levels of NK cell infiltration in coinfection compared to *Mtb* single

infection were observed in an earlier study[24].  The distribution of cell type weights of myeloid

lineage cells such as alveolar macrophages, CCL3$^+$ alveolar macrophages, classical monocytes,

monocyte derived macrophages, interstitial macrophages was similar across all groups suggesting

that infection status has minimal or no effect in the infiltration of these cells (Figure S1).

Altogether, these results suggest that lung tissue infiltration of various immune cell types depend

on the type of infection.

**Lung immune cell infiltration patterns differ based on infection status as well as local**

**histological milieu**

To understand the distinct patterns of immune cell infiltration into lungs, we performed

clustering of Visium spots across all 4 infection states together using Uniform Manifold

Approximation and Projection (UMAP) based on 50 principal components at a resolution of 0.9.

We observed a total of 14 clusters across 4 samples with no infection (12), HIV (11), *Mtb* (12),

and coinfection (12) (Figure 2A, S2A). Clusters 0-11 were present in all 4 samples while cluster

12 was detected neither in HIV nor in coinfection. Interestingly, cluster 13 was detected

exclusively in the coinfection group (Figure 3, S2A).

Interestingly, UMAP projection (Figure 2A) revealed a broad separation of all 14 clusters

into 3 distinct groups. These three cluster groups are represented by all four infection groups

(Figure 2A). Majority of the clusters, including clusters 0-3, 5-8, 10, 11 and 13 (total 11 clusters)

were placed together on the right on UMAP projection as a single large group (Figure 2B). These

spots correspond to alveolar/interstitial compartment of the lung across all 4 infection categories

(Figure 2B, S2B). Therefore, we aggregated these 11 clusters into one large cluster namely

'alveolar/interstitial' cluster for further analysis. Clusters 4 and 12 were placed separately together

and far from alveolar/interstitial cluster on UMAP projection (Figure 2C; represented in green and

blue, respectively). These spots belong to areas spanning airways in all four lungs (Figure S2C;

represented in green and blue, respectively). Therefore, we combined clusters 4 and 12 into one

cluster called 'airway_1' (Figure S2C; represented in red). Similarly, cluster 9, another distinct

cluster far away from the rest on the UMAP projection also harbored spots representing airways

across all four groups. It was also renamed as 'airway_2' for further analysis. Although, both

airway-1 and airway_2 clusters represent airways, they were not combined because they appeared

distinct from each other on UMAP projection. Interestingly, clusters were separated based on tissue

location and histological features rather than infection status. Altogether, these results suggest that

immune infiltration is strongly influenced by histological features compared to infection status in

our study.

To investigate the immune infiltration pattern differences between different clusters

associated with different histological location, we used the cell type weights obtained in

deconvolution analysis described in Figure 1. Abundance of $CD4^+$ T cells varied marginally

between alveolar/interstitial, airway_1, and airway_2 clusters with alveolar/interstitial and

airway_2 clusters exhibiting high abundance compared to airway_1 (Figure S2D). In contrast, the

abundance of monocyte derived macrophages was high in airway-1 compared to the rest (Figure

S2D). Interestingly, airway-2 cluster is enriched with $CCL3^+$ alveolar macrophages which are

known to be effective in phagocytosis of bacteria and other antigens (Figure 2D)[25] whereas

alveolar macrophages without high expression of CCL3 were marginally more abundant in

airway_1 cluster (Figure 2D). In contrast, airway_2 cluster and alveolar/interstitium clusters

exhibited a lower abundance of nonclassical monocytes compared to the airway_1 (Figure 2D).

Alveolar/interstium cluster is enriched with $CD8^+$ T cells and classical monocytes while

type 2 dendritic cells were less (Figure 2D). B cell, mast cells and type 1 dendritic cells did not

show any difference in abundance among the three clusters (Figure S2E). Altogether, these results

suggest that different histological regions may harbor district immune cell types. However, it is

not clear whether this differential abundance contributes to different immune responses.

**Cellular organization and composition of lymphoid cell aggregates markedly different from typical mature tuberculosis granuloma**

Several previous studies investigated the cellular architecture of mature tuberculosis granuloma. Granuloma may appear as a tiny lymphoid cell aggregate at initial stages and will progress into a non-necrotic or central necrotic mature granuloma. Our understanding of the cellular architecture of lymphoid cell aggregates is limited compared to mature granuloma. Examination of hematoxylin (H&E) staining revealed the presence of a lymphoid aggregate in the region of tissue subjected to spatial transcriptome profiling from coinfection mice. Focusing exclusively on this region, we investigated 1) whether the cellular architecture of lymphoid aggregate is similar or different from the mature granuloma, and 2) whether the lymphoid aggregates show any transcriptomic changes characteristic of mature granuloma.

Surprisingly, we noticed that spots belonging to cluster 13, the only cluster on UMAP that was exclusively detected in coinfection, spanned the entirety of the lymphoid cell aggregate in this sample, suggesting that this lymphoid aggregate exhibit distinct transcription program compared to the rest of the tissues across all 4 infection groups (Fig 3A). This lymphoid aggregate is present in the interstitial region adjacent to a large airway and appears to have densely populated infiltrated lymphoid cells on H& E staining (Fig 3B). On UMAP projection, cluster 13 is located together within all clusters spanning interstitial region yet appears distinct from the rest of all interstitial clusters (Figure 3C).

Necrotic granulomas are structured around a central core of necrotic cell debris in which much of the bacteria are concentrated. The layers adjacent to necrotic core are mainly composed of macrophages[2]. Layers of epithelioid macrophages surround the necrotic core interspersed with other macrophage populations such as classical alveolar macrophages, monocyte derive

macrophages, lipid laden foamy macrophages, and giant cell[2]. A diversity of other immune cell

types is seen at the periphery as well as within the epithelioid layers of the granuloma. Non-

necrotic granulomas have similar structural arrangement except that they do not have necrotic

core[2]. However, cellular architecture and dynamics of these lymphoid aggregates are poorly

understood.

To decipher the cell type abundances in the Visium spots spanning the lymphoid cell

aggregate, we extracted cell type frequency predictions obtained from RCTD deconvolution

analysis. We labeled spots spanning the lymphoid cell aggregate as inner/core region, edge and

periphery, beginning from the center to the outer region (Figure 3D). To compare the abundance

of different immune cells predicted based on deconvolution analysis, we plotted a heatmap with

hierarchical clustering gene wise and spot wise (Figure 3E). Interestingly, we observed that the

spot wise clustering as shown by the column dendrogram separated peripheral spots from the inner

and edge spots accurately suggesting that cell distribution in these compartments is stratified based

on the location. In mature granuloma, B cells are present in the periphery while being absent or

minimal in the core. In contrast, we observed that in lymphoid aggregate, inner/core region spot is

highly abundant in B cells (Figure 3E). High abundance of B cells in the core region of the

lymphoid aggregate was also confirmed by the highest abundance of the mature B cell markers

like CD19 (Figure 3F), CD20 (MS4A1) (Figure S3A), and IgM (IGHM) (Figure S3B).

Plasmacytoid DCs are also highly enriched in the core and edge regions as shown in the heatmap

(Figure 3E), and corroborating with expression of plasmacytoid markers such as CD123 (IL3RA)

(Figure 3G), and CD303 (CLEC4C) (Figure S3C). Similarly, lymphoid aggregate harbors

nonclassical monocytes abundantly (Figure 3E). Nonclassical monocytes express high levels of

CD97 (ADGRE5), P2RX1, and Siglec10[26]. As expected, we observed a high abundance of CD97

(ADGRE5) (Figure S3D), P2RX1 (Figure S3E), and Siglec10 (Figure S3F) in the core or edge of

the lymphoid aggregate while their abundance being less in the periphery. We also observed

enrichment of plasma cells, DC2, CD4 T cells and nonclassical monocytes in the core of the lesion

(Figure 3E). In contrast, we observed NK cells, mast cells, macrophages, CD8 T cells, DC1 cells,

and classical monocytes on the periphery (Figure 3E).

Altogether, these results suggest that lymphoid cell aggregates may not essentially

originate with macrophage cells predominantly at the core. We speculate that granuloma

architecture may evolve into macrophage laden core and lymphocyte abundant periphery overtime.

**Lymphoid cell aggregates exhibited gene expression changes characteristic of granulomas**

Given that lymphoid cell aggregates did not show cellular architecture akin to granuloma,

we checked whether this lesion exhibits any transcriptomic changes seen in mature granulomas.

We examined the expression of genes that are part of gene sets related to granuloma in Molecular

Signatures Database (MSigDB). We analyzed the HP_GRANULOMA gene set consisting of

ACP5, CYBB, CYBC1, DNASE2, DOCK2, IRF8, NF1, and PIK3CG. Aggregated expression of

these genes was higher in the lymphoid aggregate region (Figure 4A) compared to the rest of the

lung tissue while individual genes also exhibited relatively high expression in the lymphoid

aggregate region (Figure S4). Similarly, another gene set named HP_ GRANULOMATOSIS

consisting of ASAH1, CTLA4, CYBA, CYBB, HLA-DPA1, HLA-DPB1, NCF1, NCF2, PRTN3,

and PTPN22 also exhibited higher expression in lymphoid aggregate compared to rest of the tissue

(Figure 4A, S5A). These results suggest that gene expression changes characteristic of mature

granulomas appear to emerge early into the development of the granuloma from lymphoid

aggregate.

Encouraged by these results, to understand the molecular changes occurring early stages

of the granuloma development, we performed pathway analysis to identify signaling pathways

enriched in the lymphoid cell aggregate region. We performed differential expression analysis

using Seurat 'find-markers' function to identify genes that were up or downregulated in cluster 13.

At a log2FC cut off 2 and p.adj value of 0.05, we detected 383 upregulated and 20 downregulated

genes. Pathway analysis was performed independently for both up and down-regulated genes.

Multiple pathway terms representing immune response to viral, *Mtb*, and bacterial infections, B

cell and NK cell activity, cytokine signaling, glucose metabolism and oxidative phosphorylation,

ROS generation, interferon signaling and activation of several Interferon regulatory factor

transcriptional factors were significantly enriched in genes upregulated in the lymphoid cell

aggregate (Figure 4B, Supplementary Table 1). Interestingly, oxidative phosphorylation and ROS

generation are crucial for pathogen killing inside granulomas[27]. Transcriptional targets of several

interferon regulatory factors (IRF5, IRF7, and IRF9) were enriched in upregulated genes (Figure

4B-C, Supplementary Table 1). Type I-interferon signaling is known to promote *Mtb* infection by

facilitating the release of neutrophil extracellular traps[28] and suppressing interferon gamma and

interleukin 1α/β signaling[29].

Intriguingly, similar analysis on downregulated genes revealed enrichment of multiple

pathways related to downregulation of inhibitory SMAD expression in lymphoid cell aggregate

suggesting that this lesion may harbor activated TGFβ signaling (Figure 4D, Supplementary Table

2). This was corroborated by decreased expression of SMAD6 which inhibits TGFβ signaling and

upregulation of TGFβ signaling ligands namely TGFβ1/2 in the same tissue region, suggesting that

TGFβ signaling is activated (Figure 4E). Altogether, these results suggest that lymphoid cell

aggregates may show several characteristics of granulomas even before they evolve into mature

granulomas.

**Discussion**

Novel spatial transcriptomic technologies  provide a powerful tool to assess molecular and

immunological changes in situ in various tissues, thus understanding the effects of infection on the

host and allowing for in-depth knowledge of how disease is developed or avoided[30,31]. In the

present study, these technologies were used to evaluate the early changes caused by HIV and *Mtb*

infections and coinfection in humanized mouse lung tissues. Although lung is the main target organ

for *Mtb*, it's also greatly affected by the generalized immunosuppression caused by HIV[9] due to

massive depletion of $CD4^+$ T cell population[8,9]. The increase in $CD4^+$ T cells after *Mtb* single

infection relates to increased migration to the affected sites to control the pathogens[4]. Meanwhile,

the severe local depletion of this CD4+ T cell subset in HIV-infected hu-mice (with or without *Mtb*

coinfection) is a staple of HIV that has been previously described[8] and found in the present work.

The approach used in the present study allowed for the detection of alterations in less studied

populations, mainly non-classical monocytes and NK cells. Interestingly, even though HIV and

*Mtb* single infection increased the NK cell population, when compared to the uninfected control,

coinfection had the opposite effect. A similar pattern was reported for certain phenotypes of NK

cells in HIV/*Mtb* coinfected patients and it is likely that alterations in this cell subset vary in

accordance with the phenotypes differentiated in the subject[32].

A considerable portion of the differences evidenced in cell populations presented in the

tissue corresponded not only with infection, but rather with different histological regions, with

areas including bronchi and major airways being clearly differentiated from interstitial tissue.

However, even within the clustered airway regions, viral and bacterial infection modified the

phenotypical landscape. This is shown by the high presence of alveolar macrophages expressing

CCL3 in the airway regions from all the infected animals, in contrast with the uninfected controls.

CCL3 is known to aid in the activation of alveolar macrophages and efforts to control bacterial

and viral pathogens[25,33]. Recently, it was also found to be increased in COVID-19 patients suffering

alveolar damage, supporting the important role of these cells for pulmonary immunity[34].

The finding of a structurally organized and clearly differentiated cell cluster with a unique

transcriptomic profile, in the sample from HIV/*Mtb* coinfected mouse, provides insight into the

early phenotypical and signaling changes induced by this type of infection. Spatial analysis showed

the presence of both lymphoid and myeloid cells in the lymphoid aggregates. Remarkably, despite

being infected with HIV and showing generalized CD4$^+$ T cell depletion in the tissue, this cell

subset was found within the structure, mainly in the inner part and accompanied by B cells and

plasmacytoid DCs. It should be noted that the manner in which the cell subsets are organized in

the lymphoid structure found in the HIV/*Mtb* coinfected sample is not compatible with that found

in mature granulomas, which generally have a necrotic core, surrounded by macrophages and

epithelioid cells with lymphocytes in the outer layer[2,3,7]. Conversely, the unique structure found in

the HIV/*Mtb* coinfected hu-mouse tissue lacks evidence of necrosis and the monocytic cells within

it are in the outer layer. This is likely due to the fact that this cell aggregate is in the early stages

of forming a granulomatous structure and the time after infection has not yet been sufficient to

generate necrosis. This is in line with previous findings in murine *Mtb* model which detected the

presence of inflammatory cell aggregates, mainly adjacent to circulation or bronchi as early as 2

weeks after infection, with the first necrotic changes being detected around 1 month after

infection[35,36]. Moreover, our results show that the transcriptomic alterations found in this

premature granuloma-like structure are compatible with the signaling reported for mature

granulomas, supporting the idea that the lymphoid aggregate found in this sample is in the early

stages of granuloma formation.

The process of granuloma establishment involves precise modulation of the interferon

pathway, mainly through the upregulation of multiple IRFs, as shown in the cells present in this

granuloma-like structure. Some of the IRF genes found to be upregulated in the lymphoid

aggregate from the present study have been previously reported to be crucial for granuloma

formation and maintenance, as IRF8 deficient mice proved unable to form these structures and

limit *Mtb* growth after infection[37]. However, the increased transcription of different IRF genes

suggests regulation of the IFN response using multiple mechanisms, including cell differentiation

and innate immunity. This is likely a response to the initial induction of type-I IFNs and activation

of the IFN pathway and IFN stimulated genes (ISGs) after infection, but the relevance of this

multiple approach to IFN modulation in more advanced stages of structural remodeling remains to

be elucidated. Reports suggest that mature granulomas (after 9 weeks post infection) do not show

upregulation of these pathways, while others report increased IRF transcription in these structures,

though mainly in non-immune cell types (fibroblasts)[6,38].

Consequently, the activation of immune response by the IFN pathway leads to increased

energy requirements in the immune cells present in the lymphoid aggregate, as exemplified by the

upregulation of oxidative phosphorylation related genes. This has been established as an important

metabolic consequence of both HIV and *Mtb* infections, as well as HIV/*Mtb* coinfection[39-41]. Our

results show that the generalized metabolic changes induced by HIV/*Mtb* coinfection also occurs

locally in the affected tissues and, specifically in the affected areas, starting at an early stage of the

infection.

In addition, contrasting with the regulation of the IFN pathway, our results show a lack of

TGFβ regulation. TGFβ inhibition of *Mtb*-specific CD4$^+$ T cells has been found to play a role in

the maintenance of *Mtb* within mature granuloma structures in rhesus macaque and mice models[42].

This cytokine and its associated pathways has also been proposed as a determinant of myeloid cell

differentiation in TB granulomas[43], while virtual modeling results have suggested an enhancement

of bacterial clearance in absence of TGFβ[44].

**Conclusions**

The modulation of immune response by pathogens is an important determinant for the

outcome of infection. This can take the form of generalized immunosuppression, as caused by

HIV, or more nuanced modulation of immune response, as induced by *Mtb*. The results from the

present study show the effects of these pathogens, whether in single infection or coinfection, in

lung tissue of a humanized mouse model at an early timepoint after inoculation. Our findings

suggest that alterations in cellular infiltration and structural remodeling of the tissue begin to take

place early after infection and closely resemble those that have been previously reported for

humans and other animal models. This further validates the use of the NSG-SGM3-IL15 mouse

strain as a reliable animal model that can reproduce multiple aspects of HIV and *Mtb* infections,

as well as HIV/*Mtb* coinfection. Nevertheless, future studies, using higher resolution spatial

transcriptomics, will further elucidate the transcriptomic changes in lung immune cell populations

at a single cell level. This information will provide further detail into specific changes in different

cell populations, both in terms of phenotype and signaling and open up new avenues of research

to generate intervention strategies that improve infection outcome in patients infected with HIV

and *Mtb*.

**Conflict of interest:**

Authors declare no conflicts of interest.

**Acknowledgements:**

We thank the technical support from the Cancer Prevention and Research Institute of Texas (CPRIT

RP240610). We thank the Texas Advanced Computing Center (TACC) at The University of Texas

at Austin for providing computational resources that have contributed to the research results

reported within this paper. We'd like to thank Joshua Kleam at the Histopathology core at UTHCT,

for his help with this study and Drs. Buka Samten and Amy Tvinnereim for their aid in the animal

experiment.

**Funding**

This work was supported by a NIH Common funds/National Institute of Allergy and Infectious

Diseases grant UG3AI150550, a National Institute of Allergy and Infectious Diseases grant

R01AI184551, and a National Heart, Lung, and Blood Institute grant R01HL125016 to GY

**References**

1    WHO. Global Tuberculosis Report. (WHO, Geneva, Switzerland, 2023).
2    Cronan, M. R. In the Thick of It: Formation of the Tuberculous Granuloma and Its
Effects on Host and Therapeutic Responses. *Frontiers in Immunology* **13** (2022).
https://doi.org:10.3389/fimmu.2022.820134
3    McCaffrey, E. F. *et al*. The immunoregulatory landscape of human tuberculosis
granulomas. *Nat Immunol* **23**, 318-329 (2022). https://doi.org:10.1038/s41590-021-
01121-x
4    Bromley, J. D. *et al*. CD4(+) T cells re-wire granuloma cellularity and regulatory
networks to promote immunomodulation following Mtb reinfection. *Immunity* **57**,
2380-2398 e2386 (2024). https://doi.org:10.1016/j.immuni.2024.08.002
5    Carow, B. *et al*. Spatial and temporal localization of immune transcripts defines
hallmarks and diversity in the tuberculosis granuloma. *Nat Commun* **10**, 1823 (2019).
https://doi.org:10.1038/s41467-019-09816-4
6    Krausgruber, T. *et al*. Single-cell and spatial transcriptomics reveal aberrant lymphoid
developmental programs driving granuloma formation. *Immunity* **56**, 289-306 e287
(2023). https://doi.org:10.1016/j.immuni.2023.01.014

7    Qiu, X. *et al*. Spatial transcriptomic sequencing reveals immune microenvironment
features of Mycobacterium tuberculosis granulomas in lung and omentum.
*Theranostics* **14**, 6185-6201 (2024). https://doi.org:10.7150/thno.99038
8    Foreman, T. W. *et al*. CD4 T cells are rapidly depleted from tuberculosis granulomas
following acute SIV co-infection. *Cell Rep* **39**, 110896 (2022).
https://doi.org:10.1016/j.celrep.2022.110896
9    Kaushal, D., Singh, D. K. & Mehra, S. Immune Responses in Lung Granulomas during
Mtb/HIV Co-Infection: Implications for Pathogenesis and Therapy. *Pathogens* **12**
(2023). https://doi.org:10.3390/pathogens12091120
10   Azevedo-Pereira, J. M. *et al*. HIV/Mtb Co-Infection: From the Amplification of Disease
Pathogenesis to an "Emerging Syndemic". *Microorganisms 2023, Vol. 11, Page 853* **11**
(2023-03-27). https://doi.org:10.3390/microorganisms11040853
11   Bohorquez, J. A., Jagannath, C., Xu, H., Wang, X. & Yi, G. T Cell Responses during
Human Immunodeficiency Virus/Mycobacterium tuberculosis Coinfection. *Vaccines*
*(Basel)* **12** (2024). https://doi.org:10.3390/vaccines12080901
12   Sawyer, A. J. *et al*. Spatial mapping reveals granuloma diversity and histopathological
superstructure in human tuberculosis. *J Exp Med* **220** (2023).
https://doi.org:10.1084/jem.20221392
13   Bohorquez, J. A. *et al*. A Novel Humanized Mouse Model for HIV and Tuberculosis Co-
infection Studies. *bioRxiv*, 2024.2003.2005.583545 (2024).
https://doi.org:10.1101/2024.03.05.583545
14   Lepard, M. *et al*. Comparing Current and Next-Generation Humanized Mouse Models
for Advancing HIV and HIV/Mtb Co-Infection Studies. *Viruses* **14** (2022).
https://doi.org:10.3390/v14091927
15   Feng, Y. *et al*. Exposure to Cigarette Smoke Inhibits the Pulmonary T-Cell Response to
Influenza Virus and<i>Mycobacterium tuberculosis</i>. *Infection and Immunity* **79**,
229-237 (2011). https://doi.org:10.1128/iai.00709-10
16   Butler, S. L., Hansen, M. S. T. & Bushman, F. D. A quantitative assay for HIV DNA
integration in vivo. *Nature Medicine* **7**, 631-634 (2001). https://doi.org:10.1038/87979
17   Morton, J. & Snider, T. A. Guidelines for collection and processing of lungs from aged
mice for histological studies. *Pathobiology of Aging & Age-related Diseases* **7**,
1313676 (2017). https://doi.org:10.1080/20010001.2017.1313676
18   Davenport, M. L., Sherrill, T. P., Blackwell, T. S. & Edmonds, M. D. Perfusion and
Inflation of the Mouse Lung for Tumor Histology. *J Vis Exp* (2020).
https://doi.org:10.3791/60605
19   Hao, Y. *et al*. Dictionary learning for integrative, multimodal and scalable single-cell
analysis. *Nat Biotechnol* **42**, 293-304 (2024). https://doi.org:10.1038/s41587-023-
01767-y
20   Kolberg, L., Raudvere, U., Kuzmin, I., Vilo, J. & Peterson, H. gprofiler2 -- an R package
for gene list functional enrichment analysis and namespace conversion toolset
g:Profiler. *F1000Res* **9** (2020). https://doi.org:10.12688/f1000research.24956.2
21   Cable, D. M. *et al*. Robust decomposition of cell type mixtures in spatial
transcriptomics. *Nat Biotechnol* **40**, 517-526 (2022). https://doi.org:10.1038/s41587-
021-00830-w

22    Wang, L. et al. Single-cell RNA-sequencing reveals heterogeneity and intercellular
crosstalk in human tuberculosis lung. *J Infect* **87**, 373-384 (2023).
https://doi.org:10.1016/j.jinf.2023.09.004

23    Hao, Y. et al. Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587
e3529 (2021). https://doi.org:10.1016/j.cell.2021.04.048

24    Xiao, G. et al. Uncovering the Bronchoalveolar Single-Cell Landscape of Patients With
Pulmonary Tuberculosis With Human Immunodeficiency Virus Type 1 Coinfection. *J
Infect Dis* **230**, e524-e535 (2024). https://doi.org:10.1093/infdis/jiae042

25    Lindell, D. M., Standiford, T. J., Mancuso, P., Leshen, Z. J. & Huffnagle, G. B.
Macrophage inflammatory protein 1alpha/CCL3 is required for clearance of an acute
Klebsiella pneumoniae pulmonary infection. *Infect Immun* **69**, 6364-6369 (2001).
https://doi.org:10.1128/IAI.69.10.6364-6369.2001

26    Wong, K. L. et al. Gene expression profiling reveals the defining features of the
classical, intermediate, and nonclassical human monocyte subsets. *Blood* **118**, e16-
31 (2011). https://doi.org:10.1182/blood-2010-12-326355

27    Song, E. et al. Chronic granulomatous disease: a review of the infectious and
inflammatory complications. *Clin Mol Allergy* **9**, 10 (2011).
https://doi.org:10.1186/1476-7961-9-10

28    Chowdhury, C. S. et al. Type I IFN-mediated NET release promotes Mycobacterium
tuberculosis replication and is associated with granuloma caseation. *Cell Host
Microbe* **32**, 2092-2111 e2097 (2024). https://doi.org:10.1016/j.chom.2024.11.008

29    Donovan, M. L., Schultz, T. E., Duke, T. J. & Blumenthal, A. Type I Interferons in the
Pathogenesis of Tuberculosis: Molecular Drivers and Immunological Consequences.
*Front Immunol* **8**, 1633 (2017). https://doi.org:10.3389/fimmu.2017.01633

30    Marx, V. Method of the Year: spatially resolved transcriptomics. *Nat Methods* **18**, 9-14
(2021). https://doi.org:10.1038/s41592-020-01033-y

31    Williams, C. G., Lee, H. J., Asatsuma, T., Vento-Tormo, R. & Haque, A. An introduction
to spatial transcriptomics for biomedical research. *Genome Med* **14**, 68 (2022).
https://doi.org:10.1186/s13073-022-01075-1

32    Gao, J. et al. Mtb/HIV co-infection immune microenvironment subpopulations
heterogeneity. *Int Immunopharmacol* **143**, 113341 (2024).
https://doi.org:10.1016/j.intimp.2024.113341

33    Sadee, W. et al. Human alveolar macrophage response to Mycobacterium
tuberculosis: immune characteristics underlying large inter-individual variability. *Res
Sq* (2023). https://doi.org:10.21203/rs.3.rs-2986649/v1

34    Garcia-Prieto, C. A. et al. Spatial transcriptomics unveils the *in situ* cellular and
molecular hallmarks of the lung in fatal COVID-19. *bioRxiv*, 2024.2007.2003.601404
(2024). https://doi.org:10.1101/2024.07.03.601404

35    Tsai, M. C. et al. Characterization of the tuberculous granuloma in murine and human
lungs: cellular composition and relative tissue oxygen tension. *Cell Microbiol* **8**, 218-
232 (2006). https://doi.org:10.1111/j.1462-5822.2005.00612.x

36    Driver, E. R. et al. Evaluation of a mouse model of necrotic granuloma formation using
C3HeB/FeJ mice for testing of drugs against Mycobacterium tuberculosis. *Antimicrob
Agents Chemother* **56**, 3181-3195 (2012). https://doi.org:10.1128/AAC.00217-12

Rocca, S. *et al.* Interferon regulatory factor 8-deficiency determines massive neutrophil recruitment but T cell defect in fast growing granulomas during tuberculosis. *PLoS One* **8**, e62751 (2013). https://doi.org:10.1371/journal.pone.0062751

Mehra, S. *et al.* Granuloma correlates of protection against tuberculosis and mechanisms of immune modulation by Mycobacterium tuberculosis. *J Infect Dis* **207**, 1115-1127 (2013). https://doi.org:10.1093/infdis/jis778

Herbert, C., Luies, L., Loots, D. T. & Williams, A. A. The metabolic consequences of HIV/TB co-infection. *BMC Infectious Diseases* **23** (2023). https://doi.org:10.1186/s12879-023-08505-4

Masson, J. J. R. *et al.* Assessment of metabolic and mitochondrial dynamics in CD4+ and CD8+ T cells in virologically suppressed HIV-positive individuals on combination antiretroviral therapy. *PLOS ONE* **12** (30 ago 2017). https://doi.org:10.1371/journal.pone.0183931

Singh, V. *et al.* Mycobacterium tuberculosis-Driven Targeted Recalibration of Macrophage Lipid Homeostasis Promotes the Foamy Phenotype. *Cell Host & Microbe* **12** (2012/11/15). https://doi.org:10.1016/j.chom.2012.09.012

Gern, B. H. *et al.* TGFβ restricts expansion, survival, and function of T cells within the tuberculous granuloma. *Cell Host & Microbe* **29**, 594-606.e596 (2021). https://doi.org:10.1016/j.chom.2021.02.005

Peters, J. M. *et al.* Systematic deconstruction of myeloid cell signaling in tuberculosis granulomas reveals IFN-γ, TGF-β, and time are associated with conserved myeloid diversity. *bioRxiv*, 2024.2005.2024.595747 (2024). https://doi.org:10.1101/2024.05.24.595747

Warsinske, H. C., Pienaar, E., Linderman, J. J., Mattila, J. T. & Kirschner, D. E. Deletion of TGF-beta1 Increases Bacterial Clearance by Cytotoxic T Cells in a Tuberculosis Granuloma Model. *Front Immunol* **8**, 1843 (2017). https://doi.org:10.3389/fimmu.2017.01843

 **Figure legends**

 **Figure 1. Cell type abundance variations between control, HIV, Mtb and coinfection mice as**

 **predicted by cell specific marker analysis and cell type deconvolution analysis.**

A) The experimental protocol. The number of mice was four per group. B) Expression levels of CD4 genes were shown across the four groups. D) Cell type weights for each spot in the samples based on RCTD analysis were plotted for each infection types. X axis shows cell types weights which are proportional to the cell type abundance in the tissue. Difference in cell weight distributions was computed using Kruskal Wallis test followed by Dunn's test for pairwise comparisons. P were adjusted for multiple hypothesis testing using Bonferroni correction method. Significance levels are indicated as *P < .05, **P < .01, ***P < .001, and ****P < .0001. Figure created using Biorender.com

**Figure 2. Identification clusters based on gene expression of all tissue spots from no infection, HIV, Mtb, and coinfection mice lungs using Seurat package.**

A) Clusters from all four samples projected on UMAP and color based on cluster identifier (left) and sample type (right). B) Alveolar/interstitium cluster (shown in blue), airway_1 (shown in blue/green) and airway-2 (shown in red) clusters on UMAP projection. D) Cell type weights obtained in RCTD deconvolution analysis were plotted for each cluster. Y axis shows cell types weights which are proportional to the cell type abundance in the corresponding cluster. Color key indicated the cluster identifier. Difference in cell weight distributions was computed using Kruskal Wallis test followed by Dunn's test for pairwise comparisons. P values were adjusted for multiple hypothesis testing using Bonferroni correction method. Significance levels are indicated as *P < .05, **P < .01, ***P < .001, and ****P < .0001.

**Figure 3: Characterization of the unique cluster (cluster 13) detected in HIV/Mtb coinfection**

**mouse lung.**

A) Spatial mapping of cluster 13 which was detected exclusively in coinfection lung. Clusters were

identified on UMPA projection of all Visium spots passing quality control from all four infection

groups combined. Spots representing cluster 13 were highlighted in blue. B) Hematoxylin staining

of lung region matching with cluster 13 location in coinfection lung. This region contains a circular

lymphoid aggregate formed by dense lymphocyte infiltration located within the interstitial region

and close to two airways. C) UMAP projection of all Visium spots passing quality control from all

four infection groups highlighting the location of cluster 13. D) Projection of the Visium spots

overlapping with lymphoid aggregate shown in Figure 3B. The spots were labeled as core/inner

region, edge, and periphery depending on the location and were highlighted according to the label

with different colors. E) Heatmap showing the cell type abundances all the spots highlighted in

Figure 3D. Columns show individual spots spanning the lymphoid aggregate. The rows show the

immune cells detected in the spots. Color intensities represent the abundance of each cell in each

spot. Row and column dendrograms represent the similarity between cell abundances and

individual spots, respectively, as computed using hierarchical clustering. F) Spatial map of

expression of CD19, a mature B cell marker within and around the lymphoid aggregate. Spot-wise

gene expression data was projected on hematoxylin staining image of the lung tissue. Color key

represent the level of transcript expression. G) Expression of CD123 (IL3RA), plasmacytoid

markers in the same region as described for Figure 3F.

**Figure 4: gene expression changes and pathway analysis of lymphoid aggregate**

A) Aggregated gene expression of granuloma and granulomatosis gene sets from MSigDB

projected on hematoxylin stain of lungs from coinfection mice. Spot-wise gene expression

aggregate was projected on lymphoid cell aggregate and its surround tissue. Color key represents

aggregate of expression values of all the genes in the granuloma gene set (DOCK2, CYBC1,

PIK3CG, CYBB, IRF8, NF1, DNASE2, and ACP5) and granulomatosis gene set (DOCK2,

CYBC1, PIK3CG, CYBB, IRF8, NF1, DNASE2, and ACP5). B) Top 20 pathways based on p.adj

identified in over representation analysis on genes upregulated in cluster 13 spanning lymphoid

cell aggregate in coinfection mice. Gene ratios were computed with the percentage of total DEGs

in the given pathway term. C) Expression levels of interferon regulatory factors enriched among

upregulated genes in lymphoid cell aggregate were projected individually on lymphoid cell

aggregate and its surround tissue. D) Top 20 pathways identified in over representation analysis

on gene down regulated in cluster 13 as shown like figure 4B. E) Down regulation of SMAD6, an

inhibitory SMAD and upregulation of TGFB1/2 in lymphoid cell aggregate. Gene expression

levels quantified in spatial transcriptomics analysis were projected on hematoxylin-stained tissue

section. A, C, E) Color key represents aggregate/individual gene expression values in log2 scale.

**Figure 1:**

623     **Figure 2:**

**Figure 3:**

627 **Figure 4:**

628